

**Lemma 3.1.7.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $f \in L^1_{loc}(\Omega)$  such that, for all  $\varphi \in \mathcal{D}(\Omega)$ ,  $\int f(x)\varphi(x)dx = 0$ . Then we have  $f = 0$ .*

*Proof.* Let  $K$  be a compact subset of  $\Omega$  and  $\chi \in \mathcal{D}(\Omega)$  equal to 1 on a neighborhood of  $K$  as in the lemma 3.1.3. With  $\phi$  as in the proposition 3.1.1, we get that  $\lim_{\epsilon \rightarrow 0_+} \phi_\epsilon * (\chi f) = \chi f$  in  $L^1(\mathbb{R}^n)$ . We have

$$(\phi_\epsilon * (\chi f))(x) = \int f(y) \underbrace{\chi(y)\phi((x-y)\epsilon^{-1})\epsilon^{-n}}_{=\varphi_x(y)} dy, \quad \text{supp } \varphi_x \subset K, \varphi_x \in \mathcal{D}(\Omega),$$

and from the assumption of the lemma, we obtain  $(\phi_\epsilon * (\chi f))(x) = 0$  for all  $x$ , implying  $\chi f = 0$  from the convergence result; the conclusion follows.  $\square$

We note that it makes sense to restrict a distribution  $T \in \mathcal{D}'(\Omega)$  to an open subset  $U \subset \Omega$ : just define

$$\langle T|_U, \varphi \rangle_{\mathcal{D}'(U), \mathcal{D}(U)} = \langle T, \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)}, \quad (3.1.7)$$

and  $T|_U$  is obviously a distribution on  $U$ . With this in mind, we can define the support of a distribution exactly as in (3.1.8).

**Definition 3.1.8.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$  and  $T \in \mathcal{D}'(\Omega)$ . We define the support of  $T$  as*

$$\text{supp } T = \{x \in \Omega, \forall U \text{ open } \in \mathcal{V}_x, T|_U \neq 0\}. \quad (3.1.8)$$

We define the  $C^\infty$  singular support of  $T$  as

$$\text{singsupp } T = \{x \in \Omega, \forall U \text{ open } \in \mathcal{V}_x, T|_U \notin C^\infty(U)\}. \quad (3.1.9)$$

Note that the support and the singular support are closed subset of  $\Omega$  since their complements in  $\Omega$  are open: we have

$$(\text{supp } T)^c = \{x \in \Omega, \exists U \text{ open } \in \mathcal{V}_x, T|_U = 0\}, \quad (3.1.10)$$

$$(\text{singsupp } T)^c = \{x \in \Omega, \exists U \text{ open } \in \mathcal{V}_x, T|_U \in C^\infty(U)\}. \quad (3.1.11)$$

A simple consequence of that definition is that, for  $T \in \mathcal{D}'(\Omega)$ ,  $\varphi \in \mathcal{D}(\Omega)$ ,

$$\text{supp } \varphi \subset (\text{supp } T)^c \implies \langle T, \varphi \rangle = 0. \quad (3.1.12)$$

### 3.1.3 First examples of distributions

#### The Dirac mass

We define for  $\varphi \in C_c^0(\mathbb{R}^n)$ ,  $\langle \delta_0, \varphi \rangle = \varphi(0)$ ; the property (3.1.5) is satisfied with  $C_K = 1, N_K = 0$ . We have  $\text{supp } \delta_0 = \{0\}$ . From this, the Dirac mass cannot be an  $L^1_{loc}$  function, otherwise, since it is 0 a.e., it would be 0. Let  $\phi, \epsilon$  as in the proposition 3.1.1: then we have from that proposition

$$\lim_{\epsilon \rightarrow 0_+} \int \phi_\epsilon(x)\varphi(x)dx = \varphi(0),$$

so that the Dirac mass appears as the weak limit of  $\epsilon^{-n}\phi(x\epsilon^{-1})$ .

### The simple layer

We consider in  $\mathbb{R}^n$  the hypersurface  $\Sigma = \{(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}, x_n = f(x')\}$ , where  $f \in C^1(\mathbb{R}^{n-1})$ . We define for  $\varphi \in C_c^0(\mathbb{R}^n)$ ,

$$\langle \delta_\Sigma, \varphi \rangle = \int_{\mathbb{R}^{n-1}} \varphi(x', f(x')) (1 + |\nabla f(x')|^2)^{1/2} dx'.$$

The property (3.1.5) is satisfied with  $C_K = \text{area}(\Sigma \cap K)$ ,  $N_K = 0$ ,  $\text{supp } \delta_\Sigma = \Sigma$ , and since  $\Sigma$  has Lebesgue measure 0 in  $\mathbb{R}^n$ , the simple layer potential cannot be an  $L_{\text{loc}}^1$  function.

### The principal value of $1/x$

We define for  $\varphi \in C_c^1(\mathbb{R})$ ,

$$\langle \text{pv } \frac{1}{x}, \varphi \rangle = \lim_{\epsilon \rightarrow 0^+} \int_{|x| \geq \epsilon} \frac{\varphi(x)}{x} dx. \quad (3.1.13)$$

Let us check that this limit exists. We have for parity reasons,

$$\begin{aligned} \int_{|x| \geq \epsilon} \frac{\varphi(x)}{x} dx &= \int_{\epsilon}^{+\infty} (\varphi(x) - \varphi(-x)) \frac{dx}{x} \\ &= [\ln x (\varphi(x) - \varphi(-x))]_{x=\epsilon}^{x=+\infty} - \int_{\epsilon}^{+\infty} (\varphi'(x) + \varphi'(-x)) \ln x dx \end{aligned}$$

and thus, using that  $\lim_{\epsilon \rightarrow 0^+} \epsilon \ln \epsilon = 0$ ,  $\ln |x| \in L_{\text{loc}}^1(\mathbb{R})$ , we get

$$\langle \text{pv } \frac{1}{x}, \varphi \rangle = - \int_0^{+\infty} (\varphi'(x) + \varphi'(-x)) \ln x dx = - \int_{\mathbb{R}} \varphi'(x) (\ln |x|) dx,$$

yielding  $|\langle \text{pv } \frac{1}{x}, \varphi \rangle| \leq \int_{\text{supp } \varphi'} |\ln |x|| dx \|\varphi'\|_{L^\infty}$ .

### 3.1.4 Continuity properties

**Definition 3.1.9.** Let  $\Omega$  be an open subset of  $\mathbb{R}^n$  and let  $(\varphi_j)_{j \geq 1}$  be a sequence of functions in  $C_c^\infty(\Omega)$ . We shall say that  $\lim_j \varphi_j = 0$  in  $C_c^\infty(\Omega)$  when the two following conditions are satisfied:

- (1) there exists a compact set  $K \subset \Omega$ , such that  $\forall j \geq 1, \text{supp } \varphi_j \subset K$ ,
- (2)  $\lim_j \varphi_j = 0$  in the Fréchet space  $C_K^\infty(\Omega)$ , i.e.  $\forall \alpha \in \mathbb{N}^n, \lim_j (\sup_{x \in K} |(\partial_x^\alpha \varphi_j)(x)|) = 0$ .

**Proposition 3.1.10.** Let  $\Omega$  be an open subset of  $\mathbb{R}^n$  and  $T$  be a linear form defined on  $C_c^\infty(\Omega)$ . The linear form  $T$  is a distribution on  $\Omega$  if and only if it is sequentially continuous.

*Proof.* Assuming  $|\langle T, \varphi \rangle| \leq C_K \max_{|\alpha| \leq N_K} \|\partial_x^\alpha \varphi\|_{L^\infty}$  for all  $\varphi \in C_K^\infty(\Omega)$  and all  $K$  compact  $\subset \Omega$  implies readily the sequential continuity. Conversely, if  $T$  does not satisfy (3.1.5), we have

$$\exists K_0 \text{ compact } \subset \Omega, \forall k \geq 1, \forall N \in \mathbb{N}, \exists \varphi_{k,N} \in C_{K_0}^\infty(\Omega), |\langle T, \varphi_{k,N} \rangle| > k \max_{|\alpha| \leq N} \|\partial_x^\alpha \varphi_{k,N}\|_{L^\infty}.$$

From the strict inequality, we infer that the function  $\varphi_{k,N}$  is not identically 0, and we may define

$$\psi_k = \frac{\varphi_{k,k}}{k \max_{|\alpha| \leq k} \|\partial_x^\alpha \varphi_{k,k}\|_{L^\infty}}, \quad \text{so that } |\langle T, \psi_k \rangle| > 1.$$

But the sequence  $(\psi_k)_{k \geq 1}$  converges to 0 since  $\text{supp } \psi_k \subset K_0$  and for  $|\beta| \leq k$ ,  $\|\partial_x^\beta \psi_k\|_{L^\infty} \leq 1/k$ , implying for each multi-index  $\beta$  that  $\lim_k \|\partial_x^\beta \psi_k\|_{L^\infty} = 0$ . The sequential continuity is violated since  $|\langle T, \psi_k \rangle| > 1$  and the converse is proven.  $\square$

**Definition 3.1.11.** Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $T \in \mathcal{D}'(\Omega)$  and  $N \in \mathbb{N}$ . The distribution  $T$  will be said of finite order  $N$  if

$$\exists N \in \mathbb{N}, \forall K \text{ compact } \subset \Omega, \exists C_K > 0, \forall \varphi \in C_K^\infty(\Omega), |\langle T, \varphi \rangle| \leq C_K \sup_{\substack{|\alpha| \leq N \\ x \in \mathbb{R}^n}} |(\partial_x^\alpha \varphi)(x)|. \quad (3.1.14)$$

The vector space of distributions of order  $N$  on  $\Omega$  will be denoted by  $\mathcal{D}'^N(\Omega)$ . The vector space  $\mathcal{D}'^0(\Omega)$  is called the space of Radon measures on  $\Omega$ .

**Proposition 3.1.12.** Let  $\Omega$  be an open subset of  $\mathbb{R}^n$  and  $m \in \mathbb{N}$ . The vector space  $\mathcal{D}'^m(\Omega)$  is equal to the sequentially continuous<sup>1</sup> linear forms on  $C_c^m(\Omega)$ : if  $T \in \mathcal{D}'^m(\Omega)$ , it can be extended to a sequentially continuous linear form on  $C_c^m(\Omega)$ . If  $T$  is a sequentially continuous linear form on  $C_c^m(\Omega)$ , then  $T \in \mathcal{D}'^m(\Omega)$ .

*Proof.* Let us first consider  $T \in \mathcal{D}'^m(\Omega)$ ,  $\varphi \in C_c^m(\Omega)$ . Applying the proposition 3.1.1, we find a sequence  $(\varphi_k)_{k \geq 1}$  in  $C_c^\infty(\Omega)$ , converging in  $C_c^m(\Omega)$  with limit  $\varphi$ . Since we may assume that all the functions  $\varphi_k$  and  $\varphi$  are supported in a fixed compact subset  $K$  of  $\Omega$ , we have, according to the estimate (3.1.14),

$$|\langle T, \varphi_k - \varphi_l \rangle| \leq C \max_{|\alpha| \leq m} \|\partial_x^\alpha (\varphi_k - \varphi_l)\|_{L^\infty} = Cp(\varphi_k - \varphi_l),$$

where  $p$  is the norm in the Banach space  $C_K^m(\Omega)$ . Since the sequence  $(\varphi_k)_{k \geq 1}$  converges in  $C_K^m(\Omega)$ , we get that the sequence  $(\langle T, \varphi_k \rangle)_{k \geq 1}$  is a Cauchy sequence in  $\mathbb{C}$ , thus converges; moreover, if for some compact subset  $L$  of  $\Omega$ ,  $(\psi_k)_{k \geq 1}$  is another sequence of  $C_L^m(\Omega)$  converging to  $\varphi$ , we have

$$|\langle T, \psi_k - \varphi_k \rangle| \leq C' \max_{|\alpha| \leq m} \|\partial_x^\alpha (\varphi_k - \psi_k)\|_{L^\infty} = C'p(\varphi_k - \psi_k) \leq C'p(\varphi_k - \varphi) + C'p(\varphi - \psi_k)$$

and  $\lim_k \langle T, \psi_k - \varphi_k \rangle = 0$  so that, we can extend the linear form to  $C_c^m(\Omega)$  by defining  $\langle T, \varphi \rangle = \lim_k \langle T, \varphi_k \rangle$ . We get also immediately that (3.1.14) holds with  $N = m$  and  $C_K^\infty(\Omega)$  replaced by  $C_K^m(\Omega)$ , so that  $T$  is obviously sequentially continuous.

Let us now consider a sequentially continuous linear form  $T$  on  $C_c^m(\Omega)$ ; reproducing the proof of the proposition 3.1.10, we get that the estimate (3.1.14) holds with  $N = m$ , proving that  $T \in \mathcal{D}'^m(\Omega)$ . The proof of the proposition is complete.  $\square$

**Remark 3.1.13.** We have already proven directly that functions in  $L_{\text{loc}}^1(\Omega)$  (see (3.1.6)), the Dirac mass and a simple layer (see the section 3.1.3) are distributions of order 0. It is an exercise left to the reader to prove that the distribution  $\text{pv } \frac{1}{x}$  defined in (3.1.13) is of order 1 and not of order 0.

<sup>1</sup>The convergence of a sequence in  $C_c^m(\Omega)$  is analogous to the convergence given in the definition 3.1.9, except that (2) is required in the Banach space  $C_K^m(\Omega)$ , i.e.  $|\alpha| \leq m$ .

### 3.1.5 Partitions of unity and localization

**Theorem 3.1.14** (Partition of unity). *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $K$  a compact subset of  $\Omega$  and  $\Omega_1, \dots, \Omega_m$  open subsets of  $\Omega$  such that  $K \subset \Omega_1 \cup \dots \cup \Omega_m$ . Then for  $1 \leq j \leq m$ , there exists  $\psi_j \in C_c^\infty(\Omega_j; [0, 1])$  and  $V$  open such that*

$$\Omega \supset V \supset K, \quad \forall x \in V, \quad \sum_{1 \leq j \leq m} \psi_j(x) = 1,$$

and for all  $x \in \Omega$ ,  $\sum_{1 \leq j \leq m} \psi_j(x) \in [0, 1]$ .

*Proof.* The case  $m = 1$  of the theorem is proven in the lemma 3.1.3. We consider now  $m > 1$  and we note that, since  $x \in K$  implies  $x \in$  one of the  $\Omega_j$ ,

$$K \subset \cup_{x \in K} B(x, r_x), \quad \bar{B}(x, r_x) \subset \text{one of the } \Omega_j, \quad r_x > 0.$$

From the compactness of  $K$ , we get that  $K \subset \cup_{1 \leq l \leq N} B(x_l, r_{x_l})$  and we may assume that

$$\begin{aligned} \bar{B}(x_l, r_{x_l}) &\subset \Omega_1, & \text{for } 1 \leq l \leq N_1, \\ \bar{B}(x_l, r_{x_l}) &\subset \Omega_2, & \text{for } N_1 < l \leq N_2, \\ &\dots\dots\dots \\ \bar{B}(x_l, r_{x_l}) &\subset \Omega_m, & \text{for } N_{m-1} < l \leq N_m = N. \end{aligned}$$

We define then the compact sets

$$K_1 = \cup_{1 \leq l \leq N_1} \bar{B}(x_l, r_{x_l}), \quad \dots, \quad K_m = \cup_{N_{m-1} < l \leq N_m} \bar{B}(x_l, r_{x_l}),$$

and we have  $K \subset \cup_{1 \leq j \leq m} K_j$ , and for each  $j$ ,  $K_j \subset \Omega_j$ . Using the lemma 3.1.3, we find  $\varphi_j \in C_c^\infty(\Omega_j; [0, 1])$  such that  $\varphi_j = 1$  on a neighborhood  $V_j (\subset \Omega_j)$  of  $K_j$ . We define then

$$\begin{aligned} \psi_1 &= \varphi_1, \\ \psi_2 &= \varphi_2(1 - \varphi_1), \\ &\dots\dots\dots \\ \psi_j &= \varphi_j(1 - \varphi_1) \dots (1 - \varphi_{j-1}), \end{aligned}$$

so that  $\psi_j \in C_c^\infty(\Omega_j; [0, 1])$  and we have

$$\sum_{1 \leq j \leq m} \psi_j = \sum_{1 \leq j \leq m} \varphi_j \left( \prod_{1 \leq k < j} (1 - \varphi_k) \right) = 1 - \prod_{1 \leq k \leq m} (1 - \varphi_k), \quad (3.1.15)$$

since the formula (second equality above) is true for  $m = 1$  and inductively,

$$\begin{aligned} \sum_{1 \leq j \leq m+1} \varphi_j \left( \prod_{1 \leq k < j} (1 - \varphi_k) \right) &= 1 - \prod_{1 \leq k \leq m} (1 - \varphi_k) + \varphi_{m+1} \prod_{1 \leq k \leq m} (1 - \varphi_k) \\ &= 1 - (1 - \varphi_{m+1}) \prod_{1 \leq k \leq m} (1 - \varphi_k) = 1 - \prod_{1 \leq k \leq m+1} (1 - \varphi_k). \end{aligned}$$

We have thus for  $x \in \cup_{1 \leq j \leq m} V_j$  (which is a neighborhood of  $K$  in  $\Omega$ ), using (3.1.15) and  $\varphi_j = 1$  on  $V_j$ ,  $\sum_{1 \leq j \leq m} \psi_j(x) = 1$ . On the other hand, (3.1.15) and  $\varphi_j$  valued in  $[0, 1]$  show that  $\sum_{1 \leq j \leq m} \psi_j(x) \in [0, 1]$  for all  $x$ . The proof is complete.  $\square$

**Theorem 3.1.15.** *Let  $\Omega$  be an open set of  $\mathbb{R}^n$  and  $(\Omega_j)_{j \in J}$  be an open covering of  $\Omega$ : each  $\Omega_j$  is open and  $\cup_{j \in J} \Omega_j = \Omega$ . Let us assume that for each  $j \in J$ , we are given  $T_j \in \mathcal{D}'(\Omega_j)$  in such a way that*

$$T_j|_{\Omega_j \cap \Omega_k} = T_k|_{\Omega_j \cap \Omega_k}. \quad (3.1.16)$$

*Then there exists a unique  $T \in \mathcal{D}'(\Omega)$  such that for all  $j \in J$ ,  $T|_{\Omega_j} = T_j$ .*

*Proof.* Uniqueness: if  $T, S$  are such distributions, we get that  $(T - S)|_{\Omega_j} = 0$ , so that for all  $j \in J$ ,  $\Omega_j \subset (\text{supp}(T - S))^c$  and thus  $\Omega = \cup_{j \in J} \Omega_j \subset (\text{supp}(T - S))^c$ , i.e.  $T - S = 0$ .

Existence: let  $\varphi \in \mathcal{D}(\Omega)$  and let us consider the compact set  $K = \text{supp} \varphi$ . We have  $K \subset \cup_{j \in M} \Omega_j$  with  $M$  a finite subset of  $J$ . Using the theorem on partitions of unity, we find some function  $\psi_j \in C_c^\infty(\Omega_j)$  for  $j \in M$  such that  $\sum_{j \in M} \psi_j = 1$  on a neighborhood of  $K$ . As a consequence, we have  $\varphi = \sum_{j \in M} \psi_j \varphi$  and we define

$$\langle T, \varphi \rangle = \sum_{j \in M} \langle T_j, \psi_j \varphi \rangle.$$

The required estimates (3.1.5) are easily checked, but the linearity and the independence with respect to the decomposition deserve some attention. Assume that we have  $\varphi = \sum_{k \in N} \phi_k \varphi$ , where  $N$  is a finite subset of  $J$  and  $\phi_k \in C_c^\infty(\Omega_k)$ : we have

$$\sum_{k \in N} \langle T_k, \phi_k \varphi \rangle = \sum_{j \in M, k \in N} \langle T_k, \phi_k \psi_j \varphi \rangle \underbrace{=}_{\text{from (3.1.16)}} \sum_{j \in M, k \in N} \langle T_j, \phi_k \psi_j \varphi \rangle = \sum_{j \in M} \langle T_j, \psi_j \varphi \rangle,$$

proving that  $T$  is defined independently of the decomposition. The linearity follows at once. The proof is complete.  $\square$

### 3.1.6 Weak convergence of distributions

We have not defined a topology on the space of test functions  $\mathcal{D}(\Omega)$ , although we gave the definition of convergence of a sequence (see the definition 3.1.9); we shall need also a simple notion of weak-dual convergence of a sequence of distributions, which is the  $\sigma(\mathcal{D}', \mathcal{D})$  convergence.

**Definition 3.1.16.** *Let  $\Omega$  be an open set of  $\mathbb{R}^n$ ,  $(T_j)_{j \geq 1}$  be a sequence of  $\mathcal{D}'(\Omega)$  and  $T \in \mathcal{D}'(\Omega)$ . We shall say that  $\lim_j T_j = T$  in the weak-dual topology if*

$$\forall \varphi \in \mathcal{D}(\Omega), \quad \lim_j \langle T_j, \varphi \rangle = \langle T, \varphi \rangle. \quad (3.1.17)$$

**Remark 3.1.17.** We have already seen (see the section 3.1.3) that for  $\rho \in C_c^\infty(\mathbb{R}^n)$ ,  $\epsilon > 0$ ,  $\rho_\epsilon(x) = \epsilon^{-n} \rho(x \epsilon^{-1})$ ,  $\lim_{\epsilon \rightarrow 0+} \rho_\epsilon = \delta_0 \int \rho(t) dt$ . Moreover, on  $\mathcal{D}'(\mathbb{R})$ , we have with  $T_\lambda(x) = e^{i\lambda x}$ ,  $\lim_{\lambda \rightarrow +\infty} T_\lambda = 0$  since for  $\varphi \in \mathcal{D}(\mathbb{R})$ ,

$$\int_{\mathbb{R}} e^{i\lambda x} \varphi(x) dx = (i\lambda)^{-1} \int_{\mathbb{R}} \frac{d}{dx} (e^{i\lambda x}) \varphi(x) dx = -(i\lambda)^{-1} \int_{\mathbb{R}} e^{i\lambda x} \varphi'(x) dx.$$

**Theorem 3.1.18.** *Let  $\Omega$  be an open set of  $\mathbb{R}^n$ ,  $(T_j)_{j \geq 1}$  be a sequence of  $\mathcal{D}'(\Omega)$  such that, for all  $\varphi \in \mathcal{D}(\Omega)$ , the (numerical) sequence  $(\langle T_j, \varphi \rangle)_{j \geq 1}$  converges. Defining the linear form  $T$  on  $\mathcal{D}(\Omega)$ , by  $\langle T, \varphi \rangle = \lim_j \langle T_j, \varphi \rangle$ , we obtain that  $T$  belongs to  $\mathcal{D}'(\Omega)$ .*

*Proof.* This is an important consequence of the Banach-Steinhaus theorem 2.1.8; let us consider a compact subset  $K$  of  $\Omega$ . Then defining  $T_{j,K}$  as the restriction of  $T_j$  to the Fréchet space  $\mathcal{D}_K(\Omega)$ , we see that the assumptions of the corollary 2.1.8 are satisfied since  $T_{j,K}$  belongs to the topological dual of  $\mathcal{D}_K(\Omega)$ , according to the remark 3.1.6. As a consequence the restriction of  $T$  to  $\mathcal{D}_K(\Omega)$  belongs to the topological dual of  $\mathcal{D}_K(\Omega)$  and from the same remark 3.1.6, it gives that  $T \in \mathcal{D}'(\Omega)$ .  $\square$

**N.B.** The reader may note that we have used  $E = \mathcal{D}(\Omega) = \cup_{j \in \mathbb{N}} \mathcal{D}_{K_j}(\Omega) = \cup_j E_j$ , and that our definition of the topological dual of  $E$  as linear forms  $T$  on  $E$  such that, for all  $j$ ,  $T|_{E_j} \in$  the topological dual of the Fréchet space  $E_j$ . This structure allows us to use the Banach-Steinhaus theorem, although we have not defined a topology on  $E$ ; this observation is a good introduction to the more abstract setting of  $LF$  spaces, the so-called inductive limits of Fréchet spaces.

## 3.2 Differentiation of distributions, multiplication by $C^\infty$ functions

### 3.2.1 Differentiation

**Definition 3.2.1.** *Let  $\Omega$  be an open set of  $\mathbb{R}^n$  and  $T \in \mathcal{D}'(\Omega)$ . We define the distributions  $\partial_{x_j} T$  and for a multi-index  $\alpha \in \mathbb{N}^n$  (see (2.3.6)),  $\partial_x^\alpha T$  by*

$$\langle \partial_{x_j} T, \varphi \rangle = -\langle T, \partial_{x_j} \varphi \rangle, \quad \langle \partial_x^\alpha T, \varphi \rangle = (-1)^{|\alpha|} \langle T, \partial_x^\alpha \varphi \rangle. \quad (3.2.1)$$

We note that  $\partial_x^\alpha T$  is indeed a distribution on  $\Omega$ , since the mappings  $\varphi \mapsto \partial_x^\alpha \varphi$  are continuous on each Fréchet space  $\mathcal{D}_K(\Omega)$ .

**Remark 3.2.2.** If  $\lim_j T_j = T$  in the weak-dual topology of  $\mathcal{D}'(\Omega)$ , then, for all multi-indices  $\alpha$ ,  $\lim_j \partial_x^\alpha T_j = \partial_x^\alpha T$  (in the weak-dual topology): we have, for each  $\varphi \in \mathcal{D}(\Omega)$ ,

$$\langle \partial_x^\alpha T_j, \varphi \rangle = (-1)^{|\alpha|} \langle T_j, \partial_x^\alpha \varphi \rangle \longrightarrow_{j \rightarrow +\infty} (-1)^{|\alpha|} \langle T, \partial_x^\alpha \varphi \rangle = \langle \partial_x^\alpha T, \varphi \rangle.$$

**Remark 3.2.3.** If  $u \in C^1(\Omega)$ , its derivative  $\partial_{x_j} u$  as a distribution coincides with the distribution defined by the continuous function  $\partial u / \partial x_j$ ; for  $\varphi \in \mathcal{D}(\Omega)$ ,

$$\langle \partial_{x_j} u, \varphi \rangle = -\langle u, \partial_{x_j} \varphi \rangle = -\int u(x) \frac{\partial \varphi}{\partial x_j}(x) dx = \int \frac{\partial u}{\partial x_j}(x) \varphi(x) dx = \langle \frac{\partial u}{\partial x_j}, \varphi \rangle.$$

Also, if  $u, v \in C^0(\Omega)$  are such that  $\partial_{x_1} u = v$  in  $\mathcal{D}'(\Omega)$ , then the function  $u$  admits  $v$  as a partial derivative with respect to  $x_1$ . To prove this, we may assume that  $u, v$  are both compactly supported in  $\Omega$ : in fact it is enough to prove that for  $\chi \in C_c^\infty(\Omega)$

identically equal to 1 near a point  $x_0$ , the function  $\chi u$  (compactly supported) has a partial derivative with respect to  $x_1$  which is  $\chi v + u\partial_{x_1}\chi$  (compactly supported) and we know that in  $\mathcal{D}'(\Omega)$  we have

$$\langle \partial_{x_1}(\chi u), \varphi \rangle = -\langle u, \chi \partial_{x_1} \varphi \rangle = -\langle u, \partial_{x_1}(\chi \varphi) \rangle + \langle u, \varphi \partial_{x_1} \chi \rangle = \langle \partial_{x_1} u, \chi \varphi \rangle + \langle u \partial_{x_1} \chi, \varphi \rangle$$

which implies a particular case of Leibniz' formula  $\partial_{x_1}(\chi u) = \chi \partial_{x_1} u + u \partial_{x_1} \chi = \chi v + u \partial_{x_1} \chi$ . Assuming then that  $u, v$  are compactly supported, we have from the proposition 3.1.1,  $u = \lim_{\epsilon} (u * \phi_{\epsilon})$  in  $C_c^0(\Omega)$  and the functions  $u * \phi_{\epsilon} \in C_c^{\infty}(\Omega)$ . Also we have, with the ordinary differentiation,

$$(\partial_{x_1}(u * \phi_{\epsilon}))(x) = \int u(y) (\partial_{x_1} \phi_{\epsilon})(x-y) dy = \langle u(\cdot), -\partial_{y_1}(\phi_{\epsilon}(x-\cdot)) \rangle = \int v(y) \phi_{\epsilon}(x-y) dy,$$

and  $\lim_{\epsilon} (v * \phi_{\epsilon}) = v$  in  $C_c^0(\Omega)$ . As a result the sequences  $(u * \phi_{\epsilon}), (\partial_{x_1}(u * \phi_{\epsilon}))$  are both uniformly converging sequences of (compactly supported) continuous functions with respective limits  $u, v$ , and this implies that the continuous function  $u$  has  $v$  as a partial derivative with respect to  $x_1$ .

### 3.2.2 Examples

Defining the Heaviside function  $H$  as  $\mathbf{1}_{\mathbb{R}_+}$ , we get

$$H' = \delta_0 \tag{3.2.2}$$

since for  $\varphi \in \mathcal{D}(\mathbb{R})$ , we have  $\langle H', \varphi \rangle = -\langle H, \varphi' \rangle = -\int_0^{+\infty} \varphi'(t) dt = \varphi(0)$ . Still in one dimension, we have

$$\langle \delta_0^{(k)}, \varphi \rangle = (-1)^k \varphi^{(k)}(0), \tag{3.2.3}$$

since it is true for  $k = 0$  and inductively  $\langle \delta_0^{(k+1)}, \varphi \rangle = -\langle \delta_0^{(k)}, \varphi' \rangle = -(-1)^k \varphi^{(k)}(0) = (-1)^{k+1} \varphi^{(k+1)}(0)$ . Looking at the definition (3.1.13), we see that we have proven

$$\text{pv}\left(\frac{1}{x}\right) = \frac{d}{dx}(\ln|x|), \quad (\text{distribution derivative}). \tag{3.2.4}$$

Let  $f$  be a finitely-piecewise  $C^1$  function defined on  $\mathbb{R}$ : it means that there is an increasing finite sequence of real numbers  $(a_n)_{1 \leq n \leq N}$ , so that  $f$  is  $C^1$  on all closed intervals  $[a_n, a_{n+1}]$  for  $1 \leq n < N$  and on  $]-\infty, a_1]$  and  $[a_N, +\infty[$ . In particular, the function  $f$  has a left-limit  $f(a_n^-)$  and a right-limit  $f(a_n^+)$  which may be different. Let us compute the distribution derivative of  $f$ ; for  $\varphi \in \mathcal{D}(\mathbb{R})$ , since  $f$  is locally integrable, we have, setting  $a_0 = -\infty, a_{N+1} = +\infty$ ,

$$\begin{aligned} \langle f', \varphi \rangle &= -\langle f, \varphi' \rangle = -\int_{\mathbb{R}} f(x) \varphi'(x) dx = -\sum_{0 \leq n \leq N} \int_{a_n}^{a_{n+1}} f(x) \varphi'(x) dx \\ &= \sum_{0 \leq n \leq N} \int_{a_n}^{a_{n+1}} \frac{df}{dx}(x) \varphi(x) dx + \sum_{0 \leq n \leq N} (f(a_n^+) \varphi(a_n) - f(a_{n+1}^-) \varphi(a_{n+1})) \\ &= \int \varphi(x) \left( \sum_{0 \leq n \leq N} \frac{df}{dx}(x) \mathbf{1}_{[a_n, a_{n+1}]}(x) \right) + \sum_{1 \leq n \leq N} f(a_n^+) \varphi(a_n) - \sum_{1 \leq n \leq N} f(a_n^-) \varphi(a_n), \end{aligned}$$

so that we have obtained the so-called *formula of jumps*

$$f' = \sum_{0 \leq n \leq N} \frac{df}{dx} \mathbf{1}_{[a_n, a_{n+1}]} + \sum_{1 \leq n \leq N} (f(a_n^+) - f(a_n^-)) \delta_{a_n}, \quad (3.2.5)$$

where  $\delta_{a_n}$  is the Dirac mass at  $a_n$ , defined by  $\langle \delta_{a_n}, \varphi \rangle = \varphi(a_n)$ .

We consider now the following determination of the logarithm given for  $z \in \mathbb{C} \setminus \mathbb{R}_-$  by

$$\text{Log } z = \oint_{[1, z]} \frac{d\xi}{\xi}, \quad (3.2.6)$$

which makes sense since  $\mathbb{C} \setminus \mathbb{R}_-$  is star-shaped with respect to 1, i.e. the segment  $[1, z] \subset \mathbb{C} \setminus \mathbb{R}_-$  for  $z \in \mathbb{C} \setminus \mathbb{R}_-$ . Since the function  $\text{Log}$  coincides with  $\ln$  on  $\mathbb{R}_+^*$  and is holomorphic on  $\mathbb{C} \setminus \mathbb{R}_-$ , we get by analytic continuation that

$$e^{\text{Log } z} = z, \quad \text{for } z \in \mathbb{C} \setminus \mathbb{R}_-. \quad (3.2.7)$$

Also by analytic continuation, we have for  $|\text{Im } z| < \pi$ ,  $\text{Log}(e^z) = z$ . We want now to study the distributions on  $\mathbb{R}$ ,

$$u_y(x) = \text{Log}(x + iy), \quad \text{where } y \neq 0 \text{ is a real parameter.}$$

We leave as an exercise for the reader to prove that

$$\lim_{y \rightarrow 0_{\pm}} \text{Log}(x + iy) = \ln |x| \pm i\pi(1 - H(x)), \quad (3.2.8)$$

where the limits are taken in the sense of the definition 3.1.16; also the reader can check

$$\frac{1}{x \pm i0} = \text{pv}\left(\frac{1}{x}\right) \mp i\pi\delta_0, \quad (3.2.9)$$

where we have defined

$$\left\langle \frac{1}{x \pm i0}, \varphi \right\rangle = \lim_{\epsilon \rightarrow 0_{+}} \int \frac{\varphi(x)}{x \pm i\epsilon} dx \quad (3.2.10)$$

(part of the exercise is to prove that these limits exist for  $\varphi \in \mathcal{D}(\mathbb{R})$ ). We conclude that section of examples with a more general lemma on a simple ODE.

**Lemma 3.2.4.** *Let  $I$  be an open interval of  $\mathbb{R}$ . The solutions in  $\mathcal{D}'(I)$  of  $u' = 0$  are the constants. The solutions in  $\mathcal{D}'(I)$  of  $u' = f$  make a one-dimensional affine subspace of  $\mathcal{D}'(I)$ .*

*Proof.* We assume first that  $f = 0$ ; if  $u$  is a constant, then it is of course a solution. Conversely, let us assume that  $u \in \mathcal{D}'(I)$  satisfies  $u' = 0$ . Let  $\chi_0 \in C_c^\infty(I)$  such that  $\int_{\mathbb{R}} \chi_0(x) dx = 1$ ; then we have for any  $\varphi \in C_c^\infty(I)$ , with  $J(\varphi) = \int_{\mathbb{R}} \varphi(x) dx$ ,  $\psi(x) = \int_{-\infty}^x (\varphi(t) - J(\varphi)\chi_0(t)) dt$ , noting that  $\psi$  belongs<sup>2</sup> to  $C_c^\infty(I)$ ,

$$\langle u, \varphi - J(\varphi)\chi_0 \rangle = \langle u, \psi' \rangle = -\langle u', \psi \rangle = 0,$$

<sup>2</sup>The function  $\psi$  is obviously smooth and if  $\varphi, \chi_0$  are both supported in  $\{a \leq x \leq b\}$ ,  $a, b \in I$ , so is  $\psi$ , thanks to the condition  $\int \chi_0 = 1$ .

which gives  $\langle u, \varphi \rangle = J(\varphi)\langle u, \chi_0 \rangle$ , i.e.  $u = \langle u, \chi_0 \rangle$  proving that  $u$  is indeed a constant. We have proven that the solutions  $u \in \mathcal{D}'(I)$  of  $u' = 0$  are simply the constants. If  $f \in \mathcal{D}'(I)$ , we need only to construct a solution  $v_0$  of  $v_0' = f$  and then use the previous result to obtain that the set of solutions of  $u' = f$  is  $v_0 + \mathbb{R}$ . Let us construct such a solution  $v_0$ . For  $\varphi \in \mathcal{D}(I)$ , we define with the same  $\psi$  as above,

$$\langle v_0, \varphi \rangle = -\langle f, \psi \rangle. \tag{3.2.11}$$

It is a distribution since for  $\text{supp } \varphi$  compact  $\subset I$ , we define (the compact set)  $K_1 = \text{supp } \varphi \cup \text{supp } \chi_0$ , and we have

$$|\langle v_0, \varphi \rangle| = |\langle f, \psi \rangle| \leq C_{K_1} \max_{0 \leq j \leq N_{K_1}} \|\psi^{(j)}\|_{L^\infty} \leq C \max_{0 \leq j \leq (N_{K_1}-1)_+} \|\varphi^{(j)}\|_{L^\infty}.$$

Moreover the formula (3.2.11) implies the sought result

$$\langle v_0', \varphi \rangle = -\langle v_0, \varphi' \rangle = \langle f, \psi_{\varphi'} \rangle = \langle f, \varphi \rangle,$$

since  $\psi_{\varphi'}(x) = \int_{-\infty}^x (\varphi'(t) - J(\varphi')\chi_0(t))dt = \varphi(x)$  because  $J(\varphi') = 0$ . The proof of the lemma is complete. □

### 3.2.3 Product by smooth functions

We define now the product of a  $C^\infty$  (resp.  $C^N$ ) function by a distribution (resp. of order  $N$ ).

**Definition 3.2.5.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$  and  $u \in \mathcal{D}'(\Omega)$ . For  $f \in C^\infty(\Omega)$ , we define the product  $f \cdot u$  as the distribution defined by*

$$\langle f \cdot u, \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \langle u, f\varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)}. \tag{3.2.12}$$

If  $u$  is of order  $N$  and  $f \in C^N(\Omega)$ , we define the product  $f \cdot u$  as the distribution of order  $N$  defined by

$$\langle f \cdot u, \varphi \rangle_{\mathcal{D}'^N(\Omega), C_c^N(\Omega)} = \langle u, f\varphi \rangle_{\mathcal{D}'^N(\Omega), C_c^N(\Omega)}. \tag{3.2.13}$$

**Remark 3.2.6.** Since the multiplication by a  $C^\infty(\Omega)$  (resp.  $C^N(\Omega)$ ) function is a continuous linear operator from  $C_c^\infty(\Omega)$  (resp.  $C_c^N(\Omega)$ ) into itself, we get that the above formulas actually define the products as distributions on  $\Omega$  with the right order (see the proposition 3.1.12). Also the product defined in the second part coincides with the first definition whenever  $f \in C_c^\infty(\Omega)$  and if  $u \in L_{\text{loc}}^1(\Omega)$ ,  $f \in C^0(\Omega)$ , the usual product  $fu$  coincides with the  $f \cdot u$  defined here, thanks to the lemma 3.1.7.

The next theorem is providing an extension to the classical Leibniz' formula for the derivatives of a product.

**Theorem 3.2.7.** *Let  $\Omega$  be an open set of  $\mathbb{R}^n$ ,  $u \in \mathcal{D}'(\Omega)$ ,  $f \in C^\infty(\Omega)$  and  $\alpha \in \mathbb{N}^n$  be a multi-index (see (2.3.6)). Then we have*

$$\frac{\partial_x^\alpha (fu)}{\alpha!} = \sum_{\substack{\beta, \gamma \in \mathbb{N}^n \\ \beta + \gamma = \alpha}} \frac{\partial_x^\beta (f)}{\beta!} \frac{\partial_x^\gamma (u)}{\gamma!}. \tag{3.2.14}$$

*Proof.* We get immediately by induction on  $|\alpha|$  the formula

$$\frac{\partial_x^\alpha(fu)}{\alpha!} = \sum_{\substack{\beta, \gamma \in \mathbb{N}^n \\ \beta + \gamma = \alpha}} \sigma_{\beta, \gamma} \frac{\partial_x^\beta(f)}{\beta!} \frac{\partial_x^\gamma(u)}{\gamma!}, \quad \text{with } \sigma_{\beta, \gamma} \in \mathbb{R}_+.$$

To find the  $\sigma_{\beta, \gamma}$ , we choose  $f(x) = e^{x \cdot \xi}$ ,  $u(x) = e^{x \cdot \eta}$ , with  $\xi, \eta \in \mathbb{R}^n$ . We find then for all  $\xi, \eta \in \mathbb{R}^n$ , the identity

$$\frac{(\xi + \eta)^\alpha}{\alpha!} = \frac{\partial_x^\alpha(e^{x \cdot (\xi + \eta)})}{\alpha!} \Big|_{x=0} = \sum_{\substack{\beta, \gamma \in \mathbb{N}^n \\ \beta + \gamma = \alpha}} \sigma_{\beta, \gamma} \frac{\partial_x^\beta(e^{x \cdot \xi})}{\beta!} \frac{\partial_x^\gamma(e^{x \cdot \eta})}{\gamma!} \Big|_{x=0} = \sum_{\substack{\beta, \gamma \in \mathbb{N}^n \\ \beta + \gamma = \alpha}} \sigma_{\beta, \gamma} \frac{\xi^\beta \eta^\gamma}{\beta! \gamma!},$$

and the formula (2.3.7) shows that for  $\beta, \gamma$  such that  $\beta + \gamma = \alpha$

$$\sigma_{\beta, \gamma} = \partial_\xi^\beta \partial_\eta^\gamma \left( \frac{(\xi + \eta)^\alpha}{\alpha!} \right) \Big|_{\xi = \eta = 0} = 1,$$

completing the proof of the theorem.  $\square$

**Examples.** Let  $f$  be a continuous function on  $\mathbb{R}$  and  $\delta_0$  be the Dirac mass at 0. The product  $f \cdot \delta_0$  is equal to  $f(0)\delta_0$ : since  $\delta_0$  is a distribution of order 0, we can multiply it by a continuous function and if  $\varphi \in C_c^0(\mathbb{R})$ , we have

$$\langle f \cdot \delta_0, \varphi \rangle = \langle \delta_0, f\varphi \rangle = f(0)\varphi(0) = \langle f(0)\delta_0, \varphi \rangle \implies f \cdot \delta_0 = f(0)\delta_0. \quad (3.2.15)$$

On the other hand if  $f \in C^1(\mathbb{R})$  we have

$$f \cdot \delta'_0 = f(0)\delta'_0 - f'(0)\delta_0, \quad (3.2.16)$$

since the Leibniz' formula (3.2.14) gives  $f(0)\delta'_0 = (f \cdot \delta_0)' = f' \cdot \delta_0 + f \cdot \delta'_0 = f'(0)\delta_0 + f \cdot \delta'_0$ . In particular  $x\delta'_0 = -\delta_0$ .

### 3.2.4 Division of distribution on $\mathbb{R}$ by $x^m$

We want now to address the question of division of a function (or a distribution) by a polynomial; a typical example is the division of 1 by the linear function  $x$  expressed by the identity

$$x \operatorname{pv}(1/x) = 1 \quad (3.2.17)$$

which is an immediate consequence of (3.1.13). We note also from the previous examples that, for any constant  $c$ , we have  $x(\operatorname{pv}(1/x) + c\delta_0) = 1$ . The next theorem shows that  $T = \operatorname{pv}(1/x) + c\delta_0$  are the only distributions solutions of the equation  $xT = 1$ .

**Theorem 3.2.8.** *Let  $m \geq 1$  be an integer.*

- (1) *If  $u \in \mathcal{D}'(\mathbb{R})$  is such that  $x^m u = 0$ , then  $u = \sum_{0 \leq j < m} c_j \delta_0^{(j)}$ .*
- (2) *Let  $v \in \mathcal{D}'(\mathbb{R})$ ; there exists  $u \in \mathcal{D}'(\mathbb{R})$  such that  $v = x^m u$ .*

*Proof.* Let us first prove (1). For  $\varphi, \chi_0 \in C_c^\infty(\mathbb{R})$  with  $\chi_0 = 1$  near 0, we have

$$\varphi(x) = \underbrace{\sum_{0 \leq j < m} \frac{\varphi^{(j)}(0)}{j!} x^j}_{p_{\varphi, m}(x)} + \underbrace{\int_0^1 \frac{(1-t)^{m-1}}{(m-1)!} \varphi^{(m)}(tx) dt x^m}_{\psi_{m, \varphi}(x)}, \quad \psi_{m, \varphi} \in C^\infty(\mathbb{R}),$$

and thus, since  $x^m u = 0$ ,

$$\begin{aligned} \langle u, \varphi \rangle &= \overbrace{\langle x^m u, x^{-m}(1 - \chi_0)\varphi \rangle}^{=0} + \langle u, \chi_0 \varphi \rangle = \langle u, \chi_0 p_{m, \varphi} \rangle + \overbrace{\langle x^m u, \chi_0 \psi_{m, \varphi} \rangle}^{=0} \\ &= \sum_{0 \leq j < m} \frac{\varphi^{(j)}(0)}{j!} \langle u, \chi_0 \rangle = \sum_{0 \leq j < m} \langle c_j \delta_0^{(j)}, \varphi \rangle, \end{aligned}$$

which is the sought result. To obtain (2), for  $\varphi \in C_c^\infty(\mathbb{R})$ , and a given  $v_0 \in \mathcal{D}'(\mathbb{R})$ , we define, using the above notations,

$$\langle u, \varphi \rangle = \langle v_0, \chi_0 \psi_{m, \varphi} \rangle + \langle v_0, x^{-m}(1 - \chi_0)\varphi \rangle.$$

This defines obviously a distribution on  $\mathbb{R}$  and  $\langle x^m u, \varphi \rangle = \langle u, x^m \varphi \rangle$ ; for the function  $\phi(x) = x^m \varphi(x)$ , we have  $p_{\phi, m} = 0$ ,  $x^m \psi_{m, \phi}(x) = x^m \varphi(x)$ , so that the smooth functions  $\psi_{m, \phi} = \varphi$ ,

$$\langle x^m u, \varphi \rangle = \langle v_0, \chi_0 \varphi \rangle + \langle v_0, x^{-m}(1 - \chi_0)x^m \varphi \rangle = \langle v_0, \varphi \rangle. \quad \square$$

### 3.3 Distributions with compact support

#### 3.3.1 Identification with $\mathcal{E}'$

Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ . We have already seen that the space  $C^\infty(\Omega)$  (also denoted by  $\mathcal{E}(\Omega)$ ) is a Fréchet space. Denoting by  $\mathcal{E}'(\Omega)$  the topological dual of  $\mathcal{E}(\Omega)$ , we can consider  $T \in \mathcal{E}'(\Omega)$  as a distribution  $\tilde{T}$  on  $\Omega$  by defining

$$\langle \tilde{T}, \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \langle T, \varphi \rangle_{\mathcal{E}'(\Omega), \mathcal{E}(\Omega)} \quad (\text{this makes sense since } \mathcal{D}(\Omega) \subset \mathcal{E}(\Omega)).$$

The linearity is obvious and the continuity of  $T$  as a linear form on the Fréchet space  $\mathcal{E}(\Omega)$  implies that there exists  $C > 0, N \in \mathbb{N}, K$  compact subset of  $\Omega$  such that

$$\forall \varphi \in \mathcal{E}(\Omega), \quad |\langle T, \varphi \rangle_{\mathcal{E}'(\Omega), \mathcal{E}(\Omega)}| \leq C \sup_{|\alpha| \leq N, x \in K} |(\partial_x^\alpha \varphi)(x)|.$$

This estimate also proves that  $\tilde{T}$  belongs to  $\mathcal{D}'(\Omega)$ ; moreover, it has compact support in the sense of the definition (3.1.8): we have  $\langle \tilde{T}, \varphi \rangle = 0$  for  $\varphi \in C_c^\infty(\Omega)$ ,  $\text{supp } \varphi \subset K^c$ , so that  $\tilde{T}|_{K^c} = 0$  and thus  $\text{supp } \tilde{T} \subset K$ . The next theorem proves that we can identify the space  $\mathcal{E}'(\Omega)$  with the distributions on  $\Omega$  with compact support, denoted by  $\mathcal{D}'_{\text{comp}}(\Omega)$ .

**Theorem 3.3.1.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ . The mapping  $\iota : \mathcal{E}'(\Omega) \rightarrow \mathcal{D}'_{\text{comp}}(\Omega)$ , defined as above by  $\iota(T) = \tilde{T}$  is bijective.*

*Proof.* The mapping  $\iota$  is linear and if  $\iota(T) = 0$ , we know that  $T$  vanishes on all functions of  $\mathcal{D}(\Omega)$ .

**Lemma 3.3.2.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ . The space  $\mathcal{D}(\Omega)$  is dense in  $\mathcal{E}(\Omega)$ .*

*Proof of the lemma.* We consider a sequence  $(K_j)_{j \geq 1}$  of compact subsets of  $\Omega$  such that the lemma 2.3.1 is satisfied. For each  $j \geq 1$ , we may use the lemma 3.1.3 to construct a function  $\chi_j \in \mathcal{D}(\Omega)$  with  $\chi_j = 1$  near  $K_j$ . For a given  $\varphi \in \mathcal{E}(\Omega)$ , the sequence  $(\varphi\chi_j)_{j \geq 1}$  of functions in  $\mathcal{D}(\Omega)$  converges in  $\mathcal{E}(\Omega)$  to  $\varphi$ , thanks to the last property of the lemma 2.3.1, proving the lemma.  $\square$

Since  $T$  is continuous on  $\mathcal{E}(\Omega)$ ,  $\langle T, \varphi \rangle_{\mathcal{E}'(\Omega), \mathcal{E}(\Omega)} = \lim_j \langle T, \varphi\chi_j \rangle_{\mathcal{E}'(\Omega), \mathcal{E}(\Omega)} = 0$  since  $T$  vanishes on  $\mathcal{D}(\Omega)$ . Let us consider now  $T \in \mathcal{D}'_{\text{comp}}(\Omega)$  with  $\text{supp } T = L$  (compact subset of  $\Omega$ ). Using the lemma 3.1.3, we consider  $\chi_0 \in \mathcal{D}(\Omega)$  such that  $\chi_0 = 1$  on a neighborhood of  $L$ . For  $\varphi \in \mathcal{E}(\Omega)$ , we define  $S \in \mathcal{E}'(\Omega)$  by

$$\langle S, \varphi \rangle_{\mathcal{E}'(\Omega), \mathcal{E}(\Omega)} = \langle T, \chi_0 \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} \quad (\text{note that } |\langle S, \varphi \rangle| \leq C \sup_{|\alpha| \leq N, x \in \text{supp } \chi_0} |\partial_x^\alpha \varphi|),$$

We have  $\iota(S) = T$  because

$$\langle \iota(S), \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \langle S, \varphi \rangle_{\mathcal{E}'(\Omega), \mathcal{E}(\Omega)} = \langle T, \chi_0 \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)} = \langle \chi_0 T, \varphi \rangle_{\mathcal{D}'(\Omega), \mathcal{D}(\Omega)},$$

and since for  $\varphi \in \mathcal{D}(\Omega)$ , the function  $(1 - \chi_0)\varphi$  vanishes on an open neighborhood  $V$  of  $L$  implying

$$\text{supp}((1 - \chi_0)\varphi) \subset V^c \subset L^c \implies \langle T, (1 - \chi_0)\varphi \rangle = 0,$$

so that  $\iota(S) = \chi_0 T = \chi_0 T + \underbrace{(1 - \chi_0)T}_{=0} = T$ . The proof of the theorem is complete.  $\square$

**Remark 3.3.3.** We can then identify  $\mathcal{D}'_{\text{comp}}(\Omega)$  with  $\mathcal{E}'(\Omega)$ , and we may note that for  $T \in \mathcal{D}'_{\text{comp}}(\Omega)$  with  $\text{supp } T = L$ ,  $T$  is of finite order  $N$ , and for all neighborhoods  $K$  of  $L$ , there exists  $C > 0$  such that, for all  $\varphi \in \mathcal{E}(\Omega)$ ,

$$|\langle T, \varphi \rangle| \leq C \sup_{|\alpha| \leq N, x \in K} |(\partial_x^\alpha \varphi)(x)|. \quad (3.3.1)$$

In general, it is not possible to take  $K = L$  in the above estimate.

### 3.3.2 Distributions with support at a point

The next theorem characterizes the distributions supported in  $\{0\}$ .

**Theorem 3.3.4.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $x_0 \in \Omega$  and let  $u \in \mathcal{D}'(\Omega)$  such that  $\text{supp } u = \{x_0\}$ . Then  $u = \sum_{|\alpha| \leq N} c_\alpha \delta_{x_0}^{(\alpha)}$ , where the  $c_\alpha$  are some constants.*

*Proof.* Let  $\varphi \in C^\infty(\Omega)$ ; we have for  $x \in V_0 \subset$  open neighborhood of  $x_0$  (included in  $\Omega$ ),  $N_0$  the order of  $u$ ,

$$\varphi(x) = \sum_{|\alpha| \leq N_0} \frac{(\partial_x^\alpha \varphi)(x_0)}{\alpha!} (x-x_0)^\alpha + \underbrace{\int_0^1 \frac{(1-\theta)^{N_0}}{N_0!} \varphi^{(N_0+1)}(x_0 + \theta(x-x_0)) d\theta}_{\psi(x), \psi \in C^\infty(V_0)} (x-x_0)^{N_0+1},$$

and thus for  $\chi_0 \in C_c^\infty(V_0)$ ,  $\chi_0 = 1$  near  $x_0$ ,

$$\langle u, \varphi \rangle = \langle u, \chi_0 \varphi \rangle = \sum_{|\alpha| \leq N_0} \frac{(\partial_x^\alpha \varphi)(x_0)}{\alpha!} \langle u, \chi_0(x) (x-x_0)^\alpha \rangle + \langle u, \chi_0(x) \psi(x) (x-x_0)^{N_0+1} \rangle. \quad (3.3.2)$$

We have also

$$|\langle u, \chi_0(x) \psi(x) (x-x_0)^{N_0+1} \rangle| \leq C_0 \sup_{|\alpha| \leq N_0} |\partial_x^\alpha (\chi_0(x) \psi(x) (x-x_0)^{N_0+1})|. \quad (3.3.3)$$

We can take  $\chi_0(x) = \rho(\frac{x-x_0}{\epsilon})$ , where  $\rho \in C_c^\infty(\mathbb{R}^n)$  is supported in the unit ball  $B_1$ ,  $\rho = 1$  in  $\frac{1}{2}B_1$  and  $\epsilon > 0$ . We have then

$$\begin{aligned} \chi_0(x) \psi(x) (x-x_0)^{N_0+1} &= \epsilon^{N_0+1} \rho\left(\frac{x-x_0}{\epsilon}\right) \psi\left(x_0 + \epsilon \frac{x-x_0}{\epsilon}\right) \frac{(x-x_0)^{N_0+1}}{\epsilon^{N_0+1}} \\ &= \epsilon^{N_0+1} \rho_1\left(\frac{x-x_0}{\epsilon}\right) \end{aligned}$$

with  $\rho_1(t) = \rho(t) \psi(x_0 + \epsilon t) t^{N_0+1}$ , so that  $\rho_1 \in C_c^\infty(\mathbb{R}^n)$  is supported in the unit ball  $B_1$  has all its derivatives bounded independently of  $\epsilon$ . From (3.3.3), we get for all  $\epsilon > 0$ ,

$$|\langle u, \chi_0(x) \psi(x) (x-x_0)^{N_0+1} \rangle| \leq C_0 \sup_{|\alpha| \leq N_0} \epsilon^{N_0+1-|\alpha|} |(\partial_t^\alpha \rho_1)\left(\frac{x-x_0}{\epsilon}\right)| \leq C_1 \epsilon,$$

which implies that the left-hand-side of (3.3.3) is zero. On the other hand, for  $\chi_1 \in C_c^\infty(V_0)$ ,  $\chi_1 = 1$  near the support of  $\chi_0$ , we have

$$\begin{aligned} \langle u, \chi_1(x) (x-x_0)^\alpha \rangle &= \langle u, \underbrace{\chi_1(x) \chi_0(x)}_{=\chi_0(x)} (x-x_0)^\alpha \rangle + \langle u, \underbrace{\chi_1(x) (1-\chi_0(x))}_{\text{supported in } (\text{supp } u)^c} (x-x_0)^\alpha \rangle \\ &= \langle u, \chi_0(x) (x-x_0)^\alpha \rangle \end{aligned}$$

so that the latter does not depend on  $\epsilon$  for  $\epsilon$  small enough. The result of the theorem follows from (3.3.2).  $\square$

### 3.4 Tensor products

Let  $X$  be an open subset of  $\mathbb{R}^m$ ,  $Y$  be an open subset of  $\mathbb{R}^n$  and  $f \in C_c^\infty(X)$ ,  $g \in C_c^\infty(Y)$ . The tensor product  $f \otimes g$  is defined by  $(f \otimes g)(x, y) = f(x)g(y)$  and belongs