

THE AFFINE GRASSMANNIAN AS A PRESHEAF QUOTIENT

KĘSTUTIS ČESNAVIČIUS

ABSTRACT. For a reductive group G over a ring A , its affine Grassmannian Gr_G plays important roles in a wide range of subjects and is typically defined as the étale sheafification of the presheaf quotient LG/L^+G of the loop group LG by its positive loop subgroup L^+G . We show that the Zariski sheafification gives the same result. Moreover, for totally isotropic G (for instance, for quasi-split G), we show that no sheafification is needed at all: Gr_G is already the presheaf quotient LG/L^+G , which seems new already in the classical case of G over $\mathbb{C}$. For totally isotropic G , we also show that the affine Grassmannian may be formed using polynomial loops. We deduce all of these results from the study of G -torsors on $\mathbb{P}_A^1$ that is ultimately built on the geometry of Bun_G .

1. The affine Grassmannian and its modular description	2
2. The affine Grassmannian as a Zariski quotient	6
3. The affine Grassmannian as a presheaf quotient	8
Acknowledgements	10
References	10

The affine Grassmannian Gr_G of a reductive group G originated in Lusztig’s [Lus83, Section 11] (see also Beilinson–Drinfeld’s [BD, Section 4.5]) and is instrumental in the geometric Langlands program and other subjects that study G -torsors and their moduli. The goal of this article is to show that Gr_G admits a simpler definition than previously thought: for most G , it is simply the presheaf quotient LG/L^+G of the loop group LG by its positive loop subgroup L^+G , so that the fpqc or étale sheafifications of this quotient that were used previously are not needed, see Theorems 12 and 17 for precise statements.

1. Conventions. As in [SGA 3_{III new}, exposé XIX, définition 2.7], a *reductive group scheme* over a scheme S is a smooth, affine S -group scheme whose geometric S -fibers are connected reductive groups. An S -scheme is *locally constant* if fpqc locally on S it becomes isomorphic to some $\bigsqcup_{i \in I} S$.

CNRS, UNIVERSITÉ PARIS-SACLAY, LABORATOIRE DE MATHÉMATIQUES D’ORSAY, F-91405, ORSAY, FRANCE
E-mail address: `kestutis@math.u-psud.fr`.

Date: May 18, 2025.

2020 Mathematics Subject Classification. Primary 14L15; Secondary 14M17.

Key words and phrases. Affine Grassmannian, loop group, reductive group, torsor.

1. THE AFFINE GRASSMANNIAN AND ITS MODULAR DESCRIPTION

In this section, we fix a base ring A and aim to review the definition and the modular interpretation of the affine Grassmannian Gr_G for a smooth, quasi-affine¹ $A[[t]]$ -group scheme $\mathcal{G}$, see Proposition 7.

2. The loop functor. For a functor $\mathcal{X}$ on the category of $A((t))$ -algebras (resp., $A[[t]]$ -algebras), its *loop functor* $L\mathcal{X}$ (resp., *positive loop functor* $L^+\mathcal{X}$) is defined on the category of A -algebras by

$$L\mathcal{X}: B \mapsto \mathcal{X}(B((t))) \quad (\text{resp., } L^+\mathcal{X}: B \mapsto \mathcal{X}(B[[t]])).$$

The morphism $L^+\mathcal{X} \rightarrow L\mathcal{X}$ is often an inclusion, for instance, this is so whenever $\mathcal{X}$ is a subfunctor of a separated $A[[t]]$ -scheme (see [EGA I, corollaire 9.5.6]).

If $\mathcal{X}$ is an $A[[t]]$ -scheme, then, by [Bha16, Theorem 4.1 and Remark 4.6], $L^+\mathcal{X}$ is an A -scheme because

$$L^+\mathcal{X} \cong \varprojlim_{n>0} \mathrm{Res}_{(A[t]/(t^n))/A}(\mathcal{X}_{A[t]/(t^n)}), \quad \text{equivalently, } (L^+\mathcal{X})(B) = \varprojlim_{n>0} \mathcal{X}(B[t]/(t^n))$$

for every A -algebra B , compare with [CLNS18, Chapter 3, Corollary 3.3.7 b)]. Similarly, if $\mathcal{X}$ is a quasi-compact and quasi-separated $A[[t]]$ -algebraic space, then, by [Bha16, Theorem 4.1] and [SP, Proposition 05YF and Lemmas 07SF and 05YD], $L^+\mathcal{X}$ is a quasi-compact and quasi-separated A -algebraic space. If $\mathcal{X}$ is even an affine $A[[t]]$ -scheme, then, by considering a presentation of its coordinate ring in terms of generators and relations, $L^+\mathcal{X}$ is an affine A -scheme and $L\mathcal{X}$ is an ind-affine A -ind-scheme, more precisely, there are affine A -schemes X_n and closed immersions

$$X_0 \hookrightarrow X_1 \hookrightarrow \cdots \quad \text{such that} \quad L^+\mathcal{X} = X_0 \quad \text{and} \quad L\mathcal{X} = \bigcup_{n \geq 0} X_n \quad \text{as functors on } A\text{-algebras } B.$$

For an A -algebra B , let $B\{t\}$ be the Henselization of $B[t]$ with respect to the ideal $tB[t]$, see [BC22, Section 2.1.2] or [SP, Lemma 0A02]. It is useful to consider Henselian (resp., algebraic; resp., polynomial) variants $L_h\mathcal{X}$ and $L_h^+\mathcal{X}$ (resp., $L_{\mathrm{alg}}\mathcal{X}$ and $L_{\mathrm{alg}}^+\mathcal{X}$; resp., $L_{\mathrm{poly}}\mathcal{X}$ and $L_{\mathrm{poly}}^+\mathcal{X}$):

$$\begin{aligned} L_h\mathcal{X}: B &\mapsto \mathcal{X}(B\{t\}[\tfrac{1}{t}]) & \text{and} & & L_h^+\mathcal{X}: B &\mapsto \mathcal{X}(B\{t\}), \\ L_{\mathrm{alg}}\mathcal{X}: B &\mapsto \mathcal{X}((B[t]_{1+tB[t]})[\tfrac{1}{t}]) & \text{and} & & L_{\mathrm{alg}}^+\mathcal{X}: B &\mapsto \mathcal{X}(B[t]_{1+tB[t]}), \\ L_{\mathrm{poly}}\mathcal{X}: B &\mapsto \mathcal{X}(B[t, t^{-1}]) & \text{and} & & L_{\mathrm{poly}}^+\mathcal{X}: B &\mapsto \mathcal{X}(B[t]), \end{aligned}$$

granted that $\mathcal{X}$ begins its life over $A\{t\}[\frac{1}{t}]$ (resp., over $(A[t]_{1+tA[t]})[\frac{1}{t}]$; resp., over $A[t, t^{-1}]$) and, for $L_{\mathrm{poly}}^+\mathcal{X}$, even already over $A\{t\}$ (resp., over $A[t]_{1+tA[t]}$; resp., over $A[t]$). These variant functors are sometimes easier to handle, for instance, they all commute with filtered direct limits in B granted that so does $\mathcal{X}$, moreover, $L_{\mathrm{poly}}^+\mathcal{X}$ and $L_{\mathrm{poly}}\mathcal{X}$ are nothing else but restrictions of scalars.

By the following proposition, for many $\mathcal{X}$, the functors $L_*^{(+)}\mathcal{X}$ cannot be improved by sheafifying.

Proposition 3. *For a scheme $\mathcal{X}$ over $A((t))$, or over $A\{t\}[\frac{1}{t}]$, or over $(A[t]_{1+tA[t]})[\frac{1}{t}]$, or over $A[t, t^{-1}]$ (resp., over $A[[t]]$, or over $A\{t\}$, or over $A[t]_{1+tA[t]}$, or over $A[t]$) as in §2 such that every quasi-compact open of $\mathcal{X}$ is quasi-affine, the functor $L_*\mathcal{X}$ (resp., and also its subfunctor $L_*^+\mathcal{X}$) is an fpqc sheaf on the category of A -algebras.*

Proof. Since $\mathcal{X}$ is separated, [EGA I, corollaire 9.5.6] ensures that the map $L_*^+\mathcal{X} \rightarrow L_*\mathcal{X}$ is indeed an inclusion. Thus, all we need to show is that for an fpqc cover $B \rightarrow B'$ of A -algebras, the sequence $(L_*\mathcal{X})(B) \rightarrow (L_*\mathcal{X})(B') \rightrightarrows (L_*\mathcal{X})(B' \otimes_B B')$ (resp., $(L_*^+\mathcal{X})(B) \rightarrow (L_*^+\mathcal{X})(B') \rightrightarrows (L_*^+\mathcal{X})(B' \otimes_B B')$)

¹Of course, over a field every quasi-affine group scheme of finite type is affine, see [SGA 3I_{new}, exposé VI_B, proposition 11.11], and likewise for flat, finite type, separated groups over Dedekind rings with affine (or merely quasi-affine) generic fibers, see [SGA 3II, exposé XVII, proposition C.2.1 (3)] and [Ana73, proposition 2.3.1]. Over higher-dimensional base rings, however, there exist quasi-affine groups that are not affine, see [Ray70, chapitre VII, section 3] for such an example over $\mathbb{C}[x, y]$.

is exact. The case of the polynomial variant follows from fpqc descent [SP, Lemma 023Q] because

$$B[t] \rightarrow B'[t] \rightrightarrows (B' \otimes_B B')[t] \quad \text{and} \quad B[t, t^{-1}] \rightarrow B'[t, t^{-1}] \rightrightarrows (B' \otimes_B B')[t, t^{-1}]$$

are both fpqc cover sequences. For other variants, since each ring-valued point of $\mathcal{X}$ factors through a quasi-compact open, we may assume that $\mathcal{X}$ is quasi-affine. The maps

$$B\{t\} \rightarrow B'\{t\} \quad \text{and} \quad B[t]_{1+tB[t]} \rightarrow B'[t]_{1+tB'[t]}$$

are both faithfully flat, in particular, they induce surjections on spectra, so the Henselian and algebraic variants reduce further to affine $\mathcal{X}$. In the power series case, the reduction to affine $\mathcal{X}$ is more subtle if B and B' are not Noetherian (because then we do not know whether the map $\text{Spec } B'((t)) \rightarrow \text{Spec } B((t))$ is surjective) and will use ideas from [BČ22, Lemma 2.2.9 (i)] as follows.

By realizing $\mathcal{X}$ as the complement of the vanishing locus of a finitely generated ideal in an affine $A((t))$ -scheme (resp., $A[[t]]$ -scheme), to reduce to affine $\mathcal{X}$ we need to show that elements $b_1, \dots, b_n$ in $B((t))$ (resp., in $B[[t]]$) generate the unit ideal as soon as their images do so in $B'((t))$ (resp., in $B'[[t]]$). The case of $B[[t]]$ follows by checking modulo t and using the faithful flatness of $B \rightarrow B'$. To treat $B((t))$, we may assume that $b_1, \dots, b_n \in B[[t]]$ and use the faithfully flat cover $B[[t]] \rightarrow (B[[t]] \otimes_B B')_{(t)}^h$ (Henselization with respect to the ideal generated by t) to reduce to showing that $b_1, \dots, b_n$ generate the unit ideal in $(B[[t]] \otimes_B B')_{(t)}^h[\frac{1}{t}]$. The t -adic completion of $(B[[t]] \otimes_B B')_{(t)}^h$ is $B'[[t]]$ and, by assumption, there are $a_1, \dots, a_n \in B'[[t]]$ with $a_1 b_1 + \dots + a_n b_n = t^N$ in $B'[[t]]$ for some $N > 0$. By approximating modulo t^{N+1} , therefore, there are $a'_1, \dots, a'_n \in (B[[t]] \otimes_B B')_{(t)}^h$ such that

$$a'_1 b_1 + \dots + a'_n b_n \in t^N + t^{N+1} (B[[t]] \otimes_B B')_{(t)}^h.$$

This means that $b_1, \dots, b_n$ indeed generate the unit ideal in $(B[[t]] \otimes_B B')_{(t)}^h[\frac{1}{t}]$, as desired.

Now that $\mathcal{X}$ is affine, the functor $\mathcal{X}(-)$ turns fiber products of rings into fiber products of sets. Thus, all we need to show is the exactness of the horizontal sequences in the commutative diagram

$$\begin{array}{ccccc} B[t]_{1+tB[t]} & \longrightarrow & B'[t]_{1+tB'[t]} & \rightrightarrows & (B' \otimes_B B')[t]_{1+t(B' \otimes_B B')[t]} \\ \downarrow & & \downarrow & & \downarrow \\ B\{t\} & \longrightarrow & B'\{t\} & \rightrightarrows & (B' \otimes_B B')\{t\} \\ \downarrow & & \downarrow & & \downarrow \\ B[[t]] & \longrightarrow & B'[[t]] & \rightrightarrows & (B' \otimes_B B')[[t]], \end{array}$$

which implies the corresponding exactness after further inverting t ; here the bottom vertical maps are injective by [BČ22, Section 2.1.2] (a limit argument to reduce to finite type $\mathbb{Z}$ -algebras). The exactness of the bottom row is seen coefficientwise. Thus, by fpqc descent, it is enough to show that

$$\begin{aligned} B'\{t\} \otimes_{B\{t\}} B'\{t\} &\hookrightarrow (B' \otimes_B B')\{t\}, \\ B'[t]_{1+tB'[t]} \otimes_{B[t]_{1+tB[t]}} B'[t]_{1+tB'[t]} &\hookrightarrow (B' \otimes_B B')[t]_{1+t(B' \otimes_B B')[t]}. \end{aligned}$$

The injectivity of the second map is evident because $B'[t] \otimes_{B[t]} B'[t] \xrightarrow{\sim} (B' \otimes_B B')[t]$ and the elements of $1 + t(B' \otimes_B B')[t]$ are nonzerodivisors. As for the first map, if the B -algebra B' is finitely presented, then we could use a limit argument to reduce to a Noetherian situation and then check the injectivity after passing to completions. In general, since every étale $(B' \otimes_B B')[t]$ -algebra $(B' \otimes_B B')$ -fiberwise has no embedded associated primes, both of the maps in question are injections by [RG71, première partie, corollaire 3.2.6] and a limit argument. $\square$

Remark 4. In Proposition 3, the assumption on quasi-compact opens holds if $\mathcal{X}$ is a subscheme of an affine scheme. In general, a scheme whose quasi-compact opens are all quasi-affine is *ind-quasi-affine*, see [BČ22, Definition 2.2.5] and the references to [SP] given there for details. We avoid this

terminology here because it may be confusing in the context of ind-schemes. The automorphism scheme of a reductive group is such an $\mathcal{X}$ that need not be quasi-affine, see [Čes22b, Section 1.3.7]. Another useful example is locally constant schemes $\mathcal{X}$, for which even $L\mathcal{X} = L^+\mathcal{X}$ as follows.

Corollary 5. *Let A be a ring.*

(a) *In the following diagram, the squares are Cartesian and the maps are bijective on idempotents:*

$$\begin{array}{ccccccccc} A & \hookrightarrow & A[t] & \hookrightarrow & A[t]_{1+tA[t]} & \hookrightarrow & A\{t\} & \hookrightarrow & A[[t]] \\ & \searrow & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ & & A[t, t^{-1}] & \hookrightarrow & (A[t]_{1+tA[t]})[\frac{1}{t}] & \hookrightarrow & A\{t\}[\frac{1}{t}] & \hookrightarrow & A((t)). \end{array}$$

(b) *For every scheme $\mathcal{X}$ over $A[[t]]$, or over $A\{t\}$, or over $A[t]_{1+tA[t]}$, or over $A[t]$ as in §2 such that $\mathcal{X}$ is locally constant (see §1), we have $L_*^+\mathcal{X} \xrightarrow{\sim} L_*\mathcal{X}$.*

Proof.

- (a) As indicated, the maps are all injective, either by inspection, or by faithful flatness, or by a limit argument given in [BČ22, Section 2.1.2]. The squares are Cartesian by [SP, Lemma 0BNR]. As for idempotents, since the maps are injective, it suffices to recall from [BČ22, Corollary 2.1.19] that the map $A \rightarrow A((t))$ is bijective on idempotents (as one may also show directly).
- (b) By [SP, Lemma 0AP8], we may check fpqc locally on the base that our locally constant $\mathcal{X}$ satisfies the assumptions of Proposition 3, more precisely, that every quasi-compact open of $\mathcal{X}$ is quasi-affine. In particular, in the case when $\mathcal{X}$ begins life over $A\{t\}$, the square

$$\begin{array}{ccc} \mathcal{X}(A\{t\}) & \hookrightarrow & \mathcal{X}(A\{t\}[\frac{1}{t}]) \\ \downarrow & & \downarrow \\ \mathcal{X}(A[[t]]) & \hookrightarrow & \mathcal{X}(A((t))) \end{array}$$

is Cartesian by [BČ22, Proposition 2.2.12], and likewise with $A[t]_{1+tA[t]}$ or $A[t]$ in place of $A\{t\}$. Since we may vary A , this reduces us to showing that the inclusion $\mathcal{X}(A[[t]]) \subset \mathcal{X}(A((t)))$ is an equality. For this, since *loc. cit.* also implies that for every prime $\mathfrak{p} \subset A$ the square

$$\begin{array}{ccc} \mathcal{X}(A[[t]] \otimes_A A_{\mathfrak{p}}) & \hookrightarrow & \mathcal{X}(A((t)) \otimes_A A_{\mathfrak{p}}) \\ \downarrow & & \downarrow \\ \mathcal{X}(A_{\mathfrak{p}}[[t]]) & \hookrightarrow & \mathcal{X}(A_{\mathfrak{p}}((t))) \end{array}$$

is Cartesian, by combining this square with spreading out, we may assume further that A is local. At this point, Proposition 3 allows us replace A by its strict Henselization.

Once A is strictly Henselian local, we fix any $A((t))$ -point of $\mathcal{X}$ and form its schematic image to get a closed subscheme $Z \subset \mathcal{X}$, see [SP, Definition 01R7]. It is enough to show that

$$Z \stackrel{?}{\cong} \mathrm{Spec}(A[[t]]).$$

We first claim that the map $Z \rightarrow \mathrm{Spec}(A[[t]])$ is surjective. For this, since $\mathcal{X}$ is locally constant, we may choose a faithfully flat $A[[t]]$ -algebra B such that $\mathcal{X}_B \cong \bigsqcup_{i \in I} \mathrm{Spec} B$. By [SP, Lemma 081I], the formation of Z commutes with base change to B , so we need to show that the schematic image of any $B[\frac{1}{t}]$ -point of $\mathcal{X}_B$ surjects onto $\mathrm{Spec} B$. It does because

otherwise, by the explicit nature of $\mathcal{X}_B$, the open immersion $\mathrm{Spec}(B[\frac{1}{t}]) \subset \mathrm{Spec} B$ would factor through some proper closed subscheme, contradicting its schematic density.

Since $A[[t]]$ is strictly Henselian local and $\mathcal{X}$ is étale, [EGA IV₄, théorème 18.5.11] ensures that for every $z \in Z$ above the maximal ideal of $A[[t]]$, we have

$$\mathcal{X} \cong \mathrm{Spec}(\mathcal{O}_{\mathcal{X},z}) \sqcup \mathcal{X}' \quad \text{with} \quad \mathcal{O}_{\mathcal{X},z} \cong A[[t]].$$

We have already argued that such a z exists, and (a) shows that $\mathrm{Spec}(A((t)))$ inherits connectedness from $\mathrm{Spec} A$, and so ensures that $Z \cap \mathcal{X}' = \emptyset$. The desired $Z \cong \mathrm{Spec}(A[[t]])$ now follows from the schematic density of the open immersion $\mathrm{Spec}(A((t))) \subset \mathrm{Spec}(A[[t]])$. $\square$

6. The affine Grassmannian. In the setting of §2, if the functor $\mathcal{X}$ is group valued, then so are the functors $L^+_*\mathcal{X}$ and $L^+_*\mathcal{X}$. With this in mind, for a smooth, quasi-affine $A[[t]]$ -group scheme $\mathcal{G}$, its *affine Grassmannian* is the pointed set valued functor defined by

$$\mathrm{Gr}_{\mathcal{G}} := (L\mathcal{G}/L^+\mathcal{G})_{\mathrm{\acute{e}t}}, \tag{6.1}$$

where, as indicated, the sheafification of the presheaf quotient $L\mathcal{G}/L^+\mathcal{G}$ is formed in the étale topology. If instead $\mathcal{G}$ begins life as a smooth, quasi-affine $A\{t\}$ -group scheme, then [BČ22, Example 2.2.19] (which is based on approximation and algebraization techniques) ensures that we may use Henselian loops instead, more precisely, that we have an identification of presheaf quotients

$$L\mathcal{G}/L^+\mathcal{G} \cong L_h\mathcal{G}/L_h^+\mathcal{G}$$

In many situations, the affine Grassmannian $\mathrm{Gr}_{\mathcal{G}}$ is represented by an A -ind-scheme, which is often even ind-projective over A , see, for instance, [PR08]. This proceeds by embedding $\mathcal{G}$ into some GL_n , which is not always possible with our general assumptions, so we will not use representability results.

In the literature one sometimes finds the affine Grassmannian defined as the fpqc sheafification $(L\mathcal{G}/L^+\mathcal{G})_{\mathrm{fpqc}}$. A priori this makes no mathematical sense: even for a field, isomorphism classes of its fpqc covers form a proper class, so the fpqc sheafification may not exist; however, by the following proposition, this “fpqc sheafification approach” gives the same result for our $\mathcal{G}$ as above.

Proposition 7. *For a ring A and a smooth, quasi-affine $A[[t]]$ -group scheme $\mathcal{G}$, the affine Grassmannian $\mathrm{Gr}_{\mathcal{G}}$ is an fpqc sheaf and has the following modular description on the category of A -algebras B :*

$$\mathrm{Gr}_{\mathcal{G}}(B) \cong \left\{ (\mathcal{E}, \iota) \mid \begin{array}{l} \mathcal{E} \text{ is a } \mathcal{G}\text{-torsor over } B[[t], \\ \iota \in \mathcal{E}(B((t))) \text{ is a trivialization over } B((t)) \end{array} \right\} / \sim,$$

while the presheaf quotient $L\mathcal{G}/L^+\mathcal{G}$ is a subfunctor of $\mathrm{Gr}_{\mathcal{G}}$ parametrizing those $(\mathcal{E}, \iota)$ with $\mathcal{E}$ trivial.

Proof. For the trivial $\mathcal{G}$ -torsor, $L^+\mathcal{G}$ parametrizes its $\mathcal{G}$ -torsor automorphisms over $(-)[[t]]$ while $L\mathcal{G}$ parametrizes its trivializations ι over $(-)((t))$, so the presheaf quotient $L\mathcal{G}/L^+\mathcal{G}$ is the claimed subfunctor of $\mathrm{Gr}_{\mathcal{G}}$, granted that the latter has the displayed modular description. Moreover, by the smoothness and quasi-affineness of $\mathcal{E}$ inherited from $\mathcal{G}$ (see [SP, Lemma 0247]) and by the infinitesimal lifting of sections (compare also with [BČ22, Proposition 2.1.4]), each $(\mathcal{E}, \iota)$ lands in this subfunctor étale locally on B , so the sought modular description will follow once we argue that it defines an fpqc sheaf. For this, since the pairs $(\mathcal{E}, \iota)$ have no nontrivial automorphisms, all that remains is to show that the groupoid-valued functor

$$B \mapsto \left\{ (\mathcal{E}, \iota) \mid \begin{array}{l} \mathcal{E} \text{ is a } \mathcal{G}\text{-torsor over } B[[t], \\ \iota \in \mathcal{E}(B((t))) \text{ is a trivialization over } B((t)) \end{array} \right\}$$

is a stack for the fpqc topology on A -algebras B . For this, let $B \rightarrow B'$ be an fpqc cover of A -algebras and let $(\mathcal{E}', \iota')$ be a pair for B' equipped with a descent datum with respect to this cover. It suffices to uniquely descend $\mathcal{E}'$ to a $\mathcal{G}$ -torsor $\mathcal{E}$ over $B[[t]]$: the trivialization ι' will then also descend because $L\mathcal{E}$ is an fpqc sheaf by Proposition 3. The usual fpqc descent for $\mathcal{G}$ -torsors (see [SP, Lemma 0247]) gives us a unique compatible system of $\mathcal{G}$ -torsor descents $\mathcal{E}_n$ over $B[[t]]/(t^{n+1})$ for $n \geq 0$. It then remains to uniquely algebraize this sequence to a $\mathcal{G}$ -torsor $\mathcal{E}$ over $B[[t]]$, and there are several ways to do this (with even more possible arguments if $\mathcal{G}$ is affine). Perhaps the most direct is to apply the algebraization result [BHL17, Theorem 8.1] to the classifying stack $\mathbf{B}\mathcal{G}$. A somewhat more elementary approach is to first use [BČ22, Theorem 2.1.6] to lift $\mathcal{E}_0$ to a $\mathcal{G}$ -torsor $\mathcal{E}$ over $B[[t]]$ and to then inductively build a compatible sequence of isomorphisms

$$\mathcal{E}|_{B[[t]]/(t^{n+1})} \simeq \mathcal{E}_n,$$

by using *loc. cit.* and the smoothness of the quasi-affine $B[[t]]$ -group $\underline{\mathrm{Aut}}_{\mathcal{G}}(\mathcal{E})$. $\square$

2. THE AFFINE GRASSMANNIAN AS A ZARISKI QUOTIENT

Even though the affine Grassmannian of $\mathcal{G}$ is defined as the étale sheafification of the presheaf quotient $L\mathcal{G}/L\mathcal{G}^+$ (see (6.1)), in Theorem 12 below we show that the Zariski sheafification suffices when $\mathcal{G}$ is reductive and descends to A . For intuition for why the Zariski topology may be enough, we recall that the inclusion $L^+\mathcal{G} \subset L\mathcal{G}$ is morally similar to the inclusion $P \subset G$ of a parabolic subgroup of a reductive group scheme, and that $(G/P)_{\mathrm{fpqc}} \cong (G/P)_{\mathrm{Zar}}$ because over any semilocal ring parabolic subgroups of the same type are conjugate, see [SGA 3_{III new}, exposé XXVI, corollaire 5.2].

We will deduce that the Zariski sheafification is enough from the following result about torsors over $\mathbb{P}_A^1$ that is proved by studying the geometry of the algebraic stack $\mathrm{Bun}_{\mathcal{G}}$ that parametrizes such torsors, or in [PS24] by a different approach. Various weaker and more technical earlier variants of this result would suffice as well, for instance, [Fed22, Theorem 6] or [Čes22b, Proposition 5.3.6]. In effect, in some sense, we obtain the main results of this article by ascending geometric information along the uniformization map $\mathrm{Gr}_{\mathcal{G}} \rightarrow \mathrm{Bun}_{\mathcal{G}}$.

Theorem 8 ([ČF23, Theorem 3.6]). *Let G be a reductive group scheme over a semilocal ring A . Every G -torsor E over $\mathbb{P}_A^1$ is A -sectionwise constant, equivalently, $E|_{\{t=0\}} \simeq E|_{\{t=\infty\}}$. $\square$*

We will use Theorem 8 through its following consequence for reductive group torsors over $A[[t]]$.

Proposition 9. *Let A be a semilocal ring and let $\mathcal{G}$ be an $A[[t]]$ -group scheme that is an extension of an $A[[t]]$ -group $\mathcal{G}^{\mathrm{ét}}$ that is locally constant (see §1) by a reductive $A[[t]]$ -group scheme $\mathcal{G}^0$ such that $\mathcal{G}_{A((t))}^0$ descends to a reductive A -group scheme. No nontrivial $\mathcal{G}$ -torsor over $A[[t]]$ trivializes over $A((t))$, that is, we have*

$$\mathrm{Ker}(H^1(A[[t]], \mathcal{G}) \rightarrow H^1(A((t)), \mathcal{G})) = \{*\};$$

in particular, $\mathcal{G}^0$ itself descends to a reductive A -group scheme.

Proof. Let $\mathcal{E}$ be a $\mathcal{G}$ -torsor over $A[[t]]$ that trivializes over $A((t))$. By Corollary 5 (b), every $e \in \mathcal{E}(A((t)))$ gives rise to an $A((t))$ -point of the $\mathcal{G}^{\mathrm{ét}}$ -torsor $\mathcal{E}/\mathcal{G}^0$ that extends uniquely to an $A[[t]]$ -point $\bar{e} \in (\mathcal{E}/\mathcal{G}^0)(A[[t]])$. The preimage of $\bar{e}$ in $\mathcal{E}$ is a $\mathcal{G}^0$ -torsor that trivializes over $A((t))$. We may replace $\mathcal{E}$ by this preimage to reduce to the case when $\mathcal{G} = \mathcal{G}^0$.

Now that $\mathcal{G} = \mathcal{G}^0$, suppose first that $\mathcal{G}$ descends to a reductive A -group G . Due to the triviality over $A((t))$, we may patch $\mathcal{E}$ with the trivial G -torsor over $\mathbb{P}_A^1 \setminus \{t = 0\}$ (see, for instance, [BČ22, Lemma 2.2.11 (b)]) to build a G -torsor E over $\mathbb{P}_A^1$ such that $E|_{\{t=0\}} \simeq \mathcal{E}|_{\{t=0\}}$ and $E|_{\{t=\infty\}}$ is trivial.

By Theorem 8, then $\mathcal{E}|_{\{t=0\}}$ is also trivial, to the effect that, by the infinitesimal lifting of sections due to smoothness, $\mathcal{E}$ is trivial, too, as desired (compare also with [BČ22, Proposition 2.1.4]).

To complete the proof, it remains to show that the reductive A -group scheme G for which we have $G_{A((t))} \simeq \mathcal{G}_{A((t))}$ also descends $\mathcal{G}$, that is, that already $G_{A[[t]]} \simeq \mathcal{G}_{A[[t]]}$. By Corollary 5 (a) and the classification of reductive group schemes (recalled in [Čes22b, Sections 1.3.1 and 1.3.7]), our $\mathcal{G}$ corresponds to an $\underline{\text{Aut}}_{\text{gp}}(G)$ -torsor over $A[[t]]$ that trivializes over $A((t))$. However, as recalled in *loc. cit.*, $\underline{\text{Aut}}_{\text{gp}}(G)$ is an extension of a locally constant A -group by the reductive A -group G^{ad} . In particular, the case settled in the first two paragraphs of the proof applies and shows that the $\underline{\text{Aut}}_{\text{gp}}(G)$ -torsor in question is trivial already over $A[[t]]$, so that $G_{A[[t]]} \simeq \mathcal{G}$, as desired. $\square$

Corollary 10. *For a reductive group scheme G over a semilocal ring A , we have*

$$H^1(A, G) \hookrightarrow H^1(A((t)), G).$$

Proof. By twisting (see [Čes22b, equation (1.2.1.1)]), it suffices to show that the map in question has trivial kernel. This follows from Proposition 9 because, by [BČ22, Theorem 2.1.6], we have

$$H^1(A, G) \xrightarrow{\sim} H^1(A[[t]], G). \quad \square$$

Remark 11. Results of [FG21] suggest that the reductivity assumption may be nonessential for Corollary 10. It would therefore be interesting to find a more general result of this type.

We are ready for the promised sufficiency of the Zariski sheafification for $\text{Gr}_{\mathcal{G}}$ in the case when $\mathcal{G}$ is (constant) reductive. One may compare this to its earlier variant [Bac19, Proposition 14] that restricted to smooth A over a field and deduced the conclusion from the Grothendieck–Serre conjecture. For us, it is the study of the latter that has indirectly led to the results of this article.

Theorem 12. *For a ring A and an $A[[t]]$ -group scheme $\mathcal{G}$ that is an extension of a finite étale $A[[t]]$ -group $\mathcal{G}^{\text{ét}}$ by a reductive $A[[t]]$ -group scheme $\mathcal{G}^0$ that descends to a reductive A -group scheme, the affine Grassmannian $\text{Gr}_{\mathcal{G}}$ is the Zariski sheafification of the presheaf quotient $L\mathcal{G}/L^+\mathcal{G}$, that is,*

$$\text{Gr}_{\mathcal{G}} \cong (L\mathcal{G}/L^+\mathcal{G})_{\text{Zar}}.$$

Proof. By the modular interpretation supplied by Proposition 7, the possibility of varying A , and the infinitesimal lifting of sections due to smoothness, all we need to show is that for any $\mathcal{G}$ -torsor $\mathcal{E}$ over $A[[t]]$ that trivializes over $A((t))$, the $\mathcal{G}$ -torsor $\mathcal{E}|_{\{t=0\}}$ trivializes Zariski locally on A . However, Proposition 9 ensures that $\mathcal{E}|_{\{t=0\}}$ trivializes even Zariski semilocally on A . $\square$

Remark 13.

- (1) We do not know whether the reductivity assumption is critical for Theorem 12, for instance, whether the Zariski sheafification also suffices for parahoric groups. Certainly, it does in the case when $\mathcal{G}$ is a reductive $A[[t]]$ -group scheme and $\mathcal{P}$ is a smooth, quasi-affine $A[[t]]$ -group scheme equipped with an $A[[t]]$ -morphism $\mathcal{P} \rightarrow \mathcal{G}$ that modulo t reduces to an inclusion of a parabolic subgroup: indeed, by [BČ22, Theorem 2.1.6] and [Čes22b, equation (1.3.5.2)], these assumptions ensure that $H^1(B[[t]], \mathcal{P}) \subset H^1(B[[t]], \mathcal{G})$ for any semilocal A -algebra B , so

$$\text{Ker}(H^1(B[[t]], \mathcal{G}) \rightarrow H^1(B((t)), \mathcal{G})) = \{*\} \implies \text{Ker}(H^1(B[[t]], \mathcal{P}) \rightarrow H^1(B((t)), \mathcal{P})) = \{*\},$$

to the effect that Proposition 7 and Theorem 12 imply the claimed $\text{Gr}_{\mathcal{P}} \cong (L\mathcal{P}/L^+\mathcal{P})_{\text{Zar}}$.

- (2) We do not know whether Theorem 12 admits a version for the Witt vector affine Grassmannian. For a version of Theorem 12 for the B_{dR}^+ -affine Grassmannian, see [ČY24, Theorem 3.1].

3. THE AFFINE GRASSMANNIAN AS A PRESHEAF QUOTIENT

For most reductive group schemes $\mathcal{G}$, even the Zariski sheafification is not needed when forming the affine Grassmannian $\text{Gr}_{\mathcal{G}}$: in Theorem 17 below, we show that the latter often agrees already with the presheaf quotient $L\mathcal{G}/L^+\mathcal{G}$. This appears to be new already for reductive groups G over $\mathbb{C}$, although for GL_n it essentially follows from [BČ22, Theorem 2.1.24], and is based on the finer variant of Theorem 8 recorded in Theorem 16 below, which, in addition to the geometry of Bun_G , uses Quillen patching for torsors over $\mathbb{A}_A^1$ to progress beyond semilocal A . For this variant, the relevant condition on $\mathcal{G}$ is the following.

Definition 14 ([Čes22a, Definition 8.1]). A semisimple group scheme G over a scheme S is *totally isotropic* if in the canonical decomposition of [SGA 3III new, exposé XXIV, proposition 5.10 (i)]:

$$G^{\text{ad}} \cong \prod_{i \in \{A_n, B_n, \dots, G_2\}} \text{Res}_{S_i/S}(G_i)$$

of its adjoint quotient G^{ad} , where i ranges over the types of connected Dynkin diagrams, S_i is a finite étale S -scheme, and G_i is an adjoint semisimple S_i -group with simple geometric fibers of type i , Zariski locally on S each G_i has a parabolic S_i -subgroup that contains no S_i -fiber of G_i (equivalently, Zariski locally on S each $\text{Res}_{S_i/S}(G_i)$ contains a nontrivial split torus $\mathbb{G}_{m,S}$, see [SGA 3III new, exposé XXVI, corollaire 6.12] and [Čes22b, end of Section 1.3.4]).

Example 15. Slightly informally, G is totally isotropic if Zariski locally on S it has a parabolic subgroup containing no factor of the adjoint group G^{ad} . To see this, recall that parabolic subgroups of G correspond to those of G^{ad} , which correspond to collections of parabolic subgroups of G_i , one for each i , see [Čes22b, end of Section 1.3.4]. Certainly, every quasi-split (so also every split) semisimple group is totally isotropic.

Theorem 16 ([ČF23, Theorem 4.2]). *Let G be a totally isotropic reductive group scheme over a ring A . For a G -torsor E over $\mathbb{P}_A^1$, if $E|_{\{t=\infty\}}$ is trivial, then $E|_{\mathbb{A}_A^1}$ is also trivial.* $\square$

Theorem 17. *For a ring A and an $A[[t]]$ -group scheme $\mathcal{G}$ that is an extension of a finite étale $A[[t]]$ -group $\mathcal{G}^{\text{ét}}$ by a reductive $A[[t]]$ -group scheme $\mathcal{G}^0$ that descends to a reductive A -group scheme whose adjoint quotient is totally isotropic, the affine Grassmannian $\text{Gr}_{\mathcal{G}}$ is the presheaf quotient $L\mathcal{G}/L^+\mathcal{G}$, that is,*

$$\text{Gr}_{\mathcal{G}} \cong L\mathcal{G}/L^+\mathcal{G}.$$

Proof. As in the proof of Theorem 12, by the modular description of Proposition 7 and the possibility of varying A , we need to show that no nontrivial $\mathcal{G}$ -torsor $\mathcal{E}$ over $A[[t]]$ trivializes over $A((t))$. For this, as in the proof of Proposition 9, Corollary 5 (b) immediately reduces us to the case when $\mathcal{G} = \mathcal{G}_0$ and $\mathcal{G}$ is the base change of a reductive A -group scheme G whose adjoint quotient is totally isotropic.

To treat this case, we again patch $\mathcal{E}$ with the trivial G -torsor over $\mathbb{P}_A^1 \setminus \{t = 0\}$ to build a G -torsor E over $\mathbb{P}_A^1$ such that $E|_{A[[t]]} \simeq \mathcal{E}$ and $E|_{\{t=\infty\}}$ is trivial. By Theorem 16, this last condition forces $E|_{\mathbb{A}_A^1}$ to be trivial. However, then $\mathcal{E}$ is trivial, too, as desired. $\square$

We recall from [BČ22, Theorems 2.1.24 and 3.1.7] that the following global variant of Corollary 10 was known when G is either a pure inner form of GL_n or a torus.

Corollary 18. *For a reductive group scheme G over a ring A with G^{ad} totally isotropic, no nontrivial G -torsor over A trivializes over $A((t))$, in other words, we have*

$$\text{Ker}(H^1(A, G) \rightarrow H^1(A((t)), G)) = \{*\}.$$

Proof. Proposition 7 and Theorem 17 show that no nontrivial G -torsor over $A[[t]]$ trivializes over $A((t))$. Thus, it suffices to recall from [BČ22, Theorem 2.1.6] that

$$H^1(A, G) \xrightarrow{\sim} H^1(A[[t]], G). \quad \square$$

Finally, we note that the preceding proof shows that the affine Grassmannian in the totally isotropic case may even be formed using polynomial loops as follows.

Theorem 19. *For a reductive group scheme G over a ring A with G^{ad} totally isotropic, its affine Grassmannian Gr_G may be formed as the presheaf quotient using the polynomial loops, more precisely, we have*

$$\text{Gr}_G \cong L_{\text{poly}}G / L_{\text{poly}}^+G,$$

explicitly, we have

$$\begin{aligned} G(A((t)))/G(A[[t]]) &\cong G(A\{t\}[\tfrac{1}{t}])/G(A\{t\}) \\ &\cong G((A[t]_{1+tA[t]})[\tfrac{1}{t}])/G(A[t]_{1+tA[t]}) \cong G(A[t, t^{-1}])/G(A[t]), \end{aligned}$$

equivalently,

$$\begin{aligned} G(A((t))) &= G(A[t, t^{-1}])G(A[[t]]), \\ G(A\{t\}[\tfrac{1}{t}]) &= G(A[t, t^{-1}])G(A\{t\}), \\ G((A[t]_{1+tA[t]})[\tfrac{1}{t}]) &= G(A[t, t^{-1}])G(A[t]_{1+tA[t]}). \end{aligned}$$

These equalities seem elementary, but we do not know how to argue them directly even for $G = \text{GL}_n$. For instance, in terms of Beauville–Laszlo patching used in the proof below, one would need to argue that every finite projective $A[t]$ -module that is free both over $A[t, t^{-1}]$ and over $A[[t]]$ is free.

Proof. Since G is affine, its functor of points preserves the Cartesianness of the squares from Corollary 5 (a). In particular, the map

$$G(A[t, t^{-1}])/G(A[t]) \hookrightarrow G(A((t)))/G(A[[t]])$$

is injective, and so are its counterparts for algebraic or Henselian loops in place of polynomial loops. Thus, all we need to show is that this map is also surjective or, equivalently, that

$$G(A[t, t^{-1}]) \backslash G(A((t)))/G(A[[t]]) = \{*\}.$$

However, if this double quotient was nontrivial, then we could use patching (for instance, [BČ22, Lemma 2.2.11 (b)]) to build a nontrivial G -torsor over $\mathbb{A}_A^1$ that would trivialize both over $\mathbb{G}_{m, A}$ and also over the formal completion along $\{t = 0\}$. We could then extend this G -torsor to all of $\mathbb{P}_A^1$ by patching with the trivial torsor at infinity, and thus obtain a contradiction to Theorem 16. $\square$

Remark 20. We do not know the extent to which the assumptions of Theorems 17 and 19 are optimal because they are imposed by our proofs, for instance, we do not know whether the affine Grassmannian Gr_G agrees with the presheaf quotient LG/L^+G for every (possibly not totally isotropic) reductive A -group G , although we expect that it does not. On the other hand, since our proofs are reductions to general results about torsors over $\mathbb{P}_A^1$, the reader will have no trouble adapting them to various close variants of the affine Grassmannian that are sometimes considered

in the literature, for instance, to affine Grassmannians constructed using general relative Cartier divisors in $\mathbb{A}_A^1$ in place of $\{t = 0\}$.

Acknowledgements. This article was written in relation to the Jacques Herbrand prize of the French Academy of Sciences. I thank the Academy for the prize and for the invitation to submit to the Comptes Rendus de l'Académie des Sciences on this occasion. I thank the referee for helpful remarks and suggestions. I thank Alexis Bouthier, Ofer Gabber, and Alex Youcis for helpful conversations and correspondence. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 851146). This project is based upon work supported by the National Science Foundation under Grant No. DMS-1926686.

REFERENCES

- [Ana73] Sivaramakrishna Anantharaman, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, Sur les groupes algébriques, Soc. Math. France, Paris, 1973, pp. 5–79. Bull. Soc. Math. France, Mém. 33 (French). MR0335524 (49 #305)
- [Bac19] Tom Bachmann, *Affine Grassmannians in $\mathbb{A}^1$ -homotopy theory*, Selecta Math. (N.S.) **25** (2019), no. 2, Paper No. 25, 14, DOI 10.1007/s00029-019-0471-1. MR3925100
- [BČ22] Alexis Bouthier and Kęstutis Česnavičius, *Torsors on loop groups and the Hitchin fibration*, Ann. Sci. École Norm. Sup. **55** (2022), no. 3, 791–864.
- [BD] A. Beilinson and V. Drinfeld, *Quantization of Hitchin's integrable system and Hecke eigensheaves*, preprint. Accessed at <https://math.uchicago.edu/~drinfeld/langlands/QuantizationHitchin.pdf> on November 27, 2023.
- [Bha16] Bhargav Bhatt, *Algebraization and Tannaka duality*, Camb. J. Math. **4** (2016), no. 4, 403–461, DOI 10.4310/CJM.2016.v4.n4.a1. MR3572635
- [BHL17] Bhargav Bhatt and Daniel Halpern-Leistner, *Tannaka duality revisited*, Adv. Math. **316** (2017), 576–612, DOI 10.1016/j.aim.2016.08.040. MR3672914
- [CLNS18] Antoine Chambert-Loir, Johannes Nicaise, and Julien Sebag, *Motivic integration*, Progress in Mathematics, vol. 325, Birkhäuser/Springer, New York, 2018. MR3838446
- [Čes22a] Kęstutis Česnavičius, *Grothendieck–Serre in the quasi-split unramified case*, Forum Math. Pi **10** (2022), Paper No. e9, 30, DOI 10.1017/fmp.2022.5. MR4403660
- [Čes22b] Kęstutis Česnavičius, *Problems About Torsors over Regular Rings*, Acta Math. Vietnam. **47** (2022), no. 1, 39–107, DOI 10.1007/s40306-022-00477-y. MR4406561
- [ČF23] Kęstutis Česnavičius and Roman Fedorov, *Unramified Grothendieck–Serre for isotropic groups*, preprint (2023). Available at <https://arxiv.org/abs/2311.08660>.
- [ČY24] Kęstutis Česnavičius and Alex Youcis, *The analytic topology suffices for the B_{dR}^+ -Grassmannian*, Proceedings of the Simons Symposium on p-adic Hodge theory (2022), to appear (2024). Available at <https://arxiv.org/abs/2303.11710v3>.
- [EGA I] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique. I. Le langage des schémas*, Inst. Hautes Études Sci. Publ. Math. **4** (1960), 228. MR0217083 (36 #177a)
- [EGA IV₄] ———, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas*, Inst. Hautes Études Sci. Publ. Math. **32** (1967), 361 (French). MR0238860 (39 #220)
- [Fed22] Roman Fedorov, *On the Grothendieck–Serre conjecture about principal bundles and its generalizations*, Algebra Number Theory **16** (2022), no. 2, 447–465, DOI 10.2140/ant.2022.16.447. MR4412579
- [FG21] Mathieu Florence and Philippe Gille, *Residues on affine Grassmannians*, J. Reine Angew. Math. **776** (2021), 119–150, DOI 10.1515/crelle-2021-0007. MR4279105
- [Lus83] George Lusztig, *Singularities, character formulas, and a q -analog of weight multiplicities*, Analysis and topology on singular spaces, II, III (Luminy, 1981), Astérisque, vol. 101-102, Soc. Math. France, Paris, 1983, pp. 208–229. MR0737932
- [PR08] G. Pappas and M. Rapoport, *Twisted loop groups and their affine flag varieties*, Adv. Math. **219** (2008), no. 1, 118–198, DOI 10.1016/j.aim.2008.04.006. With an appendix by T. Haines and Rapoport. MR2435422
- [PS24] Ivan Panin and Anastasia Stavrova, *On the Gille theorem for the relative projective line*, preprint (2024). Available at <https://arxiv.org/abs/2305.16627v2>.

- [Ray70] Michel Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics, Vol. 119, Springer-Verlag, Berlin, 1970 (French). MR0260758 (41 #5381)
- [RG71] Michel Raynaud and Laurent Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), 1–89, DOI 10.1007/BF01390094 (French). MR0308104
- [SGA 3_I new] Philippe Gille and Patrick Polo (eds.), *Schémas en groupes (SGA 3). Tome I. Propriétés générales des schémas en groupes*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 7, Société Mathématique de France, Paris, 2011 (French). Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64]; A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M. Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre; Revised and annotated edition of the 1970 French original. MR2867621
- [SGA 3_{II}] *Schémas en groupes. II: Groupes de type multiplicatif, et structure des schémas en groupes généraux*, Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 152, Springer-Verlag, Berlin-New York, 1970 (French). MR0274459 (43 #223b)
- [SGA 3_{III} new] Philippe Gille and Patrick Polo (eds.), *Schémas en groupes (SGA 3). Tome III. Structure des schémas en groupes réductifs*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 8, Société Mathématique de France, Paris, 2011 (French). Séminaire de Géométrie Algébrique du Bois Marie 1962–64. [Algebraic Geometry Seminar of Bois Marie 1962–64]; A seminar directed by M. Demazure and A. Grothendieck with the collaboration of M. Artin, J.-E. Bertin, P. Gabriel, M. Raynaud and J-P. Serre; Revised and annotated edition of the 1970 French original. MR2867622
- [SP] A. J. de Jong et al., *The Stacks Project*. Available at <http://stacks.math.columbia.edu>.