

**Multiplicity and vanishing lemmas
for differential and q -difference equations
in the Siegel-Shidlovsky theory.**

D. Bertrand (Paris VI)

Gel'fond 100, Moscow, Jan. 07

Abstract

We present a general multiplicity estimate for linear forms in solutions of various type of functional equations, which covers and extends the zero estimates used in recent work on the Siegel-Shidlovsky theorem and its q -analogues. We also present a dual version of this estimate, as well as a new interpretation of Siegel's theorem itself in terms of periods of Deligne's irregular Hodge theory.

Plan

- 1. A bit of history on Siegel-Shidlovsky**
- 2. Yet another multiplicity estimate ...
What for ?**
- 3. Generalized Shidlovsky lemmas**
- 3. Vanishing lemmas**
- 4. Deligne's periods**

XXth century

$n > 0$, $[K : \mathbf{Q}] = \kappa$, $K \subset \mathbf{C}$; $K \ni \gamma \rightarrow 1$

$$\frac{d}{dz} \begin{pmatrix} \mathcal{E}_1 \\ \vdots \\ \mathcal{E}_n \end{pmatrix} = A(z) \begin{pmatrix} \mathcal{E}_1 \\ \vdots \\ \mathcal{E}_n \end{pmatrix} \quad (*)$$

where $A(z) \in gl_n(K(z) \cap K[[z-1]])$.

$\mathcal{E} = (\mathcal{E}_1, \dots, \mathcal{E}_n)$, KE -functions, generating a $\mathbf{C}(z)$ -vector space of dimension $n(\mathcal{E})$.

$\mathcal{E}(1) = (\mathcal{E}_1(1), \dots, \mathcal{E}_n(1))$, "generating" a K -vector space W_1 of dimension $r := r_1(\mathcal{E})$.

Theorem (Siegel-Shidlovsky) : $r_1(\mathcal{E}) \geq \frac{n(\mathcal{E})}{\kappa}$.

Nesterenko-Shidlovsky (1996) : if $K \rightarrow \overline{\mathbf{Q}}$, then $r_\gamma(\mathcal{E}) = n(\mathcal{E})$ for a.a. γ 's $\in \overline{\mathbf{Q}}$.

XXI th century

Y. André (2000) : new proof of S-Sh. The fundamental lemma is : *let f be a QE -function, and let $\mathcal{L} \in \mathbb{C}(z)[d/dz]$ of minimal order such that $\mathcal{L}(f) = 0$. If $f(1) = 0$, then, all solutions of \mathcal{L} vanish at $z = 1$.* Then, as in the Gel'fond-Dèbes method from the theory of G -functions, construct an auxiliary KE -function with high multiplicity at $z = 1$, rather than at 0. Take the product of its conjugates to get a QE -function ($\Rightarrow \frac{1}{\kappa}$).

D.B. (2004) : new proof of S-Sh., based on Laurent interpolation determinants. Requires a new type of multiplicity (or vanishing) lemma, more on this later. No auxiliary function, and the roles of 0 and 1 are parallel. Cf. A. Sert (1999) in the Lindemann-Weierstrass case.

F. Beukers (2006) : $r_1(\mathcal{E}) = n(\mathcal{E}) !!!$

In other words, S-Sh. is valid over $\overline{\mathbf{Q}}$. The proof is based on André's lemma and on differential Galois theory. The output is that André's lemma is valid for KE -functions, hence no loss of $\frac{1}{\kappa}$ in the final estimate.

Meanwhile, in the q -difference world :

$$\begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} (qz) = A(z) \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} (z) \quad (*_q)$$

where $A(z) \in GL_n(K(z))$.

$Y := (y_1, \dots, y_n)$ analytic at 0 with $n(Y) = n$, $0 \neq s = (p_1, \dots, p_n) \in (\mathbf{C}[z])^n$, $\deg(s) \leq L$, $s.Y = p_1 y_1 + \dots + p_n y_n$, $s_k.Y(z) = (s.Y)(q^k z)$, generating a $\mathbf{C}(z)$ v.-s. of dimension ν . Then :

M. Amou, T. Mataha-Alo, K. Väänänen (2003, 2006) : $ord_0(s.Y) \leq \nu L + c$.

Applications in the style of Siegel-Shidlovsky : see Keijo's talk on Wednesday.

D.B. (2006) : new type of multiplicity estimates, involving 0 and $q^{\mathbf{N}}$ -orbits. No application yet.

What for ?

Recall $W_1 =$ smallest K -v-s. through $\mathcal{E}(1) = (\mathcal{E}_1(1), \dots, \mathcal{E}_n(1))$, of dimension $r := r_1(\mathcal{E})$, assume $n(\mathcal{E}) = n$, and let $\mathcal{Z}_1, \dots, \mathcal{Z}_r$ be a basis of solutions of $(*)$ whose values at 1 lie in W_1 . Fix parameters $L, T_0, T_1 \in \mathbf{N}$, and consider the linear map (with $\partial = d/dz$) :

$$\phi : (\mathbf{C}[z]_{\leq L})^n \rightarrow \mathbf{C}^{T_0} \oplus \mathbf{C}^{rT_1}$$

$$\dim = n(L + 1) \qquad \dim = T_0 + rT_1$$

$$s = (p_1, \dots, p_n) \mapsto (\partial^t(s.\mathcal{E})(0)_{t < T_0}; \partial^t(s.\mathcal{Z}_\rho)(1)_{t < T_1})$$

represented by the $(T_0 + rT_1) \times n(L + 1)$ matrix $\Phi =$

$$\left(\begin{array}{c} \Phi_0 = \left(\partial^t(s_i.\mathcal{E})(0) \right)_{0 \leq t \leq T_0 - 1; 1 \leq i \leq L + 1} \\ \dots \\ \Phi_\rho = \left(\partial^t(s_i.\mathcal{Z}_\rho)(1) \right)_{0 \leq t \leq T_1 - 1; 1 \leq i \leq L + 1} \\ \dots \qquad \qquad \qquad (\rho = 1, \dots, r) \end{array} \right).$$

where $s_i, i \leq (L + 1)^n$ is a basis of $(\mathbf{C}[z]_{\leq L})^n$.

If we knew that

" $n(L + 1) < T_0 + rT_1 \Rightarrow \phi$ injective",
 or " $n(L + 1) > T_0 + rT_1 \Rightarrow \phi$ surjective",

then the proof would consist of two words :
just look !

$$\left(\begin{array}{c} \Phi_0 = \left(\partial^t \left(\frac{1}{\ell!} z^\ell \mathcal{E}_\iota \right) (0) \right)_{0 \leq t \leq T_0 - 1; 1 \leq \iota \leq n, 0 \leq \ell \leq L} \\ \dots \\ \Phi_\rho = \left(\partial^t \left(\frac{1}{\ell!} z^\ell \mathcal{Z}_{\rho, \iota} \right) (1) \right)_{0 \leq t \leq T_1 - 1; 1 \leq \iota \leq n, 0 \leq \ell \leq L} \\ \dots \qquad \qquad \qquad (\rho=1, \dots, r) \end{array} \right) \cdot$$

(and extract a $n(L + 1)$ - (or $T_0 + rT_1$ -) minor determinant $\Delta \in K^*$, whose height forces

$$\boxed{T_0 T_1 \leq r\kappa L T_1 + r(\kappa + 1) T_1^2 + O(L^2 / \text{Log} L),}$$

hence $n \leq r\kappa$, if $T_0 = (n - \epsilon)L, T_1$ small.)

For Lindemann-Weierstrass, one can also use :

$$\left(\begin{array}{c} \Phi_0 = \left(\partial^t \left(\frac{1}{\ell!} (z - 1)^\ell \mathcal{E}_\iota \right) (0) \right)_{0 \leq t \leq T_0 - 1; 1 \leq \iota \leq n, 0 \leq \ell \leq L} \\ \dots \\ \Phi_\rho = \left(\partial^t \left(\frac{1}{\ell!} (z - 1)^\ell \mathcal{Z}_{\rho, \iota} \right) (1) \right)_{0 \leq t \leq T_1 - 1; 1 \leq \iota \leq n, 0 \leq \ell \leq L} \\ \dots \qquad \qquad \qquad (\rho=1, \dots, r) \end{array} \right)$$

(and conclude that $\boxed{T_0 T_1 \leq \kappa T_0 L + O(L^2 / \text{Log} L),}$
 hence $n \leq r\kappa$, if $T_1 = \left(\frac{n}{r} - \epsilon\right)L, T_0$ small.)

Generalized Shidlovsky lemmas

Write $(\mathcal{M} = \mathbf{C}(z)^n, \nabla)$ for $(*)$, with set of singularities S . Let $\mathcal{R} \subset \mathbf{C}$ be a finite set, and for all $\alpha \in \mathcal{R}$, let $\hat{\mathcal{W}}_\alpha$ be a \mathbf{C} -subspace of $\hat{\mathcal{M}}_\alpha = (K[[z - \alpha]])^n$ formed by solutions of ∇ . A linear form s in $\mathcal{M}^*(L) = (\mathbf{C}[z]_{\leq L})^n$ vanishes to an order $\geq T$ along $\hat{\mathcal{W}}_\alpha$ if for all $\mathcal{Z} \in \hat{\mathcal{W}}_\alpha$, $s \cdot \mathcal{Z}$ vanishes to an order $\geq T$.

Differential multiplicity lemma : $\exists c(\nabla)$, computable in terms of \mathcal{M}, ∇ and $\text{card}(\mathcal{R})$, such that : let $\{T_\alpha, \alpha \in \mathcal{R}; L\} \in \mathbf{N}$, and $0 \neq s \in \mathcal{M}^*(L)$ vanishing to an order $\geq T_\alpha$ along $\hat{\mathcal{W}}_\alpha$, for all $\alpha \in \mathcal{R}$. Then, there exists a subspace \mathcal{M}' in $\text{Ker}(s) \subset \mathcal{M}$ stable under ∇ , such that

$$\sum_{\alpha \in \mathcal{R}} \dim(\hat{\mathcal{W}}_\alpha / \hat{\mathcal{W}}_\alpha \cap \hat{\mathcal{M}}'_\alpha) \cdot T_\alpha \leq \text{rk}(\mathcal{M} / \mathcal{M}') \cdot L + c(\nabla).$$

[And we may in fact take for \mathcal{M}' the maximal ∇ -stable subspace of $\text{Ker}(s)$.]

$\mathcal{R} = \{0, 1\}$, $\dim(\widehat{\mathcal{W}}_0) = 1$, $r = \dim(\widehat{\mathcal{W}}_1)$. Say that $\widehat{\mathcal{W}}_1$ is *non degenerate* if for all $\mathcal{M}' \neq \mathcal{M}$ stable under ∇ , we have :

$$\frac{r'}{n'} := \frac{\dim(\widehat{\mathcal{W}}_1 / \widehat{\mathcal{W}}_1 \cap \widehat{\mathcal{M}}'_1)}{\text{rk}(\mathcal{M} / \mathcal{M}')} \geq \frac{\dim(\widehat{\mathcal{W}}_1)}{\text{rk}(\mathcal{M})} := \frac{r}{n}$$

(NB : $n(\mathcal{E}) = n \Leftrightarrow \widehat{\mathcal{W}}_0$ non-degenerate.)

Corollary : let $T_0, T_1, L \in \mathbb{N}$, let $s \in \mathcal{M}^*(L)$ vanishing to an order $\geq T_\alpha$ along $\widehat{\mathcal{W}}_\alpha$, $\alpha = 0, 1$. Assume the $\widehat{\mathcal{W}}_\alpha$'s are non-degenerate, and that $T_0 + rT_1 > nL + nc(\nabla)$. Then, $s = 0$. In other words, ϕ is injective.

(NB : could replace the non-degeneracy of $\widehat{\mathcal{W}}_1$ by $L > T_1$.) Forgetting $\alpha = 1$, this implies Shidlovsky's original lemma that if the order of $s.\mathcal{E}$ at $\alpha = 0$ is almost nL , then, the linear forms $s = s_1, \nabla^* s = s_2, \dots, s_n$ are linearly independent.

In the q -difference world

Let $|q| < 1$. For $\alpha \in \mathbf{C}^*$, the positive (resp. negative) orbit of α is $\{q^n \alpha, n \geq 0\}$ (resp. $n \leq 0$).

$f(z)$ in the Nielsen class (of quasiunipotent type) means : a polynomial in a fractional power of z and in $\text{Log}z$, whose coefficients are meromorphic functions near 0. Given $\alpha \in \mathbf{C}^*$ and some determination of $\text{Log}z$ such that f is defined on the positive orbit of α , set :

$$\text{ord}_\alpha^q(f) = \sup\{t \in \mathbf{N}, f(\alpha) = \dots = f(q^{t-1}\alpha) = 0\}.$$

When $f \neq 0$, this is a finite number := the *order* of f at α relatively to the q -difference operator $\delta_q : f \rightarrow \delta_q f$, where $\delta_q f(z) = \frac{f(qz) - f(z)}{qz - z}$.

If $\alpha = 0$ and f is analytic at 0, $\text{ord}_0^q(f) := \text{ord}_0(f)$ is the order of f at 0 in the usual sense, i.e. relatively to $\delta_q.(0) := \frac{d}{dz}|_0$; indeed, $\frac{d}{dz}f(0)$ is the limit of $\delta_q(f)(\alpha)$ when α tends to 0.

Write $M = (\mathbf{C}(z))^n, \Psi$, $\Psi Y(z) = A(z)^{-1}Y(qz)$ for $(*_q)$, and assume that Ψ is *regular singular at 0*, with quasi-unipotent local monodromy. No assumption at ∞ (e.g. regular and confluent q -hypergeometric equations). Then, the Nielsen type solutions of Ψ form a \mathbf{C} -vector space M^Ψ of dimension n .

For $\alpha \neq 0, \alpha \notin \text{Sing}(A)$, let W_α be a \mathbf{C} -subspace of M^Ψ and let $s = (p_1, \dots, p_n) \in (\mathbf{C}[z])^n$ be a linear form on M . For any $Y = (y_1, \dots, y_n)^t \in W_\alpha$, the Nielsen type function

$$s.Y(z) = p_1(z)y_1(z) + \dots + p_n(z)y_n(z)$$

is defined on the positive orbit of α , and we may speak of its q -order $ord_\alpha^q(s.Y)$ at α . We then set :

$$ord_{W_\alpha}^q(s) = \min(ord_\alpha^q(s.Y); Y \in W_\alpha).$$

This expression still makes sense if $\alpha = 0$, as long as the \mathbf{C} -subspace W_0 consists of solutions all of whose coordinates are analytic at 0 : then, $ord_{W_0}^q(s)$ is the order of s along W_0 in the previous (differential) sense.

Let $\mathcal{R} = \{\alpha_1, \dots, \alpha_r\}$ be a finite set of complex nbs, possibly including 0 but not meeting the negative q -orbit of $Sing(A)$, and whose classes modulo $q^{\mathbb{Z}}$ are distinct. For all $\alpha \in \mathcal{R}$, let $W_\alpha \subset M^\Psi$ be a \mathbb{C} -subspace of solutions of Ψ (analytic at 0 if $\alpha = 0$).

q -difference multiplicity lemma : $\exists c(\Psi)$, depending only on (M, Ψ) and $card(\mathcal{R})$, such that : let $\{T_\alpha, \alpha \in \mathcal{R}; L\} \in \mathbb{N}$, and $0 \neq s \in M^*(L)$ vanishing to an order $\geq T_\alpha$ along W_α , for all $\alpha \in \mathcal{R}$. Then, the maximal subspace $M' \subset Ker(s) \subset M$ stable under Ψ satisfies :

$$\sum_{\alpha \in \mathcal{R}} dim(W_\alpha / W_\alpha \cap M') \cdot T_\alpha \leq rk(M/M') \cdot L + c(\Psi).$$

Same corollaries as earlier, e.g. :

(Väänänen's "Shidlovsky lemma") : the dimension ν of the $\mathbb{C}(z)$ -subspace of $M^*(L)$ generated by $s = s_1, \Psi^*s = s_2, \dots, s_n$ satisfies : $ord_0(s.Y) \leq \nu L + c$.

\Rightarrow non-vanishing of the n -order determinant \Rightarrow independence results.

Also : assume $\mathcal{R} = \{0, 1\}$, $\dim W_0 = 1$, $\dim W_1 = r$, $\text{ord}_{W_0}^q(s) \geq T_0$, $\text{ord}_{W_1}^q(s) \geq T_1$, $L > T_1$, and $T_0 + rT_1 > nL + c(\Psi)$. Then $s = 0$.

\Rightarrow non vanishing of the $n(L + 1)$ -order determinant $\Rightarrow ?$

Proof of the multiplicity lemmas

As in Shidklovsky, the crucial point is that the $\mathbf{C}(z)$ -subspaces of \mathcal{M} (resp. M) stable under ∇ (resp. Ψ) are definable by linear forms with degrees bounded by a constant γ depending only on ∇ (resp. Ψ). However, while Fuchs's relation (or methods from symbolic algebra) provides effective estimates for $\gamma(\nabla)$ in terms of the coefficients of the matrix $A(z)$, the present status of $\gamma(\Psi)$ seems ineffective. The problem reduces to finding a priori upper bounds for the degree of the rational solutions of a linear q -difference operator $\mathcal{L}y = y(q^\mu z) + a_{\mu-1}y(q^{\mu-1}z) + \dots + a_0y(z)$ with coefficients in $\mathbf{C}(z)$, regular singular at 0.

Vanishing lemmas

These are “interpolation lemmas”, which imply the *surjectivity* of ϕ , and can therefore be viewed as vanishing criteria for the H^1 of certain sheaves (hence their name). They should be easier to prove than the multiplicity lemmas, but for the moment, the deduction goes the reverse way, following a method of D. Masser and S. Fischler. Here is an example in the differential case.

On top of the previous assumption that the line $\widehat{\mathcal{W}}_0$ and the subspace $\widehat{\mathcal{W}}_1$ are non-degenerate, we suppose that $\mathcal{E}(0) \neq 0$, and that 1 is not a singularity of ∇

Differential vanishing lemma : $\exists \widehat{c}(\nabla)$ computable in terms of (\mathcal{M}, ∇) such that : let $\{a_{0,t}, 0 \leq t \leq T_0 - 1, a_{\rho,t}, 1 \leq \rho \leq r, 0 \leq t \leq T_1 - 1\}$ be a $(T_0 + rT_1)$ -uple of complex numbers. Let further $T_0, T_1, L \in \mathbb{N}$ satisfy $nL \geq T_0 + rT_1 + \widehat{c}(\nabla)$. Then, there exists a linear form $s \in \mathcal{M}^*(L)$ such that $\partial^t(s.\mathcal{E})(0) = a_{0,t}$ for all $t \leq T_0 - 1$ and $\partial^t(s.\mathcal{Z}_\rho)(1) = a_{\rho,t}$ for all $\rho = 1, \dots, r, t \leq T_1 - 1$.

Deligne's periods

Irregular singularities provide theorems : Siegel-Shidlovsky's !

Regular singularities provide conjectures : Grothendieck's on periods.

Deligne's "irregular periods" : in the case of e^{-z^2} , set

$$H_{dR}^1 = \{e^{-z^2} \mathbf{Q}[z] dz\} / d(\{(e^{-z^2} \mathbf{Q}[z])\}) \simeq \mathbf{Q} e^{-z^2} dz$$

$$H_1^B = \mathbf{Z} \cdot \gamma, \gamma = \text{the real line } \mathbf{R}.$$

Period : $\int_{-\infty}^{+\infty} e^{-z^2} dz = \sqrt{\pi}$ (not a period in the motivic sense).

Irregular periods in a family : consider $e^{z+\lambda/z}$, $\lambda \in K$ (a "Legendre" parameter)

$$H_{dR}^1 = \{P(z, z^{-1}) e^{z+\lambda/z} \frac{dz}{z} / \text{exact forms} \}$$

$$\simeq K\omega \oplus K\eta, \omega = e^{z+\lambda/z} \frac{dz}{z}, \eta = e^{z+\lambda/z} dz$$

$H_1^B = \mathbf{Z}\gamma_1 \oplus \mathbf{Z}\gamma_2, \gamma_1 = \{|z| = 1\}, \gamma_2 = \mathbf{R}^-$ (if $\lambda \in \mathbf{R}^+$).

H_{dR}^1 is a $\mathbf{C}(\lambda)$ -vector space with a connexion, whose dual admits γ_1 and γ_2 as horizontal vectors (see also Bloch-Esnault). Therefore, the family of periods

$$\begin{aligned} \omega_1(\lambda) &= \int_{\gamma_1} \omega = \int_{|z|=1} e^{z+\lambda/z} \frac{dz}{z} \\ &= 2i\pi \sum_{n \geq 0} \frac{\lambda^n}{(n!)^2} = 2i\pi J_0(\lambda) \end{aligned}$$

is a solution of a 2nd order differential equation (Bessel!), whose derivative $J_1(\lambda)$ is essentially given by $\eta_1(\lambda) = \int_{\gamma_1} \eta$. The second period

$$\omega_2(\lambda) = \int_{\gamma_2} \omega = \int_{-\infty}^0 e^{z+\lambda/z} \frac{dz}{z}$$

(essentially $Y_0(\lambda)$) has a logarithmic singularity at $\lambda = 0$.

Now, Siegel's theorem on the algebraic independence of $J_0(\lambda)$ and $J'_0(\lambda)$ implies : *for any $\lambda \in \overline{\mathbf{Q}}, \lambda \neq 0$, the periods $\omega_1(\lambda)$ and $\omega_2(\lambda)$ are linearly independent over $\overline{\mathbf{Q}}$.* In particular, the slope $\tau(\lambda) = \frac{\omega_1(\lambda)}{\omega_2(\lambda)}$ never vanishes.

Questions :

i) what can be said of the "quasi-periods" $\eta_i(\lambda)$, which involve *E- and G-functions*? (NB : there is a Legendre relation, since the wronskian of the Bessel equation is rational).

ii) what is the analogue of Grothendieck's conjecture for these irregular periods?

Many other irregular periods can be studied, using Shidlovsky's theorem on hypergeometric equations. In a sense, we have a theorem waiting for a ... conjecture!