

The role of complex conjugation in transcendental number theory

Michel Waldschmidt

Institut de Mathématiques de Jussieu + CIMPA
<http://www.math.jussieu.fr/~miw/>

December 17, 2005

DION 2005

TIFR

December 17, 2005

Hermite–Lindemann’s Theorem as a consequence of
Gel’fond–Schneider’s Theorem

The Six Exponentials Theorem and the Four Exponentials
Conjecture

The strong Six Exponentials Theorem and the strong Four
Exponentials Conjecture

Recent results

Product of logarithms of algebraic numbers

References

Hermite–Lindemann’s Theorem

- ▶ *Let α be a nonzero algebraic number and let $\log \alpha$ be any nonzero logarithm of α . Then $\log \alpha$ is transcendental.*

Hermite–Lindemann’s Theorem

- ▶ *Let α be a nonzero algebraic number and let $\log \alpha$ be any nonzero logarithm of α . Then $\log \alpha$ is transcendental.*
- ▶ **Notations.** Denote by $\overline{\mathbb{Q}}$ the field of algebraic numbers and by \mathcal{L} the \mathbb{Q} -vector space of logarithms of algebraic numbers :

$$\mathcal{L} = \{\lambda \in \mathbb{C} ; e^\lambda \in \overline{\mathbb{Q}}^\times\} = \exp^{-1}(\overline{\mathbb{Q}}^\times) = \{\log \alpha ; \alpha \in \overline{\mathbb{Q}}^\times\}.$$

Hermite–Lindemann's Theorem

- ▶ *Let α be a nonzero algebraic number and let $\log \alpha$ be any nonzero logarithm of α . Then $\log \alpha$ is transcendental.*
- ▶ **Notations.** Denote by $\overline{\mathbb{Q}}$ the field of algebraic numbers and by \mathcal{L} the \mathbb{Q} -vector space of logarithms of algebraic numbers :

$$\mathcal{L} = \{\lambda \in \mathbb{C} ; e^\lambda \in \overline{\mathbb{Q}}^\times\} = \exp^{-1}(\overline{\mathbb{Q}}^\times) = \{\log \alpha ; \alpha \in \overline{\mathbb{Q}}^\times\}.$$

- ▶ **Alternative statement of Hermite–Lindemann's Theorem :**

$$\mathcal{L} \cap \overline{\mathbb{Q}} = \{0\}.$$

Hermite–Lindemann’s Theorem (continued)

- ▶ Another alternative statement of
Hermite–Lindemann’s Theorem : *Let β be a nonzero algebraic number. Then e^β is transcendental.*

Hermite–Lindemann’s Theorem (continued)

- ▶ **Another alternative statement of Hermite–Lindemann’s Theorem** : *Let β be a nonzero algebraic number. Then e^β is transcendental.*
- ▶ **Question (G. Diaz)** : *Let t be a non-zero real number and β a non-zero algebraic number. Is it true that $e^{t\beta}$ is transcendental?*

Hermite–Lindemann’s Theorem (continued)

- ▶ Another alternative statement of Hermite–Lindemann’s Theorem : *Let β be a nonzero algebraic number. Then e^β is transcendental.*
- ▶ Question (G. Diaz) : *Let t be a non-zero real number and β a non-zero algebraic number. Is it true that $e^{t\beta}$ is transcendental?*
- ▶ Answer (G. Diaz) : **No!**

Hermite–Lindemann's Theorem (continued)

- ▶ Another alternative statement of Hermite–Lindemann's Theorem : *Let β be a nonzero algebraic number. Then e^β is transcendental.*
- ▶ Question (G. Diaz) : *Let t be a non-zero real number and β a non-zero algebraic number. Is it true that $e^{t\beta}$ is transcendental?*
- ▶ Answer (G. Diaz) : **No!**
- ▶ First example : assume $\beta \in \mathbb{R}$. Take $t = (\log 2)/\beta$.

Hermite–Lindemann's Theorem (continued)

- ▶ Another alternative statement of Hermite–Lindemann's Theorem : *Let β be a nonzero algebraic number. Then e^β is transcendental.*
- ▶ Question (G. Diaz) : *Let t be a non-zero real number and β a non-zero algebraic number. Is it true that $e^{t\beta}$ is transcendental?*
- ▶ Answer (G. Diaz) : **No!**
- ▶ First example : assume $\beta \in \mathbb{R}$. Take $t = (\log 2)/\beta$.
- ▶ Second example : assume $\beta \in i\mathbb{R}$. Take $t = i\pi/\beta$.

Diaz’ Theorem

- ▶ *Let $\beta \in \overline{\mathbb{Q}}$ and $t \in \mathbb{R}^\times$. Assume $\beta \notin \mathbb{R} \cup i\mathbb{R}$. Then $e^{t\beta}$ is transcendental.*

Diaz' Theorem

- ▶ Let $\beta \in \overline{\mathbb{Q}}$ and $t \in \mathbb{R}^\times$. Assume $\beta \notin \mathbb{R} \cup i\mathbb{R}$. Then $e^{t\beta}$ is transcendental.
- ▶ **Equivalently** : for $\lambda \in \mathcal{L}$ with $\lambda \notin \mathbb{R} \cup i\mathbb{R}$,

$$\mathbb{R}\lambda \cap \overline{\mathbb{Q}} = \{0\}.$$

Diaz' Theorem

- ▶ Let $\beta \in \overline{\mathbb{Q}}$ and $t \in \mathbb{R}^\times$. Assume $\beta \notin \mathbb{R} \cup i\mathbb{R}$. Then $e^{t\beta}$ is transcendental.
- ▶ **Equivalently** : for $\lambda \in \mathcal{L}$ with $\lambda \notin \mathbb{R} \cup i\mathbb{R}$,

$$\mathbb{R}\lambda \cap \overline{\mathbb{Q}} = \{0\}.$$

- ▶ **Proof.** Set $\alpha = e^{t\beta}$. The complex conjugate $\bar{\alpha}$ of α is $e^{t\bar{\beta}} = \alpha^{\bar{\beta}/\beta}$. Since $\beta \notin \mathbb{R} \cup i\mathbb{R}$, the algebraic number $\bar{\beta}/\beta$ is not real (its modulus is 1 and it is not ± 1), hence not rational. **Gel'fond-Schneider's Theorem** implies that α and $\bar{\alpha}$ cannot be both algebraic. Hence they are both transcendental. □

Gel’fond-Schneider implies Hermite-Lindemann (almost)

- ▶ Gel’fond-Schneider’s Theorem implies : *there exists*
 $\beta_0 \in \mathbb{R} \cup i\mathbb{R}$ such that

$$\{\beta \in \overline{\mathbb{Q}} ; e^\beta \in \overline{\mathbb{Q}}\} = \mathbb{Q}\beta_0.$$

Gel’fond-Schneider implies Hermite-Lindemann (almost)

- ▶ Gel’fond-Schneider’s Theorem implies : *there exists*
 $\beta_0 \in \mathbb{R} \cup i\mathbb{R}$ such that

$$\{\beta \in \overline{\mathbb{Q}} ; e^\beta \in \overline{\mathbb{Q}}\} = \mathbb{Q}\beta_0.$$

- ▶ **Remark.** Hermite–Lindemann’s Theorem tells us that
in fact $\beta_0 = 0$.

Gel’fond-Schneider implies Hermite-Lindemann (almost)

- ▶ Gel’fond-Schneider’s Theorem implies : *there exists*
 $\beta_0 \in \mathbb{R} \cup i\mathbb{R}$ such that

$$\{\beta \in \overline{\mathbb{Q}} ; e^\beta \in \overline{\mathbb{Q}}\} = \mathbb{Q}\beta_0.$$

- ▶ **Remark.** Hermite–Lindemann’s Theorem tells us that in fact $\beta_0 = 0$.
- ▶ **Proof.** From Gel’fond-Schneider’s Theorem one deduces that the \mathbb{Q} -vector-space $\{\beta \in \overline{\mathbb{Q}} ; e^\beta \in \overline{\mathbb{Q}}\}$ has dimension ≤ 1 and is contained in $\mathbb{R} \cup i\mathbb{R}$. \square

Gel’fond-Schneider implies Hermite-Lindemann (almost)

- ▶ Gel’fond-Schneider’s Theorem implies : *there exists*
 $\beta_0 \in \mathbb{R} \cup i\mathbb{R}$ such that

$$\{\beta \in \overline{\mathbb{Q}} ; e^\beta \in \overline{\mathbb{Q}}\} = \mathbb{Q}\beta_0.$$

- ▶ **Remark.** Hermite–Lindemann’s Theorem tells us that
in fact $\beta_0 = 0$.
- ▶ **Proof.** From Gel’fond-Schneider’s Theorem one
deduces that the \mathbb{Q} -vector-space $\{\beta \in \overline{\mathbb{Q}} ; e^\beta \in \overline{\mathbb{Q}}\}$ has
dimension ≤ 1 and is contained in $\mathbb{R} \cup i\mathbb{R}$. \square
- ▶ Schneider’s method : proof without derivatives.

Reference



G. DIAZ – « Utilisation de la conjugaison complexe dans l’étude de la transcendance de valeurs de la fonction exponentielle », *J. Théor. Nombres Bordeaux* **16** (2004), p. 535–553.

The Six Exponentials Theorem

- ▶ Selberg, Siegel, Lang, Ramachandra.

The Six Exponentials Theorem

- ▶ Selberg, Siegel, Lang, Ramachandra.
- ▶ **Theorem** : *If x_1, x_2 are \mathbb{Q} -linearly independent complex numbers and y_1, y_2, y_3 are \mathbb{Q} -linearly independent complex numbers, then one at least of the six numbers*

$$e^{x_1 y_1}, e^{x_1 y_2}, e^{x_1 y_3}, e^{x_2 y_1}, e^{x_2 y_2}, e^{x_2 y_3}$$

is transcendental.

The Six Exponentials Theorem

References :



S. LANG – *Introduction to transcendental numbers*,
Addison-Wesley Publishing Co., Reading,
Mass.-London-Don Mills, Ont., 1966.



K. RAMACHANDRA – « Contributions to the theory of
transcendental numbers. I, II », *Acta Arith.* 14
(1967/68), 65–72; *ibid.* 14 (1967/1968), p. 73–88.

Corollary

- ▶ **Example :** Take $x_1 = 1$, $x_2 = \pi$, $y_1 = \log 2$, $y_2 = \pi \log 2$, $y_3 = \pi^2 \log 2$, the six exponentials are respectively

$$2, 2^\pi, 2^{\pi^2}, 2^\pi, 2^{\pi^2}, 2^{\pi^3},$$

hence one at least of the three numbers

$$2^\pi, 2^{\pi^2}, 2^{\pi^3}$$

is transcendental

Corollary

- ▶ **Example** : Take $x_1 = 1$, $x_2 = \pi$, $y_1 = \log 2$, $y_2 = \pi \log 2$, $y_3 = \pi^2 \log 2$, the six exponentials are respectively

$$2, 2^\pi, 2^{\pi^2}, 2^\pi, 2^{\pi^2}, 2^{\pi^3},$$

hence one at least of the three numbers

$$2^\pi, 2^{\pi^2}, 2^{\pi^3}$$

is transcendental



- ▶ **Shorey** : lower bound for

$$|2^\pi - \alpha_1| + |2^{\pi^2} - \alpha_2| + |2^{\pi^3} - \alpha_3|$$

for algebraic $\alpha_1, \alpha_2, \alpha_3$. The estimate depends on the heights and degrees of these algebraic numbers.




Relevant references

References :

-  T. N. SHOREY – « On a theorem of Ramachandra », *Acta Arith.* **20** (1972), p. 215–221.
-  T. N. SHOREY – « On the sum $\sum_{k=1}^3 2^{\pi^k} - \alpha_k$, α_k algebraic numbers », *J. Number Theory* **6** (1974), p. 248–260.

S. Srinivasan contributions

Further references :

-  S. SRINIVASAN – « On algebraic approximations to 2^{π^k} ($k = 1, 2, 3, \dots$) », *Indian J. Pure Appl. Math.* **5** (1974), no. 6, p. 513–523.
-  S. SRINIVASAN – « On algebraic approximations to 2^{π^k} ($k = 1, 2, 3, \dots$). II », *J. Indian Math. Soc. (N.S.)* **43** (1979), no. 1-4, p. 53–60 (1980).
-  K. RAMACHANDRA & S. SRINIVASAN – « A note to a paper : “Contributions to the theory of transcendental numbers. I, II” by Ramachandra on transcendental numbers », *Hardy-Ramanujan J.* **6** (1983), p. 37–44.

Conjectures

- ▶ **Remark** : It is unknown whether one of the two numbers

$$2^\pi, 2^{\pi^2}$$

is transcendental. One **conjectures** (Schanuel) that each of the three numbers $2^\pi, 2^{\pi^2}, 2^{\pi^3}$ is transcendental

Conjectures

- ▶ **Remark** : It is unknown whether one of the two numbers

$$2^\pi, 2^{\pi^2}$$

is transcendental. One **conjectures** (Schanuel) that each of the three numbers $2^\pi, 2^{\pi^2}, 2^{\pi^3}$ is transcendental

- ▶ and that the numbers

$$\pi, \log 2, 2^\pi, 2^{\pi^2}, 2^{\pi^3}$$

are algebraically independent.

The Four Exponentials Conjecture

- ▶ Selberg, Siegel, Schneider, Lang, Ramachandra.

The Four Exponentials Conjecture

- ▶ Selberg, Siegel, Schneider, Lang, Ramachandra.
- ▶ **Conjecture.** *If x_1, x_2 are \mathbb{Q} -linearly independent complex numbers and y_1, y_2 are \mathbb{Q} -linearly independent complex numbers, then one at least of the four numbers*

$$e^{x_1 y_1}, e^{x_1 y_2}, e^{x_2 y_1}, e^{x_2 y_2}$$

is transcendental.

Ramachandra's trick

- ▶ **Remark** : Let x and y be two **real** numbers.
The following properties are equivalent :
 - (i) one at least of the two numbers x , y is transcendental.
 - (ii) the **complex** number $x + iy$ is transcendental.

Ramachandra’s trick

- ▶ **Remark** : Let x and y be two **real** numbers.
The following properties are equivalent :
 - (i) one at least of the two numbers x, y is transcendental.
 - (ii) the **complex** number $x + iy$ is transcendental.
- ▶ **Example** : (H.W. Lenstra) if γ is Euler’s constant, then the number $\gamma + ie^\gamma$ is transcendental.

Ramachandra's trick

- ▶ **Remark** : Let x and y be two **real** numbers.
The following properties are equivalent :
 - (i) one at least of the two numbers x, y is transcendental.
 - (ii) the **complex** number $x + iy$ is transcendental.
- ▶ **Example** : (H.W. Lenstra) if γ is Euler's constant, then the number $\gamma + ie^\gamma$ is transcendental.
- ▶ **Proof** : check $\gamma \neq 0$ and use Hermite–Lindemann's Theorem. □

Ramachandra’s trick

Other example.

- ▶ Let x_1, x_2 be two elements in $\mathbb{R} \cup i\mathbb{R}$ which are \mathbb{Q} -linearly independent. Let y_1, y_2 be two complex numbers. Assume that the three numbers $y_1, y_2, \overline{y_2}$ are \mathbb{Q} -linearly independent. Then one at least of the four numbers

$$e^{x_1 y_1}, e^{x_1 y_2}, e^{x_2 y_1}, e^{x_2 y_2}$$

is transcendental.

Ramachandra’s trick

Other example.

- ▶ Let x_1, x_2 be two elements in $\mathbb{R} \cup i\mathbb{R}$ which are \mathbb{Q} -linearly independent. Let y_1, y_2 be two complex numbers. Assume that the three numbers $y_1, y_2, \overline{y_2}$ are \mathbb{Q} -linearly independent. Then one at least of the four numbers

$$e^{x_1 y_1}, e^{x_1 y_2}, e^{x_2 y_1}, e^{x_2 y_2}$$

is transcendental.

- ▶ **Proof** : Set $y_3 = \overline{y_2}$. Then $e^{x_j y_3} = \overline{e^{\pm x_j y_2}}$ for $j = 1, 2$ and $\overline{\mathbb{Q}}$ is stable under complex conjugation. □

Logarithms of algebraic numbers

Rank of matrices. An alternate form of the Six Exponentials Theorem (resp. the Four Exponentials Conjecture) is the fact that a 2×3 (resp. 2×2) matrix with entries in \mathcal{L}

$$\begin{pmatrix} \lambda_{11} & \lambda_{12} & \lambda_{13} \\ \lambda_{21} & \lambda_{22} & \lambda_{23} \end{pmatrix} \quad (\text{resp.} \quad \begin{pmatrix} \lambda_{11} & \lambda_{12} \\ \lambda_{21} & \lambda_{22} \end{pmatrix} \quad),$$

the rows of which are linearly independent over \mathbb{Q} and the columns of which are also linearly independent over \mathbb{Q} , has maximal rank 2.

A lemma on the rank of matrices

Remark. A $d \times \ell$ matrix M has rank ≤ 1 if and only if there exist x_1, \dots, x_d and y_1, \dots, y_ℓ such that

$$M = \begin{pmatrix} x_1 y_1 & x_1 y_2 & \dots & x_1 y_\ell \\ x_2 y_1 & x_2 y_2 & \dots & x_2 y_\ell \\ \vdots & \vdots & \ddots & \vdots \\ x_d y_1 & x_d y_2 & \dots & x_d y_\ell \end{pmatrix}.$$

Linear combinations of logarithms of algebraic numbers

Denote by $\tilde{\mathcal{L}}$ the $\overline{\mathbb{Q}}$ -vector space spanned by 1 and \mathcal{L} :
hence $\tilde{\mathcal{L}}$ is the set of linear combinations with algebraic coefficients of logarithms of algebraic numbers :

$$\tilde{\mathcal{L}} = \{\beta_0 + \beta_1 \lambda_1 + \cdots + \beta_n \lambda_n ; n \geq 0, \beta_i \in \overline{\mathbb{Q}}, \lambda_i \in \mathcal{L}\}.$$

The strong Six Exponentials Theorem

Theorem (D.Roy). *If x_1, x_2 are $\overline{\mathbb{Q}}$ -linearly independent complex numbers and y_1, y_2, y_3 are $\overline{\mathbb{Q}}$ -linearly independent complex numbers, then one at least of the six numbers*

$$x_1y_1, x_1y_2, x_1y_3, x_2y_1, x_2y_2, x_2y_3$$

is not in $\tilde{\mathcal{L}}$.

The strong Four Exponentials Conjecture

Conjecture. *If x_1, x_2 are $\overline{\mathbb{Q}}$ -linearly independent complex numbers and y_1, y_2 are $\overline{\mathbb{Q}}$ -linearly independent complex numbers, then one at least of the four numbers*

$$x_1y_1, x_1y_2, x_2y_1, x_2y_2$$

is not in $\tilde{\mathcal{L}}$.

Lower bound for the rank of matrices

- ▶ **Rank of matrices.** An alternate form of the strong Six Exponentials Theorem (resp. the strong Four Exponentials Conjecture) is the fact that a 2×3 (resp. 2×2) matrix with entries in $\tilde{\mathcal{L}}$

$$\begin{pmatrix} \Lambda_{11} & \Lambda_{12} & \Lambda_{13} \\ \Lambda_{21} & \Lambda_{22} & \Lambda_{23} \end{pmatrix} \quad (\text{resp.} \quad \begin{pmatrix} \Lambda_{11} & \Lambda_{12} \\ \Lambda_{21} & \Lambda_{22} \end{pmatrix} \quad),$$

the rows of which are linearly independent over $\overline{\mathbb{Q}}$ and the columns of which are also linearly independent over $\overline{\mathbb{Q}}$, has maximal rank 2.

Lower bound for the rank of matrices

- ▶ **Rank of matrices.** An alternate form of the strong Six Exponentials Theorem (resp. the strong Four Exponentials Conjecture) is the fact that a 2×3 (resp. 2×2) matrix with entries in $\tilde{\mathcal{L}}$



$$\begin{pmatrix} \Lambda_{11} & \Lambda_{12} & \Lambda_{13} \\ \Lambda_{21} & \Lambda_{22} & \Lambda_{23} \end{pmatrix} \quad (\text{resp.} \quad \begin{pmatrix} \Lambda_{11} & \Lambda_{12} \\ \Lambda_{21} & \Lambda_{22} \end{pmatrix} \quad),$$

the rows of which are linearly independent over $\overline{\mathbb{Q}}$ and the columns of which are also linearly independent over $\overline{\mathbb{Q}}$, has maximal rank 2.

- ▶ **Remark :** Under suitable conditions one can show that a $d \times \ell$ matrix with entries in $\tilde{\mathcal{L}}$ has rank $\geq d\ell/(d + \ell)$. This is a consequence of the *Linear Subgroup Theorem*.

The strong Six Exponentials Theorem

References :

-  D. ROY – « Matrices whose coefficients are linear forms in logarithms », *J. Number Theory* **41** (1992), no. 1, p. 22–47.
-  M. WALDSCHMIDT – *Diophantine approximation on linear algebraic groups*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. **326**, Springer-Verlag, Berlin, 2000.

Alternate form of the strong Four Exponentials Conjecture

- ▶ **Conjecture.** *Let $\Lambda_1, \Lambda_2, \Lambda_3$ be nonzero elements in $\tilde{\mathcal{L}}$. Assume the numbers Λ_2/Λ_1 and Λ_3/Λ_1 are both transcendental. Then the number $\Lambda_2\Lambda_3/\Lambda_1$ is not in $\tilde{\mathcal{L}}$.*

Alternate form of the strong Four Exponentials Conjecture

- ▶ **Conjecture.** *Let $\Lambda_1, \Lambda_2, \Lambda_3$ be nonzero elements in $\tilde{\mathcal{L}}$. Assume the numbers Λ_2/Λ_1 and Λ_3/Λ_1 are both transcendental. Then the number $\Lambda_2\Lambda_3/\Lambda_1$ is not in $\tilde{\mathcal{L}}$.*
- ▶ **Equivalence between both statements :** the matrix

$$\begin{pmatrix} \Lambda_1 & \Lambda_2 \\ \Lambda_3 & \Lambda_2\Lambda_3/\Lambda_1 \end{pmatrix}$$

has rank 1. □

Consequences of the strong Four Exponentials Conjecture

Assume the strong Four Exponentials Conjecture.

- ▶ If Λ is in $\tilde{\mathcal{L}} \setminus \overline{\mathbb{Q}}$ then the quotient $1/\Lambda$ is not in $\tilde{\mathcal{L}}$.

Consequences of the strong Four Exponentials Conjecture

Assume the strong Four Exponentials Conjecture.

- ▶ If Λ is in $\tilde{\mathcal{L}} \setminus \overline{\mathbb{Q}}$ then the quotient $1/\Lambda$ is not in $\tilde{\mathcal{L}}$.
- ▶ If Λ_1 and Λ_2 are in $\tilde{\mathcal{L}} \setminus \overline{\mathbb{Q}}$, then the product $\Lambda_1\Lambda_2$ is not in $\tilde{\mathcal{L}}$.

Consequences of the strong Four Exponentials Conjecture

Assume the strong Four Exponentials Conjecture.

- ▶ If Λ is in $\tilde{\mathcal{L}} \setminus \overline{\mathbb{Q}}$ then the quotient $1/\Lambda$ is not in $\tilde{\mathcal{L}}$.
- ▶ If Λ_1 and Λ_2 are in $\tilde{\mathcal{L}} \setminus \overline{\mathbb{Q}}$, then the product $\Lambda_1\Lambda_2$ is not in $\tilde{\mathcal{L}}$.
- ▶ If Λ_1 and Λ_2 are in $\tilde{\mathcal{L}}$ with Λ_1 and Λ_2/Λ_1 transcendental, then this quotient Λ_2/Λ_1 is not in $\tilde{\mathcal{L}}$.

Example where the strong Four Exponentials Conjecture is true

- ▶ **Theorem** (G. Diaz). *Let x_1 and x_2 be two elements of $\mathbb{R} \cup i\mathbb{R}$ which are $\overline{\mathbb{Q}}$ -linearly independent. Let y_1, y_2 be two complex numbers such that the three numbers $y_1, y_2, \overline{y_2}$ are $\overline{\mathbb{Q}}$ -linearly independent. Then one at least of the four numbers*

$$x_1y_1, x_1y_2, x_2y_1, x_2y_2$$

is not in $\tilde{\mathcal{L}}$.

Example where the strong Four Exponentials Conjecture is true

- ▶ **Theorem** (G. Diaz). *Let x_1 and x_2 be two elements of $\mathbb{R} \cup i\mathbb{R}$ which are $\overline{\mathbb{Q}}$ -linearly independent. Let y_1, y_2 be two complex numbers such that the three numbers $y_1, y_2, \overline{y_2}$ are $\overline{\mathbb{Q}}$ -linearly independent. Then one at least of the four numbers*

$$x_1 y_1, x_1 y_2, x_2 y_1, x_2 y_2$$

is not in $\tilde{\mathcal{L}}$.

- ▶ **Proof** : Set $y_3 = \overline{y_2}$. Then $e^{x_j y_3} = \overline{e^{\pm x_j y_2}}$ for $j = 1, 2$ and $\tilde{\mathcal{L}}$ is stable under complex conjugation. □

Example where the strong Four Exponentials Conjecture is true

- ▶ **Corollary** of Diaz’ Theorem. *Let $\Lambda_1, \Lambda_2, \Lambda_3$ be three elements in $\tilde{\mathcal{L}}$. Assume that the three numbers $\Lambda_1, \Lambda_2, \overline{\Lambda_2}$ are linearly independent over $\overline{\mathbb{Q}}$. Further assume $\Lambda_3/\Lambda_1 \in (\mathbb{R} \cup i\mathbb{R}) \setminus \overline{\mathbb{Q}}$. Then*

$$\Lambda_2\Lambda_3/\Lambda_1 \notin \overline{\mathbb{Q}}.$$

Example where the strong Four Exponentials Conjecture is true

- ▶ **Corollary** of Diaz' Theorem. *Let $\Lambda_1, \Lambda_2, \Lambda_3$ be three elements in $\tilde{\mathcal{L}}$. Assume that the three numbers $\Lambda_1, \Lambda_2, \overline{\Lambda_2}$ are linearly independent over $\overline{\mathbb{Q}}$. Further assume $\Lambda_3/\Lambda_1 \in (\mathbb{R} \cup i\mathbb{R}) \setminus \overline{\mathbb{Q}}$. Then*

$$\Lambda_2\Lambda_3/\Lambda_1 \notin \overline{\mathbb{Q}}.$$

- ▶ **Proof** : set $x_1 = 1, x_2 = \Lambda_3/\Lambda_1, y_1 = \Lambda_1, y_2 = \Lambda_2$. □

Example where the strong Four Exponentials Conjecture is true

Consequence : one deduces examples where one can actually prove that numbers like

$$1/\Lambda, \quad \Lambda_1\Lambda_2, \quad \Lambda_2/\Lambda_1$$

(with $\Lambda, \Lambda_1, \Lambda_2$ in $\tilde{\mathcal{L}}$) are not in $\tilde{\mathcal{L}}$.

Transcendence of e^{π^2}

- ▶ **Open problem** : *is the number e^{π^2} transcendental?*

Transcendence of e^{π^2}

- ▶ **Open problem** : *is the number e^{π^2} transcendental?*
- ▶ **More generally** : *for $\lambda \in \mathcal{L} \setminus \{0\}$, is it true that $\lambda\bar{\lambda} \notin \mathcal{L}$?*

Transcendence of e^{π^2}

- ▶ **Open problem** : *is the number e^{π^2} transcendental?*
- ▶ **More generally** : *for $\lambda \in \mathcal{L} \setminus \{0\}$, is it true that $\lambda\bar{\lambda} \notin \mathcal{L}$?*
- ▶ **More generally** : *for λ_1 and λ_2 in $\mathcal{L} \setminus \{0\}$, is it true that $\lambda_1\lambda_2 \notin \mathcal{L}$?*

Transcendence of e^{π^2}

- ▶ **Open problem** : *is the number e^{π^2} transcendental?*
- ▶ **More generally** : *for $\lambda \in \mathcal{L} \setminus \{0\}$, is it true that $\lambda\bar{\lambda} \notin \mathcal{L}$?*
- ▶ **More generally** : *for λ_1 and λ_2 in $\mathcal{L} \setminus \{0\}$, is it true that $\lambda_1\lambda_2 \notin \mathcal{L}$?*
- ▶ *For λ_1 and λ_2 in $\mathcal{L} \setminus \{0\}$, is it true that $\lambda_1\lambda_2 \notin \tilde{\mathcal{L}}$?*

Product of logarithms of algebraic numbers

- ▶ **Theorem (Diaz).** *Let λ_1 and λ_2 be in $\mathcal{L} \setminus \{0\}$. Assume $\lambda_1 \in \mathbb{R} \cup i\mathbb{R}$ and $\lambda_2 \notin \mathbb{R} \cup i\mathbb{R}$. Then $\lambda_1 \lambda_2 \notin \tilde{\mathcal{L}}$.*

Product of logarithms of algebraic numbers

- ▶ **Theorem (Diaz).** *Let λ_1 and λ_2 be in $\mathcal{L} \setminus \{0\}$. Assume $\lambda_1 \in \mathbb{R} \cup i\mathbb{R}$ and $\lambda_2 \notin \mathbb{R} \cup i\mathbb{R}$. Then $\lambda_1 \lambda_2 \notin \tilde{\mathcal{L}}$.*
- ▶ **Proof.** Apply the strong Six Exponentials Theorem to

$$\begin{pmatrix} 1 & \lambda_2 & \overline{\lambda_2} \\ \lambda_1 & \Lambda & \overline{\Lambda} \end{pmatrix}$$

with $\Lambda \in \tilde{\mathcal{L}}$.



Product of logarithms of algebraic numbers

- ▶ **Theorem (Diaz).** Let λ_1 and λ_2 be in $\mathcal{L} \setminus \{0\}$. Assume $\lambda_1 \in \mathbb{R} \cup i\mathbb{R}$ and $\lambda_2 \notin \mathbb{R} \cup i\mathbb{R}$. Then $\lambda_1 \lambda_2 \notin \tilde{\mathcal{L}}$.
- ▶ **Proof.** Apply the strong Six Exponentials Theorem to

$$\begin{pmatrix} 1 & \lambda_2 & \overline{\lambda_2} \\ \lambda_1 & \Lambda & \overline{\Lambda} \end{pmatrix}$$

with $\Lambda \in \tilde{\mathcal{L}}$. □

- ▶ **Diaz’ Conjecture.** Let $u \in \mathbb{C}^\times$. Assume $|u|$ is algebraic. Then e^u is transcendental.

Recent results





G. DIAZ – « Utilisation de la conjugaison complexe dans l’étude de la transcendance de valeurs de la fonction exponentielle », *J. Théor. Nombres Bordeaux* **16** (2004), p. 535–553.



G. DIAZ – « Produits et quotients de combinaisons linéaires de logarithmes de nombres algébriques : conjectures et résultats partiels », Submitted (2005), 19 p.

Recent results

-  M. WALDSCHMIDT – « Transcendence results related to the six exponentials theorem », *Hindustan Book Agency* (2005), p. 338–355.
Appendix by H. Shiga : Periods of the Kummer surface, p. 356–358.
-  M. WALDSCHMIDT – « Further variations on the Six Exponentials Theorem », *The Hardy-Ramanujan Journal*, vol. **28**, to appear on **december 22**, 2005.

Further result

Let M be a 2×3 matrix with entries in $\tilde{\mathcal{L}}$:

$$M = \begin{pmatrix} \Lambda_{11} & \Lambda_{12} & \Lambda_{13} \\ \Lambda_{21} & \Lambda_{22} & \Lambda_{23} \end{pmatrix}.$$

Assume that the five rows of the matrix

$$\begin{pmatrix} M \\ I_3 \end{pmatrix} = \begin{pmatrix} \Lambda_{11} & \Lambda_{12} & \Lambda_{13} \\ \Lambda_{21} & \Lambda_{22} & \Lambda_{23} \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

are linearly independent over $\overline{\mathbb{Q}}$ and that the five columns of the matrix

$$(I_2, M) = \begin{pmatrix} 1 & 0 & \Lambda_{11} & \Lambda_{12} & \Lambda_{13} \\ 0 & 1 & \Lambda_{21} & \Lambda_{22} & \Lambda_{23} \end{pmatrix}$$

are linearly independent over $\overline{\mathbb{Q}}$.

Further result

Then one at least of the three numbers

$$\Delta_1 = \begin{vmatrix} \Lambda_{12} & \Lambda_{13} \\ \Lambda_{22} & \Lambda_{23} \end{vmatrix}, \quad \Delta_2 = \begin{vmatrix} \Lambda_{13} & \Lambda_{11} \\ \Lambda_{23} & \Lambda_{21} \end{vmatrix}, \quad \Delta_3 = \begin{vmatrix} \Lambda_{11} & \Lambda_{12} \\ \Lambda_{21} & \Lambda_{22} \end{vmatrix}$$

is not in $\tilde{\mathcal{L}}$.

Higher rank : an example

Let $M = (\Lambda_{ij})_{1 \leq i \leq m; 1 \leq j \leq \ell}$ be a $m \times \ell$ matrix with entries in $\tilde{\mathcal{L}}$. Denote by I_m the identity $m \times m$ matrix and assume that the $m + \ell$ column vectors of the matrix (I_m, M) are linearly independent over $\overline{\mathbb{Q}}$. Let $\Lambda_1, \dots, \Lambda_m$ be elements of $\tilde{\mathcal{L}}$. Assume that the numbers $1, \Lambda_1, \dots, \Lambda_m$ are $\overline{\mathbb{Q}}$ -linearly independent. Assume further $\ell > m^2$. Then one at least of the ℓ numbers

$$\Lambda_1 \Lambda_{1j} + \dots + \Lambda_m \Lambda_{mj} \quad (j = 1, \dots, \ell)$$

is not in $\tilde{\mathcal{L}}$.

Janam din diyan wadhayian, Tarlok !

<http://www.math.jussieu.fr/~miw/>