



The Institute of Mathematical Sciences  
Mathematics Colloquium

December 9, 2009

<http://www.imsc.res.in/>

## **Recent Diophantine results on zeta values: a survey**

*Michel Waldschmidt*

Institut de Mathématiques de Jussieu & Paris VI

<http://www.math.jussieu.fr/~miw/>

# Abstract

After the proof by R. Apéry of the irrationality of  $\zeta(3)$  in 1976, a number of papers have been devoted to the study of Diophantine properties of values of the Riemann zeta function at positive integers.

A survey has been written by S. Fischler for the Bourbaki Seminar in November 2002.

We review more recent results including contributions by S. Fischler, M. Hata, C. Krattenthaler, R. Marcovecchio, R. Murty, G. Rhin, T. Rivoal, C. Viola, W. Zudilin.

We plan also to say a few words on the analog of this theory in finite characteristic, with works of W.D. Brownawell, M. Papanikolas, D. Thakur, Chieh-Yu Chang, Jing Yu and others.

# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .

# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .

# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .

# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .

# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .

# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .



# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .


# Zeta

- Riemann zeta values
- Multizeta values
- Weierstraß zeta function
- Fibonacci zeta values
- Hurwitz zeta function
- Carlitz zeta values
- (Other zeta functions : Dedekind, Hasse-Weil, Lerch, Selberg, Witten, Milnor, dynamical systems. . .)
- $L$ - functions. . .

# Riemann zeta function



$$\begin{aligned}\zeta(s) &= \sum_{n \geq 1} \frac{1}{n^s} \\ &= \prod_p \frac{1}{1 - p^{-s}}\end{aligned}$$



Euler :  $s \in \mathbf{R}$ .

Riemann :  $s \in \mathbf{C}$ .

# Special values of Riemann zeta function

Leonard Euler (1739)

$\zeta(s)$  for  $s \in \mathbf{Z}$



$$\zeta(2) = \frac{\pi^2}{6},$$

$$\zeta(4) = \frac{\pi^4}{90},$$

$$\zeta(6) = \frac{\pi^6}{945},$$

$$\zeta(8) = \frac{\pi^8}{9450}.$$

$$\pi^{-2k} \zeta(2k) \in \mathbf{Q} \quad \text{for } k \geq 1$$

# Bernoulli numbers

Jacques Bernoulli  
(1654–1705),



Bernoulli numbers :

$$B_2 = 1/6$$

$$B_4 = -1/30$$

$$B_6 = 1/42$$

$$B_8 = -1/30$$


$$B_{10} = 5/66$$

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} B_n \frac{z^n}{n!},$$

$$\zeta(2k) = (-1)^{k-1} 2^{2k-1} \frac{B_{2k}}{(2k)!} \pi^{2k} \quad (k \geq 1).$$

# Transcendence of even zeta values

- F. Lindemann :  $\pi$  is transcendental, hence  $\zeta(2k)$  also for  $k \geq 1$ .




Theorem (Hermite–Lindemann).

*For any non-zero complex number  $z$ , one at least of the two numbers  $z$  and  $e^z$  is transcendental.*

*Corollaries.* Transcendence of  $\log \alpha$  and of  $e^\beta$  for  $\alpha$  and  $\beta$  non-zero algebraic complex numbers, provided  $\log \alpha \neq 0$ .

# Transcendence of even zeta values

- F. Lindemann :  $\pi$  is transcendental, hence  $\zeta(2k)$  also for  $k \geq 1$ .




Theorem (Hermite–Lindemann).

*For any non-zero complex number  $z$ , one at least of the two numbers  $z$  and  $e^z$  is transcendental.*

*Corollaries.* Transcendence of  $\log \alpha$  and of  $e^\beta$  for  $\alpha$  and  $\beta$  non-zero algebraic complex numbers, provided  $\log \alpha \neq 0$ .

# Transcendence of even zeta values

- F. Lindemann :  $\pi$  is transcendental, hence  $\zeta(2k)$  also for  $k \geq 1$ .



Theorem (Hermite–Lindemann).

*For any non-zero complex number  $z$ , one at least of the two numbers  $z$  and  $e^z$  is transcendental.*

*Corollaries.* Transcendence of  $\log \alpha$  and of  $e^\beta$  for  $\alpha$  and  $\beta$  non-zero algebraic complex numbers, provided  $\log \alpha \neq 0$ .



# Diophantine question

Odd positive integers :  $\zeta(2k + 1)$ ,  $k \geq 1$ ?

Question. For  $n \geq 1$ , is the number

$$\frac{\zeta(2k + 1)}{\pi^{2k+1}}$$

rational?

Describe all algebraic relations among the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

**Conjecture.** There is no relation at all : the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

are algebraically independent.

In particular the numbers  $\zeta(2k + 1)$  and  $\zeta(2k + 1)/\pi^{2k+1}$  for  $k \geq 1$  are conjectured to be transcendental.

# Diophantine question

Odd positive integers :  $\zeta(2k + 1)$ ,  $k \geq 1$ ?

Question. For  $n \geq 1$ , is the number

$$\frac{\zeta(2k + 1)}{\pi^{2k+1}}$$

rational?

Describe all algebraic relations among the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

**Conjecture.** There is no relation at all : the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

are algebraically independent.

In particular the numbers  $\zeta(2k + 1)$  and  $\zeta(2k + 1)/\pi^{2k+1}$  for  $k \geq 1$  are conjectured to be transcendental.

# Diophantine question

Odd positive integers :  $\zeta(2k + 1)$ ,  $k \geq 1$ ?

Question. For  $n \geq 1$ , is the number

$$\frac{\zeta(2k + 1)}{\pi^{2k+1}}$$

rational?

Describe all algebraic relations among the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

Conjecture. There is no relation at all : the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

are algebraically independent.

In particular the numbers  $\zeta(2k + 1)$  and  $\zeta(2k + 1)/\pi^{2k+1}$  for  $k \geq 1$  are conjectured to be transcendental.

# Diophantine question

Odd positive integers :  $\zeta(2k + 1)$ ,  $k \geq 1$ ?

Question. For  $n \geq 1$ , is the number

$$\frac{\zeta(2k + 1)}{\pi^{2k+1}}$$

rational?

Describe all algebraic relations among the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

**Conjecture.** There is no relation at all : the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

are algebraically independent.

In particular the numbers  $\zeta(2k + 1)$  and  $\zeta(2k + 1)/\pi^{2k+1}$  for  $k \geq 1$  are conjectured to be transcendental.

# Diophantine question

Odd positive integers :  $\zeta(2k + 1)$ ,  $k \geq 1$ ?

Question. For  $n \geq 1$ , is the number

$$\frac{\zeta(2k + 1)}{\pi^{2k+1}}$$

rational?

Describe all algebraic relations among the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$


**Conjecture.** There is no relation at all : the numbers

$$\zeta(2), \quad \zeta(3), \quad \zeta(5), \quad \zeta(7), \dots$$

are algebraically independent.

In particular the numbers  $\zeta(2k + 1)$  and  $\zeta(2k + 1)/\pi^{2k+1}$  for  $k \geq 1$  are conjectured to be transcendental.

# Values of $\zeta$ at odd positive integers




- Apéry (1978) : *The number*

$$\zeta(3) = \sum_{n \geq 1} \frac{1}{n^3} = 1.202\,056\,903\,159\,594\,285\,399\,738\,161\,511 \dots$$

*is irrational.*

- Rivoal (2000) & Ball, Zudilin. . . *Infinitely many  $\zeta(2k+1)$  are irrational & lower bound for the dimension of the  $\mathbb{Q}$ -span.*

# Values of $\zeta$ at odd positive integers



- Apéry (1978) : *The number*

$$\zeta(3) = \sum_{n \geq 1} \frac{1}{n^3} = 1.202\,056\,903\,159\,594\,285\,399\,738\,161\,511 \dots$$

*is irrational.*

- Rivoal (2000) & Ball, Zudilin... *Infinitely many  $\zeta(2k + 1)$  are irrational & lower bound for the dimension of the  $\mathbb{Q}$ -span.*

# Infinitely many odd zeta are irrational

Tanguy Rivoal (2000)

*Let  $\epsilon > 0$ . For any sufficiently large odd integer  $a$ , the dimension of the  $\mathbb{Q}$ -vector space spanned by the numbers  $1, \zeta(3), \zeta(5), \dots, \zeta(a)$  is at least*


$$\frac{1 - \epsilon}{1 + \log 2} \log a.$$






# Wadim Zudilin

- *At least one of the four numbers  $\zeta(5)$ ,  $\zeta(7)$ ,  $\zeta(9)$ ,  $\zeta(11)$  is irrational.*
- *There exists an odd integer  $j$  in the range  $[5, 69]$  such that the three numbers  $1$ ,  $\zeta(3)$ ,  $\zeta(j)$  are  $\mathbb{Q}$ -linearly independent.*



# Wadim Zudilin

- *At least one of the four numbers  $\zeta(5)$ ,  $\zeta(7)$ ,  $\zeta(9)$ ,  $\zeta(11)$  is irrational.*
- *There exists an odd integer  $j$  in the range  $[5, 69]$  such that the three numbers  $1$ ,  $\zeta(3)$ ,  $\zeta(j)$  are  $\mathbb{Q}$ -linearly independent.*



# Zudilin's home page <http://wain.mi.ras.ru/zw/index.html>

## *References to works on zeta values by*

2000 M. Hata, T. Rivoal

2001 K. Ball and T. Rivoal, L.A. Gutnik, G. Rhin and C. Viola, T. Vasilyev, W. Zudilin

2002 T. Rivoal, V.N. Sorokin, W. Zudilin

2003 Yu.V. Nesterenko, T. Rivoal, J. Sondow, C. Viola, W. Zudilin

2004 **S. Fischler**, W. Zudilin

2005 F. Calegari, S. Zlobin

2006 M. Huttner, C. Krattenthaler, T. Rivoal and Zudilin

2007 C. Krattenthaler and T. Rivoal

2008 F. Beukers

2009 S. Fischler and W. Zudilin

*Last modified on September 19, 2009*

# Irrationality of zeta values

S. Fischler

*Irrationalité de valeurs de zêta,*  
(d'après Apéry, Rivoal, ...),  
Sém. Nicolas Bourbaki, 2002-2003,  
N° 910 (Novembre 2002).  
Astérisque **294** (2004), 27-62

<http://www.math.u-psud.fr/~fischler/publi.html>



# Christian Krattenthaler and Tanguy Rivoal

<http://www-fourier.ujf-grenoble.fr/~rivoal/articles.html>



C. Krattenthaler et T. Rivoal,  
*Hypergéométrie et fonction  
zêta de Riemann*, Mem.  
Amer. Math. Soc. **186**  
(2007), 93 p.



# Irrationality measures : the state of the art

$\vartheta \in \mathbf{R}$

$$\left| \vartheta - \frac{p}{q} \right| \geq \frac{1}{q^{\mu+\epsilon}}$$

$\mu(\vartheta) < +\infty \iff \vartheta$  is not a Liouville number

$\vartheta$	year	author	$\mu(\vartheta) <$
$\pi$	2008	V.Kh. Salikhov	7.6063085
$\zeta(2) = \pi^2/6$	1996	G. Rhin and C. Viola	5.441243
$\zeta(3)$	2001	G. Rhin and C. Viola	5.513891
$\log 2$	2008	R. Marcovecchio	3.57455391

# Irrationality measure for $\pi$ : history

K. Mahler 1953 :  $\pi$  is not a Liouville number and  $\mu(\pi) \leq 30$

M. Mignotte 1974 :  $\mu(\pi) \leq 20$

G.V. Chudnovsky 1984 :  $\mu(\pi) \leq 14.5$

M. Hata 1992 :  $\mu(\pi) \leq 8.0161$

V.Kh. Salikhov 2008 :  $\mu(\pi) \leq 7.6063$

A bound  $\mu(\vartheta^2) \leq \kappa$  for some  $\vartheta \in \mathbf{R}$  implies  $\mu(\vartheta) \leq 2\kappa$  .

Hence the result of Rhin and Viola  $\mu(\zeta(2)) \leq 5.441\dots$  implies only  $\mu(\pi) \leq 11.882\dots$

Conversely, a bound for the irrationality exponent of  $\vartheta$  does not yield any bound for  $\mu(\vartheta^2)$  !

# Irrationality measure for $\zeta(2)$ and $\zeta(3)$ : history

## $\zeta(2)$

R. Apéry 1978, F. Beukers 1979

$$\mu(\zeta(2)) < 11.85$$

R. Dvornicich and C. Viola 1987

$$\mu(\zeta(2)) < 10.02$$

M. Hata 1990

$$\mu(\zeta(2)) < 7.52$$

G. Rhin and C. Viola 1993

$$\mu(\zeta(2)) < 7.39$$

G. Rhin and C. Viola 1996

$$\mu(\zeta(2)) < 5.44$$

## $\zeta(3)$

R. Apéry 1978, F. Beukers 1979

$$\mu(\zeta(3)) < 13.41$$

R. Dvornicich and C. Viola 1987

$$\mu(\zeta(3)) < 12.74$$

M. Hata 1990

$$\mu(\zeta(3)) < 8.83$$

G. Rhin and C. Viola 2001

$$\mu(\zeta(3)) < 5.51$$



# Irrationality measure for $\log 2$ : history

$\log 2$

Hermite–Lindemann, Mahler, Baker, Gel'fond, Feldman, . . . :  
transcendence measures

G. Rhin 1987

$$\mu(\log 2) < 4.07$$

E.A. Rukhadze 1987


$$\mu(\log 2) < 3.89$$

R. Marcovecchio 2008

$$\mu(\log 2) < 3.57$$

*Reference* : R. Marcovecchio, The Rhin-Viola method for  $\log 2$ , Acta Arithmetica vol. **139** no.2 (2009), 147–184.

# Georges Rhin and Carlo Viola



On a permutation group related to  $\zeta(2)$ , *Acta Arith.* **77** (1996), no.1, 23–56.

The group structure for  $\zeta(3)$ , *Acta Arith.* **97** (2001), no.3, 269–293.

The permutation group method for the dilogarithm, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (**5**) **4** (2005), no.3, 389–437.

# Criterion of Yu. V. Nesterenko (qualitative)

Let  $\vartheta_1, \dots, \vartheta_m$  be complex numbers.



Yu.V.Nesterenko (1985)

Let  $m$  be a positive integer and  $\alpha$  a positive real number satisfying  $\alpha > m - 1$ . Assume there is a sequence  $(L_n)_{n \geq 0}$  of linear forms in

$\mathbf{Z}X_0 + \mathbf{Z}X_1 + \dots + \mathbf{Z}X_m$  of height  $\leq e^n$  such that

$$|L_n(1, \vartheta_1, \dots, \vartheta_m)| = e^{-\alpha n + o(n)}.$$

Then  $1, \vartheta_1, \dots, \vartheta_m$  are linearly independent over  $\mathbf{Q}$ .

Example :  $m = 1$  – irrationality criterion.

# Simplified proof of Nesterenko's Theorem



Francesco Amoroso



Pierre Colmez

Refinements : Raffaele Marcovecchio, Pierre Bel.

# Recent developments



Stéphane Fischler and Wadim Zudilin, *A refinement of Nesterenko's linear independence criterion with applications to zeta values.*  
To appear in Math. Annalen.

Preprint MPIM 2009-35.

# Fischler and Zudilin, 2009

There exist positive odd integers  $i \leq 139$  and  $j \leq 1961$  such that the numbers  $1, \zeta(3), \zeta(i), \zeta(j)$  are linearly independent over  $\mathbb{Q}$ .

There exist positive odd integers  $i \leq 93$  and  $j \leq 1151$  such that the numbers  $1, \log 2, \zeta(i), \zeta(j)$  are linearly independent over  $\mathbb{Q}$ .

# Fischler and Zudilin, 2009

There exist positive odd integers  $i \leq 139$  and  $j \leq 1961$  such that the numbers  $1, \zeta(3), \zeta(i), \zeta(j)$  are linearly independent over  $\mathbb{Q}$ .

There exist positive odd integers  $i \leq 93$  and  $j \leq 1151$  such that the numbers  $1, \log 2, \zeta(i), \zeta(j)$  are linearly independent over  $\mathbb{Q}$ .

# Multizeta values

For  $s_1, \dots, s_k$  positive integers with  $s_1 \geq 2$ ,

$$\zeta(s_1, \dots, s_k) = \sum_{n_1 > \dots > n_k \geq 1} \frac{1}{n_1^{s_1} \cdots n_k^{s_k}}$$

P. Cartier. –

*Fonctions polylogarithmes,  
nombres polyzêtas et groupes  
pro-unipotents.*

Sém. Bourbaki no. 885

Astérisque **282** (2002), 137-173.





# M. Hoffman's web site

<http://www.usna.edu/Users/math/meh/biblio.html>

## *References on multizeta values and Euler sums*

A	Double harmonic series	48 references
B	Triple harmonic series	8 references
C	Multiple harmonic series/multiple zeta values	137 references
D	Multiple zeta values over function fields	5 references
E	Alternating series	16 references
F	Multiple polylogarithms/nested sums	46 references
G	Finite multiple harmonic sums	25 references

In 2008 : 62 references

In 2009 : 30 references

+ preprints : 66 references

*Last modified on August 18, 2009*

# EZFace calculator at CECM



<http://oldweb.cecm.sfu.ca/projects/EZFace/>

## Centre for Experimental and Constructive Mathematics at Simon Fraser University

The calculator gives numerical values of MZVs with up to 100 decimal places accuracy.

The calculator also has a function to look for relations of linear dependence ;

$\text{lindep}([a, b, c])$  looks for a vanishing linear combination of  $a$ ,  $b$ ,  $c$  with integer coefficients.

This makes it easy (EZ ?) to discover new identities !

J. Blümlein, D.J. Broadhurst, J.A.M. Vermaseren

*The Multiple Zeta Value Data Mine*

arXiv :0907.2557v1 [math-ph]

# Gamma and Beta values




$$\begin{aligned}\Gamma(z) &= \int_0^{\infty} e^{-t} t^z \cdot \frac{dt}{t} \\ &= e^{-\gamma z} z^{-1} \prod_{n=1}^{\infty} \left(1 + \frac{z}{n}\right)^{-1} e^{z/n}.\end{aligned}$$

$$\Gamma(n+1) = n!, \quad (n \geq 0); \quad \Gamma(1/2) = \sqrt{\pi}, \quad \Gamma'(1) = -\gamma.$$

$$\begin{aligned}B(a, b) &= \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} \\ &= \int_0^1 x^{a-1} (1-x)^{b-1} dx.\end{aligned}$$

# Weierstraß functions



Let  $\Omega = \mathbf{Z}\omega_1 + \mathbf{Z}\omega_2$  be a lattice in  $\mathbf{C}$ .

The *canonical product* attached to  $\Omega$  is the Weierstraß sigma function

$$\sigma(z) = \sigma_{\Omega}(z) = z \prod_{\omega \in \Omega \setminus \{0\}} \left(1 - \frac{z}{\omega}\right) e^{(z/\omega) + (z^2/2\omega^2)}$$

The logarithmic derivative of the sigma function is the *Weierstraß zeta function*

$$\frac{\sigma'}{\sigma} = \zeta$$

and the derivative of  $\zeta$  is  $-\wp$ , where  $\wp$  is the Weierstraß elliptic function :

$$\wp'^2 = 4\wp^3 - g_2\wp - g_3,$$

$$\wp(z + \omega) = \wp(z), \quad \zeta(z + \omega) = \zeta(z) + \eta.$$

# Complex multiplication : $\mathbf{Q}(i)$

$$\wp'^2 = 4\wp^3 - 4\wp, \quad g_2 = 4, \quad g_3 = 0,$$

$$\omega_1 = \int_0^1 \frac{dx}{\sqrt{x-x^3}} = \frac{1}{2}B(1/4, 1/2) = \frac{\Gamma(1/4)^2}{\sqrt{8\pi}} = 2.622\,057\,554\,2\dots$$

$$\eta_1 = \frac{\pi}{\omega_1} = \frac{(2\pi)^{3/2}}{\Gamma(1/4)^2}, \quad \eta_2 = -i\eta_1.$$

# Complex multiplication : $\mathbf{Q}(\varrho)$

$$\varrho = e^{2i\pi/3}$$

$$\varphi'^2 = 4\varphi^3 - 4, \quad g_2 = 0, \quad g_3 = 4,$$

$$\omega_1 = \int_0^1 \frac{dx}{\sqrt{1-x^3}} = \frac{1}{3}B(1/3, 1/2) = \frac{\Gamma(1/3)^3}{2^{4/3}\pi} = 2.428\,650\,648\dots$$

$$\eta_1 = \frac{2\pi}{\sqrt{3}\omega_1} = \frac{2^{7/3}\pi^2}{3^{1/2}\Gamma(1/3)^3}, \quad \eta_2 = \varrho^2\eta_1.$$

# Transcendence of special values of Weierstraß functions



Th. Schneider (1934). *The numbers*

$$\Gamma(1/4)^4/\pi^3$$

*and*

$$\Gamma(1/3)^3/\pi^2$$

*are transcendental.*

# Diophantine approximation

$\Gamma(1/4)^4/\pi^3$  and  $\Gamma(1/3)^3/\pi^2$  are not Liouville numbers.

Lower bounds for linear combinations of elliptic logarithms :  
Baker, Coates, Anderson ...in the CM case,  
Philippon-Waldschmidt in the general case, refinements by  
N. Hirata Kohno, S. David, É. Gaudron - use Arakhelov's  
Theory (J-B. Bost : *slopes inequalities*).

*Motivation* : method of S. Lang for solving Diophantine  
equations (integer points on elliptic curves).



# Diophantine approximation

$\Gamma(1/4)^4/\pi^3$  and  $\Gamma(1/3)^3/\pi^2$  are not Liouville numbers.

Lower bounds for linear combinations of elliptic logarithms :  
Baker, Coates, Anderson ...in the CM case,  
Philippon-Waldschmidt in the general case, refinements by  
N. Hirata Kohno, S. David, É. Gaudron - use Arakhelov's  
Theory (J-B. Bost : *slopes inequalities*).

*Motivation* : method of S. Lang for solving Diophantine  
equations (integer points on elliptic curves).

# Diophantine approximation

$\Gamma(1/4)^4/\pi^3$  and  $\Gamma(1/3)^3/\pi^2$  are not Liouville numbers.

Lower bounds for linear combinations of elliptic logarithms :  
Baker, Coates, Anderson . . . in the CM case,  
Philippon-Waldschmidt in the general case, refinements by  
N. Hirata Kohno, S. David, É. Gaudron - use Arakhelov's  
Theory (J-B. Bost : *slopes inequalities*).

*Motivation* : method of S. Lang for solving Diophantine  
equations (integer points on elliptic curves).

# Sinnou David and Noriko Hirata



David, Sinnou ; Hirata-Kohno, Noriko  
Linear forms in elliptic logarithms.

J. Reine Angew. Math. **628** (2009), 37–89.

# Abelian varieties


Th. Schneider (1948). *For  $a$  and  $b$  in  $\mathbf{Q}$  with  $a$ ,  $b$  and  $a + b$  not in  $\mathbf{Z}$ , the number*

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a + b)}$$

*is transcendental.*

The proof involves Abelian integrals of higher genus, related with the Jacobian of a Fermat curve.

# Chudnovsky's algebraic independence Theorem



G.V. Chudnovsky (1978)

**Theorem** *Two at least of the numbers*

$$g_2, g_3, \omega_1, \omega_2, \eta_1, \eta_2$$

*are algebraically independent.*

**Corollary :**  $\pi$  and  $\Gamma(1/4) = 3.625\,609\,908\,2\dots$  are algebraically independent. Also  $\pi$  and  $\Gamma(1/3) = 2.678\,938\,534\,7\dots$  are algebraically independent.

# Diophantine approximation

## Transcendence measures for $\Gamma(1/4)$

(P. Philippon, S. Bruilhet)

For  $P \in \mathbf{Z}[X, Y]$  with degree  $d$  and height  $H$ ,

$$\log |P(\pi, \Gamma(1/4))| > -10^{326} ((\log H + d \log(d+1)) \cdot d^2 (\log(d+1))^2)$$

**Corollary :**  $\Gamma(1/4)$  is not a Liouville number :

$$\left| \Gamma(1/4) - \frac{p}{q} \right| > \frac{1}{q^{10^{330}}}.$$

# Chudnovsky's method


(K.G. Vasil'ev 1996, P. Grinspan 2002). Two at least of the three numbers  $\pi$ ,  $\Gamma(1/5)$  and  $\Gamma(2/5)$  are algebraically independent.

The proof involves a simple factor of dimension 2 of the Jacobian of the Fermat curve

$$X^5 + Y^5 = Z^5$$

which is an Abelian variety of dimension 6.

# Ramanujan Functions



$$P(q) = 1 - 24 \sum_{n=1}^{\infty} \frac{nq^n}{1 - q^n},$$

$$Q(q) = 1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n},$$

$$R(q) = 1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n}.$$



# Eisenstein Series

$$E_{2k}(z) = 1 + (-1)^k \frac{4k}{B_k} \sum_{n=1}^{\infty} \frac{n^{2k-1} z^n}{1 - z^n}.$$



F. G. M. Eisenstein  
(1823 - 1852)

$$P(z) = E_2(z),$$

$$Q(z) = E_4(z),$$

$$R(z) = E_6(z).$$

# Special values

$$\tau = i, \quad q = e^{-2\pi}, \quad \omega_1 = \frac{\Gamma(1/4)^2}{\sqrt{8\pi}} = 2.622\,057\,554\,2\dots$$

$$P(q) = \frac{3}{\pi}, \quad Q(q) = 3 \left(\frac{\omega_1}{\pi}\right)^4, \quad R(q) = 0.$$

$$\tau = \rho, \quad q = -e^{-\pi\sqrt{3}}, \quad \omega_1 = \frac{\Gamma(1/3)^3}{2^{4/3}\pi} = 2.428\,650\,648\dots$$

$$P(q) = \frac{2\sqrt{3}}{\pi}, \quad Q(q) = 0, \quad R(q) = \frac{27}{2} \left(\frac{\omega_1}{\pi}\right)^6.$$

# Yu. V. Nesterenko

**Theorem** (Nesterenko, 1996).  
*For any  $q \in \mathbf{C}$  with  $0 < |q| < 1$ ,  
three at least of the four numbers  
 $q$ ,  $P(q)$ ,  $Q(q)$ ,  $R(q)$   
are algebraically independent.*



**Tools** : The functions  $P$ ,  $Q$ ,  $R$  are algebraically independent over  $\mathbf{C}(q)$  (K. Mahler) and satisfy a system of differential equations for  $D = q d/dq$  :

$$12 \frac{DP}{P} = P - \frac{Q}{P}, \quad 3 \frac{DQ}{Q} = P - \frac{R}{Q}, \quad 2 \frac{DR}{R} = P - \frac{Q^2}{R}.$$

# Consequences of Nesterenko's Theorem

*The three numbers*

$$\pi, \quad e^\pi, \quad \Gamma(1/4)$$


*are algebraically independent.*

*The three numbers*

$$\pi, \quad e^{\pi\sqrt{3}}, \quad \Gamma(1/3)$$

*are algebraically independent.*

# Special values of Weierstraß sigma functions



*The number*

$$\sigma_{\mathbf{Z}[i]}(1/2) = 2^{5/4} \pi^{1/2} e^{\pi/8} \Gamma(1/4)^{-2}$$

*is transcendental.*

# Fibonacci zeta values

$F_0 = 0, F_1 = 1, F_{n+1} = F_n + F_{n-1}$

$$\zeta_F(s) = \sum_{n \geq 1} \frac{1}{F_n^s}$$

Akita Shiokawa (joint works  
with Carsten Elsner and Shun  
Shimomura, 2006)



$\zeta_F(2), \zeta_F(4), \zeta_F(6)$  are algebraically independent.  
Consequence of Nesterenko's Theorem.

Fibonacci zeta values  $\zeta_F(s) = \sum_{n \geq 1} \frac{1}{F_n^s}$

$$u = \zeta_F(2), \quad v = \zeta_F(4)$$

$\zeta_F(4s+2) \in \mathbf{Q}(u, v)$  for  $s \geq 0, s \in \mathbf{Z}$ .

$\zeta_F(4s) - r_s \zeta_F(4) \in \mathbf{Q}(u, v)$  for  $s \geq 2, s \in \mathbf{Z}$ , with some  $r_s \in \mathbf{Q} \setminus \{0\}$ .

For  $s_1, s_2, s_3$ , distinct positive integers, the numbers  $\zeta_F(2s_1), \zeta_F(2s_2), \zeta_F(2s_3)$  are algebraically dependent if and only if the three integers  $s_i$  are odd.

Fibonacci zeta values  $\zeta_F(s) = \sum_{n \geq 1} \frac{1}{F_n^s}$

$$u = \zeta_F(2), \quad v = \zeta_F(4)$$

$\zeta_F(4s + 2) \in \mathbf{Q}(u, v)$  for  $s \geq 0$ ,  $s \in \mathbf{Z}$ .

$\zeta_F(4s) - r_s \zeta_F(4) \in \mathbf{Q}(u, v)$  for  $s \geq 2$ ,  $s \in \mathbf{Z}$ , with some  $r_s \in \mathbf{Q} \setminus \{0\}$ .

For  $s_1, s_2, s_3$ , distinct positive integers, the numbers  $\zeta_F(2s_1), \zeta_F(2s_2), \zeta_F(2s_3)$  are algebraically dependent if and only if the three integers  $s_i$  are odd.



Fibonacci zeta values  $\zeta_F(s) = \sum_{n \geq 1} \frac{1}{F_n^s}$

$$u = \zeta_F(2), \quad v = \zeta_F(4)$$

$\zeta_F(4s + 2) \in \mathbf{Q}(u, v)$  for  $s \geq 0$ ,  $s \in \mathbf{Z}$ .

$\zeta_F(4s) - r_s \zeta_F(4) \in \mathbf{Q}(u, v)$  for  $s \geq 2$ ,  $s \in \mathbf{Z}$ , with some  $r_s \in \mathbf{Q} \setminus \{0\}$ .

For  $s_1, s_2, s_3$ , distinct positive integers, the numbers  $\zeta_F(2s_1), \zeta_F(2s_2), \zeta_F(2s_3)$  are algebraically dependent if and only if the three integers  $s_i$  are odd.

Fibonacci zeta values  $\zeta_F(s) = \sum_{n \geq 1} \frac{1}{F_n^s}$

$$u = \zeta_F(2), \quad v = \zeta_F(4)$$

$\zeta_F(4s + 2) \in \mathbf{Q}(u, v)$  for  $s \geq 0$ ,  $s \in \mathbf{Z}$ .

$\zeta_F(4s) - r_s \zeta_F(4) \in \mathbf{Q}(u, v)$  for  $s \geq 2$ ,  $s \in \mathbf{Z}$ , with some  $r_s \in \mathbf{Q} \setminus \{0\}$ .

For  $s_1, s_2, s_3$ , distinct positive integers, the numbers  $\zeta_F(2s_1), \zeta_F(2s_2), \zeta_F(2s_3)$  are algebraically dependent if and only if the three integers  $s_i$  are odd.

# Standard relations among Gamma values

Translation :

$$\Gamma(a + 1) = a\Gamma(a)$$

Reflexion :

$$\Gamma(a)\Gamma(1 - a) = \frac{\pi}{\sin(\pi a)}$$

Multiplication : for any positive integer  $n$ ,

$$\prod_{k=0}^{n-1} \Gamma\left(a + \frac{k}{n}\right) = (2\pi)^{(n-1)/2} n^{-na+(1/2)} \Gamma(na).$$

# Rohrlich's Conjecture

**Conjecture** (D. Rohrlich) *Any multiplicative relation*

$$\pi^{b/2} \prod_{a \in \mathbf{Q}} \Gamma(a)^{m_a} \in \overline{\mathbf{Q}}$$

*with  $b$  and  $m_a$  in  $\mathbf{Z}$  lies in the ideal generated by the standard relations.*

Examples :

$$\Gamma\left(\frac{1}{14}\right) \Gamma\left(\frac{9}{14}\right) \Gamma\left(\frac{11}{14}\right) = 4\pi^{3/2}$$

$$\prod_{\substack{1 \leq k \leq n \\ (k,n)=1}} \Gamma(k/n) = \begin{cases} (2\pi)^{\varphi(n)/2} / \sqrt{p} & \text{if } n = p^r \text{ is a prime power,} \\ (2\pi)^{\varphi(n)/2} & \text{otherwise.} \end{cases}$$

# Rohrlich's Conjecture

**Conjecture** (D. Rohrlich) *Any multiplicative relation*

$$\pi^{b/2} \prod_{a \in \mathbf{Q}} \Gamma(a)^{m_a} \in \overline{\mathbf{Q}}$$

*with  $b$  and  $m_a$  in  $\mathbf{Z}$  lies in the ideal generated by the standard relations.*

Examples :

$$\Gamma\left(\frac{1}{14}\right) \Gamma\left(\frac{9}{14}\right) \Gamma\left(\frac{11}{14}\right) = 4\pi^{3/2}$$

$$\prod_{\substack{1 \leq k \leq n \\ (k,n)=1}} \Gamma(k/n) = \begin{cases} (2\pi)^{\varphi(n)/2} / \sqrt{p} & \text{if } n = p^r \text{ is a prime power,} \\ (2\pi)^{\varphi(n)/2} & \text{otherwise.} \end{cases}$$

# Rohrlich's Conjecture

**Conjecture** (D. Rohrlich) *Any multiplicative relation*

$$\pi^{b/2} \prod_{a \in \mathbf{Q}} \Gamma(a)^{m_a} \in \overline{\mathbf{Q}}$$

*with  $b$  and  $m_a$  in  $\mathbf{Z}$  lies in the ideal generated by the standard relations.*

Examples :

$$\Gamma\left(\frac{1}{14}\right) \Gamma\left(\frac{9}{14}\right) \Gamma\left(\frac{11}{14}\right) = 4\pi^{3/2}$$

$$\prod_{\substack{1 \leq k \leq n \\ (k,n)=1}} \Gamma(k/n) = \begin{cases} (2\pi)^{\varphi(n)/2} / \sqrt{p} & \text{if } n = p^r \text{ is a prime power,} \\ (2\pi)^{\varphi(n)/2} & \text{otherwise.} \end{cases}$$

# Small Gamma Products with Simple Values

The two previous examples are due respectively to

Albert Nijenhuis, *Small Gamma products with Simple Values*  
<http://arxiv.org/abs/0907.1689>, July 9, 2009.

and to

Greg Martin, *A product of Gamma function values at fractions with the same denominator*  
<http://arxiv.org/abs/0907.4384>, July 24, 2009.

# Small Gamma Products with Simple Values

The two previous examples are due respectively to


Albert Nijenhuis, *Small Gamma products with Simple Values*  
<http://arxiv.org/abs/0907.1689>, July 9, 2009.

and to

Greg Martin, *A product of Gamma function values at fractions with the same denominator*  
<http://arxiv.org/abs/0907.4384>, July 24, 2009.



# Lang's Conjecture



**Conjecture** (S. Lang) *Any algebraic dependence relation among the numbers  $(2\pi)^{-1/2}\Gamma(a)$  with  $a \in \mathbf{Q}$  lies in the ideal generated by the standard relations.*  
(Universal odd distribution).

# Consequence of the Rohrlich–Lang Conjecture

As an example, the Rohrlich–Lang Conjecture implies that for any  $q > 1$ , the transcendence degree of the field generated by numbers

$$\pi, \quad \Gamma(a/q) \quad 1 \leq a \leq q, \quad (a, q) = 1$$

is  $1 + \varphi(q)/2$ .

# Variant of the Rohrlich–Lang Conjecture

Conjecture of S. Gun, R. Murty, P. Rath (2009) : for any  $q > 1$ , the numbers

$$\log \Gamma(a/q) \quad 1 \leq a \leq q, (a, q) = 1$$

are linearly independent over the field  $\overline{\mathbb{Q}}$  of algebraic numbers.

A consequence is that for any  $q > 1$ , there is at most one primitive odd character  $\chi$  modulo  $q$  for which

$$L'(1, \chi) = 0.$$

# Variant of the Rohrlich–Lang Conjecture

Conjecture of S. Gun, R. Murty, P. Rath (2009) : for any  $q > 1$ , the numbers

$$\log \Gamma(a/q) \quad 1 \leq a \leq q, (a, q) = 1$$

are linearly independent over the field  $\overline{\mathbb{Q}}$  of algebraic numbers.

A consequence is that for any  $q > 1$ , there is at most one primitive odd character  $\chi$  modulo  $q$  for which

$$L'(1, \chi) = 0.$$

# Ram and Kumar Murty (2009)

Ram Murty




Kumar Murty



Transcendental values of class group  $L$ -functions.

# Peter Bundschuh (1979)



For  $p/q \in \mathbf{Q}$  with  
 $0 < |p/q| < 1$ ,

$$\sum_{n=2}^{\infty} \zeta(n) (p/q)^n$$

is transcendental.

For  $p/q \in \mathbf{Q} \setminus \mathbf{Z}$ ,

$$\frac{\Gamma'}{\Gamma} \left( \frac{p}{q} \right) + \gamma$$

is transcendental

# Peter Bundschuh (1979)

(P. Bundschuh) : *As a consequence of Nesterenko's Theorem, the number*

$$\sum_{n=0}^{\infty} \frac{1}{n^2 + 1} = \frac{1}{2} + \frac{\pi}{2} \cdot \frac{e^{\pi} + e^{-\pi}}{e^{\pi} - e^{-\pi}} = 2.076\,674\,047\,4\dots$$

*is transcendental, while*

$$\sum_{n=0}^{\infty} \frac{1}{n^2 - 1} = \frac{3}{4}$$

(telescoping series).

Hence the number

$$\sum_{n=2}^{\infty} \frac{1}{n^s - 1}$$

is transcendental over  $\mathbb{Q}$  for  $s = 4$ . The transcendence of this number for even integers  $s \geq 4$  would follow as a consequence of Schanuel's Conjecture.

# Peter Bundschuh (1979)

(P. Bundschuh) : *As a consequence of Nesterenko's Theorem, the number*

$$\sum_{n=0}^{\infty} \frac{1}{n^2 + 1} = \frac{1}{2} + \frac{\pi}{2} \cdot \frac{e^{\pi} + e^{-\pi}}{e^{\pi} - e^{-\pi}} = 2.076\,674\,047\,4\dots$$

*is transcendental, while*

$$\sum_{n=0}^{\infty} \frac{1}{n^2 - 1} = \frac{3}{4}$$

(telescoping series).

Hence the number

$$\sum_{n=2}^{\infty} \frac{1}{n^s - 1}$$

is transcendental over  $\mathbf{Q}$  for  $s = 4$ . The transcendence of this number for even integers  $s \geq 4$  would follow as a consequence of Schanuel's Conjecture.



$$\sum_{n \geq 1} A(n)/B(n)$$

Arithmetic nature of

$$\sum_{n \geq 1} \frac{A(n)}{B(n)}$$

where

$$A/B \in \mathbf{Q}(X).$$

In case  $B$  has distinct zeroes, by decomposing  $A/B$  in simple fractions one gets linear combinations of logarithms of algebraic numbers (Baker's method).

The example  $A(X)/B(X) = 1/X^3$  shows that the general case is hard.

Work by S.D. Adhikari, N. Saradha, T.N. Shorey and R. Tijdeman (2001),

Sanoli Gun, Ram Murty and Purusottam Rath (2009).

# Adolf Hurwitz (1859 - 1919)



Hurwitz zeta function :  
for  $z \in \mathbf{C}$   $z \neq 0$  and  
 $\Re(s) > 1$ ,

$$\zeta(s, z) = \sum_{n=0}^{\infty} \frac{1}{(n+z)^s}$$

$$\zeta(s, 1) = \zeta(s)$$


(Riemann zeta function)

# Conjecture of Chowla and Milnor

Sarvadaman Chowla  
(1907 - 1995)



John Willard Milnor  
(1931 - )



For  $k$  and  $q$  integers  $> 1$ , the  $\varphi(q)$  numbers

$$\zeta(k, a/q), \quad 1 \leq a \leq q, \quad (a, q) = 1$$

are linearly independent over  $\mathbf{Q}$ .

# Sanoli Gun, Ram Murty and Purusottam Rath

The Chowla-Milnor Conjecture for  $q = 4$  implies the irrationality of the numbers  $\zeta(2n + 1)/\pi^{2n+1}$  for  $n \geq 1$ .

**Strong Chowla-Milnor Conjecture (2009)** : For  $k$  and  $q$  integers  $> 1$ , the  $1 + \varphi(q)$  numbers

$$1 \quad \text{and} \quad \zeta(k, a/q), \quad 1 \leq a \leq q, \quad (a, q) = 1$$

are linearly independent over  $\mathbb{Q}$ .

For  $k > 1$  odd, the number  $\zeta(k)$  is irrational if and only if the strong Chowla-Milnor Conjecture holds for this value of  $k$  and for at least one of the two values  $q = 3$  and  $q = 4$ .

Hence the strong Chowla-Milnor Conjecture holds for  $k = 3$  (Apéry) and also for infinitely many  $k$  (Rivoal).

# Sanoli Gun, Ram Murty and Purusottam Rath

The Chowla-Milnor Conjecture for  $q = 4$  implies the irrationality of the numbers  $\zeta(2n + 1)/\pi^{2n+1}$  for  $n \geq 1$ .

**Strong Chowla-Milnor Conjecture (2009)** : For  $k$  and  $q$  integers  $> 1$ , the  $1 + \varphi(q)$  numbers

$$1 \quad \text{and} \quad \zeta(k, a/q), \quad 1 \leq a \leq q, \quad (a, q) = 1$$

are linearly independent over  $\mathbf{Q}$ .

For  $k > 1$  odd, the number  $\zeta(k)$  is irrational if and only if the strong Chowla-Milnor Conjecture holds for this value of  $k$  and for at least one of the two values  $q = 3$  and  $q = 4$ .

Hence the strong Chowla-Milnor Conjecture holds for  $k = 3$  (Apéry) and also for infinitely many  $k$  (Rivoal).

# Sanoli Gun, Ram Murty and Purusottam Rath

The Chowla-Milnor Conjecture for  $q = 4$  implies the irrationality of the numbers  $\zeta(2n + 1)/\pi^{2n+1}$  for  $n \geq 1$ .

**Strong Chowla-Milnor Conjecture (2009)** : For  $k$  and  $q$  integers  $> 1$ , the  $1 + \varphi(q)$  numbers

$$1 \quad \text{and} \quad \zeta(k, a/q), \quad 1 \leq a \leq q, \quad (a, q) = 1$$

are linearly independent over  $\mathbf{Q}$ .

For  $k > 1$  odd, the number  $\zeta(k)$  is irrational if and only if the strong Chowla-Milnor Conjecture holds for this value of  $k$  and for at least one of the two values  $q = 3$  and  $q = 4$ .

Hence the strong Chowla-Milnor Conjecture holds for  $k = 3$  (Apéry) and also for infinitely many  $k$  (Rivoal).

# Sanoli Gun, Ram Murty and Purusottam Rath

The Chowla-Milnor Conjecture for  $q = 4$  implies the irrationality of the numbers  $\zeta(2n + 1)/\pi^{2n+1}$  for  $n \geq 1$ .

**Strong Chowla-Milnor Conjecture (2009)** : For  $k$  and  $q$  integers  $> 1$ , the  $1 + \varphi(q)$  numbers

$$1 \quad \text{and} \quad \zeta(k, a/q), \quad 1 \leq a \leq q, \quad (a, q) = 1$$

are linearly independent over  $\mathbf{Q}$ .

For  $k > 1$  odd, the number  $\zeta(k)$  is irrational if and only if the strong Chowla-Milnor Conjecture holds for this value of  $k$  and for at least one of the two values  $q = 3$  and  $q = 4$ .

Hence the strong Chowla-Milnor Conjecture holds for  $k = 3$  (Apéry) and also for infinitely many  $k$  (Rivoal).

# Linear independence of polylogarithms

For  $k \geq 1$  and  $|z| < 1$ ,

$$\text{Li}_k(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^k}.$$

Thus  $\text{Li}_1(z) = \log(1 - z)$  and  $\text{Li}_k(1) = \zeta(k)$  for  $k \geq 2$ .

Polylog Conjecture of S. Gun, R. Murty, P. Rath : *Let  $k > 1$  be an integer and  $\alpha_1, \dots, \alpha_n$  algebraic numbers such that  $\text{Li}_k(\alpha_1), \dots, \text{Li}_k(\alpha_n)$  are linearly independent over  $\mathbb{Q}$ . Then these numbers  $\text{Li}_k(\alpha_1), \dots, \text{Li}_k(\alpha_n)$  are linearly independent over the field  $\overline{\mathbb{Q}}$  of algebraic numbers.*

S. Gun, R. Murty, P. Rath : *if the polylog Conjecture is true, then the Chowla-Milnor Conjecture is true for all  $k$  and all  $q$ .*



# Linear independence of polylogarithms

For  $k \geq 1$  and  $|z| < 1$ ,

$$\text{Li}_k(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^k}.$$

Thus  $\text{Li}_1(z) = \log(1 - z)$  and  $\text{Li}_k(1) = \zeta(k)$  for  $k \geq 2$ .

Polylog Conjecture of S. Gun, R. Murty, P. Rath : *Let  $k > 1$  be an integer and  $\alpha_1, \dots, \alpha_n$  algebraic numbers such that  $\text{Li}_k(\alpha_1), \dots, \text{Li}_k(\alpha_n)$  are linearly independent over  $\mathbb{Q}$ . Then these numbers  $\text{Li}_k(\alpha_1), \dots, \text{Li}_k(\alpha_n)$  are linearly independent over the field  $\overline{\mathbb{Q}}$  of algebraic numbers.*

S. Gun, R. Murty, P. Rath : *if the polylog Conjecture is true, then the Chowla-Milnor Conjecture is true for all  $k$  and all  $q$ .*

# Linear independence of polylogarithms

For  $k \geq 1$  and  $|z| < 1$ ,

$$\operatorname{Li}_k(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^k}.$$

Thus  $\operatorname{Li}_1(z) = \log(1 - z)$  and  $\operatorname{Li}_k(1) = \zeta(k)$  for  $k \geq 2$ .

*Polylog Conjecture of S. Gun, R. Murty, P. Rath : Let  $k > 1$  be an integer and  $\alpha_1, \dots, \alpha_n$  algebraic numbers such that  $\operatorname{Li}_k(\alpha_1), \dots, \operatorname{Li}_k(\alpha_n)$  are linearly independent over  $\mathbf{Q}$ . Then these numbers  $\operatorname{Li}_k(\alpha_1), \dots, \operatorname{Li}_k(\alpha_n)$  are linearly independent over the field  $\overline{\mathbf{Q}}$  of algebraic numbers.*

*S. Gun, R. Murty, P. Rath : if the polylog Conjecture is true, then the Chowla-Milnor Conjecture is true for all  $k$  and all  $q$ .*

# The digamma function

For  $z \in \mathbf{C} \setminus \{0, -1, -2, \dots\}$ ,

$$\psi(x) = \frac{d}{dz} \log \Gamma(z) = \frac{\Gamma'(z)}{\Gamma(z)}.$$

$$\psi(z) = -\gamma - \frac{1}{z} - \sum_{n=1}^{\infty} \left( \frac{1}{n+z} - \frac{1}{n} \right)$$

$$\psi(1+z) = -\gamma + \sum_{n=2}^{\infty} (-1)^n \zeta(n) z^{n-1}.$$

# The digamma function

For  $z \in \mathbf{C} \setminus \{0, -1, -2, \dots\}$ ,

$$\psi(x) = \frac{d}{dz} \log \Gamma(z) = \frac{\Gamma'(z)}{\Gamma(z)}.$$

$$\psi(z) = -\gamma - \frac{1}{z} - \sum_{n=1}^{\infty} \left( \frac{1}{n+z} - \frac{1}{n} \right)$$

$$\psi(1+z) = -\gamma + \sum_{n=2}^{\infty} (-1)^n \zeta(n) z^{n-1}.$$

# The digamma function

For  $z \in \mathbf{C} \setminus \{0, -1, -2, \dots\}$ ,

$$\psi(x) = \frac{d}{dz} \log \Gamma(z) = \frac{\Gamma'(z)}{\Gamma(z)}.$$

$$\psi(z) = -\gamma - \frac{1}{z} - \sum_{n=1}^{\infty} \left( \frac{1}{n+z} - \frac{1}{n} \right)$$

$$\psi(1+z) = -\gamma + \sum_{n=2}^{\infty} (-1)^n \zeta(n) z^{n-1}.$$

# Special values of the digamma function

$$\psi(1) = -\gamma, \quad \psi\left(\frac{1}{2}\right) = -2\log(2) - \gamma,$$

$$\psi\left(2k - \frac{1}{2}\right) = -2\log(2) - \gamma + \sum_{n=1}^{2k-1} \frac{1}{n + 1/2},$$

$$\psi\left(\frac{1}{4}\right) = -\frac{\pi}{2} - 3\log(2) - \gamma,$$

$$\psi\left(\frac{3}{4}\right) = \frac{\pi}{2} - 3\log(2) - \gamma.$$

Hence

$$\psi(1) + \psi(1/4) - 3\psi(1/2) + \psi(3/4) = 0.$$

# Special values of the digamma function

$$\psi(1) = -\gamma, \quad \psi\left(\frac{1}{2}\right) = -2\log(2) - \gamma,$$

$$\psi\left(2k - \frac{1}{2}\right) = -2\log(2) - \gamma + \sum_{n=1}^{2k-1} \frac{1}{n + 1/2},$$

$$\psi\left(\frac{1}{4}\right) = -\frac{\pi}{2} - 3\log(2) - \gamma,$$

$$\psi\left(\frac{3}{4}\right) = \frac{\pi}{2} - 3\log(2) - \gamma.$$

Hence

$$\psi(1) + \psi(1/4) - 3\psi(1/2) + \psi(3/4) = 0.$$

# Special values of the digamma function

$$\psi(1) = -\gamma, \quad \psi\left(\frac{1}{2}\right) = -2\log(2) - \gamma,$$

$$\psi\left(2k - \frac{1}{2}\right) = -2\log(2) - \gamma + \sum_{n=1}^{2k-1} \frac{1}{n + 1/2},$$

$$\psi\left(\frac{1}{4}\right) = -\frac{\pi}{2} - 3\log(2) - \gamma,$$

$$\psi\left(\frac{3}{4}\right) = \frac{\pi}{2} - 3\log(2) - \gamma.$$

Hence

$$\psi(1) + \psi(1/4) - 3\psi(1/2) + \psi(3/4) = 0.$$



# Special values of the digamma function

$$\psi(1) = -\gamma, \quad \psi\left(\frac{1}{2}\right) = -2\log(2) - \gamma,$$

$$\psi\left(2k - \frac{1}{2}\right) = -2\log(2) - \gamma + \sum_{n=1}^{2k-1} \frac{1}{n + 1/2},$$

$$\psi\left(\frac{1}{4}\right) = -\frac{\pi}{2} - 3\log(2) - \gamma,$$

$$\psi\left(\frac{3}{4}\right) = \frac{\pi}{2} - 3\log(2) - \gamma.$$

Hence

$$\psi(1) + \psi(1/4) - 3\psi(1/2) + \psi(3/4) = 0.$$

# Special values of the digamma function

$$\psi(1) = -\gamma, \quad \psi\left(\frac{1}{2}\right) = -2\log(2) - \gamma,$$

$$\psi\left(2k - \frac{1}{2}\right) = -2\log(2) - \gamma + \sum_{n=1}^{2k-1} \frac{1}{n + 1/2},$$

$$\psi\left(\frac{1}{4}\right) = -\frac{\pi}{2} - 3\log(2) - \gamma,$$

$$\psi\left(\frac{3}{4}\right) = \frac{\pi}{2} - 3\log(2) - \gamma.$$

Hence

$$\psi(1) + \psi(1/4) - 3\psi(1/2) + \psi(3/4) = 0.$$

# Ram Murty and N. Saradha

Conjecture (2007) : Let  $K$  be a number field over which the  $q$ -th cyclotomic polynomial is irreducible. Then the  $\varphi(q)$  numbers  $\psi(a/q)$  with  $1 \leq a \leq q$  and  $(a, q) = 1$  are linearly independent over  $K$ .



# Ram Murty and N. Saradha

Baker periods : elements of the  $\overline{\mathbb{Q}}$ -vector space spanned by the logarithms of algebraic numbers.

A Baker period is a period in the sense of Kontsevich and Zagier, and is either zero or else transcendental, by Baker's Theorem.


Murty and Saradha : one at least of the two following statements is true :

- Euler's Constant  $\gamma$  is not a Baker period
- the  $\varphi(q)$  numbers  $\psi(a/q)$  with  $1 \leq a \leq q$  and  $(a, q) = 1$  are linearly independent over  $K$ , whenever  $K$  be a number field over which the  $q$ -th cyclotomic polynomial is irreducible.

# Euler constant

## Euler–Mascheroni constant

$$\gamma = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n \right) = 0.577\,215\,664\,9\dots$$



Neil J. A. Sloane's encyclopaedia

<http://www.research.att.com/~njas/sequences/A001620>



$$\gamma = \int_0^{\infty} \sum_{k=2}^{\infty} \frac{1}{k^2 \binom{t+k}{k}} dt$$

$$\gamma = \lim_{s \rightarrow 1^+} \sum_{n=1}^{\infty} \left( \frac{1}{n^s} - \frac{1}{s^n} \right)$$

$$\gamma = \int_1^{\infty} \frac{1}{2t(t+1)} F \left( \begin{matrix} 1, & 2, & 2 \\ 3, & t+2 \end{matrix} \right) dt.$$

# Euler's constant $\gamma$

A.I. Aptekarev (2007) : Approximation to Euler's constant.

Tanguy Rivoal (2009) : Approximation to the function  $\gamma + \log x$ .

*Consequence* : approximation to  $\gamma$  and to  $\zeta(2) - \gamma^2$ .

## Open Problems.

- *Is the Euler constant  $\gamma$  irrational?*
- *Is  $\gamma$  transcendental?*
- *Kontsevich – Zagier :  $\gamma$  is not a period*

# Euler's constant $\gamma$

A.I. Aptekarev (2007) : Approximation to Euler's constant.

Tanguy Rivoal (2009) : Approximation to the function  $\gamma + \log x$ .

*Consequence* : approximation to  $\gamma$  and to  $\zeta(2) - \gamma^2$ .

## Open Problems.

- *Is the Euler constant  $\gamma$  irrational?*
- *Is  $\gamma$  transcendental?*
- *Kontsevich – Zagier :  $\gamma$  is not a period*



# Euler's constant $\gamma$

A.I. Aptekarev (2007) : Approximation to Euler's constant.

Tanguy Rivoal (2009) : Approximation to the function  $\gamma + \log x$ .

*Consequence* : approximation to  $\gamma$  and to  $\zeta(2) - \gamma^2$ .

## Open Problems.

- *Is the Euler constant  $\gamma$  irrational?*
- *Is  $\gamma$  transcendental?*
- *Kontsevich – Zagier :  $\gamma$  is not a period*

# Euler's constant $\gamma$

A.I. Aptekarev (2007) : Approximation to Euler's constant.

Tanguy Rivoal (2009) : Approximation to the function  $\gamma + \log x$ .


*Consequence* : approximation to  $\gamma$  and to  $\zeta(2) - \gamma^2$ .

## Open Problems.

- *Is the Euler constant  $\gamma$  irrational?*
- *Is  $\gamma$  transcendental?*
- *Kontsevich – Zagier :  $\gamma$  is not a period*

# Carlitz zeta values

Leonard Carlitz (1907 - 1999)



$$A = \mathbf{F}_q[t],$$

$A_+ \subset A$  monic polynomials,

$P =$  prime polynomials in  $A_+$ ,

$$K = \mathbf{F}_q(t),$$

$$K_\infty = \mathbf{F}_q((1/t)),$$

Carlitz zeta values : for  $s \in \mathbf{Z}$ ,

$$\zeta_A(s) = \sum_{a \in A_+} \frac{1}{a^s} = \prod_{p \in P} (1 - p^{-s})^{-1} \in K_\infty.$$

# Thakur Gamma function

Dinesh Thakur



$$\Gamma(z) = \frac{1}{z} \prod_{a \in A_+} \left(1 + \frac{z}{a}\right)$$

# Thakur Gamma values

Independence of Gamma values in positive characteristic :  
Linear relations ( W.D. Brownawell and M. Papanikolas, 2002)  
and algebraic relations (with G. Anderson, 2004).



Dale Brownawell



Matt Papanikolas

# Carlitz zeta values at even $A$ -integers

Define

$$\tilde{\pi} = (t - t^q)^{1/(q-1)} \prod_{n=1}^{\infty} \left( 1 - \frac{t^{q^n} - t}{t^{q^{n+1}} - t} \right)$$

For  $m$  a multiple of  $q - 1$ ,

$$\tilde{\pi}^{-m} \zeta_A(m) \in A$$

Carlitz – Bernoulli numbers.

# Carlitz zeta values at even $A$ -integers

Define

$$\tilde{\pi} = (t - t^q)^{1/(q-1)} \prod_{n=1}^{\infty} \left( 1 - \frac{t^{q^n} - t}{t^{q^{n+1}} - t} \right)$$

For  $m$  a multiple of  $q - 1$ ,

$$\tilde{\pi}^{-m} \zeta_A(m) \in A$$

Carlitz – Bernoulli numbers.

# Greg Anderson, Dinesh Thakur, Jing Yu

For  $m$  a positive integer,  $\zeta_A(m)$  is transcendental over  $K$ .

For  $m$  a positive integer not a multiple of  $q - 1$ ,  $\zeta_A(m)/\tilde{\pi}^m$  is transcendental over  $K$ .

Dinesh Thakur



Jing Yu





# Bourbaki Seminar

Federico PELLARIN

*Aspects de l'indépendance algébrique en caractéristique non nulle*

*Aspects of algebraic independence in non-zero characteristic*

Séminaire Bourbaki - Volume 2006/2007 - Exposés 967-981

Astérisque **317** (2008), 205–242

Title : *Geometric Gamma values and zeta values in positive characteristic*

arXiv :0905.2876

Abstract : In analogy with values of the classical Euler Gamma-function at rational numbers and the Riemann zeta-function at positive integers, we consider Thakur's geometric Gamma-function evaluated at rational arguments and Carlitz zeta-values at positive integers. We prove that, when considered together, all of the algebraic relations among these special values arise from the standard functional equations of the Gamma-function and from the Euler-Carlitz relations and Frobenius  $p$ -th power relations of the zeta-function.

# Chieh-Yu Chang, Matthew A. Papanikolas, Dinesh S. Thakur, Jing Yu

Title : *Algebraic independence of arithmetic gamma values and Carlitz zeta values* [arXiv :0909.0096](https://arxiv.org/abs/0909.0096)

Abstract : We consider the values at proper fractions of the arithmetic gamma function and the values at positive integers of the zeta function for  $\mathbf{F}_q[\theta]$  and provide complete algebraic independence results for them.

# Chieh-Yu Chang



Title : *Periods of third kind for rank 2 Drinfeld modules and algebraic independence of logarithms*

arXiv :0909.0101

**Abstract** : In analogy with the periods of abelian integrals of differentials of third kind for an elliptic curve defined over a number field, we introduce a notion of periods of third kind for a rank 2 Drinfeld  $\mathbf{F}_q[t]$ -module  $\rho$  defined over an algebraic function field and derive explicit formulae for them. When  $\rho$  has complex multiplication by a separable extension, we prove the algebraic independence of  $\rho$ -logarithms of algebraic points that are linearly independent over the CM field of  $\rho$ . Together with the main result in [CP08], we completely determine all the algebraic relations among the periods of first, second and third kinds for rank 2 Drinfeld  $\mathbf{F}_q[t]$ -modules in odd characteristic.



The Institute of Mathematical Sciences  
Mathematics Colloquium

December 9, 2009

<http://www.imsc.res.in/>

## Recent Diophantine results on zeta values: a survey

*Michel Waldschmidt*

Institut de Mathématiques de Jussieu & Paris VI

<http://www.math.jussieu.fr/~miw/>