

THE VALUATION PAIRING ON AN UPPER CLUSTER ALGEBRA

PEIGEN CAO, BERNHARD KELLER, AND FAN QIN

ABSTRACT. It is known that many (upper) cluster algebras are not unique factorization domains. We exhibit the local factorization properties with respect to any given seed t : any non-zero element in a full rank upper cluster algebra can be uniquely written as the product of a cluster monomial in t and another element not divisible by the cluster variables in t . Our approach is based on introducing the valuation pairing on an upper cluster algebra: it counts the maximal multiplicity of a cluster variable among the factorizations of any given element.

We apply the valuation pairing to obtain many results concerning factoriality, d -vectors, F -polynomials and the combinatorics of cluster Poisson variables. In particular, we obtain that full rank and primitive upper cluster algebras are factorial; an explanation of d -vectors using valuation pairing; a cluster monomial in non-initial cluster variables is determined by its F -polynomial; the F -polynomials of non-initial cluster variables are irreducible; and the cluster Poisson variables parametrize the exchange pairs of the corresponding upper cluster algebra.

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. The valuation pairing on an upper cluster algebra	13
4. Application to factoriality of upper cluster algebras	16
5. Applications to d -vectors	22
6. Application to F -polynomials	25
7. Application to combinatorics of cluster Poisson variables	28
References	36

1. INTRODUCTION

1.1. **Background.** Around the year 2000, Fomin and Zelevinsky introduced [FZ02] cluster algebras with the aim of developing a combinatorial approach to the theory of canonical bases in quantum groups and the closely related theory of total positivity in algebraic groups. Since then, cluster algebras have been linked to numerous other subjects and their study has flourished, cf. for example the surveys [Lec10, Fom10, Kel12, KD20].

2010 *Mathematics Subject Classification.* 13F60; 05E40.

Key words and phrases. Cluster algebra, valuation pairing, d -vectors, F -polynomials, cluster Poisson variables, unique factorization domains.

A cluster algebra \mathcal{A} is a subalgebra of an ambient field \mathcal{F} generated by certain combinatorially defined generators (called *cluster variables*), which are grouped into overlapping sets (called *clusters*) of constant cardinality n . Different clusters are obtained from each other by a sequence of *mutations*. One remarkable feature of cluster algebras is the Laurent phenomenon [FZ02], that is, for any given cluster $A_{t_0} = (A_{1;t_0}, \dots, A_{n;t_0})$, each cluster variable $A_{k;t}$ can be written as

$$A_{k;t} = \frac{P(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{d_1} \cdots A_{n;t_0}^{d_n}},$$

where P is a polynomial in variables from A_{t_0} such that $A_{i;t_0}$ does not divide P for any i . The vector

$$\mathbf{d}^{t_0}(A_{k;t}) = (d_1, \dots, d_n)^T$$

is called the *d-vector* of $A_{k;t}$ with respect to A_{t_0} and the polynomial P is called the *numerator polynomial* of $A_{k;t}$ with respect to A_{t_0} .

For each cluster algebra \mathcal{A} , Fock–Goncharov defined [FG09] a pair of varieties: the *cluster K_2 -variety* and the *cluster Poisson variety* (we follow the terminology of the appendix to [SW20]). The *upper cluster algebra* \mathcal{U} is defined to be the algebra of global functions on the cluster K_2 -variety, and the *cluster Poisson algebra* \mathcal{X} is defined to be the algebra of the global functions on the cluster Poisson variety. The clusters of \mathcal{U} correspond to local toric charts and the cluster variables correspond to local coordinates (which happen to be global functions) of the cluster K_2 -variety. The cluster Poisson variety admits a canonical atlas of dual toric charts whose coordinates are the *cluster Poisson variables* (but they are not global) of \mathcal{X} .

1.2. Main results. An upper cluster algebra is constructed from seeds. Each seed contains a cluster and a skew-symmetrizable matrix B called its exchange matrix, see Section 2. Moreover, when the upper cluster algebra has geometric type, we can associate to t the extended exchange matrix \tilde{B} , see Definition 2.10. The upper cluster algebra is said to be full rank or primitive if \tilde{B} is. Such properties are independent of the choice of the seed.

Valuation pairings, factoriality and the ray fish theorem. It is known from [GLS13] that many (upper) cluster algebras are not unique factorization domains. In order to study the local factorization properties of upper cluster algebras, we introduce the *valuation pairing* (see Definition 3.1) on any upper cluster algebra \mathcal{U} . To each pair $(A_{k;t}, M)$ consisting of a cluster variable $A_{k;t}$ and an element M in \mathcal{U} , it associates the largest integer s (possibly infinity) such that $M/A_{k;t}^s$ still belongs to \mathcal{U} . We write $(A_{k;t} || M)_v = s$. Using the valuation pairing we prove that any full rank upper cluster algebra has the following *local unique factorization property*: For each seed t of \mathcal{U} , any non-zero element M can be uniquely factorized as $M = N \cdot L$, where N is a cluster monomial in t and L is an element in \mathcal{U} not divisible by any cluster variable in t . We will give many applications to *d*-vectors, *F*-polynomials, factoriality of upper cluster algebras and combinatorics of cluster Poisson variables.

As an application to factoriality of upper cluster algebras, we prove that a full rank upper cluster algebra \mathcal{U} with initial seed t_0 is factorial if and only if each exchange binomial of t_0 is irreducible in the corresponding polynomial ring (see Theorem 4.4). In particular, full rank, primitive upper cluster algebras are factorial (see Theorem 4.8 (i)). These include principal coefficient upper cluster algebras as a special case. For full rank, primitive upper cluster algebras, we also prove that the

numerator polynomials of non-initial cluster variables are irreducible (see Theorem 4.8 (ii)).

The starfish theorem in [BFZ05], cf. also Theorem 2.16 of the present paper, plays a very important role in this paper. It states that any full rank upper cluster algebra \mathcal{U} can be written as the intersection of $n+1$ Laurent polynomial rings. Here n is the rank of \mathcal{U} . Thanks to the results on factoriality of upper cluster algebras, we show that any full rank, primitive upper cluster algebra can be written as the intersection of *two* Laurent polynomial rings (see Theorem 4.18). We call this the *ray fish theorem*.

Application to d -vectors. In [CL20], Li and the first author of the present paper proved that there exists a well-defined function $(-||-)_d$ on the set of cluster variables, which is called the *d -compatibility degree*. The values of the d -compatibility degree $(-||-)_d$ are given by the components of the d -vectors. One remarkable property of the d -compatibility degree is that it uniquely determines how the set of cluster variables is grouped into clusters. As an application to d -vectors, we show how to express the d -compatibility degree and the d -vectors using the valuation pairing for full rank upper cluster algebras (see Theorem 5.1). As an application, in the full rank, primitive case, we prove that if M is a monomial in non-initial cluster variables, then M and its d -vector are uniquely determined by the numerator polynomial P_M of M (see Proposition 5.6).

Application to F -polynomials. Let B be an $n \times n$ skew-symmetrizable integer matrix. The F -polynomial $F_{k;t}^{B;t_0}$ associated with $(B, t_0; k, t)$ may be constructed by an explicit recursion or using the cluster algebra \mathcal{A} with principal coefficients [FZ07] associated with (B, t_0) : Let $A_{t_0} = (A_{1;t_0}, \dots, A_{n;t_0})$ be the initial cluster of \mathcal{A} and $A_{k;t}$ a cluster variable of \mathcal{A} . In this case, the cluster variable $A_{k;t}$ can be written as a Laurent polynomial in $\mathbb{Z}[Z_1, \dots, Z_n][A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$. Then the F -polynomial of $A_{k;t}$ is the specialization given by

$$F_{k;t}^{B;t_0}(Z_1, \dots, Z_n) := A_{k;t}|_{A_{1;t_0}=\dots=A_{n;t_0}=1} \in \mathbb{Z}[Z_1, \dots, Z_n].$$

When B and t_0 is clear, we just write $F_{k;t}^{B;t_0}$ for $F_{k;t}$.

Let $M = \prod_{i=1}^s M_i$ be a monomial in cluster variables, where each M_i is a cluster variable of \mathcal{A} . The F -polynomial of M is the polynomial $F_M := \prod_{i=1}^s F_{M_i}$, where F_{M_i} is the F -polynomial of the cluster variable M_i .

The F -polynomials are the non tropical ingredients of the canonical expressions [NZ12] for both, the cluster variables and the cluster Poisson variables. They are fundamental in the additive categorification of cluster algebras (see for example [CC06, DWZ10] and the surveys [BM06, GLS08, Kel10, Kel12, Pla18]) and in their link to Donaldson–Thomas theory (see for example [KS08, Nag13, Bri17]). It is known that F -polynomials enjoy many nice properties, for example, they have positive coefficients and constant term 1. We refer the readers to [LS15, DWZ10, GHKK18] for these results and to [GY20, Gyo21, FG19a, Fei19a, Fei19b, LP22] for some recent work on F -polynomials.

As an application to F -polynomials, we prove that if M is a monomial in non-initial cluster variables, then M is uniquely determined by its F -polynomial F_M

(see Theorem 6.1). We also prove that the F -polynomials of non-initial cluster variables are irreducible (see Theorem 6.2).

Application to cluster Poisson variables. As an application to combinatorics of cluster Poisson variables, we give several equivalent characterizations of when two cluster Poisson variables are equal (see Theorem 7.5). Recall that each mutation $t' = \mu_k(t)$ gives an \mathcal{A} -exchange pair $(A_{k;t}, A_{k;t'})$ of the upper cluster algebra \mathcal{U} and an \mathcal{X} -exchange pair $(X_{k;t}, X_{k;t'})$ of the cluster Poisson algebra \mathcal{X} .

As the first application of Theorem 7.5, we prove that the cluster Poisson variables of a cluster Poisson algebra \mathcal{X} parametrize the \mathcal{A} -exchange pairs of the upper cluster algebra \mathcal{U} of the same type as \mathcal{X} (see Theorem 7.6). This extends the corresponding result by Sherman-Bennett [SB19] from the finite type case to full generality.

As the second application of Theorem 7.5, we prove that the \mathcal{X} -seeds of \mathcal{X} whose Poisson clusters contain particular cluster Poisson variables form a connected subgraph of the exchange graph of \mathcal{X} (see Theorem 7.10). This is analogous to the result on the exchange graphs of cluster algebras given in [CL20], cf. also Theorem 2.17 of this paper.

1.3. Contents. This paper is organized as follows: In Section 2, some basic definitions, notations and known results are introduced. In Section 3, we introduce the valuation pairing $(-||-)_v$ on any upper cluster algebra and prove the local unique factorization property for full rank upper cluster algebras (see Theorem 3.7). In Sections 4, 5, 6, 7, we give the applications to d -vectors, F -polynomials, factoriality of upper cluster algebras and combinatorics of cluster Poisson variables. To be more precise:

In Section 4.1, we prove that a full rank upper cluster algebra \mathcal{U} with initial seed t_0 is factorial if and only if each exchange binomial of t_0 is irreducible in the corresponding polynomial ring (see Theorem 4.4). In particular, we show in Section 4.2 that full rank, primitive upper cluster algebras are factorial (see Theorem 4.8 (i)). Moreover, for these upper cluster algebras, we also show that the numerator polynomials of non-initial cluster variables are irreducible (see Theorem 4.8 (ii)). In Section 4.3, we give some examples of non-factorial upper cluster algebras. In Section 4.4, we prove the ray fish theorem, which states that any full rank, primitive upper cluster algebra \mathcal{U} can be written as the intersection of *two* Laurent polynomial rings (see Theorem 4.18).


In Section 5.1, we show how to express the d -compatibility degree and the d -vectors using the valuation pairing for full rank upper cluster algebras (see Theorem 5.1). In Section 5.2, we give a local factorization for cluster monomials (see Theorem 5.4). As an application, in the full rank, primitive case, we prove that if M is a monomial in non-initial cluster variables, then M and its d -vector are uniquely determined by the numerator polynomial P_M of M (see Proposition 5.6).

In Section 6.1, we prove that if M is a monomial in non-initial cluster variables, then M is uniquely determined by its F -polynomial F_M (see Theorem 6.1). In Section 6.2, we prove that the F -polynomials of non-initial cluster variables are irreducible (see Theorem 6.2).

In Section 7, we give several equivalent characterizations of when two cluster Poisson variables are equal (see Theorem 7.5). As the first application, we prove

that the cluster Poisson variables of a cluster Poisson algebra \mathcal{X} parametrize the \mathcal{A} -exchange pairs of the upper cluster algebra \mathcal{U} of the same type as \mathcal{X} (see Theorem 7.6). As the second application, we prove that the \mathcal{X} -seeds of \mathcal{X} whose Poisson clusters contain particular cluster Poisson variables form a connected subgraph of the exchange graph of \mathcal{X} (see Theorem 7.10).

The following diagram gives the logical dependence among the proofs of the main theorems in this paper.



1.4. Convention and assumption. Throughout this article, \mathbb{K} is a field of characteristic 0 or $\mathbb{K} = \mathbb{Z}$ and all upper cluster algebras are considered as algebras over $\mathbb{K}\mathbb{P}$, where \mathbb{P} is some abelian multiplicative group and $\mathbb{K}\mathbb{P}$ the corresponding group ring.

We always assume that the *exchange binomials* of upper cluster algebras in this article are *not invertible in $\mathbb{K}\mathbb{P}$* . Note that when $\mathbb{K} = \mathbb{Z}$, this condition is always satisfied. When \mathbb{K} is a field and we have a *trivial exchange relation*

$$A_k A'_k = P_k \in (\mathbb{K}\mathbb{P})^\times$$

in an upper cluster algebra \mathcal{U} , we can always freeze the cluster variable A_k and consider the upper cluster algebra \mathcal{U}^\dagger with smaller rank over $\mathbb{K}\mathbb{P}^\dagger = \mathbb{K}\mathbb{P}[A_k^\pm]$. Note that \mathcal{U}^\dagger and \mathcal{U} are isomorphic as \mathbb{K} -algebras.

Acknowledgements. The authors are very grateful to Luc Pirio, whose conjectures have inspired a good part of the results of this paper. P. Cao would like to thank Xiaofa Chen, Yu Wang and Yilin Wu for their helpful discussions during his stay in Paris. B. Keller is indebted to Lauren Williams and Melissa Sherman-Bennett for their interest and for stimulating conversations.

P. Cao thanks the Fondation Sciences Mathématiques de Paris (FSMP) for funding his postdoctoral stay (2019–2020) at the Université de Paris during which the first two authors began work on this paper. He also thanks the ERC (Grant No. 669655) for funding his postdoctoral stay (2021–present) at The Hebrew University of Jerusalem.

2. PRELIMINARIES

2.1. Basics on cluster algebras and cluster Poisson algebras. An $n \times n$ integer matrix B is *skew-symmetrizable* if there is an integer diagonal matrix S with strictly positive diagonal entries such that SB is skew-symmetric. Such an S is said to be a *skew-symmetrizer* of B .

Definition 2.1 (Matrix mutation). Let $\tilde{B} = \begin{pmatrix} B_{n \times n} \\ C_{m \times n} \end{pmatrix} = (b_{ij})$ be an $(n+m) \times n$ integer matrix with B skew-symmetrizable. The mutation $\mu_k(\tilde{B})$ of \tilde{B} at $k \in$

$\{1, \dots, n\}$ is the new matrix $\mu_k(\tilde{B}) = \tilde{B}' = \begin{pmatrix} B'_{n \times n} \\ C'_{m \times n} \end{pmatrix} = (b'_{ij})$ given by

$$b'_{ij} = \begin{cases} -b_{ij}, & \text{if } i = k \text{ or } j = k; \\ b_{ij} + \operatorname{sgn}(b_{ik}) \max(b_{ik} b_{kj}, 0), & \text{otherwise.} \end{cases}$$

It is not hard to check that the submatrix B' of \tilde{B}' is still skew-symmetrizable with the same skew-symmetrizer as B and μ_k is an involution.

Proposition 2.2. [BFZ05, Lemma 3.2] *Matrix mutations preserve the rank of \tilde{B} .*

Recall that $(\mathbb{P}, \oplus, \cdot)$ is a *semifield* if (\mathbb{P}, \cdot) is an abelian multiplicative group endowed with a binary operation of auxiliary addition \oplus which is commutative, associative and satisfies that the multiplication distributes over the auxiliary addition.

The *tropical semifield* $\mathbb{P} = \operatorname{Trop}(Z_1, \dots, Z_m)$ is the free (multiplicative) abelian group generated by Z_1, \dots, Z_m with auxiliary addition \oplus defined by

$$\prod_i Z_i^{a_i} \oplus \prod_i Z_i^{b_i} = \prod_i Z_i^{\min(a_i, b_i)}.$$

Let $\mathbb{Q}_{\text{sf}}(Z_1, \dots, Z_m)$ be the set of all non-zero rational functions in m independent variables Z_1, \dots, Z_m , which can be written as subtraction-free rational expressions in Z_1, \dots, Z_m . The set $\mathbb{Q}_{\text{sf}}(Z_1, \dots, Z_m)$ is a semifield with respect to the usual operations of multiplication and addition. It is called an *universal semifield*.

Definition 2.3 (\mathcal{X} -seed and cluster Poisson seed). (i) A (*labeled*) \mathcal{X} -seed over a semifield \mathbb{P} is a pair (B, X) , where

- $B = (b_{ij})$ is an $n \times n$ skew-symmetrizable integer matrix, called an *exchange matrix*;
- $X = (X_1, \dots, X_n)$ is an n -tuple of elements in \mathbb{P} . We call X the \mathcal{X} -cluster and X_1, \dots, X_n the \mathcal{X} -variables of (B, X) .

(ii) Let (B, X) be an \mathcal{X} -seed over a semifield \mathbb{P} . If $\mathbb{P} = \mathbb{Q}_{\text{sf}}(X_1, \dots, X_n)$, we call (B, X) a *cluster Poisson seed*, X the *Poisson cluster*, and X_1, \dots, X_n the *cluster Poisson variables* of (B, X) .

Throughout this article, we assume that \mathbb{K} is a field of characteristic 0 or $\mathbb{K} = \mathbb{Z}$ and we take the ambient field \mathcal{F} to be the field of rational functions in n independent variables with coefficients in $\mathbb{K}\mathbb{P}$.

Definition 2.4 (\mathcal{A} -seed). A (*labeled*) \mathcal{A} -seed over \mathbb{P} is a triple (B, X, A) , where

- (B, X) forms an \mathcal{X} -seed over \mathbb{P} ;
- $A = (A_1, \dots, A_n)$ is an n -tuple such that $\{A_1, \dots, A_n\}$ is a free generating set of \mathcal{F} over $\mathbb{K}\mathbb{P}$. We call A the *cluster* and A_1, \dots, A_n the *cluster variables* of (B, X, A) ;

Definition 2.5 (\mathcal{X} -seed mutation and \mathcal{X} -exchange pair). Let (B, X) be an \mathcal{X} -seed over \mathbb{P} . Define the \mathcal{X} -seed mutation of (B, X) at $k \in \{1, \dots, n\}$ as a new pair $\mu_k(B, X) = (B', X')$, where $B' = \mu_k(B)$ and $X' = (X'_1, \dots, X'_n)$ is given by

$$X'_i = \begin{cases} X_k^{-1}, & \text{if } i = k; \\ X_i X_k^{\max(b_{ki}, 0)} (1 \oplus X_k)^{-b_{ki}}, & \text{if } i \neq k. \end{cases}$$

In this case, (X_k, X'_k) is called an \mathcal{X} -exchange pair.

Definition 2.6 (\mathcal{A} -seed mutation and \mathcal{A} -exchange pair). Let (B, X, A) be an \mathcal{A} -seed over \mathbb{P} . Define the *mutation* of (B, X, A) at $k \in \{1, \dots, n\}$ as a new triple $\mu_k(B, X, A) = (B', X', A')$, where $(B', X') = \mu_k(B, X)$ and $A' = (A'_1, \dots, A'_n)$ is given by

$$A'_i = \begin{cases} A_i, & \text{if } i \neq k; \\ A_k^{-1} \left(\frac{X_k}{1 \oplus X_k} \prod_{b_{jk} > 0} A_j^{b_{jk}} + \frac{1}{1 \oplus X_k} \prod_{b_{jk} < 0} A_j^{-b_{jk}} \right), & \text{if } i = k. \end{cases}$$

In this case, (A_k, A'_k) is called an \mathcal{A} -*exchange pair*. The binomial

$$\frac{X_k}{1 \oplus X_k} \prod_{b_{jk} > 0} A_j^{b_{jk}} + \frac{1}{1 \oplus X_k} \prod_{b_{jk} < 0} A_j^{-b_{jk}}$$

is called the k -th *exchange binomial* of (B, X, A) .

It is not hard to check that each mutation μ_k maps a seed (\mathcal{X} -seed or cluster Poisson seed or \mathcal{A} -seed) to a new seed of the same type and that μ_k is an involution.

Let \mathbb{T}_n be the n -regular tree. Let us label the edges of \mathbb{T}_n by $1, \dots, n$ such that the n different edges adjacent to the same vertex of \mathbb{T}_n receive different labels.

Definition 2.7 (Seed pattern). A *seed pattern* \mathcal{S} over \mathbb{P} is an assignment of a seed Σ_t (\mathcal{X} -seed or cluster Poisson seed or \mathcal{A} -seed) to every vertex t of the n -regular tree \mathbb{T}_n such that $\Sigma_{t'} = \mu_k(\Sigma_t)$ for any edge $t \xrightarrow{k} t'$.

Note that the datum of an initial seed at a vertex t_0 determines a unique seed pattern. Now we give some symbols which are used in the sequel. We always write $B_t = (b_{ij}^t)$, $X_t = (X_{1;t}, \dots, X_{n;t})$ and $A_t = (A_{1;t}, \dots, A_{n;t})$. For simplicity, we will also use t to denote the seed at $t \in \mathbb{T}_n$.

Two (labeled) seeds are *equivalent* if they are the same up to relabeling.

Definition 2.8 (Exchange graph). Let \mathcal{S} be a seed pattern. The *exchange graph* $\mathbf{EG}(\mathcal{S})$ of \mathcal{S} is a graph whose vertices are in bijection with the seeds (up to equivalence) of \mathcal{S} and whose edges correspond to the seed mutations.

Proposition 2.9. [FZ07, Proposition 3.9] *Let \mathcal{S} be an \mathcal{A} -seed pattern. With each \mathcal{A} -seed $t = (B_t, X_t, A_t)$, we associate an \mathcal{X} -seed (B_t, \hat{X}_t) , where $\hat{X}_t = (\hat{X}_{1;t}, \dots, \hat{X}_{n;t})$ is given by $\hat{X}_{j;t} := X_{j;t} \prod_{i=1}^n A_{i;t}^{b_{ij}^t}$. Then $\hat{\mathcal{S}} = \{(B_t, \hat{X}_t)\}_{t \in \mathbb{T}_n}$ forms an \mathcal{X} -seed pattern.*

Now we recall some seed patterns (\mathcal{X} -seed patterns or \mathcal{A} -seed patterns) over particular semifields.

- A seed pattern \mathcal{S} over \mathbb{P} is said to be of *geometric type*, if \mathbb{P} is a tropical semifield, say $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$.
- A seed pattern \mathcal{S} over \mathbb{P} is said to be with *principal coefficients* at t_0 , if $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_n)$ and $X_{i;t_0} = Z_i$ for $i = 1, \dots, n$.
- A seed pattern \mathcal{S} over \mathbb{P} is said to be with *universal coefficient semifield*, if $\mathbb{P} = \mathbb{Q}_{\text{st}}(X_{1;t_0}, \dots, X_{n;t_0})$ for some seed t_0 of \mathcal{S} .

Definition 2.10 (Coefficient matrices and extended exchange matrices). Let \mathcal{S} be a seed pattern of geometric type with coefficient semifield $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$. We know that each \mathcal{X} -variable $X_{k;t}$ has the form

$$X_{k;t} = Z_1^{c_{1k}^t} \dots Z_m^{c_{mk}^t}.$$

The matrix $C_t := (c_{ij}^t)_{m \times n}$ is called the *coefficient matrix* at t and the $(n+m) \times n$ matrix $\tilde{B}_t = \begin{pmatrix} B_t \\ C_t \end{pmatrix}$ is called the *extended exchange matrix* at t .

Proposition 2.11. [FZ02, Proposition 5.8] *Let \mathcal{S} be a seed pattern of geometric type. Then for any edge $t \xrightarrow{k} t'$ in \mathbb{T}_n , we have $\tilde{B}_{t'} = \mu_k(\tilde{B}_t)$, where \tilde{B}_t and $\tilde{B}_{t'}$ are the extended exchange matrices at t and t' .*

Definition 2.12 (Cluster Poisson algebra). Let \mathcal{S}_{uc} be an \mathcal{X} -seed pattern with universal coefficient semifield. The *cluster Poisson algebra* $\mathcal{X} = \mathcal{X}(\mathcal{S}_{\text{uc}})$ associated with \mathcal{S}_{uc} is the intersection

$$\mathcal{X} = \bigcap_{t \in \mathbb{T}_n} \mathcal{L}(t),$$

where $\mathcal{L}(t) = \mathbb{K}[X_{1;t}^{\pm 1}, \dots, X_{n;t}^{\pm 1}]$.

Definition 2.13 (Cluster algebra and upper cluster algebra). Let \mathcal{S} be an \mathcal{A} -seed pattern over a semifield \mathbb{P} .

(i) The *cluster algebra* $\mathcal{A} = \mathcal{A}(\mathcal{S})$ associated with \mathcal{S} is the $\mathbb{K}\mathbb{P}$ -subalgebra of \mathcal{F} generated by the cluster variables of \mathcal{S} , namely, $\mathcal{A} = \mathbb{K}\mathbb{P}[A_{1;t}, \dots, A_{n;t} | t \in \mathbb{T}_n]$.

(ii) The *upper cluster algebra* $\mathcal{U} = \mathcal{U}(\mathcal{S})$ associated with \mathcal{S} is the intersection

$$\mathcal{U} = \bigcap_{t \in \mathbb{T}_n} \mathcal{L}(t),$$

where $\mathcal{L}(t) = \mathbb{K}\mathbb{P}[A_{1;t}^{\pm 1}, \dots, A_{n;t}^{\pm 1}]$.

Since cluster Poisson algebras and (upper) cluster algebras are defined from seed patterns, we can talk about the exchange graphs of these algebras. We can also talk about the (upper) cluster algebras of geometric type, with principal coefficients and with universal coefficient semifield.

Theorem 2.14 (Laurent phenomenon and positivity). *Let \mathcal{A} be a cluster algebra with coefficient semifield \mathbb{P} and initial seed t_0 . The following statements hold.*

- (i) [FZ02] *Each cluster variable $A_{k;t}$ of \mathcal{A} can be written as a Laurent polynomial in $\mathbb{Z}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$.*
- (ii) [GHKK18] *The coefficients in above Laurent polynomial are positive.*
- (iii) [FZ03] *If $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$, then each cluster variable $A_{k;t}$ of \mathcal{A} can be written as a Laurent polynomial in $\mathbb{Z}[Z_1, \dots, Z_m][A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$.*

A geometric upper cluster algebra with initial seed t_0 is called a *full rank upper cluster algebra* if its initial extended exchange matrix \tilde{B}_{t_0} has full rank.

Remark 2.15. For a full rank upper cluster algebra \mathcal{U} , we know that every extended exchange matrix \tilde{B}_t of \mathcal{U} has full rank, since mutations preserve the rank of the extended exchange matrices (see Proposition 2.2).

Theorem 2.16 (Starfish theorem). *Let \mathcal{U} be a full rank upper cluster algebra and t_0 a seed of \mathcal{U} . Then we have*

$$\mathcal{U} = \bigcap_{i=0}^n \mathcal{L}(t_i),$$

where $t_j = \mu_j(t_0)$ for $j = 1, \dots, n$ and $\mathcal{L}(t_i) = \mathbb{K}\mathbb{P}[A_{1;t_i}^{\pm 1}, \dots, A_{n;t_i}^{\pm 1}]$ for $i = 0, 1, \dots, n$.

Proof. For the case $\mathbb{K} = \mathbb{Z}$, the proof can be found in [BFZ05] and the proof there also works for the case where \mathbb{K} is a field of characteristic 0. \square

Theorem 2.17. [CL20] *Let \mathcal{A} be a cluster algebra with coefficient semifield \mathbb{P} . Then the seeds of \mathcal{A} whose clusters contain particular cluster variables form a connected subgraph of the exchange graph of \mathcal{A} .*

The following result is a direct corollary.

Corollary 2.18. [CL20] *Let \mathcal{A} be a cluster algebra with coefficient semifield \mathbb{P} and t_1, t_2 two seeds of \mathcal{A} . If t_1 and t_2 have at least $n - 1$ common cluster variables, then in the exchange graph of \mathcal{A} either t_1 and t_2 represent the same vertex or there is an edge between t_1 and t_2 .*

Recall that a cluster algebra is *acyclic* if it has an acyclic seed: a seed with an exchange matrix $B = (b_{ij})$, such that there is no sequence of indices $i_1, \dots, i_{\ell+1} \in \{1, \dots, n\}$ with $b_{i_j, i_{j+1}} > 0$ for all j and $i_{\ell+1} = i_1$.

Proposition 2.19. [BFZ05, Mul14] *Let \mathcal{A} be an acyclic cluster algebra and \mathcal{U} the corresponding upper cluster algebra. Then $\mathcal{A} = \mathcal{U}$.*

2.2. d -vectors, f -vectors, g -vectors, c -vectors and F -polynomials. Let M be any non-zero element in $\mathcal{L}(t_0) = \mathbb{K}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$. Then M has the form

$$M = \frac{P_M(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{d_1} \cdots A_{n;t_0}^{d_n}},$$

where $P_M \in \mathbb{K}\mathbb{P}[A_{1;t_0}, \dots, A_{n;t_0}]$ with $A_{j;t_0} \nmid P_M$ for $j = 1, \dots, n$. The vector

$$\mathbf{d}^{t_0}(M) = (d_1, \dots, d_n)^T$$

is called the d -vector of M with respect to $A_{t_0} = (A_{1;t_0}, \dots, A_{n;t_0})$ and the polynomial P_M is called the *numerator polynomial* of M with respect to A_{t_0} .

Remark 2.20. Notice that for any $0 \neq M, N \in \mathcal{L}(t_0)$, we have

$$\mathbf{d}^{t_0}(M \cdot N) = \mathbf{d}^{t_0}(M) + \mathbf{d}^{t_0}(N).$$

Definition 2.21 (F -polynomial). Let B be an $n \times n$ skew-symmetrizable integer matrix and \mathcal{A} a principal coefficient cluster algebra at t_0 with $B_{t_0} = B$. Let $A_{k;t}$ be a cluster variable of \mathcal{A} , which can be written as a Laurent polynomial in

$$\mathbb{Z}[Z_1, \dots, Z_n][A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}],$$

by the Laurent phenomenon. The polynomial

$$F_{k;t}^{B;t_0}(Z_1, \dots, Z_n) := A_{k;t} \big|_{A_{1;t_0} = \dots = A_{n;t_0} = 1} \in \mathbb{Z}[Z_1, \dots, Z_n],$$

which only depends on $(B, t_0; k, t)$, is called an F -polynomial of B . Note that we always view $F_{k;t}^{B;t_0}$ as a polynomial in new variables Y_1, \dots, Y_n in the sequel.

Proposition 2.22. [FZ07, Proposition 5.2] *The F -polynomial $F_{k;t}^{B;t_0}(Y_1, \dots, Y_n)$ is not divisible by any Y_j .*

Let B be an $n \times n$ skew-symmetrizable matrix and t_0 a vertex of \mathbb{T}_n . Now we define two families of integer matrices $\{D_t^{B;t_0}\}_{t \in \mathbb{T}_n}$ and $\{F_t^{B;t_0}\}_{t \in \mathbb{T}_n}$.

Definition 2.23 (*D*-matrices and *F*-matrices). Let B be an $n \times n$ skew-symmetrizable matrix and t_0 a vertex of \mathbb{T}_n .

(i) [FZ07] The matrix $D_t^{B;t_0} = (d_{ij;t}^{B;t_0})$ is uniquely determined by the initial condition $B_{t_0} = B$ and $D_{t_0}^{B;t_0} = -I_n$, together with the following recurrence relations:

$$d_{ij;t'}^{B;t_0} = \begin{cases} d_{ij;t}^{B;t_0}, & \text{if } j \neq k; \\ -d_{ik;t}^{B;t_0} + \max\left\{ \sum_{b_{lk}^t > 0} d_{il;t}^{B;t_0} b_{lk}^t, \sum_{b_{lk}^t < 0} -d_{il;t}^{B;t_0} b_{lk}^t \right\}, & \text{if } j = k, \end{cases}$$

for any edge $t \xrightarrow{k} t'$ in \mathbb{T}_n . The matrices $\{D_t^{B;t_0}\}_{t \in \mathbb{T}_n}$ are called the *D*-matrices of B .

(ii) [FG19b] Let $F_{k;t}^{B;t_0}(Z_1, \dots, Z_n)$ be the *F*-polynomial given in Definition 2.21, and $f_{ik;t}^{B;t_0}$ be the maximal exponent of Z_i appearing in $F_{k;t}^{B;t_0}$. The vector

$$\mathbf{f}_{k;t}^{B;t_0} = (f_{1k;t}^{B;t_0}, \dots, f_{nk;t}^{B;t_0})^\top$$

is called an *f*-vector of B and the matrix $F_t^{B;t_0} = (f_{ij;t}^{B;t_0})_{n \times n}$ is called an *F*-matrix of B .

Remark 2.24. Let \mathcal{A} be a cluster algebra with initial exchange matrix B at t_0 . Then the k -th column vector $\mathbf{d}_{k;t}^{B;t_0}$ of $D_t^{B;t_0}$ is exactly the d -vector $\mathbf{d}^{t_0}(A_{k;t})$ of $A_{k;t}$ with respect to A_{t_0} , by [FZ07, (7.7)].

Proposition 2.25. *Let \mathcal{U} be an upper cluster algebra and M a non-zero element in \mathcal{U} . Let t_0, t_1 be two seeds of \mathcal{U} with $t_1 = \mu_k(t_0)$. Let $\mathbf{d} = (d_1, \dots, d_n)^\top$ and $\mathbf{d}' = (d'_1, \dots, d'_n)^\top$ be the d -vectors of M with respect to t_0 and t_1 respectively. Then $d_i = d'_i$ for any $i \neq k$.*

Proof. The proof is the same as that of [RS18, Proposition 2.5]. \square

Theorem 2.26 (Canonical expressions). *Let B be an $n \times n$ skew-symmetrizable matrix and*

$$\{F_{1;t}^{B;t_0}, \dots, F_{n;t}^{B;t_0}\}_{t \in \mathbb{T}_n}$$

*the *F*-polynomials given in Definition 2.21.*

(i) [FZ07] *Let \mathcal{A} be a cluster algebra with principal coefficients at t_0 and with $B_{t_0} = B$. Then for any cluster variable $A_{k;t}$ of \mathcal{A} , there exists a unique vector $\mathbf{g}_{k;t}^{B;t_0} = (g_{1k;t}^{B;t_0}, \dots, g_{nk;t}^{B;t_0})^\top \in \mathbb{Z}^n$ such that*

$$A_{k;t} = A_{t_0}^{\mathbf{g}_{k;t}^{B;t_0}} \cdot F_{k;t}^{B;t_0}(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}),$$

$$\text{where } A_{t_0}^{\mathbf{g}_{k;t}^{B;t_0}} = \prod_{i=1}^n A_{i;t}^{g_{ik;t}^{B;t_0}}, \hat{X}_{j;t_0} = X_{j;t_0} \prod_{i=1}^n A_{i;t_0}^{b_{ij}^{t_0}} = Z_j \prod_{i=1}^n A_{i;t_0}^{b_{ij}^{t_0}}.$$

(ii) [NZ12] *Let \mathcal{X} be the cluster Poisson algebra with $B_{t_0} = B$. Then for any cluster Poisson variable $X_{k;t}$ of \mathcal{X} , there exists a unique vector $\mathbf{c}_{k;t}^{B;t_0} = (c_{1k;t}^{B;t_0}, \dots, c_{nk;t}^{B;t_0})^\top \in \mathbb{Z}^n$ such that*

$$X_{k;t} = X_{t_0}^{\mathbf{c}_{k;t}^{B;t_0}} \cdot \prod_{i=1}^n (F_{i;t}^{B;t_0}(X_{1;t_0}, \dots, X_{n;t_0}))^{b_{ik}^t}.$$

Definition 2.27 (*g*-vectors and *c*-vectors). Keep the notations of Theorem 2.26.

(i) The vector $\mathbf{g}_{k;t}^{B;t_0} = (g_{1k;t}^{B;t_0}, \dots, g_{nk;t}^{B;t_0})^T \in \mathbb{Z}^n$, which only depends on $(B, t_0; k, t)$, is called a *g*-vector of B . The matrix $G_t^{B;t_0} = (\mathbf{g}_{ij;t}^{B;t_0})_{n \times n}$ is called a *G*-matrix of B at t .

(ii) The vector $\mathbf{c}_{k;t}^{B;t_0} = (c_{1k;t}^{B;t_0}, \dots, c_{nk;t}^{B;t_0})^T \in \mathbb{Z}^n$, which only depends on $(B, t_0; k, t)$, is called a *c*-vector of B . The matrix $C_t^{B;t_0} = (\mathbf{c}_{ij;t}^{B;t_0})_{n \times n}$ is called a *C*-matrix of B at t .

Theorem 2.28. [NZ12, CL18] *Let B be an $n \times n$ skew-symmetrizable matrix and S a skew-symmetrizer of B . Then for any vertices $t_0, t \in \mathbb{T}_n$, we have*

$$SC_t^{B;t_0} S^{-1} (G_t^{B;t_0})^T = I_n.$$

2.3. Compatibility degrees on the set of cluster variables. In this subsection, we recall that the *d*-compatibility degree given in [CL20] and *f*-compatibility degree given in [FG19a].

Definition 2.29 (Compatibility degrees). Let \mathcal{A} be a cluster algebra and \mathbb{A} the set of cluster variables of \mathcal{A} .

(i) [CL20] The *d*-compatibility degree $(-||-)_d : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{Z}_{\geq -1}$ is defined by

$$(A_{i;t_0} || A_{j;t})_d := d_{ij;t}^{B_{t_0};t_0},$$

where $d_{ij;t}^{B_{t_0};t_0}$ is the (i, j) -entry of the *D*-matrix $D_t^{B_{t_0};t_0}$.

(ii) [FG19a] The *f*-compatibility degree $(-||-)_f : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{N}$ is defined by

$$(A_{i;t_0} || A_{j;t})_f := f_{ij;t}^{B_{t_0};t_0},$$

where $f_{ij;t}^{B_{t_0};t_0}$ is the (i, j) -entry of the *F*-matrix $F_t^{B_{t_0};t_0}$.

Now we summarize some properties of the *d*-compatibility degree $(-||-)_d$ and the *f*-compatibility degree $(-||-)_f$. We mainly refer to [CL20] and [FG19a].

Proposition 2.30. [CL20, FG19a] *The *d*-compatibility degree $(-||-)_d$ and the *f*-compatibility degree $(-||-)_f$ have the following properties.*

- (1) *They are well-defined.*
- (2) *The following statements are equivalent.*
 - (i) *There exists a cluster $A_{t'}$ containing both $A_{i;t_0}$ and $A_{j;t}$;*
 - (ii) $(A_{i;t_0} || A_{j;t})_d \leq 0$;
 - (iii) $(A_{j;t} || A_{i;t_0})_d \leq 0$;
 - (iv) $(A_{i;t_0} || A_{j;t})_f = 0$;
 - (v) $(A_{j;t} || A_{i;t_0})_f = 0$.
- (3) *The following statements are equivalent.*
 - (i) *There exists no cluster $A_{t'}$ containing both $A_{i;t_0}$ and $A_{j;t}$;*
 - (ii) $(A_{i;t_0} || A_{j;t})_d > 0$;
 - (iii) $(A_{j;t} || A_{i;t_0})_d > 0$;
 - (iv) $(A_{i;t_0} || A_{j;t})_f > 0$;
 - (v) $(A_{j;t} || A_{i;t_0})_f > 0$.
- (4) *The following statements are equivalent.*
 - (i) $A_{j;t} = A_{i;t_0}$;
 - (ii) $(A_{i;t_0} || A_{j;t})_d < 0$;
 - (iii) $(A_{j;t} || A_{i;t_0})_d < 0$;

- (iv) $(A_{i;t_0} || A_{j;t})_d = -1$;
- (v) $(A_{j;t} || A_{i;t_0})_d = -1$.
- (5) *The following statements are equivalent.*
 - (i) $A_{i;t_0} \neq A_{j;t}$ and there exists a cluster $A_{t'}$ containing both $A_{i;t_0}$ and $A_{j;t}$;
 - (ii) $(A_{i;t_0} || A_{j;t})_d = 0$;
 - (iii) $(A_{j;t} || A_{i;t_0})_d = 0$;

Remark 2.31. (i) Roughly speaking, the integer $(A_{i;t_0} || A_{j;t})_d$ is defined to be the i -th component of the d -vector of $A_{j;t}$ with respect to A_{t_0} . By Proposition 2.25 and Theorem 2.17, the d -compatibility degree $(-||-)_d$ is actually a well-defined function on $\mathbb{A} \times (\mathcal{U} \setminus \{0\})$, where \mathcal{U} is the corresponding upper cluster algebra.

(ii) By Remark 2.20, for any $0 \neq M, N \in \mathcal{U}$, we have

$$(A_{i;t_0} || M \cdot N)_d = (A_{i;t_0} || M)_d + (A_{i;t_0} || N)_d.$$

Recall that a *cluster monomial* in seed t is a monomial in cluster variables in t .

Corollary 2.32. Let \mathcal{A} be a cluster algebra with coefficient semifield \mathbb{P} and $A_{k;t_0}$ a cluster variable of \mathcal{A} . Let t be a seed of \mathcal{A} and $M = A_{1;t}^{c_1} \cdots A_{n;t}^{c_n}$ a cluster monomial in t . Denote $I = \{i | c_i > 0\}$. The following statements hold.

- (i) If $(A_{k;t_0} || M)_d < 0$, then there exists $j_0 \in I$ such that $A_{k;t_0} = A_{j_0;t}$. Namely, $A_{k;t_0}$ appears in M .
- (ii) If $(A_{k;t_0} || M)_d \leq 0$, then $(A_{k;t_0} || A_{i;t})_d \leq 0$ for any $i \in I$.

Proof. (i) By $c_1, \dots, c_n \geq 0$ and $(A_{k;t_0} || M)_d < 0$, we have

$$(A_{k;t_0} || M)_d = \sum_{i=1}^n c_i (A_{k;t_0} || A_{i;t})_d = \sum_{i \in I} c_i (A_{k;t_0} || A_{i;t})_d < 0.$$

So there must exist some $j_0 \in I$ such that $(A_{k;t_0} || A_{j_0;t})_d < 0$. Then by Proposition 2.30 (4), we get $A_{k;t_0} = A_{j_0;t}$.

(ii) Assume by contradiction there exists some $i_0 \in I$ such that

$$(A_{k;t_0} || A_{i_0;t})_d > 0.$$

Then by $(A_{k;t_0} || M)_d = \sum_{i \in I} c_i (A_{k;t_0} || A_{i;t})_d \leq 0$, there must exist some $i_1 \in I$ such that $(A_{k;t_0} || A_{i_1;t})_d < 0$. So we get $A_{k;t_0} = A_{i_1;t}$, by Proposition 2.30 (4). Thus $A_{i_0;t}$ and $A_{k;t_0} = A_{i_1;t}$ are in the same cluster. Then by Proposition 2.30 (2), we get $(A_{k;t_0} || A_{i_0;t})_d \leq 0$. This contradicts $(A_{k;t_0} || A_{i_0;t})_d > 0$. So $(A_{k;t_0} || A_{i;t})_d \leq 0$ for any $i \in I$. \square

Let \mathcal{A} be a cluster algebra and \mathbb{A} the set of cluster variables of \mathcal{A} . We say that two cluster variables $A_{i;t_0}$ and $A_{j;t}$ of \mathcal{A} are *compatible* if there exists a cluster $A_{t'}$ of \mathcal{A} containing both $A_{i;t_0}$ and $A_{j;t}$, which is equivalent to $(A_{i;t_0} || A_{j;t})_d \leq 0$, by Proposition 2.30 (2). A subset U of \mathbb{A} is called a *compatible set* if any two cluster variables in U are compatible.

Theorem 2.33. [CL20] *Let \mathcal{A} be a cluster algebra and \mathbb{A} the set of cluster variables of \mathcal{A} . Then*

- (i) *a subset U of \mathbb{A} is a compatible set if and only if it is a subset of some cluster of \mathcal{A} ;*
- (ii) *a subset U of \mathbb{A} is a maximal compatible set if and only if it is a cluster of \mathcal{A} .*

2.4. Unique factorization domains. Recall that an integral domain is a non zero commutative ring without zero divisors. From now on, we always assume that R is an integral domain and we denote by R^\times the set of invertible elements in R .

Two non-zero elements $r_1, r_2 \in R$ are *associate* if there exists some invertible element $s \in R^\times$ such that $r_2 = sr_1$. A non-invertible element r in R is *irreducible*, if any factorization $r = r_1r_2$ with $r_1, r_2 \in R$ implies that either r_1 or r_2 belongs to R^\times . A non-invertible element r in R is *prime*, if whenever $r \mid r_1r_2$ for some $r_1, r_2 \in R$, then $r \mid r_1$ or $r \mid r_2$. Every prime element is irreducible, but the converse is not true in general.

Definition 2.34. An integral domain R is *factorial* if the following hold:

- (i) Every non-zero non-invertible element $r \in R$ can be written as a product $r = a_1 \cdots a_s$ of irreducible elements $a_i \in R$.
- (ii) If $a_1 \cdots a_s = b_1 \cdots b_t$ with $a_i, b_j \in R$ irreducible for all i and j , then $s = t$ and there is a bijection $\sigma : \{1, \dots, s\} \rightarrow \{1, \dots, s\}$ such that a_i and $b_{\sigma(i)}$ are associate for all $1 \leq i \leq s$.

Factorial domains are often also called *unique factorization domains*.

It is easy to see that in a factorial domain, all irreducible elements are prime.

Theorem 2.35. Let \mathcal{A} be a geometric cluster algebra with coefficient semifield $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$ and \mathcal{U} the corresponding upper cluster algebra. Then we have

- (i) [GLS13, Theorem 1.3]
 - (a) $\mathcal{A}^\times = \{\lambda Z_1^{c_1} \cdots Z_m^{c_m} \mid \lambda \in \mathbb{K}^\times, c_1, \dots, c_m \in \mathbb{Z}\}$.
 - (b) any cluster variable is irreducible in \mathcal{A} .
- (ii) (a) $\mathcal{U}^\times = \{\lambda Z_1^{c_1} \cdots Z_m^{c_m} \mid \lambda \in \mathbb{K}^\times, c_1, \dots, c_m \in \mathbb{Z}\}$.
 - (b) any cluster variable is irreducible in \mathcal{U} .

Proof. The proof of (i) can be found in [GLS13] and the proof of (ii) is the same as that of (i). \square

3. THE VALUATION PAIRING ON AN UPPER CLUSTER ALGEBRA

In this section, we introduce the valuation pairing $(-||-)_v$ on an upper cluster algebra and prove the local unique factorization property for full rank upper cluster algebras.

Definition 3.1 (Valuation pairing). Let \mathcal{U} be an upper cluster algebra and \mathbb{A} the set of cluster variables of \mathcal{U} . The pairing

$$\begin{aligned} (-||-)_v : \mathbb{A} \times \mathcal{U} &\rightarrow \mathbb{N} \cup \{\infty\} \\ (A_{k;t}, M) &\mapsto \max\{s \in \mathbb{N} \mid M/A_{k;t}^s \in \mathcal{U}\} \end{aligned}$$

is called the *valuation pairing* on \mathcal{U} .

It is easy to see that $(A_{k;t}||M)_v = 0$ if and only if $M/A_{k;t} \notin \mathcal{U}$.

Example 3.2. Let \mathcal{U} be an upper cluster algebra with a seed t and M a cluster variable in t . Then

$$(A_{k;t}||M)_v = \begin{cases} 1, & \text{if } M = A_{k;t}; \\ 0, & \text{if } M = A_{j;t} \text{ and } j \neq k. \end{cases}$$

Now we summarize some useful and easy facts on the valuation pairing in the following proposition.

Proposition 3.3. *Let \mathcal{U} be an upper cluster algebra and t_0 a seed of \mathcal{U} . Let M and L be two elements in \mathcal{U} . The following statements hold.*

- (i) *If $M/A_{k;t_0}^m \notin \mathcal{U}$ for some $m \in \mathbb{Z}$, then we also have $M/A_{k;t_0}^{m+1} \notin \mathcal{U}$.*
- (ii) *If $M \neq 0$, then there exists $m > 0$ such that*

$$M/A_{k;t_0}^m \notin \mathcal{U},$$

In particular, $(A_{k;t_0} || M)_v < \infty$.

- (iii) *$(A_{k;t_0} || M)_v = \infty$ if and only if $M = 0$.*
- (iv) *$(A_{k;t_0} || M + L)_v \geq \min\{(A_{k;t_0} || M)_v, (A_{k;t_0} || L)_v\}$.*
- (v) *For any $s \geq 0$, we have $(A_{k;t_0} || A_{k;t_0}^s \cdot M)_v = s + (A_{k;t_0} || M)_v$.*

Proof. (i) Assume by contradiction that $M/A_{k;t_0}^{m+1} \in \mathcal{U}$, then

$$M/A_{k;t_0}^m = A_{k;t_0} \cdot (M/A_{k;t_0}^{m+1}) \in \mathcal{U}.$$

This is a contradiction. So $M/A_{k;t_0}^{m+1} \notin \mathcal{U}$.

(ii) Without loss of generality, we can assume that $k = n$. As an element in

$$\mathcal{L}(t_0) = \mathbb{K}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}] = \mathbb{K}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n-1;t_0}^{\pm 1}][A_{n;t_0}^{\pm 1}],$$

M can be written as

$$M = \sum_{s \in \mathbb{Z}} A_{n;t_0}^s L_s,$$

where L_s is a Laurent polynomial in $\mathbb{K}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n-1;t_0}^{\pm 1}]$, i.e., L_s does not contain any $A_{n;t_0}^{\pm 1}$. By $M \neq 0$, we know that there exists some s_0 such that $L_{s_0} \neq 0$. Let $t_1 = \mu_n(t_0)$ and thus $t_0 = \mu_n(t_1)$. Applying the exchange relation, we have

$$A_{i;t_0} = A_{i;t_1} \quad \text{and} \quad A_{n;t_0} = A_{n;t_1}^{-1} P$$

where $i = 1, \dots, n-1$ and $P \in \mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}]$ is the n -th exchange binomial of the seed t_1 . Now each L_s can be also viewed as a Laurent polynomial in $\mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}]$. By our assumption in Subsection 1.4, we know that P is not invertible in $\mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}]$. Then by the fact that $\mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}]$ is factorial and L_{s_0} belongs to $\mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}]$, we know that there must exist some m_0 large enough such that

$$L_{s_0}/P^{m_0} \notin \mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}].$$

Set $m = m_0 + s_0$. Then we know that

$$M/A_{n;t_0}^m = \sum_{s \in \mathbb{Z}} A_{n;t_0}^{s-m} L_s = \sum_{s \in \mathbb{Z}} A_{n;t_1}^{m-s} (L_s/P^{m-s})$$

cannot be a Laurent polynomial in $\mathbb{K}\mathbb{P}[A_{1;t_1}^{\pm 1}, \dots, A_{n-1;t_1}^{\pm 1}]$. So $M/A_{n;t_0}^m$ is not in \mathcal{U} . Then by (i), we know that $(A_{k;t_0} || M)_v < m < \infty$.

(iii) follows from (ii) and the fact $(A_{k;t_0} || 0)_v = \infty$.

(iv) and (v) follow from the definition of the valuation pairing. \square

Definition 3.4 (Local factorization). Let \mathcal{U} be an upper cluster algebra and t a seed of \mathcal{U} . Let $M = N \cdot L$ be a factorization in \mathcal{U} . If N is a cluster monomial in t and $L \in \mathcal{U}$ satisfies $(A_{k;t} || L)_v = 0$, that is, $L/A_{k;t} \notin \mathcal{U}$ for $k = 1, \dots, n$, we call $M = N \cdot L$ a *local factorization* of M with respect to t .

Proposition 3.5 (Existence of local factorization). *Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and t_0 a seed of \mathcal{U} . Then any $0 \neq M \in \mathcal{U}$ admits a local factorization with respect to t_0 .*

Proof. We first define a cluster monomial $N = A_{1;t_0}^{m_1} \cdots A_{n;t_0}^{m_n}$ in t_0 . Let $m_1 := (A_{1;t_0} \| M)_v$ and m_2, \dots, m_n are defined by induction. If m_k has been defined, m_{k+1} is defined as follows:

$$m_{k+1} := (A_{k+1;t_0} \| \frac{M}{A_{1;t_0}^{m_1} \cdots A_{k;t_0}^{m_k}})_v.$$

Thus the non-negative integers m_1, \dots, m_n are defined. So we get a cluster monomial $N = A_{1;t_0}^{m_1} \cdots A_{n;t_0}^{m_n}$ in t_0 and we know that $L := M/N \in \mathcal{U}$.

Now we show that $L = M/N$ satisfies $(A_{k;t_0} \| L)_v = 0$, that is, $L/A_{k;t_0} \notin \mathcal{U}$ for $k = 1, \dots, n$. Assume by contradiction that there exists some $k \in \{1, \dots, n\}$ such that

$$L/A_{k;t_0} = \frac{M}{A_{1;t_0}^{m_1} \cdots A_{k-1;t_0}^{m_{k-1}} A_{k;t_0}^{m_k+1} A_{k+1;t_0}^{m_{k+1}} \cdots A_{n;t_0}^{m_n}} \in \mathcal{U}.$$

Thus we get

$$\frac{M}{A_{1;t_0}^{m_1} \cdots A_{k-1;t_0}^{m_{k-1}} A_{k;t_0}^{m_k+1}} = A_{k+1;t_0}^{m_{k+1}} \cdots A_{n;t_0}^{m_n} (L/A_{k;t_0}) \in \mathcal{U},$$

which contradicts the choice of m_k . So $L/A_{k;t_0} \notin \mathcal{U}$ for $k = 1, \dots, n$. Thus $M = N \cdot L$ is a local factorization of M with respect to t_0 . This completes the proof. \square

Lemma 3.6 (Reduction Lemma). *Let \mathcal{U} be a full rank upper cluster algebra and t_0 a seed of \mathcal{U} . Let L be a non-zero element in \mathcal{U} and $N = A_{1;t_0}^{m_1} \cdots A_{n;t_0}^{m_n}$ a cluster monomial in t_0 . Then the following statements hold.*

(i) $L/A_{k;t_0}^s \in \mathcal{U}$ if and only if $L/A_{k;t_0}^s \in \mathcal{L}(t_k)$, where $s \in \mathbb{N}$, $t_k = \mu_k(t_0)$ and

$$\mathcal{L}(t_k) = \mathbb{K}\mathbb{P}[A_{1;t_k}^{\pm 1}, \dots, A_{n;t_k}^{\pm 1}].$$

(ii) $(A_{k;t_0} \| L)_v = \max\{s \in \mathbb{N} \mid L/A_{k;t_0}^s \in \mathcal{L}(t_k)\}$.

(iii) $(A_{k;t_0} \| N \cdot L)_v = m_k + (A_{k;t_0} \| L)_v$.

Proof. (i) By the starfish Theorem 2.16, we know that $\mathcal{U} = \bigcap_{i=0}^n \mathcal{L}(t_i)$. Since $A_{k;t_0}$ is a cluster variable in $\mathcal{L}(t_i)$ for any $i \in \{0, 1, \dots, n\} \setminus \{k\}$, we know that $L/A_{k;t_0}^s \in \mathcal{L}(t_i)$ for any $i \in \{0, 1, \dots, n\} \setminus \{k\}$. Thus we get that $L/A_{k;t_0}^s \in \mathcal{U}$ if and only if $L/A_{k;t_0}^s \in \mathcal{L}(t_k)$.

(ii) follows from (i) and the definition of the valuation pairing.

(iii) Without loss of generality, we assume $k = 1$. Since $A_{2;t_0}^{m_2} \cdots A_{n;t_0}^{m_n}$ is invertible in $\mathcal{L}(t_1)$ and by (ii), we have

$$(A_{1;t_0} \| N \cdot L)_v = (A_{1;t_0} \| \frac{N \cdot L}{A_{2;t_0}^{m_2} \cdots A_{n;t_0}^{m_n}})_v = (A_{1;t_0} \| A_{1;t_0}^{m_1} \cdot L)_v.$$

Then by Proposition 3.3 (v), we get

$$(A_{1;t_0} \| A_{1;t_0}^{m_1} \cdot L)_v = m_1 + (A_{1;t_0} \| L)_v.$$

This completes the proof. \square

Theorem 3.7 (Local unique factorization property). *Let \mathcal{U} be a full rank upper cluster algebra and t_0 a seed of \mathcal{U} . Then any $0 \neq M \in \mathcal{U}$ admits a unique local factorization with respect to t_0 .*

Proof. The existence of local factorization of M with respect to t_0 is known from Proposition 3.5. Now we show the uniqueness. Let $M = N \cdot L$ be a local factorization of M with respect to t_0 , where $N = A_{1;t_0}^{m_1} \cdots A_{n;t_0}^{m_n}$ is a cluster monomial in t_0 and $L \in \mathcal{U}$ satisfies $(A_{k;t_0} || L)_v = 0$ for $k = 1, \dots, n$. By the Reduction Lemma 3.6 (iii), we have

$$(A_{k;t_0} || M)_v = (A_{k;t_0} || N \cdot L)_v = m_k + (A_{k;t_0} || L)_v = m_k.$$

So $m_k = (A_{k;t_0} || M)_v$ is uniquely determined by M . Namely, N is uniquely determined by M . This completes the proof. \square

Now we give a counter-example to the local unique factorization property in the case where \tilde{B}_{t_0} is *not full rank*.

Example 3.8. Let \mathcal{U} be a geometric upper cluster algebra with initial extended exchange matrix \tilde{B}_{t_0} given by

$$\tilde{B}_{t_0} = B_{t_0} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Clearly, \tilde{B}_{t_0} is not full rank. Denote $t_1 = \mu_1(t_0)$ and $t_3 = \mu_3(t_0)$. Applying the exchange relations, we have the following equality.

$$(3.1) \quad A_{1;t_0} A_{1;t_1} = A_{2;t_0} + 1 = A_{3;t_0} A_{3;t_3}.$$

It is easy to check that both $A_{1;t_1}$ and $A_{3;t_3}$ are not divisible by any cluster variable in t_0 (alternatively, one can also refer to Theorem 5.4 (ii)). So the equality (3.1) gives two different local factorizations of $A_{2;t_0} + 1$ with respect to the initial seed t_0 .

4. APPLICATION TO FACTORIALITY OF UPPER CLUSTER ALGEBRAS

In this section we give several equivalent characterizations for the factoriality of upper cluster algebras. As an application, we show that full rank, primitive upper cluster algebras are factorial.

4.1. Characterizations for the factoriality of upper cluster algebras. We first give an observation which shows why the valuation pairing can be used to study the factoriality of upper cluster algebras.

Proposition 4.1 (Observation). *Let \mathcal{U} be a geometric upper cluster algebra and $A_{k;t}$ a cluster variable of \mathcal{U} . Then the following two statements are equivalent.*

- (i) $A_{k;t}$ is prime in \mathcal{U} .
- (ii) For any non-zero elements M and L in \mathcal{U} , we have the following equality:

$$(4.1) \quad (A_{k;t} || M \cdot L)_v = (A_{k;t} || M)_v + (A_{k;t} || L)_v.$$

Proof. (i) \Rightarrow (ii): This is clear from the definition of the valuation pairing.

(ii) \Rightarrow (i): For any $0 \neq M, L \in \mathcal{U}$ with $A_{k;t}|M \cdot L$ in \mathcal{U} , we claim that either $A_{k;t}|M$ in \mathcal{U} or $A_{k;t}|L$ in \mathcal{U} . Otherwise, we have that both M and L are not divisible by $A_{k;t}$ in \mathcal{U} . Then we have

$$(A_{k;t}||M)_v = 0 = (A_{k;t}||L)_v.$$

Now by our assumption, we have

$$(A_{k;t}||M \cdot L)_v = (A_{k;t}||M)_v + (A_{k;t}||L)_v = 0.$$

This contradicts $A_{k;t}|M \cdot L$ in \mathcal{U} . So either $A_{k;t}|M$ or $A_{k;t}|L$ in \mathcal{U} .

By Theorem 2.35 (ii), we know that $A_{k;t}$ is not invertible in \mathcal{U} . Hence, $A_{k;t}$ is prime in \mathcal{U} . \square

It is natural to ask under which conditions the equality (4.1) always holds. Thanks to Reduction Lemma 3.6, we can answer this question for *full rank upper cluster algebras*.

Proposition 4.2. *Let \mathcal{U} be a full rank upper cluster algebra and t a seed of \mathcal{U} . Put $t_k = \mu_k(t)$ and let $P_{k;t}$ be the k -th exchange binomial of t . Then the following statements are equivalent.*

- (i) $(A_{k;t}||M \cdot L)_v = (A_{k;t}||M)_v + (A_{k;t}||L)_v$ holds for any non-zero elements M and L in \mathcal{U} .
- (ii) $A_{k;t}$ is prime in $\mathcal{L}(t_k) = \mathbb{K}[Z_1^{\pm 1}, \dots, Z_m^{\pm 1}][A_{1;t_k}^{\pm 1}, \dots, A_{n;t_k}^{\pm 1}]$.
- (iii) $P_{k;t}$ is prime in $\mathcal{L}(t_k)$.
- (iv) $P_{k;t}$ is prime in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}]$, which is actually equal to $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_k}, \dots, \hat{A}_{k;t_k}, \dots, A_{n;t_k}]$.
- (v) $P_{k;t}$ is irreducible in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}]$.

Proof. (i) \Rightarrow (ii): By our assumption in Subsection 1.4, we know that $A_{k;t}$ is not invertible in $\mathcal{L}(t_k)$. Now assume by contradiction that $A_{k;t}$ is not prime in $\mathcal{L}(t_k)$. Then there exists $M', L' \in \mathcal{L}(t_k)$ such that both M' and L' are not divisible by $A_{k;t}$ in $\mathcal{L}(t_k)$, but the product $M' \cdot L'$ is divisible by $A_{k;t}$ in $\mathcal{L}(t_k)$. Since M' and L' are in

$$\mathcal{L}(t_k) = \mathbb{K}[Z_1^{\pm 1}, \dots, Z_m^{\pm 1}][A_{1;t_k}^{\pm 1}, \dots, A_{n;t_k}^{\pm 1}],$$

there exists a cluster monomial $A_{t_k}^{\mathbf{v}}$ in t_k such that $M := A_{t_k}^{\mathbf{v}} M'$ and $L := A_{t_k}^{\mathbf{v}} L'$ belong to \mathcal{U} .

Because $A_{t_k}^{\mathbf{v}}$ is invertible in $\mathcal{L}(t_k)$, we know that both M and L are not divisible by $A_{k;t}$ in $\mathcal{L}(t_k)$, but the product $M \cdot L$ is divisible by $A_{k;t}$ in $\mathcal{L}(t_k)$. By the Reduction Lemma 3.6 (ii), we have

$$(A_{k;t}||M)_v = 0 = (A_{k;t}||L)_v \text{ but } (A_{k;t}||M \cdot L)_v > 0.$$

This contradicts that $(A_{k;t}||M \cdot L)_v = (A_{k;t}||M)_v + (A_{k;t}||L)_v$ holds for any non-zero elements M and L in \mathcal{U} .

(ii) \Rightarrow (i): This follows from the Reduction Lemma 3.6 (ii).

(ii) \Leftrightarrow (iii): Applying the exchange relation, we have $A_{k;t} A_{k;t_k} = P_{k;t}$. Since $A_{k;t_k}$ is invertible in $\mathcal{L}(t_k)$, we know that $A_{k;t}$ is prime in $\mathcal{L}(t_k)$ if and only if $P_{k;t}$ is prime in $\mathcal{L}(t_k)$.

(iii) \Leftrightarrow (iv): Notice that $\mathcal{L}(t_k)$ is a localization of $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_k}, \dots, A_{n;t_k}]$ at $Z_1, \dots, Z_m, A_{1;t_k}, \dots, A_{n;t_k}$. Since $P_{k;t} = P_{k;t_k}$ is a binomial in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_k}, \dots, A_{n;t_k}]$$

and it is not divisible by any Z_i and $A_{j;t_k}$, we know that $P_{k;t}$ is prime in $\mathcal{L}(t_k)$ if and only if it is prime in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_k}, \dots, A_{n;t_k}]$. Because $P_{k;t}$ is actually a binomial in variables from

$$\{Z_1, \dots, Z_m, A_{1;t_k}, \dots, \hat{A}_{k;t_k}, \dots, A_{n;t_k}\} = \{Z_1, \dots, Z_m, A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}\},$$

we know that $P_{k;t}$ is prime in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, A_{n;t}]$ if and only if it is prime in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}].$$

Hence, $P_{k;t}$ is prime in $\mathcal{L}(t_k)$ if and only if it is prime in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}].$$

(iv) \iff (v): This follows from the fact that the polynomial ring

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}]$$

is factorial. □

Lemma 4.3. [GELS19, Corollary 1.23] *Let R be a domain and $S \subseteq R \setminus \{0\}$ a multiplicative set generated by prime elements. Then R is factorial if and only if the localization $S^{-1}R$ is.*

Theorem 4.4. *Let \mathcal{U} be a full rank upper cluster algebra with initial seed t_0 . Then the following statements are equivalent.*

- (i) \mathcal{U} is factorial.
- (ii) Any initial cluster variable $A_{k;t_0}$ is prime in \mathcal{U} .
- (iii) Any initial cluster variable $A_{k;t_0}$ satisfies

$$(A_{k;t_0} \parallel M \cdot L)_v = (A_{k;t_0} \parallel M)_v + (A_{k;t_0} \parallel L)_v,$$

for any non-zero elements M and L in \mathcal{U} .

- (iv) Any exchange binomial $P_{k;t_0}$ of t_0 is irreducible in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, \hat{A}_{k;t_0}, \dots, A_{n;t_0}].$$

Proof. (i) \implies (ii): By Theorem 2.35 (ii), we know that any cluster variable is irreducible in \mathcal{U} . Because \mathcal{U} is factorial, we get that any irreducible element is prime in \mathcal{U} . In particular, any initial cluster variable $A_{k;t_0}$ is prime in \mathcal{U} .

(ii) \implies (i): Let S be the multiplicative set generated by initial cluster variables. By the Laurent phenomenon, we have

$$S^{-1}\mathcal{U} = \mathcal{L}(t_0) = \mathbb{Z}[Z_1^{\pm 1}, \dots, Z_n^{\pm 1}][A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}],$$

which is factorial. Since the initial cluster variables are prime in \mathcal{U} and by Lemma 4.3, we get that \mathcal{U} is factorial.

The equivalence of (ii), (iii) and (iv) follows from Proposition 4.2. □

Remark 4.5. Garcia Elsener *et al.* in [GELS19, Theorem 5.1] prove the equivalence of (i) and (iv) in Theorem 4.4 for geometric cluster algebra \mathcal{A} (possibly not full rank) with an *acyclic initial seed*. Note that in acyclic case, $\mathcal{A} = \mathcal{U}$, by Proposition 2.19. Our method here is very different with that in [GELS19].

4.2. Full rank, primitive upper cluster algebras are factorial. In this subsection, we show that full rank and primitive upper cluster algebras are factorial.

Recall that a vector $(b_1, \dots, b_{n+m})^\top \in \mathbb{Z}^{n+m}$ is primitive if the greatest common divisor $d \in \mathbb{N}$ of b_1, \dots, b_{n+m} is 1. An extended exchange matrix \tilde{B}_t is *primitive* if each column vector of \tilde{B}_t is primitive.

Definition 4.6 (Primitive upper cluster algebra). Let \mathcal{U} be a geometric upper cluster algebra with initial seed t_0 . We say that \mathcal{U} is a *primitive upper cluster algebra*, if the initial extended exchange matrix \tilde{B}_{t_0} is primitive.

Proposition 4.7. *Let \mathcal{U} be a primitive upper cluster algebra. Then for any seed t , the following statements hold.*

- (i) [LP16, Lemma 4.3] \tilde{B}_t is primitive.
- (ii) [LP16, Lemma 4.1] The k -th exchange binomial of t is irreducible in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t}, \dots, \hat{A}_{k;t}, \dots, A_{n;t}],$$

where $k = 1, \dots, n$.

Theorem 4.8. *Let \mathcal{U} be a full rank, primitive upper cluster algebra with initial seed t_0 , and let*

$$M = \frac{P_M(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{d_1} \cdots A_{n;t_0}^{d_n}}$$

be a non-initial cluster variable of \mathcal{U} , where $\mathbf{d} := (d_1, \dots, d_n)^\top$ is the d -vector of M with respect to t_0 . Then

- (i) \mathcal{U} is factorial.
- (ii) the numerator polynomial P_M of M is irreducible in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}].$$

Proof. (i) By Proposition 4.7 (ii), we know that every exchange binomial of t_0 is irreducible in the corresponding polynomial ring. Then the result follows from Theorem 4.4 (i)(iv).

(ii) We first show that P_M is not invertible in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}]$. Since M is a non-initial cluster variable and by Proposition 2.30 (4), we have $\mathbf{d} \in \mathbb{N}^n$ and thus $A_{t_0}^{\mathbf{d}} \in \mathcal{U}$. By $M = P_M/A_{t_0}^{\mathbf{d}}$, we get $MA_{t_0}^{\mathbf{d}} = P_M$. By Theorem 2.35 (2), we know that both M and $A_{t_0}^{\mathbf{d}}$ are not invertible in \mathcal{U} . So $P_M = MA_{t_0}^{\mathbf{d}}$ is not invertible in \mathcal{U} . Namely, we have

$$P_M \notin \mathcal{U}^\times = \{\lambda Z_1^{c_1} \cdots Z_m^{c_m} \mid \lambda \in \mathbb{K}^\times, c_1, \dots, c_m \in \mathbb{Z}\}.$$

In particular, $P_M \notin \mathbb{K}^\times$. So P_M is not invertible in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}]$.

We claim that P_M is not divisible by any Z_i and $A_{j;t_0}$ in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}].$$

Since P_M is a numerator polynomial, it is not divisible by any $A_{j;t_0}$. There are two approaches to prove $Z_i \nmid P_M$. The first one is that we prove the result by induction on the minimal length of a sequence of mutations from the initial seed to the final seed. The second one is that we just view Z_i as an exchangeable cluster variable. Then by the fact that M and Z_i are in the same cluster and are not equal, we get $(Z_i \parallel M)_d = 0$, by Proposition 2.30 (5). This implies that P_M is not divisible by Z_i .

We claim that P_M is not invertible in

$$\mathcal{L}(t_0) = \mathbb{K}[Z_1^{\pm 1}, \dots, Z_m^{\pm 1}][A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}].$$

This follows from the two facts that P_M is not invertible in

$$\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}]$$

and P_M is not divisible by any Z_i and $A_{j;t_0}$ in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}]$.

Now we show that P_M is irreducible in $\mathcal{L}(t_0)$. Assume that $P_M = L_1 L_2$ for some $L_1, L_2 \in \mathcal{L}(t_0)$. We know that there exist vectors $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{N}^n$ such that

$$U_1 := A_{t_0}^{\mathbf{v}_1} L_1 \in \mathcal{U} \quad \text{and} \quad U_2 := A_{t_0}^{\mathbf{v}_2} L_2 \in \mathcal{U}.$$

By $M = \frac{P_M}{A_{t_0}^{\mathbf{d}}} = \frac{L_1 L_2}{A_{t_0}^{\mathbf{d}}} = \frac{U_1 U_2}{A_{t_0}^{\mathbf{d} + \mathbf{v}_1 + \mathbf{v}_2}}$, we get

$$(4.2) \quad M A_{t_0}^{\mathbf{d} + \mathbf{v}_1 + \mathbf{v}_2} = U_1 U_2,$$

which is an equality in \mathcal{U} , since $\mathbf{d}, \mathbf{v}_1, \mathbf{v}_2 \in \mathbb{N}^n$. Because \mathcal{U} is factorial and M is irreducible in \mathcal{U} , we get that either U_1 or U_2 is divisible by M in \mathcal{U} . Without loss of generality, we assume that $U_1' := U_1/M \in \mathcal{U}$. Then by the equality (4.2), we get $U_1' U_2 = A_{t_0}^{\mathbf{d} + \mathbf{v}_1 + \mathbf{v}_2}$, which implies that both U_1' and U_2 are invertible in $\mathcal{L}(t_0)$. So $L_2 = U_2/A_{t_0}^{\mathbf{v}_2}$ is invertible in $\mathcal{L}(t_0)$. Hence, P_M is irreducible in $\mathcal{L}(t_0)$.

Since P_M is irreducible in $\mathcal{L}(t_0)$ and by the fact that P_M is not divisible by any Z_i and $A_{j;t_0}$ in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}]$, we get that P_M is irreducible in $\mathbb{K}[Z_1, \dots, Z_m][A_{1;t_0}, \dots, A_{n;t_0}]$. This completes the proof. \square

Remark 4.9. (i) Note that any principal coefficient upper cluster algebra is primitive and has full rank. So principal coefficient upper cluster algebras are always factorial.

(ii) A different proof of Theorem 4.8 (i) via the factoriality of Cox rings can be found in [GHK15, Corollary 4.7].

(iii) Garcia Elsener *et al.* in [GELS19] prove that any principal coefficient cluster algebra \mathcal{A} with an *acyclic initial exchange matrix* is factorial. Note that in the acyclic case, we have $\mathcal{A} = \mathcal{U}$, by Proposition 2.19.

Corollary 4.10. Let \mathcal{U} be a full rank, primitive upper cluster algebra and $A_{k;t_0}$ a cluster variable of \mathcal{U} . Then for any non-zero elements M and L in \mathcal{U} , we have

$$(A_{k;t_0} \parallel M \cdot L)_v = (A_{k;t_0} \parallel M)_v + (A_{k;t_0} \parallel L)_v.$$

Proof. This follows from Theorem 4.4 (i)(iii) and Theorem 4.8 (i). \square

4.3. Examples of non-factorial upper cluster algebras.

Proposition 4.11. [GLS13, Corollary 6.5] *Let \mathcal{A} be a geometric cluster algebra with initial extended exchange matrix*

$$\tilde{B}_{t_0} = B_{t_0} = \begin{pmatrix} 0 & -c \\ d & 0 \end{pmatrix},$$

where $c \geq 1$ and $d \geq 3$ an odd number. Then \mathcal{A} is not factorial.

Remark 4.12. Note that in Proposition 4.11, \mathcal{A} is an acyclic cluster algebra. In this case, \mathcal{A} coincides with its upper cluster algebra \mathcal{U} , by Proposition 2.19. So Proposition 4.11 provides many upper cluster algebras which have full rank but are not factorial.

Let \mathcal{U} be the geometric upper cluster algebra with initial extended exchange matrix

$$\tilde{B}_{t_0} = B_{t_0} = \begin{pmatrix} 0 & -1 \\ 3 & 0 \end{pmatrix}.$$

We know that \mathcal{U} is a full rank cluster algebra. It is non-factorial, by Remark 4.12. In the view point of Theorem 4.4, the non-factoriality of \mathcal{U} is due to the fact that the first exchange binomial $A_{2;t_0}^3 + 1 = (A_{2;t_0} + 1)(A_{2;t_0}^2 - A_{2;t_0} + 1)$ of t_0 is not irreducible.

The following is a concrete example where \mathcal{U} is primitive but not factorial.

Example 4.13. Let \mathcal{A} be a geometric cluster algebra with the following initial extended exchange matrix

$$\tilde{B}_{t_0} = B_{t_0} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

It is easy to see that \tilde{B}_{t_0} is primitive and acyclic but not full rank. It is known from [GLS13, Proposition 6.1] that \mathcal{A} is not factorial. Since \mathcal{A} is acyclic, \mathcal{A} coincides with its upper cluster algebra \mathcal{U} , by Proposition 2.19. So $\mathcal{U} = \mathcal{A}$ is not factorial.

4.4. Ray fish theorem. Recall that the starfish theorem states that any full rank upper cluster algebra \mathcal{U} can be written as the intersection of $n + 1$ Laurent polynomial rings. In this subsection, we show that any full rank, primitive upper cluster algebra can be written as the intersection of two Laurent polynomial rings. We call this the ray fish theorem.

Proposition 4.14. [GLS13, Corollary 4.2] *Let \mathcal{A} be a geometric cluster algebra with coefficient semifield $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$ and \mathcal{U} its upper cluster algebra. Let t, t_0 be two seeds of \mathcal{A} with no common cluster variables. If \mathcal{A} is factorial, then*

$$\mathcal{A} = \mathcal{U} = \mathcal{L}(t) \cap \mathcal{L}(t_0),$$

where $\mathcal{L}(t) = \mathbb{K}\mathbb{P}[A_{1;t}^{\pm 1}, \dots, A_{n;t}^{\pm 1}]$ and $\mathcal{L}(t_0) = \mathbb{K}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$.

Remark 4.15. By Proposition 4.14, a factorial geometric cluster algebra must coincide with its upper cluster algebra. This provides a reason why we focus on the factoriality of upper cluster algebras rather than cluster algebras.

Corollary 4.16. Let \mathcal{U} be a full rank, primitive upper cluster algebra and \mathcal{A} the corresponding cluster algebra. Then $\mathcal{A} = \mathcal{U}$ if and only if \mathcal{A} is factorial.

Proof. “ \implies ”: Since \mathcal{U} is full rank and primitive and by Theorem 4.8, we know that \mathcal{U} is factorial. Then the factoriality of \mathcal{A} follows from $\mathcal{A} = \mathcal{U}$.

“ \impliedby ”: This follows from Proposition 4.14. \square

Lemma 4.17. *Let \mathcal{U} be a geometric upper cluster algebra with coefficient semifield $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$ and $A_{i;t_0}, A_{j;t}$ two cluster variables of \mathcal{U} . Then $A_{i;t_0}$ and $A_{j;t}$ are associate in \mathcal{U} if and only if $A_{i;t_0} = A_{j;t}$.*

Proof. Assume that $A_{i;t_0}$ and $A_{j;t}$ are associate in \mathcal{U} . Then

$$(4.3) \quad A_{j;t} = \lambda Z_1^{c_1} \cdots Z_m^{c_m} A_{i;t_0}$$

for some $\lambda \in \mathbb{K}^\times$ and $c_1, \dots, c_m \in \mathbb{Z}$, by Theorem 2.35 (2). Notice that the equality (4.3) can be viewed as the Laurent expansion of $A_{j;t}$ with respect to t_0 . It is easy to see that $(A_{i;t_0} \parallel A_{j;t})_d = -1$. Then by Proposition 2.30 (4), we get $A_{j;t} = A_{i;t_0}$.

Conversely, assume $A_{i;t_0} = A_{j;t}$. Clearly, we have that $A_{i;t_0}$ and $A_{j;t}$ are associate in \mathcal{U} . This completes the proof. \square

Inspired by Proposition 4.14, we give the following result.

Theorem 4.18 (Ray fish theorem). *Let \mathcal{U} be a full rank, primitive upper cluster algebra with coefficient semifield $\mathbb{P} = \text{Trop}(Z_1, \dots, Z_m)$ and t, t_0 two seeds of \mathcal{U} with no common cluster variables. Then $\mathcal{U} = \mathcal{L}(t) \cap \mathcal{L}(t_0)$, where $\mathcal{L}(t) = \mathbb{K}\mathbb{P}[A_{1;t}^{\pm 1}, \dots, A_{n;t}^{\pm 1}]$ and $\mathcal{L}(t_0) = \mathbb{K}\mathbb{P}[A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$.*

Proof. The proof is the same as that of [GLS13, Corollary 4.2]. For convenience of the reader, we repeat it here.

Notice that the inclusion $\mathcal{U} \subseteq \mathcal{L}(t) \cap \mathcal{L}(t_0)$ is clear. Now we show the converse inclusion. We know that any $M \in \mathcal{L}(t) \cap \mathcal{L}(t_0)$ has the following form:

$$M = \frac{P(A_{1;t}, \dots, A_{n;t})}{A_{1;t}^{v_1} \cdots A_{n;t}^{v_n}} = \frac{Q(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{s_1} \cdots A_{n;t_0}^{s_n}},$$

where $P \in \mathbb{K}\mathbb{P}[A_{1;t}, \dots, A_{n;t}]$, $Q \in \mathbb{K}\mathbb{P}[A_{1;t_0}, \dots, A_{n;t_0}] \subseteq \mathcal{U}$ and $v_i, s_i \geq 0$ for any $i = 1, \dots, n$. Thus we have $A_{1;t_0}^{s_1} \cdots A_{n;t_0}^{s_n} P = Q A_{1;t}^{v_1} \cdots A_{n;t}^{v_n}$.

By Theorem 2.35 (2), we know that $A_{i;t_0}$ and $A_{j;t}$ are irreducible in \mathcal{U} for any i and j . By Lemma 4.17 and the fact that t and t_0 have no common cluster variables, we know that $A_{i;t_0}$ and $A_{j;t}$ are non-associate for all $1 \leq i, j \leq n$. Since \mathcal{U} is full rank and primitive and by Theorem 4.8, we know that \mathcal{U} is factorial. So the equality $A_{1;t_0}^{s_1} \cdots A_{n;t_0}^{s_n} P = Q A_{1;t}^{v_1} \cdots A_{n;t}^{v_n}$ implies that P is divisible by $A_{1;t}^{v_1} \cdots A_{n;t}^{v_n}$ in \mathcal{U} . Namely, there exists $P' \in \mathcal{U}$ such that $P = A_{1;t}^{v_1} \cdots A_{n;t}^{v_n} P'$. Thus

$$M = \frac{P}{A_{1;t}^{v_1} \cdots A_{n;t}^{v_n}} = P' \in \mathcal{U}.$$

So we have $\mathcal{L}(t) \cap \mathcal{L}(t_0) \subseteq \mathcal{U}$ and thus $\mathcal{U} = \mathcal{L}(t) \cap \mathcal{L}(t_0)$. \square

5. APPLICATIONS TO d -VECTORS

5.1. d -compatibility degree and d -vectors via the valuation pairing. In this subsection, we show how to express d -compatibility and d -vectors using the valuation pairing for full rank upper cluster algebras.

Theorem 5.1. *Let \mathcal{U} be a full rank upper cluster algebra with initial seed t_0 . Let*

$$M = \frac{P_M(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{d_1} \cdots A_{n;t_0}^{d_n}}$$

be a non-zero element in \mathcal{U} , where $\mathbf{d}_M := (d_1, \dots, d_n)^\top$ is the d -vector of M with respect to A_{t_0} . Then the following statements hold.

(i) *For any $k = 1, \dots, n$, we have*

$$d_k = (A_{k;t_0} \| M)_d = (A_{k;t_0} \| P_M)_v - (A_{k;t_0} \| M)_v.$$

(ii) *If $(A_{k;t_0} \| M)_v = 0$ for some k , that is, $M/A_{k;t_0} \notin \mathcal{U}$, then*

$$d_k = (A_{k;t_0} \| M)_d = (A_{k;t_0} \| P_M)_v \geq 0.$$

(iii) *If $(A_{k;t_0} \| M)_v = 0$ for $k = 1, \dots, n$, then \mathbf{d}_M is uniquely determined by P_M .*

Proof. (i) We know that there exist two vectors

$$\mathbf{d}^+ = (d_1^+, \dots, d_n^+)^\top, \quad \mathbf{d}^- = (d_1^-, \dots, d_n^-)^\top \in \mathbb{N}^n$$

such that

$$\mathbf{d}_M = \mathbf{d}^+ - \mathbf{d}^-.$$

By $M = P_M/A_{t_0}^{\mathbf{d}^M} = P_M/A_{t_0}^{\mathbf{d}^+ - \mathbf{d}^-}$, we have $A_{t_0}^{\mathbf{d}^+} M = A_{t_0}^{\mathbf{d}^-} P_M$. By the Reduction Lemma 3.6 (iii), we get

$$(A_{k;t_0} \| M)_v + d_k^+ = (A_{k;t_0} \| A_{t_0}^{\mathbf{d}^+} M)_v = (A_{k;t_0} \| A_{t_0}^{\mathbf{d}^-} P_M)_v = (A_{k;t_0} \| P_M)_v + d_k^-.$$

So we have

$$d_k = d_k^+ - d_k^- = (A_{k;t_0} \| P_M)_v - (A_{k;t_0} \| M)_v.$$

(ii) and (iii) follow from (i). \square

5.2. Local factorizations of cluster monomials. Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and initial seed t_0 . In this subsection, we prove that if M is a cluster monomial in non-initial cluster variables, then $M/A_{k;t_0} \notin \mathcal{U}$, that is, $(A_{k;t_0} \| M)_v = 0$ for $k = 1, \dots, n$. In particular, we can give a local factorization for any cluster monomial.

Lemma 5.2. *Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and t a seed of \mathcal{U} . If $N = A_{1;t}^{b_1} A_{2;t}^{b_2} \cdots A_{n;t}^{b_n} \in \mathcal{U}$ for some $b_1, b_2, \dots, b_n \in \mathbb{Z}$, then $b_1, b_2, \dots, b_n \geq 0$.*

Proof. Assume by contradiction that there exists some $b_{i_0} < 0$. Without loss of generality, we just assume $b_1 < 0$. Let $t' = \mu_1(t)$ and thus $t = \mu_1(t')$. Applying the exchange relation, we know that

$$A_{1;t} A_{1;t'} = P \in \mathbb{K}\mathbb{P}[A_{2;t'}, \dots, A_{n;t'}],$$

where P is the first exchange binomial of t' . Then we know that

$$N = A_{1;t}^{b_1} A_{2;t}^{b_2} \cdots A_{n;t}^{b_n} = \frac{A_{1;t'}^{-b_1} A_{2;t'}^{b_2} \cdots A_{n;t'}^{b_n}}{P^{-b_1}},$$

which is the expansion of N with respect to $A_{t'}$. By $N \in \mathcal{U}$, we get that

$$N = \frac{A_{1;t'}^{-b_1} A_{2;t'}^{b_2} \cdots A_{n;t'}^{b_n}}{P^{-b_1}} \in \mathcal{L}(t') = \mathbb{K}\mathbb{P}[A_{1;t'}^{\pm 1}, \dots, A_{n;t'}^{\pm 1}].$$

Thus we have $1/P^{-b_1} \in \mathcal{L}(t')$. Because both P^{-b_1} and $1/P^{-b_1}$ are in $\mathcal{L}(t')$, we must have $P^{-b_1} = \lambda A_{1;t'}^{c_1} \cdots A_{n;t'}^{c_n}$ for some invertible element λ in $\mathbb{K}\mathbb{P}$ and $c_1, \dots, c_n \in \mathbb{Z}$. Since P is a polynomial in $\mathbb{K}\mathbb{P}[A_{2;t'}, \dots, A_{n;t'}]$ with $A_{i;t'} \nmid P$ for any i , we get that $c_1 = \dots = c_n = 0$ and $P^{-b_1} = \lambda$ is invertible in $\mathbb{K}\mathbb{P}$. By our assumption in Subsection 1.4, we know that $P^{-b_1} = \lambda$ is not invertible in $\mathbb{K}\mathbb{P}$. This concludes a contradiction. So we must have $b_1, \dots, b_n \geq 0$. \square

Proposition 5.3. *Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and t_0, t two seeds of \mathcal{U} . Let $M = A_{1;t_0}^{c_1} \cdots A_{n;t_0}^{c_n}$ be a cluster monomial in A_{t_0} and $N = A_{1;t}^{b_1} \cdots A_{n;t}^{b_n}$ a Laurent monomial in A_t . Let*

$$I = \{i | c_i > 0\} \quad \text{and} \quad J = \{j | b_j > 0\}.$$

If there exists an invertible element λ in $\mathbb{K}\mathbb{P}$ such that $M = \lambda N$, then

- (i) *N is a cluster monomial in A_t , i.e., we have $b_1, \dots, b_n \geq 0$;*
- (ii) *there exists a bijection $\sigma : I \rightarrow J$ such that $A_{i;t_0} = A_{\sigma(i);t}$ and $c_i = b_{\sigma(i)}$. In particular, we have $M = N$.*

Proof. (i) By $M = \lambda N$, we know that $N = \lambda^{-1}M \in \mathcal{U}$. Then by Lemma 5.2, we know that N is a cluster monomial in A_t .

(ii) Let $A_{j;t}$ be a cluster variable appearing in N . Then by (i), we have $b_j > 0$ and $j \in J$. Because we have

$$\sum_{i \in I}^n c_i (A_{j;t} \| A_{i;t_0})_d = (A_{j;t} \| M)_d = (A_{j;t} \| \lambda N)_d = (A_{j;t} \| N)_d = -b_j < 0,$$

there must exist some $j' \in I$ such that

$$(A_{j;t} \| A_{j';t_0})_d < 0.$$

Then by Proposition 2.30 (4), we have $A_{j;t} = A_{j';t_0}$. So we have

$$c_{j'} = -(A_{j';t_0} \| M)_d = -(A_{j;t} \| M)_d = -(A_{j;t} \| N)_d = b_j.$$

Similarly, we can show that any cluster variable $A_{i;t_0}$ appearing in M also appears in N , and the multiplicity of $A_{i;t_0}$ in N is equal to the multiplicity of $A_{i;t_0}$ in M . Then the result follows. \square

Theorem 5.4 (Local factorization of cluster monomials). *Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and initial seed t_0 . Then the following statements hold.*

- (i) *If L is a cluster monomial of \mathcal{U} which does not contain the initial cluster variable $A_{k;t_0}$ for some k , then $L/A_{k;t_0} \notin \mathcal{U}$, that is, $(A_{k;t_0} \| L)_v = 0$.*
- (ii) *If L is a cluster monomial in non-initial cluster variables, then $L/A_{k;t_0} \notin \mathcal{U}$, that is, $(A_{k;t_0} \| L)_v = 0$ for $k = 1, \dots, n$.*
- (iii) *Let $M = \prod_{i=1}^n A_{i;t}^{b_i}$ be a cluster monomial in a seed t of \mathcal{U} and $I = \{i | A_{i;t} \in A_{t_0}\}$, then $M = \prod_{i \in I} A_{i;t}^{b_i} \cdot \prod_{j \notin I} A_{j;t}^{b_j}$ is a local factorization of M with respect to t_0 .*

Proof. (i) Assume by contradiction that $L/A_{k;t_0} \in \mathcal{U}$. In the following proof, we will deduce a contradiction.

Because the initial cluster variable $A_{k;t_0}$ does not appear in L , there exists a seed t such that L is a cluster monomial in A_t and $A_{k;t_0}$ does not belong to A_t . Consider the Laurent expansion of $A_{k;t_0}$ with respect to A_t , say

$$A_{k;t_0} = \frac{P(A_{1;t}, \dots, A_{n;t})}{A_{1;t}^{d'_1} \cdots A_{n;t}^{d'_n}},$$

where $P \in \mathbb{K}\mathbb{P}[A_{1;t}, \dots, A_{n;t}]$ with $A_{i;t} \nmid P$ for $i = 1, \dots, n$.

Now we have that

$$L/A_{k;t_0} = \frac{L \cdot (A_{1;t}^{d'_1} \cdots A_{n;t}^{d'_n})}{P},$$

which is the expansion of $L/A_{k;t_0}$ with respect to A_t . By $L/A_{k;t_0} \in \mathcal{U}$, we have

$$\frac{L \cdot (A_{1;t}^{d'_1} \cdots A_{n;t}^{d'_n})}{P} \in \mathcal{L}(t) := \mathbb{K}\mathbb{P}[A_{1;t}^{\pm 1}, \dots, A_{n;t}^{\pm 1}].$$

Because $L \cdot (A_{1;t}^{d'_1} \cdots A_{n;t}^{d'_n})$ is a monic Laurent monomial in $\mathcal{L}(t)$, we get $1/P \in \mathcal{L}(t)$. Since both P and $1/P$ are in $\mathcal{L}(t)$, we must have $P = \lambda A_{1;t}^{c_1} \cdots A_{n;t}^{c_n}$ for some invertible element λ in $\mathbb{K}\mathbb{P}$ and $c_1, \dots, c_n \in \mathbb{Z}$. Because P is a polynomial in

$\mathbb{K}\mathbb{P}[A_{1;t}, \dots, A_{n;t}]$ with $A_{i;t} \nmid P$ for any i , we get $c_1 = \dots = c_n = 0$ and thus $P = \lambda$ is invertible in $\mathbb{K}\mathbb{P}$. Applying Proposition 5.3 to

$$A_{k;t_0} = \frac{P}{A_{1;t}^{d'_1} \cdots A_{n;t}^{d'_n}} = \lambda \cdot A_{1;t}^{-d'_1} \cdots A_{n;t}^{-d'_n},$$

we get $A_{k;t_0} \in A_t$, which contradicts $A_{k;t_0} \notin A_t$. So $L/A_{k;t_0} \notin \mathcal{U}$.

(ii) follows from (i).

(iii) follows from (ii) and the definition of local factorization. \square

Corollary 5.5. Let \mathcal{U} be a full rank upper cluster algebra with initial seed t_0 . Let

$$M = \frac{P_M(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{d_1} \cdots A_{n;t_0}^{d_n}}$$

be a cluster monomial of \mathcal{U} , where $\mathbf{d}_M := (d_1, \dots, d_n)^T$ is the d -vector of M with respect to A_{t_0} . Then the following statements hold.

(i) If M does not contain the initial cluster variable $A_{k;t_0}$ for some k , then we have

$$d_k = (A_{k;t_0} \| M)_d = (A_{k;t_0} \| P_M)_v \geq 0.$$

(ii) If M is a cluster monomial in non-initial cluster variables, then \mathbf{d}_M and M are uniquely determined by the numerator polynomial P_M .

Proof. (i) By Theorem 5.4 (i), we know that $(A_{k;t_0} \| M)_v = 0$. Then by Theorem 5.1 (ii), we get $d_k = (A_{k;t_0} \| M)_d = (A_{k;t_0} \| P_M)_v \geq 0$.

(ii) This follows from (i). \square

Actually, the above result can be extended from cluster monomials to monomials in cluster variables if we further assume that \mathcal{U} is primitive or factorial.

Proposition 5.6. Let \mathcal{U} be a full rank, primitive upper cluster algebra with initial seed t_0 . Let

$$M = \frac{P_M(A_{1;t_0}, \dots, A_{n;t_0})}{A_{1;t_0}^{d_1} \cdots A_{n;t_0}^{d_n}}$$

be a monomial in cluster variables of \mathcal{U} (not necessarily a cluster monomial), where $\mathbf{d}_M := (d_1, \dots, d_n)^T$ is the d -vector of M with respect to A_{t_0} . Then the following statements hold.

(i) If M does not contain the initial cluster variable $A_{k;t_0}$ for some k , then we have $(A_{k;t_0} \| M)_v = 0$ and $d_k = (A_{k;t_0} \| M)_d = (A_{k;t_0} \| P_M)_v \geq 0$.

(ii) If M is a monomial in non-initial cluster variables, then \mathbf{d}_M and M are uniquely determined by the numerator polynomial P_M .

Proof. (i) Since M is a monomial in cluster variables and it does not contain $A_{k;t_0}$, we can assume that $M = \prod_{i=1}^s M_i$, where each M_i is a cluster variable of \mathcal{U} different from $A_{k;t_0}$. By Theorem 5.4 (i), we have $(A_{k;t_0} \| M_i)_v = 0$, for $i = 1, \dots, s$.

Since \mathcal{U} is a full rank, primitive upper cluster algebra, we know that \mathcal{U} is factorial, by Theorem 4.8. Then by Theorem 4.4 (i)(iii), we know that

$$(A_{k;t_0} \| M)_v = \sum_{i=1}^s (A_{k;t_0} \| M_i)_v = 0.$$

Then by Theorem 5.1 (ii), we get $d_k = (A_{k;t_0} \| M)_d = (A_{k;t_0} \| P_M)_v \geq 0$.

(ii) This follows from (i). \square

6. APPLICATION TO F -POLYNOMIALS

In this section, we always assume that \mathcal{U} is an upper cluster algebra with principal coefficients at t_0 .

6.1. From F -polynomials to monomials in non-initial cluster variables. In this subsection, we prove that if M is a monomial in non-initial cluster variables, then M is uniquely determined by its F -polynomial.

Let M be a monomial in cluster variables of \mathcal{U} , say $M = \prod_{i=1}^s M_i$, where each M_i is a cluster variable of \mathcal{U} . The F -polynomial F_M of M is defined to be the polynomial $F_M := \prod_{i=1}^s F_{M_i}$, where F_{M_i} is the F -polynomial of the cluster variable M_i . The g -vector of M is defined to be the vector $\mathbf{g}_M := \sum_{i=1}^s \mathbf{g}_{M_i}$, where \mathbf{g}_{M_i} is the g -vector of M_i .

Theorem 6.1. *Let \mathcal{U} be an upper cluster algebra with principal coefficients at t_0 . Let M, N be two monomials in non-initial cluster variables of \mathcal{U} and F_M, F_N their F -polynomials. If $F_M = F_N$, then $M = N$. In particular, $\mathbf{g}_M = \mathbf{g}_N$, where \mathbf{g}_M and \mathbf{g}_N are g -vectors of M and N .*

Proof. Let \mathbf{d}_M and \mathbf{d}_N be the d -vectors of M and N with respect to A_{t_0} . We know that M and N have the form

$$(6.1) \quad M = P_M/A_{t_0}^{\mathbf{d}_M} \quad \text{and} \quad N = P_N/A_{t_0}^{\mathbf{d}_N},$$

where $P_M, P_N \in \mathbb{Z}[Z_1, \dots, Z_n; A_{1;t_0}, \dots, A_{n;t_0}]$ with $A_{i;t_0} \nmid P_M$ and $A_{i;t_0} \nmid P_N$ for any $i = 1, \dots, n$.

By Theorem 2.26 (i), we know that

$$\begin{aligned} M &= A_{t_0}^{\mathbf{g}_M} F_M(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}), \\ N &= A_{t_0}^{\mathbf{g}_N} F_N(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}), \end{aligned}$$

where $\hat{X}_{j;t_0} = Z_j \prod_{i=1}^n A_{i;t_0}^{b_{ij}^{t_0}}$. So we can get the following equalities:

$$\begin{aligned} F_M(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}) &= P_M/A_{t_0}^{\mathbf{d}_M + \mathbf{g}_M}, \\ F_N(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}) &= P_N/A_{t_0}^{\mathbf{d}_N + \mathbf{g}_N}. \end{aligned}$$

Because P_M and P_N are not divisible by any $A_{i;t_0}$, we know that $\mathbf{d}_M + \mathbf{g}_M$ and $\mathbf{d}_N + \mathbf{g}_N$ are the d -vectors of $F_M(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0})$ and $F_N(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0})$ with respect to t_0 . By $F_M = F_N$, we have

$$F_M(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}) = F_N(\hat{X}_{1;t_0}, \dots, \hat{X}_{n;t_0}),$$

which implies that $\mathbf{d}_M + \mathbf{g}_M = \mathbf{d}_N + \mathbf{g}_N$ and $P_M = P_N$.

By $P_M = P_N$ and applying Proposition 5.6 to M and N , we get $\mathbf{d}_M = \mathbf{d}_N$. Then by the equality (6.1), we know that $M = N$. In particular, we have $\mathbf{g}_M = \mathbf{g}_N$. This completes the proof. \square

6.2. F -polynomials of non-initial cluster variables are irreducible. In this subsection, we prove that the F -polynomials of non-initial cluster variables are irreducible in $\mathbb{K}[Z_1, \dots, Z_n]$.

Theorem 6.2. *Let \mathcal{U} be an upper cluster algebra with principal coefficients at t_0 and $A_{k;t}$ a non-initial cluster variable of \mathcal{U} . Then the F -polynomial $F_{k;t}$ of $A_{k;t}$ is irreducible in $\mathbb{K}[Z_1, \dots, Z_n]$.*

Proof. We claim that $F_{k;t}$ is not a constant. Otherwise, the f -compatibility degree $(A_{i;t_0} || A_{k;t})_f = 0$ for $i = 1, \dots, n$. Then by Proposition 2.30 (ii), we get that $A_{k;t}$ is compatible with any cluster variable in A_{t_0} . Thus $A_{k;t} \in A_{t_0}$, by Theorem 2.33. This contradicts that $A_{k;t}$ is a non-initial cluster variable of \mathcal{U} . So $F_{k;t}$ is not a constant.

Since $F_{k;t}$ is not a constant, it can not be invertible in $\mathbb{K}[Z_1, \dots, Z_n]$. So we can discuss its irreducibility. Assume there exist two polynomials $F_1, F_2 \in \mathbb{K}[Z_1, \dots, Z_n]$ such that

$$(6.2) \quad F_{k;t}(Z_1, \dots, Z_n) = F_1(Z_1, \dots, Z_n)F_2(Z_1, \dots, Z_n).$$

We will show that either $F_1 \in \mathbb{K}^\times$ or $F_2 \in \mathbb{K}^\times$.

For $1 \leq j \leq n$, put $\hat{X}_j = Z_j \prod_{i=1}^n A_{i;t_0}^{b_{ij}^{t_0}}$. Consider the Laurent polynomials $F_1(\hat{X}_1, \dots, \hat{X}_n)$ and $F_2(\hat{X}_1, \dots, \hat{X}_n)$ in $\mathbb{K}[Z_1, \dots, Z_n][A_{1;t_0}^{\pm 1}, \dots, A_{n;t_0}^{\pm 1}]$. We choose two vectors $\mathbf{d}_1, \mathbf{d}_2$ in \mathbb{N}^n such that

$$M_1 := A_{t_0}^{\mathbf{d}_1} F_1(\hat{X}_1, \dots, \hat{X}_n) \quad \text{and} \quad M_2 := A_{t_0}^{\mathbf{d}_2} F_2(\hat{X}_1, \dots, \hat{X}_n)$$

are polynomials in $\mathbb{K}[Z_1, \dots, Z_n][A_{1;t_0}, \dots, A_{n;t_0}] \subseteq \mathcal{U}$. By the local unique factorization property, there exist unique cluster monomials $N_1 = A_{t_0}^{\mathbf{v}_1}, N_2 = A_{t_0}^{\mathbf{v}_2}$ such that

$$L_1 := M_1/N_1 = A_{t_0}^{\mathbf{d}_1 - \mathbf{v}_1} F_1(\hat{X}_1, \dots, \hat{X}_n)$$

and

$$L_2 := M_2/N_2 = A_{t_0}^{\mathbf{d}_2 - \mathbf{v}_2} F_2(\hat{X}_1, \dots, \hat{X}_n)$$

are in \mathcal{U} and satisfy

$$(A_{j;t_0} || L_1)_v = 0 = (A_{j;t_0} || L_2)_v$$

for $j = 1, \dots, n$. Then by Corollary 4.10, we have

$$(A_{j;t_0} || L_1 L_2)_v = (A_{j;t_0} || L_1)_v + (A_{j;t_0} || L_2)_v = 0$$

for $j = 1, \dots, n$.

By Theorem 2.26 (i), we know that the cluster variable $A_{k;t}$ has the following form:

$$A_{k;t} = A_{t_0}^{\mathbf{g}} \cdot F_{k;t}(\hat{X}_1, \dots, \hat{X}_n),$$

where $\mathbf{g} \in \mathbb{Z}^n$. By the equality (6.2), we have that

$$F_{k;t}(\hat{X}_1, \dots, \hat{X}_n) = F_1(\hat{X}_1, \dots, \hat{X}_n)F_2(\hat{X}_1, \dots, \hat{X}_n).$$

So we know that

$$A_{k;t}/(L_1 L_2) = A_{t_0}^{\mathbf{g} + \mathbf{v}_1 + \mathbf{v}_2 - \mathbf{d}_1 - \mathbf{d}_2}.$$

Thus we have $A_{k;t} = A_{t_0}^{\mathbf{y}} (L_1 L_2)$, where $\mathbf{v} = \mathbf{g} + \mathbf{v}_1 + \mathbf{v}_2 - \mathbf{d}_1 - \mathbf{d}_2$.

Now we show that $\mathbf{v} = 0$ and $A_{k;t} = L_1 L_2$. We choose two vectors \mathbf{v}^+ and \mathbf{v}^- in \mathbb{N}^n such that $\mathbf{v} = \mathbf{v}^+ - \mathbf{v}^-$. Then by

$$A_{k;t} = A_{t_0}^{\mathbf{v}} (L_1 L_2) = A_{t_0}^{\mathbf{v}^+ - \mathbf{v}^-} (L_1 L_2),$$

we get that

$$(6.3) \quad M := A_{t_0}^{\mathbf{v}^-} A_{k;t} = A_{t_0}^{\mathbf{v}^+} L_1 L_2.$$

Since $F_{k;t}$ is not a constant, we know that $A_{k;t}$ is not an initial cluster variable. Then by Theorem 5.4, we have $(A_{j;t_0} || A_{k;t})_v = 0$ for $j = 1, \dots, n$. On the other hand, we know that $L_1 L_2$ also satisfies that $(A_{j;t_0} || L_1 L_2)_v = 0$ for $j = 1, \dots, n$. So the equality (6.3) gives two local factorizations of M with respect to t_0 . Then by the uniqueness in Theorem 3.7, we get that

$$\mathbf{v}^+ = \mathbf{v}^- \quad \text{and} \quad A_{k;t} = L_1 L_2.$$

By Theorem 2.35 (ii), we know that $\mathcal{U}^\times = \{\lambda Z_1^{c_1} \cdots Z_n^{c_n} | \lambda \in \mathbb{K}^\times, c_1, \dots, c_n \in \mathbb{Z}\}$ and that $A_{k;t}$ is irreducible in \mathcal{U} . So the factorization $A_{k;t} = L_1 L_2$ implies either $L_1 = \lambda Z_1^{c_1} \cdots Z_n^{c_n}$ or $L_2 = \lambda Z_1^{c_1} \cdots Z_n^{c_n}$ for some $\lambda \in \mathbb{K}^\times$ and $(c_1, \dots, c_n) \in \mathbb{Z}^n$. So either

$$F_1(Z_1, \dots, Z_n) = L_1|_{A_{1;t_0}=\dots=A_{n;t_0}=1} = \lambda Z_1^{c_1} \cdots Z_n^{c_n}$$

or

$$F_2(Z_1, \dots, Z_n) = L_2|_{A_{1;t_0}=\dots=A_{n;t_0}=1} = \lambda Z_1^{c_1} \cdots Z_n^{c_n}$$

for some $\lambda \in \mathbb{K}^\times$ and $(c_1, \dots, c_n) \in \mathbb{Z}^n$. Because both F_1 and F_2 are polynomials in $\mathbb{K}[Z_1, \dots, Z_n]$, we must have $c_1, \dots, c_n \geq 0$. Now by Proposition 2.22, we know that $F_{k;t} = F_1 F_2$ is not divisible by any Z_j . So we must have $c_1 = \dots = c_n = 0$. Thus either $F_1 = \lambda$ or $F_2 = \lambda$ for some $\lambda \in \mathbb{K}^\times$. So $F_{k;t}$ is irreducible in $\mathbb{K}[Z_1, \dots, Z_n]$. \square

Remark 6.3. Garcia Elsener *et al.* in [GELS19] prove that the F -polynomials of non-initial cluster variables are irreducible for *factorial principal coefficient cluster algebras*. In fact, their proof still works for factorial principal coefficient upper cluster algebras. Note that the factoriality of principal coefficient upper cluster algebras is no longer a problem, thanks to Theorem 4.8. Thus one can also use the method in [GELS19] to show the irreducibility of F -polynomials of non-initial cluster variables.

7. APPLICATION TO COMBINATORICS OF CLUSTER POISSON VARIABLES

7.1. Parametrization of \mathcal{A} -exchange pairs by cluster Poisson variables.

In this subsection, we give several equivalent characterizations of when two cluster Poisson variables are equal. As an application, we prove that the \mathcal{A} -exchange pairs are parameterized by the cluster Poisson variables.

Proposition 7.1. *Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and initial seed t_0 . Let M_1, M_2 be two cluster monomials of \mathcal{U} and N_1, N_2 two different cluster monomials in t_0 . If*

$$p_1 M_1 + p_2 M_2 = p'_1 N_1 + p'_2 N_2$$

holds for some $p_1, p_2, p'_1, p'_2 \in \mathbb{P}$, then either $(M_1, M_2, p_1, p_2) = (N_1, N_2, p'_1, p'_2)$ or $(M_1, M_2, p_1, p_2) = (N_2, N_1, p'_2, p'_1)$ holds.

Proof. Since both N_1 and N_2 are cluster monomials in A_{t_0} , we have

$$(A_{i;t_0} || p'_1 N_1)_d = (A_{i;t_0} || N_1)_d \leq 0$$

and

$$(A_{i;t_0} || p'_2 N_2)_d = (A_{i;t_0} || N_2)_d \leq 0$$

for $i = 1, \dots, n$. By the positivity of the Laurent phenomenon and since the coefficients in $p_1M_1 + p_2M_2$ and $p'_1N_1 + p'_2N_2$ are positive, we have

$$\begin{aligned} & \max\{(A_{i;t_0}||M_1)_d, (A_{i;t_0}||M_2)_d\} \\ &= \max\{(A_{i;t_0}||p_1M_1)_d, (A_{i;t_0}||p_2M_2)_d\} \\ &= (A_{i;t_0}||p_1M_1 + p_2M_2)_d = (A_{i;t_0}||p'_1N_1 + p'_2N_2)_d \\ &= \max\{(A_{i;t_0}||p'_1N_1)_d, (A_{i;t_0}||p'_2N_2)_d\} \leq 0, \end{aligned}$$

for $i = 1, \dots, n$.

We claim that M_1 and M_2 are cluster monomials in t_0 . Let $A_{k;t}$ be a cluster variable appearing in M_1 or M_2 . By Corollary 2.32 (ii) and

$$\max\{(A_{i;t_0}||M_1)_d, (A_{i;t_0}||M_2)_d\} \leq 0,$$

we know that $A_{k;t}$ is compatible with $A_{i;t_0}$ for any $i = 1, \dots, n$. Then by Theorem 2.33, we get $A_{k;t} \in A_{t_0}$. So M_1 and M_2 are cluster monomials in t_0 .

Hence, the polynomial $p_1M_1 + p_2M_2$ is actually a polynomial in variables from A_{t_0} . Since N_1 and N_2 are two different cluster monomials in t_0 , we must have $M_1 \neq M_2$. Otherwise, $p'_1N_1 + p'_2N_2$ can not be equal to $p_1M_1 + p_2M_2$. Now the result follows from the algebraic independence of $A_{1;t_0}, \dots, A_{n;t_0}$. \square

Proposition 7.2. *Let \mathcal{U} be an upper cluster algebra with coefficient semifield \mathbb{P} and initial seed t_0 . Let $A_{k;t}$ be a cluster variable of \mathcal{U} . Then the following two statements are equivalent.*

- (i) $(A_{j;t_0}, A_{k;t})$ is an \mathcal{A} -exchange pair associated with the mutation $t_1 := \mu_j(t_0)$;
- (ii) $(A_{j;t_0}||A_{k;t})_d > 0$ and $(A_{i;t_0}||A_{k;t})_d = 0$ for any $i \neq j$.

Proof. (i) \implies (ii): This is clear.

(ii) \implies (i): By Proposition 2.30 (4) and the assumption, we get that $A_{k;t} \notin A_{t_0}$ and $\{A_{k;t}\} \cup (A_{t_0} \setminus \{A_{j;t_0}\})$ is a compatible set with n elements. Then by Theorem 2.33, we know that $\{A_{k;t}\} \cup (A_{t_0} \setminus \{A_{j;t_0}\})$ is a cluster of \mathcal{U} , say $A_{t_2} = \{A_{k;t}\} \cup (A_{t_0} \setminus \{A_{j;t_0}\})$. Since the seeds t_2 and t_0 have exactly $n - 1$ common cluster variables, we get that there is an edge between t_2 and t_0 in the exchange graph of \mathcal{U} , by Corollary 2.18. Then the result follows. \square

Proposition 7.3. *Let \mathcal{U} and \mathcal{U}' be any two upper cluster algebras with the same initial exchange matrix B_{t_0} at t_0 . For any vertex t of \mathbb{T}_n , denote A_t (respectively, A'_t) be the cluster of \mathcal{U} (respectively, \mathcal{U}') at t . Then we have*

- (i) $A_{i;t_1} = A_{j;t_2}$ in \mathcal{U} if and only if $A'_{i;t_1} = A'_{j;t_2}$ in \mathcal{U}' ;
- (ii) The map $\rho : A_{k;t} \mapsto A'_{k;t}$ gives a bijection from the set of cluster variables of \mathcal{U} to that of \mathcal{U}' ;
- (iii) ρ induces a bijection from the set of \mathcal{A} -exchange pairs of \mathcal{U} to that of \mathcal{U}' .

Proof. (i) Because \mathcal{U} and \mathcal{U}' have the same initial exchange matrix B_{t_0} at t_0 , we know that they have the same exchange matrix at each vertex t of \mathbb{T}_n . Notice that both $(A_{i;t_1}||A_{j;t_2})_d$ and $(A'_{i;t_1}||A'_{j;t_2})_d$ are defined by the (i, j) -entry of the D -matrix $D_{t_2}^{B_{t_1}; t_1}$, which is independent of the choice of coefficient semifield. So we have $(A_{i;t_1}||A_{j;t_2})_d = (A'_{i;t_1}||A'_{j;t_2})_d$. By Proposition 2.30 (4), we know that $A_{i;t_1} = A_{j;t_2}$ in \mathcal{U} if and only if $(A_{i;t_1}||A_{j;t_2})_d = -1$ if and only if $(A'_{i;t_1}||A'_{j;t_2})_d = -1$ if and only if $A'_{i;t_1} = A'_{j;t_2}$ in \mathcal{U}' .

(ii) follows from (i) and (iii) follows from (ii) and the characterization of \mathcal{A} -exchange pairs in Proposition 7.2. \square

Lemma 7.4. *Let \mathcal{U}_{uc} be an upper cluster algebra with universal coefficient semifield and t, t_0 two seeds of \mathcal{U}_{uc} . Let $t' = \mu_k(t)$ and $t_1 = \mu_j(t_0)$. Let $\frac{X_{k;t}}{1+X_{k;t}}M_1 + \frac{1}{1+X_{k;t}}M_2$ be the k -th exchange binomial of t and $\frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2$ the j -th exchange binomial of t_0 . If*

$$\frac{X_{k;t}}{1+X_{k;t}}M_1 + \frac{1}{1+X_{k;t}}M_2 = \frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2,$$

then either

$$(A_{k;t}, A_{k;t'}) = (A_{j;t_0}, A_{j;t_1}) \quad \text{or} \quad (A_{k;t}, A_{k;t'}) = (A_{j;t_1}, A_{j;t_0}).$$

Proof. By the assumption, we know that

$$A_{k;t}A_{k;t'} = \frac{X_{k;t}}{1+X_{k;t}}M_1 + \frac{1}{1+X_{k;t}}M_2 = \frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2.$$

The above equality can be viewed as the Laurent expansion of $A_{k;t}A_{k;t'}$ with respect to A_{t_0} . Clearly, this Laurent expansion is a polynomial in A_{t_0} .

In the following proof, we distinguish two cases. Case (a): $(A_{i;t_0} || A_{k;t})_d \leq 0$ for any $i = 1, \dots, n$. Case (b): there exists some i_0 such that $(A_{i_0;t_0} || A_{k;t})_d > 0$.

Case (a): If $(A_{i;t_0} || A_{k;t})_d \leq 0$ for any $i = 1, \dots, n$, then $A_{k;t}$ is compatible with any cluster variable in A_{t_0} . By Theorem 2.33, we know that A_{t_0} is a maximal compatible set. So $A_{k;t} \in A_{t_0}$, say $A_{k;t} = A_{i_0;t_0}$. We will show that

$$i_0 = j \quad \text{and} \quad A_{i_0;t_0} = A_{j;t_0}.$$

We know that

$$A_{k;t'} = A_{k;t}^{-1} \cdot \left(\frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2 \right) = A_{i_0;t_0}^{-1} \cdot \left(\frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2 \right),$$

which is the Laurent expansion of $A_{k;t'}$ with respect to A_{t_0} . So we have

$$(A_{i_0;t_0} || A_{k;t'})_d = 1 > 0 \quad \text{and} \quad (A_{i;t_0} || A_{k;t'})_d = 0$$

for any $i \neq i_0$. Then by Proposition 7.2, we get that

$$A_{k;t'} = A_{i_0;t_2},$$

where $t_2 = \mu_{i_0}(t_0)$.

Let $\frac{X_{i_0;t_0}}{1+X_{i_0;t_0}}L_1 + \frac{1}{1+X_{i_0;t_0}}L_2$ be the i_0 -th exchange binomial of t_0 . By Proposition 7.1 and the following equality,

$$\frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2 = A_{k;t}A_{k;t'} = A_{i_0;t_0}A_{i_0;t_2} = \frac{X_{i_0;t_0}}{1+X_{i_0;t_0}}L_1 + \frac{1}{1+X_{i_0;t_0}}L_2,$$

we get that either $X_{j;t_0} = X_{i_0;t_0}$ or $X_{j;t_0} = X_{i_0;t_0}^{-1}$. Since $X_{1;t_0}, \dots, X_{n;t_0}$ are algebraically independent, we must have $X_{j;t_0} = X_{i_0;t_0}$ and $j = i_0$. So

$$A_{k;t} = A_{i_0;t_0} = A_{j;t_0}.$$

Then by the following equality,

$$A_{k;t}A_{k;t'} = \frac{X_{k;t}}{1+X_{k;t}}M_1 + \frac{1}{1+X_{k;t}}M_2 = \frac{X_{j;t_0}}{1+X_{j;t_0}}N_1 + \frac{1}{1+X_{j;t_0}}N_2 = A_{j;t_0}A_{j;t_1},$$

we get $A_{k;t'} = A_{j;t_1}$ and thus $(A_{k;t}, A_{k;t'}) = (A_{j;t_0}, A_{j;t_1})$.

Case (b): There exists some i_0 such that $(A_{i_0;t_0} \| A_{k;t})_d > 0$. By the fact that the expansion of $A_{k;t} A_{k;t'}$ with respect to A_{t_0} is a polynomial, we know that

$$0 \geq (A_{i_0;t_0} \| A_{k;t} A_{k;t'}) = (A_{i_0;t_0} \| A_{k;t})_d + (A_{i_0;t_0} \| A_{k;t'})_d.$$

So we must have $(A_{i_0;t_0} \| A_{k;t'})_d < 0$. Then by Proposition 2.30 (4), we get

$$A_{k;t'} = A_{i_0;t_0}.$$

Similarly to the proof in Case (a), we can prove that $i_0 = j$ and thus

$$A_{k;t'} = A_{i_0;t_0} = A_{j;t_0}.$$

Then $A_{k;t} = A_{j;t_1}$ follows from the following equality

$$A_{k;t} A_{k;t'} = \frac{X_{k;t}}{1+X_{k;t}} M_1 + \frac{1}{1+X_{k;t}} M_2 = \frac{X_{j;t_0}}{1+X_{j;t_0}} N_1 + \frac{1}{1+X_{j;t_0}} N_2 = A_{j;t_0} A_{j;t_1}.$$

So we have $(A_{k;t}, A_{k;t'}) = (A_{j;t_1}, A_{j;t_0})$. This completes the proof. \square

Theorem 7.5. *Let \mathcal{U}_{uc} be an upper cluster algebra with universal coefficient semifield and t, t_0 two seeds of \mathcal{U}_{uc} . Let $\frac{X_{k;t}}{1+X_{k;t}} M_1 + \frac{1}{1+X_{k;t}} M_2$ be the k -th exchange binomial of t and $\frac{X_{j;t_0}}{1+X_{j;t_0}} N_1 + \frac{1}{1+X_{j;t_0}} N_2$ the j -th exchange binomial of t_0 . Then the following statements are equivalent.*

- (i) $X_{k;t} = X_{j;t_0}$;
- (ii) $(A_{k;t}, M_1, M_2) = (A_{j;t_0}, N_1, N_2)$;
- (iii) $(A_{k;t}, \{A_{i;t} | b_{ik}^t \neq 0\}) = (A_{j;t_0}, \{A_{i;t_0} | b_{ij}^{t_0} \neq 0\})$;
- (iv) $A_{k;t} = A_{j;t_0}$ and $\{A_{i;t} | b_{ik}^t \neq 0\} \subseteq A_{t_0}$;
- (v) $(A_{k;t}, A_{k;t'}) = (A_{j;t_0}, A_{j;t_1})$, where $t' = \mu_k(t)$ and $t_1 = \mu_j(t_0)$;
- (vi) $A_{k;t} = A_{j;t_0}$ and $\frac{X_{k;t}}{1+X_{k;t}} M_1 + \frac{1}{1+X_{k;t}} M_2 = \frac{X_{j;t_0}}{1+X_{j;t_0}} N_1 + \frac{1}{1+X_{j;t_0}} N_2$.

Proof. (i) \implies (ii): By Theorem 2.26 (ii), we know that

$$X_{k;t} = X_{t_0}^{\mathbf{c}_{k;t}^{B_{t_0};t_0}} \cdot \prod_{i=1}^n (F_{i;t}^{B_{t_0};t_0}(X_{1;t_0}, \dots, X_{n;t_0}))^{b_{ik}^t}.$$

By $X_{k;t} = X_{j;t_0}$ and Proposition 2.22, we get that $\mathbf{c}_{k;t}^{B_{t_0};t_0}$ is equal to the j -th column vector of I_n and

$$\prod_{i=1}^n (F_{i;t}^{B_{t_0};t_0}(X_{1;t_0}, \dots, X_{n;t_0}))^{b_{ik}^t} = 1.$$

So we get that

$$(7.1) \quad \prod_{b_{ik}^t > 0} (F_{i;t}^{B_{t_0};t_0})^{b_{ik}^t} = \prod_{b_{ik}^t < 0} (F_{i;t}^{B_{t_0};t_0})^{-b_{ik}^t}.$$

Let $\bar{\mathcal{U}}$ be the upper cluster algebra with principal coefficients at t_0 and with initial exchange matrix B_{t_0} . We use $(\bar{B}_t, \bar{X}_t, \bar{A}_t)$ to denote the seed of $\bar{\mathcal{U}}$ at vertex t . By $\bar{B}_{t_0} = B_{t_0}$, we know that $\bar{B}_t = B_t$ for any vertex t . For any cluster monomial $L = A_{1;t}^{v_1} \cdots A_{n;t}^{v_n}$ of \mathcal{U}_{uc} , we denote by

$$\bar{L} = \bar{A}_{1;t}^{v_1} \cdots \bar{A}_{n;t}^{v_n}$$

the corresponding cluster monomial of \bar{U} and we denote by $F_{\bar{L}}^{B_{t_0}; t_0}$ the F -polynomial of \bar{L} . By the assumption, we know that $M_1 = \prod_{b_{i_k}^t > 0} A_{i;t}^{b_{i_k}^t}$ and $M_2 = \prod_{b_{i_k}^t < 0} A_{i;t}^{-b_{i_k}^t}$. So we have

$$\bar{M}_1 = \prod_{b_{i_k}^t > 0} \bar{A}_{i;t}^{b_{i_k}^t} \quad \text{and} \quad \bar{M}_2 = \prod_{b_{i_k}^t < 0} \bar{A}_{i;t}^{-b_{i_k}^t}.$$

We claim that M_1 and M_2 are cluster monomials in A_{t_0} . In order to prove this claim, we first show that \bar{M}_1 and \bar{M}_2 are cluster monomials in \bar{A}_{t_0} .

By Theorem 5.4 (iii), we can always write the cluster monomial \bar{M}_s ($s = 1, 2$) as

$$\bar{M}_s = \bar{M}_s'' \cdot \bar{M}_s',$$

where \bar{M}_s'' is a cluster monomial in initial cluster variables of \bar{U} and \bar{M}_s' is a cluster monomial in non-initial cluster variables of \bar{U} . By the equality (7.1), we know that $F_{\bar{M}_1}^{B_{t_0}; t_0} = F_{\bar{M}_2}^{B_{t_0}; t_0}$. Because \bar{M}_1'' and \bar{M}_2'' are cluster monomials in initial cluster variables, we have $F_{\bar{M}_1''}^{B_{t_0}; t_0} = F_{\bar{M}_2''}^{B_{t_0}; t_0} = 1$. So

$$F_{\bar{M}_1'}^{B_{t_0}; t_0} = F_{\bar{M}_1}^{B_{t_0}; t_0} = F_{\bar{M}_2}^{B_{t_0}; t_0} = F_{\bar{M}_2'}^{B_{t_0}; t_0}.$$

Then by Theorem 6.1 and the fact that \bar{M}_1' and \bar{M}_2' are two cluster monomials in non-initial cluster variables, we get $\bar{M}_1' = \bar{M}_2'$. Because \bar{M}_1 and \bar{M}_2 have no common factor in $\mathbb{Z}[\bar{A}_{1;t}, \dots, \bar{A}_{n;t}]$, the same holds for \bar{M}_1' and \bar{M}_2' . So $\bar{M}_1' = \bar{M}_2'$ implies that

$$\bar{M}_1' = \bar{M}_2' = 1.$$

Thus $\bar{M}_1 = \bar{M}_1''$ and $\bar{M}_2 = \bar{M}_2''$ are cluster monomials in \bar{A}_{t_0} . Then by Proposition 7.3 (i), we get that M_1 and M_2 are cluster monomials in A_{t_0} .

By Proposition 2.9, we know that $\hat{S} = \{(B_t, \hat{X}_t)\}_{t \in \mathbb{T}_n}$ forms an \mathcal{X} -seed pattern, where $\hat{X}_t = (\hat{X}_{1;t}, \dots, \hat{X}_{n;t})$ is given by $\hat{X}_{k;t} := X_{k;t} \prod_{i=1}^n A_{i;t}^{b_{i_k}^t}$. Since \mathcal{U}_{uc} is an upper cluster algebra with universal coefficient semifield, $X_{k;t} = X_{j;t_0}$ implies that $\hat{X}_{k;t} = \hat{X}_{j;t_0}$. So we have $X_{k;t} \prod_{i=1}^n A_{i;t}^{b_{i_k}^t} = X_{j;t_0} \prod_{i=1}^n A_{i;t_0}^{b_{i_j}^{t_0}}$. Thus we get that

$$\frac{M_1}{M_2} = \prod_{i=1}^n A_{i;t}^{b_{i_k}^t} = \prod_{i=1}^n A_{i;t_0}^{b_{i_j}^{t_0}} = \frac{N_1}{N_2}.$$

Since M_1, M_2, N_1, N_2 are monomials in $\mathbb{Z}[A_{1;t_0}, \dots, A_{n;t_0}]$ and (M_1, M_2) and (N_1, N_2) are relatively prime in $\mathbb{Z}[A_{1;t_0}, \dots, A_{n;t_0}]$, we get

$$M_1 = N_1 \quad \text{and} \quad M_2 = N_2.$$

Then by $X_{k;t} = X_{j;t_0}$, we have

$$\frac{X_{k;t}}{1 + X_{k;t}} M_1 + \frac{1}{1 + X_{k;t}} M_2 = \frac{X_{j;t_0}}{1 + X_{j;t_0}} N_1 + \frac{1}{1 + X_{j;t_0}} N_2.$$

By Lemma 7.4, we get that

$$A_{k;t} = A_{j;t_0} \quad \text{or} \quad A_{k;t} = A_{j;t_1},$$

where $t_1 = \mu_j(t_0)$.

Now we rule out the possibility of $A_{k;t} = A_{j;t_1}$ with the help of the condition $X_{k;t} = X_{j;t_0}$. Assume by contradiction that $A_{k;t} = A_{j;t_1}$. Then the k -th column of the G -matrix $G_t^{B_{t_0};t_0}$ equals the j -th column of $G_{t_1}^{B_{t_0};t_0}$. In particular, we have $g_{jk;t}^{B_{t_0};t_0} = g_{jj;t_1}^{B_{t_0};t_0}$. By $t_1 = \mu_j(t_0)$, we get that

$$g_{jk;t}^{B_{t_0};t_0} = g_{jj;t_1}^{B_{t_0};t_0} = -1.$$

By Theorem 2.28, we know that

$$(7.2) \quad S^{-1}(G_t^{B_{t_0};t_0})^T S \cdot C_t^{B_{t_0};t_0} = I_n,$$

where $S = \text{diag}(s_1, \dots, s_n)$ is a skew-symmetrizer of B_{t_0} . By $X_{k;t} = X_{j;t_0}$, we know that the k -th column of the C -matrix $C_t^{B_{t_0};t_0}$ equals the j -th column of I_n . By comparing the k -th column of the two sides of the equality (7.2), we get that the j -th column of $S^{-1}(G_t^{B_{t_0};t_0})^T S$ equals the k -th column of I_n . In particular, the (k, j) -entry $s_k^{-1} \cdot g_{jk;t}^{B_{t_0};t_0} \cdot s_j$ of $S^{-1}(G_t^{B_{t_0};t_0})^T S$ equals 1, and thus

$$g_{jk;t}^{B_{t_0};t_0} = \frac{s_k}{s_j} > 0.$$

This contradicts $g_{jk;t}^{B_{t_0};t_0} = -1$. So $A_{k;t} \neq A_{j;t_1}$ and thus $A_{k;t} = A_{j;t_0}$.

Hence, $X_{k;t} = X_{j;t_0}$ implies $(A_{k;t}, M_1, M_2) = (A_{j;t_0}, N_1, N_2)$.

(ii) \implies (iii): It suffices to show that

$$\{A_{i;t} | b_{ik}^t \neq 0\} = \{A_{i;t_0} | b_{ij}^{t_0} \neq 0\}.$$

Let \mathbb{A} be the set of cluster variables of \mathcal{U}_{uc} and $P \in \mathbb{A}$. We first claim that $(P || M_1 \cdot M_2)_d < 0$ if and only if P is a cluster variable appearing in the cluster monomial $M_1 \cdot M_2$.

If P is a cluster variable appearing in the cluster monomial $M_1 \cdot M_2$, then it is easy to see that $(P || M_1 \cdot M_2)_d < 0$. Conversely, if $(P || M_1 \cdot M_2)_d < 0$, then by Corollary 2.32 (i), we know that P is a cluster variable appearing in the cluster monomial $M_1 \cdot M_2$.

By the discussion above, we get

$$\{A_{i;t} | b_{ik}^t \neq 0\} = \{P \in \mathbb{A} | (P || M_1 \cdot M_2)_d < 0\}.$$

Similarly, we have

$$\{A_{i;t_0} | b_{ij}^{t_0} \neq 0\} = \{P \in \mathbb{A} | (P || N_1 \cdot N_2)_d < 0\}.$$

By $(A_{k;t}, M_1, M_2) = (A_{j;t_0}, N_1, N_2)$, we have $M_1 \cdot M_2 = N_1 \cdot N_2$. Thus we get

$$\{A_{i;t} | b_{ik}^t \neq 0\} = \{A_{i;t_0} | b_{ij}^{t_0} \neq 0\}.$$

(iii) \implies (iv): This is clear.

(iv) \implies (v): We know that

$$A_{k;t'} = A_{k;t}^{-1} \left(\frac{X_{k;t}}{1 + X_{k;t}} M_1 + \frac{1}{1 + X_{k;t}} M_2 \right).$$

By $A_{k;t} = A_{j;t_0}$ and the fact that M_1, M_2 are monomials in variables from

$$\{A_{i;t} | b_{ik}^t \neq 0\} \subseteq A_{t_0},$$

we know that

$$A_{k;t'} = A_{k;t}^{-1} \left(\frac{X_{k;t}}{1 + X_{k;t}} M_1 + \frac{1}{1 + X_{k;t}} M_2 \right) = A_{j;t_0}^{-1} \left(\frac{X_{k;t}}{1 + X_{k;t}} M_1 + \frac{1}{1 + X_{k;t}} M_2 \right)$$

can be viewed as the Laurent expansion of $A_{k;t'}$ with respect to A_{t_0} . Thus we know that

$$(A_{j;t_0} || A_{k;t'})_d = 1 > 0 \quad \text{and} \quad (A_{i;t_0} || A_{k;t'})_d = 0,$$

for any $i \neq j$. Then by Proposition 7.2, we get $A_{k;t'} = A_{j;t_1}$. So

$$(A_{k;t}, A_{k;t'}) = (A_{j;t_0}, A_{j;t_1}).$$

(v) \implies (vi): This is clear.

(vi) \implies (i): If the j -th column of B_{t_0} is zero, then the same holds for any exchange matrices of \mathcal{U}_{uc} . In this case, the action of μ_j on any seed is very clear. We can see that $A_{k;t} = A_{j;t_0}$ implies that $k = j$ and $X_{k;t} = X_{j;t_0}$.

If the j -th column of B_{t_0} is non-zero, then N_1 and N_2 are two different cluster monomials in A_{t_0} . Then by Proposition 7.1 and the following equality,

$$\frac{X_{j;t_0}}{1 + X_{j;t_0}} N_1 + \frac{1}{1 + X_{j;t_0}} N_2 = \frac{X_{k;t}}{1 + X_{k;t}} M_1 + \frac{1}{1 + X_{k;t}} M_2,$$

we get $\frac{X_{k;t}}{1 + X_{k;t}} = \frac{X_{j;t_0}}{1 + X_{j;t_0}}$ or $\frac{X_{k;t}}{1 + X_{k;t}} = \frac{1}{1 + X_{j;t_0}}$. Namely, we get that

$$X_{k;t} = X_{j;t_0} \quad \text{or} \quad X_{k;t} = X_{j;t_0}^{-1}.$$

Similar to the argument for the statement (i) \implies (ii), now we rule out the possibility of $X_{k;t} = X_{j;t_0}^{-1}$ with the help of the condition $A_{k;t} = A_{j;t_0}$. Assume by contradiction that $X_{k;t} = X_{j;t_0}^{-1}$. Then the k -th column of $C_t^{B_{t_0};t_0}$ is the j -th column of $-I_n$, by Theorem 2.26 (ii). So we have

$$c_{jk;t}^{B_{t_0};t_0} = -1 < 0.$$

By Theorem 2.28, we know that

$$(7.3) \quad (G_t^{B_{t_0};t_0})^T (S C_t^{B_{t_0};t_0} S^{-1}) = I_n,$$

where $S = \text{diag}(s_1, \dots, s_n)$ is a skew-symmetrizer of B_{t_0} .

By $A_{k;t} = A_{j;t_0}$, we know that the k -th column of $G_t^{B_{t_0};t_0}$ equals the j -th column vector of I_n . So the k -th row of $(G_t^{B_{t_0};t_0})^T$ equals the j -th row of I_n . By comparing the k -th row of the two sides of the equality (7.3), we know that the j -th row of $S C_t^{B_{t_0};t_0} S^{-1}$ equals the k -th row of I_n . In particular, the (j, k) -entry $s_j c_{jk;t}^{B_{t_0};t_0} s_k^{-1}$ of $S C_t^{B_{t_0};t_0} S^{-1}$ equals 1 and thus

$$c_{jk;t}^{B_{t_0};t_0} = \frac{s_k}{s_j} > 0.$$

This contradicts $c_{jk;t}^{B_{t_0};t_0} = -1 < 0$. So $X_{k;t} \neq X_{j;t_0}^{-1}$ and thus $X_{k;t} = X_{j;t_0}$. \square

Let \mathcal{U} be an upper cluster algebra and \mathcal{X} a cluster Poisson algebra. We say that \mathcal{U} and \mathcal{X} have the *same type* if they have the same exchange matrix at a vertex t_0 of \mathbb{T}_n .

Theorem 7.6. *Let \mathcal{X} be a cluster Poisson algebra and \mathcal{U} an upper cluster algebra of the same type as \mathcal{X} . Denote \overline{X}_t the Poisson cluster of \mathcal{X} at t to distinguish the \mathcal{X} -cluster X_t of \mathcal{U} at t . Each mutation $t_k = \mu_k(t)$ gives an \mathcal{X} -exchange pair $(\overline{X}_{k;t}, \overline{X}_{k;t_k})$ of \mathcal{X} and an \mathcal{A} -exchange pair $(A_{k;t}, A_{k;t_k})$ of \mathcal{U} . Then*

- (i) *The map $\psi : \overline{X}_{k;t} \mapsto (A_{k;t}, A_{k;t_k})$ gives a bijection from the set of cluster Poisson variables of \mathcal{X} to the set of \mathcal{A} -exchange pairs of \mathcal{U} ;*

- (ii) The map $\psi_1 : (\overline{X}_{k;t}, \overline{X}_{k;t_k}) \mapsto (A_{k;t}, A_{k;t_k})$ gives a bijection from the set of \mathcal{X} -exchange pairs of \mathcal{X} to the set of \mathcal{A} -exchange pairs of \mathcal{U} .

Proof. (i) Thanks to Proposition 7.3, we can just assume that $\mathcal{U} = \mathcal{U}_{\text{uc}}$ is an upper cluster algebra with universal coefficient semifield. In this case, the cluster Poisson variables of \mathcal{X} can be viewed as the \mathcal{X} -variables of $\mathcal{U} = \mathcal{U}_{\text{uc}}$. Then (i) follows from Theorem 7.5 (i)(v).

(ii) follows from (i) and the fact that in any \mathcal{X} -exchange pair $(\overline{X}_{k;t}, \overline{X}_{k;t_k})$, one has $\overline{X}_{k;t_k} = \overline{X}_{k;t}^{-1}$. \square

Remark 7.7. For (upper) cluster algebras of finite type, a different proof of Theorem 7.6 can be found in [SB19].

7.2. Exchange graphs of cluster Poisson algebras. Let \mathcal{X} be a cluster Poisson algebra. In this subsection, we prove that the seeds of \mathcal{X} whose Poisson clusters contain particular cluster Poisson variables form a connected subgraph of the exchange graph of \mathcal{X} .

Lemma 7.8. *Let \mathcal{U}_{uc} be an upper cluster algebra with universal coefficient semifield, and let t_0 and t_1 be two seeds of \mathcal{U}_{uc} with $t_1 = \mu_k(t_0)$. For a given $1 \leq j \leq n$, denote $I(t_0) = \{j\} \sqcup \{i | b_{ij}^{t_0} \neq 0\}$ and $I(t_1) = \{j\} \sqcup \{i | b_{ij}^{t_1} \neq 0\}$. If $k \notin I(t_0)$, then we have $X_{j;t_0} = X_{j;t_1}$ and $I(t_0) = I(t_1)$.*

Proof. The results follow from $k \notin I(t_0)$ and the definition of mutation. \square

Lemma 7.9. *Let \mathcal{U}_{uc} be an upper cluster algebra with universal coefficient semifield and t, t_0 two seeds of \mathcal{U}_{uc} . Then $X_{j;t_0} \in X_t$ if and only if $\{A_{j;t_0}\} \sqcup \{A_{i;t_0} | b_{ij}^{t_0} \neq 0\} \subseteq A_t$.*

Proof. \implies : This follows from Theorem 7.5 (i)(iii).

\impliedby : By $\{A_{j;t_0}\} \sqcup \{A_{i;t_0} | b_{ij}^{t_0} \neq 0\} \subseteq A_t$, we know that

$$U := \{A_{j;t_0}\} \sqcup \{A_{i;t_0} | b_{ij}^{t_0} \neq 0\}$$

is a common subset of A_t and A_{t_0} . Then by Theorem 2.17, there exists a sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$ such that the \mathcal{A} -seed at $t_s = \mu_{k_s} \cdots \mu_{k_1}(t_0)$ is equivalent to the \mathcal{A} -seed at t and the cluster variables in U remain unchanged when we do each mutation along the sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$. Set $I = \{j\} \sqcup \{i | b_{ij}^{t_0} \neq 0\}$. We know that $k_\ell \notin I$ for any $\ell = 1, \dots, s$. Applying Lemma 7.8 iteratively, we get $X_{j;t_0} \in X_{t_s}$. Because the two seeds t and t_s are equivalent, we get $X_{j;t_0} \in X_t$. \square

Theorem 7.10. *Let \mathcal{X} be a cluster Poisson algebra. Then the seeds of \mathcal{X} whose Poisson clusters contain particular cluster Poisson variables form a connected subgraph of the exchange graph of \mathcal{X} .*

Proof. Let $\mathcal{U} = \mathcal{U}_{\text{uc}}$ be the upper cluster algebra with universal coefficient semifield of the same type as \mathcal{X} . In this case, the cluster Poisson variables (respectively, Poisson clusters) of \mathcal{X} can be viewed as the \mathcal{X} -variables (respectively, \mathcal{X} -clusters) of $\mathcal{U} = \mathcal{U}_{\text{uc}}$. Now we work on the upper cluster algebra \mathcal{U} .

Fix a subset V of X_{t_0} . Let t be a seed of \mathcal{U} , whose \mathcal{X} -cluster X_t satisfies $V \subseteq X_t$. It suffices for us to show that there exists a sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$ such that the \mathcal{X} -seed at $t_s = \mu_{k_s} \cdots \mu_{k_1}(t_0)$ is equivalent to the \mathcal{X} -seed at t and the \mathcal{X} -variables in V remain unchanged when we do each mutation along the sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$.

Let $U := \bigcup_{X_j; t_0 \in V} (\{A_{j; t_0}\} \sqcup \{A_{i; t_0} | b_{ij}^{t_0} \neq 0\})$. By Lemma 7.9, we know that for a seed t' of \mathcal{U} , its \mathcal{X} -cluster $X_{t'}$ satisfies $V \subseteq X_{t'}$ if and only if its cluster $A_{t'}$ satisfies $U \subseteq A_{t'}$.

By $X_{t_0} \supseteq V$ and $X_t \supseteq V$, we get that $A_{t_0} \supseteq U$ and $A_t \supseteq U$. Then by Theorem 2.17, there exists a sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$ such that the \mathcal{A} -seed at $t_s = \mu_{k_s} \cdots \mu_{k_1}(t_0)$ is equivalent to the \mathcal{A} -seed at t and the cluster variables in U remain unchanged when we do each mutation along the sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$. So the \mathcal{X} -seed at $t_s = \mu_{k_s} \cdots \mu_{k_1}(t_0)$ is equivalent to the \mathcal{X} -seed at t and the \mathcal{X} -variables in V remain unchanged when we do each mutation along the sequence of mutations $(\mu_{k_1}, \dots, \mu_{k_s})$. This completes the proof. \square

REFERENCES

- [BFZ05] A. Berenstein, S. Fomin, and A. Zelevinsky. Cluster algebras III: upper bounds and double Bruhat cells. *Duke Math. J.*, 126(1):1–52, 2005.
- [BM06] Aslak Bakke Buan and Robert Marsh. Cluster-tilting theory. In *Trends in representation theory of algebras and related topics*, Contemp. Math., pages 1–30. Amer. Math. Soc., Providence, RI, 2006.
- [Bri17] T. Bridgeland. Scattering diagrams, Hall algebras and stability conditions. *Algebr. Geom.*, 4(5):523–561, 2017.
- [CC06] P. Caldero and F. Chapoton. Cluster algebras as Hall algebras of quiver representations. *Comment. Math. Helv.*, 81(3):595–616, 2006.
- [CL18] P. Cao and F. Li. Some conjectures on generalized cluster algebras via the cluster formula and D-matrix pattern. *J. Algebra*, 493:57–78, 2018.
- [CL20] P. Cao and F. Li. The enough g-pairs property and denominator vectors of cluster algebras. *Math. Ann.*, 377:1547–1572, 2020.
- [DWZ10] H. Derksen, J. Weyman, and A. Zelevinsky. Quivers with potentials and their representations II: Applications to cluster algebras. *J. Amer. Math. Soc.*, 23(3):749–790, 2010.
- [Fei19a] J. Fei. Combinatorics of F -polynomials. Preprint, arXiv:1909.10151, 2019.
- [Fei19b] J. Fei. Tropical F -polynomials and general presentations. Preprint, arXiv:1911.10513, 2019.
- [FG09] Vladimir V. Fock and Alexander B. Goncharov. Cluster ensembles, quantization and the dilogarithm. *Ann. Sci. Éc. Norm. Supér. (4)*, 42(6):865–930, 2009.
- [FG19a] C. Fu and Y. Gyoda. Compatibility degree of cluster complexes. Preprint, arXiv:1911.07193, 2019.
- [FG19b] S. Fujiwara and Y. Gyoda. Duality between final-seed and initial-seed mutations in cluster algebras. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 15:Paper No. 040, 24 pp., 2019.
- [Fom10] S. Fomin. Total positivity and cluster algebras. In *Proceedings of the International Congress of Mathematicians*, volume II, page 125–145. Hindustan Book Agency, New Delhi, 2010.
- [FZ02] S. Fomin and A. Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [FZ03] S. Fomin and A. Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [FZ07] S. Fomin and A. Zelevinsky. Cluster algebras. IV. Coefficients. *Compos. Math.*, 143:112–164, 2007.
- [GELS19] A. Garcia Elsener, P. Lampe, and D. Smertnig. Factoriality and class groups of cluster algebras. *Advances in Mathematics*, 358:Paper No. 106858, 48 pp., 2019.
- [GHK15] M. Gross, P. Hacking, and S. Keel. Birational geometry of cluster algebras. *Algebr. Geom.*, 2(2):137–175, 2015.
- [GHKK18] M. Gross, P. Hacking, S. Keel, and M. Kontsevich. Canonical bases for cluster algebras. *J. Amer. Math. Soc.*, 31:497–608, 2018.

- [GLS08] C. Geiß, B. Leclerc, and J. Schröer. Preprojective algebras and cluster algebras. In *Trends in representation theory of algebras and related topics*, EMS Ser. Congr. Rep., pages 253–283. Eur. Math. Soc., Zürich, 2008.
- [GLS13] C. Geiß, B. Leclerc, and J. Schröer. Factorial cluster algebras. *Doc. Math.*, 18:249–274, 2013.
- [GY20] Y. Gyoda and T. Yurikusa. F -matrices of cluster algebras from triangulated surfaces. *Ann. Comb.*, 24(4):649–695, 2020.
- [Gyo21] Y. Gyoda. Relation between f -vectors and d -vectors in cluster algebras of finite type or rank 2. *Ann. Comb.*, 25(3):573–594, 2021.
- [KD20] B. Keller and L. Demonet. A survey on maximal green sequences. In *Representation theory and beyond*, Contemp. Math., 758, pages 267–286. Amer. Math. Soc., [Providence], RI, 2020.
- [Kel10] B. Keller. Cluster algebras, quiver representations and triangulated categories. In *Triangulated categories*, London Math. Soc. Lecture Note Ser., pages 76–160. Cambridge Univ. Press, Cambridge, 2010.
- [Kel12] B. Keller. Cluster algebras and derived categories. In *Derived categories in algebraic geometry*, EMS Ser. Congr. Rep., pages 123–183. Eur. Math. Soc., Zürich, 2012.
- [KS08] M. Kontsevich and Y. Soibelman. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations. Preprint, arXiv:0811.2435, 2008.
- [Lec10] B. Leclerc. Cluster algebras and representation theory. In *Proceedings of the International Congress of Mathematicians*, volume IV, pages 2471–2488. Hindustan Book Agency, New Delhi, 2010.
- [LP16] T. Lam and P. Pylyavskyy. Laurent phenomenon algebras. *Cambridge Journal of Mathematics*, 4(1):121–162, 2016.
- [LP22] F. Li and J. Pan. Recurrence formula, positivity and polytope basis in cluster algebras via Newton polytopes. Preprint, arXiv:2201.01440, 2022.
- [LS15] K. Lee and R. Schiffler. Positivity for cluster algebras. *Ann. of Math. (2)*, 182(1):73–125, 2015.
- [Mul14] G. Muller. $\mathcal{A} = \mathcal{U}$ for locally acyclic cluster algebras. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 10:Paper No. 094, 8 pp., 2014.
- [Nag13] K. Nagao. Donaldson-Thomas theory and cluster algebras. *Duke Math. J.*, 162(7):1313–1367, 2013.
- [NZ12] T. Nakanishi and A. Zelevinsky. On tropical dualities in cluster algebras. In *Algebraic groups and quantum groups*, Contemp. Math., pages 217–226. Amer. Math. Soc., Providence, RI, 2012.
- [Pla18] Pierre-Guy Plamondon. Cluster characters. In *Homological methods, representation theory, and cluster algebras*, CRM Short Courses, pages 101–125. Springer, Cham, 2018.
- [RS18] N. Reading and S. Stella. Initial-seed recursions and dualities for d -vectors. *Pacific J. Math.*, 293(1):179–206, 2018.
- [SB19] M. Sherman-Bennett. Combinatorics of \mathcal{X} -variables in finite type cluster algebras. *J. Combin. Theory Ser. A*, 165:273–298, 2019.
- [SW20] L. Shen and D. Weng. Cyclic sieving and cluster duality of Grassmannian. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 16:Paper No. 067, 41 pp., 2020.

UNIVERSITÉ PARIS CITÉ, UFR DE MATHÉMATIQUES, CNRS, INSTITUT DE MATHÉMATIQUES DE
JUSSIEU-PARIS RIVE GAUCHE, IMJ-PRG, BÂTIMENT SOPHIE GERMAIN, 75205 PARIS CEDEX 13,
FRANCE

PRESENT ADDRESS: EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, THE
HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM 91904, ISRAEL

Email address: peigencao@126.com

UNIVERSITÉ PARIS CITÉ, UFR DE MATHÉMATIQUES, CNRS, INSTITUT DE MATHÉMATIQUES DE
JUSSIEU-PARIS RIVE GAUCHE, IMJ-PRG, BÂTIMENT SOPHIE GERMAIN, 75205 PARIS CEDEX 13,
FRANCE

Email address: bernhard.keller@imj-prg.fr

URL: <https://webusers.imj-prg.fr/~bernhard.keller/>

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, SHANGHAI 200240,
CHINA

Email address: qin.fan.math@gmail.com