

Updated: 09/11/2017

**Linear recurrence sequences,  
exponential polynomials  
and Diophantine approximation**

*Michel Waldschmidt*

Institut de Mathématiques de Jussieu — Paris VI

<http://webusers.imj-prg.fr/~michel.waldschmidt/>

# Abstract

In the first part :

*Linear recurrence sequences : an introduction*

we gave a number of examples and we stated some properties of linear recurrence sequences.

Here we give more information on this topic and we include new results, arising from a joint work with [Claude Levesque](#), involving families of Diophantine equations, with explicit examples related to some units of [L. Bernstein](#) and [H. Hasse](#).

# Linear recurrence sequences : definitions

A *linear recurrence sequence* is a sequence of numbers  $\mathbf{u} = (u_0, u_1, u_2, \dots)$  for which there exist a positive integer  $d$  together with numbers  $a_1, \dots, a_d$  with  $a_d \neq 0$  such that, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Here, a *number* means an element of a field  $\mathbb{K}$  of zero characteristic.

Given  $\underline{a} = (a_1, \dots, a_d) \in \mathbb{K}^d$ , the set  $E_{\underline{a}}$  of linear recurrence sequences  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfying  $(\star)$  is a  $\mathbb{K}$ -vector subspace of dimension  $d$  of the space  $\mathbb{K}^{\mathbb{N}}$  of all sequences .

The characteristic (or companion) polynomial of the linear recurrence is

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

# Linear recurrence sequences : definitions

A *linear recurrence sequence* is a sequence of numbers  $\mathbf{u} = (u_0, u_1, u_2, \dots)$  for which there exist a positive integer  $d$  together with numbers  $a_1, \dots, a_d$  with  $a_d \neq 0$  such that, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Here, a *number* means an element of a field  $\mathbb{K}$  of zero characteristic.

Given  $\underline{a} = (a_1, \dots, a_d) \in \mathbb{K}^d$ , the set  $E_{\underline{a}}$  of linear recurrence sequences  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfying  $(\star)$  is a  $\mathbb{K}$ -vector subspace of dimension  $d$  of the space  $\mathbb{K}^{\mathbb{N}}$  of all sequences .

The characteristic (or companion) polynomial of the linear recurrence is

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

# Linear recurrence sequences : definitions

A *linear recurrence sequence* is a sequence of numbers  $\mathbf{u} = (u_0, u_1, u_2, \dots)$  for which there exist a positive integer  $d$  together with numbers  $a_1, \dots, a_d$  with  $a_d \neq 0$  such that, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Here, a *number* means an element of a field  $\mathbb{K}$  of zero characteristic.

Given  $\underline{a} = (a_1, \dots, a_d) \in \mathbb{K}^d$ , the set  $E_{\underline{a}}$  of linear recurrence sequences  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfying  $(\star)$  is a  $\mathbb{K}$ -vector subspace of dimension  $d$  of the space  $\mathbb{K}^{\mathbb{N}}$  of all sequences .

The characteristic (or companion) polynomial of the linear recurrence is

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

# Linear recurrence sequences : definitions

A *linear recurrence sequence* is a sequence of numbers  $\mathbf{u} = (u_0, u_1, u_2, \dots)$  for which there exist a positive integer  $d$  together with numbers  $a_1, \dots, a_d$  with  $a_d \neq 0$  such that, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Here, a *number* means an element of a field  $\mathbb{K}$  of zero characteristic.

Given  $\underline{a} = (a_1, \dots, a_d) \in \mathbb{K}^d$ , the set  $E_{\underline{a}}$  of linear recurrence sequences  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfying  $(\star)$  is a  $\mathbb{K}$ -vector subspace of dimension  $d$  of the space  $\mathbb{K}^{\mathbb{N}}$  of all sequences .

The characteristic (or companion) polynomial of the linear recurrence is

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

# Linear recurrence sequences : examples

- Constant sequence :  $u_n = u_0$ .

Linear recurrence sequence of order 1 :  $u_{n+1} = u_n$ .

Characteristic polynomial :  $f(X) = X - 1$ .

Generating series :

$$\sum_{n \geq 0} X^n = \frac{1}{1 - X}.$$

- Geometric progression :  $u_n = u_0 \gamma^n$ .

Linear recurrence sequence of order 1 :  $u_n = \gamma u_{n-1}$ .

Characteristic polynomial  $f(X) = X - \gamma$ .

Generating series :

$$\sum_{n \geq 0} u_0 \gamma^n X^n = \frac{u_0}{1 - \gamma X}.$$

# Linear recurrence sequences : examples

- Constant sequence :  $u_n = u_0$ .

Linear recurrence sequence of order 1 :  $u_{n+1} = u_n$ .

Characteristic polynomial :  $f(X) = X - 1$ .

Generating series :

$$\sum_{n \geq 0} X^n = \frac{1}{1 - X}.$$

- Geometric progression :  $u_n = u_0 \gamma^n$ .

Linear recurrence sequence of order 1 :  $u_n = \gamma u_{n-1}$ .

Characteristic polynomial  $f(X) = X - \gamma$ .

Generating series :

$$\sum_{n \geq 0} u_0 \gamma^n X^n = \frac{u_0}{1 - \gamma X}.$$



# Linear recurrence sequences : examples

- $u_n = n$ . This is a linear recurrence sequence of order 2 :

$$n + 2 = 2(n + 1) - n.$$

Characteristic polynomial

$$f(X) = X^2 - 2X + 1 = (X - 1)^2.$$

Generating series

$$\sum_{n \geq 0} nX^n = \frac{1}{1 - 2X + X^2}.$$

Power of matrices :

$$\begin{pmatrix} 0 & 1 \\ -1 & 2 \end{pmatrix}^n = \begin{pmatrix} -n + 1 & n \\ -n & n + 1 \end{pmatrix}.$$

# Linear recurrence sequences : examples

- $u_n = f(n)$ , where  $f$  is polynomial of degree  $d$ . This is a linear recurrence sequence of order  $d + 1$ .

**Proof.** The sequences

$$(f(n))_{n \geq 0}, \quad (f(n+1))_{n \geq 0}, \quad \dots, \quad (f(n+k))_{n \geq 0}$$

are  $\mathbb{K}$ -linearly independent in  $\mathbb{K}^{\mathbb{N}}$  for  $k = d - 1$  and linearly dependent for  $k = d$ .

A basis of the space of polynomials of degree  $d$  is given by the  $d + 1$  polynomials

$$f(X), f(X+1), \dots, f(X+d).$$

Question : *which is the characteristic polynomial ?*

# Linear recurrence sequences : examples

- $u_n = f(n)$ , where  $f$  is polynomial of degree  $d$ . This is a linear recurrence sequence of order  $d + 1$ .

**Proof.** The sequences

$$(f(n))_{n \geq 0}, \quad (f(n+1))_{n \geq 0}, \quad \dots, \quad (f(n+k))_{n \geq 0}$$

are  $\mathbb{K}$ -linearly independent in  $\mathbb{K}^{\mathbb{N}}$  for  $k = d - 1$  and linearly dependent for  $k = d$ .

A basis of the space of polynomials of degree  $d$  is given by the  $d + 1$  polynomials

$$f(X), f(X+1), \dots, f(X+d).$$

Question : *which is the characteristic polynomial ?*

# Linear recurrence sequences : examples

- $u_n = f(n)$ , where  $f$  is polynomial of degree  $d$ . This is a linear recurrence sequence of order  $d + 1$ .

**Proof.** The sequences

$$(f(n))_{n \geq 0}, \quad (f(n+1))_{n \geq 0}, \quad \dots, \quad (f(n+k))_{n \geq 0}$$

are  $\mathbb{K}$ -linearly independent in  $\mathbb{K}^{\mathbb{N}}$  for  $k = d - 1$  and linearly dependent for  $k = d$ .

A basis of the space of polynomials of degree  $d$  is given by the  $d + 1$  polynomials

$$f(X), f(X+1), \dots, f(X+d).$$

Question : *which is the characteristic polynomial ?*

# Linear sequences which are ultimately recurrent

The sequence

$$(1, 0, 0, \dots)$$

is not a linear recurrence sequence.

The condition

$$u_{n+1} = u_n$$

is satisfied only for  $n \geq 1$ .

The relation

$$u_{n+2} = u_{n+1} + 0u_n$$

with  $d = 2$ ,  $a_d = 0$  does not fulfill the requirement  $a_d \neq 0$ .

# Linear sequences which are ultimately recurrent

The sequence

$$(1, 0, 0, \dots)$$

is not a linear recurrence sequence.

The condition

$$u_{n+1} = u_n$$

is satisfied only for  $n \geq 1$ .

The relation

$$u_{n+2} = u_{n+1} + 0u_n$$

with  $d = 2$ ,  $a_d = 0$  does not fulfill the requirement  $a_d \neq 0$ .

# Linear sequences which are ultimately recurrent

The sequence

$$(1, 0, 0, \dots)$$

is not a linear recurrence sequence.

The condition

$$u_{n+1} = u_n$$

is satisfied only for  $n \geq 1$ .

The relation

$$u_{n+2} = u_{n+1} + 0u_n$$

with  $d = 2$ ,  $a_d = 0$  does not fulfill the requirement  $a_d \neq 0$ .

# Order of a linear recurrence sequence

If  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfies the linear recurrence, the characteristic polynomial of which is  $f$ , then, for any monic polynomial  $g \in \mathbb{K}[X]$  with  $g(0) \neq 0$ , this sequence  $\mathbf{u}$  also satisfies the linear recurrence, the characteristic polynomial of which is  $fg$ .

Example : for  $g(X) = X - \gamma$  with  $\gamma \neq 0$ , from

$$(\star) \quad u_{n+d} - a_1 u_{n+d-1} - \cdots - a_d u_n = 0$$

we deduce

$$\begin{aligned} &u_{n+d+1} - a_1 u_{n+d} - \cdots - a_d u_{n+1} \\ &\quad - \gamma(u_{n+d} - a_1 u_{n+d-1} - \cdots - a_d u_n) = 0. \end{aligned}$$

The *order* of a linear recurrence sequence is the smallest  $d$  such that  $(\star)$  holds for all  $n \geq 0$ .



# Order of a linear recurrence sequence

If  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfies the linear recurrence, the characteristic polynomial of which is  $f$ , then, for any monic polynomial  $g \in \mathbb{K}[X]$  with  $g(0) \neq 0$ , this sequence  $\mathbf{u}$  also satisfies the linear recurrence, the characteristic polynomial of which is  $fg$ .

Example : for  $g(X) = X - \gamma$  with  $\gamma \neq 0$ , from

$$(\star) \quad u_{n+d} - a_1 u_{n+d-1} - \cdots - a_d u_n = 0$$

we deduce

$$\begin{aligned} u_{n+d+1} - a_1 u_{n+d} - \cdots - a_d u_{n+1} \\ - \gamma(u_{n+d} - a_1 u_{n+d-1} - \cdots - a_d u_n) = 0. \end{aligned}$$

The *order* of a linear recurrence sequence is the smallest  $d$  such that  $(\star)$  holds for all  $n \geq 0$ .

# Generating series of a linear recurrence sequence

Let  $\mathbf{u} = (u_n)_{n \geq 0}$  be a linear recurrence sequence

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0$$

with characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \cdots - a_d.$$

Denote by  $f^-$  the reciprocal polynomial of  $f$  :

$$f^-(X) = X^d f(X^{-1}) = 1 - a_1 X - \cdots - a_d X^d.$$

Then

$$\sum_{n=0}^{\infty} u_n X^n = \frac{r(X)}{f^-(X)},$$

where  $r$  is a polynomial of degree less than  $d$  determined by the initial values of  $\mathbf{u}$ .

# Generating series of a linear recurrence sequence

Assume

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0.$$

Then

$$\sum_{n=0}^{\infty} u_n X^n = \frac{r(X)}{f^-(X)}.$$

**Proof.** Comparing the coefficients of  $X^n$  for  $n \geq d$  shows that

$$f^-(X) \sum_{n=0}^{\infty} u_n X^n$$

is a polynomial of degree less than  $d$

# Generating series of a linear recurrence sequence

Assume

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0.$$

Then

$$\sum_{n=0}^{\infty} u_n X^n = \frac{r(X)}{f^-(X)}.$$

**Proof.** Comparing the coefficients of  $X^n$  for  $n \geq d$  shows that

$$f^-(X) \sum_{n=0}^{\infty} u_n X^n$$

is a polynomial of degree less than  $d$

# Taylor coefficients of rational functions

Conversely, the coefficients the Taylor expansion of any rational fraction  $a(X)/b(X)$  with  $\deg a < \deg b$  and  $b(0) \neq 0$  satisfies the recurrence relation with characteristic polynomial  $f \in K[X]$  given by  $f(X) = b^-(X)$ .

Therefore a sequence  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfies the recurrence relation  $(\star)$  with characteristic polynomial  $f \in K[X]$  if and only if

$$\sum_{n=0}^{\infty} u_n X^n = \frac{r(X)}{f^-(X)},$$

where  $r$  is a polynomial of degree less than  $d$  determined by the initial values of  $\mathbf{u}$ .

# Taylor coefficients of rational functions

Conversely, the coefficients the Taylor expansion of any rational fraction  $a(X)/b(X)$  with  $\deg a < \deg b$  and  $b(0) \neq 0$  satisfies the recurrence relation with characteristic polynomial  $f \in K[X]$  given by  $f(X) = b^-(X)$ .

Therefore a sequence  $\mathbf{u} = (u_n)_{n \geq 0}$  satisfies the recurrence relation  $(\star)$  with characteristic polynomial  $f \in K[X]$  if and only if

$$\sum_{n=0}^{\infty} u_n X^n = \frac{r(X)}{f^-(X)},$$

where  $r$  is a polynomial of degree less than  $d$  determined by the initial values of  $\mathbf{u}$ .

# Linear differential equations

Given a sequence  $(u_n)_{n \geq 0}$  of numbers, its exponential generating power series is

$$f(z) = \sum_{n \geq 0} u_n \frac{z^n}{n!}.$$

For  $k \geq 0$ , the  $k$ -th derivative  $f^{(k)}$  of  $f$  satisfies

$$f^{(k)}(z) = \sum_{n \geq 0} u_{n+k} \frac{z^n}{n!}.$$

Hence the sequence satisfies the linear recurrence relation

$$(*) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0$$

if and only if  $f$  satisfies the homogeneous linear differential equation

$$y^{(d)} = a_1 y^{(d-1)} + \cdots + a_d y.$$

# Linear differential equations

Given a sequence  $(u_n)_{n \geq 0}$  of numbers, its exponential generating power series is

$$f(z) = \sum_{n \geq 0} u_n \frac{z^n}{n!}.$$

For  $k \geq 0$ , the  $k$ -th derivative  $f^{(k)}$  of  $f$  satisfies

$$f^{(k)}(z) = \sum_{n \geq 0} u_{n+k} \frac{z^n}{n!}.$$

Hence the sequence satisfies the linear recurrence relation

$$(*) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0$$

if and only if  $f$  satisfies the homogeneous linear differential equation

$$y^{(d)} = a_1 y^{(d-1)} + \cdots + a_d y.$$



# Linear differential equations

Given a sequence  $(u_n)_{n \geq 0}$  of numbers, its exponential generating power series is

$$f(z) = \sum_{n \geq 0} u_n \frac{z^n}{n!}.$$

For  $k \geq 0$ , the  $k$ -th derivative  $f^{(k)}$  of  $f$  satisfies

$$f^{(k)}(z) = \sum_{n \geq 0} u_{n+k} \frac{z^n}{n!}.$$

Hence the sequence satisfies the linear recurrence relation

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0$$

if and only if  $f$  satisfies the homogeneous linear differential equation

$$y^{(d)} = a_1 y^{(d-1)} + \cdots + a_d y.$$

# Matrix notation for a linear recurrence sequence

The linear recurrence sequence

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n \quad \text{for } n \geq 0$$

can be written

$$\begin{pmatrix} u_{n+1} \\ u_{n+2} \\ \vdots \\ u_{n+d} \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ a_d & a_{d-1} & a_{d-2} & \cdots & a_1 \end{pmatrix} \begin{pmatrix} u_n \\ u_{n+1} \\ \vdots \\ u_{n+d-1} \end{pmatrix}.$$

# Matrix notation for a linear recurrence sequence

$$U_{n+1} = AU_n$$

with

$$U_n = \begin{pmatrix} u_n \\ u_{n+1} \\ \vdots \\ u_{n+d-1} \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ a_d & a_{d-1} & a_{d-2} & \cdots & a_1 \end{pmatrix}.$$

The determinant of  $I_d X - A$  (the characteristic polynomial of  $A$ ) is nothing but

$$f(X) = X^d - a_1 X^{d-1} - \cdots - a_d,$$

the characteristic polynomial of the linear recurrence sequence.  
By induction

$$U_n = A^n U_0.$$

# Matrix notation for a linear recurrence sequence

$$U_{n+1} = AU_n$$

with

$$U_n = \begin{pmatrix} u_n \\ u_{n+1} \\ \vdots \\ u_{n+d-1} \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ a_d & a_{d-1} & a_{d-2} & \cdots & a_1 \end{pmatrix}.$$

The determinant of  $I_d X - A$  (the characteristic polynomial of  $A$ ) is nothing but

$$f(X) = X^d - a_1 X^{d-1} - \cdots - a_d,$$

the characteristic polynomial of the linear recurrence sequence.

By induction

$$U_n = A^n U_0.$$

# Matrix notation for a linear recurrence sequence

$$U_{n+1} = AU_n$$

with

$$U_n = \begin{pmatrix} u_n \\ u_{n+1} \\ \vdots \\ u_{n+d-1} \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ a_d & a_{d-1} & a_{d-2} & \cdots & a_1 \end{pmatrix}.$$

The determinant of  $I_d X - A$  (the characteristic polynomial of  $A$ ) is nothing but

$$f(X) = X^d - a_1 X^{d-1} - \cdots - a_d,$$

the characteristic polynomial of the linear recurrence sequence.  
By induction

$$U_n = A^n U_0.$$

# Powers of matrices

Let  $A = (a_{ij})_{1 \leq i, j \leq d} \in \text{GL}_{d \times d}(\mathbb{K})$  be a  $d \times d$  matrix with coefficients in  $\mathbb{K}$  and nonzero determinant. For  $n \geq 0$ , define

$$A^n = (a_{ij}^{(n)})_{1 \leq i, j \leq d}.$$

Then each of the  $d^2$  sequences  $(a_{ij}^{(n)})_{n \geq 0}$ ,  $(1 \leq i, j \leq d)$  is a linear recurrence sequence. The roots of the characteristic polynomial of these linear recurrences are the eigenvalues of  $A$ .

In particular the sequence  $(\text{Tr}(A^n))_{n \geq 0}$  satisfies the linear recurrence, the characteristic polynomial of which is the characteristic polynomial of the matrix  $A$ .

# Powers of matrices

Let  $A = (a_{ij})_{1 \leq i, j \leq d} \in \text{GL}_{d \times d}(\mathbb{K})$  be a  $d \times d$  matrix with coefficients in  $\mathbb{K}$  and nonzero determinant. For  $n \geq 0$ , define

$$A^n = (a_{ij}^{(n)})_{1 \leq i, j \leq d}.$$

Then each of the  $d^2$  sequences  $(a_{ij}^{(n)})_{n \geq 0}$ ,  $(1 \leq i, j \leq d)$  is a linear recurrence sequence. The roots of the characteristic polynomial of these linear recurrences are the eigenvalues of  $A$ .

In particular the sequence  $(\text{Tr}(A^n))_{n \geq 0}$  satisfies the linear recurrence, the characteristic polynomial of which is the characteristic polynomial of the matrix  $A$ .

# Powers of matrices

Let  $A = (a_{ij})_{1 \leq i, j \leq d} \in \text{GL}_{d \times d}(\mathbb{K})$  be a  $d \times d$  matrix with coefficients in  $\mathbb{K}$  and nonzero determinant. For  $n \geq 0$ , define

$$A^n = (a_{ij}^{(n)})_{1 \leq i, j \leq d}.$$

Then each of the  $d^2$  sequences  $(a_{ij}^{(n)})_{n \geq 0}$ ,  $(1 \leq i, j \leq d)$  is a linear recurrence sequence. The roots of the characteristic polynomial of these linear recurrences are the eigenvalues of  $A$ .

In particular the sequence  $(\text{Tr}(A^n))_{n \geq 0}$  satisfies the linear recurrence, the characteristic polynomial of which is the characteristic polynomial of the matrix  $A$ .



## Conversely :

Given a linear recurrence sequence  $\mathbf{u} \in \mathbb{K}^{\mathbb{N}}$ , there exist an integer  $d \geq 1$  and a matrix  $A \in \text{GL}_d(\mathbb{K})$  such that, for each  $n \geq 0$ ,

$$u_n = a_{11}^{(n)}.$$

The characteristic polynomial of  $A$  is the characteristic polynomial of the linear recurrence sequence.

EVEREST G., VAN DER POORTEN A., SHPARLINSKI I., WARD T. – *Recurrence sequences*, Mathematical Surveys and Monographs (AMS, 2003), volume 104.

## Conversely :

Given a linear recurrence sequence  $\mathbf{u} \in \mathbb{K}^{\mathbb{N}}$ , there exist an integer  $d \geq 1$  and a matrix  $A \in \text{GL}_d(\mathbb{K})$  such that, for each  $n \geq 0$ ,

$$u_n = a_{11}^{(n)}.$$

The characteristic polynomial of  $A$  is the characteristic polynomial of the linear recurrence sequence.

EVEREST G., VAN DER POORTEN A., SHPARLINSKI I., WARD T. – *Recurrence sequences*, Mathematical Surveys and Monographs (AMS, 2003), volume 104.

# Linear recurrence sequences : simple roots

A basis of  $E_{\underline{a}}$  over  $\mathbb{K}$  is obtained by attributing to the initial values  $u_0, \dots, u_{d-1}$  the values given by the canonical basis of  $\mathbb{K}^d$ .

Given  $\gamma$  in  $\mathbb{K}^\times$ , a necessary and sufficient condition for a sequence  $(\gamma^n)_{n \geq 0}$  to satisfy  $(\star)$  is that  $\gamma$  is a root of the characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

If this polynomial has  $d$  distinct roots  $\gamma_1, \dots, \gamma_d$  in  $\mathbb{K}$ ,

$$f(X) = (X - \gamma_1) \cdots (X - \gamma_d), \quad \gamma_i \neq \gamma_j,$$

then a basis of  $E_{\underline{a}}$  over  $\mathbb{K}$  is given by the  $d$  sequences  $(\gamma_i^n)_{n \geq 0}$ ,  $i = 1, \dots, d$ .

## Linear recurrence sequences : simple roots

A basis of  $E_{\underline{a}}$  over  $\mathbb{K}$  is obtained by attributing to the initial values  $u_0, \dots, u_{d-1}$  the values given by the canonical basis of  $\mathbb{K}^d$ .

Given  $\gamma$  in  $\mathbb{K}^\times$ , a necessary and sufficient condition for a sequence  $(\gamma^n)_{n \geq 0}$  to satisfy  $(\star)$  is that  $\gamma$  is a root of the characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

If this polynomial has  $d$  distinct roots  $\gamma_1, \dots, \gamma_d$  in  $\mathbb{K}$ ,

$$f(X) = (X - \gamma_1) \cdots (X - \gamma_d), \quad \gamma_i \neq \gamma_j,$$

then a basis of  $E_{\underline{a}}$  over  $\mathbb{K}$  is given by the  $d$  sequences  $(\gamma_i^n)_{n \geq 0}$ ,  $i = 1, \dots, d$ .

# Linear recurrence sequences : simple roots

A basis of  $E_{\underline{a}}$  over  $\mathbb{K}$  is obtained by attributing to the initial values  $u_0, \dots, u_{d-1}$  the values given by the canonical basis of  $\mathbb{K}^d$ .

Given  $\gamma$  in  $\mathbb{K}^\times$ , a necessary and sufficient condition for a sequence  $(\gamma^n)_{n \geq 0}$  to satisfy  $(\star)$  is that  $\gamma$  is a root of the characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d.$$

If this polynomial has  $d$  distinct roots  $\gamma_1, \dots, \gamma_d$  in  $\mathbb{K}$ ,

$$f(X) = (X - \gamma_1) \cdots (X - \gamma_d), \quad \gamma_i \neq \gamma_j,$$

then a basis of  $E_{\underline{a}}$  over  $\mathbb{K}$  is given by the  $d$  sequences  $(\gamma_i^n)_{n \geq 0}$ ,  $i = 1, \dots, d$ .

# Linear recurrence sequences : double roots

The characteristic polynomial of the linear recurrence  $u_n = 2\gamma u_{n-1} - \gamma^2 u_{n-2}$  is  $X^2 - 2\gamma X + \gamma^2 = (X - \gamma)^2$  with a double root  $\gamma$ .

The sequence  $(n\gamma^n)_{n \geq 0}$  satisfies

$$n\gamma^n = 2\gamma(n-1)n\gamma^{n-1} - \gamma^2(n-2)\gamma^{n-2}.$$

A basis of  $E_{\underline{a}}$  for  $a_1 = 2\gamma$ ,  $a_2 = -\gamma^2$  is given by the two sequences  $(\gamma^n)_{n \geq 0}$ ,  $(n\gamma^n)_{n \geq 0}$ .

Given  $\gamma \in \mathbb{K}^\times$ , a necessary and sufficient condition for the sequence  $n\gamma^n$  to satisfy the linear recurrence relation  $(\star)$  is that  $\gamma$  is a root of multiplicity  $\geq 2$  of  $f(X)$ .

# Linear recurrence sequences : double roots

The characteristic polynomial of the linear recurrence  $u_n = 2\gamma u_{n-1} - \gamma^2 u_{n-2}$  is  $X^2 - 2\gamma X + \gamma^2 = (X - \gamma)^2$  with a double root  $\gamma$ .

The sequence  $(n\gamma^n)_{n \geq 0}$  satisfies

$$n\gamma^n = 2\gamma(n-1)n\gamma^{n-1} - \gamma^2(n-2)\gamma^{n-2}.$$

A basis of  $E_a$  for  $a_1 = 2\gamma$ ,  $a_2 = -\gamma^2$  is given by the two sequences  $(\gamma^n)_{n \geq 0}$ ,  $(n\gamma^n)_{n \geq 0}$ .

Given  $\gamma \in \mathbb{K}^\times$ , a necessary and sufficient condition for the sequence  $n\gamma^n$  to satisfy the linear recurrence relation  $(\star)$  is that  $\gamma$  is a root of multiplicity  $\geq 2$  of  $f(X)$ .

# Linear recurrence sequences : double roots

The characteristic polynomial of the linear recurrence  $u_n = 2\gamma u_{n-1} - \gamma^2 u_{n-2}$  is  $X^2 - 2\gamma X + \gamma^2 = (X - \gamma)^2$  with a double root  $\gamma$ .

The sequence  $(n\gamma^n)_{n \geq 0}$  satisfies

$$n\gamma^n = 2\gamma(n-1)n\gamma^{n-1} - \gamma^2(n-2)\gamma^{n-2}.$$

A basis of  $E_{\underline{a}}$  for  $a_1 = 2\gamma$ ,  $a_2 = -\gamma^2$  is given by the two sequences  $(\gamma^n)_{n \geq 0}$ ,  $(n\gamma^n)_{n \geq 0}$ .

Given  $\gamma \in \mathbb{K}^\times$ , a necessary and sufficient condition for the sequence  $n\gamma^n$  to satisfy the linear recurrence relation  $(\star)$  is that  $\gamma$  is a root of multiplicity  $\geq 2$  of  $f(X)$ .



# Linear recurrence sequences : double roots

The characteristic polynomial of the linear recurrence  $u_n = 2\gamma u_{n-1} - \gamma^2 u_{n-2}$  is  $X^2 - 2\gamma X + \gamma^2 = (X - \gamma)^2$  with a double root  $\gamma$ .

The sequence  $(n\gamma^n)_{n \geq 0}$  satisfies

$$n\gamma^n = 2\gamma(n-1)n\gamma^{n-1} - \gamma^2(n-2)\gamma^{n-2}.$$

A basis of  $E_{\underline{a}}$  for  $a_1 = 2\gamma$ ,  $a_2 = -\gamma^2$  is given by the two sequences  $(\gamma^n)_{n \geq 0}$ ,  $(n\gamma^n)_{n \geq 0}$ .

Given  $\gamma \in \mathbb{K}^\times$ , a necessary and sufficient condition for the sequence  $n\gamma^n$  to satisfy the linear recurrence relation  $(\star)$  is that  $\gamma$  is a root of multiplicity  $\geq 2$  of  $f(X)$ .

# Linear recurrence sequences : multiple roots

In general, when the characteristic polynomial splits as

$$X^d - a_1X^{d-1} - \dots - a_d = \prod_{i=1}^{\ell} (X - \gamma_i)^{t_i},$$

a basis of  $E_a$  is given by the  $d$  sequences

$$(n^k \gamma_i^n)_{n \geq 0}, \quad 0 \leq k \leq t_i - 1, \quad 1 \leq i \leq \ell.$$

# Polynomial combinations of powers

The sum and the product of any two linear recurrence sequences are linear recurrence sequences.

The set  $\cup_a E_a$  of all linear recurrence sequences with coefficients in  $\mathbb{K}$  is a sub- $\mathbb{K}$ -algebra of  $\mathbb{K}^{\mathbb{N}}$ .

Given polynomials  $p_1, \dots, p_\ell$  in  $\mathbb{K}[X]$  and elements  $\gamma_1, \dots, \gamma_\ell$  in  $\mathbb{K}^\times$ , the sequence

$$(p_1(n)\gamma_1^n + \dots + p_\ell(n)\gamma_\ell^n)_{n \geq 0}$$

is a linear recurrence sequence.

Conversely, any linear recurrence sequence is of this form.

# Polynomial combinations of powers

The sum and the product of any two linear recurrence sequences are linear recurrence sequences.

The set  $\cup_a E_a$  of all linear recurrence sequences with coefficients in  $\mathbb{K}$  is a sub- $\mathbb{K}$ -algebra of  $\mathbb{K}^{\mathbb{N}}$ .

Given polynomials  $p_1, \dots, p_\ell$  in  $\mathbb{K}[X]$  and elements  $\gamma_1, \dots, \gamma_\ell$  in  $\mathbb{K}^\times$ , the sequence

$$(p_1(n)\gamma_1^n + \dots + p_\ell(n)\gamma_\ell^n)_{n \geq 0}$$

is a linear recurrence sequence.

Conversely, any linear recurrence sequence is of this form.

# Polynomial combinations of powers

The sum and the product of any two linear recurrence sequences are linear recurrence sequences.

The set  $\cup_a E_a$  of all linear recurrence sequences with coefficients in  $\mathbb{K}$  is a sub- $\mathbb{K}$ -algebra of  $\mathbb{K}^{\mathbb{N}}$ .

Given polynomials  $p_1, \dots, p_\ell$  in  $\mathbb{K}[X]$  and elements  $\gamma_1, \dots, \gamma_\ell$  in  $\mathbb{K}^\times$ , the sequence

$$(p_1(n)\gamma_1^n + \dots + p_\ell(n)\gamma_\ell^n)_{n \geq 0}$$

is a linear recurrence sequence.

Conversely, any linear recurrence sequence is of this form.

# Polynomial combinations of powers

The sum and the product of any two linear recurrence sequences are linear recurrence sequences.

The set  $\cup_a E_a$  of all linear recurrence sequences with coefficients in  $\mathbb{K}$  is a sub- $\mathbb{K}$ -algebra of  $\mathbb{K}^{\mathbb{N}}$ .

Given polynomials  $p_1, \dots, p_\ell$  in  $\mathbb{K}[X]$  and elements  $\gamma_1, \dots, \gamma_\ell$  in  $\mathbb{K}^\times$ , the sequence

$$(p_1(n)\gamma_1^n + \dots + p_\ell(n)\gamma_\ell^n)_{n \geq 0}$$

is a linear recurrence sequence.

Conversely, any linear recurrence sequence is of this form.

# Consequence

- When  $f$  is a polynomial of degree  $< d$ , the characteristic polynomial of the sequence  $u_n = f(n)$  divides  $(X - 1)^d$ .

Proof.

Set

$$A = \begin{pmatrix} 1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix} = I_d + N$$

where  $I_d$  is the  $d \times d$  identity matrix and  $N$  is nilpotent :  
 $N^d = 0$ .

# Consequence

The characteristic polynomial of  $A$  is  $(X - 1)^d$ . Hence for  $1 \leq i, j \leq d$ , the sequence  $u_n$  of the coefficient  $a_{ij}^{(n)}$  of  $A^n$  satisfies the linear recurrence relation

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n,$$

that is

$$u_{n+d} = d u_{n+d-1} - \binom{d}{2} u_{n+d-2} + \cdots + (-1)^{d-2} d u_{n+1} + (-1)^{d-1} u_n.$$

The characteristic polynomial of this recurrence relation is  $(X - 1)^d$ .



# Characteristic polynomial of the recurrence sequence $f(n)$ .

Since, for  $1 \leq i, j \leq d$  and  $n \geq 0$ , we have

$$a_{ij}^{(n)} = \binom{n}{j-i}$$

(where we agree that  $\binom{n}{k} = 0$  for  $k < 0$  and for  $k > n$ , while  $\binom{d}{0} = \binom{d}{d} = 1$ ), we deduce that each of the  $d$  polynomials

$$1, \frac{X(X+1)\cdots(X+k-1)}{k!} \quad k = 1, 2, \dots, d-1$$

namely

$$1, X, \frac{X(X+1)}{2}, \dots, \frac{X(X+1)\cdots(X+d-2)}{(d-1)!},$$

satisfies the recurrence  $(\star)$ . These  $d$  polynomials constitute a basis of the space of polynomials of degree  $\leq d$ .

# Sum of polynomial combinations of powers

If  $\mathbf{u}_1$  and  $\mathbf{u}_2$  are two linear recurrence sequences of characteristic polynomials  $f_1$  and  $f_2$  respectively, then  $\mathbf{u}_1 + \mathbf{u}_2$  satisfies the linear recurrence, the characteristic polynomial of which is

$$\frac{f_1 f_2}{\gcd(f_1, f_2)}.$$

# Product of polynomial combinations of powers

If the characteristic polynomials of the two linear recurrence sequences  $\mathbf{u}_1$  and  $\mathbf{u}_2$  are respectively

$$f_1(T) = \prod_{j=1}^{\ell} (T - \gamma_j)^{t_j} \quad \text{and} \quad f_2(T) = \prod_{k=1}^{\ell'} (T - \gamma'_k)^{t'_k},$$

then  $\mathbf{u}_1 \mathbf{u}_2$  satisfies the linear recurrence, the characteristic polynomial of which is

$$\prod_{j=1}^{\ell} \prod_{k=1}^{\ell'} (T - \gamma_j \gamma'_k)^{t_j + t'_k - 1}.$$

# Linear recurrence sequences and Brahmagupta–Pell–Fermat Equation

Let  $d$  be a positive integer, not a square. The solutions  $(x, y) \in \mathbb{Z} \times \mathbb{Z}$  of the Brahmagupta–Pell–Fermat Equation

$$x^2 - dy^2 = \pm 1$$

form a sequence  $(x_n, y_n)_{n \in \mathbb{Z}}$  defined by

$$x_n + \sqrt{d}y_n = (x_1 + \sqrt{d}y_1)^n.$$

From

$$2x_n = (x_1 + \sqrt{d}y_1)^n + (x_1 - \sqrt{d}y_1)^n$$

we deduce that  $(x_n)_{n \geq 0}$  is a linear recurrence sequence. Same for  $y_n$ , and also for  $n \leq 0$ .

# Linear recurrence sequences and Brahmagupta–Pell–Fermat Equation

Let  $d$  be a positive integer, not a square. The solutions  $(x, y) \in \mathbb{Z} \times \mathbb{Z}$  of the Brahmagupta–Pell–Fermat Equation

$$x^2 - dy^2 = \pm 1$$

form a sequence  $(x_n, y_n)_{n \in \mathbb{Z}}$  defined by

$$x_n + \sqrt{d}y_n = (x_1 + \sqrt{d}y_1)^n.$$

From

$$2x_n = (x_1 + \sqrt{d}y_1)^n + (x_1 - \sqrt{d}y_1)^n$$

we deduce that  $(x_n)_{n \geq 0}$  is a linear recurrence sequence. Same for  $y_n$ , and also for  $n \leq 0$ .

# Doubly infinite linear recurrence sequences

A sequence  $(u_n)_{n \in \mathbb{Z}}$  indexed by  $\mathbb{Z}$  is a linear recurrence sequence if it satisfies

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n.$$

for all  $n \in \mathbb{Z}$ .

Recall  $a_d \neq 0$ .

Such a sequence is determined by  $d$  consecutive values.

# Doubly infinite linear recurrence sequences

A sequence  $(u_n)_{n \in \mathbb{Z}}$  indexed by  $\mathbb{Z}$  is a linear recurrence sequence if it satisfies

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n.$$

for all  $n \in \mathbb{Z}$ .

Recall  $a_d \neq 0$ .

Such a sequence is determined by  $d$  consecutive values.

# Doubly infinite linear recurrence sequences

A sequence  $(u_n)_{n \in \mathbb{Z}}$  indexed by  $\mathbb{Z}$  is a linear recurrence sequence if it satisfies

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n.$$

for all  $n \in \mathbb{Z}$ .

Recall  $a_d \neq 0$ .

Such a sequence is determined by  $d$  consecutive values.



# Discrete version of linear differential equations

A sequence  $\mathbf{u} \in \mathbb{K}^{\mathbb{N}}$  can be viewed as a linear map  $\mathbb{N} \rightarrow \mathbb{K}$ .  
Define the discrete derivative  $\mathcal{D}$  by

$$\begin{aligned} \mathcal{D}\mathbf{u} : \mathbb{N} &\longrightarrow \mathbb{K} \\ n &\longmapsto u_{n+1} - u_n. \end{aligned}$$

A sequence  $\mathbf{u} \in \mathbb{K}^{\mathbb{N}}$  is a linear recurrence sequence if and only if there exists  $Q \in \mathbb{K}[T]$  with  $Q(1) \neq 1$  such that

$$Q(\mathcal{D})\mathbf{u} = 0.$$

Linear recurrence sequences are a discrete version of linear differential equations with constant coefficients.

The condition  $Q(1) \neq 0$  reflects  $a_d \neq 0$  – otherwise one gets *ultimately* recurrent sequences.

# 97th Indian Science Congress, 2010



A.K. Agarwal


Invited by [Ashok Agrawal](#) to  
the 97th Indian Science  
Congress in  
Thiruvananthapuram  
(Trivandrum, Kerala), January  
3-7, 2010.

- Lecture on *Number Theory  
Challenges of 21st Century*

# A. P. J. Abdul Kalam (1931-2015)

Public Lecture during the 97th Indian Science Congress,  
Thiruvananthapuram – 4 January 2010 Thiruvananthapuram

*Basic research is vital for  
enhancing national and  
international competitiveness*



[http://www.abdulkalam.com/kalam/theme/jsp/guest/  
content-display.jsp](http://www.abdulkalam.com/kalam/theme/jsp/guest/content-display.jsp)

# Kerala 2010

Sudhir Ghorpade



Jugal K. Verma




Ambar Vijayatkumar



January 9-10, 2010, Cochin = Kochi (Kerala) Department of Mathematics, Cochin University of Science and Technology CUSAT

# KSOM 2010

January 8, 2010, Calicut = Kozhikode (Kerala) The Kerala School of Mathematics (KSoM)



A. J. Parameswaran, Director of the Kerala School of Mathematical Science (KSOM) in Kozikhode (Calicut)

# KSOM 2010

Work on dynamical systems by A. J. Parameswaran and S.G. Dani



A. J. Parameswaran



S.G. Dani

# A dynamical system

Let  $V$  be a finite dimensional vector space over a field of zero characteristic,  $H$  an hyperplane of  $V$ ,  $f : V \rightarrow V$  an endomorphism (linear map) and  $x$  an element in  $V$ .

**Theorem.** *If there exist infinitely many  $n \geq 1$  such that  $f^n(x) \in H$ , then there is an (infinite) arithmetic progression of  $n$  for which it is so.*

# A dynamical system

Let  $V$  be a finite dimensional vector space over a field of zero characteristic,  $H$  an hyperplane of  $V$ ,  $f : V \rightarrow V$  an endomorphism (linear map) and  $x$  an element in  $V$ .

**Theorem.** *If there exist infinitely many  $n \geq 1$  such that  $f^n(x) \in H$ , then there is an (infinite) arithmetic progression of  $n$  for which it is so.*



# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Linear recurrence sequence :

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n, \quad n \geq 0 \quad (a_d \neq 0).$$

Characteristic polynomial :

$$X^d - a_1 X^{d-1} - \cdots - a_d = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

$$u_n = \sum_{j=1}^{\ell} \sum_{i=0}^{t_j-1} c_{ij} n^i \gamma_j^n.$$

# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Linear recurrence sequence :

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n, \quad n \geq 0 \quad (a_d \neq 0).$$

Characteristic polynomial :

$$X^d - a_1 X^{d-1} - \cdots - a_d = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

$$u_n = \sum_{j=1}^{\ell} \sum_{i=0}^{t_j-1} c_{ij} n^i \gamma_j^n.$$

# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Linear recurrence sequence :

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n, \quad n \geq 0 \quad (a_d \neq 0).$$

Characteristic polynomial :

$$X^d - a_1 X^{d-1} - \cdots - a_d = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

$$u_n = \sum_{j=1}^{\ell} \sum_{i=0}^{t_j-1} c_{ij} n^i \gamma_j^n.$$

# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Linear recurrence sequence :

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n, \quad n \geq 0 \quad (a_d \neq 0).$$

Characteristic polynomial :

$$X^d - a_1 X^{d-1} - \cdots - a_d = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

$$u_n = \sum_{j=1}^{\ell} \sum_{i=0}^{t_j-1} c_{ij} n^i \gamma_j^n.$$

# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Linear recurrence sequence :

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n, \quad n \geq 0 \quad (a_d \neq 0).$$

Characteristic polynomial :

$$X^d - a_1 X^{d-1} - \cdots - a_d = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

$$u_n = \sum_{j=1}^{\ell} \sum_{i=0}^{t_j-1} c_{ij} n^i \gamma_j^n.$$

# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Linear recurrence sequence :

$$u_{n+d} = a_1 u_{n+d-1} + \cdots + a_d u_n, \quad n \geq 0 \quad (a_d \neq 0).$$

Characteristic polynomial :

$$X^d - a_1 X^{d-1} - \cdots - a_d = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

$$u_n = \sum_{j=1}^{\ell} \sum_{i=0}^{t_j-1} c_{ij} n^i \gamma_j^n.$$

# Wolfgang M. Schmidt

Thue – Siegel – Roth – Schmidt,

Schmidt's Subspace Theorem. The generalized  $S$ -unit Theorem

Let  $\mathbb{K}$  be a field of characteristic zero, let  $G$  be a finitely multiplicative subgroup of the multiplicative group  $\mathbb{K}^\times = \mathbb{K} \setminus \{0\}$  and let  $n \geq 2$ . Then the equation

$$u_1 + u_2 + \cdots + u_n = 1,$$

where the values of the unknowns  $u_1, u_2, \dots, u_n$  are in  $G$  for which no nontrivial subsum

$$\sum_{i \in I} u_i \quad \emptyset \neq I \subset \{1, \dots, n\}$$

vanishes, has only finitely many solutions.

# Wolfgang M. Schmidt

Thue – Siegel – Roth – Schmidt,

Schmidt's Subspace Theorem. The generalized  $S$ -unit Theorem

Let  $\mathbb{K}$  be a field of characteristic zero, let  $G$  be a finitely multiplicative subgroup of the multiplicative group  $\mathbb{K}^\times = \mathbb{K} \setminus \{0\}$  and let  $n \geq 2$ . Then the equation

$$u_1 + u_2 + \cdots + u_n = 1,$$

where the values of the unknowns  $u_1, u_2, \dots, u_n$  are in  $G$  for which no nontrivial subsum

$$\sum_{i \in I} u_i \quad \emptyset \neq I \subset \{1, \dots, n\}$$

vanishes, has only finitely many solutions.



# Wolfgang M. Schmidt

Thue – Siegel – Roth – Schmidt,

Schmidt's Subspace Theorem. The generalized  $S$ -unit Theorem

Let  $\mathbb{K}$  be a field of characteristic zero, let  $G$  be a finitely multiplicative subgroup of the multiplicative group  $\mathbb{K}^\times = \mathbb{K} \setminus \{0\}$  and let  $n \geq 2$ . Then the equation

$$u_1 + u_2 + \cdots + u_n = 1,$$

where the values of the unknowns  $u_1, u_2, \dots, u_n$  are in  $G$  for which no nontrivial subsum

$$\sum_{i \in I} u_i \quad \emptyset \neq I \subset \{1, \dots, n\}$$

vanishes, has only finitely many solutions.

# Wolfgang M. Schmidt

Thue – Siegel – Roth – Schmidt,

Schmidt's Subspace Theorem. The generalized  $S$ -unit Theorem

Let  $\mathbb{K}$  be a field of characteristic zero, let  $G$  be a finitely multiplicative subgroup of the multiplicative group  $\mathbb{K}^\times = \mathbb{K} \setminus \{0\}$  and let  $n \geq 2$ . Then the equation

$$u_1 + u_2 + \cdots + u_n = 1,$$

where the values of the unknowns  $u_1, u_2, \dots, u_n$  are in  $G$  for which no nontrivial subsum

$$\sum_{i \in I} u_i \quad \emptyset \neq I \subset \{1, \dots, n\}$$

vanishes, has only finitely many solutions.

# Schmidt's subspace Theorem



Wolfgang M. Schmidt



Pietro Corvaja



Umberto Zannier

# Balu's 60's Birthday, 2011

December 15 - 20, 2011 : HRI : International Meeting on Number Theory 2011 celebrating the 60th Birthday of Professor R. Balasubramanian.

Pietro Corvaja




M. Manickam, director of KSOM.



December 16, 2011 : lecture on *Families of Thue-Mahler equations*.

# Joint work with Claude Levesque

<http://arxiv.org/abs/1505.06653>



*Solving simultaneously Thue Diophantine equations : almost totally imaginary case*  
Proceedings of the International Meeting on Number Theory HRI 2011, in honor of [R. Balasubramanian](#).


Ramanujan Mathematical Society, Lecture Notes Series **23**,  
*Highly composite : papers in number theory*, (2016), 137–156.  
Editors [Kumar Murty](#), [Ravindranathan Thangadurai](#).

<http://www.ramanujanmathsociety.org/publications/rms-lecture-notes-series>

<http://www.ramanujanmathsociety.org/publications/rms-lecture-notes-series>

## Number-20: The Legacy of Srinivasa Ramanujan

posted Nov 22, 2013, 6:32 AM by RMS Administrator [ updated Nov 23, 2013, 10:55 AM ]



**Title:**


Number 20 - The Legacy of Srinivasa Ramanujan

**Volume Editors:**

Bruce C. Berndt, Dipendra Prasad

# KSOM 2013

Workshop *number theory and dynamical systems* in KSOM  
(Director [M. Manickam](#)) in February 2013



Yann Bugeaud



Pietro Corvaja



S.G. Dani

# Reference

M. WALDSCHMIDT. *Diophantine approximation with applications to dynamical systems*. Proceedings of the International Conference on Pure and Applied Mathematics ICPAM–LAE 2013, South Pacific Journal of Pure and Applied Mathematics, vol. 1, No 2 (2014), 1–18.



# Skolem – Mahler – Lech Theorem

**Theorem** (Skolem 1934 – Mahler 1935 – Lech 1953). *Given a linear recurrence sequence, the set of indices  $n \geq 0$  such that  $u_n = 0$  is a finite union of arithmetic progressions.*

Thoralf Albert Skolem  
(1887 – 1963)



Kurt Mahler  
(1903 – 1988)



Christer Lech

An *arithmetic progression* is a set of positive integers of the form  $\{n_0, n_0 + k, n_0 + 2k, \dots\}$ . Here, we allow  $k = 0$ .

# A dynamical system

Let  $V$  be a finite dimensional vector space over a field of zero characteristic,  $W$  a subspace of  $V$ ,  $f : V \rightarrow V$  an endomorphism (linear map) and  $x$  an element in  $V$ .

**Corollary of the Skolem – Mahler – Lech Theorem.** *The set of  $n \geq 0$  such that  $f^n(x) \in W$  is a finite union of arithmetic progressions.*

By induction, it suffices to consider the case where  $W = H$  is an hyperplane of  $V$ .

# A dynamical system

Let  $V$  be a finite dimensional vector space over a field of zero characteristic,  $W$  a subspace of  $V$ ,  $f : V \rightarrow V$  an endomorphism (linear map) and  $x$  an element in  $V$ .

**Corollary of the Skolem – Mahler – Lech Theorem.** *The set of  $n \geq 0$  such that  $f^n(x) \in W$  is a finite union of arithmetic progressions.*

By induction, it suffices to consider the case where  $W = H$  is an hyperplane of  $V$ .

# A dynamical system

Let  $V$  be a finite dimensional vector space over a field of zero characteristic,  $W$  a subspace of  $V$ ,  $f : V \rightarrow V$  an endomorphism (linear map) and  $x$  an element in  $V$ .

**Corollary of the Skolem – Mahler – Lech Theorem.** *The set of  $n \geq 0$  such that  $f^n(x) \in W$  is a finite union of arithmetic progressions.*

By induction, it suffices to consider the case where  $W = H$  is an hyperplane of  $V$ .

# Idea of the proof of the corollary

Choose a basis of  $V$ . The endomorphism  $f$  is given by a square  $d \times d$  matrix  $A$ , where  $d$  is the dimension of  $V$ . Consider the characteristic polynomial of  $A$ , say

$$X^d - a_{d-1}X^{d-1} - \cdots - a_1X - a_0.$$

By the Theorem of Cayley – Hamilton,

$$A^d = a_{d-1}A^{d-1} + \cdots + a_1A + a_0I_d$$

where  $I_d$  is the identity  $d \times d$  matrix.

# Theorem of Cayley – Hamilton

Arthur Cayley  
(1821 – 1895)



Sir William Rowan Hamilton  
(1805 – 1865)



Hence, for  $n \geq 0$ ,

$$A^{n+d} = a_{d-1}A^{n+d-1} + \dots + a_1A^{n+1} + a_0A^n.$$

It follows that each entry  $a_{ij}(n)$ ,  $1 \leq i, j \leq d$ , satisfies a linear recurrence relation, the same for all  $i, j$ .

# Theorem of Cayley – Hamilton

Arthur Cayley  
(1821 – 1895)



Sir William Rowan Hamilton  
(1805 – 1865)




Hence, for  $n \geq 0$ ,

$$A^{n+d} = a_{d-1}A^{n+d-1} + \cdots + a_1A^{n+1} + a_0A^n.$$

It follows that each entry  $a_{ij}(n)$ ,  $1 \leq i, j \leq d$ , satisfies a linear recurrence relation, the same for all  $i, j$ .

# Theorem of Cayley – Hamilton

Arthur Cayley  
(1821 – 1895)



Sir William Rowan Hamilton  
(1805 – 1865)



Hence, for  $n \geq 0$ ,

$$A^{n+d} = a_{d-1}A^{n+d-1} + \cdots + a_1A^{n+1} + a_0A^n.$$

It follows that each entry  $a_{ij}(n)$ ,  $1 \leq i, j \leq d$ , satisfies a linear recurrence relation, the same for all  $i, j$ .



# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the Skolem – Mahler – Lech Theorem applies.

# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the Skolem – Mahler – Lech Theorem applies.

# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the Skolem – Mahler – Lech Theorem applies.

# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the Skolem – Mahler – Lech Theorem applies.

# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the Skolem – Mahler – Lech Theorem applies.

# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the Skolem – Mahler – Lech Theorem applies.

# Hyperplane membership

Let  $b_1x_1 + \cdots + b_dx_d = 0$  be an equation of the hyperplane  $H$  in the selected basis of  $V$ . Let  ${}^t\underline{b}$  denote the  $1 \times d$  matrix  $(b_1, \dots, b_d)$  (transpose of a column matrix  $\underline{b}$ ). Using the notation  $\underline{v}$  for the  $d \times 1$  (column) matrix given by the coordinates of an element  $v$  in  $V$ , the condition  $v \in H$  can be written  ${}^t\underline{b}\underline{v} = 0$ .

Let  $x$  be an element in  $V$  and  $\underline{x}$  the  $d \times 1$  (column) matrix given by its coordinates. The condition  $f^n(x) \in H$  can now be written

$${}^t\underline{b}A^n\underline{x} = 0.$$

The entry  $u_n$  of the  $1 \times 1$  matrix  ${}^t\underline{b}A^n\underline{x}$  satisfies a linear recurrence relation, hence, the **Skolem – Mahler – Lech** Theorem applies.

# Remark on the theorem of Skolem–Mahler–Lech

T.A. Skolem treated the case  $K = \mathbb{Q}$  of in 1934

K. Mahler the case  $\mathbb{K} = \overline{\mathbb{Q}}$ , the algebraic closure of  $\mathbb{Q}$ , in 1935

The general case was settled by C. Lech in 1953.




# Finite characteristic

C. Lech pointed out in 1953 that such a result may not hold if the characteristic of  $\mathbb{K}$  is positive : he gave as an example the sequence  $u_n = (1 + x)^n - x^n - 1$ , a third-order linear recurrence over the field of rational functions in one variable over the field  $\mathbb{F}_p$  with  $p$  elements, where  $u_n = 0$  for  $n \in \{1, p, p^2, p^3, \dots\}$ . A substitute is provided by a result of Harm Derksen (2007), who proved that the zero set in characteristic  $p$  is a  $p$ -automatic sequence. Further results by Boris Adamczewski and Jason Bell.



Harm Derksen



Boris Adamczewski



Jason Bell

# Polynomial-linear recurrence relation

A generalization of the Theorem of Skolem–Mahler–Lech has been achieved by Jason P. Bell, Stanley Burris and Karen Yeats who prove that the same conclusion holds if the sequence  $(u_n)_{n \geq 0}$  satisfies a polynomial-linear recurrence relation

$$u_n = \sum_{i=1}^d P_i(n) u_{n-i}$$

where  $d$  is a positive integer and  $P_1, \dots, P_d$  are polynomials with coefficient in the field  $\mathbb{K}$  of zero characteristic, provided that  $P_d(x)$  is a nonzero constant.

# Algebraic maps, algebraic groups

There are also analogues of the Theorem of **Skolem–Mahler–Lech** for algebraic maps on varieties (**Jason Bell**).

A version of the **Skolem–Mahler–Lech** Theorem for any algebraic group is due to **Umberto Zannier**.



Jason Bell



Umberto Zannier

# Open problem

One main open problem related with Theorem of **Skolem–Mahler–Lech** is that it is not effective : explicit upper bounds for the number of arithmetic progressions, depending only on the order  $d$  of the linear recurrence sequence, are known (**W.M. Schmidt**, **U. Zannier**), but no upper bound for the arithmetic progressions themselves is known. A related open problem raised by **T.A. Skolem** and **C. Pisot** is :

*Given an integer linear recurrence sequence, is the truth of the statement “ $x_n \neq 0$  for all  $n$ ” decidable in finite time?*

**T. TAO**, *Effective Skolem Mahler Lech theorem*. In “Structure and Randomness : pages from year one of a mathematical blog”, American Mathematical Society (2008), 298 pages.

<http://terrytao.wordpress.com/2007/05/25/open-question-effective-skolem-mahler-lech-theorem/>

# Zeros of linear recurrence sequences

Jean Berstel et Maurice Mignotte. – Deux propriétés décidables des suites récurrentes linéaires Bulletin de la S.M.F., tome 104 (1976), p. 175-184.

[http://www.numdam.org/item?id=BSMF\\_1976\\_\\_104\\_\\_175\\_0](http://www.numdam.org/item?id=BSMF_1976__104__175_0)

*Given a linear recurrence sequence with integer coefficients ; are there finitely or infinitely many zeroes ?*

Philippe Robba. – Zéros de suites récurrentes linéaires. Groupe de travail d'analyse ultramétrique (1977-1978) Volume : 5, page 1-5.

L. Cerlienco, M. Mignotte, F. Piras. Suites récurrentes linéaires. Propriétés algébriques et arithmétiques. L'Enseignement Mathématique **33** (1987).

# Zeros of linear recurrence sequences

Jean Berstel et Maurice Mignotte. – Deux propriétés décidables des suites récurrentes linéaires Bulletin de la S.M.F., tome 104 (1976), p. 175-184.

[http://www.numdam.org/item?id=BSMF\\_1976\\_\\_104\\_\\_175\\_0](http://www.numdam.org/item?id=BSMF_1976__104__175_0)

*Given a linear recurrence sequence with integer coefficients ; are there finitely or infinitely many zeroes ?*

Philippe Robba. – Zéros de suites récurrentes linéaires. Groupe de travail d'analyse ultramétrique (1977-1978) Volume : 5, page 1-5.

L. Cerlienco, M. Mignotte, F. Piras. Suites récurrentes linéaires. Propriétés algébriques et arithmétiques. L'Enseignement Mathématique **33** (1987).

# Zeros of linear recurrence sequences

Jean Berstel et Maurice Mignotte. – Deux propriétés décidables des suites récurrentes linéaires Bulletin de la S.M.F., tome 104 (1976), p. 175-184.

[http://www.numdam.org/item?id=BSMF\\_1976\\_\\_104\\_\\_175\\_0](http://www.numdam.org/item?id=BSMF_1976__104__175_0)

*Given a linear recurrence sequence with integer coefficients; are there finitely or infinitely many zeroes?*

Philippe Robba. – Zéros de suites récurrentes linéaires. Groupe de travail d'analyse ultramétrique (1977-1978) Volume : 5, page 1-5.

L. Cerlienco, M. Mignotte, F. Piras. Suites récurrentes linéaires. Propriétés algébriques et arithmétiques. L'Enseignement Mathématique **33** (1987).

# Zeros of linear recurrence sequences

Maurice Mignotte Propriétés arithmétiques des suites récurrentes linéaires. Besançon, 1989  
<http://pmb.univ-fcomte.fr/1989/Mignotte.pdf>

E. Bavencoffe and J-P. Bézivin *Une famille remarquable de suites récurrentes linéaires*. – Monatshefte für Mathematik, (1995) **120** 3, 189–203

Karim Samake. – Suites récurrentes linéaires, problème d'effectivité. Inst. de Recherche Math. Avancée, 1996 - 62 pages



# Zeros of linear recurrence sequences

Maurice Mignotte Propriétés arithmétiques des suites récurrentes linéaires. Besançon, 1989  
<http://pmb.univ-fcomte.fr/1989/Mignotte.pdf>

E. Bavencoffe and J-P. Bézivin *Une famille remarquable de suites récurrentes linéaires*. – Monatshefte für Mathematik, (1995) **120** 3, 189–203

Karim Samake. – Suites récurrentes linéaires, problème d'effectivité. Inst. de Recherche Math. Avancée, 1996 - 62 pages

# Zeros of linear recurrence sequences

Maurice Mignotte Propriétés arithmétiques des suites récurrentes linéaires. Besançon, 1989  
<http://pmb.univ-fcomte.fr/1989/Mignotte.pdf>

E. Bavencoffe and J-P. Bézivin *Une famille remarquable de suites récurrentes linéaires*. – Monatshefte für Mathematik, (1995) **120** 3, 189–203

Karim Samake. – Suites récurrentes linéaires, problème d'effectivité. Inst. de Recherche Math. Avancée, 1996 - 62 pages

# Reference

EVEREST, GRAHAM ; VAN DER POORTEN, ALF ; SHPARLINSKI, IGOR ; WARD, TOM – *Recurrence sequences*, Mathematical Surveys and Monographs (AMS, 2003), volume 104. 1290 references.



Graham Everest



Alf van der Poorten



Igor Shparlinski



Tom Ward

# Berstel's sequence

<http://oeis.org/A007420>

0, 0, 1, 2, 0, -4, 0, 16, 16, -32, -64, 64, 256, 0, -768, ...



Jean Berstel

$$b_0 = b_1 = 0, b_2 = 1,$$
$$b_{n+3} = 2b_{n+2} - 4b_{n+1} + 4b_n$$

for  $n \geq 0$ .

Linear recurrence sequence of  
order 3 with exactly 6 zeros :  
 $n = 0, 1, 4, 6, 13, 52$ .

<http://www-igm.univ-mlv.fr/~berstel/>

# Ternary linear recurrences

Berstel's sequence is a linear recurrence sequence of order 3 with 6 zeroes.



Frits Beukers

Frits Beukers (1991) : up to trivial transformation, any other linear recurrence of order 3 with finitely many zeroes has at most 5 zeros.

# Edgard Bavencoffe and Jean-Paul Bézivin

Let  $n \geq 2$ . The sequence with initial values

$$u_0 = 1, u_1 = \cdots = u_{n-1} = 0$$

satisfying the recurrence relation of order  $n$  with characteristic polynomial

$$\frac{X^{n+1} - (-2)^{n-1}X + (-2)^n}{X + 2}$$

has at least

$$\frac{n(n+1)}{2} - 1$$

zeroes.

# Edgard Bavencoffe and Jean-Paul Bézivin

For  $n = 3$  one obtains Berstel's sequence which happens to have an extra zero.

$$\frac{X^4 + 4X - 8}{X + 2} = X^3 - 2X^2 + 4X - 4.$$



Edgard Bavencoffe




Jean-Paul Bézivin

# Berstel's sequence

0, 0, 1, 2, 0, -4, 0, 16, 16, -32, -64, 64, 256, 0, -768, ...

$b_0 = b_1 = 0$ ,  $b_2 = 1$ ,  $b_{n+3} = 2b_{n+2} - 4b_{n+1} + 4b_n$  for  $n \geq 0$ .




Maurice Mignotte

The equation  $b_m = \pm b_n$  has exactly 21 solutions  $(m, n)$  with  $m \neq n$ .

The equation  $b_n = \pm 2^r 3^s$  has exactly 44 solutions  $(n, r, s)$ .



# Joint work with Claude Levesque



*Linear recurrence sequences  
and twisted binary forms.*

Proceedings of the  
International Conference on  
Pure and Applied  
Mathematics

ICPAM-GOROKA 2014.

South Pacific Journal of Pure  
and Applied Mathematics.

<http://webusers.imj-prg.fr/~michel.waldschmidt//articles/pdf/ProcConfPNG2014.pdf>

# Families of binary forms

Consider a binary form  $F_0(X, Y) \in \mathbb{C}[X, Y]$  which satisfies  $F_0(1, 0) = 1$ . We write it as

$$F_0(X, Y) = X^d + a_1 X^{d-1} Y + \cdots + a_d Y^d = \prod_{i=1}^d (X - \alpha_i Y).$$

Let  $\epsilon_1, \dots, \epsilon_d$  be  $d$  nonzero complex numbers not necessarily distinct. Twisting  $F_0$  by the powers  $\epsilon_1^n, \dots, \epsilon_d^n$  ( $n \in \mathbb{Z}$ ) boils down to considering the family of binary forms

$$F_n(X, Y) = \prod_{i=1}^d (X - \alpha_i \epsilon_i^n Y),$$

which we write as

$$X^d - U_1(n) X^{d-1} Y + \cdots + (-1)^d U_d(n) Y^d.$$

Therefore

$$U_h(0) = (-1)^h a_h \quad (1 \leq h \leq d).$$

# Families of binary forms

Consider a binary form  $F_0(X, Y) \in \mathbb{C}[X, Y]$  which satisfies  $F_0(1, 0) = 1$ . We write it as

$$F_0(X, Y) = X^d + a_1 X^{d-1} Y + \cdots + a_d Y^d = \prod_{i=1}^d (X - \alpha_i Y).$$

Let  $\epsilon_1, \dots, \epsilon_d$  be  $d$  nonzero complex numbers not necessarily distinct. Twisting  $F_0$  by the powers  $\epsilon_1^n, \dots, \epsilon_d^n$  ( $n \in \mathbb{Z}$ ) boils down to considering the family of binary forms

$$F_n(X, Y) = \prod_{i=1}^d (X - \alpha_i \epsilon_i^n Y),$$

which we write as

$$X^d - U_1(n) X^{d-1} Y + \cdots + (-1)^d U_d(n) Y^d.$$

Therefore

$$U_h(0) = (-1)^h a_h \quad (1 \leq h \leq d).$$

# Families of binary forms

Consider a binary form  $F_0(X, Y) \in \mathbb{C}[X, Y]$  which satisfies  $F_0(1, 0) = 1$ . We write it as

$$F_0(X, Y) = X^d + a_1 X^{d-1} Y + \cdots + a_d Y^d = \prod_{i=1}^d (X - \alpha_i Y).$$

Let  $\epsilon_1, \dots, \epsilon_d$  be  $d$  nonzero complex numbers not necessarily distinct. Twisting  $F_0$  by the powers  $\epsilon_1^n, \dots, \epsilon_d^n$  ( $n \in \mathbb{Z}$ ) boils down to considering the family of binary forms

$$F_n(X, Y) = \prod_{i=1}^d (X - \alpha_i \epsilon_i^n Y),$$

which we write as

$$X^d - U_1(n) X^{d-1} Y + \cdots + (-1)^d U_d(n) Y^d.$$

Therefore

$$U_h(0) = (-1)^h a_h \quad (1 \leq h \leq d).$$

# Families of binary forms

Consider a binary form  $F_0(X, Y) \in \mathbb{C}[X, Y]$  which satisfies  $F_0(1, 0) = 1$ . We write it as

$$F_0(X, Y) = X^d + a_1 X^{d-1} Y + \cdots + a_d Y^d = \prod_{i=1}^d (X - \alpha_i Y).$$

Let  $\epsilon_1, \dots, \epsilon_d$  be  $d$  nonzero complex numbers not necessarily distinct. Twisting  $F_0$  by the powers  $\epsilon_1^n, \dots, \epsilon_d^n$  ( $n \in \mathbb{Z}$ ) boils down to considering the family of binary forms

$$F_n(X, Y) = \prod_{i=1}^d (X - \alpha_i \epsilon_i^n Y),$$

which we write as

$$X^d - U_1(n) X^{d-1} Y + \cdots + (-1)^d U_d(n) Y^d.$$

Therefore

$$U_h(0) = (-1)^h a_h \quad (1 \leq h \leq d).$$

# Families of Diophantine equations

With **Claude Levesque**, we considered some families of diophantine equations

$$F_n(x, y) = m$$

obtained in the same way from a given irreducible form  $F(X, Y)$  with coefficients in  $\mathbb{Z}$ , when  $\epsilon_1, \dots, \epsilon_d$  are algebraic units and when the algebraic numbers  $\alpha_1\epsilon_1, \dots, \alpha_d\epsilon_d$  are **Galois** conjugates with  $d \geq 3$ .

**Theorem.** *Let  $\mathbb{K}$  be a number field of degree  $d \geq 3$ ,  $S$  a finite set of places of  $\mathbb{K}$  containing the places at infinity. Denote by  $\mathcal{O}_S$  the ring of  $S$ -integers of  $\mathbb{K}$  and by  $\mathcal{O}_S^\times$  the group of  $S$ -units of  $\mathbb{K}$ . Assume  $\alpha_1, \dots, \alpha_d, \epsilon_1, \dots, \epsilon_d$  belong to  $\mathbb{K}^\times$ . Then there are only finitely many  $(x, y, n)$  in  $\mathcal{O}_S \times \mathcal{O}_S \times \mathbb{Z}$  satisfying*

$$F_n(x, y) \in \mathcal{O}_S^\times, \quad xy \neq 0 \quad \text{and} \quad \text{Card}\{\alpha_1\epsilon_1^n, \dots, \alpha_d\epsilon_d^n\} \geq 3.$$

# Families of Diophantine equations

With **Claude Levesque**, we considered some families of diophantine equations

$$F_n(x, y) = m$$

obtained in the same way from a given irreducible form  $F(X, Y)$  with coefficients in  $\mathbb{Z}$ , when  $\epsilon_1, \dots, \epsilon_d$  are algebraic units and when the algebraic numbers  $\alpha_1\epsilon_1, \dots, \alpha_d\epsilon_d$  are **Galois** conjugates with  $d \geq 3$ .

**Theorem.** *Let  $\mathbb{K}$  be a number field of degree  $d \geq 3$ ,  $S$  a finite set of places of  $\mathbb{K}$  containing the places at infinity. Denote by  $\mathcal{O}_S$  the ring of  $S$ -integers of  $\mathbb{K}$  and by  $\mathcal{O}_S^\times$  the group of  $S$ -units of  $\mathbb{K}$ . Assume  $\alpha_1, \dots, \alpha_d, \epsilon_1, \dots, \epsilon_d$  belong to  $\mathbb{K}^\times$ . Then there are only finitely many  $(x, y, n)$  in  $\mathcal{O}_S \times \mathcal{O}_S \times \mathbb{Z}$  satisfying*

$$F_n(x, y) \in \mathcal{O}_S^\times, \quad xy \neq 0 \quad \text{and} \quad \text{Card}\{\alpha_1\epsilon_1^n, \dots, \alpha_d\epsilon_d^n\} \geq 3.$$

# Families of Diophantine equations

Each of the sequences  $(U_h(n))_{n \in \mathbb{Z}}$  coming from the coefficients of the relation

$$F_n(X, Y) = X^d - U_1(n)X^{d-1}Y + \cdots + (-1)^d U_d(n)Y^d$$

is a linear recurrence sequence.

For example, for  $n \in \mathbb{Z}$ ,

$$U_1(n) = \sum_{i=1}^d \alpha_i \epsilon_i^n, \quad U_d(n) = \prod_{i=1}^d \alpha_i \epsilon_i^n.$$

For  $1 \leq h \leq d$ , the sequence  $(U_h(n))_{n \in \mathbb{Z}}$  is a linear combination of the sequences

$$((\epsilon_{i_1} \cdots \epsilon_{i_h})^n)_{n \in \mathbb{Z}}, \quad (1 \leq i_1 < \cdots < i_h \leq d).$$



# Families of Diophantine equations

Each of the sequences  $(U_h(n))_{n \in \mathbb{Z}}$  coming from the coefficients of the relation

$$F_n(X, Y) = X^d - U_1(n)X^{d-1}Y + \cdots + (-1)^d U_d(n)Y^d$$

is a linear recurrence sequence.

For example, for  $n \in \mathbb{Z}$ ,

$$U_1(n) = \sum_{i=1}^d \alpha_i \epsilon_i^n, \quad U_d(n) = \prod_{i=1}^d \alpha_i \epsilon_i^n.$$

For  $1 \leq h \leq d$ , the sequence  $(U_h(n))_{n \in \mathbb{Z}}$  is a linear combination of the sequences

$$((\epsilon_{i_1} \cdots \epsilon_{i_h})^n)_{n \in \mathbb{Z}}, \quad (1 \leq i_1 < \cdots < i_h \leq d).$$

# Families of Diophantine equations

Each of the sequences  $(U_h(n))_{n \in \mathbb{Z}}$  coming from the coefficients of the relation

$$F_n(X, Y) = X^d - U_1(n)X^{d-1}Y + \cdots + (-1)^d U_d(n)Y^d$$

is a linear recurrence sequence.

For example, for  $n \in \mathbb{Z}$ ,

$$U_1(n) = \sum_{i=1}^d \alpha_i \epsilon_i^n, \quad U_d(n) = \prod_{i=1}^d \alpha_i \epsilon_i^n.$$

For  $1 \leq h \leq d$ , the sequence  $(U_h(n))_{n \in \mathbb{Z}}$  is a linear combination of the sequences

$$((\epsilon_{i_1} \cdots \epsilon_{i_h})^n)_{n \in \mathbb{Z}}, \quad (1 \leq i_1 < \cdots < i_h \leq d).$$

# Some units of Bernstein and Hasse

Let  $t$  and  $s$  be two positive integers,  $D$  an integer  $\geq 1$ , and  $c \in \{-1, +1\}$ . Let  $\omega > 1$  satisfy

$$\omega^{st} = D^{st} + c,$$

where it is assumed that  $\mathbb{Q}(\omega)$  is of degree  $st$ .

Consider

$$\alpha = D - \omega, \quad \epsilon = D^t - \omega^t.$$

L. Bernstein and H. Hasse noticed that  $\alpha$  and  $\epsilon$  are units of degree  $st$  and  $s$  respectively, and showed that these units can be obtained from the Jacobi–Perron algorithm. H.-J. Stender proved that for  $s = t = 2$ ,  $\{\alpha, \epsilon\}$  is a fundamental system of units of the quartic field  $\mathbb{Q}(\omega)$ .

# Some units of Bernstein and Hasse

Let  $t$  and  $s$  be two positive integers,  $D$  an integer  $\geq 1$ , and  $c \in \{-1, +1\}$ . Let  $\omega > 1$  satisfy

$$\omega^{st} = D^{st} + c,$$

where it is assumed that  $\mathbb{Q}(\omega)$  is of degree  $st$ .

Consider

$$\alpha = D - \omega, \quad \epsilon = D^t - \omega^t.$$

L. Bernstein and H. Hasse noticed that  $\alpha$  and  $\epsilon$  are units of degree  $st$  and  $s$  respectively, and showed that these units can be obtained from the Jacobi–Perron algorithm. H.-J. Stender proved that for  $s = t = 2$ ,  $\{\alpha, \epsilon\}$  is a fundamental system of units of the quartic field  $\mathbb{Q}(\omega)$ .

# Helmut Hasse (1898-1979)

$$D > 0, s \geq 1, t \geq 1,$$
$$c \in \{-1, +1\}, \omega > 0,$$

$$\omega^{st} = D^{st} + c,$$

$$\alpha = D - \omega,$$

$$\epsilon = D^t - \omega^t.$$



$$(\alpha - D)^{st} = (-1)^{st}(D^{st} + c).$$

# Diophantine equations associated with some units of Bernstein and Hasse

The irreducible polynomial of  $\alpha$  is  $F_0(X, 1)$ , with

$$F_0(X, Y) = (X - DY)^{st} - (-1)^{st}(D^{st} + c)Y^{st}.$$

For  $n \in \mathbb{Z}$ , the binary form  $F_n(X, Y)$ , obtained by twisting  $F_0(X, Y)$  with the powers  $\epsilon^n$  of  $\epsilon$ , is the homogeneous version of the irreducible polynomial  $F_n(X, 1)$  of  $\alpha\epsilon^n$ . So  $F_n$  depends of the parameters  $n, D, s, t$  and  $c$ .

**Theorem (LW).** *Suppose  $st \geq 3$ . There exists an effectively computable constant  $\kappa$ , depending only on  $D, s$  and  $t$ , with the following property. Let  $m, a, x, y$  be rational integers satisfying  $m \geq 2, xy \neq 0, [\mathbb{Q}(\alpha\epsilon^a) : \mathbb{Q}] = st$  and*

$$|F_n(x, y)| \leq m.$$

*Then*

$$\max\{\log |x|, \log |y|, |n|\} \leq \kappa \log m.$$

# Diophantine equations associated with some units of Bernstein and Hasse

The irreducible polynomial of  $\alpha$  is  $F_0(X, 1)$ , with

$$F_0(X, Y) = (X - DY)^{st} - (-1)^{st}(D^{st} + c)Y^{st}.$$

For  $n \in \mathbb{Z}$ , the binary form  $F_n(X, Y)$ , obtained by twisting  $F_0(X, Y)$  with the powers  $\epsilon^n$  of  $\epsilon$ , is the homogeneous version of the irreducible polynomial  $F_n(X, 1)$  of  $\alpha\epsilon^n$ . So  $F_n$  depends of the parameters  $n$ ,  $D$ ,  $s$ ,  $t$  and  $c$ .

**Theorem (LW).** *Suppose  $st \geq 3$ . There exists an effectively computable constant  $\kappa$ , depending only on  $D$ ,  $s$  and  $t$ , with the following property. Let  $m$ ,  $a$ ,  $x$ ,  $y$  be rational integers satisfying  $m \geq 2$ ,  $xy \neq 0$ ,  $[\mathbb{Q}(\alpha\epsilon^a) : \mathbb{Q}] = st$  and*

$$|F_n(x, y)| \leq m.$$

*Then*

$$\max\{\log |x|, \log |y|, |n|\} \leq \kappa \log m.$$

# Diophantine equations associated with some units of Bernstein and Hasse

The irreducible polynomial of  $\alpha$  is  $F_0(X, 1)$ , with

$$F_0(X, Y) = (X - DY)^{st} - (-1)^{st}(D^{st} + c)Y^{st}.$$

For  $n \in \mathbb{Z}$ , the binary form  $F_n(X, Y)$ , obtained by twisting  $F_0(X, Y)$  with the powers  $\epsilon^n$  of  $\epsilon$ , is the homogeneous version of the irreducible polynomial  $F_n(X, 1)$  of  $\alpha\epsilon^n$ . So  $F_n$  depends of the parameters  $n$ ,  $D$ ,  $s$ ,  $t$  and  $c$ .

**Theorem (LW).** *Suppose  $st \geq 3$ . There exists an effectively computable constant  $\kappa$ , depending only on  $D$ ,  $s$  and  $t$ , with the following property. Let  $m$ ,  $a$ ,  $x$ ,  $y$  be rational integers satisfying  $m \geq 2$ ,  $xy \neq 0$ ,  $[\mathbb{Q}(\alpha\epsilon^a) : \mathbb{Q}] = st$  and*

$$|F_n(x, y)| \leq m.$$

Then

$$\max\{\log |x|, \log |y|, |n|\} \leq \kappa \log m.$$



# Hankel determinants

To test an arbitrary sequence  $\mathbf{u} = (u_n)_{n \geq 0}$  of elements of a field  $\mathbb{K}$  for the property of being a linear recurrence sequence, consider the **Hankel** determinants

$$\Delta_{N,d}(\mathbf{u}) = \det (u_{d+i+j})_{0 \leq i,j \leq N}.$$

The sum



Hermann Hankel  
(1839–1873)

$$f(z) = \sum_{n=0}^{\infty} u_n z^n$$

represents a rational function if and only if for some  $d$ ,  $\Delta_{N,d}(\mathbf{u}) = 0$  for all sufficiently large  $N$


# Hankel determinants

Alan Haynes, Wadim Zudilin. – Hankel determinants of zeta values  
(Submitted on 7 Oct 2015)

Abstract: *We study the asymptotics of **Hankel** determinants constructed using the values  $\zeta(an + b)$  of the **Riemann** zeta function at positive integers in an arithmetic progression. Our principal result is a Diophantine application of the asymptotics.*




Alan Haynes



Wadim Zudilin

# Perfect powers in the Fibonacci sequence

Yann Bugeaud, Maurice Mignotte, Samir Siksek (2004) : The only perfect powers (squares, cubes, etc.) in the Fibonacci sequence are 1, 8 and 144.



Y. Bugeaud



M. Mignotte



S. Siksek

# Powers in recurrence sequences



Mike Bennett

M. A. Bennett, Powers in recurrence sequences : Pell equations, Trans. Amer. Math. Soc. **357** (2005), 1675-1691.

<http://www.math.ubc.ca/~bennett/paper31.pdf>

# Bases of the space of linear recurrence sequences

Given  $a_1, \dots, a_d$  with  $a_d \neq 0$ , consider the vector space of linear recurrence sequences satisfying, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Assuming the characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d$$

of the recurrence splits completely in  $\mathbb{K}$ ,

$$f(X) = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

we have two bases. The first one given by the initial conditions  $(u_0, \dots, u_{d-1})$ , and the second one is given by the sequences

$$(n^i \gamma_j^n)_{n \geq 0}, \quad 0 \leq i \leq t_j - 1, \quad 1 \leq j \leq \ell.$$

# Bases of the space of linear recurrence sequences

Given  $a_1, \dots, a_d$  with  $a_d \neq 0$ , consider the vector space of linear recurrence sequences satisfying, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Assuming the characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d$$

of the recurrence splits completely in  $\mathbb{K}$ ,

$$f(X) = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

we have two bases. The first one given by the initial conditions  $(u_0, \dots, u_{d-1})$ , and the second one is given by the sequences

$$(n^i \gamma_j^n)_{n \geq 0}, \quad 0 \leq i \leq t_j - 1, \quad 1 \leq j \leq \ell.$$

# Bases of the space of linear recurrence sequences

Given  $a_1, \dots, a_d$  with  $a_d \neq 0$ , consider the vector space of linear recurrence sequences satisfying, for  $n \geq 0$ ,

$$(\star) \quad u_{n+d} = a_1 u_{n+d-1} + \dots + a_d u_n.$$

Assuming the characteristic polynomial

$$f(X) = X^d - a_1 X^{d-1} - \dots - a_d$$

of the recurrence splits completely in  $\mathbb{K}$ ,

$$f(X) = \prod_{j=1}^{\ell} (X - \gamma_j)^{t_j}$$

we have two bases. The first one given by the initial conditions  $(u_0, \dots, u_{d-1})$ , and the second one is given by the sequences

$$(n^i \gamma_j^n)_{n \geq 0}, \quad 0 \leq i \leq t_j - 1, \quad 1 \leq j \leq \ell.$$

# Change of basis

The matrix of change of bases is

$$M = \begin{pmatrix} M_1 \\ \vdots \\ M_\ell \end{pmatrix}$$

where

$$M_j = \begin{pmatrix} 1 & \gamma_j & \gamma_j^2 & \cdots & \gamma_j^{t_j-1} & \gamma_j^{t_j} & \cdots & \gamma_j^{d-1} \\ 0 & 1 & \binom{2}{1}\gamma_j & \cdots & \binom{t_j-1}{1}\gamma_j^{t_j-2} & \binom{t_j}{1}\gamma_j^{t_j-1} & \cdots & \binom{d-1}{1}\gamma_j^{d-2} \\ 0 & 0 & 1 & \cdots & \binom{t_j-1}{2}\gamma_j^{t_j-3} & \binom{t_j}{2}\gamma_j^{t_j-2} & \cdots & \binom{d-1}{2}\gamma_j^{d-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & \binom{t_j}{t_j-1}\gamma_j & \cdots & \binom{d-1}{t_j-1}\gamma_j^{d-t_j} \end{pmatrix}$$



# Exponential polynomials

The sequence of derivatives of an exponential polynomial evaluated at one point satisfies a linear recurrence relation.

Let  $p_1(z), \dots, p_\ell(z)$  be nonzero polynomials of  $\mathbb{C}[z]$  of degrees smaller than  $t_1, \dots, t_\ell$  respectively. Let  $\gamma_1, \dots, \gamma_\ell$  be distinct complex numbers. Suppose that the function

$$F(z) = p_1(z)e^{\gamma_1 z} + \dots + p_\ell(z)e^{\gamma_\ell z}$$

is not identically 0. Then its vanishing order at a point  $z_0$  is smaller than or equal to  $t_1 + \dots + t_\ell - 1$ .

In other terms, when the complex numbers  $\gamma_j$  are distinct, the determinant

$$\left| \left( \frac{d}{dz} \right)^a (z^i e^{\gamma_j z}) \Big|_{z=0} \right|_{\substack{0 \leq i \leq t_j - 1, 1 \leq j \leq \ell \\ 0 \leq a \leq d-1}}$$

is different from 0. This is no surprise that we come across the determinant of the matrix  $M$ .

# Exponential polynomials

The sequence of derivatives of an exponential polynomial evaluated at one point satisfies a linear recurrence relation. Let  $p_1(z), \dots, p_\ell(z)$  be nonzero polynomials of  $\mathbb{C}[z]$  of degrees smaller than  $t_1, \dots, t_\ell$  respectively. Let  $\gamma_1, \dots, \gamma_\ell$  be distinct complex numbers. Suppose that the function

$$F(z) = p_1(z)e^{\gamma_1 z} + \dots + p_\ell(z)e^{\gamma_\ell z}$$

is not identically 0. Then its vanishing order at a point  $z_0$  is smaller than or equal to  $t_1 + \dots + t_\ell - 1$ .

In other terms, when the complex numbers  $\gamma_j$  are distinct, the determinant

$$\left| \left( \frac{d}{dz} \right)^a (z^i e^{\gamma_j z}) \Big|_{z=0} \right|_{\substack{0 \leq i \leq t_j - 1, 1 \leq j \leq \ell \\ 0 \leq a \leq d-1}}$$

is different from 0. This is no surprise that we come across the determinant of the matrix  $M$ .

# Exponential polynomials

The sequence of derivatives of an exponential polynomial evaluated at one point satisfies a linear recurrence relation. Let  $p_1(z), \dots, p_\ell(z)$  be nonzero polynomials of  $\mathbb{C}[z]$  of degrees smaller than  $t_1, \dots, t_\ell$  respectively. Let  $\gamma_1, \dots, \gamma_\ell$  be distinct complex numbers. Suppose that the function

$$F(z) = p_1(z)e^{\gamma_1 z} + \dots + p_\ell(z)e^{\gamma_\ell z}$$

is not identically 0. Then its vanishing order at a point  $z_0$  is smaller than or equal to  $t_1 + \dots + t_\ell - 1$ .

In other terms, when the complex numbers  $\gamma_j$  are distinct, the determinant

$$\left| \left( \frac{d}{dz} \right)^a (z^i e^{\gamma_j z}) \Big|_{z=0} \right|_{\substack{0 \leq i \leq t_j - 1, 1 \leq j \leq \ell \\ 0 \leq a \leq d-1}}$$

is different from 0. This is no surprise that we come across the determinant of the matrix  $M$ .

# Exponential polynomials

The sequence of derivatives of an exponential polynomial evaluated at one point satisfies a linear recurrence relation. Let  $p_1(z), \dots, p_\ell(z)$  be nonzero polynomials of  $\mathbb{C}[z]$  of degrees smaller than  $t_1, \dots, t_\ell$  respectively. Let  $\gamma_1, \dots, \gamma_\ell$  be distinct complex numbers. Suppose that the function

$$F(z) = p_1(z)e^{\gamma_1 z} + \dots + p_\ell(z)e^{\gamma_\ell z}$$

is not identically 0. Then its vanishing order at a point  $z_0$  is smaller than or equal to  $t_1 + \dots + t_\ell - 1$ .

In other terms, when the complex numbers  $\gamma_j$  are distinct, the determinant

$$\left| \left( \frac{d}{dz} \right)^a (z^i e^{\gamma_j z}) \Big|_{z=0} \right|_{\substack{0 \leq i \leq t_j - 1, 1 \leq j \leq \ell \\ 0 \leq a \leq d-1}}$$

is different from 0. This is no surprise that we come across the determinant of the matrix  $M$ .

# The matrix $M$

The determinant of  $M$  is

$$\det M = \prod_{1 \leq i < j \leq \ell} (\gamma_j - \gamma_i)^{t_i t_j}.$$

For  $1 \leq j \leq \ell$ ,  $0 \leq i \leq t_j - 1$ ,  $0 \leq k \leq d - 1$ , the  $(s_j + i, k)$  entry of the matrix  $M$  is

$$\frac{1}{i!} \left( \frac{d}{dT} \right)^i T^k \Big|_{T=\gamma_j} = \binom{k}{i} \gamma_j^{k-i}.$$

The matrix  $M$  is associated with the linear system of  $d$  equations in  $d$  unknowns which amounts to finding a polynomial  $f \in K[z]$  of degree  $< d$  for which the  $d$  numbers

$$\frac{d^i f}{dz^i}(\gamma_j), \quad (1 \leq j \leq \ell, 0 \leq i \leq t_j - 1)$$

take prescribed values.

# The matrix $M$

The determinant of  $M$  is

$$\det M = \prod_{1 \leq i < j \leq \ell} (\gamma_j - \gamma_i)^{t_i t_j}.$$

For  $1 \leq j \leq \ell$ ,  $0 \leq i \leq t_j - 1$ ,  $0 \leq k \leq d - 1$ , the  $(s_j + i, k)$  entry of the matrix  $M$  is

$$\frac{1}{i!} \left( \frac{d}{dT} \right)^i T^k \Big|_{T=\gamma_j} = \binom{k}{i} \gamma_j^{k-i}.$$

The matrix  $M$  is associated with the linear system of  $d$  equations in  $d$  unknowns which amounts to finding a polynomial  $f \in K[z]$  of degree  $< d$  for which the  $d$  numbers

$$\frac{d^i f}{dz^i}(\gamma_j), \quad (1 \leq j \leq \ell, 0 \leq i \leq t_j - 1)$$

take prescribed values.

# The matrix $M$

The determinant of  $M$  is

$$\det M = \prod_{1 \leq i < j \leq \ell} (\gamma_j - \gamma_i)^{t_i t_j}.$$

For  $1 \leq j \leq \ell$ ,  $0 \leq i \leq t_j - 1$ ,  $0 \leq k \leq d - 1$ , the  $(s_j + i, k)$  entry of the matrix  $M$  is

$$\frac{1}{i!} \left( \frac{d}{dT} \right)^i T^k \Big|_{T=\gamma_j} = \binom{k}{i} \gamma_j^{k-i}.$$

The matrix  $M$  is associated with the linear system of  $d$  equations in  $d$  unknowns which amounts to finding a polynomial  $f \in K[z]$  of degree  $< d$  for which the  $d$  numbers

$$\frac{d^i f}{dz^i}(\gamma_j), \quad (1 \leq j \leq \ell, 0 \leq i \leq t_j - 1)$$

take prescribed values.

# Interpolation

Let  $\gamma_j$  ( $1 \leq j \leq \ell$ ) be distinct elements in  $\mathbb{K}$ ,  $t_j$  ( $1 \leq j \leq \ell$ ) be positive integers,  $\eta_{ij}$  ( $1 \leq j \leq \ell$ ,  $0 \leq i \leq t_j - 1$ ) be elements in  $\mathbb{K}$ . Set  $d = t_1 + \cdots + t_\ell$ . There exists a unique polynomial  $f \in \mathbb{K}[z]$  of degree  $< d$  satisfying

$$\frac{d^i f}{dz^i}(\gamma_j) = \eta_{ij}, \quad (1 \leq j \leq \ell, 0 \leq i \leq t_j - 1).$$



# Truncated Taylor expansion

Let  $g \in \mathbb{K}(z)$ , let  $z_0 \in \mathbb{K}$  and let  $t \geq 1$ . Assume  $z_0$  is not a pole of  $g$ . We set

$$T_{g,z_0,t}(z) = \sum_{i=0}^{t-1} \frac{d^i g}{dz^i}(z_0) \frac{(z - z_0)^i}{i!}.$$

In other words,  $T_{g,z_0,t}$  is the unique polynomial in  $\mathbb{K}[z]$  of degree  $< t$  such that there exists  $r(z) \in \mathbb{K}(z)$  having no pole at  $z_0$  with

$$g(z) = T_{g,z_0,t}(z) + (z - z_0)^t r(z).$$

Notice that if  $g$  is a polynomial of degree  $< t$ , then  $g = T_{g,z_0,t}$  for any  $z_0 \in \mathbb{K}$ .

# Truncated Taylor expansion

Let  $g \in \mathbb{K}(z)$ , let  $z_0 \in \mathbb{K}$  and let  $t \geq 1$ . Assume  $z_0$  is not a pole of  $g$ . We set

$$T_{g,z_0,t}(z) = \sum_{i=0}^{t-1} \frac{d^i g}{dz^i}(z_0) \frac{(z - z_0)^i}{i!}.$$

In other words,  $T_{g,z_0,t}$  is the unique polynomial in  $\mathbb{K}[z]$  of degree  $< t$  such that there exists  $r(z) \in \mathbb{K}(z)$  having no pole at  $z_0$  with

$$g(z) = T_{g,z_0,t}(z) + (z - z_0)^t r(z).$$

Notice that if  $g$  is a polynomial of degree  $< t$ , then  $g = T_{g,z_0,t}$  for any  $z_0 \in \mathbb{K}$ .

# Truncated Taylor expansion

Let  $g \in \mathbb{K}(z)$ , let  $z_0 \in \mathbb{K}$  and let  $t \geq 1$ . Assume  $z_0$  is not a pole of  $g$ . We set

$$T_{g,z_0,t}(z) = \sum_{i=0}^{t-1} \frac{d^i g}{dz^i}(z_0) \frac{(z - z_0)^i}{i!}.$$

In other words,  $T_{g,z_0,t}$  is the unique polynomial in  $\mathbb{K}[z]$  of degree  $< t$  such that there exists  $r(z) \in \mathbb{K}(z)$  having no pole at  $z_0$  with

$$g(z) = T_{g,z_0,t}(z) + (z - z_0)^t r(z).$$

Notice that if  $g$  is a polynomial of degree  $< t$ , then  $g = T_{g,z_0,t}$  for any  $z_0 \in \mathbb{K}$ .

# Explicit solution to the interpolation problem

For  $j = 1, \dots, \ell$ , define

$$h_j(z) = \prod_{\substack{1 \leq k \leq \ell \\ k \neq j}} \left( \frac{z - \gamma_k}{\gamma_j - \gamma_k} \right)^{t_k} \quad \text{and} \quad p_j(z) = \sum_{i=0}^{t_j-1} \eta_{ij} \frac{(z - \gamma_j)^i}{i!}.$$

Then the solution  $f$  of the interpolation problem

$$\frac{d^i f}{dz^i}(\gamma_j) = \eta_{ij}, \quad (1 \leq j \leq \ell, 0 \leq i \leq t_j - 1).$$

is given by

$$f = \sum_{j=1}^{\ell} h_j T_{\frac{p_j}{h_j}, \gamma_j, t_j}.$$

# Explicit solution to the interpolation problem

For  $j = 1, \dots, \ell$ , define

$$h_j(z) = \prod_{\substack{1 \leq k \leq \ell \\ k \neq j}} \left( \frac{z - \gamma_k}{\gamma_j - \gamma_k} \right)^{t_k} \quad \text{and} \quad p_j(z) = \sum_{i=0}^{t_j-1} \eta_{ij} \frac{(z - \gamma_j)^i}{i!}.$$

Then the solution  $f$  of the interpolation problem

$$\frac{d^i f}{dz^i}(\gamma_j) = \eta_{ij}, \quad (1 \leq j \leq \ell, 0 \leq i \leq t_j - 1).$$

is given by

$$f = \sum_{j=1}^{\ell} h_j T_{\frac{p_j}{h_j}, \gamma_j, t_j}.$$

Updated: 09/11/2017

**Linear recurrence sequences,  
exponential polynomials  
and Diophantine approximation**

*Michel Waldschmidt*

Institut de Mathématiques de Jussieu — Paris VI

<http://webusers.imj-prg.fr/~michel.waldschmidt/>