

On the *abc* Conjecture and some of its consequences

by

Michel Waldschmidt

Université P. et M. Curie (Paris VI)

<http://www.imj-prg.fr/~michel.waldschmidt/>

Abstract

According to *Nature News*, 10 September 2012, quoting *Dorian Goldfeld*, the *abc* Conjecture is “the most important unsolved problem in Diophantine analysis”. It is a kind of grand unified theory of Diophantine curves : “The remarkable thing about the *abc* Conjecture is that it provides a way of reformulating an infinite number of Diophantine problems,” says *Goldfeld*, “and, if it is true, of solving them.” Proposed independently in the mid-80s by *David Masser* of the University of Basel and *Joseph Oesterlé* of Pierre et Marie Curie University (Paris 6), the *abc* Conjecture describes a kind of balance or tension between addition and multiplication, formalizing the observation that when two numbers *a* and *b* are divisible by large powers of small primes, *a + b* tends to be divisible by small powers of large primes. The *abc* Conjecture implies – in a few lines – the proofs of many difficult theorems and outstanding conjectures in Diophantine equations– including *Fermat’s Last Theorem*.

Abstract (continued)

This talk will be at an elementary level, giving a collection of consequences of the *abc* Conjecture. It will not include an introduction to the Inter-universal *Teichmüller* Theory of *Shinichi Mochizuki*.



<http://www.kurims.kyoto-u.ac.jp/~motizuki/top-english.html>

Poster with Razvan Barbulescu — Archives HAL



<https://hal.archives-ouvertes.fr/hal-01626155>

As simple as abc



American Broadcasting Company



http://fr.wikipedia.org/wiki/American_Broadcasting_Company

Annapurna Base Camp, October 22, 2014



Mt. Annapurna (8091m) is the 10th highest mountain in the world and the journey to its base camp is one of the most popular treks on earth.

<http://www.himalayanglacier.com/trekking-in-nepal/160/annapurna-base-camp-trek.htm>

The radical of a positive integer

According to the fundamental theorem of arithmetic, any integer $n \geq 2$ can be written as a product of prime numbers :

$$n = p_1^{a_1} p_2^{a_2} \cdots p_t^{a_t}.$$

The *radical* (also called *kernel*) $\text{Rad}(n)$ of n is the product of the distinct primes dividing n :

$$\text{Rad}(n) = p_1 p_2 \cdots p_t.$$

$$\text{Rad}(n) \leq n.$$

Examples : $\text{Rad}(2^a) = 2,$

$$\text{Rad}(60\,500) = \text{Rad}(2^2 \cdot 5^3 \cdot 11^2) = 2 \cdot 5 \cdot 11 = 110,$$

$$\text{Rad}(82\,852\,996\,681\,926) = 2 \cdot 3 \cdot 23 \cdot 109 = 15\,042.$$

abc-triples

An *abc*-triple is a triple of three positive integers a, b, c which are coprime, $a < b$ and that $a + b = c$.

Examples:

$$\begin{aligned}1 + 2 &= 3, & 1 + 8 &= 9, \\1 + 80 &= 81, & 4 + 121 &= 125, \\2 + 3^{10} \cdot 109 &= 23^5, & 11^2 + 3^2 5^6 7^3 &= 2^{21} \cdot 23.\end{aligned}$$

13 *abc*-triples with $c < 10$

a, b, c are coprime, $1 \leq a < b$, $a + b = c$ and $c \leq 9$.

$$\begin{aligned}1 + 2 &= 3 \\1 + 3 &= 4 \\1 + 4 &= 5 & 2 + 3 &= 5 \\1 + 5 &= 6 \\1 + 6 &= 7 & 2 + 5 &= 7 & 3 + 4 &= 7 \\1 + 7 &= 8 & & & 3 + 5 &= 8 \\1 + 8 &= 9 & 2 + 7 &= 9 & & & 4 + 5 &= 9\end{aligned}$$

Radical of the *abc*-triples with $c < 10$

$$\begin{aligned}\text{Rad}(1 \cdot 2 \cdot 3) &= 6 \\ \text{Rad}(1 \cdot 3 \cdot 4) &= 6 \\ \text{Rad}(1 \cdot 4 \cdot 5) &= 10 & \text{Rad}(2 \cdot 3 \cdot 5) &= 30 \\ \text{Rad}(1 \cdot 5 \cdot 6) &= 30 \\ \text{Rad}(1 \cdot 6 \cdot 7) &= 42 & \text{Rad}(2 \cdot 5 \cdot 7) &= 70 & \text{Rad}(3 \cdot 4 \cdot 7) &= 42 \\ \text{Rad}(1 \cdot 7 \cdot 8) &= 14 & & & \text{Rad}(3 \cdot 5 \cdot 8) &= 30 \\ \text{Rad}(1 \cdot 8 \cdot 9) &= 6 & \text{Rad}(2 \cdot 7 \cdot 9) &= 54 & \text{Rad}(4 \cdot 5 \cdot 9) &= 30\end{aligned}$$

$$a = 1, b = 8, c = 9, a + b = c, \text{gcd} = 1, \text{Rad}(abc) < c.$$

abc-hits

Following F. Beukers, an *abc*-hit is an *abc*-triple such that $\text{Rad}(abc) < c$.



<http://www.staff.science.uu.nl/~beuke106/ABCpresentation.pdf>

Example: $(1, 8, 9)$ is an *abc*-hit since $1 + 8 = 9$, $\text{gcd}(1, 8, 9) = 1$ and

$$\text{Rad}(1 \cdot 8 \cdot 9) = \text{Rad}(2^3 \cdot 3^2) = 2 \cdot 3 = 6 < 9.$$

On the condition that a, b, c are relatively prime

Starting with $a + b = c$, multiply by a power of a divisor $d > 1$ of abc and get

$$ad^\ell + bd^\ell = cd^\ell.$$

The radical did not increase : the radical of the product of the three numbers ad^ℓ , bd^ℓ and cd^ℓ is nothing else than $\text{Rad}(abc)$; but c is replaced by cd^ℓ .

For ℓ sufficiently large, cd^ℓ is larger than $\text{Rad}(abc)$.

But $(ad^\ell, bd^\ell, cd^\ell)$ is not an abc -hit.

It would be too easy to get examples without the condition that a, b, c are relatively prime.

Some abc -hits

$(1, 80, 81)$ is an abc -hit since $1 + 80 = 81$, $\text{gcd}(1, 80, 81) = 1$ and

$$\text{Rad}(1 \cdot 80 \cdot 81) = \text{Rad}(2^4 \cdot 5 \cdot 3^4) = 2 \cdot 5 \cdot 3 = 30 < 81.$$

$(4, 121, 125)$ is an abc -hit since $4 + 121 = 125$, $\text{gcd}(4, 121, 125) = 1$ and

$$\text{Rad}(4 \cdot 121 \cdot 125) = \text{Rad}(2^2 \cdot 5^3 \cdot 11^2) = 2 \cdot 5 \cdot 11 = 110 < 125.$$

Further abc -hits

- $(2, 3^{10} \cdot 109, 23^5) = (2, 6\,436\,341, 6\,436\,343)$

is an abc -hit since $2 + 3^{10} \cdot 109 = 23^5$ and
 $\text{Rad}(2 \cdot 3^{10} \cdot 109 \cdot 23^5) = 15\,042 < 23^5 = 6\,436\,343$.

- $(11^2, 3^2 \cdot 5^6 \cdot 7^3, 2^{21} \cdot 23) = (121, 48\,234\,275, 48\,234\,496)$

is an abc -hit since $11^2 + 3^2 \cdot 5^6 \cdot 7^3 = 2^{21} \cdot 23$ and
 $\text{Rad}(2^{21} \cdot 3^2 \cdot 5^6 \cdot 7^3 \cdot 11^2 \cdot 23) = 53\,130 < 2^{21} \cdot 23 = 48\,234\,496$.

- $(1, 5 \cdot 127 \cdot (2 \cdot 3 \cdot 7)^3, 19^6) = (1, 47\,045\,880, 47\,045\,881)$

is an abc -hit since $1 + 5 \cdot 127 \cdot (2 \cdot 3 \cdot 7)^3 = 19^6$ and
 $\text{Rad}(5 \cdot 127 \cdot (2 \cdot 3 \cdot 7)^3 \cdot 19^6) = 5 \cdot 127 \cdot 2 \cdot 3 \cdot 7 \cdot 19 = 506\,730$.

abc -triples and abc -hits

Among $15 \cdot 10^6$ abc -triples with $c < 10^4$, we have 120 abc -hits.

Among $380 \cdot 10^6$ abc -triples with $c < 5 \cdot 10^4$, we have 276 abc -hits.

More abc -hits

Recall the abc -hit $(1, 80, 81)$, where $81 = 3^4$.

$$(1, 3^{16} - 1, 3^{16}) = (1, 43\,046\,720, 43\,046\,721)$$

is an abc -hit.

Proof.

$$\begin{aligned} 3^{16} - 1 &= (3^8 - 1)(3^8 + 1) \\ &= (3^4 - 1)(3^4 + 1)(3^8 + 1) \\ &= (3^2 - 1)(3^2 + 1)(3^4 + 1)(3^8 + 1) \\ &= (3 - 1)(3 + 1)(3^2 + 1)(3^4 + 1)(3^8 + 1) \end{aligned}$$

is divisible by 2^6 . (Quotient : 672 605).

Hence

$$\text{Rad}((3^{16} - 1) \cdot 3^{16}) \leq \frac{3^{16} - 1}{2^6} \cdot 2 \cdot 3 < 3^{16}.$$

Infinitely many abc -hits

Proposition. *There are infinitely many abc -hits.*

Take $k \geq 1$, $a = 1$, $c = 3^{2^k}$, $b = c - 1$.

Lemma. 2^{k+2} divides $3^{2^k} - 1$.

Proof : Induction on k using

$$3^{2^k} - 1 = (3^{2^{k-1}} - 1)(3^{2^{k-1}} + 1).$$

Consequence :

$$\text{Rad}((3^{2^k} - 1) \cdot 3^{2^k}) \leq \frac{3^{2^k} - 1}{2^{k+1}} \cdot 3 < 3^{2^k}.$$

Hence

$$(1, 3^{2^k} - 1, 3^{2^k})$$

is an abc -hit.

Infinitely many abc -hits

This argument shows that there exist infinitely many abc -triples such that

$$c > \frac{1}{6 \log 3} R \log R$$

with $R = \text{Rad}(abc)$.

Question : *Are there abc -triples for which $c > \text{Rad}(abc)^2$?*

We do not know the answer.

Examples

When a , b and c are three positive relatively prime integers satisfying $a + b = c$, define

$$\lambda(a, b, c) = \frac{\log c}{\log \text{Rad}(abc)}.$$


Here are the two largest known values for $\lambda(abc)$

$a + b = c$	$\lambda(a, b, c)$	authors
$2 + 3^{10} \cdot 109 = 23^5$	1.629912...	É. Reyssat
$11^2 + 3^2 5^6 7^3 = 2^{21} \cdot 23$	1.625990...	B.M. de Weger

Number of digits of the good abc -triples

At the date of September 11, 2008, 217 abc triples with $\lambda(a, b, c) \geq 1.4$ were known. <https://nitaj.users.lmno.cnrs.fr/tableabc.pdf>

At the date of August 1, 2015, 238 were known. On May 15, 2017, the total is 240. <http://www.math.leidenuniv.nl/~desmit/abc/index.php?sort=1>



The list up to 20 digits is complete.

$$\text{Eric Reyssat : } 2 + 3^{10} \cdot 109 = 23^5$$



Example of Reyssat $2 + 3^{10} \cdot 109 = 23^5$

$$a + b = c$$

$$a = 2, \quad b = 3^{10} \cdot 109, \quad c = 23^5 = 6\,436\,343,$$

$$\text{Rad}(abc) = \text{Rad}(2 \cdot 3^{10} \cdot 109 \cdot 23^5) = 2 \cdot 3 \cdot 109 \cdot 23 = 15\,042,$$

$$\lambda(a, b, c) = \frac{\log c}{\log \text{Rad}(abc)} = \frac{5 \log 23}{\log 15\,042} \simeq 1.62991.$$

Continued fraction

$$2 + 109 \cdot 3^{10} = 23^5$$

Continued fraction of $109^{1/5} : [2; 1, 1, 4, 77733, \dots]$,
approximation : $[2; 1, 1, 4] = 23/9$

$$109^{1/5} = 2.555\,555\,39\dots$$

$$\frac{23}{9} = 2.555\,555\,55\dots$$

N. A. Carella. *Note on the ABC Conjecture*

<http://arXiv.org/abs/math/0606221>

Benne de Weger : $11^2 + 3^2 \cdot 5^6 \cdot 7^3 = 2^{21} \cdot 23$

$\text{Rad}(2^{21} \cdot 3^2 \cdot 5^6 \cdot 7^3 \cdot 11^2 \cdot 23) = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 23 = 53\,130.$

$2^{21} \cdot 23 = 48\,234\,496 = (53\,130)^{1.625990\dots}$



Explicit *abc* Conjecture



According to S. Laishram and T. N. Shorey, an explicit version, due to A. Baker, of the *abc* Conjecture, yields

$$c < \text{Rad}(abc)^{7/4}$$

for any *abc*-triple (a, b, c) .

The *abc* Conjecture

Recall that for a positive integer n , the *radical* of n is

$$\text{Rad}(n) = \prod_{p|n} p.$$

abc Conjecture. Let $\varepsilon > 0$. Then the set of *abc* triples for which

$$c > \text{Rad}(abc)^{1+\varepsilon}$$

is finite.

Equivalent statement : For each $\varepsilon > 0$ there exists $\kappa(\varepsilon)$ such that, if a, b and c in $\mathbf{Z}_{>0}$ are relatively prime and satisfy $a + b = c$, then

$$c < \kappa(\varepsilon)\text{Rad}(abc)^{1+\varepsilon}.$$

Lower bound for the radical of *abc*

The *abc* Conjecture is a **lower bound** for the radical of the product *abc* :

abc Conjecture. For any $\varepsilon > 0$, there exist $\kappa(\varepsilon)$ such that, if a, b and c are relatively prime positive integers which satisfy $a + b = c$, then

$$\text{Rad}(abc) > \kappa(\varepsilon)c^{1-\varepsilon}.$$

The *abc* Conjecture of Oesterlé and Masser



The *abc* Conjecture resulted from a discussion between J. Oesterlé and D. W. Masser in the mid 1980's.


C.L. Stewart and Yu Kunrui

Best known non conditional result : C.L. Stewart and Yu Kunrui (1991, 2001) :

$$\log c \leq \kappa R^{1/3} (\log R)^3.$$

with $R = \text{Rad}(abc)$:

$$c \leq e^{\kappa R^{1/3} (\log R)^3}.$$



Lucien Szpiro

J. Oesterlé and A. Nitaj proved that the *abc* Conjecture implies a previous conjecture by L. Szpiro on the conductor of elliptic curves.



Given any $\varepsilon > 0$, there exists a constant $C(\varepsilon) > 0$ such that, for every elliptic curve with minimal discriminant Δ and conductor N ,

$$|\Delta| < C(\varepsilon) N^{6+\varepsilon}.$$

Szpiro's Conjecture

Conversely, J. Oesterlé proved in 1988 that the conjecture of L. Szpiro implies a weak form of the *abc* conjecture with $1 - \varepsilon$ replaced by $(5/6) - \varepsilon$.



Further examples

When a , b and c are three positive relatively prime integers satisfying $a + b = c$, define

$$\varrho(a, b, c) = \frac{\log abc}{\log \text{Rad}(abc)}.$$

Here are the two largest known values for $\varrho(abc)$, found by A. Nitaj.

$a + b = c$	$\varrho(a, b, c)$
$13 \cdot 19^6 + 2^{30} \cdot 5 = 3^{13} \cdot 11^2 \cdot 31$	4.41901...
$2^5 \cdot 11^2 \cdot 19^9 + 5^{15} \cdot 37^2 \cdot 47 = 3^7 \cdot 7^{11} \cdot 743$	4.26801...

On March 19, 2003, 47 abc triples were known with $0 < a < b < c$, $a + b = c$ and $\text{gcd}(a, b) = 1$ satisfying $\varrho(a, b, c) > 4$.

<https://nitaj.users.lmno.cnrs.fr/tableszipiro.pdf>

33 / 120

Abderrahmane Nitaj

عبد الرحمان نتاج

<https://nitaj.users.lmno.cnrs.fr/abc.html>




Bart de Smit

<http://www.math.leidenuniv.nl/~desmit/abc/>

35 / 120

Escher and the Droste effect



<http://escherdroste.math.leidenuniv.nl/>

36 / 120



ABC@home is an educational and non-profit distributed computing project finding abc-triples related to the ABC conjecture.

The ABC conjecture is currently one of the greatest open problems in mathematics. If it is proven to be true, a lot of other open problems can be answered directly from it.

The ABC conjecture is one of the greatest open mathematical questions, one of the holy grails of mathematics. It will teach us something about our very own numbers.

Fermat's Last Theorem $x^n + y^n = z^n$ for $n \geq 6$

Pierre de Fermat
1601 – 1665

Andrew Wiles
1953 –



Solution in 1994

Fermat's last Theorem for $n \geq 6$ as a consequence of the *abc* Conjecture

Assume $x^n + y^n = z^n$ with $\text{gcd}(x, y, z) = 1$ and $x < y$. Then (x^n, y^n, z^n) is an *abc*-triple with

$$\text{Rad}(x^n y^n z^n) \leq xyz < z^3.$$

If the explicit *abc* Conjecture $c < \text{Rad}(abc)^2$ is true, then one deduces

$$z^n < z^6,$$

hence $n \leq 5$ (and therefore $n \leq 2$).

Square, cubes...

- A **perfect power** is an integer of the form a^b where $a \geq 1$ and $b > 1$ are positive integers.
- **Squares** :
1, 4, 9, 16, 25, 36, 49, 64, 81, 100, 121, 144, 169, 196, ...
- **Cubes** :
1, 8, 27, 64, 125, 216, 343, 512, 729, 1000, 1331, ...
- **Fifth powers** :
1, 32, 243, 1024, 3125, 7776, 16807, 32768, ...

Perfect powers

1, 4, 8, 9, 16, 25, 27, 32, 36, 49, 64, 81, 100, 121, 125, 128, 144, 169, 196, 216, 225, 243, 256, 289, 324, 343, 361, 400, 441, 484, 512, 529, 576, 625, 676, 729, 784, ...



Neil J. A. Sloane's encyclopaedia
<http://oeis.org/A001597>



Two conjectures

Subbayya Sivasankaranarayana Pillai
(1901-1950)

Eugène Charles Catalan (1814 – 1894)

- **Catalan's Conjecture** : In the sequence of perfect powers, 8, 9 is the only example of consecutive integers.
- **Pillai's Conjecture** : In the sequence of perfect powers, the difference between two consecutive terms tends to infinity.

Nearly equal perfect powers

- Difference 1 : (8, 9)
- Difference 2 : (25, 27), ...
- Difference 3 : (1, 4), (125, 128), ...
- Difference 4 : (4, 8), (32, 36), (121, 125), ...
- Difference 5 : (4, 9), (27, 32), ...

Pillai's Conjecture :

- **Pillai's Conjecture** : In the sequence of perfect powers, the difference between two consecutive terms tends to infinity.
- **Alternatively** : Let k be a positive integer. The equation

$$x^p - y^q = k,$$

where the unknowns x , y , p and q take integer values, all ≥ 2 , has only finitely many solutions (x, y, p, q) .

Results

P. Mihăilescu, 2002.

Catalan was right : the equation $x^p - y^q = 1$ where the unknowns x, y, p and q take integer values, all ≥ 2 , has only one solution $(x, y, p, q) = (3, 2, 2, 3)$.



Previous work on Catalan's Conjecture



J.W.S. Cassels, Rob Tijdeman



$$x^p < y^q < \exp \exp \exp \exp(730)$$

Michel Langevin

Previous work on Catalan's Conjecture



Maurice Mignotte



Yuri Bilu

Pillai's conjecture and the abc Conjecture

There is no value of $k \geq 2$ for which one knows that Pillai's equation $x^p - y^q = k$ has only finitely many solutions.

Pillai's conjecture as a consequence of the abc Conjecture : if $x^p \neq y^q$, then

$$|x^p - y^q| \geq c(\epsilon) \max\{x^p, y^q\}^{\kappa - \epsilon}$$

with

$$\kappa = 1 - \frac{1}{p} - \frac{1}{q}$$

Lower bounds for linear forms in logarithms

- A special case of my conjectures with S. Lang for

$$|q \log y - p \log x|$$

yields

$$|x^p - y^q| \geq c(\epsilon) \max\{x^p, y^q\}^{\kappa - \epsilon}$$

with

$$\kappa = 1 - \frac{1}{p} - \frac{1}{q}.$$

Serge Lang
(1927 - 2005)



Not a consequence of the *abc* Conjecture

$$p = 3, q = 2$$

Hall's Conjecture (1971) :


if $x^3 \neq y^2$, then

$$|x^3 - y^2| \geq c \max\{x^3, y^2\}^{1/6}.$$



http://en.wikipedia.org/wiki/Marshall_Hall,_Jr

Conjecture of F. Beukers and C.L. Stewart (2010)



Let p, q be coprime integers with $p > q \geq 2$. Then, for any $c > 0$, there exist infinitely many positive integers x, y such that

$$0 < |x^p - y^q| < c \max\{x^p, y^q\}^\kappa$$

with $\kappa = 1 - \frac{1}{p} - \frac{1}{q}$.

Generalized Fermat's equation $x^p + y^q = z^r$

Consider the equation $x^p + y^q = z^r$ in positive integers (x, y, z, p, q, r) such that x, y, z relatively prime and p, q, r are ≥ 2 .

If

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} \geq 1,$$

then (p, q, r) is a permutation of one of

$$(2, 2, k), (2, 3, 3), (2, 3, 4), (2, 3, 5),$$

$$(2, 4, 4), (2, 3, 6), (3, 3, 3)$$

and in each case the set of solutions (x, y, z) is known (for some of these values there are infinitely many solutions).

Frits Beukers and Don Zagier

For

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1,$$

10 primitive solutions (x, y, z, p, q, r) (up to obvious symmetries) to the equation

$$x^p + y^q = z^r$$

are known.



Primitive solutions to $x^p + y^q = z^r$

Condition : x, y, z are relatively prime

Trivial example of a non primitive solution : $2^p + 2^p = 2^{p+1}$.

Exercise (Claude Levesque) : for any pairwise relatively prime (p, q, r) , there exist positive integers x, y, z with $x^p + y^q = z^r$.

Hint :

$$(17 \times 71^{21})^3 + (2 \times 71^9)^7 = (71^{13})^5.$$

Generalized Fermat's equation

For

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1,$$

the equation

$$x^p + y^q = z^r$$

has the following 10 solutions with x, y, z relatively prime :

$$1 + 2^3 = 3^2, \quad 2^5 + 7^2 = 3^4, \quad 7^3 + 13^2 = 2^9, \quad 2^7 + 17^3 = 71^2,$$

$$3^5 + 11^4 = 122^2, \quad 33^8 + 1\,549\,034^2 = 15\,613^3,$$

$$1\,414^3 + 2\,213\,459^2 = 65^7, \quad 9\,262^3 + 15\,312\,283^2 = 113^7,$$

$$17^7 + 76\,271^3 = 21\,063\,928^2, \quad 43^8 + 96\,222^3 = 30\,042\,907^2.$$

Conjecture of Beal, Granville and Tijdeman–Zagier




The equation $x^p + y^q = z^r$ has no solution in positive integers (x, y, z, p, q, r) with each of p, q and r at least 3 and with x, y, z relatively prime.

<http://mathoverflow.net/>

Andrew Beal

Find a solution with all exponents at least 3, or prove that there is no such solution.



<http://www.forbes.com/2009/04/03/banking-andy-beal-business-wall-street-beal.html>

Beal's Prize

Mauldin, R. D. – A generalization of *Fermat's last theorem* : the *Beal Conjecture* and prize problem. Notices Amer. Math. Soc. **44** N°11 (1997), 1436–1437.

The prize. Andrew Beal is very generously offering a prize of \$5,000 for the solution of this problem. The value of the prize will increase by \$5,000 per year up to \$50,000 until it is solved. The prize committee consists of Charles Fefferman, Ron Graham, and R. Daniel Mauldin, who will act as the chair of the committee. All proposed solutions and inquiries about the prize should be sent to Mauldin.

Beal's Prize : 1, 000, 000\$ US

An AMS-appointed committee will award this prize for either a proof of, or a counterexample to, the *Beal Conjecture* published in a refereed and respected mathematics publication. The prize money – currently US\$1,000,000 – is being held in trust by the AMS until it is awarded. Income from the prize fund is used to support the annual *Erdős* Memorial Lecture and other activities of the Society.

One of *Andrew Beal's* goals is to inspire young people to think about the equation, think about winning the offered prize, and in the process become more interested in the field of mathematics.

<http://www.ams.org/profession/prizes-awards/ams-supported/beal-prize>

Henri Darmon, Andrew Granville

"Fermat-Catalan" Conjecture (H. Darmon and A. Granville), consequence of the *abc Conjecture* : *the set of solutions* (x, y, z, p, q, r) to $x^p + y^q = z^r$ with x, y, z relatively prime and $(1/p) + (1/q) + (1/r) < 1$ is finite.



Hint: $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$ implies $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} \leq \frac{41}{42}$.

1995 (H. Darmon and A. Granville) : unconditionally, for fixed (p, q, r) , only finitely many (x, y, z) .

Henri Darmon, Loïc Merel : $(p, p, 2)$ and $(p, p, 3)$

Unconditional results by H. Darmon and L. Merel (1997) :

For $p \geq 4$, the equation $x^p + y^p = z^2$ has no solution in relatively prime positive integers x, y, z .

For $p \geq 3$, the equation $x^p + y^p = z^3$ has no solution in relatively prime positive integers x, y, z .



Fermat's Little Theorem

For $a > 1$, any prime p not dividing a divides $a^{p-1} - 1$.

Hence if p is an odd prime, then p divides $2^{p-1} - 1$.



Wieferich primes (1909) : p^2 divides $2^{p-1} - 1$

The only known **Wieferich** primes are 1093 and 3511. These are the only ones below $4 \cdot 10^{12}$.

Infinitely many primes are not **Wieferich** assuming abc



Joseph H. Silverman

J.H. Silverman : if the abc Conjecture is true, given a positive integer $a > 1$, there exist infinitely many primes p such that p^2 does not divide $a^{p-1} - 1$.

Nothing is known about the finiteness of the set of **Wieferich** primes.

Consecutive integers with the same radical

Notice that

$$75 = 3 \cdot 5^2 \quad \text{and} \quad 1215 = 3^5 \cdot 5$$

hence

$$\text{Rad}(75) = \text{Rad}(1215) = 3 \cdot 5 = 15.$$

But also

$$76 = 2^2 \cdot 19 \quad \text{and} \quad 1216 = 2^6 \cdot 19$$

have the same radical

$$\text{Rad}(76) = \text{Rad}(1216) = 2 \cdot 19 = 38.$$

Consecutive integers with the same radical

For $k \geq 1$, the two numbers

$$x = 2^k - 2 = 2(2^{k-1} - 1)$$

and

$$y = (2^k - 1)^2 - 1 = 2^{k+1}(2^{k-1} - 1)$$

have the same radical, and also

$$x + 1 = 2^k - 1 \quad \text{and} \quad y + 1 = (2^k - 1)^2$$

have the same radical.

Consecutive integers with the same radical

Are there further examples of $x \neq y$ with

$$\text{Rad}(x) = \text{Rad}(y) \quad \text{and} \quad \text{Rad}(x + 1) = \text{Rad}(y + 1)?$$

Is it possible to find two distinct integers x, y such that


$$\text{Rad}(x) = \text{Rad}(y),$$

$$\text{Rad}(x + 1) = \text{Rad}(y + 1)$$

and

$$\text{Rad}(x + 2) = \text{Rad}(y + 2)?$$

Erdős – Woods Conjecture



<http://school.maths.uwa.edu.au/~woods/>

There exists an absolute constant k such that, if x and y are positive integers satisfying

$$\text{Rad}(x + i) = \text{Rad}(y + i)$$

for $i = 0, 1, \dots, k - 1$, then $x = y$.

Erdős – Woods as a consequence of abc

M. Langevin : The abc Conjecture implies that there exists an absolute constant k such that, if x and y are positive integers satisfying

$$\text{Rad}(x + i) = \text{Rad}(y + i)$$

for $i = 0, 1, \dots, k - 1$, then $x = y$.



Already in 1975 M. Langevin studied the radical of $n(n + k)$ with $\gcd(n, k) = 1$ using lower bounds for linear forms in logarithms (Baker's method).

A factorial as a product of factorials

For $n > a_1 \geq a_2 \geq \dots \geq a_t > 1$, $t > 1$, consider

$$a_1! a_2! \cdots a_t! = n!$$

Trivial solutions :

$$2^r! = (2^r - 1)! 2!^r \text{ with } r \geq 2.$$

Non trivial solutions :

$$7! 3! 22! = 9!, \quad 7! 6! = 10!, \quad 7! 5! 3! = 10!, \quad 14! 5! 2! = 16!.$$

Saranya Nair and Tarlok Shorey : The effective *abc* conjecture implies Hickerson's conjecture that the largest non-trivial solution is given by $n = 16$.



69 / 120

Erdős Conjecture on $2^n - 1$

In 1965, P. Erdős conjectured that the greatest prime factor $P(2^n - 1)$ satisfies

$$\frac{P(2^n - 1)}{n} \rightarrow \infty \text{ when } n \rightarrow \infty.$$



In 2002, R. Murty and S. Wong proved that this is a consequence of the *abc* Conjecture.

In 2012, C.L. Stewart proved Erdős Conjecture (in a wider context of Lucas and Lehmer sequences) :

$$P(2^n - 1) > n \exp(\log n / 104 \log \log n).$$

70 / 120

Is *abc* Conjecture optimal ?



Let $\delta > 0$. In 1986, C.L. Stewart and R. Tijdeman proved that there are infinitely many *abc*-triples for which

$$c > R \exp\left((4 - \delta) \frac{(\log R)^{1/2}}{\log \log R}\right).$$

Better than $c > R \log R$.

71 / 120

Conjectures by Machiel van Frankenhuijsen, Olivier Robert, Cam Stewart and Gérald Tenenbaum

Let $\varepsilon > 0$. There exists $\kappa(\varepsilon) > 0$ such that for any *abc* triple with $R = \text{Rad}(abc) > 8$,


$$c < \kappa(\varepsilon) R \exp\left((4\sqrt{3} + \varepsilon) \left(\frac{\log R}{\log \log R}\right)^{1/2}\right).$$

Further, there exist infinitely many *abc*-triples for which

$$c > R \exp\left((4\sqrt{3} - \varepsilon) \left(\frac{\log R}{\log \log R}\right)^{1/2}\right).$$

72 / 120

Machiel van Frankenhuijsen, Olivier Robert, Cam Stewart and Gérald Tenenbaum



Heuristic assumption

Whenever a and b are coprime positive integers, $R(a + b)$ is independent of $R(a)$ and $R(b)$.

O. Robert, C.L. Stewart and G. Tenenbaum, *A refinement of the abc conjecture*, Bull. London Math. Soc., Bull. London Math. Soc. (2014) **46** (6) : 1156-1166.

<http://blms.oxfordjournals.org/content/46/6/1156.full.pdf>

http://iecl.univ-lorraine.fr/~Gerald.Tenenbaum/PUBLIC/Prepublications_et_publications/abc.pdf

Waring's Problem

In 1770, a few months before J.L. Lagrange solved a conjecture of Bachet (1621) and Fermat (1640) by proving that every positive integer is the sum of at most four squares of integers, E. Waring wrote :



Edward Waring
(1736 - 1798)

"Omnis integer numerus vel est cubus, vel e duobus, tribus, 4, 5, 6, 7, 8, vel novem cubis compositus, est etiam quadrato-quadratus vel e duobus, tribus, & usque ad novemdecim compositus, & sic deinceps"

"Every integer is a cube or the sum of two, three, . . . nine cubes; every integer is also the square of a square, or the sum of up to nineteen such; and so forth. Similar laws may be affirmed for the correspondingly defined numbers of quantities of any like degree."

Waring's functions $g(k)$ and $G(k)$

- Waring's function g is defined as follows : For any integer $k \geq 2$, $g(k)$ is the least positive integer s such that any positive integer N can be written $x_1^k + \dots + x_s^k$.
- Waring's function G is defined as follows : For any integer $k \geq 2$, $G(k)$ is the least positive integer s such that any sufficiently large positive integer N can be written $x_1^k + \dots + x_s^k$.

J.L. Lagrange : $g(2) = 4$.

$g(2) \leq 4$: any positive number is a sum of at most 4 squares :

$$n = x_1^2 + x_2^2 + x_3^2 + x_4^2.$$

$g(2) \geq 4$: there are positive numbers (for instance 7) which are not sum of 3 squares.



Joseph-Louis Lagrange
(1736 – 1813)

Lower bounds are easy, not upper bounds.

$g(4) \geq 19$.

We want to write 79 as sum $a_1^4 + a_2^4 + \dots + a_s^4$ with s as small as possible.

Since $79 < 81$, we cannot use 3^4 . Hence we can use only $2^4 = 16$ and $1^4 = 1$.

Since $79 < 5 \times 16$, we can use at most 4 terms 2^4 .

Now

$$79 = 64 + 15 = 4 \times 2^4 + 15 \times 1^4$$

with 4 + 15 terms a^4 (namely 4 with 2^4 and 15 with 1^4).

The number of terms is 19.

$n = x_1^4 + \dots + x_{19}^4 : g(4) = 19$

Any positive integer is the sum of at most 19 biquadrates
R. Balasubramanian, J-M. Deshouillers, F. Dress (1986).



François Dress, R. Balasubramanian, Jean-Marc Deshouillers

Evaluations of $g(k)$ for $k = 2, 3, 4, \dots$

$g(2) = 4$	Lagrange	1770
$g(3) = 9$	Kempner	1912
$g(4) = 19$	Balasubramanian, Dress, Deshouillers	1986
$g(5) = 37$	Chen Jingrun	1964
$g(6) = 73$	Pillai	1940
$g(7) = 143$	Dickson	1936

Lower bound for $g(k)$

Let $k \geq 2$. Select $N < 3^k$ of the form $N = 2^k q - 1$. Since $N < 3^k$, writing N as a sum of k -th powers can involve no term 3^k , and since $N < 2^k q$, it involves at most $(q - 1)$ terms 2^k , all others being 1^k ; so the most economical way of writing N as a sum of k -th powers is

$$N = (q - 1)2^k + (2^k - 1)1^k$$

which requires a total number of $(q - 1) + (2^k - 1)$ terms. The largest value is obtained by taking for q the largest integer with $2^k q < 3^k$. Since $(3/2)^k$ is not an integer, this integer q is $\lfloor (3/2)^k \rfloor$ (quotient of the division of 3^k by 2^k).

$$g(k) \geq I(k)$$

For each integer $k \geq 2$, define $I(k) = 2^k + \lfloor (3/2)^k \rfloor - 2$. Then $g(k) \geq I(k)$.



(J. A. Euler, son of Leonhard Euler).

The ideal Waring's "Theorem" : $g(k) = I(k)$

Recall

$$I(k) = 2^k + \lfloor (3/2)^k \rfloor - 2.$$

Conjecture (C.A. Bretschneider, 1853) : $g(k) = I(k)$ for any $k \geq 2$.

Divide 3^k by 2^k :

$$3^k = 2^k q + r \quad \text{with} \quad 0 < r < 2^k, \quad q = \lfloor (3/2)^k \rfloor$$

The remainder $r = 3^k - 2^k q$ satisfies $r < 2^k$. A slight improvement of this upper bound would yield the desired result. L.E. Dickson and S.S. Pillai proved independently in 1936 that $g(k) = I(k)$, provided that $r = 3^k - 2^k q$ satisfies

$$r \leq 2^k - q - 2 \quad \text{with} \quad q = \lfloor (3/2)^k \rfloor.$$

The condition $r \leq 2^k - q - 2$

The condition $r \leq 2^k - q - 2$ is satisfied for $4 \leq k \leq 471\,600\,000$.

If, for some k , the condition $r \leq 2^k - q - 2$ is not satisfied, then $(3/2)^k$ is extremely close to an integer :

$$q + 1 - \frac{q - 2}{2^k} < \left(\frac{3}{2}\right)^k < q + 1,$$

which is unlikely : one expects that the numbers $(3/2)^k$ are well distributed modulo 1.

Mahler's contribution

- The estimate

$$r \leq 2^k - q - 2$$

is valid for all sufficiently large k .

Kurt Mahler
(1903 - 1988)



Hence the ideal Waring's Theorem

$$g(k) = 2^k + \lfloor (3/2)^k \rfloor - 2$$

holds for all sufficiently large k .

Waring's Problem and the abc Conjecture

S. David : the estimate

$$r \leq 2^k - q - 2$$

for sufficiently large k follows from the abc Conjecture.



S. Laishram : the ideal Waring's Theorem

$g(k) = 2^k + \lfloor (3/2)^k \rfloor - 2$ follows from the explicit abc Conjecture.

Conjecture of Alan Baker (1996)

Let (a, b, c) be an abc -triple and let $\epsilon > 0$. Then

$$c \leq \kappa (\epsilon^{-\omega} R)^{1+\epsilon}$$

where κ is an absolute constant, $R = \text{Rad}(abc)$ and $\omega = \omega(abc)$ is the number of distinct prime factors of abc .

Remark of Andrew Granville : the minimum of the function on the right hand side over $\epsilon > 0$ occurs essentially with $\epsilon = \omega / \log R$. This yields a slightly sharper form of the conjecture :

$$c \leq \kappa R \frac{(\log R)^\omega}{\omega!}.$$

Alan Baker : explicit abc Conjecture (2004)

Let (a, b, c) be an abc -triple. Then

$$c \leq \frac{6}{5} R \frac{(\log R)^\omega}{\omega!}$$

with $R = \text{Rad}(abc)$ the radical of abc and $\omega = \omega(abc)$ the number of distinct prime factors of abc .



Shanta Laishram and Tarlok Shorey



The Nagell–Ljunggren equation is the equation

$$y^q = \frac{x^n - 1}{x - 1}$$

in integers $x > 1$, $y > 1$,
 $n > 2$, $q > 1$.

This means that in basis x , all the digits of the perfect power y^q are 1.

If the explicit abc -conjecture of Baker is true, then the only solutions are

$$11^2 = \frac{3^5 - 1}{3 - 1}, \quad 20^2 = \frac{7^4 - 1}{7 - 1}, \quad 7^3 = \frac{18^3 - 1}{18 - 1}.$$


The abc conjecture for number fields

P. Vojta (1987) - variants due to D.W. Masser and K. Györy



The abc conjecture for number fields (continued)

Survey by J. Browkin.



Jerzy Browkin
(1934 – 2015)

The abc -conjecture for
Algebraic Numbers
Acta Mathematica Sinica,
Jan., 2006, Vol. 22, No. 1,
pp. 211–222

<http://dx.doi.org/10.1007/s10114-005-0624-3>

Mordell's Conjecture (Faltings's Theorem)

Using an effective extension of the abc Conjecture for a number field, N. Elkies deduces an effective version of Faltings's Theorem on the finiteness of the set of rational points on an algebraic curve of genus ≥ 2 over the same number field.

L.J. Mordell (1922)

G. Faltings (1984)

N. Elkies (1991)



<http://www.math.harvard.edu/~elkies/>

The *abc* conjecture for number fields



Andrea Surroca

The effective *abc* Conjecture implies an effective version of Siegel's Theorem on the finiteness of the set of integer points on a curve.

A. Surroca, *Méthodes de transcendance et géométrie diophantienne*, Thèse, Université de Paris 6, 2003.

Thue–Siegel–Roth Theorem (Bombieri)

Using the *abc* Conjecture for number fields, E. Bombieri (1994) deduces a refinement of the Thue–Siegel–Roth Theorem on the rational approximation of algebraic numbers

$$\left| \alpha - \frac{p}{q} \right| > \frac{1}{q^{2+\varepsilon}}$$

where he replaces ε by

$$\kappa(\log q)^{-1/2}(\log \log q)^{-1}$$

where κ depends only on the algebraic number α .



Siegel's zeroes (A. Granville and H.M. Stark)

The uniform *abc* Conjecture for number fields implies a lower bound for the class number of an imaginary quadratic number field, and K. Mahler has shown that this implies that the associated *L*-function has no Siegel zero.



abc and Vojta's height Conjecture



Paul Vojta

Vojta stated a conjectural inequality on the height of algebraic points of bounded degree on a smooth complete variety over a global field of characteristic zero which implies the *abc* Conjecture.

Further consequences of the *abc* Conjecture

- Erdős's Conjecture on consecutive powerful numbers.
- Dressler's Conjecture : between two positive integers having the same prime factors, there is always a prime (Cochrane and textcolormacouleurDressler 1999).
- Squarefree and powerfree values of polynomials (Browkin, Filaseta, Greaves and Schinzel, 1995).
- Lang's conjectures : lower bounds for heights, number of integral points on elliptic curves (Frey 1987, Hindry Silverman 1988).
- Bounds for the order of the Tate–Shafarevich group (Goldfeld and Szpiro 1995).
- Greenberg's Conjecture on Iwasawa invariants λ and μ in cyclotomic extensions (Ichimura 1998).
- Lower bound for the class number of imaginary quadratic fields (Granville and Stark 2000), hence no Siegel zero for the associated L -function (Mahler).
- Fundamental units of certain quadratic and biquadratic fields (Katayama 1999).
- The height conjecture and the degree conjecture (Frey 1987, Mai and Murty 1996)

The n -Conjecture



Nils Bruin, Generalization of the ABC-conjecture, Master Thesis, Leiden University, 1995.

<http://www.cecm.sfu.ca/~nbruin/scriptie.pdf>

Let $n \geq 3$. There exists a positive constant κ_n such that, if x_1, \dots, x_n are relatively prime rational integers satisfying $x_1 + \dots + x_n = 0$ and if no proper subsum vanishes, then

$$\max\{|x_1|, \dots, |x_n|\} \leq \text{Rad}(x_1 \cdots x_n)^{\kappa_n}.$$

? Should hold for all but finitely many (x_1, \dots, x_n) with $\kappa_n = 2n - 5 + \epsilon$?

A consequence of the n -Conjecture

Open problem : for $k \geq 5$, no positive integer can be written in two essentially different ways as sum of two k -th powers.

It is not even known whether such a k exists.

Reference : Hardy and Wright : §21.11

For $k = 4$ (Euler) :

$$59^4 + 158^4 = 133^4 + 134^4 = 635\,318\,657$$

A parametric family of solutions of $x_1^4 + x_2^4 = x_3^4 + x_4^4$ is known

Reference : <http://mathworld.wolfram.com/DiophantineEquation4thPowers.html>

abc and meromorphic function fields



Nevanlinna value distribution theory.

Recent work of Hu, Pei-Chu, Yang, Chung-Chun and P. Vojta.

ABC Theorem for polynomials

Let K be an algebraically closed field. The *radical* of a monic polynomial

$$P(X) = \prod_{i=1}^n (X - \alpha_i)^{a_i} \in K[X]$$

with α_i pairwise distinct is defined as

$$\text{Rad}(P)(X) = \prod_{i=1}^n (X - \alpha_i) \in K[X].$$

ABC Theorem for polynomials

ABC Theorem (A. Hurwitz, W.W. Stothers, R. Mason).

Let A, B, C be three relatively prime polynomials in $K[X]$ with $A + B = C$ and let $R = \text{Rad}(ABC)$. Then

$$\max\{\deg(A), \deg(B), \deg(C)\} < \deg(R).$$



Adolf Hurwitz (1859–1919)

This result can be compared with the *abc* Conjecture, where the degree replaces the logarithm.

The radical of a polynomial as a gcd

The common zeroes of

$$P(X) = \prod_{i=1}^n (X - \alpha_i)^{a_i} \in K[X]$$

and P' are the α_i with $a_i \geq 2$. They are zeroes of P' with multiplicity $a_i - 1$. Hence

$$\text{Rad}(P) = \frac{P}{\gcd(P, P')}.$$

Proof of the ABC Theorem for polynomials

Now suppose $A + B = C$ with A, B, C relatively prime.

Notice that

$$\text{Rad}(ABC) = \text{Rad}(A)\text{Rad}(B)\text{Rad}(C).$$

We may suppose A, B, C to be monic and, say, $\deg(A) \leq \deg(B) \leq \deg(C)$.

Write

$$A + B = C, \quad A' + B' = C',$$

and

$$AB' - A'B = AC' - A'C.$$

Proof of the ABC Theorem for polynomials

Recall $\gcd(A, B, C) = 1$. Since $\gcd(C, C')$ divides $AC' - A'C = AB' - A'B$, it divides also

$$\frac{AB' - A'B}{\gcd(A, A') \gcd(B', B')}$$

which is a polynomial of degree

$$< \deg(\text{Rad}(A)) + \deg(\text{Rad}(B)) = \deg(\text{Rad}(AB)).$$

Hence

$$\deg(\gcd(C, C')) < \deg(\text{Rad}(AB))$$

and

$$\deg(C) < \deg(\text{Rad}(C)) + \deg(\text{Rad}(AB)) = \deg(\text{Rad}(ABC)).$$

Shinichi Mochizuki



INTER-UNIVERSAL
TEICHMÜLLER THEORY
IV :
LOG-VOLUME
COMPUTATIONS AND
SET-THEORETIC
FOUNDATIONS
by
Shinichi Mochizuki

<http://www.kurims.kyoto-u.ac.jp/~motizuki/top-english.html>

Shinichi Mochizuki@RIMS

<http://www.kurims.kyoto-u.ac.jp/~motizuki/top-english.html>

EXIT



Papers of Shinichi Mochizuki

- General Arithmetic Geometry
- Intrinsic Hodge Theory
- p -adic Teichmüller Theory
- Anabelian Geometry, the Geometry of Categories
- The Hodge-Arakelov Theory of Elliptic Curves
- Inter-universal Teichmüller Theory

Shinichi Mochizuki

[1] Inter-universal Teichmüller Theory I : Construction of Hodge Theaters. PDF

[2] Inter-universal Teichmüller Theory II : Hodge-Arakelov-theoretic Evaluation. PDF

[3] Inter-universal Teichmüller Theory III : Canonical Splittings of the Log-theta-lattice. PDF

[4] Inter-universal Teichmüller Theory IV : Log-volume Computations and Set-theoretic Foundations. PDF

https://en.wikipedia.org/wiki/Abc_conjecture

In August 2012, Shinichi Mochizuki released a series of four preprints announcing a proof of the *abc* Conjecture.




When an error in one of the articles was pointed out by Vesselin Dimitrov and Akshay Venkatesh in October 2012, Mochizuki posted a comment on his website acknowledging the mistake, stating that it would not affect the result, and promising a corrected version in the near future. He proceeded to post a series of corrected papers of which the latest dated November 2017.

<http://www.kurims.kyoto-u.ac.jp/~motizuki/top-english.html>

<https://www.maths.nottingham.ac.uk/personal/ibf/files/symcor.iut.html>

Inter-universal Teichmüller Theory

- [1] Inter-universal Teichmüller Theory I: Construction of Hodge Theaters. PDF **NEW !! (2017-08-18)**
- [2] Inter-universal Teichmüller Theory II: Hodge-Arakelov-theoretic Evaluation. PDF **NEW !! (2017-08-18)**
- [3] Inter-universal Teichmüller Theory III: Canonical Splittings of the Log-theta-lattice. PDF **NEW !! (2017-11-01)**
- [4] Inter-universal Teichmüller Theory IV: Log-volume Computations and Set-theoretic Foundations. PDF **NEW !! (2017-11-01)**



Workshop on IUT Theory of Shinichi Mochizuki, December 7-11 2015

CMI Workshop supported by Clay Math Institute and Symmetries and Correspondences

Organisers : Ivan Fesenko, Minhyong Kim, Kobi Kremnitzer
Finding the speakers and the program of the workshop : Ivan Fesenko

CMI Workshop supported by Clay Math Institute and Symmetries and Correspondences

The work (currently being refereed) of SHINICHI MOCHIZUKI on inter-universal Teichmüller theory (also known as arithmetic deformation theory) and its application to famous conjectures in diophantine geometry became publicly available in August 2012. This theory, developed over 20 years, introduces a vast collection of novel ideas, methods and objects. Aspects of the theory extend arithmetic geometry to a non-scheme-theoretic setting and, more generally, have the potential to open new fundamental areas of mathematics.

The workshop aims to present and analyse key principles, concepts, objects and proofs of the theory of Mochizuki and study its relations with existing theories in different areas, to help to increase the number of experts in the theory of Mochizuki and stimulate its further applications.

Speakers

Shinichi Mochizuki will answer questions during skype sessions of the workshop. He also responds directly to emailed questions.

Invited speakers : Oren Ben-Bassat, Weronika Czerniawska, Yuichiro Hoshi, Ariyan Javanpeykar, Kiran Kedlaya, Robert Kucharczyk, Ulf Kühn, Lars Kuehne, Emmanuel Lepage, Chung Pang Mok, Jakob Stix, Tamás Szamuely, Fucheng Tan, Go Yamashita, Shou-Wu Zhang.

Participants

Participants:

Julio Andrade (Univ. Oxford), Federico Bambozzi (Univ. Regensburg), Alexander Beilinson (Univ. Chicago), Oren Ben-Bassat (Univ. Haifa), Brian Birch (Univ. Oxford), Francis Brown (Univ. Oxford), Martin Bridson (Univ. Oxford), Olivia Caramello (Univ. Paris 7), Brian Conrad (Stanford Univ.), Weronika Czerniawska (Univ. Nottingham), Ishai Dan-Cohen (Univ. Duisburg-Essen), Jamshid Derakhshan (Univ. Oxford), Taylor Dupuy (Univ. California Los Angeles), Gerd Faltings (MPIM, Bonn), Ivan Fesenko (Univ. Nottingham), Gerhard Frey (Univ. Duisburg-Essen), Adam Gai (Univ. Oxford), Lena Gai (Univ. Oxford), Dorian Goldfeld (Columbia Univ.), Nigel Hitchin (Univ. Oxford), Yuichiro Hoshi (RIMS, Kyoto Univ.), Alexander Ivanov (Techn. Univ. München), Artur Jackson (Purdue Univ.), Ariyan Javanpeykar (Univ. Mainz), Kiran Kedlaya (Univ. California San Diego), Minhyong Kim (Univ. Oxford), Kobi Kremnitzer (Univ. Oxford), Robert Kucharczyk (ETH, Zurich), Ulf Kühn (Univ. Hamburg), Lars Kuehne (MPIM, Bonn), Laurent Lafforgue (IHES, Bures-sur-Yvette), Emmanuel Lepage (Univ. Paris 7), Junghwan Lim (Univ. Oxford), Angus Macintyre (Univ. Oxford), Nils Matthes (Univ. Hamburg), Chung Pang Mok (Morningside Center Mathematics Beijing and Purdue Univ.), Alexander Cruz Morales (MPIM, Bonn), Sergey Olbezin (Univ. Nottingham), Alexander G. Oldenziel (Utrecht Univ.), Thomas Oliver (Univ. Bristol), Florian Pop (Univ. Pennsylvania at Philadelphia), Damian Rössler (Univ. Oxford), Thomas Scanlon (Univ. California Berkeley), Francisco Simkievich (Univ. Oxford), Jakob Stix (Univ. Frankfurt), Tamás Szamuely (Rényi Inst. Math., Budapest), Fucheng Tan (Shanghai Cent. Math. Sc. & Shanghai Jiao Tong Univ.), Dinesh Thakur (Rochester Univ.), Ulrike Tillmann (Univ. Oxford), Wester van Urk (Univ. Nottingham), Felipe Voloch (Univ. Texas Austin), Matthew Waller (Univ. Nottingham), Andrew Wiles (Univ. Oxford), Bora Yalkinoglu (Univ. Strasbourg), Go Yamashita (RIMS, Kyoto Univ.), Fernando Garcia Yamauti (Univ. Sao Paulo), Shou-Wu Zhang (Princeton Univ.), Boris Zilber (Univ. Oxford), Lorenzo Lane (Univ. Edinburgh)

State of the art concerning Mochizuki's contribution (Nov. 2017)

The submitted proof is more than 500 pages long and is currently being peer-reviewed.

Ivan Fesenko estimates that the proof has been verified at least 30 times in §3.1 of the most recent updated version <https://www.maths.nottingham.ac.uk/personal/ibf/notesoniut.pdf> of his survey.

Links

Not Even Wrong

Latest on abc

Posted on December 16, 2017 by [PETER WOIT](#)

<http://www.math.columbia.edu/~woit/wordpress/?p=9871>

The ABC conjecture has (still) not been proved

Posted on December 17, 2017 by [FRANK CALEGARI](#)

<https://galoisrepresentations.wordpress.com/2017/12/17/the-abc-conjecture-has-still-not-been-proved/>

[HECTOR PASTEN](#)

Shimura curves and the abc conjecture

<https://arxiv.org/abs/1705.09251>

Reference

[Michel Waldschmidt](#)

On the *abc* conjecture and some of its consequences.

Mathematics in 21st Century, 6th World Conference, Lahore, March 2013,

([P. Cartier](#), [A.D.R. Choudary](#), [M. Waldschmidt](#) Editors),

Springer Proceedings in Mathematics and Statistics **98** (2015), 211–230.

<http://www.imj-prg.fr/~michel.waldschmidt/articles/pdf/abcLahoreProceedings.pdf>

Poster with Razvan Barbulescu — Archives HAL

February 15, 2018



<https://hal.archives-ouvertes.fr/hal-01626155>

On the *abc* Conjecture and some of its consequences

by

[Michel Waldschmidt](#)

Université P. et M. Curie (Paris VI)

<http://www.imj-prg.fr/~michel.waldschmidt/>