

Abstract

We prove an equidistribution theorem à la Bader-Muchnik ([3]) for operator-valued measures associated with boundary representations in the context of discrete groups of isometries of CAT(-1) spaces thanks to an equidistribution theorem of T. Roblin ([22]). This result can be viewed as a von Neumann's mean ergodic theorem for quasi-invariant measures. In particular, this approach gives a dynamical proof of the fact that boundary representations are irreducible. Moreover, we prove some equidistribution results for conformal densities using elementary techniques from harmonic analysis.

AMS subject classifications: Primary 37A25, 37A30; Secondary 43A65, 43A90.

AMS keywords: Conformal densities, boundary representations, ergodic theorems, irreducibility, equidistribution.

Adrien Boyer, Technion, Haifa, Israel.

E-mail address: adrienboyer@technion.ac.il.

Equidistribution, Ergodicity and Irreducibility in CAT(-1) spaces

Adrien Boyer *

August 25, 2016

1 Introduction

Any action of a locally compact group G on a measure space (X, μ) where μ is a G -quasi-invariant measure gives rise to a unitary representation, after renormalization with the square root of the Radon-Nikodym derivative of the action of G on (X, μ) . This unitary representation is called a *quasi-regular representation*, and generalizes the standard notion of quasi-regular representations given by $G \curvearrowright G/H$ where H is a closed subgroup of G , and G/H always carries a G -quasi-invariant measure.

The dynamical properties of the action $G \curvearrowright (X, \mu)$ can be reflected in a such representation.

In the context of fundamental groups of compact negatively curved manifolds, U. Bader and R. Muchnik prove in [3, Theorem 3] an equidistribution theorem for some operator-valued measures. This theorem can be thought as a generalization of von Neumann's mean ergodic theorem for quasi-invariant measures for fundamental groups acting on the Gromov boundary of universal covers of compact negatively curved manifolds endowed with the Patterson-Sullivan measures. These quasi-regular representations are called *boundary representations*. It turns out that the irreducibility of boundary representations follows from this generalization of von Neumann's ergodic theorem. We refer to [4],[3],[11],[12],[14] and [19] for examples of natural irreducible quasi-regular representations which are related to the following conjecture:

Conjecture 1. *For a locally compact group G and a spread-out probability measure μ on G , the quasi-regular representation associated to a μ -boundary of G is irreducible.*

In this paper, we generalize the work of U. Bader and R. Muchnik to convex cocompact groups of isometries of a CAT(-1) space with a non-arithmetic spectrum and to (non-uniform) lattices of a non-compact connected semisimple Lie group of rank one. Our results are based on the fundamental work of T. Roblin in [22]. The main tool of this paper is an equidistribution theorem of T. Roblin (see Subsection 4.3) which is inspired by the ideas of G. Margulis (see [20]), based on the mixing property of the geodesic flow. Following the technical ideas developed in [3] and using Roblin's equidistribution theorem, we obtain a dynamical explanation of irreducibility of boundary representations in the context of CAT(-1) spaces: it comes from the mixing property of the geodesic flow. Nevertheless this approach does not work in the context of general hyperbolic groups and we refer to [11], [12], [19] and more recently [14] for different approaches.

*The author is supported by ERC Grant 306706.

Moreover, we prove two equidistribution results for densities associated with the Poisson kernel and the square root of the Poisson Kernel in CAT(-1) spaces with respect to the weak* convergence of the dual space L^1 functions on the boundary.

Main Results

The Banach space of finite signed measures on a topological compact space Z is, by the Riesz representation theorem, the dual of the space of the continuous functions $C(Z)$. The Banach space of bounded linear operators from the Banach space of continuous functions to the Banach space of bounded operators on a Hilbert space will be denoted by $\mathcal{L}(C(Z), \mathcal{B}(\mathcal{H}))$. Observe that $\mathcal{L}(C(Z), \mathcal{B}(\mathcal{H}))$ is isomorphic as a Banach space to the dual of the Banach space $C(Z) \widehat{\otimes} \mathcal{H} \widehat{\otimes} \overline{\mathcal{H}}$ where $\overline{\mathcal{H}}$ denotes the conjugate Hilbert space of the complex Hilbert space \mathcal{H} , and $\widehat{\otimes}$ denotes the projective tensor product. Hence $\mathcal{L}(C(Z), \mathcal{B}(\mathcal{H}))$ will be called the space of *operator-valued measures*.

Let Γ be a non-elementary discrete group of isometries of (X, d) a proper CAT(-1) metric space (i.e. the balls are relatively compact). We denote by ∂X its Gromov boundary, and let \overline{X} be the topological space $X \cup \partial X$ endowed with its usual topology that makes \overline{X} compact. Recall the critical exponent $\alpha(\Gamma)$ of Γ :

$$\alpha(\Gamma) := \inf \left\{ s \in \mathbb{R}_+^* \mid \sum_{\gamma \in \Gamma} e^{-sd(\gamma x, x)} < \infty \right\}.$$

Notice that the definition of $\alpha(\Gamma)$ does not depend on x . We assume from now on that $\alpha(\Gamma) < \infty$.

The limit set of Γ denoted by Λ_Γ is the set of all accumulation points in ∂X of an orbit. Namely $\Lambda_\Gamma := \overline{\Gamma x} \cap \partial X$, with the closure in \overline{X} . Notice that the limit set does not depend on the choice of $x \in X$. Following the notations in [8], define the geodesic hull $GH(\Lambda_\Gamma)$ as the union of all geodesics in X with both endpoints in Λ_Γ . The convex hull of Λ_Γ denoted by $CH(\Lambda_\Gamma)$, is the smallest subset of X containing $GH(\Lambda_\Gamma)$ with the property that every geodesic segment between any pair of points $x, y \in CH(\Lambda_\Gamma)$ also lies in $CH(\Lambda_\Gamma)$. We say that Γ is *convex cocompact* if it acts cocompactly on $CH(\Lambda_\Gamma)$.

The translation length of an element $\gamma \in \Gamma$ is defined as $t(\gamma) := \inf \{d(x, \gamma x), x \in X\}$. The *spectrum* of Γ is defined as the subgroup of \mathbb{R} generated by $t(\gamma)$ where γ ranges over the hyperbolic isometries in Γ . We say that Γ has an arithmetic spectrum if its spectrum is a discrete subgroup of \mathbb{R} . We are interested in discrete groups with a *non-arithmetic spectrum* because they guarantee the mixing property of the geodesic flow (see Subsection 2.2), and this condition is verified in the following cases: for isometries group of Riemannian surfaces, hyperbolic spaces and isometries groups of a CAT(-1) space such that the limit set has a non-trivial connected component. We refer to [10] and to [22, Proposition 1.6, Chapitre 1] for more details.

A Riemannian symmetric space X of non-compact type of rank one endowed with its natural Riemannian metric is a particular case of CAT(-1) space. The space X as well as its boundary ∂X can be described by the quotients $X = G/K$ and $\partial X = G/Q$ where G is a non-compact connected semisimple Lie group of real rank one, K a maximal compact subgroup and Q a minimal parabolic subgroup of G . A *lattice* Γ is a discrete subgroup of G such that the quotient $\Gamma \backslash G$ has finite volume with respect to the Haar measure. In this case $\Lambda_\Gamma = \partial X$ and $CH(\Lambda_\Gamma) = X$. If $\Gamma \backslash G$ is a compact, we say that Γ is a uniform lattice and this is a particular case of convex compact groups. Otherwise we say that Γ is a *non-uniform lattice*.

The foundations of Patterson-Sullivan measures theory are in the important papers [21], [27]. See [6],[7], and [22] for more general results in the context of CAT(-1) spaces. These measures are also called *conformal densities*.

We denote by $M(Z)$ the Banach space of Radon measures on a locally compact space Z , which is identified with the dual space of compactly supported functions denoted by $C_c(Z)^*$, endowed with the norm $\|\mu\| = \sup\{|\int_Z f d\mu|, \|f\|_\infty \leq 1, f \in C_c(Z)\}$ where $\|f\|_\infty = \sup_{z \in Z} |f(z)|$. Recall that $\gamma_*\mu$ means $\gamma_*\mu(B) = \mu(\gamma^{-1}B)$ where γ is in Γ and B is a borel subset of Z .

We say that μ is a Γ -invariant conformal density of dimension $\alpha \geq 0$, if μ is a map which satisfies the following conditions:

1. μ is a map from $x \in X \mapsto \mu_x \in M(\overline{X})$, i.e. μ_x is a positive finite measure (density).
2. For all x and y in X , μ_x and μ_y are equivalent, and we have

$$\frac{d\mu_y}{d\mu_x}(v) = \exp(\alpha\beta_v(x, y))$$

(conformal of dimension α).

3. For all $\gamma \in \Gamma$, and for all $x \in X$ we have $\gamma_*\mu_x = \mu_{\gamma x}$ (invariant),

where $\beta_v(x, y)$ denotes the horospherical distance from x to y relative to v (see Subsection 2.1).

If X is a CAT(-1) space and if Γ is a discrete group of isometries of X , then there exists a Γ -invariant conformal density of dimension $\alpha(\Gamma)$ whose support is Λ_Γ . A proof can be found in [21] and [27] for the case of hyperbolic spaces and see [3] and [6] for the case of CAT(-1) spaces.

A conformal density μ gives rise to unitary representations $(\pi_x)_{x \in X}$ defined for $x \in X$ as:

$$\begin{aligned} \pi_x : \Gamma &\rightarrow \mathcal{U}(L^2(\partial X, \mu_x)) \\ (\pi_x(\gamma)\xi)(v) &= \xi(\gamma^{-1}v) \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right), \end{aligned} \tag{1.1}$$

where $\xi \in L^2(\partial X, \mu_x)$ and $v \in \partial X$.

These representations are unitarily equivalent: the multiplication operator

$$U_{xy} : \xi \in L^2(\partial X, \mu_x) \rightarrow (m_{xy} \cdot \xi) \in L^2(\partial X, \mu_y)$$

defined by the function

$$m_{xy}(v) = \exp\left(-\frac{\alpha}{2}\beta_v(x, y)\right),$$

is a unitary operator which intertwines the unitary representations π_x and π_y .

The matrix coefficient

$$\phi_x : \Gamma \rightarrow \langle \pi_x(\gamma)\mathbf{1}_{\partial X}, \mathbf{1}_{\partial X} \rangle \in \mathbb{R}^+, \tag{1.2}$$

where $\mathbf{1}_{\partial X}$ is the characteristic function of ∂X , is called the *Harish-Chandra function*.

Pick x in X , and a positive real number ρ and define for all integers n such that $n \geq \rho$ the annulus

$$C_n(x, \rho) = \{\gamma \in \Gamma | n - \rho \leq d(\gamma x, x) < n + \rho\}.$$

Assume that $C_n(x, \rho)$ is not empty for $n \geq N_{x, \rho}$ where $N_{x, \rho}$ denotes some non-negative integer. Let $|C_n(x, \rho)|$ be the cardinality of $C_n(x, \rho)$, let D_y be the unit Dirac mass centered at a point $y \in X$ and consider the sequence of operator-valued measures defined for all $n \geq N_{x, \rho}$ as:

$$\mathcal{M}_{x, \rho}^n : f \in C(\overline{X}) \mapsto \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x}(f) \frac{\pi_x(\gamma)}{\phi_x(\gamma)} \in \mathcal{B}(L^2(\partial X, \mu_x)). \quad (1.3)$$

If $f \in C(\overline{X})$, we denote by $f|_{\partial X}$ its continuous restriction to the space ∂X . Consider also the operator-valued measure \mathcal{M}_x defined as:

$$\mathcal{M}_x : f \in C(\overline{X}) \mapsto \left(\mathcal{M}_x(f) : \xi \mapsto \left(\int_{\partial X} \xi \frac{d\mu_x}{\|\mu_x\|} \right) \frac{1}{\|\mu_x\|} f|_{\partial X} \right) \in \mathcal{B}(L^2(\partial X, \mu_x)). \quad (1.4)$$

The class \mathcal{C} :

Let Γ be a discrete group of isometries of a CAT(-1) space X . The boundary possesses a natural structure of metric space and more specifically the boundary carries a family of visual metric $(d_x)_{x \in X}$ (see Section 2.1). Let $\mu = (\mu_x)_{x \in X}$ be a conformal densities of dimension α .

Let (Z, d) be a compact metric measure space with a metric d and a measure μ . We denote by $\text{Diam}(Z)$ the diameter of Z . We say that the metric measure space (Z, d, μ) is Ahlfors α -regular for some $\alpha > 0$ if there exists a positive constant $C > 0$ such that for all z in Z and $0 < r < \text{Diam}(Z)$ we have

$$C^{-1}r^\alpha \leq \mu(B(z, r)) \leq Cr^\alpha.$$

Definition 1.1. *Then we say that Γ is in \mathcal{C} if Γ is a discrete group of isometries of a CAT(-1) space such that:*

1. Γ has non-arithmetic spectrum.
2. The metric measure space $(\Lambda_\Gamma, d_x, \mu_x)$ is α -Ahlfors regular for some x in X .

In particular the class \mathcal{C} contains the convex cocompact groups of isometries of a CAT(-1) space with a non-arithmetic spectrum and the non-uniform lattices in noncompact semisimple Lie group of rank one acting by isometries on their rank one symmetric spaces of noncompact type where d is a left invariant Riemannian metric. Moreover the class \mathcal{C} contains groups which are neither convex cocompact nor lattices, see Remark 3.2.

The main result of this paper is the following theorem:

Theorem A. *(Equidistribution à la Bader-Muchnik)*

Let Γ be in \mathcal{C} and let μ be a Γ -invariant conformal density of dimension $\alpha(\Gamma)$. Then for each x in X there exists $\rho > 0$ such that

$$\mathcal{M}_{x, \rho}^n \rightharpoonup \mathcal{M}_x$$

as $n \rightarrow +\infty$ with respect to the weak topology of the Banach space $\mathcal{L}(C(\overline{X}), \mathcal{B}(L^2(\partial X, \mu_x)))$.*

With the same hypothesis of the above theorem, we deduce immediately an ergodic theorem à la von Neumann for the Γ -quasi-invariant measures μ_x on ∂X .

Let $x \in X$, and denote by \mathcal{Q}_x the orthogonal projection onto the subspace of constant functions of $L^2(\partial X, \mu_x)$.

Corollary B. (*Ergodicity à la von Neumann*)
For all $x \in X$ there exists $\rho > 0$ such that

$$\frac{\|\mu_x\|^2}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\pi_x(\gamma)}{\phi_x(\gamma)} \rightarrow \mathcal{Q}_x$$

as $n \rightarrow +\infty$ with respect to the weak operator topology in $\mathcal{B}(L^2(\partial X, \mu_x))$.

Remark 1.1. Consider an action of \mathbb{Z} on a finite measure space (X, μ) by measure preserving transformations. Von Neumann's very well-known ergodic theorem states, in the functional analytic framework, that the ergodicity of the action is equivalent to the convergence

$$\frac{1}{2n+1} \sum_{k=-n}^n \pi(k) \rightarrow \mathcal{Q}$$

with respect to the weak operator topology, where π is the quasi-regular representation obtained from the action of \mathbb{Z} , and where \mathcal{Q} is the orthogonal projection onto the space of constant functions of the space $L^2(X, \mu)$. This theorem belongs to the foundation of ergodic theory and remains an important source of inspiration (see for example [18]).

With the same hypothesis of Theorem A we have:

Corollary C. (*Irreducibility*)

For all $x \in X$, the representations $\pi_x : \Gamma \rightarrow \mathcal{U}(L^2(\partial X, \mu_x))$ are irreducible.

Notice that Corollary C for lattices is well known, see [9] and the method of [14] applies to the case of convex-cocompact groups. Nevertheless, this approach based on Roblin's theorem unify the irreducibility via an ergodic theorem for quasi-invariant measures and gives precise asymptotic limit of operators for the groups in the class \mathcal{C} . Moreover this approach seems to be the right approach to prove irreducibility in the more general context of boundary representations associated with discrete groups with parabolic elements. Note that for some discrete groups with parabolic elements acting on the hyperbolic plane \mathbb{H}^n we obtain obtain the irreducibility of boundary representation (see Remark 3.2). The approach developped in [14] will not be fruitful whenever the group possesses parabolic elements since it works only for hyperbolic groups and this strengthens the dynamical approach of Bader and Muchnik developed here with Roblin's equidistribution theorem.

The Poisson kernel

Recall the definition of the Poisson kernel in the context of CAT(-1) spaces. Let μ be a Γ -invariant conformal density of dimension α . First define

$$p : (x, y, v) \in X \times X \times \partial X \mapsto p(x, y, v) = \exp(\beta_v(x, y)) \in \mathbb{R}^+. \quad (1.5)$$

Fix $x \in X$ a base point and define the *Poisson kernel associated to the measure μ_x* as:

$$P : (y, v) \in X \times \partial X \mapsto P(y, v) = p^\alpha(x, y, v) = \exp(\alpha\beta_v(x, y)) \in \mathbb{R}^+. \quad (1.6)$$

We follow the notations of Sjögren ([24]) and we define for $\lambda \in \mathbb{R}$ and $f \in L^1(\partial X, \mu_x)$:

$$P_\lambda f(y) = \int_{\partial X} P(y, v)^{\lambda+1/2} f(v) d\mu_x(v).$$

Furthermore we denote by ν_y the measure associated to P_0 defined as

$$d\nu_y(v) = \frac{P(y, v)^{1/2}}{P_0 \mathbf{1}_{\partial X}(y)} d\mu_x(v). \quad (1.7)$$

Observe that the measure ν_y is a probability measure. We refer to Subsection 2.2 for the definition of the Bowen-Margulis-Sullivan measure occurring in the following statement:

Theorem D. (*Equidistribution*)

• Let Γ be a discrete group of isometries of a CAT(-1) space X with a non-arithmetic spectrum. Let μ be a Γ -invariant conformal density of dimension $\alpha(\Gamma)$ the critical exponent of the group. Assume that Γ has a finite Bowen-Margulis-Sullivan measure and assume that there exists a constant $C > 0$ such that for all x, y in X we have

$$\frac{\|\mu_x\|}{\|\mu_y\|} \leq C.$$

Then for all $x \in X$ and for all $\rho > 0$ we have

$$\frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\mu_{\gamma x}}{\|\mu_{\gamma x}\|} \rightharpoonup \frac{\mu_x}{\|\mu_x\|}$$

with respect to the weak* convergence of $L^1(\partial X, \mu_x)^*$.

• Let Γ be in \mathcal{C} , then for all x in $CH(\Lambda_\Gamma)$ there exists ρ such that

$$\frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \nu_{\gamma x} \rightharpoonup \nu_x$$

with respect to the weak* convergence of $L^1(\partial X, \nu_x)^*$.

The method of proofs of Theorem A and D consists of two steps: given a sequence of functionals of the dual of a separable Banach space:

Step 1: The sequence is uniformly bounded: existence of accumulation points (by the Banach-Alaoglu theorem).

Step 2: Identification of the limit using equidistribution theorems (only one accumulation point).

Structure of the paper

In Section 2 we remind the reader of some standard facts about CAT(-1) spaces as well as the definition of Bowen-Margulis-Sullivan measures and Roblin's equidistribution theorem. We recall also some general facts about Banach spaces and projective tensor products, and we give a general construction of operator-valued measures that we investigate in the context of CAT(-1) spaces.

In Section 3 we prove uniform boundedness for two sequences of functions, and we deduce **Step 1** of our results.

In Section 4 we use Roblin's equidistribution theorem to achieve **Step 2** of our main result.

In Section 5 we prove Theorem A, Corollary B and Corollary C.

In Section 6 we prove Theorem D using the dual inequality established in Section 3.

Acknowledgements

I would like to thank Christophe Pittet and Uri Bader for useful discussions and criticisms. I would also like to thank Peter Haïssinsky, Marc Bourdon and Davis Simmons for discussions about Ahlfors regularity. I am grateful to Felix Pogorzelski, Dustin Mayeda and Antoine Pinochet Lobos for their remarks on this work. Finally, I would like to thank the referee for carefully reading the paper.

2 Preliminaries

2.1 CAT(-1) spaces

In this section we survey the geometry of CAT(-1) spaces. We freely rely on [6] where the reader could consult for further details.

A CAT(-1) space is a metric geodesic space such that every geodesic triangle is thinner than its comparison triangle in the hyperbolic plane, see [5, Introduction]. Let (X, d) be a proper CAT(-1) space. A geodesic ray of (X, d) is an isometry:

$$r : I \rightarrow X,$$

where $I = [0, +\infty) \subset \mathbb{R}$. Two geodesic rays are equivalent if the Hausdorff distance between their images are bounded, equivalently $\sup_{t \in I} d(r_1(t), r_2(t)) < +\infty$. If r is a geodesic ray, $r(+\infty)$ denotes its equivalence class. The boundary ∂X is defined as the set of equivalence classes of geodesic rays.

A geodesic segment of (X, d) is an isometry:

$$r : I \rightarrow X,$$

where $I = [0, a]$ with $a < \infty$.

Fix a base point x . We denote by $\mathcal{R}(x)$ the set of geodesic rays and of geodesic segments starting at x with the following convention: if r is a geodesic segment defined on $[0, a]$, we set $r(t) = r(a)$ for all $t > a$. Hence we have a natural map

$$\begin{aligned} \mathcal{R}(x) &\rightarrow \overline{X} = X \cup \partial X \\ r &\mapsto r(+\infty), \end{aligned}$$

which is surjective. The set $\mathcal{R}(x)$ is endowed with the topology of uniform convergence on compact subsets of $[0, +\infty)$. By the Arzela-Ascoli theorem, $\mathcal{R}(x)$ is a compact space. Hence, endowed with the quotient topology, \overline{X} is compact. Notice that the topology on \overline{X} does not depend on the choice of x , see [5, 3.7 Proposition (1), p. 429].

Since the CAT(-1) spaces are a particular class of general δ -hyperbolic spaces we have the following inequality: for all $x, y, z, t \in \overline{X}$

$$(x, z)_t \geq \min\{(x, y)_t, (y, z)_t\} - \delta, \tag{2.1}$$

see [5, 3.17 Remarks (4), p. 433].

Let x be in X , and let r be a geodesic ray. By the triangle inequality the function $t \mapsto d(x, r(t)) - t$ is decreasing and bounded below. Recall that the Busemann function associated to a geodesic ray r , is defined as the function

$$b_r(x) = \lim_{t \rightarrow \infty} d(x, r(t)) - t.$$

Let x and y be in X , and let v be in ∂X . Let r be a geodesic ray whose extremity is v , namely $r(+\infty) = v$. The limit $\lim_{t \rightarrow \infty} d(x, r(t)) - d(y, r(t))$ exists, is equal to $b_r(x) - b_r(y)$, and is independent of the choice of r . The horospherical distance from x to y relative to v is defined as

$$\beta_v(x, y) = \lim_{t \rightarrow \infty} d(x, r(t)) - d(y, r(t)). \quad (2.2)$$

It satisfies for all $v \in \partial X$, and for all $x, y \in X$ that

$$\beta_v(x, y) = -\beta_v(y, x) \quad (2.3)$$

$$\beta_v(x, y) + \beta_v(y, z) = \beta_v(x, z) \quad (2.4)$$

$$\beta_v(x, y) \leq d(x, y). \quad (2.5)$$

If γ is an isometry of X we have

$$\beta_{\gamma v}(\gamma x, \gamma y) = \beta_v(x, y). \quad (2.6)$$

Recall that the Gromov product of two points $a, b \in X$ relative to $x \in X$ is

$$(a, b)_x = \frac{1}{2}(d(x, a) + d(x, b) - d(a, b)).$$

Let v, w be in ∂X such that $v \neq w$. If $a_n \rightarrow v \in \partial X$, $b_n \rightarrow w \in \partial X$, then

$$(v, w)_x = \lim_{n \rightarrow \infty} (a_n, b_n)_x$$

exists and does not depend on v and w .

Let r be a geodesic ray which represents v . We have

$$(v, y)_x = \lim_{t \rightarrow +\infty} \frac{1}{2}(d(x, r(t)) + d(x, y) - d(r(t), y)),$$

then we obtain:

$$\beta_v(x, y) = 2(v, y)_x - d(x, y). \quad (2.7)$$

Besides, if $q \in X$ is a point of the geodesic defined by v and w , then we also have:

$$(v, w)_x = \frac{1}{2}(\beta_v(x, q) + \beta_w(x, q)).$$

The formula

$$d_x(v, w) = \exp(- (v, w)_x) \quad (2.8)$$

defines a distance on ∂X (we set $d_x(v, v) = 0$). This is due to M. Bourdon in the context of CAT(-1) spaces, we refer to [6, 2.5.1 Théorème] for more details. We have the following comparison formula:

$$d_y(v, w) = \exp\left(\frac{1}{2}(\beta_v(x, y) + \beta_w(x, y))\right) d_x(v, w).$$

We say that $(d_x)_{x \in X}$ is a family of visual metrics. A ball of radius r centered at $v \in \partial X$ with respect to d_x is denoted by $B(v, r)$. A ball of radius r centered at $y \in X$ is denoted by $B_X(y, r)$.

If γ is an isometry of (X, d) , its conformal factor at $v \in \partial X$ is:

$$\lim_{w \rightarrow v} \frac{d_x(\gamma v, \gamma w)}{d_x(v, w)} = \exp(\beta_v(x, \gamma^{-1}x)),$$

(see [6, 2.6.3 Corollaire]).

If x and y are points of X and R is a positive real number, we define the shadow

$$\mathcal{O}_R(x, y)$$

to be the set of v in ∂X such that the geodesic ray issued from x with limit point v hits the closed ball of center y with radius $R > 0$.

The Sullivan shadow lemma is a very useful tool in ergodic theory of discrete groups acting on a CAT(-1) space. See for example [22, Lemma 1.3] for a proof.

Lemma 2.1. (*D. Sullivan*) *Let Γ be a discrete group of isometries of X . Let $\mu = (\mu_x)_{x \in X}$ a Γ -invariant conformal density of dimension α . Let x be in X . Then for R large enough there exists $C > 0$ such that for all $\gamma \in \Gamma$:*

$$\frac{1}{C} \exp(-\alpha d(x, \gamma x)) \leq \mu_x(\mathcal{O}_R(x, \gamma x)) \leq C \exp(-\alpha d(x, \gamma x)).$$

2.2 Bowen-Margulis-Sullivan measures and Roblin's equidistribution theorem

We follow [22, Chapitre 1 Préliminaires, 1C. Flot géodésique] where the reader could find more details.

In [27], D. Sullivan constructs measures on the unit tangent bundle of X where X is the n -dimensional real hyperbolic space, and proves striking results for this new class of measures. We refer to [27] for more details about these measures. We recall the definitions of these analogous measures in CAT(-1) spaces.

Let SX be the set of isometries from \mathbb{R} to (X, d) endowed with the topology of uniform convergence on compact subsets of \mathbb{R} . In other words, SX is the set of geodesics of X parametrized by \mathbb{R} . We have a canonical "projection" from SX to X , playing the role of the projection from the unit tangent of bundle of a manifold to the manifold, which associates to $u \in SX$ a point in X . Indeed, notice this map in the setting of CAT(-1) spaces may be not surjective since all the geodesics are not bi-infinite.

The trivial flow on \mathbb{R} induces a continuous flow $(g_t)_{t \in \mathbb{R}}$ on SX , called the geodesic flow. For $u \in SX$, we will denote by $g_{+\infty}(u)$ the end of the geodesic determined by u for the positive time and $g_{-\infty}(u)$ the end of the geodesic for the negative time. Let $\partial^2 X$ be the set: $\partial X \times \partial X - \{(x, x) | x \in \partial X\}$. We recall now the so-called *Hopf parametrization* in CAT(-1) spaces and to do so we fix an origin $x \in X$. We have an identification of SX with $\partial^2 X \times \mathbb{R}$ via

$$u \mapsto (g_{-\infty}(u), g_{+\infty}(u), \beta_{g_{-\infty}(u)}(u, x)).$$

Observe that Γ acts on $\partial^2 X \times \mathbb{R}$ by $\gamma \cdot (v, w, s) = (\gamma v, \gamma w, s + \beta_v(x, \gamma^{-1}x))$, and \mathbb{R} acts on $\partial^2 X \times \mathbb{R}$ by translation $t \cdot (v, w, s) = g_t((v, w, s)) = (v, w, s + t)$. Notice these actions commute on SX .

Let μ be a Γ -invariant conformal density of dimension α . The Bowen-Margulis-Sullivan measure which is referred to as the *BMS measure* m on SX is defined as:

$$dm(u) = \frac{d\mu_x(v)d\mu_x(w)ds}{d_x(v,w)^{2\alpha}}.$$

The measure m is invariant by the action of the geodesic flow, and observe also that m is a Γ -invariant measure. We denote by m_Γ the measure on the quotient SX/Γ . More precisely if D is a fundamental domain for the action of Γ on SX , if h is a compactly supported function in $C_c(SX/\Gamma)$ and if \tilde{h} denotes the lift of h in $C(SX)$ we have $\int_D \tilde{h} dm = \int_{SX/\Gamma} h dm_\Gamma$. Moreover the quantity $\int_D \tilde{h} dm$ does not depend on the choice of D .

We say that Γ admits a BMS finite measure if m_Γ is finite. We denote by g_Γ^t the geodesic flow on SX/Γ . We say that g_Γ^t is mixing on SX/Γ with respect to m_Γ if for all bounded Borel subsets $A, B \subset SX/\Gamma$ we have $\lim_{t \rightarrow +\infty} m_\Gamma(A \cap g_\Gamma^t(B)) = m_\Gamma(A)m_\Gamma(B)$.

The assumption of non-arithmeticity of the spectrum of Γ guarantees that the geodesic flow on SX satisfies the mixing property with respect to BMS measures. We refer to [2, Proposition 7.7] for a proof of this fact in the case of negatively curved manifold. We refer to [22, Chapitre 3] for a general proof in CAT(-1) spaces.

In [22, Théorème 4.1.1, Chapitre 4], T. Roblin proves the following theorem based on the mixing property of the geodesic flow on $SX \setminus \Gamma$ with respect to BMS measures:

Theorem 2.1. (*T. Roblin*) *Let Γ be a discrete group of isometries of X with a non-arithmetic spectrum. Assume that Γ admits a finite BMS measure associated to a Γ -invariant conformal density μ of dimension $\alpha = \alpha(\Gamma)$. Then for all $x, y \in X$ we have:*

$$\alpha e^{-\alpha n} \|m_\Gamma\| \sum_{\{\gamma \in \Gamma | d(x, \gamma y) < n\}} D_{\gamma^{-1}x} \otimes D_{\gamma y} \rightharpoonup \mu_x \otimes \mu_y$$

as $n \rightarrow +\infty$ with respect to the weak* convergence of $C(\overline{X} \times \overline{X})^*$.

2.3 Operator-valued measures

2.3.1 The space of operator-valued measures as a dual space of a Banach space

We remind to the reader why the Banach space $\mathcal{L}(C(Z), \mathcal{B}(\mathcal{H}))$ is naturally isomorphic to the dual of the Banach space $C(\partial X) \widehat{\otimes} \mathcal{H} \widehat{\otimes} \overline{\mathcal{H}}$ where $\widehat{\otimes}$ denotes the projective tensor product:

Let E and F be Banach spaces with norms $\|\cdot\|_E$ and $\|\cdot\|_F$. We consider the algebraic tensor product $E \otimes_{alg} F$. The projective norm of an element g in $E \otimes_{alg} F$ is defined by

$$\|g\|_p := \inf \left\{ \sum_{\text{finite}} \|e_i\|_E \|f_i\|_F, \text{ such that } g = \sum_{\text{finite}} e_i \otimes f_i \right\}.$$

The projective tensor product is defined as the completion of the algebraic tensor product for the projective norm $\|\cdot\|_p$, and it is denoted by

$$E \widehat{\otimes} F := \overline{E \otimes_{alg} F}^{\|\cdot\|_p}.$$

Recall also that we have the Banach isomorphism

$$\mathcal{L}(E, F^*) \rightarrow (E \widehat{\otimes} F)^* \tag{2.9}$$

given by:

$$\mathcal{M} \mapsto (e \otimes f \mapsto \mathcal{M}(e)f).$$

See [23, p. 24] for more details.

Let $\langle \cdot, \cdot \rangle$ be the inner product on \mathcal{H} which is antilinear on the second variable. Define for $\xi \in \mathcal{H}$ the map $\xi^* \in \mathcal{H}^*$ which satisfies $\xi^*(\zeta) = \langle \zeta, \xi \rangle$ for $\zeta \in \mathcal{H}$. The canonical isomorphism between a conjugate Hilbert space and its dual is given by:

$$\xi \in \overline{\mathcal{H}} \mapsto \xi^* \in \mathcal{H}^*$$

Define the map

$$\xi \otimes \eta^* \in \mathcal{H} \otimes \mathcal{H}^* \mapsto t_{\xi, \eta} \in \mathcal{B}(\mathcal{H})$$

where

$$\forall \zeta \in \mathcal{H}, t_{\xi, \eta}(\zeta) = \eta^*(\zeta)\xi = \langle \zeta, \eta \rangle \xi.$$

Let Tr be the usual semi-finite trace on $\mathcal{B}(\mathcal{H})$ and let T be an operator in $\mathcal{B}(\mathcal{H})$. Notice that for all ξ and η in \mathcal{H} :

$$\langle T\xi, \eta \rangle = Tr(Tt_{\xi, \eta}). \quad (2.10)$$

It is well known that we have the isomorphism

$$\xi \otimes \eta \in \mathcal{H} \widehat{\otimes} \overline{\mathcal{H}} \mapsto t_{\xi, \eta} \in L^1(\mathcal{H}), \quad (2.11)$$

where $L^1(\mathcal{H})$ denotes the space of *Trace class* operators.

Recall that we have also an isomorphism

$$T \in \mathcal{B}(\mathcal{H}) \mapsto Tr_T \in L^1(\mathcal{H})^*, \quad (2.12)$$

where $Tr_T(S) = Tr(TS)$ for all $S \in L^1(\mathcal{H})$.

Recall that $T_n \rightarrow T$ with respect to the weak operator topology if for all ξ and η in \mathcal{H} we have $\langle T_n \xi, \eta \rangle \rightarrow \langle T \xi, \eta \rangle$ as $n \rightarrow +\infty$.

An explicit isomorphism

Let Z be a compact space. The space $\mathcal{L}(C(Z), \mathcal{B}(\mathcal{H}))$ is a Banach space with the norm $\|\mathcal{M}\| = \sup\{\|\mathcal{M}(f)\|_{\mathcal{B}(\mathcal{H})}, \text{ with } \|f\|_\infty \leq 1\}$. Combining isomorphisms (2.9), (2.11), and (2.12) with the observation (2.10) we obtain that the map $\mathcal{M} \in \mathcal{L}(C(Z), \mathcal{B}(\mathcal{H})) \mapsto \widetilde{\mathcal{M}} \in (C(Z) \widehat{\otimes} \mathcal{H} \widehat{\otimes} \overline{\mathcal{H}})^*$ is a Banach isomorphism and satisfies for all $(f, \xi, \eta) \in (C(Z) \times \mathcal{H} \times \overline{\mathcal{H}})$:

$$\widetilde{\mathcal{M}}(f \otimes \xi \otimes \eta) = Tr(\mathcal{M}(f)t_{\xi, \eta}) = \langle \mathcal{M}(f)\xi, \eta \rangle. \quad (2.13)$$

2.3.2 General construction of Operator-valued measures

We give in this section a general construction of “ergodic” operator-valued measures that we are interested in.

Quasi-regular representations

Let (Y, μ) be a measure space. Consider an action $\Gamma \curvearrowright (Y, \mu)$ such that μ is a finite Γ -quasi-invariant measure (i.e. μ and $\gamma_*\mu$ are in the same measure class). We denote by

$$\frac{d\gamma_*\mu}{d\mu}(y)$$

the Radon-Nikodym derivative of $\gamma_*\mu$ with respect to μ at a point y , with γ in Γ . Let \mathcal{H} be $L^2(Y, \mu)$ and for all $\xi \in \mathcal{H}$ and for all γ in Γ define π to be:

$$(\pi(\gamma)\xi)(y) = \left(\frac{d\gamma_*\mu}{d\mu} \right)^{\frac{1}{2}}(y) \xi(\gamma^{-1}y).$$

The representation $\pi : \Gamma \rightarrow \mathcal{U}(\mathcal{H})$ is a unitary representation on the Hilbert space \mathcal{H} , and is called a *quasi-regular* representation. Observe that π is a *positive representation* in the sense that π preserves the cone of positive functions.

Notice that π extends to a representation of the group algebra denoted $\mathbb{C}\Gamma$ by

$$\pi : \sum c_\gamma \gamma \in \mathbb{C}\Gamma \mapsto \sum c_\gamma \pi(\gamma) \in \mathcal{B}(\mathcal{H}).$$

Define also the following matrix coefficient

$$\phi : \gamma \in \Gamma \mapsto \langle \pi(\gamma)\mathbf{1}_Y, \mathbf{1}_Y \rangle \in \mathbb{R}^+,$$

where $\mathbf{1}_Y$ denotes the characteristic function of the measure space Y .

An ergodic operator-valued measure

Let Z be a topological space and consider the space of continuous functions on Z denoted by $C(Z)$. Consider a family of linear forms $(\ell_\gamma)_{\gamma \in \Gamma}$ on $C(Z)^*$. Assume that Γ acts isometrically on a metric space (X, d) . Let $x \in X$ and $\rho > 0$. Define for all $n \geq \rho$ the annulus

$$C_n(x, \rho) := \{n - \rho \leq d(\gamma x, x) < n + \rho\}.$$

Assume that there exists an integer $N_{x, \rho}$ that for all $n \geq N_{x, \rho}$ the annulus $C_n(x, \rho)$ is not empty. Define the sequence of operator-valued measures $(\mathcal{M}_{x, \rho}^n)_{n \geq N_{x, \rho}}$ as:

$$\mathcal{M}_{x, \rho}^n : f \in C(Z) \mapsto \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \ell_\gamma(f) \frac{\pi(\gamma)}{\phi(\gamma)}$$

and observe

$$\mathcal{M}_{x, \rho}^n(f) = \pi \left(\frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \ell_\gamma(f) \frac{\gamma}{\phi(\gamma)} \right) \in \mathcal{B}(\mathcal{H}).$$

Properties

Let T be a bounded operator on a Hilbert space and T^* is its adjoint. Let $\mathbf{1}_Z$ and $\mathbf{1}_Y$ be the constant functions which are equal to 1 on Z and on Y . The Banach space $L^\infty(Y)$ is a Banach space with its usual norm $\|\cdot\|_\infty$. We denote by $\mathcal{L}(L^\infty(Y), L^\infty(Y))$ the Banach space of operators from $L^\infty(Y)$ to itself with the norm $\|\cdot\|_{\mathcal{L}(L^\infty(Y), L^\infty(Y))}$.

We state some fundamental properties of the sequence $(\mathcal{M}_{x,\rho}^n)_{n \geq N_{x,\rho}}$.

Proposition 2.1. *Let n be in a non-negative integer. Assume that ℓ_γ are positive linear forms (i.e. $f \geq 0$ implies $\ell_\gamma(f) \geq 0$ and for all $\gamma \in \Gamma$). We have:*

1. For all $f \in C(Z)$, we have

$$(\mathcal{M}_{x,\rho}^n(f))^* = \left(\frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} \ell_\gamma(f) \frac{\rho(\gamma)}{\phi(\gamma)} \right)^* = \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} \overline{\ell_{\gamma^{-1}}(f)} \frac{\rho(\gamma)}{\phi(\gamma)}.$$

2. $\|\mathcal{M}_{x,\rho}^n\|_{\mathcal{L}(C(Z), \mathcal{B}(\mathcal{H}))} \leq \|\mathcal{M}_{x,\rho}^n(\mathbf{1}_Z)\|_{\mathcal{B}(\mathcal{H})}$.

3. $\|\mathcal{M}_{x,\rho}^n(\mathbf{1}_Z)\|_{\mathcal{L}(L^\infty(Y), L^\infty(Y))} \leq \|\mathcal{M}_{x,\rho}^n(\mathbf{1}_Z)\mathbf{1}_Y\|_\infty$.

The proofs are easy and left to the reader.

3 Uniform boundedness

In this section a point x in X is fixed.

3.1 Useful functions

Let μ be a Γ -invariant conformal density of dimension α and let $L^\infty(\mu)$ be the Banach space of essentially bounded functions endowed with its usual norm denoted by $\|\cdot\|_\infty$. Let $\rho > 0$ and assume that there exists $N_{x,\rho}$ such that $|C_n(x,\rho)| > 0$ for all $n \geq N_{x,\rho}$. Consider the sequence of *positive functions* $F_{x,\rho}^n$ defined for all $n \geq N_{x,\rho}$ as:

$$F_{x,\rho}^n : v \in \partial X \mapsto \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} \frac{\exp\left(\frac{\alpha}{2}\beta_v(x,\gamma x)\right)}{\phi_x(\gamma)} \in \mathbb{R}^+, \quad (3.1)$$

where $\phi_x = \langle \pi_x(\gamma)\mathbf{1}_{\partial X}, \mathbf{1}_{\partial X} \rangle$ is the Harish-Chandra function defined in the introduction. Observe that $F_{x,\rho}^n$ is nothing but

$$F_{x,\rho}^n = \mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})\mathbf{1}_{\partial X}. \quad (3.2)$$

Consider also the sequence of positive functions $H_{x,\rho}^n$ defined for all $n \geq N_{x,\rho}$ as:

$$H_{x,\rho}^n : v \in \partial X \mapsto \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} \frac{\exp(\alpha\beta_v(x,\gamma x))}{\|\mu_{\gamma x}\|} \in \mathbb{R}^+. \quad (3.3)$$

We shall prove that $F_{x,\rho}^n$ and $H_{x,\rho}^n$ are uniformly bounded in the $L^\infty(\mu)$ norm. The fact that $F_{x,\rho}^n$ is uniformly bounded is the first step in the proof of Theorem A.

The proof of uniform boundedness for $(F_{x,\rho}^n)_{n \geq N_{x,\rho}}$ consists in two parts: we shall obtain sharp estimates of Busemann functions on shadows, then use Ahlfors regularity condition to estimate the Harish-Chandra function ϕ_x .

The method of the proof of the uniform boundedness of $(F_{x,\rho}^n)_{n \geq N_{x,\rho}}$ applies for showing the uniform boundedness of $(H_{x,\rho}^n)_{n \geq N_{x,\rho}}$ with suitable hypothesis.

3.2 Estimates for Busemann functions

These techniques using the hyperbolic inequality (2.1) extended to the whole space \bar{X} are very powerful. See for example [8] and [3] where these techniques are used.

Lemma 3.1. *Let $R > 0$ and let $v \in \partial X$. We have for all $y \in X$ and for all w in $\mathcal{O}_R(x, y)$:*

$$\min\{(w, v)_x, d(x, y)\} - R - \delta \leq (v, y)_x \leq (v, w)_x + R + \delta.$$

Proof. Recall that $d(x, y) - 2R \leq \beta_w(x, y) \leq d(x, y)$ for all $w \in \mathcal{O}_R(x, y)$. Hence, by equation (2.7), we have

$$d(x, y) - R \leq (w, y)_x \leq d(x, y). \quad (3.4)$$

On one hand, using first the hyperbolic property (2.1), then the observation (3.4) we have

$$\begin{aligned} (v, y)_x &\geq \min\{(v, w)_x, (w, y)_x\} - \delta \\ &\geq \min\{(w, v)_x, d(x, y)\} - R - \delta. \end{aligned}$$

On the other hand, using $(v, y)_x \leq d(x, y)$ we have

$$\begin{aligned} (v, w)_x &\geq \min\{(v, y)_x, (y, w)_x\} - \delta \\ &\geq \min\{(v, y)_x, d(x, y) - R\} - \delta \\ &\geq (v, y)_x - R - \delta. \end{aligned}$$

□

Proposition 3.1. *Let $R > 0$ and let n be a non-negative integer such that $n \geq \rho$ and let $v \in \partial X$. There exists q_v in X satisfying $d(x, q_v) = n + \rho$, such that for all y in X with $n - \rho \leq d(x, y) < n + \rho$ and for all w in $\mathcal{O}_R(x, y)$ we have*

$$\beta_v(x, y) \leq \beta_w(x, q_v) + 2(R + \rho) + 4\delta.$$

Proof. Define q_v as the point on the unique geodesic passing through v and x such that $d(x, q_v) = n + \rho$.

Since $(v, y)_x \leq d(x, y)$, the right hand side inequality of Lemma 3.1, the definition of q_v combined with the hyperbolic inequality (2.1) imply for all w in $\mathcal{O}_R(x, y)$ that

$$\begin{aligned} (v, y)_x &\leq \min\{(v, w)_x, d(x, y)\} + R + \delta \\ &\leq \min\{(v, w)_x, d(x, q_v)\} + R + \delta \\ &= \min\{(v, w)_x, (v, q_v)_x\} + R + \delta \\ &\leq (w, q_v)_x + R + 2\delta. \end{aligned}$$

Since y satisfies $n - \rho \leq d(x, y) < n + \rho$ and $d(x, q_v) = n + \rho$ the above inequality implies

$$\beta_v(x, y) \leq \beta_w(x, q_v) + 2(\rho + R) + 4\delta.$$

□

3.3 Ahlfors regularity and Harish-Chandra functions

Let Γ be a discrete group of isometries of a CAT(-1) space X and let μ be a Γ -invariant conformal density of dimension α . Fix a point x in X and define the function

$$\varphi_x : y \in X \mapsto \int_{\partial X} \exp\left(\frac{\alpha}{2}\beta_v(x, y)\right) d\mu_x(v). \quad (3.5)$$

Observe that ϕ_x is the restriction of φ_x to the orbit Γx .

Let \mathcal{Y} be a subset of X . We say that φ_x satisfies the *Harish-Chandra estimates on \mathcal{Y}* if there exist two polynomials Q_1 and Q_2 of *degree one* such that for all $y \in \mathcal{Y}$ we have $Q_1(d(x, y)) > 0$ and

$$Q_1(d(x, y)) \exp\left(-\frac{\alpha}{2}d(x, y)\right) \leq \varphi_x(y) \leq Q_2(d(x, y)) \exp\left(-\frac{\alpha}{2}d(x, y)\right). \quad (3.6)$$

Let $R > 0$ and such that for all x and y in X the shadows $\mathcal{O}_R(x, y)$ are not empty. Pick a point w_x^y in $\mathcal{O}_R(x, y)$. In the context of negatively curved manifolds, we can think about w_x^y as the ending point of the geodesic passing through x and y , oriented from x to y .

Lemma 3.2. *Let $v \in \partial X$ and $y \in X$. Let w_x^y be a point in $\mathcal{O}_R(x, y)$. Then, we have*

$$\exp\left(\frac{\alpha}{2}\beta_v(x, y)\right) \leq \exp(\alpha(\delta + R)) \exp\left(-\frac{\alpha}{2}d(x, y)\right) \frac{1}{d_x^\alpha(v, w_x^y)},$$

and

$$\exp\left(\frac{\alpha}{2}\beta_v(x, y)\right) \geq \exp(-\alpha(\delta + R)) \exp\left(-\frac{\alpha}{2}d(x, y)\right) \left(\min\left\{\frac{1}{d_x^\alpha(v, w_x^y)}, \exp(\alpha d(x, y))\right\}\right).$$

Proof. We prove the first inequality. The right hand side inequality of Lemma 3.1 leads to

$$(v, y)_x \leq (v, w_x^y)_x + R + \delta.$$

Combining this inequality with equation (2.7), we have

$$\exp\left(\frac{\alpha}{2}\beta_v(x, y)\right) \leq \exp(\alpha(\delta + R)) \exp\left(\alpha(v, w_x^y)_x\right) \exp\left(-\frac{\alpha}{2}d(x, y)\right).$$

The definition (2.8) of the visual metric completes the proof.

The left hand side of the inequality of Lemma 3.1 gives the other inequality. \square

Proposition 3.2. *Let μ be a Γ -invariant conformal density of dimension α . Assume that $(\Lambda_\Gamma, d_x, \mu_x)$ is Ahlfors α -regular. Then there exists $R_x > 0$ such that the function φ_x satisfies the Harish-Chandra estimates on $\Gamma x \setminus B_X(x, R_x)$.*

Moreover, if Γ is convex cocompact there exists $R_x > 0$ such that the function φ_x satisfies the Harish-Chandra estimates on $CH(\Lambda_\Gamma) \setminus B_X(x, R_x)$.

Proof. We first prove the right hand side inequality of (3.6) on $\mathcal{Y} = \Gamma x$. Let $\gamma \in \Gamma$, and consider a point $w_x^{\gamma x} \in \mathcal{O}_R(x, \gamma x) \cap \Lambda_\Gamma$. Consider the ball of ∂X of radius $\exp(-d(x, \gamma x))$ with respect to d_x centered at $w_x^{\gamma x}$ denoted by

$$B_\gamma := B\left(w_x^{\gamma x}, \exp(-d(x, \gamma x))\right).$$

$$\begin{aligned}\phi_x(\gamma) &= \int_{\partial X} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) d\mu_x(v) \\ &= \int_{B_\gamma} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) d\mu_x(v) + \int_{\partial X \setminus B_\gamma} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) d\mu_x(v).\end{aligned}$$

Ahlfors α -regularity implies for the first term that there exists $C > 0$ such that

$$\begin{aligned}\int_{B_\gamma} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) d\mu_x(v) &\leq \mu_x(B_\gamma) \exp\left(\frac{\alpha}{2}d(x, \gamma x)\right) \\ &\leq C \exp\left(-\frac{\alpha}{2}d(x, \gamma x)\right).\end{aligned}$$

The right hand side inequality of Lemma 3.2 implies that

$$\int_{\partial X \setminus B_\gamma} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) d\mu_x(v) \leq C_{\alpha, \delta, R} \exp\left(-\frac{\alpha}{2}d(x, \gamma x)\right) \int_{\partial X \setminus B_\gamma} \frac{1}{d_x^\alpha(v, w_x^{\gamma x})} d\mu_x(v),$$

for some positive constant $C_{\alpha, \delta, R} > 0$.

Write now

$$\begin{aligned}\int_{\partial X \setminus B_\gamma} \frac{1}{d_x^\alpha(v, w_x^{\gamma x})} d\mu_x(v) &= \int_{\mathbb{R}} \mu_x\left(\left\{v \in \partial X \mid \frac{1}{d_x^\alpha(v, w_x^{\gamma x})} > t\right\}\right) dt \\ &= \int_{1/D^\alpha}^{\exp \alpha d(x, \gamma x)} \mu_x\left(\left\{v \in \partial X \mid d_x(v, w_x^{\gamma x}) < \frac{1}{t^{1/\alpha}}\right\}\right) dt \\ &\leq \sum_{n=p}^N \mu_x\left(\left\{v \in \Lambda_\Gamma \mid d_x(v, w_x^{\gamma x}) < \frac{1}{n^{1/\alpha}}\right\}\right)\end{aligned}$$

where D denotes $\text{Diam}(\partial X)$, p the integer part of $\frac{1}{D^\alpha}$ and N then integer part of $\exp(\alpha d(x, \gamma x)) + 1$. Ahlfors regularity implies that there exists $C > 0$ such that

$$\mu_x\left(\left\{v \in \Lambda_\Gamma \mid d_x(v, w_x^{\gamma x}) < \frac{1}{n^{1/\alpha}}\right\}\right) \leq \frac{C}{n}.$$

Hence there exists a constant $C' > 0$ such that

$$\int_{\partial X \setminus B_\gamma} \frac{1}{d_x^\alpha(v, w_x^{\gamma x})} d\mu_x(v) \leq \alpha d(x, \gamma x) + C'.$$

Hence, we have

$$\int_{\partial X \setminus B_\gamma} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) d\mu_x(v) \leq C_{\alpha, \delta, R} \exp\left(-\frac{\alpha}{2}d(x, \gamma x)\right) (\alpha d(x, \gamma x) + C').$$

Therefore, we have found a polynomial of degree one such that φ_x satisfies the (right hand side) Harish-Chandra estimates on Γx . The left hand side of Harish-Chandra estimates on Γx is analogous by the second inequality of Lemma 3.2, but the constant term of the polynomial Q_1

might be non-positive. Hence the Harish-Chandra estimates hold only on $\Gamma x \setminus B_X(x, R_x)$ for some positive number R_x .

Assume that Γ is convex cocompact. We shall estimate φ_x on $CH(\Lambda_\Gamma)$. Let $y \in CH(\Lambda_\Gamma)$ and pick a fundamental domain $D_\Gamma \subset CH(\Lambda_\Gamma)$ relatively compact, and consider D'_Γ a relatively compact neighborhood of x which contains D_Γ . Then there exists $\gamma \in \Gamma$ such that $y \in \gamma D'_\Gamma$. Thanks to the cocycle identity (2.4) we have

$$\varphi_x(y) = \int_{\partial X} \exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) \exp\left(\frac{\alpha}{2}\beta_v(\gamma x, y)\right) d\mu_x(v).$$

Thanks to the properties of Busemann functions (2.3) and (2.5), observe that

$$\exp\left(-\frac{\alpha}{2}\text{Diam}(D'_\Gamma)\right)\phi_x(\gamma) \leq \varphi_x(y) \leq \phi_x(\gamma) \exp\left(\frac{\alpha}{2}\text{Diam}(D'_\Gamma)\right)$$

Observe also that

$$d(x, y) - \text{Diam}(D'_\Gamma) \leq d(\gamma x, x) \leq d(x, y) + \text{Diam}(D'_\Gamma)$$

for all $y \in X$ such that $d(x, y) \geq \text{Diam}(D'_\Gamma)$. Since φ_x satisfies the Harish-Chandra estimates on $\Gamma x \setminus B_X(x, R_x)$ we have the Harish-Chandra estimates of φ_x on $CH(\Lambda_\Gamma) \setminus B_X(x, R'_x)$ where $R'_x = \max\{R_x, \text{Diam}(D'_\Gamma)\}$ and the proof is done. \square

Remark 3.1. Notice that a slight modification of the first part of this proof gives a geometrical proof of the Harish-Chandra estimates of the Ξ -Harish-Chandra function in the context of rank one semisimple Lie groups (see [1] and [13]). It would be interesting to study an analog of Harish-Chandra estimates on $CH(\Lambda_\Gamma)$ for Harish-Chandra functions associated with geometrically finite groups with parabolic elements.

Remark 3.2. In [26, Theorem 2], B. Stratmann and S.-L. Velani prove, in the context of hyperbolic plane \mathbb{H}^n , the so-called Global Measure Formula for geometrically finite groups with parabolic elements. A conformal density of a geometrically finite group with parabolic elements is Ahlfors regular if and only if all the parabolic cusps have the same rank and that rank is equal to the critical exponent of the group. Hence such geometrically finite groups belong to the class \mathcal{C} since their spectrum are non-arithmetic.

3.4 Uniform boundedness

Proposition 3.3. Let μ be a Γ -invariant conformal density of dimension $\alpha(\Gamma)$ where Γ is in \mathcal{C} . Then there exists ρ and an integer N such that for all $n \geq N$, the sequence $F_{x,\rho}^n$ is uniformly bounded in the $L^\infty(\mu)$ norm.

Proof. If two sequences u_n and v_n of positive real numbers satisfy $\lim_{n \rightarrow \infty} u_n/v_n = 1$ we write $u_n \sim v_n$.

We shall prove first that $C_n(x, \rho)$ is not empty, at least for n large enough. For a non negative integer n , set $\Gamma_n(x) := \{\gamma \in \Gamma \mid d(x, \gamma x) < n\}$. Applying Theorem 2.1 to the function $\mathbf{1}_{\overline{X}} \otimes \mathbf{1}_{\overline{X}}$ we obtain as $n \rightarrow +\infty$

$$|\Gamma_n(x)| \sim \frac{\exp(\alpha n) \|\mu_x\|^2}{\alpha \|m_\Gamma\|},$$

and thus as $n \rightarrow +\infty$

$$|C_n(x, \rho)| \sim \frac{\exp(\alpha n)(2 \sinh(\alpha \rho)) \|\mu_x\|^2}{\alpha \|m_\Gamma\|}. \quad (3.7)$$

Hence, for all ρ there exists $N_{x, \rho}$ such that for all $n \geq N_{x, \rho}$ we have $|C_n(x, \rho)| > 0$.

There are two steps:

Step 1: Assume that x is in $CH(\Lambda_\Gamma)$. Then for all $\rho > 0$, there exists $N'_{x, \rho}$ and for all $n \geq N'_{x, \rho}$ the sequence $F_{x, \rho}^n$ is uniformly bounded with respect to the $L^\infty(\mu)$ norm.

First of all let R_x be a positive real number such that the Harish-Chandra estimates hold on $CH(\Lambda_\Gamma) \setminus R_x$. Let $\rho > 0$ and let $N_{x, \rho}$ be an integer such that for all $n \geq N_{x, \rho}$ we have $C_{n, \rho}(x) \subset CH(\Lambda_\Gamma) \setminus R_x$. Let v be in Λ_Γ , then Proposition 3.1 provides $q_v \in X$ with $d(x, q_v) = n + \rho$, such that for all $\gamma \in C_{n, \rho}(x)$:

$$\exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right) \leq \frac{\exp(\alpha(2(R + \rho) + 4\delta))}{\mu_x(\mathcal{O}_R(x, \gamma x))} \int_{\mathcal{O}_R(x, \gamma x)} \exp\left(\frac{\alpha}{2}\beta_w(x, q_v)\right) d\mu_x(w). \quad (3.8)$$

We set for the following computation $C_0 := \exp(\alpha(2(R + \rho) + 4\delta))$.

Therefore we have for $v \in \Lambda_\Gamma$:

$$\begin{aligned} F_{x, \rho}^n(v) &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right)}{\phi_x(\gamma)} \\ &\leq \frac{C_0}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\int_{\mathcal{O}_R(x, \gamma x)} \exp\left(\frac{\alpha}{2}\beta_w(x, q_v)\right) d\mu_x(w)}{\mu_x(\mathcal{O}_R(x, \gamma x)) \phi_x(\gamma)} \\ &\leq C \frac{C_0}{\exp(-\alpha(n - \rho)) |C_n(x, \rho)|} \left(\sup_{\gamma \in C_n(x, \rho)} \frac{1}{\phi_x(\gamma)} \right) \sum_{\gamma \in C_n(x, \rho)} \int_{\mathcal{O}_R(x, \gamma x)} \exp\left(\frac{\alpha}{2}\beta_w(x, q_v)\right) d\mu_x(w) \\ &\leq C \frac{C_0}{\exp(-\alpha(n - \rho)) |C_n(x, \rho)|} \left(\sup_{\gamma \in C_n(x, \rho)} \frac{1}{\phi_x(\gamma)} \right) (m \times \varphi_x(q_v)), \end{aligned}$$

where the last inequality follows from the fact that there exists an integer m such that for all $w \in \partial X$ the cardinality of $\{\gamma \in C_n(x, \rho) | w \in \mathcal{O}_R(x, \gamma x)\}$ is bounded by m .

The Sullivan Shadow lemma (for R large enough) implies that there exists $c' > 0$ such that for all n big enough we have :

$$\exp(-\alpha(n - \rho)) |C_n(x, \rho)| \geq c'.$$

Since the hypothesis guarantee the Ahlfors regularity of the limit set for the groups in the class \mathcal{C} ([6, 2.7.5 Théorème] for convex cocompact groups, the case of lattices is well known) then Proposition 3.2 implies that there exists $C' > 0$, such that for $q_v \in CH(\Lambda_\Gamma) \setminus R_x$ with $d(x, q_v) = n + \rho$ we have

$$\left(\sup_{\gamma \in C_n(x, \rho)} \frac{1}{\phi_x(\gamma)} \right) \varphi_x(q_v) \leq C'.$$

Hence for $x \in CH(\Lambda_\Gamma)$ and for all $\rho > 0$, there exists $K > 0$ and $N'_{x,\rho}$ such that for all $n \geq N'_{x,\rho}$ we have

$$\|F_{x,\rho}^n\|_\infty \leq K.$$

Step 2: Assume that x is in $X \setminus CH(\Lambda_\Gamma)$. There exist $\rho'_x > 0$ and an integer N'_{x,ρ'_x} such that for all $n \geq N'_{x,\rho'_x}$ the sequence $F_{x,\rho}^n$ is uniformly bounded with respect to the $L^\infty(\mu)$ norm.

Fix $\rho > 0$ and let x_0 be the projection of x in $CH(\Lambda_\Gamma)$ and set

$$\kappa := d(x, CH(\Lambda_\Gamma)) = d(x, x_0).$$

Using the relations (2.4), (2.6), (2.3), and (2.5) we obtain

$$\phi_x(\gamma) \geq \exp(\alpha\kappa)\phi_{x_0}(\gamma).$$

Observe that $C_n(x, \rho) \subset C_n(x_0, \rho + 2\kappa)$. We have:

$$\begin{aligned} F_{x,\rho}^n(v) &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\exp\left(\frac{\alpha}{2}\beta_v(x, \gamma x)\right)}{\phi_x(\gamma)} \\ &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\exp\left(\frac{\alpha}{2}\beta_v(x, x_0)\right) \exp\left(\frac{\alpha}{2}\beta_v(x_0, \gamma x_0)\right) \exp\left(\frac{\alpha}{2}\beta_v(\gamma x_0, \gamma x)\right)}{\phi_x(\gamma)} \\ &\leq \frac{\exp(2\alpha\kappa)}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x_0, \rho+2\kappa)} \frac{\exp\left(\frac{\alpha}{2}\beta_v(x_0, \gamma x_0)\right)}{\phi_{x_0}(\gamma)} \\ &= \left(\exp(2\alpha\kappa) \frac{|C_n(x_0, \rho+2\kappa)|}{|C_n(x, \rho)|} \right) F_{x_0, \rho+2\kappa}^n, \end{aligned}$$

where the third inequality comes from the relations (2.6) and (2.5). Since $|C_n(x_0, \rho+2\kappa)|/|C_n(x, \rho)|$ is bounded above by some constant depending on ρ and κ , we apply *Step 1* to $F_{x_0, \rho+2\kappa}^n$ with x_0 and $\rho + 2\kappa$ to complete the proof. \square

Remark 3.3. *If for any choice of an origin x the metric measure space $(\Lambda_\Gamma, d_x, \mu_x)$ is Ahlfors regular and if $CH(\Lambda_\Gamma) = X$, then the above proposition holds for all $\rho > 0$ independently of the choice of x . These conditions include the case of lattices in rank one semisimple Lie groups and fundamental groups of compact negatively curved manifolds.*

Remark 3.4. *For a proof of this uniform boundedness in the context of hyperbolic groups we refer to [14, Proposition 5.2].*

Proposition 3.4. *Let μ be Γ -invariant conformal density of dimension $\alpha(\Gamma)$ the critical exponent of the group and let Γ be a discrete group of isometries of a CAT(-1) space X with a non-arithmetic spectrum with a finite BMS measure. Assume that there exists $C > 0$ such that for all $y \in X$ we have $\|\mu_y\|/\|\mu_x\| \leq C$. For all $\rho > 0$, there exists an integer N such that for all $n \geq N$ the sequence of functions $H_{x,\rho}^n$ is uniformly bounded in the $L^\infty(\mu)$ norm.*

The proof for $H_{x,\rho}^n$ follows the same method and is left to the reader. Notice that this proof is easier because it does not deal with the Harish-Chandra estimates.

4 Analysis of matrix coefficients

In this section we fix x as an origin of X .

4.1 Notation

Let Γ be a discrete group of isometries of X and let μ be a Γ -invariant conformal density of dimension α . Let A be a subset of ∂X and $a > 0$ positive real number and define $A_x(a)$ the subset of ∂X as

$$A_x(a) = \{v \mid \inf_{w \in A} d_x(v, w) < \exp(-a)\}.$$

We will write $A(a)$ instead of $A_x(a)$. Recall that $\bigcap_{a>0} A(a) = \bar{A}$.

Let R a positive real number and define the cone of base A to be

$$C_R(x, A) := \{y \in X \mid \exists v \in A \text{ satisfying } [xv] \cap B(y, R) \neq \emptyset\},$$

where $[xv]$ represents the unique geodesic passing through x with the ending point $v \in \partial X$. In other words we have:

$$C_R(x, A) := \{y \in X \mid \mathcal{O}_R(x, y) \cap A \neq \emptyset\}. \quad (4.1)$$

Define $b_x(y)$ the function

$$b_x(y) : v \in \partial X \mapsto \exp\left(\frac{\alpha}{2}\beta_v(x, y)\right). \quad (4.2)$$

Notice that $\varphi_x(y) = \int_{\partial X} b_x(y)(v) d\mu_x(v)$.

4.2 Sharp estimates

Assume that φ_x satisfies Harish-Chandra estimates on \mathcal{Y} .

Lemma 4.1. *Let A be a Borel subset of ∂X and let $a > 0$. There exists a constant C_0 such that for all y in \mathcal{Y} satisfying $\mathcal{O}_R(x, y) \cap A(a) = \emptyset$, we have*

$$\frac{\langle b_x(y), \chi_A \rangle}{\varphi_x(y)} \leq \frac{C_0 \exp(a)}{d(x, y)}.$$

Proof. Let $y \in \mathcal{Y}$ and assume that $d(x, y) < a$. It is easy to check that

$$\frac{\langle b_x(y), \chi_A \rangle}{\varphi_x(y)} \leq \frac{\exp(a)}{d(x, y)}.$$

Now assume that $d(x, y) \geq a$.

If $v \in A(a)$ and $w \in \mathcal{O}_R(x, y)$, since $\mathcal{O}_R(x, y) \cap A(a) = \emptyset$ we have $d_x(v, w) > \exp(-a)$.

Using the first inequality Lemma 3.2 and the above observation we have for all $w \in \mathcal{O}_R(x, y)$:

$$\begin{aligned} \langle b_x(y), \chi_A \rangle &\leq \exp\left(-\frac{\alpha}{2}d(x, y)\right) \int_{A(a)} \frac{1}{d_x^\alpha(v, w)} \exp\left(\alpha(R + \delta)\right) d\mu_x(v) \\ &\leq \exp\left(\alpha(R + \delta + a)\right) \|\mu\|_x \exp\left(-\frac{\alpha}{2}d(x, y)\right) \\ &\leq \exp\left(\alpha(R + \delta + a)\right) \|\mu\|_x \frac{\varphi_x(y)}{Q_1(d(x, y))} \end{aligned}$$

where the last inequality comes from the left hand side of Harish-Chandra estimates on \mathcal{Y} . Since Q_1 is a polynomial of degree one, the proof is complete. \square

Now assume that φ_x satisfies the left hand side of Harish-Chandra estimates on $\mathcal{Y} = \Gamma x$.

Proposition 4.1. *Let $\psi_t \in l^1(\Gamma)$ such that $\|\psi_t\|_1 \leq 1$, and assume that*

$$\lim_{t \rightarrow +\infty} \psi_t(\gamma) = 0,$$

for all $\gamma \in \Gamma$. Then for every Borel subset $A \subset \partial X$ we have for all $a > 0$

$$\limsup_{t \rightarrow +\infty} \sum_{\gamma \in \Gamma} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq \limsup_{t \rightarrow +\infty} \sum_{\gamma \in \Gamma} \psi_t(\gamma) D_{\gamma x}(\chi_{C_R(x, A(a))}).$$

Proof. Let A be Borel subset of ∂X and let a be a positive number. Let t_0 be another positive real number. Consider the following partition of Γ :

$$\Gamma = \Gamma_1 \sqcup \Gamma_2 \sqcup \Gamma_3$$

with

$$\Gamma_1 = \{\gamma \in \Gamma \mid d(x, \gamma x) \leq t_0\}$$

and

$$\Gamma_2 = \{\gamma \in \Gamma \mid \mathcal{O}_R(x, \gamma x) \cap A(a) \neq \emptyset\} \cap \Gamma_1^c$$

and

$$\Gamma_3 = \{\gamma \in \Gamma \mid \mathcal{O}_R(x, \gamma x) \cap A(a) = \emptyset\} \cap \Gamma_1^c.$$

Since π_x is positive, we have that

$$\sum_{\Gamma_1} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq \sum_{\Gamma_1} \psi_t(\gamma).$$

Observe that

$$\gamma \in \Gamma_2 \Leftrightarrow D_{\gamma x}(\chi_{C_R(x, A(a))}) = 1.$$

Thus

$$\sum_{\gamma \in \Gamma_2} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq \sum_{\gamma \in \Gamma_2} \psi_t(\gamma) D_{\gamma x}(\chi_{C_R(x, A(a))}).$$

Observe that

$$\langle b_x(\gamma x), \chi_A \rangle = \langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle.$$

Since $\mathcal{Y} = \Gamma x$ we can apply Lemma 4.1 via the above observation and thus for all $t_0 > 0$:

$$\sum_{\Gamma_3} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq \left(\sum_{\Gamma} \psi_t(\gamma) \right) C_0 \frac{\exp(a)}{t_0}.$$

Then, since $\|\psi_t\|_1 \leq 1$, we obtain for all $t_0 > 0$

$$\sum_{\Gamma_3} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq C_0 \frac{\exp(a)}{t_0}.$$

It follows that for all $a > 0$ and for all $t > t_0$ we have

$$\sum_{\gamma \in \Gamma} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq \sum_{\Gamma_1} \psi_t(\gamma) + \sum_{\Gamma} \psi_t(\gamma) D_{\gamma x}(\chi_{C_R(x, A(a))}) + C_0 \frac{\exp(a)}{t_0}.$$

Since $\psi_t(\gamma) \rightarrow 0$ as $t \rightarrow +\infty$, we obtain by taking the lim sup in the above inequality

$$\limsup_{t \rightarrow +\infty} \sum_{\gamma \in \Gamma} \psi_t(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)} \leq \limsup_{t \rightarrow +\infty} \sum_{\gamma \in \Gamma} \psi_t(\gamma) D_{\gamma x}(\chi_{C_R(x, A(a))}) + C_0 \frac{\exp(a)}{t_0}.$$

This inequality holds for all $t_0 > 0$, so we take $t_0 \rightarrow +\infty$ and the proof is complete. \square

4.3 A consequence of Roblin's Theorem

If $A \subset \partial X$, we denote by ∂A its frontier. We need a consequence of Theorem 2.1 which counts the points of a Γ -orbit Γx in $C_R(x, A)$ when A is a Borel subset with $\mu_x(\partial A) = 0$. This is based on the regularity of the conformal densities. We recall that the topology of \bar{X} is compatible with the metric topology defined on ∂X by the visual metrics (see [6, §1.5]). If $O \subset \bar{X}$, we denote by \bar{O} its closure in \bar{X} .

First, observe the following:

Lemma 4.2. *Let A be a closed subset of ∂X . Then $\overline{C_R(x, A)} = C_R(x, A) \sqcup A$.*

Proof. It is easy to check that $C_R(x, A) \cup A \subset \overline{C_R(x, A)}$.

Now, assume that $v \in \overline{C_R(x, A)} \cap \partial X$ (otherwise there is nothing to do). We shall prove that $v \in A$. There exists a sequence of $y_n \in C_R(x, A)$ such that $y_n \rightarrow v$ with respect to the topology of \bar{X} . Since y_n is in $C_R(x, A)$, there exists $v_n \in A \cap \mathcal{O}_R(x, y_n)$ such that $(y_n, v_n)_x \geq d(x, y_n) - R$, for all integers n . Thus, we have

$$\begin{aligned} (v_n, v)_x &\geq \min \{ (v_n, y_n)_x, (y_n, v)_x \} - \delta \\ &\geq (y_n, v)_x - R - \delta. \end{aligned}$$

where the last inequality follows from $(y_n, v)_x \leq d(x, y_n)$. Since $y_n \rightarrow v$, it follows that $(y_n, v)_x$ goes to $+\infty$, and so $v_n \rightarrow v$ with respect to d_x . Since A is closed the proof is done. \square

Then we shall give a proof the following corollary:

Corollary 4.1. *(Extracted from [22, Théorème 4.1.1, Chapitre 4]) Let Γ be a discrete group of isometries of X with a non-arithmetic spectrum. Assume that Γ admits a finite BMS measure associated with a Γ -invariant conformal density μ of dimension $\alpha = \alpha(\Gamma)$. Let A, B be two Borel subsets such that $\mu_x(\partial A) = 0 = \mu_x(\partial B)$. Then for all $\rho > 0$ we have*

$$\limsup_{n \rightarrow +\infty} \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma^{-1}x} \otimes D_{\gamma x}(\chi_{C_R(x, A)} \otimes \chi_{C_R(x, B)}) \leq \frac{\mu_x(A) \mu_x(B)}{\|\mu\|_x^2}.$$

Proof. Let ρ be a positive real number. We have for all n large enough:

$$\begin{aligned} \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma^{-1}x} \otimes D_{\gamma x} &= \frac{\alpha \|m_\Gamma\| \exp(-\alpha(n + \rho))}{|C_n(x, \rho)| \alpha \|m_\Gamma\| \exp(-\alpha(n + \rho))} \sum_{\gamma \in \Gamma_{n+\rho}(x)} D_{\gamma^{-1}x} \otimes D_{\gamma x} \\ &- \frac{\alpha \|m_\Gamma\| \exp(-\alpha(n - \rho))}{|C_n(x, \rho)| \alpha \|m_\Gamma\| \exp(-\alpha(n - \rho))} \sum_{\gamma \in \Gamma_{n-\rho}(x)} D_{\gamma^{-1}x} \otimes D_{\gamma x}. \end{aligned}$$

The estimation (3.7) for annuli implies as $n \rightarrow +\infty$

$$|C_n(x, \rho)| \alpha \|m_\Gamma\| \exp(-\alpha(n + \rho)) \sim 2 \sinh(\alpha\rho) \exp(-\alpha\rho) \|\mu\|_x^2$$

and

$$|C_n(x, \rho)| \alpha \|m_\Gamma\| \exp(-\alpha(n - \rho)) \sim 2 \sinh(\alpha\rho) \exp(\alpha\rho) \|\mu\|_x^2.$$

Therefore Theorem 2.1 implies

$$\frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma^{-1}x} \otimes D_{\gamma x} \rightarrow \frac{1}{\|\mu_x\|^2} \mu_x \otimes \mu_x, \quad (4.3)$$

with respect to the weak* topology of $C(\overline{X} \times \overline{X})^*$.

Consider a Borel subset A of ∂X such that $\mu_x(\partial A) = 0$. We have $\mu_x(\overline{A}) = \mu_x(A)$. Thus, by Lemma 4.2 we obtain

$$\mu_x(\overline{C_R(x, \overline{A})}) = \mu_x(A).$$

Let $\epsilon > 0$. Since μ_x is a regular measure there exists an open subset O_A of \overline{X} such that

$$\overline{C_R(x, \overline{A})} \subset O_A \text{ and } \mu_x(O_A) \leq \mu_x(A) + \epsilon. \quad (4.4)$$

The subset $\overline{C_R(x, \overline{A})}$ is a compact subset of \overline{X} . By Urysohn's lemma, we can find a compactly supported function f_{O_A} such that

$$\chi_{\overline{C_R(x, \overline{A})}} \leq f_{O_A} \leq \chi_{O_A}.$$

Let B be another Borel subset such that $\mu_x(\partial B) = 0$. Let f_{O_B} be the continuous function given by the above construction. Notice that for all n we have:

$$\sum_{\gamma \in C_n(x, \rho)} D_{\gamma x} \otimes D_{\gamma^{-1}x} (\chi_{C_r(x, A)} \otimes \chi_{C_r(x, B)}) \leq \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x} \otimes D_{\gamma^{-1}x} (f_{O_A} \otimes f_{O_B}).$$

The consequence of Roblin's theorem (4.3) implies:

$$\begin{aligned}
& \limsup_{n \rightarrow \infty} \frac{\|\mu_x\|^2}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x} \otimes D_{\gamma^{-1}x} (\chi_{C_R(x, A)} \otimes \chi_{C_R(x, B)}) \\
& \leq \limsup_{n \rightarrow \infty} \frac{\|\mu_x\|^2}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x} \otimes D_{\gamma^{-1}x} (f_{O_A} \otimes f_{O_B}) \\
& = \lim_{n \rightarrow \infty} \frac{\|\mu_x\|^2}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x} \otimes D_{\gamma^{-1}x} (f_{O_A} \otimes f_{O_B}) \\
& = \int_{\partial X \times \partial X} (f_{O_A} \otimes f_{O_B}) d\mu_x \otimes d\mu_x \\
& \leq \mu_x(A)\mu_x(B) + \epsilon(\mu_x(A) + \mu_x(B)) + \epsilon^2,
\end{aligned}$$

where the last inequality follows from (4.4). The above inequality holds for all $\epsilon > 0$, so the proof is done. \square

4.4 An application of Roblin's equidistribution Theorem

Let $\rho > 0$, and let $N_{x, \rho}$ be an integer such that for all $n \geq N_{x, \rho}$ the sequence $\mathcal{M}_{x, \rho}^n$ is well defined. The purpose of this section is to use Corollary 4.1 for computing the limit of the sequence of operator-valued measures $(\mathcal{M}_{x, \rho}^n)_{n \geq N_{x, \rho}}$.

We assume that φ_x satisfies the left hand side of Harish-Chandra estimates on Γx .

Proposition 4.2. *Let $A, B, U \subset \partial X$ be Borel subsets such that $\mu_x(\partial A) = \mu_x(\partial B) = \mu_x(\partial U) = 0$, let $\widehat{U} = C_R(x, U) \cup U$ be a borel subset of \overline{X} . Then we have:*

$$\lim_{n \rightarrow +\infty} \langle \mathcal{M}_{x, \rho}^n(\chi_{\widehat{U}}) \chi_A, \chi_B \rangle = \frac{\mu_x(U \cap B)\mu_x(A)}{\|\mu_x\|^2}.$$

We need some lemmas to prepare the proof of this proposition.

Lemma 4.3. *Let $B, U \subset \partial X$ be Borel subsets such that $\mu_x(\partial B) = \mu_x(\partial U) = 0$, let $\widehat{U} = C_R(x, U) \cup U$ be a borel subset of \overline{X} satisfying $U \cap B(b) = \emptyset$, for some $b > 0$. Then we have*

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x, \rho}^n(\chi_{\widehat{U}}) \mathbf{1}_{\partial X}, \chi_B \rangle = 0.$$

Proof. For all $n \geq N_{x, \rho}$ we have:

$$\begin{aligned}
\langle \mathcal{M}_{x, \rho}^n(\chi_{\widehat{U}}) \mathbf{1}_{\partial X}, \chi_B \rangle &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x}(\chi_{\widehat{U}}) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_B \rangle}{\phi_x(\gamma)} \\
&= \sum_{\gamma \in \Gamma} \psi_n(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_B \rangle}{\phi_x(\gamma)}
\end{aligned}$$

where the inequality follows from the fact that π_x is positive, and where

$$\psi_n(\gamma) := \frac{1}{|C_n(x, \rho)|} \chi_{C_n(x, \rho)}(\gamma) D_{\gamma x}(\chi_{\widehat{U}}).$$

Proposition 4.1 implies that

$$\begin{aligned}
\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \mathbf{1}_{\partial X}, \chi_B \rangle &\leq \limsup_{n \rightarrow +\infty} \sum_{\gamma \in \Gamma} \psi_n(\gamma) D_{\gamma x}(\chi_{\widehat{U}}) D_{\gamma x}(\chi_{C_R(x,B(b))}) \\
&= \limsup_{n \rightarrow +\infty} \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} D_{\gamma x}(\chi_{\widehat{U} \cap C_R(x,B(b))}) \\
&\leq \limsup_{n \rightarrow +\infty} \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} D_{\gamma x}(\chi_{C_R(x,U \cap B(b))})
\end{aligned}$$

Note the general fact $\partial(A \cap B) \subset \partial A \cup \partial B$. Corollary 4.1 implies that

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \mathbf{1}_{\partial X}, \chi_B \rangle \leq \frac{\mu_x(U \cap B(b))}{\|\mu_x\|}.$$

By hypothesis $U \cap B(b) = \emptyset$ thus we have

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \mathbf{1}_{\partial X}, \chi_B \rangle = 0.$$

□

Lemma 4.4. *Let \widehat{U} be a Borel subset of \overline{X} and let A be a Borel subset of ∂X . We have*

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \chi_A, \mathbf{1}_{\partial X} \rangle \leq \limsup_{n \rightarrow +\infty} \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} D_{\gamma^{-1}x}(\chi_{\widehat{U}}) D_{\gamma x}(\chi_{C_R(x,A(a))}).$$

Proof. We have for all $n \geq N_{x,\rho}$:

$$\langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \chi_A, \mathbf{1}_{\partial X} \rangle = \sum_{\gamma \in \Gamma} \psi_n(\gamma) \frac{\langle \pi_x(\gamma) \mathbf{1}_{\partial X}, \chi_A \rangle}{\phi_x(\gamma)},$$

with

$$\psi_n(\gamma) = \frac{1}{|C_n(x,\rho)|} \chi_{C_n(x,\rho)}(\gamma) D_{\gamma^{-1}x}(\chi_{\widehat{U}}).$$

Applying Proposition 4.1 to ψ_n defined above we obtain that:

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \chi_A, \mathbf{1}_{\partial X} \rangle \leq \limsup_{n \rightarrow +\infty} \frac{1}{|C_n(x,\rho)|} \sum_{\gamma \in C_n(x,\rho)} D_{\gamma^{-1}x}(\chi_{\widehat{U}}) D_{\gamma x}(\chi_{C_R(x,A(a))}).$$

□

Lemma 4.5. *Let $A, B, U \subset \partial X$ be Borel subsets such that $\mu_x(\partial A) = \mu_x(\partial B) = \mu_x(\partial U) = 0$ and let $\widehat{U} = C_R(x, U) \cup U$ be a Borel subset of \overline{X} with $\mu_x(\partial U) = 0$.*

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}}) \chi_A, \chi_B \rangle \leq \frac{\mu_x(U \cap B) \mu_x(A)}{\|\mu_x\|^2}.$$

Proof. Let $a > 0$ and $b > 0$, and consider $A(a)$ and $B(b)$ such that $\mu_x(\partial B(b)) = 0 = \mu_x(\partial A(a))$. Let $B(b)^c = \partial X \setminus B(b)$. Set $\widehat{U}_1 = \widehat{U} \cap \overline{C_R(x, B(b))}$ and $\widehat{U}_2 = \widehat{U} \cap \overline{X} / \overline{C_R(x, B(b))}$. Let $U_1 = \widehat{U}_1 \cap \partial X$ and $U_2 = \widehat{U}_2 \cap \partial X$ and notice that $U_1 = U \cap \overline{B(b)}$ and $U_2 = U \cap \partial X / \overline{B(b)}$. Observe that $U_2 \cap B(b) = \emptyset$. Since $\widehat{U} = \widehat{U}_1 \sqcup \widehat{U}_2$ we have:

$$\begin{aligned} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle &= \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}_1})\chi_A, \chi_B \rangle + \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}_2})\chi_A, \chi_B \rangle \\ &\leq \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}_1})\chi_A, \mathbf{1}_{\partial X} \rangle + \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}_2})\mathbf{1}_{\partial X}, \chi_B \rangle. \end{aligned}$$

Applying Lemma 4.3 to the second term and Lemma 4.4 to the first term of the right hand side above inequality, we obtain:

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle \leq \limsup_{n \rightarrow +\infty} \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma^{-1}x}(\chi_{\widehat{U}_1}) D_{\gamma x}(\chi_{C_R(x, A(a))}).$$

Since $\mu_x(\partial U_1) = 0 = \mu_x(\partial A(a))$, Roblin's corollary 4.1 leads to

$$\limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle \leq \frac{\mu_x(U \cap B(b))\mu_x(A(a))}{\|\mu_x\|^2}.$$

Because the above inequality holds for all $a, b > 0$ but at most countably many values of a and b , we obtain the required inequality. \square

Proof of Proposition 4.2. By Lemma 4.5 it is sufficient to prove that

$$\liminf_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle = \frac{\mu_x(U \cap B)\mu_x(A)}{\|\mu_x\|^2}.$$

If W is a Borel subset of ∂X (or \overline{X}), we set $W^0 = W$ and $W^1 = \partial X \setminus W$ (or $W^1 = \overline{X} \setminus W$). We have

$$\begin{aligned} 1 &= \langle \mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})\mathbf{1}_{\partial X}, \mathbf{1}_{\partial X} \rangle \\ &= \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}^0} + \chi_{\widehat{U}^1})\chi_A, \chi_B \rangle \\ &= \sum_{i,j,k} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}^i})\chi_{A^j}, \chi_{B^k} \rangle \\ &= \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle + \sum_{i,j,k \neq (0,0,0)} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}^i})\chi_{A^j}, \chi_{B^k} \rangle. \end{aligned}$$

Then

$$\begin{aligned} 1 &\leq \liminf_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle + \sum_{i,j,k \neq (0,0,0)} \limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}^i})\chi_{A^j}, \chi_{B^k} \rangle \\ &\leq \limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle + \sum_{i,j,k \neq (0,0,0)} \limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}^i})\chi_{A^j}, \chi_{B^k} \rangle \\ &\leq \frac{1}{\|\mu_x\|^2} \sum_{i,j,k} \mu_x(U^i \cap B^k) \mu_x(A^j) \\ &= 1, \end{aligned}$$

where the last inequality comes from Lemma 4.5. Hence the inequalities of the above computation are equalities, so

$$\liminf_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle = \frac{\mu_x(U \cap B)\mu_x(A)}{\|\mu_x\|^2} = \limsup_{n \rightarrow +\infty} \langle \mathcal{M}_{x,\rho}^n(\chi_{\widehat{U}})\chi_A, \chi_B \rangle$$

and the proof is done. □

5 Conclusion

5.1 Standard facts about Borel subsets of measure zero frontier

Let us recall two standard facts of measure theory that we state as lemmas:

Lemma 5.1. *Assume that (Z, d, μ) is a metric measure space. Then the σ -algebra generated by Borel subset with measure zero frontier generates the Borel σ -algebra.*

Let χ_A be the characteristic function of a Borel subset A of ∂X . We state another useful lemma (see [3, Appendix B, Lemma B.2 (1)] for a proof):

Lemma 5.2. *Assume that (Z, d, μ) is a metric measure space where μ is Radon measure. Then the closure of the subspace spanned by the characteristic functions of Borel subset having zero measure frontier is*

$$\overline{\text{Span}\{\chi_A \text{ such that } \mu(\partial A) = 0\}}^{L^2} = L^2(Z, \mu).$$

5.2 Proofs

Proof of Theorem A. Let μ be a Γ -invariant conformal density of dimension $\alpha(\Gamma)$, where Γ in \mathcal{C} . Since for all $x \in X$, the metric measure space $(\Lambda_\Gamma, d_x, \mu_x)$ is Ahlfors α -regular Proposition 3.2 ensures that the Harish-Chandra estimates hold on Γx . Hence Proposition 3.3 and 4.2 are available. The sequence $\mathcal{M}_{x,\rho}^n$ is defined for $n \geq N_{x,\rho}$ for some integer $N_{x,\rho}$. There are two steps.

Step 1: $(\mathcal{M}_{x,\rho}^n)_{n \geq N_{x,\rho}}$ **is uniformly bounded.** First of all, observe that $\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})$ is self-adjoint (see Proposition 2.1 (1)). Note that $\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})$ preserves $L^\infty(\partial X, \mu_x)$, and by duality it preserves also $L^1(\partial X, \mu_x)$.

Combining Proposition 3.3 with the fact that $\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})\mathbf{1}_{\partial X} = F_{x,\rho}^n$, we have that the sequence $(\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}}))_{n \geq N_{x,\rho}}$, with $\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})$ viewed as operators from $L^\infty(\partial X, \mu_x)$ to $L^\infty(\partial X, \mu)$, is uniformly bounded. Riesz-Thorin interpolation theorem implies the sequence $(\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}}))_{n \geq N_{x,\rho}}$, with $\mathcal{M}_{x,\rho}^n(\mathbf{1}_{\overline{X}})$ viewed as operators in $\mathcal{B}(L^2(\partial X, \mu_x))$, is uniformly bounded. Then Proposition 2.1 (2) completes **Step 1**.

Step 2: computation of the limit of $(\mathcal{M}_{x,\rho}^n)_{n \geq N_{x,\rho}}$. By the Banach-Alaoglu theorem, **Step 1** implies that $(\mathcal{M}_{x,\rho}^n)_{n \geq N_{x,\rho}}$ has accumulation points. Let \mathcal{M}_x^∞ be an accumulation point of $(\mathcal{M}_{x,\rho}^n)_{n \geq N_{x,\rho}}$ with respect to the weak* topology of $\mathcal{L}(C(\overline{X}), \mathcal{B}(L^2(\partial X, \mu_x)))$. Let $\widehat{U} = C_R(x, U) \cup U$ be Borel subset of \overline{X} with U be a Borel subset of ∂X such that $\mu_x(\partial U) = 0$. It follows from

Proposition 4.2 and from the definition (1.4) of \mathcal{M}_x that for all Borel subsets $A, B \subset \partial X$ with $\mu_x(\partial A) = \mu_x(\partial B) = 0$ we have that

$$\langle \mathcal{M}_x^\infty(\chi_{\widehat{U}})\chi_A, \chi_B \rangle = \frac{\mu_x(U \cap B)\mu_x(A)}{\|\mu_x\|^2} = \langle \mathcal{M}_x(\chi_{\widehat{U}})\chi_A, \chi_B \rangle.$$

The above equality holds for all open balls $B_X(x, r)$ of X , namely

$$\langle \mathcal{M}_x^\infty(\chi_{B_X(x, r)})\chi_A, \chi_B \rangle = \frac{\mu_x(B_X(x, r) \cap B)\mu_x(A)}{\|\mu_x\|^2} = \langle \mathcal{M}_x(\chi_{B_X(x, r)})\chi_A, \chi_B \rangle = 0.$$

Since the open balls $\{B_X(x, r), x \in X, r > 0\}$ of X together with the subsets $\widehat{U} = C_R(x, U) \cup U$, with U Borel subsets of ∂X such that $\mu_x(\partial U) = 0$ generate the Borel σ -algebra of \overline{X} , Carathéodory's extension theorem implies that for all $f \in C(\overline{X})$ and for all Borel subsets $A, B \subset \partial X$ satisfying $\mu_x(\partial A) = \mu_x(\partial B) = 0$ we have

$$\langle \mathcal{M}_x^\infty(f)\chi_A, \chi_B \rangle = \langle \mathcal{M}_x(f)\chi_A, \chi_B \rangle.$$

Lemma 5.2 combined with the above equality imply that the operators \mathcal{M}_x^∞ and \mathcal{M}_x regarded as functionals of $(C(\overline{X}) \widehat{\otimes} L^2(\partial X, \mu_x) \widehat{\otimes} L^2(\partial X, \mu_x))^*$ (see (2.13)) are equal on a dense subset of $C(\overline{X}) \widehat{\otimes} L^2(\partial X, \mu_x) \widehat{\otimes} L^2(\partial X, \mu_x)$. We deduce that \mathcal{M}_x is the unique accumulation point of the sequence $(\mathcal{M}_{x, \rho}^n)_{n \geq N_{x, \rho}}$. \square

Proof of Corollary B. Apply the definition of weak* convergence to $\mathbf{1}_{\overline{X}} \otimes \xi \otimes \eta$ for all $\xi, \eta \in L^2(\partial X, \mu_x)$, and observe that $\|\mu\|_x^2 \mathcal{M}_x(\mathbf{1}_{\overline{X}})$ is the orthogonal projection onto the space of constant functions. \square

Proof of Corollary C. Since $(\pi_x)_{x \in X}$ are unitarily equivalent, it suffices to prove irreducibility for some π_x with x in X . Theorem A shows that the vector $\mathbf{1}_{\partial X}$ is cyclic for the representation π_x . Moreover, Corollary B shows that the orthogonal projection onto the space of constant functions is in the von Neumann algebra associated with π_x . Then, a standard argument (see for example [14, Lemma 6.1]) completes the proof. \square

Remark 5.1. *The hypothesis:* Γ is convex cocompact or a lattice in a rank one semisimple Lie group guarantees the Ahlfors regularity of the limit set, that implies the Harish-Chandra estimates of φ_x on $CH(\Lambda_\Gamma)$ and on Γx . In other words, the proof of irreducibility of boundary representations for a geometrically finite group with a non-arithmetic spectrum is reduced, by this approach, to the Harish-Chandra estimates of φ_x for each $x \in X$ on $CH(\Lambda_\Gamma) \setminus B_X(x, R_x)$ and on the orbit $\Gamma x \setminus B_X(x, R_x)$ for some $R_x > 0$. And this approach should apply to some geometrically finite groups which are neither convex cocompact and nor lattices.

6 Some remarks about equidistribution results

6.1 Dirac-Weierstrass family

Let Γ be a discrete group of isometries of X . Consider $(d_x)_{x \in X}$ a visual metric on ∂X , and let μ be a Γ -invariant conformal density of dimension α . We fix $x \in X$ and we follow [17, Chapter 2, §2.1, p 46], and adapt the definition of a Dirac-Weierstrass family to the density μ :

Definition 6.1. A Dirac-Weierstrass family $(K(y, \cdot))_{y \in X}$ with respect to μ_x , is a continuous map $K : (y, v) \in X \times \partial X \mapsto K(y, v) \in \mathbb{R}$ satisfying

1. $K(y, v) \geq 0$ for all $v \in \partial X$ and $y \in X$,
2. $\int_{\partial X} K(y, v) d\mu_x(v) = 1$ for all $y \in X$,
3. for all $v_0 \in \partial X$ and for all $r_0 > 0$ we have:

$$\int_{\partial X \setminus B(v_0, r_0)} K(y, v) d\mu_x(v) \rightarrow 0 \quad \text{as } y \rightarrow v_0.$$

A Dirac-Weierstrass family yields an integral operator \mathcal{K} :

$$\mathcal{K} : f \in L^1(\partial X, \mu_x) \mapsto \mathcal{K}f \in C(X)$$

defined as :

$$\mathcal{K}f : y \in X \mapsto \int_{\partial X} f(v) K(y, v) d\mu_x(v) \in \mathbb{C}.$$

6.2 Continuity

Let f be a function on ∂X . We define the function $\bar{\mathcal{K}}f$ on \bar{X} as the following:

$$\bar{\mathcal{K}}f : y \in \bar{X} \mapsto \bar{\mathcal{K}}f(y) = \begin{cases} \mathcal{K}f(y) & \text{if } y \in X \\ f(y) & \text{if } y \in \partial X \end{cases} \quad (6.1)$$

Thus, $\bar{\mathcal{K}}$ is an operator which assigns a function defined on \bar{X} to a function defined on ∂X .

Proposition 6.1. *If f is a continuous functions on ∂X , the function $\bar{\mathcal{K}}(f)$ is a continuous function on \bar{X} .*

Proof. Observe first that since K is a continuous function on X the function $\bar{\mathcal{K}}f$ is on X . Let v_0 be in ∂X and let $\epsilon > 0$. Since f is continuous, there exists $r > 0$ such that

$$|f(v_0) - f(v)| < \frac{\epsilon}{2},$$

whenever $v \in B(v_0, r)$. Besides, by (3) in Definition 6.1, there exists a neighborhood V of v_0 such that for all $y \in V$ we have:

$$\int_{\partial X \setminus B(v_0, r)} K(y, v) d\mu(v) \leq \frac{\epsilon}{4\|f\|_\infty}.$$

We have for all $y \in V$:

$$\begin{aligned} |\bar{\mathcal{K}}f(v_0) - \bar{\mathcal{K}}f(y)| &\leq \int_{B(v_0, r)} |f(v_0) - f(v)| K(y, v) d\mu_x(v) + \int_{\partial X \setminus B(v_0, r)} |f(v_0) - f(v)| K(y, v) d\mu_x(v) \\ &\leq \frac{\epsilon}{2} + 2\|f\|_\infty \int_{\partial X \setminus B(v_0, r)} K(y, v) d\mu_x(v) \\ &\leq \epsilon. \end{aligned}$$

Hence, $\bar{\mathcal{K}}f$ is a continuous function on \bar{X} . □

6.3 Examples of Dirac-Weierstrass family

Let $R > 0$, and consider for each $y \in X$ a point $w_x^y \in \mathcal{O}_R(x, y)$. We start by a lemma:

Lemma 6.1. *Let v_0 be in ∂X . Then $d_x(v_0, w_x^y) \rightarrow 0$ as $y \rightarrow v_0$.*

Proof. Let y_n be a sequence of points of X such that $y_n \rightarrow v_0$. Apply the right hand side inequality of Lemma 3.1 to get

$$(v_0, w_x^{y_n})_x \geq (v_0, y_n)_x - R - \delta.$$

Since $y_n \rightarrow v_0$, we have $(v_0, y_n)_x$ goes to infinity, and thus $d_x(v_0, w_x^y) \rightarrow 0$ as $y \rightarrow v_0$. \square

Proposition 6.2. *Assume that there exists a polynomial Q_1 (at least of degree 1) such that for all $y \in X$ with $d(x, y)$ large enough $Q_1(d(x, y)) > 0$ and*

$$Q_1(d(x, y)) \exp\left(-\frac{\alpha}{2}d(x, y)\right) \leq P_0 \mathbf{1}_{\partial X}(y).$$

Then

$$\left(\frac{P(y, \cdot)^{1/2}}{P_0 \mathbf{1}_{\partial X}(y)}\right)_{y \in X}$$

is a Dirac-Weierstrass family.

Proof. Let $B(v_0, r_0)$ the ball of radius r_0 at v_0 in ∂X with respect to d_x .

Let $\epsilon > 0$. Since Q_1 is a polynomial at least of degree one, there exists $R' > 0$ such that for all y satisfying $d(x, y) > R'$ we have

$$\frac{C_{r_0, \alpha, \delta} \|\mu_x\|}{Q_1(d(x, y))} < \epsilon$$

where $C_{r_0, \alpha, \delta} = 2^\alpha \exp(\alpha(\delta + R))/r_0^\alpha$ is a positive constant.

Lemma 6.1 yields a neighborhood V of v_0 such that $d_x(v_0, w_x^y) \leq r_0/2$ for all $y \in V$. We have for all v in $\partial X \setminus B(v_0, r_0)$:

$$\begin{aligned} d_x(v, w_x^y) &\geq d_x(v, v_0) - d_x(v_0, w_x^y) \\ &\geq r_0 - d_x(v_0, w_x^y) \\ &\geq \frac{r_0}{2}. \end{aligned}$$

We set $V_{R'} = V \cap X \setminus B_X(x, R')$. Combining Lemma 3.2 with the above inequality we obtain for all $y \in V_{R'}$:

$$\begin{aligned} \int_{\partial X \setminus B(v_0, r_0)} \frac{P^{\frac{1}{2}}(y, v)}{P_0 \mathbf{1}_{\partial X}(y)} d\mu_x(v) &\leq \int_{\partial X \setminus B(v_0, r_0)} \frac{\exp(\alpha(\delta + R)) \exp(-\frac{\alpha}{2}d(x, y))}{d_x^\alpha(v, w_x^y) (P_0 \mathbf{1}_{\partial X}(y))} d\mu_x(v) \\ &\leq C_{r_0, \alpha, \delta} \int_{\partial X \setminus B(v_0, r_0)} \frac{\exp(-\frac{\alpha}{2}d(x, y))}{Q_1(d(x, y)) \exp(-\frac{\alpha}{2}d(x, y))} d\mu_x(v) \\ &= C_{r_0, \alpha, \delta} \int_{\partial X \setminus B(v_0, r_0)} \frac{1}{Q_1(d(x, y))} d\mu_x(v) \\ &\leq \frac{C_{r_0, \alpha, \delta} \mu_x(\partial X)}{Q_1(d(x, y))} \\ &\leq \epsilon. \end{aligned}$$

It follows that

$$\int_{\partial X \setminus B(v_0, r_0)} \frac{P^{\frac{1}{2}}(y, v)}{P_0 \mathbf{1}_{\partial X}(y)} d\mu_x(v) \rightarrow 0 \text{ as } y \rightarrow v_0.$$

□

Besides, the same method proves the following proposition:

Proposition 6.3. *Assume that there exists a constant $C > 0$ such that*

$$\frac{\|\mu_x\|}{\|\mu_y\|} \leq C,$$

for all $y \in X$. The normalized Poisson kernel

$$\left(\frac{P(y, \cdot)}{\|\mu_y\|} \right)_{y \in X}$$

is a Dirac-Weierstrass family.

6.4 Equidistribution theorems extended to $(L^1)^*$

Theorem 2.1 of T. Roblin has for immediate consequence:

Theorem 6.2. *(T. Roblin) Let Γ be a discrete subgroup of isometries of X with a non-arithmetic spectrum. Assume that Γ admits a finite BMS measure associated to a Γ -invariant conformal density μ of dimension $\alpha = \alpha(\Gamma)$. Then for each $x \in X$ and for all $\rho > 0$ we have as n goes to infinity:*

$$\frac{1}{|C_n(x, \rho)|} \sum_{C_n(x, \rho)} D_{\gamma^{-1}x} \rightarrow \frac{\mu_x}{\|\mu_x\|}$$

with respect to the weak* topology of $C(\overline{X})^*$.

We view Theorem D as new new equidistribution theorem, where the weak* convergence is not on the dual of the space of continuous functions but rather on the dual of space of L^1 functions on the boundary.

Proof of Theorem D. Let x in X and $\rho > 0$, and consider $N_{x, \rho}$ such that $n \geq N_{x, \rho}$ implies $|C_n(x, \rho)| > 0$. We give a proof for the densities $(\mu_x)_{x \in X}$. For all $n \geq N_{x, \rho}$, we denote by $\lambda_{x, \rho}^n$ the following measure

$$\lambda_{x, \rho}^n = \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\mu_{\gamma x}}{\|\mu_{\gamma x}\|}.$$

Step 1: the sequence of measures $(\lambda_{x, \rho}^n)_{n \geq N_{x, \rho}}$ is uniformly bounded.

Since the dual space of $L^1(\partial X, \mu_x)$ is $L^\infty(\partial X, \mu)$ we have for $n \geq N_{x\rho}$:

$$\begin{aligned} \|H_{x,\rho}^n\|_\infty &= \sup_{\|f\|_1 \leq 1} \left\{ \left| \int_{\partial X} H_{x,\rho}^n(v) f(v) d\mu_x(v) \right| \right\} \\ &= \sup_{\|f\|_1 \leq 1} \left\{ \left| \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\mu_{\gamma x}}{\|\mu_x\|} (f) \right| \right\} \\ &= \|\lambda_{x,\rho}^n\|_{(L^1)^*}. \end{aligned}$$

Hence Proposition 3.4 completes **Step 1**.

Step 2: computation of the limit of $(\lambda_n^x)_{n \geq N_{x,\rho}}$.

By Banach-Alaoglu's theorem, $(\lambda_{x,\rho}^n)_{n \geq N_{x,\rho}}$ has accumulation points. Denote by λ_x^∞ such accumulation point. Let $f \in C(\partial X)$, Proposition 6.3 combined with Proposition 6.1 define $\overline{\mathcal{P}}f$ as a continuous function on \overline{X} (as in (6.2) in Subsection 6.1), where \mathcal{P} is associated with the normalized Poisson kernel defined as in Proposition 6.3. We have for all $n \geq N_{x,\rho}$:

$$\begin{aligned} \lambda_{x,\rho}^n(f) &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\mu_{\gamma x}(f)}{\|\mu_{\gamma x}\|} \\ &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \mathcal{P}(f)(\gamma x) \\ &= \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} D_{\gamma x}(\overline{\mathcal{P}}(f)). \end{aligned}$$

Applying Roblin's theorem 6.2 by taking the limit in the above inequality, we obtain for all $f \in C(\partial X)$

$$\lambda_x^\infty(f) = \mu_x(\overline{\mathcal{P}}(f)) = \frac{\mu_x(f)}{\|\mu_x\|}.$$

Since $C(\partial X)$ is dense $L^1(\partial X, \mu_x)$ with respect to the L^1 norm, we deduce that $(\lambda_{x,\rho}^n)_{n \geq N_{x,\rho}}$ has only one accumulation point which is μ_x , and the proof is done.

The proof concerning $(\nu_x)_{x \in X}$ follows the same method, and uses $\varphi_x = P_0$ in order to have available Proposition 6.2 for Γ in \mathcal{C} . Indeed, since the lower bound of the Harish-Chandra estimates holds a priori only on $CH(\Lambda_\Gamma) \setminus B_X(x, R_x)$ we rather use Proposition 6.2 with $\overline{CH(\Lambda_\Gamma)} = CH(\Lambda_\Gamma) \cup \Lambda_\Gamma$ instead of $\overline{X} = X \cup \partial X$. If f is a continuous function on Λ_Γ , the function $\overline{\mathcal{P}}_0 f$ on $\overline{CH(\Lambda_\Gamma)}$ defined as

$$\overline{\mathcal{P}}_0 f : y \in \overline{CH(\Lambda_\Gamma)} \mapsto \overline{\mathcal{P}}_0 f(y) = \begin{cases} \mathcal{P}_0 f(y) & \text{if } y \in CH(\Lambda_\Gamma) \\ f(y) & \text{if } y \in \Lambda_\Gamma \end{cases} \quad (6.2)$$

is continuous on $\overline{CH(\Lambda_\Gamma)}$. □

Remark 6.1. *We may ask if an analogous theorem of Theorem 2.1 dealing with μ_x instead of the Dirac mass holds (assuming $\|\mu_x\| = 1$ for simplicity)? More precisely do we have:*

$$\frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \mu_{\gamma^{-1}x} \otimes \mu_{\gamma x} \rightharpoonup \mu_x \otimes \mu_x$$

with respect to the weak* convergence of $L^1(\partial X \times \partial X, \mu_x \otimes \mu_x)^*$ (for some ρ)? The answer is negative because a duality argument combined with Banach-Steinhaus theorem would imply that the sequence of functions

$$G_n : (v, w) \mapsto \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \exp(\alpha\beta_v(x, \gamma^{-1}x)) \exp(\alpha\beta_w(x, \gamma x))$$

is uniformly bounded with respect to the $L^\infty(\mu)$ norm. It is easy to see that this is impossible by evaluating G_n at $(v, w) \in \mathcal{O}_R(x, \gamma^{-1}x) \times \mathcal{O}_R(x, \gamma x)$ for some $\gamma \in C_n(x, \rho)$. We obtain the same answer to the same question dealing with ν_x by considering the sequence of functions

$$(v, w) \mapsto \frac{1}{|C_n(x, \rho)|} \sum_{\gamma \in C_n(x, \rho)} \frac{\exp(\frac{\alpha}{2}\beta_v(x, \gamma^{-1}x)) \exp(\frac{\alpha}{2}\beta_w(x, \gamma x))}{\phi_x^2(\gamma)}.$$

References

- [1] J.-P. Anker, *La forme exacte de l'estimation fondamentale de Harish-Chandra. (French) [The exact form of Harish-Chandra's fundamental estimate]* C. R. Acad. Sci. Paris Sr. I Math. 305 (1987), no. 9, 371-374.
- [2] M. Babillot, *Points entiers et groupes discrets: de l'analyse aux systèmes dynamiques.* Panor. Synthèses, 13, Rigidité, groupe fondamental et dynamique, 1-119, Soc. Math. France, Paris, 2002.
- [3] U. Bader, R. Muchnik, *Boundary unitary representations, irreducibility and rigidity.* J. Mod. Dyn. 5 (2011), no. 1, 49-69.
- [4] M.-E. B. Bekka, M. Cowling. *Some irreducible unitary representations of $G(K)$ for a simple algebraic group G over an algebraic number field K .* Math. Zeitschrift, 241(4):731-741, 2002.
- [5] M.R. Bridson; A. Haefliger, *Metric spaces of non-positive curvature.* Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319. Springer-Verlag, Berlin, 1999.
- [6] M. Bourdon, *Structure conforme et flot géodésique d'un CAT(-1)-espace,* L'enseignement mathématique (1995).
- [7] M. Burger, S. Mozes, *CAT(-1)-spaces, divergence groups and their commensurators.* J. Amer. Math. Soc. 9 (1996), no. 1, 57-93.
- [8] C. Connell; R. Muchnik, *Harmonicity of quasiconformal measures and Poisson boundaries of hyperbolic spaces.* Geom. Funct. Anal. 17 (2007), no. 3, 707-769.
- [9] M. Cowling, T. Steger, *The irreducibility of restrictions of unitary representations of lattices.* J. für die reine und Angew. Math., 420:85-98, 1991.
- [10] F. Dal'bo, *Remarques sur le spectre des longueurs d'une surface et comptages. (French) [Remarks on the length spectrum of a surface, and counting]* Bol. Soc. Brasil. Mat. (N.S.) 30 (1999), no. 2, 199-221.

- [11] A. Figà-Talamanca; M.-A. Picardello, *Harmonic analysis on free groups*. Lecture Notes in Pure and Applied Mathematics, 87. Marcel Dekker, Inc., New York, 1983.
- [12] A. Figà-Talamanca; T. Steger, *Harmonic analysis for anisotropic random walks on homogeneous trees*. Mem. Amer. Math. Soc. 110 (1994), no. 531, xii+68 pp.
- [13] R. Gangolli, V.S. Varadarajan, *Harmonic Analysis of Spherical Functions on Real Reductive Groups*, Springer-Verlag, New-York, 1988.
- [14] L. Garncarek, *Boundary representations of hyperbolic groups*, arXiv:1404.0903, 2014.
- [15] U. Haagerup, *An example of a nonnuclear C^* -algebra which has the metric approximation property*, Invent. Math. 50 (1978/79), no. 3, 279-293.
- [16] P. Jolissaint, *Rapidly decreasing functions in reduced C^* -algebras of groups*, Trans. Amer. Math. Soc. 317 (1990), no. 1, 167-196.
- [17] J. Jorgenson, S. Lang, *The Heat Kernel and Theta Inversion on $SL_2(\mathbb{C})$* , Springer Monographs in Mathematics. Springer, New York, 2008.
- [18] A. Gorodnik; A. Nevo, *The ergodic theory of lattice subgroups*. Annals of Mathematics Studies, 172. Princeton University Press, Princeton, NJ, 2010.
- [19] G. Kuhn, T. Steger, *More irreducible boundary representations of free groups*. Duke Math. J. 82 (1996), no. 2, 381-436.
- [20] G.-A. Margulis, *On some aspects of the theory of Anosov systems*. Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2004, With a survey by Richard Sharp: Periodic orbits of hyperbolic flows, Translated from the Russian by Valentina Vladimirovna Szulikowska.
- [21] S.-J. Patterson, *The limit set of a fuchsian group*, Acta mathematica 136, 241-273 (1976)
- [22] T. Roblin, *Ergodicité et équidistribution en courbure négative*, Mémoires de la SMF 95 (2003).
- [23] Ryan, Raymond A. *Introduction to tensor products of Banach spaces*. Springer Monographs in Mathematics. Springer-Verlag London, Ltd., London, 2002.
- [24] P. Sjögren, *Convergence for the square root of the Poisson kernel*, Pacific J. Math., 131, 2, p. 361–391, (1988).
- [25] E. Stein, *M. Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. With the assistance of Timothy S. Murphy. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [26] B. Stratmann ; S.-L. Velani, *The Patterson measure for geometrically finite groups with parabolic elements, new and old*. Proc. London Math. Soc. (3) 71 (1995), no. 1, 197-220.
- [27] D. Sullivan, *The density at infinity of a discrete group of hyperbolic motions*, Publications mathématiques de l’IHES 50 (1979), 171-202.