

SURVEY

An introduction to pressure metrics for higher Teichmüller spaces

MARTIN BRIDGEMAN[†], RICHARD CANARY[‡] and ANDRÉS SAMBARINO[§]

[†] *Boston College, Chestnut Hill, MA 02467, USA*

[‡] *University of Michigan, Ann Arbor, MI 41809, USA*

[§] *Université Pierre et Marie Curie (Paris VI), 75005 Paris, France*

(Received 14 July 2015 and accepted in revised form 7 September 2016)

Abstract. We discuss how one uses the thermodynamic formalism to produce metrics on higher Teichmüller spaces. Our higher Teichmüller spaces will be spaces of Anosov representations of a word-hyperbolic group into a semi-simple Lie group. We begin by discussing our construction in the classical setting of the Teichmüller space of a closed orientable surface of genus at least 2, then we explain the construction for Hitchin components and finally we treat the general case. This paper surveys results of Bridgeman, Canary, Labourie and Sambarino, *The pressure metric for Anosov representations*, and discusses questions and open problems which arise.

Contents

1	Introduction	1
2	The thermodynamic formalism	2
3	Basic strategy	6
4	The pressure metric for Teichmüller space	7
5	The pressure metric on the Hitchin component	11
6	Generalizations and consequences	24
7	Open problems	29
	Acknowledgements	32
	References	32

1. *Introduction*

We discuss how one uses the thermodynamic formalism to produce metrics on higher Teichmüller spaces. Our higher Teichmüller spaces will be spaces of Anosov representations of a word hyperbolic group into a semi-simple Lie group. To each such

representation we associate an Anosov flow encoding eigenvalue information, and the thermodynamic formalism gives us a way to measure the difference between two such flows. This difference gives rise to an analytic semi-norm, which in many cases turns out to be a Riemannian metric, called the *pressure metric*. This paper surveys results of Bridgeman *et al* [17] and discusses questions and open problems which arise.

We begin by discussing our construction in the classical setting of the Teichmüller space of a closed orientable surface of genus at least 2. In this setting, our construction agrees with Thurston's Riemannian metric, as re-interpreted by Bonahon [8] using geodesic currents and McMullen [60] using the thermodynamic formalism. Wolpert [87] showed that Thurston's metric is a multiple of the Weil–Petersson metric. The key difference between our approach and McMullen's is that we work directly with the geodesic flow of the surface, rather than with a Bowen–Series coding of the action of the group on the limit set. Since such a coding is not known to exist for every hyperbolic group, this approach will be crucial to generalizing our results to the setting of all hyperbolic groups.

We next discuss the construction of the pressure metric in the simplest new situation: the Hitchin component of representations of a surface group into $\mathrm{PSL}_d(\mathbb{R})$. This setting offers the cleanest results and also several simplifications of the general proof. Given a Hitchin representation, inspired by earlier work of Sambarino [72], we construct a metric Anosov flow, which we call the geodesic flow of the representation, whose periods record the spectral radii of the elements in the image. We obtain a mapping class group invariant Riemannian metric on a Hitchin component whose restriction to the Fuchsian locus is a multiple of the Weil–Petersson metric.

We hope that the discussion of the pressure metric in these two simpler settings will provide motivation and intuition for the general construction. In §6 we discuss the more general settings studied in [17] with some comments on the additional difficulties which must be overcome. We finish with a discussion of open problems.

2. The thermodynamic formalism

The thermodynamic formalism was introduced by Bowen and Ruelle [10, 12, 70] as a tool to study the ergodic theory of Anosov flows and diffeomorphisms. It was further developed by Parry and Pollicott, their monograph [64] is a standard reference for the material covered here. McMullen [60] introduced the pressure form as a tool for constructing metrics on spaces which may be mapped into Hölder potentials over a shift-space. We will give a quick summary of the basic facts we will need, but we encourage the reader to consult the original references and the more complete discussion and references in [17].

We recall that a smooth flow $\phi = (\phi_t : X \rightarrow X)_{t \in \mathbb{R}}$ on a compact Riemannian manifold is said to be *Anosov* if there is a flow-invariant splitting $TX = E^s \oplus E_0 \oplus E^u$ where E_0 is a line bundle parallel to the flow and if $t > 0$, then $d\phi_t$ is exponentially contracting on E_+ and $d\phi_{-t}$ is exponentially contracting on E_- . We will always assume that our Anosov flows are *topologically transitive* (i.e. have a dense orbit). It is a celebrated theorem of Anosov (see [47, Theorem 17.5.1]) that the geodesic flow of a closed hyperbolic surface, and more generally of a closed negatively curved manifold, is a topologically transitive Anosov flow.

2.1. *Entropy, pressure and orbit-equivalence.* Let ϕ be a topologically transitive Anosov flow on a compact Riemannian manifold X . If a is a ϕ -periodic orbit, denote by $\ell(a)$ its period and let

$$R_T = \{a \text{ closed orbit} \mid \ell(a) \leq T\}.$$

Then, following Bowen [9], we may define the *topological entropy* of ϕ to be the exponential growth rate of the number of periodic orbits whose periods are at most T , i.e.

$$h(\phi) = \limsup_{T \rightarrow \infty} \frac{\log \#R_T}{T}.$$

Moreover, if $g : X \rightarrow \mathbb{R}$ is Hölder and a is a closed orbit, let

$$\ell_g(a) = \int_0^{\ell(a)} g(\phi_s(x)) \, ds,$$

where x is any point on a . Then, following Bowen and Ruelle [12], we may define the *topological pressure* of g (or simply *pressure*) by

$$\mathbf{P}(g) = \mathbf{P}(\phi, g) = \limsup_{T \rightarrow \infty} \frac{1}{T} \log \left(\sum_{a \in R_T} e^{\ell_g(a)} \right).$$

Note that $\mathbf{P}(g)$ only depends on the periods of g , i.e. the collection of numbers $\{\ell_g(a)\}$.

Livšic provides a *pointwise* relation for two functions having the same periods: two Hölder functions $f, g : X \rightarrow \mathbb{R}$ are *Livšic cohomologous* if there exists a Hölder function $V : X \rightarrow \mathbb{R}$, which is C^1 in the direction of the flow ϕ , such that

$$f(x) - g(x) = \left. \frac{\partial}{\partial t} \right|_{t=0} V(\phi_t(x)). \tag{1}$$

Livšic [53] proved the following fundamental result.

THEOREM 2.1. (Livšic [53]) *If ϕ is a topologically transitive Anosov flow and $g : X \rightarrow \mathbb{R}$ is a Hölder function such that $\ell_g(a) = 0$ for every closed orbit a , then g is Livšic cohomologous to 0.*

Given a positive Hölder function $f : X \rightarrow (0, \infty)$ one may define a *reparametrization* of the flow so that its ‘speed’ at a point x is multiplied by $f(x)$. More formally, let

$$\kappa_f(x, t) = \int_0^t f(\phi_s(x)) \, ds,$$

and define $\phi^f = (\phi_t^f : X \rightarrow X)_{t \in \mathbb{R}}$ so that $\phi_{\kappa_f(x,t)}^f(x) = \phi_t(x)$. In particular, if a is a ϕ -closed orbit then a is also a closed orbit of the flow ϕ^f with period $\ell_f(a)$.

A *Hölder orbit equivalence* between two flows is a Hölder homeomorphism that sends orbits to orbits. If, moreover, it preserves time, it is called a *Hölder conjugacy*. In particular, the identity map is a Hölder orbit equivalence from ϕ to ϕ^f when f is a positive Hölder function. (In our setting, the inverse of any Hölder orbit equivalence is also a Hölder orbit equivalence, but we will not need this fact.)

Livšic’s theorem implies that two positive Hölder functions are Livšic cohomologous if and only if the periods of ϕ^f and ϕ^g agree. If this is the case, the function V in equation (1) provides a Hölder conjugacy between ϕ^f and ϕ^g . Moreover, one has the following standard consequence of Livšic’s theorem (see Sambarino [73, Lemma 2.11]).

LEMMA 2.2. *If ϕ is a topologically transitive, Anosov flow and there exists Hölder orbit equivalence to a Hölder-continuous flow ψ , then there exists a Hölder function $f : X \rightarrow (0, \infty)$ such that ψ is Hölder conjugate to ϕ^f .*

The flow ϕ^f remains topologically transitive and is again Anosov but in a metric sense. More specifically, ϕ^f is a Smale flow in the sense of Pollicott [66]. Pollicott shows that all the results we rely on in the ensuing discussion generalize to the setting of Smale flows, which we will refer to as metric Anosov flows. Hence, we may define the topological entropy of ϕ^f as

$$h(f) = \limsup_{T \rightarrow \infty} \frac{\log \#R_T(f)}{T},$$

where $R_T(f) = \{a \text{ closed orbit} \mid \ell_f(a) \leq T\}$. We recall the following standard lemma which relates pressure and entropy.

LEMMA 2.3. (See Sambarino [72, Lemma 2.4]) *If ϕ is a topologically transitive Anosov flow and $f : X \rightarrow (0, \infty)$ is Hölder, then $\mathbf{P}(-hf) = 0$ if and only if $h = h(f)$.*

Ruelle [70, Corollary 7.10] (see also Parry and Pollicott [64, Proposition 4.7]) proved that the pressure function $\mathbf{P}(g)$ is a real analytic function of the Hölder function g . It follows from Lemma 2.3 and the Implicit Function Theorem that entropy varies analytically in f . Ruelle [71] used a similar observation to show that the Hausdorff dimension of a quasi-Fuchsian Kleinian group varies analytically on quasi-Fuchsian space.

Ruelle [70] also showed that \mathbf{P} is a convex function and thus if $f, g : X \rightarrow \mathbb{R}$ are Hölder functions,

$$\left. \frac{\partial^2}{\partial t^2} \right|_{t=0} \mathbf{P}(f + tg) \geq 0.$$

Consider the space

$$\mathcal{P}(X) = \{\Phi : X \rightarrow \mathbb{R} \text{ Hölder} \mid \mathbf{P}(\Phi) = 0\}$$

of pressure zero Hölder functions on X . It follows immediately from the definition above that the pressure function \mathbf{P} is constant on Livšic cohomology classes, so it is natural to consider the space

$$\mathcal{H}(X) = \mathcal{P}(X) / \sim$$

of Livšic cohomology classes of pressure zero functions.

McMullen [60] defined a pressure semi-norm on the tangent space of the space of pressure zero Hölder functions on a shift space. Similarly, we define a *pressure semi-norm* on $\mathbb{T}_f \mathcal{P}(X)$, by letting

$$\|g\|_{\mathbf{P}}^2 = \left(\left. \frac{\partial^2}{\partial t^2} \right|_{t=0} \mathbf{P}(f + tg) \right) \left(\frac{-1}{\partial/\partial t|_{t=0} \mathbf{P}(f + tf)} \right)$$

for all $g \in \mathbb{T}_f \mathcal{P}(X) = \ker D_f \mathbf{P}$. (Formally, one should consider the space $\mathcal{P}^\alpha(X)$ of α -Hölder pressure zero functions for some $\alpha > 0$. In all our applications, we will consider embeddings of analytic manifolds into $\mathcal{P}(X)$ such that every point has a neighborhood which maps into $\mathcal{P}^\alpha(X)$ for some $\alpha > 0$. We will consistently suppress this technical detail.)

One obtains the following characterization of degenerate vectors, due to Ruelle, and Parry and Pollicott.

THEOREM 2.4. (Ruelle [70], see also Parry and Pollicott [64, Proposition 4.12]) *Let ϕ be a topologically transitive Anosov flow and consider $g \in \mathcal{TP}(X)$. Then, $\|g\|_{\mathbf{P}} = 0$ if and only if g is Livšic cohomologous to zero, i.e. $\ell_g(a) = 0$ for every closed orbit a .*

We make use of the following nearly immediate corollary of this characterization (see the proof of [17, Lemma 9.3]).

COROLLARY 2.5. *Let ϕ be a topologically transitive Anosov flow. Suppose that $\{f_t\}_{t \in (-1,1)} : X \rightarrow (0, \infty)$ is a smooth one parameter family of Hölder functions. Consider $\Phi : (-1, 1) \rightarrow \mathcal{P}(X)$ defined by $\Phi(t) = -h(f_t)f_t$. Then $\|\Phi_0\|_{\mathbf{P}} = 0$ if and only if*

$$\frac{\partial}{\partial t} \Big|_{t=0} h(f_t)\ell_{f_t}(a) = 0$$

for every closed orbit a of ϕ .

2.2. Intersection and pressure form. Inspired by Bonahon’s [8] intersection number, we define the *intersection number* \mathbf{I} of two positive Hölder functions $f_1, f_2 : X \rightarrow (0, \infty)$ by

$$\mathbf{I}(f_1, f_2) = \lim_{T \rightarrow \infty} \frac{1}{\#R_T(f_1)} \sum_{a \in R_T(f_1)} \frac{\ell_{f_2}(a)}{\ell_{f_1}(a)}$$

and their *renormalized intersection number* by

$$\mathbf{J}(f_1, f_2) = \frac{h(f_2)}{h(f_1)} \mathbf{I}(f_1, f_2).$$

Bowen’s equidistribution theorem on periodic orbits [9] implies that \mathbf{I} , and hence \mathbf{J} , are well defined (see [17, §3.4]). One may use the analyticity of the pressure function and results of Parry and Pollicott [64] and Ruelle [70] to check that they are analytic functions.

PROPOSITION 2.6. [17, Proposition 3.12] *Let ϕ be a topologically transitive Anosov flow and let $\{f_u\}_{u \in M}$ and $\{g_u\}_{u \in M}$ be two analytic families of positive Hölder functions on X . Then $h(f_u)$ varies analytically over M and $\mathbf{I}(f_u, g_u)$ and $\mathbf{J}(f_u, g_u)$ vary analytically over $M \times M$.*

The seminal work of Bowen and Ruelle [12] may be used to derive the following crucial rigidity property for the renormalized intersection number.

PROPOSITION 2.7. [17, Proposition 3.8] *If ϕ is a topologically transitive Anosov flow on X and f and g are positive Hölder functions on X , then*

$$\mathbf{J}(f, g) \geq 1.$$

Moreover, $\mathbf{J}(f, g) = 1$ if and only if $h(f)f$ and $h(g)g$ are Livšic cohomologous.

If $\{f_u\}_{u \in M}$ is an analytic family of positive Hölder functions on X , then, for all $u \in M$, we consider the function

$$\mathbf{J}_u : M \rightarrow \mathbb{R}$$

given by $\mathbf{J}_u(v) = \mathbf{J}(u, v)$ for all $v \in M$. Proposition 2.7 implies that the Hessian of J_u gives a non-negative bilinear form on T_uM . Lemma 2.3 allows us to define a thermodynamic mapping

$$\Phi : M \rightarrow \mathcal{P}(X)$$

by letting $\Phi(u) = -h(f_u)f_u$. The following important, but fairly simple, result shows that this bilinear form is the pullback of the pressure form.

PROPOSITION 2.8. [17, Proposition 3.11] *Let ϕ be a topologically transitive Anosov flow. If $\{f_t\}_{t \in (-1,1)}$ is a smooth one parameter family of positive Hölder functions on X and $\Phi : (-1, 1) \rightarrow \mathcal{P}(X)$ is given by $\Phi(t) = -h(f_t)f_t$, then*

$$\|\dot{\Phi}_0\|_{\mathbf{P}}^2 = \left. \frac{\partial^2}{\partial t^2} \right|_{t=0} \mathbf{J}(f_0, f_t).$$

3. Basic strategy

Our basic strategy is inspired by McMullen’s [60] re-interpretation of Thurston’s Riemannian metric on Teichmüller space and its generalization to quasi-Fuchsian space by Bridgeman [14].

We consider a family $\{\rho_u : \Gamma \rightarrow \mathbf{G}\}_{u \in M}$ of (conjugacy classes of) representations of a word hyperbolic group Γ into a semi-simple Lie group \mathbf{G} parametrized by an analytic manifold M . We recall that Gromov [36] associated a geodesic flow $\phi = \{\phi_t : U_\Gamma \rightarrow U_\Gamma\}_{t \in \mathbb{R}}$ to a hyperbolic group Γ which agrees with the geodesic flow on T^1X in the case when Γ is the fundamental group of a negatively curved manifold X (see §6 for details).

In our two basic examples $\Gamma = \pi_1(S)$ where S is a closed, oriented surface of genus at least 2, and the flow ϕ is the geodesic flow on a hyperbolic surface homeomorphic to S . The first example will be the classical Teichmüller space $\mathcal{T}(S)$ of hyperbolic structures on S , where $\mathbf{G} = \text{PSL}_2(\mathbb{R})$ and $M = \mathcal{T}(S)$. The second is the Hitchin component $\mathcal{H}_d(S)$, where $\mathbf{G} = \text{PSL}_d(\mathbb{R})$ and $M = \mathcal{H}_d(S)$.

Step 1. Associate to each representation ρ_u a topologically transitive metric Anosov flow ϕ^{ρ_u} which is Hölder orbit equivalent to the geodesic flow ϕ of Γ so that the period of the orbit associated to $\gamma \in \Gamma$ is the ‘length’ of $\rho(\gamma)$.

In the case of $\mathcal{T}(S)$, ϕ^ρ will be the geodesic flow of the surface $X_\rho = \mathbb{H}^2/\rho(S)$. In the case of a Hitchin component, we will construct a geodesic flow and our notion of length will be the logarithm of the spectral radius.

If $u \in M$, Lemma 2.2 provides a positive Hölder function $f_u : U_\Gamma \rightarrow \mathbb{R}$, well defined up to Livšic cohomology, such that ϕ^{ρ_u} is Hölder conjugate to ϕ^{f_u} .

Step 2. Define a thermodynamic mapping $\Phi : M \rightarrow \mathcal{H}(U_\Gamma)$ by letting $\Phi(u) = [-h(f_u)f_u]$ and prove that it has locally analytic lifts, i.e. if $u \in M$, then there exists a neighborhood U of u in M and an analytic map $\tilde{\Phi} : U \rightarrow \mathcal{P}(U_\Gamma)$ which is a lift of $\Phi|_U$.

We may also define a renormalized intersection number on $M \times M$, by letting $\mathbf{J}(u, v) = \mathbf{J}(f_u, f_v)$.

Step 3. Define a pressure form on M by pulling back the pressure from on $\mathcal{P}(U_\Gamma)$ by (the lifts of) $\tilde{\Phi}$.

Lemma 2.8 allows us to reinterpret the pullback of the pressure form as the Hessian of the renormalized intersection number function.

Step 4. Prove, using Corollary 2.5, that the resulting pressure form is non-degenerate so gives rise to an analytic Riemannian metric on M .

Step 4 can fail in certain situations. For example, Bridgeman’s pressure metric on quasi-Fuchsian space [14] is degenerate exactly on the set of pure bending vectors on the Fuchsian locus. However, Bridgeman’s pressure metric still gives rise to a path metric.

3.1. *Historical remarks.* Thurston’s constructed a Riemannian metric which he describes as the ‘Hessian of the length of a random geodesic.’ Wolpert’s formulation [87] of this construction agrees with the Hessian of the intersection number of the geodesic flows. From this viewpoint, one regards $\mathbf{I}(\rho, \eta)$, as the length in X_η of a random unit length geodesic on X_ρ . If one considers a sequence $\{\gamma_n\}$ of closed geodesics on X_ρ which are becoming equidistributed (in the sense that $\{\gamma_n/(\ell_\rho(\gamma_n))\}$ converges, in the space of geodesic currents on S , to the Liouville current ν_ρ of X_ρ), then

$$\mathbf{I}(f_\rho, f_\eta) = \lim \frac{\ell_\eta(\gamma_n)}{\ell_\rho(\gamma_n)}.$$

Bonahon [8] reinterprets this to say that

$$\mathbf{I}(f_\rho, f_\eta) = i(\nu_\rho, \nu_\eta),$$

where i is the geometric intersection pairing on the space of geodesic currents.

Bridgeman and Taylor [18] used Patterson–Sullivan theory to show that the Hessian of the renormalized intersection number is a non-negative form on the quasi-Fuchsian case. McMullen [60] then introduced the use of the techniques of thermodynamic formalism to interpret both of these metrics as pullbacks of the pressure metric on the space of suspension flows on the shift space associated to the Bowen-Series coding. Bridgeman [14] then showed that the resulting pressure form on quasi-Fuchsian space is degenerate exactly on the set of pure bending vectors on the Fuchsian locus.

4. The pressure metric for Teichmüller space

In this section, we survey the construction of the pressure metric for the Teichmüller space $\mathcal{T}(S)$ of a closed oriented surface S of genus $g \geq 2$.

We recall that $\mathcal{T}(S)$ may be defined as the unique connected component of

$$\text{Hom}(\pi_1(S), \text{PSL}_2(\mathbb{R}))/\text{PGL}_2(\mathbb{R})$$

which consists of discrete and faithful representations. If $\rho \in \mathcal{T}(S)$, then one obtains a hyperbolic surface $X_\rho = \mathbb{H}^2/\rho(\pi_1(S))$ by regarding $\text{PSL}_2(\mathbb{R})$ as the space of orientation-preserving isometries of the hyperbolic plane \mathbb{H}^2 .

4.1. *Basic facts.* It is useful to isolate the facts that will make the construction much simpler in this case. All of these facts will fail even in the setting of the Hitchin component.

- (1) The space $T^1\mathbb{H}^2$ is canonically identified with the space of ordered triplets on the visual boundary $\partial_\infty\mathbb{H}^2$,

$$(\partial_\infty\mathbb{H}^2)^{(3)} = \{(x, y, z) \in (\partial_\infty\mathbb{H}^2)^3 : x < y < z\},$$

where $<$ is defined by a given orientation on the topological circle $\partial_\infty\mathbb{H}^2$, and (x, y, z) is identified with the unit tangent vector to the geodesic L joining x to z at the point which is the orthogonal projection of y to L .

- (2) The surface X_ρ is closed (since it is a surface homotopy equivalent to a closed surface). In fact, by Baer’s theorem, it is diffeomorphic to S . The geodesic flow ϕ^ρ on T^1X_ρ is thus a topologically transitive Anosov flow on a closed manifold.
- (3) The topological entropy $h(\rho)$ of ϕ^ρ is equal to 1 (in particular, constant).

Fact (1) is quite straight-forward: if $(p, v) \in T^1\mathbb{H}^2$, denote by $v_\infty \in \partial_\infty\mathbb{H}^2$ the limit at $+\infty$ of the geodesic ray starting at (p, v) , then the identification is

$$(p, v) \mapsto ((-v)_\infty, (iv)_\infty, v_\infty)$$

where $iv \in T^1\mathbb{H}^2$ is such that the base $\{v, iv\}$ is orthogonal and oriented.

Fact (3) is a standard consequence of the fact, due to Manning [56], that the entropy of the geodesic flow of a negatively curved manifold agrees with the exponential rate of volume growth of a ball of radius T in its universal cover. In this setting, the universal cover is always \mathbb{H}^2 so the entropy is always 1.

4.1.1. *Conventions.* For the remainder of the section we fix $\rho_0 \in \mathcal{T}(S)$ and identify S with X_{ρ_0} . We then obtain an identification of $\partial_\infty\pi_1(S)$ with $\partial_\infty\mathbb{H}^2$ and of T^1S with $T^1X_{\rho_0}$. Let $\phi = \phi^{\rho_0}$ be the geodesic flow on S .

It will be useful to choose an analytic lift $s : \mathcal{T}(S) \rightarrow \text{Hom}(\pi_1(S), \text{PSL}_2(\mathbb{R}))$. In order to do so, we pick non-commuting elements α and β in $\pi_1(S)$ and choose a representative $\rho = s([\rho])$ of $[\rho]$ such that $\rho(\alpha)$ has attracting fixed point $+\infty \in \partial_\infty\mathbb{H}^2$ and repelling fixed point 0, while $\rho(\beta)$ has attracting fixed point 1. From now on, we will implicitly identify $\mathcal{T}(S)$ with $s(\mathcal{T}(S))$. This choice will allow us to define our thermodynamic mapping into the space $\mathcal{P}(T^1S)$ of pressure zero Hölder functions on T^1S , rather than just into the space $\mathcal{H}(T^1S)$ of Livšic cohomology classes of pressure zero Hölder functions on X .

4.2. *Analytic variation of limit maps.* It is well known that any two Fuchsian representations are conjugate by a unique Hölder map.

PROPOSITION 4.1. *If $\rho, \eta \in \mathcal{T}(S)$, then there is a unique (ρ, η) -equivariant Hölder homeomorphism $\xi_{\rho,\eta} : \partial_\infty\mathbb{H}^2 \rightarrow \partial_\infty\mathbb{H}^2$. Moreover, $\xi_{\rho,\eta}$ varies analytically in η .*

Proof. By fact (2), there exists a diffeomorphism $h : X_\rho \rightarrow X_\eta$ in the homotopy class determined by $\eta \circ \rho^{-1}$. Choose a (ρ, η) -equivariant lift $\tilde{h} : \mathbb{H}^2 \rightarrow \mathbb{H}^2$ of h . Since \tilde{h} is quasi-conformal, classical results in complex analysis (see Ahlfors and Beurling [1]), imply that \tilde{h} extends to a quasi-symmetric map $\xi_{\rho,\eta} : \partial_\infty\mathbb{H}^2 \rightarrow \partial_\infty\mathbb{H}^2$. In particular, $\xi_{\rho,\eta}$ is a Hölder homeomorphism. Since \tilde{h} is (ρ, η) -equivariant, so is ξ . The resulting map is unique, since, by equivariance, if $\gamma \in \pi_1(S)$, then $\xi_{\rho,\eta}$ must take the attracting fixed point of $\rho(\gamma)$ to the attracting fixed point of $\eta(\gamma)$.

A more modern approach to the existence of $\xi_{\rho,\eta}$ uses the fact that \mathbb{H}^2 is a proper Gromov-hyperbolic metric space with boundary $\partial_\infty\mathbb{H}^2$ and that quasi-isometries of proper Gromov-hyperbolic metric spaces extend to Hölder homeomorphisms of their boundary. Since h is a bi-Lipschitz homeomorphism, it lifts to a bi-Lipschitz homeomorphism of \mathbb{H}^2 . In particular, \tilde{h} is a quasi-isometry of \mathbb{H}^2 .

It is a classical result in Teichmüller theory that $\xi_{\rho,\eta}$ varies analytically in η . A more modern, but still complex analytic, approach uses holomorphic motions and is sketched by McMullen [60, §2]. One allows η to vary over the space $QF(S)$ of (conjugacy classes of) convex cocompact (i.e. quasi-Fuchsian) representations of $\pi_1(S)$ into $\mathrm{PSL}_2(\mathbb{C})$. (Recall that $QF(S)$ is an open neighborhood of $\mathcal{T}(S)$ in the $\mathrm{PSL}_2(\mathbb{C})$ -character variety of $\pi_1(S)$.) If $\eta \in QF(S)$, there is a (ρ, η) -equivariant embedding $\xi_{\rho,\eta} : \partial_\infty\mathbb{H}^2 \rightarrow \widehat{\mathbb{C}}$ whose image is the limit set of $\eta(\pi_1(S))$. If $z \in \partial_\infty\mathbb{H}^2$ is a fixed point of a non-trivial element $\rho(\gamma)$, then $\xi_{\rho,\eta}(z)$ varies holomorphically in η . Slodkowski’s generalized Lambda lemma [76] then implies that $\xi_{\rho,\eta}$ varies complex analytically as η varies over $QF(S)$, and hence varies real analytically as η varies over $\mathcal{T}(S)$.

One may also prove analyticity by using techniques of Hirsch *et al* [40] as discussed in the next section. □

4.3. *The thermodynamic mapping.* The next proposition allows us to construct the thermodynamic mapping we use to define the pressure metric.

PROPOSITION 4.2. *For every $\eta \in \mathcal{T}(S)$, there exists a positive Hölder function $f_\eta : \mathbb{T}^1 S \rightarrow (0, \infty)$ such that*

$$\int_{[\gamma]} f_\eta = \ell_\eta(\gamma)$$

for all $\gamma \in \pi_1(S)$. Moreover, f_η varies analytically in η .

Proof. Let $\xi_{\rho_0,\eta}$ be the (ρ_0, η) -equivariant map provided by Proposition 4.1. The identification of $\mathbb{T}^1\mathbb{H}^2$ with $\partial_\infty\pi_1(S)^{(3)}$ gives a (ρ_0, η) -equivariant Hölder homeomorphism $\tilde{\sigma} : \mathbb{T}^1\mathbb{H}^2 \rightarrow \mathbb{T}^1\mathbb{H}^2$ defined by

$$\tilde{\sigma}(x, y, z) = (\xi_{\rho_0,\eta}(x), \xi_{\rho_0,\eta}(y), \xi_{\rho_0,\eta}(z)).$$

Since $\tilde{\sigma}$ is a (ρ_0, η) -equivariant map sending geodesics to geodesics, the quotient $\sigma : \mathbb{T}^1 S \rightarrow \mathbb{T}^1 X_\eta$ is a Hölder orbit equivalence between the geodesic flows $\phi = \phi^{\rho_0}$ and ϕ^η . Lemma 2.2 gives the existence of a function f_η , but in order to establish the analytic variation we give an explicit construction.

If $a, b, c, d \in \partial_\infty\mathbb{H}^2$, then the signed-distance between the orthogonal projections of b and c onto the geodesic with endpoints a and d is $\log |B(a, b, c, d)|$ where

$$B(a, b, c, d) = \frac{(a - c)(a - d)}{(b - d)(b - c)}$$

is the cross-ratio. Let

$$\kappa_{\rho,\eta}((x, y, z), t) = \log(B(\xi_{\rho_0,\eta}(x), \xi_{\rho_0,\eta}(z), \xi_{\rho_0,\eta}(y), \xi_{\rho_0,\eta}(u_t(x, y, z))))$$

where u_t is determined by $\phi_t^{\rho_0}(x, y, z) = (x, u_t(x, y, z), z)$. We average $\kappa_{\rho, \eta}$ over intervals of length one in the flow to obtain

$$\kappa_{\rho_0, \eta}^1((x, y, z), t) = \int_0^1 \kappa_{\rho_0, \eta}((x, y, z), t + s) ds.$$

Then

$$f_\eta(x, y, z) = \frac{\partial}{\partial t} \Big|_{t=0} \kappa_{\rho_0, \eta}^1((x, y, z), t)$$

is Hölder and varies analytically in η .

One may also prove the analyticity of the reparametrizations in this setting using the techniques of Katok et al [48]. □

Since $h(\phi^\eta) = h(f_\eta) = 1$, by fact (3) in §4.1, Lemma 2.3 implies that $\mathbf{P}(-f_\eta) = 0$, where \mathbf{P} is the pressure function associated to the geodesic flow ϕ^{ρ_0} on our base surface $S = X_{\rho_0}$. Hence, Proposition 4.2 provides an analytic map $\Phi : \mathcal{T}(S) \rightarrow \mathcal{P}(\mathbb{T}^1 S)$ from the Teichmüller space $\mathcal{T}(S)$ to the space $\mathcal{P}(\mathbb{T}^1 S)$ of pressure zero functions on the unit tangent bundle $\mathbb{T}^1 S$, given by

$$\Phi(\eta) = -f_\eta.$$

We call Φ the *thermodynamic mapping*. We note that Φ depends on our choice of $\rho_0 \in \mathcal{T}(S)$ and on the lift $s : \mathcal{T}(S) \rightarrow \text{Hom}(\pi_1(S), \text{PSL}_2(\mathbb{R}))$.

4.4. *The pressure metric.* We may then define a pressure form on $\mathcal{T}(S)$ by pulling back the pressure form on $\mathcal{P}(\mathbb{T}^1 S)$. Explicitly, if $\{\eta_t\}_{t \in (-1, 1)}$ is an analytic path in $\mathcal{T}(S)$, then we define

$$\|\dot{\eta}_0\|_{\mathbf{P}}^2 = \|d\Phi(\dot{\eta}_0)\|_{\mathbf{P}}^2.$$

Proposition 2.8 will allow us to identify the pressure form with the Hessian of the intersection number \mathbf{I} .

THEOREM 4.3. (Thurston and Wolpert [87], McMullen [60]) *The pressure form is an analytic Riemannian metric on $\mathcal{T}(S)$ which is invariant under the mapping class group and independent of the reference metric ρ_0 . Moreover, the resulting pressure metric is a constant multiple of the Weil–Petersson metric on $\mathcal{T}(S)$.*

Proof. We first show that the pressure form is non-degenerate, so gives rise to a Riemannian metric. Consider an analytic path $\{\eta_t\}_{t \in (-1, 1)} \subset \mathcal{T}(S)$. If $\|d\Phi(\dot{\eta}_0)\|_{\mathbf{P}} = 0$, then Lemma 2.5 implies that if $\gamma \in \pi_1(S)$, then

$$\frac{\partial}{\partial t} \Big|_{t=0} \ell_{\eta_t}(\gamma) = 0. \tag{2}$$

However, there exist $6g - 5$ elements $\{\gamma_1, \dots, \gamma_{6g-5}\}$ of $\pi_1(S)$, so that the mapping from $\mathcal{T}(S)$ into \mathbb{R}^{6g-5} given by taking ρ to $(\ell_\rho(\gamma_i))_{i=1}^{6g-5}$ is a real analytic embedding (see Schmutz [74]). Therefore, since $\partial/\partial t|_{t=0} \ell_{\eta_t}(\gamma_i) = 0$ for all i , we conclude that $\dot{\eta}_0 = 0$. Therefore, the pressure form is non-degenerate.

If $\rho, \eta \in \mathcal{T}(S)$, the intersection number

$$\mathbf{I}(\rho, \eta) = \mathbf{I}(f_\rho, f_\eta) = \lim_{T \rightarrow \infty} \frac{1}{\#R_T(\rho)} \sum_{[\gamma] \in R_T(\rho)} \frac{\ell_\sigma(\gamma)}{\ell_\rho(\gamma)},$$

where $\ell_\rho(\gamma)$ is the translation length of $\rho(\gamma)$ and $R_T(\rho)$ is the collection of conjugacy classes of elements of $\pi_1(S)$ whose images have translation length at most T . So, \mathbf{I} is independent of the reference metric ρ_0 and invariant by the action of the mapping class group of S . Proposition 2.8 states that

$$\|\dot{\eta}_0\|_{\mathbf{P}} = \frac{\partial^2}{\partial t^2} \Big|_{t=0} \mathbf{J}(f_{\eta_0}, f_{\eta_t}) = \frac{\partial^2}{\partial t^2} \Big|_{t=0} \mathbf{I}(f_{\eta_0}, f_{\eta_t})$$

(again by fact (3)) and thus the pressure metric is mapping class group invariant.

One may interpret $\mathbf{I}(\rho, \eta)$ as the length in X_η of a random unit length geodesic on X_ρ . So, the pressure metric is given by considering the Hessian of the length of a random geodesic. Since the pressure form agrees with Thurston’s metric, Wolpert’s work [87] implies that the pressure metric is a multiple of the Weil–Petersson metric. \square

5. The pressure metric on the Hitchin component

Let V be a vector space and G be a group. Recall that a representation $\tau : G \rightarrow \text{GL}(V)$ is *irreducible* if $\tau(G)$ has no proper invariant subspaces other than $\{0\}$. Let us begin by recalling the following well-known result, see for example Humphreys [43].

PROPOSITION 5.1. *For each integer $d \geq 2$ there exists an irreducible representation $\tau_d : \text{SL}_2(\mathbb{R}) \rightarrow \text{SL}_d(\mathbb{R})$, unique up to conjugation.*

The existence of such an irreducible action is an explicit construction we will now explain. Denote by $\text{Sym}^d(\mathbb{R}^2)$ the d -dimensional vector space of homogeneous polynomials on two variables of degree $d - 1$. A base for $\text{Sym}^d(\mathbb{R}^2)$ is, for example,

$$\mathcal{B} = \{x^{d-1}, x^{d-2}y, \dots, xy^{d-2}, y^{d-1}\}.$$

We identify x with $(1, 0)$ and y with $(0, 1)$ in \mathbb{R}^2 so that if $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{R})$, then $g \cdot x = ax + cy$ and $g \cdot y = bx + dy$. The action of $\text{SL}_2(\mathbb{R})$ on $\text{Sym}^d(\mathbb{R}^2)$ is defined on the base \mathcal{B} by

$$\tau_d(g) \cdot x^k y^{d-k-1} = (g \cdot x)^k (g \cdot y)^{d-k-1}.$$

As before, let S be a closed oriented surface of genus $g \geq 2$. Hitchin [41] studied the components of the space

$$\text{Hom}(\pi_1(S), \text{PSL}_d(\mathbb{R})) / \text{PGL}_d(\mathbb{R})$$

containing an element $\rho : \pi_1(S) \rightarrow \text{PSL}_d(\mathbb{R})$ that factors as

$$\pi_1(S) \xrightarrow{\rho_0} \text{PSL}_2(\mathbb{R}) \xrightarrow{\tau_d} \text{PSL}_d(\mathbb{R}),$$

where $\rho_0 \in \mathcal{T}(S)$.

By analogy with Teichmüller space, he named these components *Teichmüller components*, but they are now known as *Hitchin components*, and denoted by $\mathcal{H}_d(S)$. Each Hitchin component contains a copy of $\mathcal{T}(S)$, known as *the Fuchsian locus*, which is an image of $\mathcal{T}(S)$ under the mapping induced by τ_d . Hitchin proved the following remarkable result.

THEOREM 5.2. (Hitchin [41]) *Each Hitchin component $\mathcal{H}_d(S)$ is an analytic manifold diffeomorphic to $\mathbb{R}^{(d^2-1)(2g-2)} = \mathbb{R}^{|\chi(S)| \dim \mathrm{PSL}_d(\mathbb{R})}$.*

Hitchin [41] commented that ‘Unfortunately, the analytical point of view used for the proofs gives no indication of the geometrical significance of the Teichmüller component.’ Labourie [49] introduced dynamical techniques to show that Hitchin representations, i.e. representations in the Hitchin component, are geometrically meaningful. In particular, Hitchin representations are discrete, faithful, quasi-isometric embeddings. Labourie’s work significantly expanded the analogy between Hitchin components and Teichmüller spaces.

We view the following result as a further step in exploring this analogy. Its proof follows the same basic strategy as in the Teichmüller space setting, although there are several additional difficulties to overcome.

THEOREM 5.3. (Bridgeman et al [17]) *There exists an analytic Riemannian metric on $\mathcal{H}_d(S)$ which is invariant under the action of the mapping class group and restricts to a multiple of the Weil–Petersson metric on the Fuchsian locus.*

Remark.

- The mapping class group, regarded as a subgroup of $\mathrm{Out}(\pi_1(S))$, acts by precomposition on $\mathcal{H}_d(S)$.
- When $d = 3$, metrics have also been constructed by Darvishzadeh and Goldman [29] and Li [52]. Li [52] showed that both her metric and the metric constructed by Darvishzadeh and Goldman have the properties obtained in our result.

5.1. *Labourie’s work.* Labourie developed the theory of Anosov representations as a tool to study Hitchin representations. This theory was further developed by Guichard and Wienhard [39] and has played a central role in the subsequent development of higher Teichmüller theory. The following theorem summarizes some of the major consequences of Labourie’s work for Hitchin representations.

THEOREM 5.4. (Labourie [49, 50]) *If $\rho \in \mathcal{H}_d(S)$ then we have the following.*

- (1) ρ is discrete and faithful.
- (2) If $\gamma \in \pi_1(S)$ is non-trivial, then $\rho(\gamma)$ is diagonalizable over \mathbb{R} with distinct eigenvalues.
- (3) ρ is a quasi-isometric embedding.
- (4) ρ is irreducible.

Theorem 5.4 is based on Labourie’s proof that Hitchin representations are Anosov with respect to a minimal parabolic subgroup for $\mathrm{PSL}_d(\mathbb{R})$, i.e. the upper triangular matrices in $\mathrm{PSL}_d(\mathbb{R})$. We will develop the terminology necessary to give a definition.

A complete flag of \mathbb{R}^d is a sequence of vector subspaces $\{V_i\}_{i=1}^d$ such that $V_i \subset V_{i+1}$ and $\dim V_i = i$ for all $i = 1, \dots, d$. Two flags $\{V_i\}$ and $\{W_i\}$ are transverse if $V_i \cap W_{d-i} = \{0\}$ for all i . Denote by \mathcal{F} the space of complete flags and by $\mathcal{F}^{(2)}$ the space of pairs of transverse flags. The following result should be viewed as the analogue of the limit map constructed in Proposition 4.1.

THEOREM 5.5. (Labourie [49]) *If $\rho \in \mathcal{H}_d(S)$, then there exists a unique ρ -equivariant Hölder map $\xi_\rho : \partial_\infty \pi_1(S) \rightarrow \mathcal{F}$ such that, if $x \neq y$, then the flags $\xi_\rho(x)$ and $\xi_\rho(y)$ are transverse.*

If $\rho \in \mathcal{H}_d(S)$ and $x \in \partial_\infty \pi_1(S)$ then we denote by $\xi_\rho^{(k)}(x)$ the k th-dimensional space in the flag $\xi_\rho(x)$. Notice that if γ_+ is an attracting fixed point of the action of $\gamma \in \pi_1(S)$ on $\partial_\infty \pi_1(S)$, then $\xi_\rho^{(k)}(\gamma_+)$ is spanned by the eigenlines of $\rho(\gamma)$ associated to the k eigenvalues of largest modulus. In particular, $\xi_\rho^{(1)}(\gamma_+)$ is the attracting fixed point for the action of $\rho(\gamma)$ on $\mathbb{P}(\mathbb{R}^d)$ and $\xi_\rho^{(d-1)}(\gamma_-)$ is its repelling hyperplane (where γ^- is the repelling fixed point for the action of γ on $\pi_1(S)$).

When ρ is Fuchsian, Labourie’s map is an explicit construction, called the *Veronese embedding*, which is moreover τ_d -equivariant. This is a map from $\partial_\infty \mathbb{H}^2 = \mathbb{P}(\mathbb{R}^2) \rightarrow \mathcal{F}$ explicitly defined, identifying \mathbb{R}^d with $\text{Sym}^d(\mathbb{R}^2)$, by

$$\mathbb{R} \cdot (ax + by) \mapsto \{p \in \text{Sym}^d(\mathbb{R}^2) : p \text{ has } (ax + by)^{d-k} \text{ as a factor}\}_{k=1}^d.$$

Conventions. As in the previous section, we fix $\rho_0 \in \mathcal{T}(S)$, so that ρ_0 identifies S with X_{ρ_0} , and hence identifies $\partial_\infty \pi_1(S)$ with $\partial_\infty \mathbb{H}^2$ and $T^1 S$ with $T^1 X_{\rho_0}$. Let $\phi = \phi^{\rho_0}$ be the geodesic flow on S . Let

$$\partial_\infty \pi_1(S)^{(2)} = \{(x, y) \in \partial_\infty \pi_1(S)^2 : x \neq y\}$$

and consider the Hopf parametrization of $T^1 \mathbb{H}^2$ by $\partial_\infty \pi_1(S)^{(2)} \times \mathbb{R}$ where (x, y, t) is the point on the geodesic L joining x to y which is a (signed) distance t from the horocycle through y and a fixed basepoint for \mathbb{H}^2 .

Labourie considers the bundle E_ρ over $T^1 S$ which is the quotient of $T^1 \mathbb{H}^2 \times \mathcal{F}$ by $\pi_1(S)$ where $\gamma \in \pi_1(S)$ acts on $T^1 \mathbb{H}^2$ by $\rho_0(\gamma)$ and acts on \mathcal{F} by $\rho(\gamma)$. There is a flow $\tilde{\psi}^\rho$ on $T^1 \mathbb{H}^2 \times \mathcal{F}$ which acts by the geodesic flow on $T^1 \mathbb{H}^2$ and acts trivially on \mathcal{F} . The flow $\tilde{\psi}^\rho$ descends to a flow ψ^ρ on E_ρ . The limit map $\xi_\rho : \partial_\infty \pi_1(S) \rightarrow \mathcal{F}$ determines a section $\tilde{\sigma}_\rho : T^1 \mathbb{H}^2 \rightarrow \tilde{E}_\rho$ given by $\tilde{\sigma}(x, y, t) = ((x, y, t), \xi_\rho(x))$ which descends to a section $\sigma : T^1 S \rightarrow E_\rho$.

A representation $\rho : \pi_1(S) \rightarrow \text{PSL}_d(\mathbb{R})$ is *Anosov with respect to a minimal parabolic subgroup* if and only if there is a limit map with the properties in Theorem 5.5 such that the inverse of the associated flow ψ^ρ is contracting on $\sigma_\rho(T^1 S)$.

5.2. The geodesic flow of a Hitchin representation. We wish to associate a topologically transitive metric Anosov flow to each Hitchin representation. Since ρ is discrete and faithful, one is tempted to consider the geodesic flow of the associated locally symmetric space

$$N_\rho = \rho(\pi_1(S)) \backslash \text{PSL}_d(\mathbb{R}) / \text{PSO}(d).$$

However, N_ρ is neither closed, nor negatively curved, so its geodesic flow will not be Anosov. Moreover, this flow does not even have a nice compact invariant set where it is metric Anosov (see Sambarino [73, Proposition 3.5]).

Sambarino [72, §5] (or more specifically [72, Theorem 3.2, Corollary 5.3 and Proposition 5.4]) constructed metric Anosov flows associated to Hitchin representations

which are Hölder orbit equivalent to a geodesic flow on a hyperbolic surface such that the closed orbit associated to $\gamma \in \pi_1(S)$ has period $\log \Lambda_\gamma(\rho)$, where $\Lambda_\gamma(\rho)$ is the spectral radius of $\rho(\gamma)$, i.e. the largest modulus of the eigenvalues of $\rho(\gamma)$.

We will use these flows to construct a thermodynamic mapping and an associated pressure metric satisfying the conclusions of Theorem 5.3.

PROPOSITION 5.6. (Sambarino [72, §5]) *For every $\rho \in \mathcal{H}_d(S)$, there exists a positive Hölder function $f_\rho : T^1S \rightarrow (0, \infty)$ such that*

$$\int_{[\gamma]} f_\rho = \log \Lambda_\gamma(\rho)$$

for every $\gamma \in \pi_1(S)$.

Notice that ϕ^{f_ρ} is a topologically transitive, metric Anosov flow, Hölder orbit equivalent to the geodesic flow, whose periods are the logarithms of the spectral radii of $\rho(\pi_1(S))$. We call this flow the *geodesic flow of the Hitchin representation*.

We will give a different construction of the geodesic flow of a Hitchin representation, from [17], which generalizes easily to the setting of projective Anosov representations of a word-hyperbolic group into $SL_d(\mathbb{R})$. If $\rho \in \mathcal{H}_d(S)$, we let $\xi_\rho : \partial_\infty \pi_1(S) \rightarrow \mathcal{F}$ be the associated limit map to the space of complete flags where $\xi_\rho(x) = \{\xi_\rho^k(x)\}_{k=1}^d$. We consider the line bundle F_ρ over $\partial_\infty \pi_1(S)^{(2)}$ whose fiber at (x, y) is

$$\begin{aligned} M(x, y) &= \{(\varphi, v) \in (\mathbb{R}^d)^* \times \mathbb{R}^d \mid \ker \varphi = \xi_\rho^{(d-1)}(x), v \in \xi_\rho^{(1)}(y), \\ &\quad \varphi(v) = 1\} / (\varphi, v) \sim (-\varphi, -v). \end{aligned}$$

Consider the flow $\tilde{\phi}^\rho = (\tilde{\phi}_t^\rho : F_\rho \rightarrow F_\rho)_{t \in \mathbb{R}}$ given by

$$\tilde{\phi}_t^\rho(\varphi, v) = (e^{-t}\varphi, e^t v).$$

Notice that the $\pi_1(S)$ -action on \tilde{F}_ρ given by

$$\gamma(x, y, (\varphi, v)) = (\gamma x, \gamma y, (\varphi \circ \rho(\gamma)^{-1}, \rho(\gamma)v))$$

is free. We further show that it is properly discontinuous and co-compact, so $\tilde{\phi}^\rho$ descends to a flow ϕ^ρ on $U_\rho = F_\rho/\pi_1(S)$, which we call the *geodesic flow* of ρ . The proof proceeds by finding a ρ -equivariant orbit equivalence between $T^1\mathbb{H}^2$ and F_ρ .

PROPOSITION 5.7. [17, Propositions 4.1 and 4.2] *The group $\pi_1(S)$ acts properly discontinuous and cocompactly on F_ρ . The quotient flow ϕ^ρ on U_ρ is Hölder orbit equivalent to the geodesic flow on T^1S . Moreover, the closed orbit associated to $\gamma \in \pi_1(S)$ has ϕ^ρ -period $\log \Lambda_\rho(\gamma)$.*

Sketch of proof. Consider the flat bundle E_ρ over T^1S which is the quotient of $T^1\mathbb{H}^2 \times \mathbb{R}^d$ by $\pi_1(S)$ where $\gamma \in \pi_1(S)$ acts on $T^1\mathbb{H}^2$ by $\rho_0(\gamma)$ and acts on \mathbb{R}^d by $\rho(\gamma)$. One considers a flow $\tilde{\psi}^\rho$ on $T^1\mathbb{H}^2 \times \mathbb{R}^d$ which acts as the geodesic flow on $T^1\mathbb{H}^2$ and acts trivially on \mathbb{R}^d . The flow $\tilde{\psi}^\rho$ preserves the $\rho(\pi_1(S))$ -invariant line sub-bundle $\tilde{\Sigma}$ whose fiber over the point (x, y, t) is $\xi_\rho^{(1)}(x)$. Thus, $\tilde{\psi}^\rho$ descends to a flow ψ^ρ on E_ρ preserving the line sub-bundle Σ which is the quotient of $\tilde{\Sigma}$. Since ρ is Anosov with respect to a minimal parabolic subgroup, ψ^ρ is contracting on Σ (see [17, Lemma 2.4]). Since ψ^ρ is contracting on Σ one may use an averaging procedure to construct a metric τ on Σ with respect to which ψ^ρ is uniformly contracting.

LEMMA 5.8. [17, Lemma 4.3] *There exists a Hölder metric τ on Σ and $\beta > 0$, so that for all $t > 0$,*

$$(\psi_t^\rho)_*(\tau) < e^{-\beta t} \tau.$$

We construct a ρ -equivariant Hölder orbit equivalence

$$\tilde{j}(x, y, t) = (x, y, u(x, y, t))$$

where $\tilde{\tau}(u(x, y, t)) = 1$ and $\tilde{\tau}$ is the lift of τ to $\tilde{\Sigma}$. The map \tilde{j} is ρ -equivariant, since ξ_ρ is, and the fact that $\tilde{\tau}$ is uniformly contracting implies that \tilde{j} is injective. It remains to prove that \tilde{j} is proper to show that it is a homeomorphism. (We refer the reader to the proof of [17, Proposition 4.2] for this relatively simple argument.) Then, \tilde{j} descends to a Hölder orbit equivalence j between T^1S and U_ρ .

In order to complete the proof, it suffices to evaluate the period of the closed orbit associated to an element $\gamma \in \pi_1(S)$. The closed orbit associated to γ is the quotient of the fiber of F_ρ over (γ^+, γ^-) . If we pick $v \in \xi_\rho^{(1)}(\gamma^+)$ and $\varphi \in \xi_\rho^{(d-1)}(\gamma^-)$ so that $\varphi(v) = 1$, then

$$\begin{aligned} \gamma(\gamma^+, \gamma^-, (\varphi, v)) &= (\gamma^-, \gamma^+, (\pm(\Lambda_\rho(\gamma))^{-1}\varphi, \pm\Lambda_\rho(\gamma)v)) \\ &= \tilde{\phi}_{\log \Lambda_\rho(\gamma)}^\rho(\gamma^+, \gamma^-, (\varphi, v)), \end{aligned} \tag{3}$$

so the closed orbit has period $\log \Lambda(\rho(\gamma))$ as claimed. □

Notice that Proposition 5.6 follows immediately from Proposition 5.7 and Lemma 2.2.

5.3. *The thermodynamic mapping.* Proposition 5.6 allows us to construct a thermodynamic mapping in the Hitchin setting. Livšic’s theorem (Theorem 2.1) guarantees that if $\rho \in \mathcal{H}_d(S)$, then the Livšic cohomology class of the reparametrization function f_ρ is well defined. So, applying Lemma 2.3, we may define a thermodynamic mapping

$$\Phi : \mathcal{H}_d(S) \rightarrow \mathcal{H}(T^1S)$$

from the Hitchin component $\mathcal{H}_d(S)$ to the space $\mathcal{H}(T^1S)$ of Livšic cohomology classes of pressure zero Hölder functions on T^1S , by letting

$$\Phi(\rho) = [-h(\rho) f_\rho].$$

In order to construct an analytic pressure form, we need to know that Φ admits local analytic lifts to the space $\mathcal{P}(T^1S)$ of pressure zero Hölder functions on T^1S .

PROPOSITION 5.9. [17, Proposition 6.2] *The mapping Φ admits local analytic lifts to the space $\mathcal{P}(T^1S)$, i.e. each $\rho \in \mathcal{H}_d(S)$ has an open neighborhood W and an analytic map $\tilde{\Phi} : W \rightarrow \mathcal{P}(T^1S)$ such that $\Phi(\rho) = [\tilde{\Phi}(\rho)]$.*

Sketch of proof. Let $\rho \in \mathcal{H}_d(S)$. Choose a neighborhood V of ρ which we may implicitly identify with a submanifold of $\text{Hom}(\pi_1(S), \text{PSL}_d(\mathbb{R}))$ (by an analytic map whose composition with the projection map is the identity). Consider the \mathcal{F} -bundle $\tilde{A} = V \times T^1\mathbb{H}^2 \times \mathcal{F}$ over $V \times T^1\mathbb{H}^2$. There is a natural action of $\pi_1(S)$ on \tilde{A} so that $\gamma \in \pi_1(S)$ takes $(\eta, (x, y, t), F)$ to $(\eta, \gamma(x, y, t), \eta(\gamma)F)$ with quotient A . The limit map ξ_ρ determines a section σ_ρ of A over $\{\rho\} \times T^1S$.

The geodesic flow on T^1S lifts to a flow $\{\Psi_t\}_{t \in \mathbb{R}}$ on A (whose lift to \tilde{A} acts trivially in the V and \mathcal{F} direction). The Anosov property of Hitchin representations implies that the inverse flow is contracting on $\sigma_\rho(\{\rho\} \times T^1S)$. One may extend σ_ρ to a section σ of A over $V \times T^1S$ which varies analytically in the V coordinate (after first possibly restricting to a smaller neighborhood of the lift of ρ). One may now apply the machinery developed by Hirsch *et al* [40] (see also Shub [75]), to find a section τ of A over $W \times T^1S$, where W is a sub-neighborhood of V , so that the inverse flow preserves and is contracting along $\tau(W \times T^1S)$. Here the main idea is to apply the contraction mapping theorem cleverly to show that one may take

$$\tau(\eta, X) = \lim \Psi_{-nt_0}(\sigma(\eta, \Psi_{nt_0}(x)))$$

for some $t_0 > 0$ so that Ψ_{-t_0} is uniformly contracting. It follows from standard techniques that τ varies smoothly in the W direction and that the restriction to $\{\eta\} \times T^1S$ is Hölder for all $\eta \in W$. One must complexify the situation by considering representations into $\mathrm{PSL}_d(\mathbb{C})$ in order to verify that τ varies analytically in the W direction. (See [17, §6] for more details.)

The section τ lifts to a section $\tilde{\tau}$ of \tilde{A} which is induced by a map

$$\hat{\xi} : W \times \partial_\infty \pi_1(S) \rightarrow \mathcal{F}$$

which varies analytically in the W direction such that

$$\hat{\xi}_\eta = \hat{\xi}(\eta, \cdot) : \partial_\infty \pi_1(S) \rightarrow \mathbb{P}(\mathbb{R}^d)$$

is η -equivariant and Hölder for all $\eta \in W$. The uniqueness of limit maps for Hitchin representations guarantees that $\hat{\xi}_\eta = \xi_\eta$. So, ξ_η varies analytically over W .

One may then examine the proof of Proposition 5.7 and apply an averaging procedure, as in the Teichmüller space case, to produce an analytically varying family of Hölder function $\{f_\eta\}_{\eta \in W}$, so that the reparametrization of the geodesic flow on T^1S by f_η has the same periods as U_η . (Again to get analytic, rather than just smooth, variation one must complexify the situation. See [17, §6] for details.) Therefore, the map

$$\tilde{\Phi} : W \rightarrow \mathcal{P}(T^1S)$$

given by

$$\tilde{\Phi}(\eta) = -h(\eta) f_\eta$$

is an analytic local lift of Φ . □

5.4. Entropy and intersection numbers. Proposition 5.6 allows us to define entropy and intersection numbers for Hitchin representations. If $\rho \in \mathcal{H}_d(S)$, let

$$R_T(\rho) = \{[\gamma] \in [\pi_1(S)] \mid \log(\Lambda_\rho(\gamma)) \leq T\}.$$

The entropy of ρ is given by

$$h(\rho) = h(f_\rho) = \lim_{T \rightarrow \infty} \frac{\log \#R_T(\rho)}{T}.$$

The intersection number of ρ and η in $\mathcal{H}_d(S)$ is given by

$$\mathbf{I}(\rho, \eta) = \mathbf{I}(f_\rho, f_\eta) = \lim_{T \rightarrow \infty} \frac{1}{\#R_T(\rho)} \sum_{[\gamma] \in R_\rho(T)} \frac{\log(\Lambda_\eta(\gamma))}{\log(\Lambda_\rho(\gamma))}$$

and their renormalized intersection number is

$$\mathbf{J}(\rho, \eta) = \mathbf{J}(f_\rho, f_\eta) = \frac{h(\eta)}{h(\rho)} \mathbf{I}(\rho, \eta).$$

Proposition 5.9 and Corollary 2.6 immediately give the following corollary.

COROLLARY 5.10. *Entropy varies analytically over $\mathcal{H}_d(S)$ and intersection \mathbf{I} and renormalized intersection \mathbf{J} vary analytically over $\mathcal{H}_d(S) \times \mathcal{H}_d(S)$.*

Remark. It follows from Bonahon’s work [8], that the intersection number is symmetric on Teichmüller space. However, it is clear that the intersection number is not symmetric on the Hitchin component. For example, one may use the work of Zhang [91, 92] to exhibit for all $K > 1$ and $d \geq 3$, $\rho_1, \rho_2 \in \mathcal{H}_d(S)$ such that $\log \Lambda(\rho_1(\gamma)) \geq K \log \Lambda(\rho_2(\gamma))$ for all $\gamma \in \pi_1(S) - \{\text{id}\}$, so $\mathbf{I}(\rho_1, \rho_2) \geq K$ and $\mathbf{I}(\rho_2, \rho_1) \leq 1/K$. One expects that the renormalized intersection number is also asymmetric.

5.5. The pressure form. We then define the analytic pressure form on $\mathcal{H}_d(S)$ as the pullback of the pressure form on $\mathcal{P}(\mathbb{T}^1 S)$ using a lift of the thermodynamic mapping Φ . Explicitly, if $\{\eta_t\}_{t \in (-1, 1)}$ is an analytic path in $\mathcal{H}_d(S)$ and $\tilde{\Phi} : U \rightarrow \mathcal{P}(\mathbb{T}^1 S)$ is an analytic lift of Φ defined on a neighborhood U of ρ , then we define

$$\|\dot{\eta}_0\|_{\mathbf{P}}^2 = \|d\tilde{\Phi}(\dot{\eta}_0)\|_{\mathbf{P}}^2.$$

If $\rho \in \mathcal{H}_d(S)$ and $\mathbf{J}_\rho : \mathcal{H}_d(S) \rightarrow \mathbb{R}$ is defined by

$$\mathbf{J}_\rho(\eta) = \mathbf{J}(\rho, \eta) = \mathbf{J}(f_\rho, f_\eta),$$

then Proposition 2.8 implies that the pressure form on $\mathbb{T}_\rho \mathcal{H}_d(S)$ is the Hessian of \mathbf{J}_ρ at ρ . Since the renormalized intersection number is mapping class group invariant by definition, it follows that the pressure form is also mapping class group invariant. Wolpert’s theorem [87] implies that the restriction of the pressure form to the Fuchsian locus is a multiple of the Weil–Peterson metric. It only remains to show that the pressure form is positive definite, so gives rise to an analytic Riemannian metric on all of $\mathcal{H}_d(S)$.

5.6. Non-degeneracy of the pressure metric. We complete the proof of Theorem 5.3 by proving the following.

PROPOSITION 5.11. *The pressure form is non-degenerate at each point in $\mathcal{H}_d(S)$.*

We note that each Hitchin component $\mathcal{H}_d(S)$ lifts to a component of

$$X(\pi_1(S), \text{SL}_d(\mathbb{R})) = \text{Hom}(\pi_1(S), \text{SL}_d(\mathbb{R}))/\text{SL}_d(\mathbb{R})$$

and we will work in this lift throughout the proof (see Hitchin [41, §10]).

In particular, this allows us to define, for all $\gamma \in \pi_1(S)$, an analytic function $\text{Tr}_\gamma : \mathcal{H}_d(S) \rightarrow \mathbb{R}$, where $\text{Tr}_\gamma(\rho)$ is the trace of $\rho(\gamma)$.

As in the Teichmüller case, the proof proceeds by applying Corollary 2.5. If $\{\eta_t\}_{(-\varepsilon, \varepsilon)} \subset \mathcal{H}_d(S)$ is a path such that $\|\dot{\eta}_0\|_{\mathbf{P}} = \|d\Phi v\|_{\mathbf{P}} = 0$, then

$$\left. \frac{\partial}{\partial t} \right|_{t=0} h(f_{\eta_t}) \ell_{f_{\eta_t}}(\gamma) = 0 \tag{4}$$

for all $\gamma \in \pi_1(S)$. The main difference is that entropy is not constant in the Hitchin component.

If $\gamma \in \pi_1(S)$, we may think of $\log \Lambda_\gamma$ as an analytic function on $\mathcal{H}_d(S)$, where we recall that $\Lambda_\gamma(\rho)$ is the spectral radius of $\rho(\gamma)$. The following lemma is an immediate consequence of equation (4) (compare with equation (2) in §4.4).

LEMMA 5.12. *If $v \in T_\rho \mathcal{H}_d(S)$ and $\|D_\rho \Phi(v)\|_{\mathbf{P}} = 0$, then*

$$D_\rho \log \Lambda_\gamma(v) = -\frac{D_\rho h(v)}{h(\rho)} \log \Lambda_\gamma(\rho)$$

for all $\gamma \in \pi_1(S)$.

Lemma 5.12 implies that if $v \in T_\rho \mathcal{H}_d(S)$ is a degenerate vector, and we set $K = -(D_\rho h(v))/h(\rho)$, then $D_\rho \log \Lambda_\gamma(v) = K \log \Lambda_\gamma(\rho)$ for all $\gamma \in \pi_1(S)$. The next proposition, which is the key step in the proof of Proposition 5.11, then guarantees that the derivative of the trace function of every element is trivial in the direction v .

PROPOSITION 5.13. *If $v \in T_\rho \mathcal{H}_d(S)$ and there exists $K \in \mathbb{R}$ such that*

$$D_\rho \log \Lambda_\gamma(v) = K \log \Lambda_\gamma(\rho)$$

for all $\gamma \in \pi_1(S)$, then $K = 0$ and $D_\rho \text{Tr}_\gamma(v) = 0$ for all $\gamma \in \pi_1(S)$.

The proof of Proposition 5.11, and hence Theorem 5.3, is then completed by applying the following standard lemma.

LEMMA 5.14. *If $\rho \in \mathcal{H}_d(S)$, then $\{D_\rho \text{Tr}_\gamma \mid \gamma \in \pi_1(S)\}$ spans the cotangent space $T_\rho^* \mathcal{H}_d(S)$.*

Since every Hitchin representation is absolutely irreducible, Schur’s lemma can be used to show that $\mathcal{H}_d(S)$ immerses in the $\text{SL}_d(\mathbb{C})$ -character variety $X(\pi_1(S), \text{SL}_d(\mathbb{C}))$. Lemma 5.14 then follows from standard facts about $X(\pi_1(S), \text{SL}_d(\mathbb{C}))$ (see Lubotzky and Magid [55]).

Proof of Proposition 5.13. It will be useful to introduce some notation. If M is a real analytic manifold, an analytic function $f : M \rightarrow \mathbb{R}$ has *log-type* K at $v \in T_u M$ if $f(u) \neq 0$ and

$$D_u \log(|f|)(v) = K \log(|f(u)|).$$

Suppose that $A \in \text{SL}_d(\mathbb{R})$ has real eigenvalues $\{\lambda_i(A)\}_{i=1}^n$ where

$$|\lambda_1(A)| > |\lambda_2(A)| > \dots > |\lambda_m(A)|.$$

If $\mathbf{p}_i(A)$ is the projection onto the $\lambda_i(A)$ -eigenspace parallel to the hyperplane spanned by the other eigenspaces, then

$$A = \sum_{k=1}^m \lambda_k(A) \mathbf{p}_k(A). \tag{5}$$

We say that two infinite order elements of $\pi_1(S)$ are *coprime* if they do not share a common power. The following lemma is an elementary computation (see Benoist [4, Corollary 1.6] or [17, Proposition 9.4]).

LEMMA 5.15. *If α and β are coprime elements of $\pi_1(S)$ and $\rho \in \mathcal{H}_d(S)$, then*

$$\text{Tr}(\mathbf{p}_1(\rho(\alpha))\mathbf{p}_1(\rho(\beta))) = \lim_{n \rightarrow \infty} \frac{\lambda_1(\rho(\alpha^n \beta^n))}{\lambda_1(\rho(\alpha^n))\lambda_1(\rho(\beta^n))} \neq 0$$

and

$$\text{Tr}(\mathbf{p}_1(\rho(\alpha))\rho(\beta)) = \lim_{n \rightarrow \infty} \frac{\lambda_1(\rho(\alpha^n \beta))}{\lambda_1(\rho(\alpha^n))} \neq 0$$

for all $\rho \in \mathcal{H}_d(S)$.

The following rather technical lemma plays a key role in the proof of Proposition 5.13.

LEMMA 5.16. *Suppose that $\{a_p\}_{p=1}^q$, $\{u_p\}_{p=1}^q$, $\{b_s\}_{s=1}^\infty$, and $\{v_s\}_{s=1}^\infty$ are collections of real numbers so that $\{|u_p|\}_{p=1}^q$ and $\{|v_s|\}_{s=1}^\infty$ are strictly decreasing, each u_p is non-zero,*

$$\sum_{p=1}^q n a_p u_p^n = \sum_{s=1}^\infty b_s v_s^n$$

for all $n > 0$, and $\sum_{s=1}^\infty b_s v_s$ is absolutely convergent. Then, $a_p = 0$ for all p .

Proof. We may assume without loss of generality that each b_s is non-zero. We divide each side of the equality by nu_1^n , to see that

$$a_1 = \lim_{n \rightarrow \infty} \sum_{s=1}^\infty \left(\frac{b_s}{n}\right) \frac{v_s^n}{u_1^n}$$

for all n . However, the right-hand side of the equation can only be bounded as $n \rightarrow \infty$, if $|v_1| \leq |u_1|$. However, if $|v_1| \leq |u_1|$, then the limit of the right-hand side, as $n \rightarrow \infty$, must be 0 and we conclude that $a_1 = 0$.

We may iterate this procedure to conclude that $a_p = 0$ for all p . □

Suppose that $\alpha, \beta \in \pi_1(S)$ are coprime. We consider the analytic function $F_n : \mathcal{H}_d(S) \rightarrow \mathbb{R}$ given by

$$F_n(\rho) = \frac{\text{Tr}(\mathbf{p}_1(\rho(\alpha))\rho(\beta^n))}{\lambda_1(\rho(\beta^n)) \text{Tr}(\mathbf{p}_1(\rho(\alpha))\mathbf{p}_1(\rho(\beta)))}.$$

Lemma 5.15 and the assumption of Proposition 5.13 imply that F_n is of log-type K at v (see the proofs of Proposition 9.4 and Lemma 9.8 in [17]). Using equation (5) we have

$$\rho(\beta^n) = \sum_{k=1}^d \lambda_k(\rho(\beta))^n \mathbf{p}_k(\rho(\beta)).$$

Thus, we can write F_n as

$$F_n(\rho) = 1 + \sum_{k=2}^d \frac{\text{Tr}(\mathbf{p}_1(\rho(\alpha))\mathbf{p}_k(\rho(\beta)))}{\text{Tr}(\mathbf{p}_1(\rho(\alpha))\mathbf{p}_1(\rho(\beta)))} \left(\frac{\lambda_k(\rho(\beta))}{\lambda_1(\rho(\beta))} \right)^n = 1 + \sum_{k=2}^d f_k t_k^n$$

where

$$f_k(\rho) = \frac{\text{Tr}(\mathbf{p}_1(\rho(\alpha))\mathbf{p}_k(\rho(\beta)))}{\text{Tr}(\mathbf{p}_1(\rho(\alpha))\mathbf{p}_1(\rho(\beta)))} \neq 0 \quad \text{and} \quad t_k(\rho) = \left(\frac{\lambda_k(\rho(\beta))}{|\lambda_1(\rho(\beta))|} \right) \neq 0.$$

Since F_n has log-type K at v and is positive in some neighborhood of ρ ,

$$D_\rho F_n(v) = \sum_{k=2}^d n f_k t_k^n \dot{t}_k + \sum_{k=2}^d \dot{f}_k t_k^n = K F_n(\rho) \log(F_n(\rho)), \tag{6}$$

where $\dot{t}_k = D_\rho t_k(v)$ and $\dot{f}_k = D_\rho f_k(v)$. In order to simplify the proof, we consider equation (6) for even powers. Using the Taylor series expansion for $\log(1+x)$ and grouping terms we have

$$F_{2n} \log(F_{2n}) = \left(1 + \sum_{k=2}^d f_k t_k^{2n} \right) \log \left(1 + \sum_{k=2}^d f_k t_k^{2n} \right) = \sum_{s=1}^\infty c_s w_s^n$$

where $\{w_s\}$ is a strictly decreasing sequence of positive terms. We may again regroup terms to obtain

$$\sum_{k=2}^d 2n \left(\frac{f_k \dot{t}_k}{t_k} \right) t_k^{2n} = \sum_{s=1}^\infty c_s w_s^n - \sum_{k=2}^d \dot{f}_k t_k^{2n} = \sum_{s=1}^\infty b_s v_s^n$$

where $\{v_s\}$ is a strictly decreasing sequence of positive terms. So, letting $u_k = t_k^2$, we see that for all n

$$\sum_{k=2}^d n \left(\frac{2 f_k \dot{t}_k}{t_k} \right) u_k^n = \sum_{s=1}^\infty b_s v_s^n.$$

Lemma 5.16 implies that $(f_k \dot{t}_k)/t_k = 0$ for all k , so $\dot{t}_k = 0$ for all k . Let $\lambda_{i,\beta}$ be the real-valued analytic function on $\mathcal{H}_d(S)$ given by $\lambda_{i,\beta}(\rho) = \lambda_i(\rho(\beta))$. Then,

$$\frac{\dot{\lambda}_{k,\beta} \lambda_{1,\beta} - \dot{\lambda}_{1,\beta} \lambda_{k,\beta}}{\lambda_{1,\beta}^2} = 0.$$

So,

$$D_\rho(\log(|\lambda_{k,\beta}|))(v) = \frac{\dot{\lambda}_{k,\beta}}{\lambda_{k,\beta}} = \frac{\dot{\lambda}_{1,\beta}}{\lambda_{1,\beta}} = D_\rho(\log |\lambda_{1,\beta}|)(v) = K \log(|\lambda_{1,\beta}(\rho)|).$$

Since $\lambda_{d,\beta} = 1/\lambda_{1,\beta-1}$,

$$\begin{aligned} K \log(|\lambda_{1,\beta-1}(\rho)|) &= D_\rho(\log(|\lambda_{1,\beta-1}|))(v) = D_\rho(\log(|\lambda_{d,\beta-1}|))(v) \\ &= -D_\rho(|\log(\lambda_{1,\beta})|)(v) = -K \log(|\lambda_{1,\beta}(\rho)|). \end{aligned}$$

Therefore, since $\log(|\lambda_{1,\beta-1}(\rho)|)$ and $\log(|\lambda_{1,\beta}(\rho)|)$ are both positive, then $K = 0$, which implies that $\dot{\lambda}_k(\beta) = 0$ for all k .

Since, $D_\rho \lambda_{i,\beta}(v) = 0$ for all i and all β , $D_\rho \text{Tr} \beta = 0$ for all $\beta \in \pi_1(S)$. This completes the proof of Proposition 5.13, and hence Proposition 5.11 and Theorem 5.3. □

5.7. Rigidity theorems for Hitchin representations.

5.7.1. Entropy rigidity. Potrie and Sambarino recently showed that entropy is maximized only along the Fuchsian locus. One may view this as an analogue of Bowen’s celebrated result [11] that the topological entropy of a quasi-Fuchsian group is at least 1 and it is 1 if and only if the group is Fuchsian.

THEOREM 5.17. (Potrie and Sambarino [69]) *If $\rho \in \mathcal{H}_d(S)$, then $h(\rho) \leq 2/(d - 1)$. Moreover, if $h(\rho) = 2/(d - 1)$, then ρ lies in the Fuchsian locus.*

Remarks. (1) Crampon [27] had earlier established that the entropy associated to Hilbert length (see §5.8) of holonomies of strictly convex projective structures on closed hyperbolic manifolds is maximal exactly at the representations into $\mathrm{SO}(d, 1)$. In particular, the Hilbert length entropy on $\mathcal{H}_3(S)$ is maximal exactly at the Fuchsian locus.

(2) Zhang [91, 92] showed that, for all d , there exist large families of sequences of Hitchin representations with entropy converging to 0. Nie [62] had earlier constructed specific examples when $d = 3$.

5.7.2. Intersection number and marked length rigidity theorems. One also obtains the following rigidity theorem for Hitchin representations with respect to the intersection number. Notice that the definition of the renormalized intersection number $\mathbf{J}(\rho_1, \rho_2)$ for $\rho_1 \in \mathcal{H}_{d_1}(S)$ and $\rho_2 \in \mathcal{H}_{d_2}(S)$ makes sense even if $d_1 \neq d_2$, see §5.4. Moreover, if $f_1 : \mathbb{T}^1 S \rightarrow \mathbb{R}$ and $f_2 : \mathbb{T}^1 S \rightarrow \mathbb{R}$ are positive Hölder functions such that $\phi^{f_1} = U_{\rho_1}$ and $\phi^{f_2} = U_{\rho_2}$, then $\mathbf{J}(\rho_1, \rho_2) = \mathbf{J}(f_1, f_2)$. In particular, see Lemma 2.7, $\mathbf{J}(\rho_1, \rho_2) \geq 1$.

THEOREM 5.18. [17, Corollary 1.5] *Let S be a closed, orientable surface and let $\rho_1 \in \mathcal{H}_{d_1}(S)$ and $\rho_2 \in \mathcal{H}_{d_2}(S)$ be two Hitchin representations such that*

$$\mathbf{J}(\rho_1, \rho_2) = 1.$$

Then, either:

- (1) $d_1 = d_2$ and $\rho_1 = \rho_2$ in $\mathcal{H}_{d_1}(S)$; or
- (2) *there exists an element ρ of the Teichmüller space $\mathcal{T}(S)$ so that $\rho_1 = \tau_{d_1}(\rho)$ and $\rho_2 = \tau_{d_2}(\rho)$.*

The proof of Theorem 5.18 makes use of general rigidity results in the thermodynamic formalism and a result of Guichard [38] classifying Zariski closures of images of Hitchin representations.

As an immediate corollary, one obtains a marked length rigidity theorem where one uses the logarithm of the spectral radius as a notion of length.

COROLLARY 5.19. *If $\rho_1, \rho_2 \in \mathcal{H}_d(S)$, then*

$$\frac{h(\rho_1)}{h(\rho_2)} \sup_{\gamma \in \pi_1(S)} \frac{\Lambda_\gamma(\rho_1)}{\Lambda_\gamma(\rho_2)} \geq 1$$

with equality if and only if there exists $g \in \mathrm{GL}_d(\mathbb{R})$ such that $g\rho_1g^{-1} = \rho_2$. In particular, if $\Lambda_\gamma(\rho_1) = \Lambda_\gamma(\rho_2)$ for all $\gamma \in \pi_1(S)$, then ρ_1 and ρ_2 are conjugate in $\mathrm{GL}_d(\mathbb{R})$.

Bridgeman, Canary and Labourie have recently established that it suffices to consider lengths of simple closed curves.

THEOREM 5.20. (Bridgeman et al [16]) *If $\rho_1, \rho_2 \in \mathcal{H}_d(S)$ and $\Lambda_\alpha(\rho_1) = \Lambda_\alpha(\rho_2)$ for all $\alpha \in \pi_1(S)$ which are represented by simple closed curves, then ρ_1 and ρ_2 are conjugate in $\text{PGL}_d(\mathbb{R})$.*

Remarks. Burger [19] introduced a renormalized intersection number between convex cocompact representations into rank-one Lie groups and proved an analogue of Theorem 5.18 in that setting. One should compare Corollary 5.19 and Theorem 5.20 with the marked length spectrum rigidity theorem of Dal’bo and Kim [28] for Zariski dense representations. Both Dal’bo and Kim [28] and Theorem 5.18 rely crucially on work of Benoist [3, Theorem 1.2]. However, the proof of Theorem 5.20 uses Labourie’s equivariant Frenet map into the flag variety, see Theorem 5.5, and the theory of positive representations developed by Fock and Goncharov [31].

5.8. *An alternate length function.* Throughout the section, we have used the logarithm of the spectral radius as a notion of length. It is also quite natural to consider the length of $\rho(\gamma)$ to be

$$\ell_H(\rho(\gamma)) = \log \Lambda(\rho(\gamma)) + \log \Lambda(\rho(\gamma^{-1})).$$

For example, if $\rho \in \mathcal{H}_3(S)$, then ρ is the holonomy of a convex projective structure on S , and $\ell_H(\rho(\gamma))$ is the translation length of γ in the associated Hilbert metric on S .

Sambarino [72] also proves that there is a reparametrization of T^1S whose periods are given by $\ell_H(\rho(\gamma))$.

PROPOSITION 5.21. (Sambarino [72, Theorem 3.2, §5]) *If $\rho \in \mathcal{H}_d(S)$, then there exists a positive Hölder function $f_\rho^H : T^1S \rightarrow (0, \infty)$ such that*

$$\int_{[\gamma]} f_\rho^H = \ell_H(\rho(\gamma))$$

for all $\gamma \in \pi_1(S)$.

We give a proof which uses a cross-ratio to construct f_ρ^H from the limit map ξ_ρ , as is done in the Teichmüller setting. It is adapted from the construction given in Labourie [50, §3].

Proof. Given linear forms $\varphi, \psi \in (\mathbb{R}^d)^*$ and vectors $v, w \in \mathbb{R}^d$ such that $v \notin \ker \psi$ and $w \notin \ker \varphi$, define the *cross-ratio*

$$[\varphi, \psi, v, w] = \frac{\varphi(v)\psi(w)}{\varphi(w)\psi(v)}.$$

Note that the cross-ratio only depends on the projective classes of φ, ψ, v , and w , and is invariant under $\text{PSL}_d(\mathbb{R})$. Moreover, if $g \in \text{PSL}_d(\mathbb{R})$ is bi-proximal and $v \notin \ker g_- \cup \ker(g^{-1})_-$, then

$$[g_-, (g^{-1})_-, v, gv] = \Lambda(g)\Lambda(g^{-1}) \tag{7}$$

where g_- is a linear functional whose kernel is the repelling hyperplane of g .

Theorem 5.5 provides a ρ -equivariant map $\xi_\rho : \partial_\infty \mathbb{H}^2 \rightarrow \mathcal{F}$. We then define $\kappa_\eta : \mathbb{T}^1 \mathbb{H}^2 \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$\kappa_\eta((x, y, z), t) = \log |[\xi_\eta^{(d-1)}(x), \xi_\eta^{(d-1)}(z), \xi_\eta^{(1)}(y), \xi_\eta^{(1)}(u_t(x, y, z))]|,$$

where u_t is determined by $\phi_t(x, y, z) = (x, u_t(x, y, z), z)$. Work of Labourie [50, §3] implies that $t \mapsto \kappa_\eta((x, y, z), t)$ is an increasing homeomorphism of \mathbb{R} , so averaging κ_η and taking derivatives as before provides the desired function $f_\eta^H : \mathbb{T}^1 S \rightarrow (0, \infty)$. Equation (7) implies that f_η^H has the desired periods. □

We may again obtain a thermodynamic mapping $\Phi^H : \mathcal{H}_d(S) \rightarrow \mathcal{H}(T^1 S)$ defined by

$$\eta \mapsto [-h(f_\eta^H) f_\eta^H].$$

One can use the same arguments as above to show that Φ^H has locally analytic lifts and one can pull back the pressure form via Φ^H to obtain an analytic pressure semi-norm $\|\cdot\|_H$ on $T\mathcal{H}_d(S)$. (Pollicott and Sharp [67] previously proved that the entropy associated to ℓ_H varies analytically over $\mathcal{H}_d(S)$.) However, this pressure form is degenerate in ways which are completely analogous to the degeneracy of the pressure metric on quasi-Fuchsian space discovered by Bridgeman [14].

Consider the *contragredient involution* $\sigma : \mathrm{PSL}_d(\mathbb{R}) \rightarrow \mathrm{PSL}_d(\mathbb{R})$ given by $g \mapsto (g^{-1})^t$, where t denotes the transpose operator associated to the standard inner product of \mathbb{R}^d . This involution induces an involution on the Hitchin component $\hat{\sigma} : \mathcal{H}_d(S) \rightarrow \mathcal{H}_d(S)$, where $\hat{\sigma}(\rho)(\gamma) = \sigma(\rho(\gamma))$ for all $\gamma \in \pi_1(S)$. If $\eta \in \mathcal{H}_d(S)$ is a representation whose image lies in (a group conjugate to) $\mathrm{Sp}(2n, \mathbb{R})$ (if $d = 2n$) or $\mathrm{SO}(n, n + 1, \mathbb{R})$ (if $d = 2n + 1$), then $\hat{\sigma}(\eta) = \eta$.

Consider the tangent vectors in $T\mathcal{H}_d(S)$ which are reversed by $D\hat{\sigma}$, i.e. let

$$B_d(S) = \{v \in T\mathcal{H}_d(S) : D\hat{\sigma}(v) = -v\}.$$

The vectors in $B_d(S)$ are degenerate for the pressure metric $\|\cdot\|_H$.

LEMMA 5.22. *If $v \in B_d(S)$, then $\|v\|_H = 0$.*

Proof. Consider a path $\{\eta_t\}_{(-1,1)} \subset \mathcal{H}_d(S)$ so that $\hat{\sigma}(\eta_t) = \eta_{-t}$ for all $t \in (-1, 1)$. Then, $\ell_H(\eta_t(\gamma)) = \ell_H(\eta_{-t}(\gamma))$ and $h(f_{\eta_t}^H) = h(f_{\eta_{-t}}^H)$ for all $t \in (-1, 1)$ and $\gamma \in \pi_1(S)$. Therefore,

$$\left. \frac{\partial}{\partial t} \right|_{t=0} h(f_{\eta_t}^H) \ell_{f_{\eta_t}^H}(\gamma) = 0$$

for all $\gamma \in \pi_1(S)$. Corollary 2.5 then implies that $\|v\|_H = 0$. □

Remark. With a little more effort one may use the techniques of [17] to show that these are the only degenerate vectors for $\|\cdot\|_H$ and that $\|\cdot\|_H$ induces a path metric on the Hitchin component.

6. Generalizations and consequences

In [17] we work in the more general setting of Anosov representations of word hyperbolic groups into semi-simple Lie groups. In this section, we will survey these more general results and discuss some of the additional difficulties which occur. The bulk of the work in [17] is done in the setting of projective Anosov representations into $SL_d(\mathbb{R})$. We note that Hitchin representations are examples of projective Anosov representations as are Benoist representations, i.e. holonomy representations of closed strictly convex (real) projective manifolds (see Guichard and Wienhard [39, Proposition 6.1]).

6.1. *Projective Anosov representations.* We first show that the pressure form gives an analytic Riemannian metric on the space of (conjugacy classes of) projective Anosov, generic, regular†, irreducible representations. In order to define projective Anosov representations, we begin by recalling basic facts about the geodesic flow of a word hyperbolic group.

Gromov [36] first established that a word hyperbolic group Γ has an associated geodesic flow U_Γ . Roughly, one considers the obvious flow on the space of all geodesics in the Cayley graph of Γ , collapses all geodesics joining two points in the Gromov boundary to a single geodesic, and considers the quotient by the action of Γ . We make use of the version due to Mineyev [61] (see also Champetier [24]). Mineyev defines a proper cocompact action of Γ on $\widetilde{U}_\Gamma = \partial_\infty \Gamma^{(2)} \times \mathbb{R}$ and a metric on \widetilde{U}_Γ , well defined only up to Hölder equivalence, so that Γ acts by isometries, every orbit of \mathbb{R} is quasi-isometrically embedded, and the \mathbb{R} -action is by Lipschitz homeomorphisms. Moreover, the \mathbb{R} -action descends to a flow on $U_\Gamma = \widetilde{U}_\Gamma / \Gamma$. In the case that Γ is the fundamental group of a negatively curved manifold M , one may take U_Γ to be the geodesic flow on $T^1 M$.

A representation $\rho : \Gamma \rightarrow SL_d(\mathbb{R})$ has *transverse projective limit maps* if there exist continuous, ρ -equivariant limit maps

$$\xi_\rho : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$$

and

$$\theta_\rho : \partial_\infty \Gamma \rightarrow Gr_{d-1}(\mathbb{R}^d) = \mathbb{P}((\mathbb{R}^d)^*)$$

so that if x and y are distinct points in $\partial_\infty \Gamma$, then

$$\xi_\rho(x) \oplus \theta_\rho(y) = \mathbb{R}^d.$$

A representation ρ with transverse projective limit maps determines a splitting $\Xi \oplus \Theta$ of the flat bundle E_ρ over U_Γ . Concretely, if \widetilde{E}_ρ is the lifted bundle over \widetilde{U}_Γ , then the lift $\widetilde{\Xi}$ of Ξ has fiber $\xi_\rho(x)$ and the lift $\widetilde{\Theta}$ of Θ has fiber $\theta_\rho(y)$ over the point (x, y, t) . The geodesic flow on U_Γ lifts to a flow on \widetilde{U}_Γ which extends, trivially in the bundle factor, to a flow on \widetilde{E}_ρ which descends to a flow on E_ρ . One says that ρ is *projective Anosov* if the resulting flow on the associated bundle $\text{Hom}(\Theta, \Xi) = \Xi \otimes \Theta^*$ is contracting.

Projective Anosov representations are discrete, well-displacing, quasi-isometric embeddings with finite kernel such that the image of each infinite order element is bi-proximal, i.e. its eigenvalues of maximal and minimal modulus have multiplicity one

† A representation $\rho : \Gamma \rightarrow SL_d(\mathbb{R})$ is *regular* if it is a smooth point of the algebraic variety $\text{Hom}(\Gamma, SL_d(\mathbb{R}))$.

(see Labourie [49, 50] and Guichard and Wienhard [39, Theorem 5.3,5.9]). However, projective Anosov representations need not be irreducible and the images of elements need not be diagonalizable over \mathbb{R} . On the other hand, Guichard and Wienhard [39, Proposition 4.10] showed that any irreducible representation with transverse projective limits maps to projective Anosov.

6.2. *Deformation spaces.* The space of all projective Anosov representations of a fixed word hyperbolic group Γ into $SL_d(\mathbb{R})$ is an open subset of $Hom(\Gamma, SL_d(\mathbb{R}))/SL_d(\mathbb{R})$ (see Labourie [49, Proposition 2.1] and Guichard and Wienhard [39, Theorem 5.13]). However, a projective Anosov representation need not be a smooth point of $Hom(\Gamma, SL_d(\mathbb{R}))/SL_d(\mathbb{R})$ (see Johnson and Millson [45]). Moreover, the set of projective Anosov representations need not be an entire component of $Hom(\Gamma, SL_d(\mathbb{R}))/SL_d(\mathbb{R})$.

In order to have the structure of a real analytic manifold, we consider the space $\tilde{\mathcal{C}}(\Gamma, d)$ of regular, projective Anosov, irreducible representations $\rho : \Gamma \rightarrow SL_d(\mathbb{R})$ and let

$$\mathcal{C}(\Gamma, d) = \tilde{\mathcal{C}}(\Gamma, d)/SL_d(\mathbb{R}).$$

If \mathbf{G} is a reductive subgroup of $SL_d(\mathbb{R})$, we can restrict the whole discussion to representations with image in \mathbf{G} , i.e. let $\tilde{\mathcal{C}}(\Gamma, \mathbf{G})$ be the space of regular, projective Anosov, irreducible representations $\rho : \Gamma \rightarrow \mathbf{G}$ and let

$$\mathcal{C}(\Gamma, \mathbf{G}) = \tilde{\mathcal{C}}(\Gamma, \mathbf{G})/\mathbf{G}.$$

We will later want to restrict to the space $\mathcal{C}_g(\Gamma, \mathbf{G})$ of \mathbf{G} -generic representation in $\mathcal{C}(\Gamma, \mathbf{G})$, i.e. representations such that the centralizer of some element in the image is a maximal torus in \mathbf{G} . In particular, in the case that $\mathbf{G} = SL_d(\mathbb{R})$, a representation is \mathbf{G} -generic if some element in the image is diagonalizable over \mathbb{C} with distinct eigenvalues. The resulting spaces are real analytic manifolds.

PROPOSITION 6.1. [17, Proposition 7.1] *If Γ is a word hyperbolic group and \mathbf{G} is a reductive subgroup of $SL_d(\mathbb{R})$, then $\mathcal{C}(\Gamma, d)$, $\mathcal{C}(\Gamma, \mathbf{G})$, $\mathcal{C}_g(\Gamma, \mathbf{G})$ and $\mathcal{C}_g(\Gamma, d) = \mathcal{C}_g(\Gamma, SL_d(\mathbb{R}))$ are all real analytic manifolds.*

6.3. *The geodesic flow, entropy and intersection number.* One new difficulty which arises, is that it is not known in general whether or not the geodesic flow of a word hyperbolic group is metric Anosov, i.e. a Smale flow in the sense of Pollicott [66]. Notice that our construction in §5.2 immediately generalizes to give, for any projective Anosov representation ρ , a geodesic flow U_ρ which is Hölder orbit equivalent to U_Γ and whose periods are exactly spectral radii of infinite order elements of Γ . In general, we must further show [17, Proposition 5.1] that U_ρ is a topologically transitive metric Anosov flow.

PROPOSITION 6.2. [17, Propositions 4.1, 5.1] *If $\rho : \Gamma \rightarrow SL_d(\mathbb{R})$ is projective Anosov, then there exists a topologically transitive, metric Anosov flow U_ρ which is Hölder orbit equivalent to U_Γ such that the orbit associated to $\gamma \in \Gamma$ has period $\Lambda(\rho(\gamma))$.*

Lemma 2.2 provides a Hölder function $f_\rho : U_\Gamma \rightarrow (0, \infty)$, well defined up to Livšic cohomology, such that U_ρ is Hölder conjugate to the reparametrization of U_Γ by f_ρ .

One may then use the thermodynamic formalism to define the entropy of a projective Anosov representation and the intersection number and renormalized intersection number of two projective Anosov representations. If ρ is projective Anosov, we define

$$R_T(\rho) = \{[\gamma] \in [\pi_1(S)] \mid \log(\Lambda_\gamma(\rho)) \leq T\}$$

and the entropy of ρ is given by

$$h(\rho) = h(f_\rho) = \lim_{T \rightarrow \infty} \frac{\log \#R_T(\rho)}{T}.$$

The intersection number of two projective Anosov representations ρ and η is given by

$$\mathbf{I}(\rho, \eta) = \mathbf{I}(f_\rho, f_\eta) = \lim_{T \rightarrow \infty} \frac{1}{\#R_T(\rho)} \sum_{[\gamma] \in R_\rho(T)} \frac{\log(\Lambda_\gamma(\eta))}{\log(\Lambda_\gamma(\rho))}$$

and their renormalized intersection number is

$$\mathbf{J}(\rho, \eta) = \frac{h(\eta)}{h(\rho)} \mathbf{I}(\rho, \eta).$$

One may use the technique of proof of Proposition 5.9 to show that all these quantities vary analytically.

THEOREM 6.3. [17, Theorem 1.3] *If Γ is a word hyperbolic group and \mathbf{G} is a reductive subgroup of $\mathrm{PSL}_d(\mathbb{R})$, then entropy varies analytically over $\mathcal{C}(\Gamma, \mathbf{G})$ and intersection number and renormalized intersection number vary analytically over $\mathcal{C}(\Gamma, \mathbf{G}) \times \mathcal{C}(\Gamma, \mathbf{G})$.*

6.4. *The pressure metric for projective Anosov representation spaces.* If \mathbf{G} is a reductive subgroup of $\mathrm{PSL}_d(\mathbb{R})$, we define a thermodynamic mapping

$$\Phi : \mathcal{C}(\Gamma, \mathbf{G}) \rightarrow \mathcal{H}(U_\Gamma)$$

by $\rho \mapsto [-h(f_\rho) f_\rho]$. We can again show that Φ has locally analytic lifts, so we can pull back the pressure norm on $\mathcal{P}(U_\Gamma)$ to obtain a pressure semi-norm $\|\cdot\|_{\mathbf{P}}$ on $\mathcal{C}(\Gamma, \mathbf{G})$. The resulting pressure semi-norm gives an analytic Riemannian metric on $\mathcal{C}_g(\Gamma, \mathbf{G})$.

THEOREM 6.4. [17, Theorem 1.4] *If Γ is a word hyperbolic group and \mathbf{G} is a reductive subgroup of $\mathrm{SL}_d(\mathbb{R})$, then the pressure form is an analytic $\mathrm{Out}(\Gamma)$ -invariant Riemannian metric on $\mathcal{C}_g(\Gamma, \mathbf{G})$. In particular, the pressure form is an analytic $\mathrm{Out}(\Gamma)$ -invariant Riemannian metric on $\mathcal{C}_g(\Gamma, d)$.*

It only remains to prove that the pressure semi-norm is non-degenerate. We follow the same outline as in the Hitchin setting, but encounter significant new technical difficulties. As before, we may use Corollary 2.5 to obtain restrictions on the derivatives of spectral length of group elements.

LEMMA 6.5. [17, Lemma 9.3] *If \mathbf{G} is a reductive subgroup of $\mathrm{PSL}_d(\mathbb{R})$, $v \in T_\rho \mathcal{C}(\Gamma, \mathbf{G})$ and $\|v\|_{\mathbf{P}} = 0$, then*

$$D_\rho \log \Lambda_\gamma(v) = -\frac{D_\rho h(v)}{h(\rho)} \log \Lambda_\gamma(\rho)$$

for all $\gamma \in \Gamma$.

We use this to establish the following analogue of Proposition 5.13 from the Hitchin setting. In order to do so, we must work in the setting of \mathbf{G} -generic representations and we can only conclude that the derivative of spectral length, rather than trace, is trivial.

PROPOSITION 6.6. [17, Proposition 9.1] *If G is a reductive subgroup of $PSL_d(\mathbb{R})$, $v \in T_\rho C_g(\Gamma, G)$ and there exists K such that*

$$D_\rho \log \Lambda_\gamma(v) = K \log \Lambda_\gamma(\rho)$$

for all $\gamma \in \Gamma$, then $K = 0$. In particular, $D_\rho \log \Lambda_\gamma(v) = 0$ for all $\gamma \in \Gamma$.

One completes the proof by showing that the derivatives of the spectral radii functions generate the cotangent space.

PROPOSITION 6.7. [17, Proposition 10.3] *If G is a reductive subgroup of $PSL_d(\mathbb{R})$ and $\rho \in C_g(\Gamma, G)$, then the set $\{D_\rho \Lambda_\gamma \mid \gamma \in \Gamma\}$ spans $T_\rho^* C_g(\Gamma, G)$.*

6.5. *Anosov representations.* We now discuss the generalizations of our work to spaces of more general Anosov representations. If G is any semisimple Lie group with finite center and P^\pm is a pair of opposite parabolic subgroups, then one may consider (G, P^\pm) -Anosov representations of a word hyperbolic group Γ into G . A (G, P^\pm) -Anosov representation $\rho : \Gamma \rightarrow G$ has limit maps

$$\xi_\rho^\pm : \partial_\infty \Gamma \rightarrow G/P^\pm$$

(which are transverse in an appropriate sense and give rise to associated flows with contracting/dilating properties). In fact, Zariski dense representations with transverse limit maps are always (G, P^\pm) -Anosov [39, Theorem 4.11].

Projective Anosov representations are (G, P^\pm) -Anosov where $G = SL_d(\mathbb{R})$, P^+ is the stabilizer of a line and P^- is the stabilizer of a complementary hyperplane [17, Propostion 2.11]. Hitchin representations are (G, P^\pm) -Anosov where $G = SL_d(\mathbb{R})$, P^+ is the group of upper triangular matrices (i.e. the stabilizer of the standard flag) and P^- is the group of lower triangular matrices (Labourie [49]).

We may think of Anosov representations as natural generalizations of Fuchsian representations, since they are discrete, faithful, quasi-isometric embeddings with finite kernel so that the image of every infinite order element is P^+ -proximal [49, 50] and [39, Theorem 5.3,5.9]. More generally, they may be thought of as generalizations of convex cocompact representations into rank-one Lie groups. See Labourie [49] and Guichard and Wienhard [39] for definitions and more detailed discussions of Anosov representations. Gueritaud *et al* [37] and Kapovich *et al* [46] have developed intriguing new viewpoints on Anosov representations and their definition.

Guichard and Wienhard [39, Proposition 4.2, Remark 4.12] (see also [17, Theorem 2.12]) showed that there exists an irreducible representation $\sigma : G \rightarrow SL(V)$ (called the *Plücker representation*) such that $\rho : \Gamma \rightarrow G$ is (G, P^\pm) -Anosov if and only if $\sigma \circ \rho$ is projective Anosov. Thus, one can often reduce the study of (G, P^\pm) -Anosov representations to the study of projective Anosov representations.

Let $\mathcal{Z}(\Gamma, G, P^\pm)$ be the space of (conjugacy classes of) regular, virtually Zariski dense (G, P^\pm) -Anosov representations. The space $\mathcal{Z}(\Gamma, G, P^\pm)$ is an analytic orbifold, which is a manifold if G is connected (see [17, Proposition 7.3]). The Plücker representation $\sigma : G \rightarrow SL_d(\mathbb{R})$ allows one to view $\mathcal{Z}(\Gamma, G, P^\pm)$ as an analytically varying family of $\sigma(G)$ -generic projective Anosov representations. One may pull back the pressure form and adapt the techniques from the projective Anosov setting to prove the following.

THEOREM 6.8. [17, Corollary 1.9] *If \mathbf{G} is semi-simple Lie group with finite center and Γ is word hyperbolic, then the pressure form is an $\text{Out}(\Gamma)$ -invariant analytic Riemannian metric on $\mathcal{Z}(\Gamma, \mathbf{G}, \mathbf{P}^\pm)$.*

6.6. Examples. There are two other important classes of higher Teichmüller spaces which are (quotients of) entire components of representation varieties.

Burger *et al* [21] have studied representations of $\pi_1(S)$ into a Hermitian Lie group \mathbf{G} of tube type with maximal Toledo invariant, i.e. maximal representations. Each maximal representation is Anosov, with respect to stabilizers of points in the Shilov boundary of the associated symmetric space [20, 22], and the space of all maximal representations is a collection of components of $\text{Hom}(\pi_1(S), \mathbf{G})$ [21]. One particularly nice case arises when $\mathbf{G} = \text{Sp}(4, \mathbb{R})$, in which case there are $2g - 3$ components which are non-simply connected manifolds consisting entirely of Zariski dense representations (see Bradlow *et al* [13]). Hence, the quotients by \mathbf{G} of all such components admit pressure metrics.

Benoist [5, 6] studied holonomies of strictly convex projective structures on a closed manifold M and showed that these consist of entire components of $\text{Hom}(\pi_1(M), \text{PSL}_d(\mathbb{R}))$. One may use his work to show that these representations, which we call Benoist representations, are projective Anosov (see Guichard and Wienhard [39, Proposition 6.1]). Johnson and Millson [45] gave examples of holonomy maps $\rho : \pi_1(M) \rightarrow \text{SO}(d - 1, 1)$ of closed hyperbolic $d - 1$ -manifold, where $d \geq 5$, such that ρ is a singular point of $\text{Hom}(\pi_1(M), \text{PSL}_d(\mathbb{R}))$.

6.7. Rank-one Lie groups. Let Γ be a word hyperbolic group and let \mathbf{G} be a rank-one semi-simple Lie group, e.g. $\text{PSL}_2(\mathbb{C})$. A representation $\rho : \Gamma \rightarrow \mathbf{G}$ is *convex cocompact* if and only if whenever one chooses a basepoint x_0 for the symmetric space $X = \mathbf{K} \backslash \mathbf{G}$ then the orbit map $\tau : \Gamma \rightarrow X$ given by $\gamma \rightarrow \gamma(x_0)$ is a quasi-isometric embedding. The limit set of $\rho(\Gamma)$ is then the set of accumulation points in $\partial_\infty X$ of the image of the orbit map and one can define the Hausdorff dimension of this set. Patterson [65], Sullivan [78], Corlette and Iozzi [26], and Yue [90] showed that the topological entropy of a convex cocompact representation agrees with the Hausdorff dimension of the limit set of its image.

A representation $\rho : \Gamma \rightarrow \mathbf{G}$ is convex cocompact if and only if it is Anosov (see Guichard and Wienhard [39, Theorem 5.15]). Since the Plücker embedding multiplies entropy by a constant depending only on \mathbf{G} (see [17, Corollary 2.14]), the analyticity of the Hausdorff dimension of the limit set follows from the analyticity of entropy for projective Anosov representations.

THEOREM 6.9. [17, Corollary 1.8] *If Γ is a word hyperbolic group and \mathbf{G} is a rank-one semi-simple Lie group, then the Hausdorff dimension of the limit set varies analytically over analytic families of convex cocompact representations of Γ into \mathbf{G} .*

Remark. When $\mathbf{G} = \text{PSL}_2(\mathbb{C})$, Ruelle [71] proved this for surface groups and Anderson and Rocha [2] proved it for free products of surface groups and free groups. Tapie [79] used work of Katok *et al* [48] to show that the Hausdorff dimension is C^1 on smooth families of convex cocompact representations.

Let $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ be the space of (conjugacy classes of) convex cocompact representation of Γ into $\mathrm{PSL}_2(\mathbb{C})$. Bers [7] showed that $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ is an analytic manifold. Recall that a convex cocompact representation is not Zariski dense if and only if it is virtually Fuchsian, i.e. contain a finite index subgroup conjugate into $\mathrm{PSL}_2(\mathbb{R})$. We may again use the Plücker representation to prove the following.

THEOREM 6.10. [17, Corollary 1.7] *If Γ is word hyperbolic, then the pressure form is $\mathrm{Out}(\Gamma)$ -invariant and analytic on $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ and is non-degenerate at any representation which is not virtually Fuchsian. In particular, if Γ is not either virtually free or virtually a surface group, then the pressure form is an analytic Riemannian metric on $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$. Moreover, the pressure form always induces a path metric on $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$.*

Bridgeman [14] had previously defined and studied the pressure metric on quasi-Fuchsian space $QF(S) = CC(\pi_1(S), \mathrm{PSL}_2(\mathbb{C}))$. He showed that the degenerate vectors in this case correspond exactly to pure bending vectors on the Fuchsian locus.

6.8. Margulis space times. A Margulis space time is a quotient of \mathbb{R}^3 by a free, non-abelian group of affine transformations which acts properly discontinuously on \mathbb{R}^3 . They were originally discovered by Margulis [57] as counterexamples to a question of Milnor. Ghosh [32] used work of Goldman, Labourie and Margulis [34, 35] to interpret holonomy maps of Margulis space times (without cusps) as ‘Anosov representations’ into the (non-semi-simple) Lie group $\mathrm{Aff}(\mathbb{R}^3)$ of affine automorphisms of \mathbb{R}^3 . Ghosh [33] was then able to adapt the techniques of [17] to produce a pressure form on the analytic manifold \mathcal{M} of (conjugacy classes of) holonomy maps of Margulis space times of fixed rank (with no cusps). This pressure form is an analytic Riemannian metric on the slice \mathcal{M}_k of \mathcal{M} consisting of holonomy maps with entropy k (see Ghosh [33, Theorem 1.0.1]), but has a degenerate direction on \mathcal{M} , so the pressure form has signature $(\dim \mathcal{M} - 1, 0)$ on \mathcal{M} .

7. Open problems

The geometry of the pressure metric is still rather mysterious and much remains to be explored. The hope is that the geometry of the pressure metric will yield insights into the nature of the Hitchin component and other higher Teichmüller spaces, in much the way that the study of the Teichmüller and Weil–Petersson metrics have been an important tool in our understanding of Teichmüller space and the mapping class group. It is natural to begin by exploring analogies with the Weil–Petersson metric on Teichmüller space. We begin the discussion by recalling some basic properties of the Weil–Petersson metric.

Properties of the Weil–Petersson metric.

- (1) The extended mapping class group is the isometry group of $\mathcal{T}(S)$ in the Weil–Petersson metric (Masur and Wolf [59]).
- (2) The Weil–Petersson metric is negatively curved, but the sectional curvature is not bounded away from either 0 or $-\infty$ (Wolpert [86], Tromba [82], Huang [42]).
- (3) If ϕ is a pseudo-Anosov mapping class, then there is a lower bound for its translation distance on Teichmüller space and there is a unique invariant geodesic axis for ϕ (Daskalopoulos and Wentworth [30]).

- (4) The Weil–Petersson metric is incomplete (Wolpert [85], Chu [25]). However, it admits a metric completion which is $CAT(0)$ and homeomorphic to the augmented Teichmüller space (see Masur [58] and Wolpert [88]).

Masur and Wolf’s result [59] on the isometry group of $\mathcal{T}(S)$ suggests the following problem.

Problem 1. Is the isometry group of a Hitchin component generated by the (extended) mapping class group and the contragredient involution? More generally, explore whether the relevant outer automorphism group is a finite index subgroup of the isometry group of a higher Teichmüller space with the pressure metric.

Bridgeman et al [16] have shown that any diffeomorphism of $\mathcal{H}_3(S)$ which preserves the intersection number is an element of the extended mapping class group or the composition of an element in the extended mapping class group with the contragredient involution. Along the way, they show that any diffeomorphism which preserves the intersection number also preserves the entropy and hence preserves the renormalized intersection number, the pressure metric, and, by work of Potrie and Sambarino [69], the Fuchsian locus. This suggests the following problem.

Problem 2. Prove that if $g : \mathcal{H}_d(S) \rightarrow \mathcal{H}_d(S)$ is an isometry with respect to the pressure metric then $\mathbf{I}(g(\rho), g(\sigma)) = \mathbf{I}(\rho, \sigma)$ for all $\rho, \sigma \in \mathcal{H}_d(S)$. It would follow that the isometry of the group of the Hitchin component $\mathcal{H}_3(S)$ is generated by the extended mapping class group of S and the contragredient involution.

Bridgeman and Canary [15] have shown that the group of diffeomorphisms of quasi-Fuchsian space $QF(S)$ which preserve the renormalized intersection number is generated by the extended mapping class group and complex conjugation. So one may also consider the corresponding analogue of Problem 2 in quasi-Fuchsian space.

It would be useful to study the curvature of the pressure metric, guided by the results of Wolpert [86], Tromba [82], and Huang [42]. Wolf’s work [84] (see also [83]) on the Hessians of length functions on Teichmüller space may offer a plan of attack here.

Problem 3. Investigate the curvature of the Hitchin component in the pressure metric.

Pollicott and Sharp [68] have investigated the curvature of the pressure metric on deformation spaces of marked metric graphs with entropy 1. In this setting, the curvature can be both positive and negative.

Labourie and Wentworth [51] have derived a formula for the pressure metric at points in the Fuchsian locus of a Hitchin component in terms of Hitchin’s parametrization of the Hitchin component by holomorphic differentials. They also obtain variational formulas which are analogues of classical results in the Teichmüller setting.

Since Labourie [50] proved that the mapping class group acts properly discontinuously on a Hitchin component, it is natural to study the geometry of this action. One specific question to start with would be the following problem.

Problem 4. Is there a lower bound for the translation distance for the action of a pseudo-Anosov mapping class on the Hitchin component?

Since the restriction of the pressure metric to the Fuchsian locus is a multiple of the Weil–Peterson metric, the Hitchin component is incomplete and the metric completion contains augmented Teichmüller space. However, very little is known about the completeness of the pressure metric in ‘other directions.’ The work of Zhang [91, 92] and Loftin [54] (when $d = 3$) should be relevant here. It may also be interesting to study the relationship between the metric completion and Parreau’s compactification [63] of the Hitchin component.

Problem 5. Investigate the metric completion of the Hitchin component or other higher Teichmüller spaces.

Xu [89] studied the pressure metric on the Teichmüller space $\mathcal{T}(S)$ where S is a surface with non-empty geodesic boundary. He shows that the pressure metric in this case is not equal to the classical Weil–Peterson metric on $\mathcal{T}(S)$. He further shows that it is not complete and that the space of marked metric graphs on a fixed graph with its pressure metric arises naturally in the completion.

The following problem indicates how little is known about the coarse geometry of the pressure metric. We recall that a subset A of a metric space X is said to be *coarsely dense* if there exists $D > 0$ such that every point in X lies within D of a point in A .

Problem 6. (a) Is the Fuchsian locus coarsely dense in a Hitchin component?

(b) Is the Fuchsian locus coarsely dense in quasi-Fuchsian space?

(c) If M is an acylindrical 3-manifold with no toroidal boundary components and $\Gamma = \pi_1(M)$, does $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ have finite diameter?

Zhang [91, 92] and Nie [62] (when $d = 3$) produce sequences in Hitchin components where entropy converges to 0. These sequences are candidates to produce points arbitrarily far from the Fuchsian locus.

In case (c), $\mathrm{Out}(\Gamma)$ is finite (see Johannson [44]) and $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ has compact closure in the $\mathrm{PSL}_2(\mathbb{C})$ -character variety (see Thurston [81]).

One may phrase all the above questions as being about the quotient of a higher Teichmüller space by its natural automorphism group. Similarly, one might ask whether the quotient of the Hitchin component by the mapping class group has finite volume.

Problem 7. Does the quotient of the Hitchin component by the action of the mapping class group have finite volume in the quotient pressure metric?

Potrie and Sambarino [69] showed that the entropy function is maximal uniquely on the Fuchsian locus of a Hitchin component, so it is natural to investigate more subtle behavior of the entropy function.

Problem 8. Investigate the critical points on the entropy function.

Bowen [11] showed that the entropy function is uniquely minimal on the Fuchsian locus in quasi-Fuchsian space $QF(S)$. Bridgeman [14] showed that the entropy function on $QF(S)$ has no local maxima and moreover the Hessian of the entropy function is positive-definite on at least a half-dimensional subspace at any critical point.

If M is an acylindrical 3-manifold with no toroidal boundary components and $\Gamma = \pi_1(M)$, then there is a unique representation in $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ where the boundary of

the limit set of the image consists of round circles (see Thurston [80]). It is conjectured that the entropy has a unique minimum at this representation (see Canary *et al* [23]). Storm [77] proved that this is the unique representation where the volume of the convex core achieves its minimum.

In the case of $CC(\Gamma, \mathrm{PSL}_2(\mathbb{C}))$ we were able to obtain a path metric, even when the pressure form is degenerate on a submanifold. One might hope to be able to do so in more general settings.

Problem 9. If Γ is a word hyperbolic group, G is a semisimple Lie group and P^\pm is a pair of opposite parabolic subgroups, can one extend the pressure metric on $\mathcal{Z}(\Gamma, G, P^\pm)$ to a path metric on the space of all (conjugacy classes of) (G, P^\pm) -Anosov representations of Γ into G ?

Acknowledgements. The paper is based on a master class given by the authors at the Centre for Quantum Geometry of Moduli Spaces in Aarhus. We thank Jorgen Anderson for the invitation to give this master class and the editors for the invitation to write this article. We thank Marc Burger, François Labourie and Adam Sikora for helpful conversations. Substantial portions of this paper were written while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, CA, during the Spring 2015 semester and were partially supported by NSF grant No. 0932078 000. Bridgeman was partially supported by NSF grant DMS—1500545, Canary was partially supported by NSF grant DMS—1306992 and Sambarino was partially supported by the European Research Council under the *European Community's* seventh Framework Programme (FP7/2007–2013)/ERC *grant agreement* n° FP7-246918. The authors also acknowledge support from US National Science Foundation grants DMS 1107452, 1107263, 1107367 ‘RNMS: GEometric structures And Representation varieties’ (the GEAR Network).

REFERENCES

- [1] L. Ahlfors and A. Beurling. The boundary correspondence under quasiconformal mappings. *Acta Math.* **96** (1956), 125–142.
- [2] J. W. Anderson and A. Rocha. Analyticity of Hausdorff dimension of limit sets of Kleinian groups. *Ann. Acad. Sci. Fenn. Math.* **22** (1997), 349–364.
- [3] Y. Benoist. Propriétés asymptotiques des groupes linéaires. *Geom. Funct. Anal.* **7** (1997), 1–47.
- [4] Y. Benoist. Propriétés asymptotiques des groupes linéaires II. *Adv. Stud. Pure Math.* **26** (2000), 33–48.
- [5] Y. Benoist. Convexes divisibles I. *Algebraic Groups and Arithmetic (Tata Institute of Fundamental Research Studies in Mathematics, 17)*. TIFR Publications, Mumbai, 2004, pp. 339–374.
- [6] Y. Benoist. Convexes divisibles III. *Ann. Sci. Éc. Norm. Supér. (4)* **38** (2005), 793–832.
- [7] L. Bers. Spaces of Kleinian groups. *Maryland Conference in Several Complex Variables I (Lecture Notes in Mathematics, 155)*. Springer, Berlin, 1970, pp. 9–34.
- [8] F. Bonahon. The geometry of Teichmüller space via geodesic currents. *Invent. Math.* **92** (1988), 139–162.
- [9] R. Bowen. Periodic orbits of hyperbolic flows. *Amer. J. Math.* **94** (1972), 1–30.
- [10] R. Bowen. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms (Lecture Notes in Mathematics, 470)*. Springer, Berlin, 1975.
- [11] R. Bowen. Hausdorff dimension of quasi-circles. *Publ. Math. Inst. Hautes Études Sci.* **50** (1979), 11–25.
- [12] R. Bowen and D. Ruelle. The ergodic theory of axiom A flows. *Invent. Math.* **29** (1975), 181–202.
- [13] S. Bradlow, O. Garcia-Prada and P. Gothen. Deformations of maximal representations in $\mathrm{Sp}(4, \mathbb{R})$. *Q. J. Math.* **63** (2012), 795–843.

- [14] M. Bridgeman. Hausdorff dimension and the Weil–Petersson extension to quasifuchsian space. *Geom. Topol.* **14** (2010), 799–831.
- [15] M. Bridgeman and R. Canary. Simple length rigidity for Kleinian surface groups and applications. *Preprint*, 2015, arXiv:1509.02510.
- [16] M. Bridgeman, R. Canary and F. Labourie. Simple length rigidity for Hitchin representations. In preparation.
- [17] M. Bridgeman, R. Canary, F. Labourie and A. Sambarino. The pressure metric for Anosov representations. *Geom. Funct. Anal.* **25** (2015), 1089–1179.
- [18] M. Bridgeman and E. Taylor. An extension of the Weil–Petersson metric to quasi-fuchsian space. *Math. Ann.* **341** (2008), 927–943.
- [19] M. Burger. Intersection, the Manhattan curve and Patterson–Sullivan theory in rank 2. *Int. Math. Res. Not. IMRN* **7** (1993), 217–225.
- [20] M. Burger, A. Iozzi, F. Labourie and A. Wienhard. Maximal representations of surface groups: symplectic Anosov structures. *Pure Appl. Math. Q.* **1** (2005), 555–601.
- [21] M. Burger, A. Iozzi and A. Wienhard. Surface group representations with maximal Toledo invariant. *Ann. of Math. (2)* **172** (2010), 517–566.
- [22] M. Burger, A. Iozzi and A. Wienhard. Maximal representations and Anosov structures. In preparation.
- [23] R. D. Canary, Y. N. Minsky and E. C. Taylor. Spectral theory, Hausdorff dimension and the topology of hyperbolic 3-manifolds. *J. Geom. Anal.* **9** (1999), 17–40.
- [24] C. Champetier. Petite simplification dans les groupes hyperboliques. *Ann. Fac. Sci. Toulouse Math. (6)* **3** (1994), 161–221.
- [25] T. Chu. The Weil–Petersson metric in moduli space. *Chinese J. Math.* **4** (1976), 29–51.
- [26] K. Corlette and A. Iozzi. Limit sets of discrete groups of isometries of exotic hyperbolic spaces. *Trans. Amer. Math. Soc.* **351** (1999), 1507–1530.
- [27] M. Crampon. Entropies of compact strictly convex projective manifolds. *J. Mod. Dynam.* **3** (2009), 511–547.
- [28] F. Dal’Bo and I. Kim. A criterion of conjugacy for Zariski dense subgroups. *C. R. Acad. Sci. Paris Sér. I Math.* **330** (2000), 647–650.
- [29] M. Darvishzadeh and W. Goldman. Deformation spaces of convex real projective structures and hyperbolic structures. *J. Korean Math. Soc.* **33** (1996), 625–639.
- [30] G. Daskalopoulos and R. Wentworth. Classification of Weil–Petersson isometries. *Amer. J. Math.* **125** (2003), 941–975.
- [31] V. Fock and A. Goncharov. Moduli spaces of local systems and higher Teichmüller theory. *Publ. Math. Inst. Hautes Études Sci.* **103** (2006), 1–211.
- [32] S. Ghosh. Anosov structure on Margulis space time. *Preprint*, available at <http://front.math.ucdavis.edu/1412.8211>.
- [33] S. Ghosh. The pressure metric on the Margulis multiverse. *Geom. Dedicata*, accepted. *Preprint*, available at <http://front.math.ucdavis.edu/1505.00534>.
- [34] W. Goldman and F. Labourie. Geodesics in Margulis spacetimes. *Ergod. Th. & Dynam. Sys.* **32** (2012), 643–651.
- [35] W. Goldman, F. Labourie and G. Margulis. Proper affine actions and geodesic flows of hyperbolic surfaces. *Ann. of Math. (2)* **170** (2009), 1051–1083.
- [36] M. Gromov. Hyperbolic groups. *Essays in Group Theory (MSRI Publications, 8)*. Springer, New York, 1987, pp. 75–263.
- [37] F. Gueritaud, O. Guichard, F. Kassel and A. Wienhard. Anosov representations and proper actions. *Geom. Topology*, accepted. *Preprint*, available at <http://front.math.ucdavis.edu/1502.03811>.
- [38] O. Guichard. In preparation.
- [39] O. Guichard and A. Wienhard. Anosov representations: domains of discontinuity and applications. *Invent. Math.* **190** (2012), 357–438.
- [40] M. W. Hirsch, C. C. Pugh and M. Shub. *Invariant Manifolds (Lecture Notes in Mathematics, 583)*. Springer, Berlin, 1977.
- [41] N. Hitchin. Lie groups and Teichmüller space. *Topology* **31** (1992), 449–473.
- [42] Z. Huang. Asymptotic flatness of the Weil–Petersson metric on Teichmüller space. *Geom. Dedicata* **110** (2005), 81–102.
- [43] J. Humphreys. *Introduction to Lie Algebras and Representation Theory*. Springer, Berlin, 1972.
- [44] K. Johannson. *Homotopy Equivalences of 3-manifolds with Boundary (Lecture Notes in Mathematics, 761)*. Springer, Berlin, 1979.
- [45] D. Johnson and J. Millson. Deformation spaces associated to compact hyperbolic manifolds. *Discrete Groups and Geometric Analysis (Progress in Mathematics, 67)*. Birkhäuser, Boston, 1987, pp. 48–106.

- [46] M. Kapovich, B. Leeb and J. Porti. Morse actions of discrete groups on symmetric spaces. *Preprint*, available at <http://front.math.ucdavis.edu/1403.7671>.
- [47] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge, 1995.
- [48] A. Katok, G. Knieper, M. Pollicott and H. Weiss. Differentiability and analyticity of topological entropy for Anosov and geodesic flows. *Invent. Math.* **98** (1989), 581–597.
- [49] F. Labourie. Anosov flows, surface groups and curves in projective space. *Invent. Math.* **165** (2006), 51–114.
- [50] F. Labourie. Cross ratios, Anosov representations and the energy functional on Teichmüller space. *Ann. Sci. Éc. Norm. Supér. (4)* **41** (2008), 437–469.
- [51] F. Labourie and R. Wentworth. Variations along the fuchsian locus. *Preprint*, available at <http://arxiv.org/1506.01686>.
- [52] Q. Li. Teichmüller space is totally geodesic in Goldman space. *Asian J. Math.* **20** (2016), 21–46.
- [53] A. N. Livšic. Cohomology of dynamical systems. *Math. USSR Izv.* **6** (1972), 1296–1320.
- [54] J. Loftin. The compactification of the moduli space of convex \mathbb{RP}^2 surfaces. I. *J. Differential Geom.* **68** (2004), 223–276.
- [55] A. Lubotzky and A. Magid. Varieties of representations of finitely generated groups. *Mem. Amer. Math. Soc.* **58** (1985), no. 336.
- [56] A. Manning. Topological entropy for geodesic flows. *Ann. of Math. (2)* **110** (1979), 567–573.
- [57] G. Margulis. Free completely discontinuous groups of affine transformations. *Sov. Math. Dokl.* **28** (1983), 435–439.
- [58] H. Masur. Extension of the Weil–Petersson metric to the boundary of Teichmüller space. *Duke Math. J.* **43** (1976), 623–635.
- [59] H. Masur and M. Wolf. The Weil–Petersson isometry group. *Geom. Dedicata* **93** (2002), 177–190.
- [60] C. McMullen. Thermodynamics, dimension and the Weil–Petersson metric. *Invent. Math.* **173** (2008), 365–425.
- [61] I. Mineyev. Flows and joins of metric spaces. *Geom. Topol.* **9** (2005), 403–482.
- [62] X. Nie. On the Hilbert geometry of simplicial Tits sets. *Ann. Inst. Fourier (Grenoble)* **65** (2015), 1005–1030.
- [63] A. Parreau. Compactification d’espaces de représentations de groupes de type fini. *Math. Z.* **272** (2012), 51–86.
- [64] W. Parry and M. Pollicott. Zeta functions and the periodic orbit structure of hyperbolic dynamics. *Astérisque* **187–188** (1990).
- [65] S. Patterson. The limit set of a Fuchsian group. *Acta Math.* **136** (1976), 241–273.
- [66] M. Pollicott. Symbolic dynamics for Smale flows. *Amer. J. Math.* **109** (1987), 183–200.
- [67] M. Pollicott and R. Sharp. Length asymptotics in higher Teichmüller theory. *Proc. Amer. Math. Soc.* **142** (2014), 101–112.
- [68] M. Pollicott and R. Sharp. A Weil–Petersson metric on spaces of metric graphs. *Geom. Dedicata* **172** (2014), 229–244.
- [69] R. Potrie and A. Sambarino. Eigenvalues and entropy of a Hitchin representation. *Preprint*, available at <http://front.math.ucdavis.edu/1411.5405>.
- [70] D. Ruelle. *Thermodynamic Formalism*. Addison-Wesley, London, 1978.
- [71] D. Ruelle. Repellers for real analytic maps. *Ergod. Th. & Dynam. Sys.* **2** (1982), 99–107.
- [72] A. Sambarino. Quantitative properties of convex representations. *Comment. Math. Helv.* **89** (2014), 443–488.
- [73] A. Sambarino. The orbital counting problem for hyperconvex representations. *Ann. Inst. Fourier (Grenoble)* **65** (2015), 1755–1797.
- [74] P. Schmutz. Die parametrisierung des Teichmüllerraumes durch geodätische Längenfunktionen. *Comment. Math. Helv.* **68** (1993), 278–288.
- [75] M. Shub. *Global Stability of Dynamical Systems*. Springer, New York, 1987.
- [76] Z. Ślodkowski. Holomorphic motions and polynomial hulls. *Proc. Amer. Math. Soc.* **111** (1991), 347–355.
- [77] P. Storm. Hyperbolic convex cores and simplicial volume. *Duke Math. J.* **140** (2007), 281–319.
- [78] D. Sullivan. The density at infinity of a discrete group of hyperbolic motions. *Publ. Math. Inst. Hautes Études Sci.* **50** (1979), 171–202.
- [79] S. Tapie. A variation formula for the topological entropy of convex-cocompact manifolds. *Ergod. Th. & Dynam. Sys.* **31** (2011), 1849–1864.
- [80] W. Thurston. Hyperbolic geometry and 3-manifolds. *Low-Dimensional Topology (Bangor, 1979)*. Cambridge University Press, Cambridge, 1982, pp. 9–25.
- [81] W. P. Thurston. Hyperbolic structures on 3-manifolds, I: deformation of acylindrical manifolds. *Ann. of Math. (2)* **124** (1986), 203–246.

- [82] A. Tromba. On a natural algebraic affine connection on the space of almost complex structures and the curvature of Teichmüller space with respect to its Weil–Petersson metric. *Manuscripta Math.* **56** (1986), 475–497.
- [83] M. Wolf. The Teichmüller theory of harmonic maps. *J. Differential Geom.* **29** (1989), 449–479.
- [84] M. Wolf. The Weil–Petersson Hessian of length on Teichmüller space. *J. Differential Geom.* **91** (2012), 129–169.
- [85] S. Wolpert. Noncompleteness of the Weil–Petersson metric for Teichmüller space. *Pacific J. Math.* **61** (1975), 573–577.
- [86] S. Wolpert. Chern forms and the Riemann tensor for the moduli space of curves. *Invent. Math.* **85** (1986), 119–145.
- [87] S. Wolpert. Thurston’s Riemannian metric for Teichmüller space. *J. Differential Geom.* **23** (1986), 143–174.
- [88] S. Wolpert. Geometry of the Weil–Petersson completion of Teichmüller space. *Surveys in Differential Geometry, Vol. VIII (Boston, MA, 2002)*. International Press of Boston, Boston, 2003, pp. 357–393.
- [89] B. Xu. Incompleteness of pressure metric on Teichmüller space of a bordered surface. *Preprint*, available at <http://front.math.ucdavis.edu/1608.03937>.
- [90] C. Yue. The ergodic theory of discrete isometry groups on manifolds of variable negative curvature. *Trans. Amer. Math. Soc.* **348** (1996), 4965–5005.
- [91] T. Zhang. The degeneration of convex \mathbb{RP}^2 structures on surfaces. *Proc. Lond. Math. Soc. (3)* **111** (2015), 967–1012.
- [92] T. Zhang. Degeneration of Hitchin sequences along internal sequences. *Geom. Funct. Anal.* **25** (2015), 1588–1645.