

# Geometrization of the local Langlands correspondence

Laurent Fargues (CNRS/IMJ) - Joint with Peter Scholze

## Local Langlands parameters

- ▶  $E$  local field residue field  $\mathbb{F}_q$ ,  $[E : \mathbb{Q}_p] < +\infty$  or  $E = \mathbb{F}_q((\pi))$
- ▶  $G$  reductive group over  $E$
- ▶  $\ell \neq p$ ,  $\Lambda \in \{\overline{\mathbb{F}}_\ell, \overline{\mathbb{Q}}_\ell\}$
- ▶  ${}^L G = \hat{G} \rtimes W_E$  Langlands dual over  $\Lambda$
- ▶  $\pi$  smooth representation of  $G(E)$  with coefficients in  $\Lambda$ , Schur irreducible i.e.  $\text{End}(\pi) = \Lambda$
- ▶ We construct

$$\varphi_\pi : W_E \rightarrow {}^L G$$

its semi-simple Langlands parameter.

- ▶ Compatible with parabolic induction and usual class field theory for tori. Usual local Langlands for  $\text{GL}_n$  (Harris-Taylor, Henniart)
- ▶ semi-simple :  $N = 0$  when  $\Lambda = \overline{\mathbb{Q}}_\ell$ . For example :  
 $\varphi_{\text{triv}} = \varphi_{\text{Steinberg}}$  for  $\text{GL}_n$

# Morphisms between centers

In fact we do much more.

- ▶ For  $\Lambda$  a  $\mathbb{Z}_\ell$ -algebra make it a **condensed ring** via

$$\Lambda := \Lambda \otimes_{\mathbb{Z}_\ell^{disc}} \mathbb{Z}_\ell$$

- ▶ There is a scheme  $/\mathbb{Z}_\ell$ ,  $\coprod_{\text{infinite}}$  affine schemes,

$$Z^1(W_E, \hat{G})$$

Value on  $\Lambda$  is condensed 1-cocycles  $W_E \rightarrow \hat{G}(\Lambda)$

- ▶ Studied in details by Dat-Helm-Kurinczuk-Moss
- ▶ Then

$$\text{LocSys}_{\hat{G}} := [Z^1(W_E, \hat{G})/\hat{G}]$$

is a **zero dimensional locally complete intersection algebraic stack**  $/\mathbb{Z}_\ell$ . Moduli of Langlands parameters.

## Morphisms between centers

- ▶ Coarse moduli space

$$Z^1(W_E, \hat{G}) // \hat{G}$$

$\coprod_{\text{infinite}}$  affine schemes finite type/ $\mathbb{Z}_\ell$ .

- ▶ Functions on it

$$\mathfrak{Z}^{\text{spec}}(G, \mathbb{Z}_\ell) = \mathcal{O}(Z^1(W_E, \hat{G}))^{\hat{G}}$$

- ▶ Example :  $G = \text{GL}_n$ ,  $\mathfrak{Z}^{\text{spec}}(G, \mathbb{Z}_\ell) \rightarrow \Lambda =$   
pseudo-representations  $W_E \rightarrow \text{GL}_n(\Lambda)$ .
- ▶  $\mathfrak{Z}(G(E), \mathbb{Z}_\ell) =$  Bernstein center = center of the category of  
smooth representations of  $G(E)$  with coefficients in  $\mathbb{Z}_\ell$
- ▶ We construct a morphism

$$\mathfrak{Z}^{\text{spec}}(G, \mathbb{Z}_\ell) \longrightarrow \mathfrak{Z}(G(E), \mathbb{Z}_\ell)$$

# The real deal : $\text{Bun}_G$

In fact we do much much more.

- ▶  $S$  an  $\overline{\mathbb{F}}_q$ -perfectoid space  $\rightsquigarrow X_S = E$ -adic space  
"the relative curve parametrized by  $S$ "
- ▶ i.e there is a way to put in family the collection of curves

$$(X_{k(s), k(s)^+})_{s \in S}$$

where  $X_{k(s), k(s)^+}$  is the curve defined and studied with Fontaine attached to the perfectoid field  $k(s)$

We will consider the  $v$ -topology on  $\overline{\mathbb{F}}_q$ -perfectoid spaces = some kind of analog of fpqc topology for schemes

$$* = \text{Spa}(\overline{\mathbb{F}}_q)$$

final object of the  $v$ -topos (not representable)

# $\text{Bun}_G$

## Theorem

The correspondence  $S \mapsto \{\text{principal } G\text{-bundles on } X_S\}$  defines a  $v$ -stack

$$\text{Bun}_G \longrightarrow *$$

that is an "Artin  $v$ -stack" ( $\ell$ -cohomologically) smooth of dimension 0.

- ▶ diagonal of  $\text{Bun}_G$  representable in locally spatial diamonds
- ▶ there is a surjection  $U \rightarrow \text{Bun}_G$  that is ( $\ell$ -coho.) smooth with  $U$  a locally spatial diamond s.t.  $U \rightarrow *$  is ( $\ell$ -coho.) smooth

## Bun<sub>G</sub> : points

- ▶ Set  $\check{E} = \widehat{E^{un}}$  with its Frobenius  $\sigma$ . One has

$$X_S = Y_S / \varphi^{\mathbb{Z}}$$

with  $Y_S \rightarrow \mathrm{Spa}(\check{E})$ ,  $\varphi$  = some Frobenius that extends  $\sigma$  on  $\check{E}$ .

- ▶ Functor

Isocrystals  $\longrightarrow$  vector bundles on  $X_S$

$$(D, \varphi) \longmapsto Y_S \times_{\varphi^{\mathbb{Z}}} D$$

- ▶  $B(G) = G(\check{E}) / \sigma$ -conjugation,  $b \sim gbg^{-\sigma}$ , **Kottwitz set** of  $G$ -isocrystals
- ▶  $b \in G(\check{E}) \rightsquigarrow \mathcal{E}_b$  principal  $G$ -bundle on  $X_S$

## $\text{Bun}_G$ : points

Theorem (Fargues-Fontaine ( $\text{GL}_n$ ), Fargues)

$F$  alg. closed

$$\begin{aligned} B(G) &\xrightarrow{\sim} H_{\text{ét}}^1(X_F, G) \\ [b] &\mapsto [\mathcal{E}_b] \end{aligned}$$

- ▶ Dictionary : **reduction theory** (Atiyah-Bott) for  $G$ -bundles / Kottwitz description of  $B(G)$ .
- ▶ *Example* :  $\mathcal{E}_b$  semi-stable  $\Leftrightarrow b$  is basic (isoclinic)

Thus, identification

$$B(G) = |\text{Bun}_G|$$



## Bun<sub>G</sub> : geometry

- ▶  $c_1 : \pi_0(\text{Bun}_G) \xrightarrow{\sim} \pi_1(G)_\Gamma$
- ▶ Nice Harder-Narasimhan stratification, in particular

$$\text{Bun}_G^{\text{ss}} \subset \text{Bun}_G \text{ is open}$$

Each connected component has a unique ss point and

$$\text{Bun}_G^{\text{ss}} = \coprod_{[b] \text{ basic}} \underbrace{[* / G_b(E)]}_{\text{classifying stack of pro-étale torsors}}$$

with  $G_b =$  inner form of  $G$  ( $G_1 = G$  for example)

- ▶ More generally for any  $[b] \in B(G)$  the associated HN strata is a classifying stack

$$[* / \tilde{G}_b]$$

with  $\tilde{G}_b = \tilde{G}_b^0 \times \underline{G_b(E)}$ ,  $\tilde{G}_b^0 =$  unipotent diamond  $G_b =$  inner form of a Levi

## The real deal : $D_{lis}(\text{Bun}_G, \Lambda)$

- ▶  $\Lambda$  any  $\mathbb{Z}_\ell$ -algebra
- ▶ We define a triangulated category

$$D_{lis}(\text{Bun}_G, \Lambda)$$

that is  $D_{\text{ét}}(\text{Bun}_G, \Lambda)$  when  $\Lambda$  is torsion and a sub-category of  $D_{\text{proét}}(\text{Bun}_G, \Lambda_{\blacksquare})$  in general

- ▶ For  $[b] \in B(G)$  inclusion of HN stratum

$$i^b : [* / \tilde{G}_b] \hookrightarrow \text{Bun}_G$$

induces

$$(i^b)^* : D_{lis}(\text{Bun}_G, \Lambda) \longrightarrow D_{lis}([* / \tilde{G}_b], \Lambda) \underbrace{=}_{\ell \neq p} D(G_b(E), \Lambda)$$

(derived category of smooth representations of  $G_b(E)$ )

## $D_{lis}(\text{Bun}_G, \Lambda)$

- ▶ In particular, via  $(i^1)_!$  and  $(i^1)^*$

$$D(G(E), \Lambda) \subset D_{lis}(\text{Bun}_G, \Lambda)$$

is a **direct factor**.

- ▶ Good object for the local Langlands program is not a smooth representation  $\pi$  or a complex in  $D(G(E), \Lambda)$  but an object of  $D_{lis}(\text{Bun}_G, \Lambda)$ !!! Have to think the local Langlands program from this point of view!
- ▶ Usual notions of **admissible, finite representations or Bernstein-Zelevinsky duality** extend to  $D_{lis}(\text{Bun}_G, \Lambda)$

# $D_{lis}(\text{Bun}_G, \Lambda)$

More precisely.

## Theorem

For  $A \in D_{lis}(\text{Bun}_G, \Lambda)$

1.  $A$  is *compact* iff it has finite support and for all  $[b] \in B(G)$ ,  $(i^b)^*A \in D(G(E), \Lambda)$  is compact (i.e. is bounded with *finite type* cohomology if  $\Lambda = \overline{\mathbb{Q}}_\ell$ )
2.  $A$  is *ULA* iff for all  $[b] \in B(G)$ ,  $(i^b)^*A \in D(G(E), \Lambda)$  is such that for all compact open  $K \subset G(E)$ ,  $((i^b)^*A)^K$  is a *perfect complex of  $\Lambda$ -modules*.
3. There is a duality functor  $\mathbb{D}_{BZ} : D_{lis}(\text{Bun}_G, \Lambda)^\omega \longrightarrow D_{lis}(\text{Bun}_G, \Lambda)^\omega$  that extends *Bernstein-Zelevinsky duality* on  $D^b(G(E), \Lambda)$

# The real deal : the spectral action

## Theorem

There is a monoidal action of  $\text{Perf}(\text{LocSys}_{\hat{G}})$  on  $D_{lis}(\text{Bun}_G, \mathbb{Z}_\ell)$ .

- ▶ This monoidal action defines the morphism between centers

$$\underbrace{\mathfrak{Z}^{spec}(G, \mathbb{Z}_\ell)}_{\text{spectral stable Bernstein center}} = \mathfrak{Z}(\text{Perf}(\text{LocSys}_{\hat{G}}))$$
$$\rightarrow \underbrace{\mathfrak{Z}(D_{lis}(\text{Bun}_G, \mathbb{Z}_\ell))}_{\text{geometric stable Bernstein center}} \rightarrow \underbrace{\mathfrak{Z}(G(E), \Lambda)}_{\text{Bernstein center}}$$

- ▶ Defined using some [geometric Satake correspondence](#) for sheaves of  $\Lambda$ -modules on the  $B_{dR}$ -affine Grassmanian + some enhanced version of Beilinson-Drinfeld/Vincent Lafforgue formalism (quantum field theory/factorization sheaves)

# The real deal : the geometrization conjecture

$G$  quasisplit. Fix  $\psi : U(E) \rightarrow \overline{\mathbb{Z}}_\ell$  non-degenerate. Let

$$\mathcal{W}_\psi = (i^1)_! (c\text{-ind}_{U(E)}^{G(E)} \psi) \in D_{lis}(\text{Bun}_G, \overline{\mathbb{Z}}_\ell)$$

be the "Whittaker sheaf".

## Conjecture

*The functor*

$$\begin{aligned} \text{Perf}(\text{LocSys}_{\hat{G}}/\overline{\mathbb{Z}}_\ell) &\longrightarrow D_{lis}(\text{Bun}_G, \overline{\mathbb{Z}}_\ell) \\ \mathcal{F} &\longmapsto \mathcal{F} * \mathcal{W}_\psi \end{aligned}$$

*extends to an equivalence compatible with the spectral action*

$$\text{Coh}_{\text{Nilp}}(\text{LocSys}_{\hat{G}}/\overline{\mathbb{Z}}_\ell) \xrightarrow{\sim} D_{lis}(\text{Bun}_G, \overline{\mathbb{Z}}_\ell)^\omega$$

# The real deal

Here  $Nilp = \text{Arinkin-Gaitsgory singular support condition}$ .  
Disappears over  $\overline{\mathbb{Q}}_\ell$  (automatic).

Thus, have to think of local Langlands as a "non-abelian Fourier transform" with "kernel given by the Whittaker representation" !!

Many other things in the article. For example :

- ▶ Notion of ULA sheaves in non-archimedean geometry (typically for spatial diamonds)
- ▶ Jacobian criterion of smoothness that allows to construct some coho smooth locally spatial diamonds using the curve (very deep and difficult theorem)