

# VOLUME ESTIMATES IN ANALYTIC AND ADELIC GEOMETRY

Antoine CHAMBERT-LOIR

Institut de recherche mathématique de Rennes, Université de Rennes 1

*Joint work with*      Yuri TSCHINKEL

Courant Institute, New-York University

*Counting points on varieties*  
Leiden, April 20–24, 2009



# CONTENTS

- ① TWO VOLUME ESTIMATES FOR SEMI-SIMPLE GROUPS
- ② CONJECTURES OF BATYREV, MANIN AND PEYRE
- ③ VOLUMES AND DISTRIBUTION OF REAL OR  $p$ -ADIC HEIGHT BALLS
- ④ TAMAGAWA MEASURES, ADELIC HEIGHT BALLS, AND THEIR VOLUMES



# TWO VOLUME ESTIMATES FOR SEMI-SIMPLE GROUPS

## 1 TWO VOLUME ESTIMATES FOR SEMI-SIMPLE GROUPS

- Real semi-simple Lie groups
- Adelic semi-simple algebraic groups

## 2 CONJECTURES OF BATYREV, MANIN AND PEYRE

## 3 VOLUMES AND DISTRIBUTION OF REAL OR $p$ -ADIC HEIGHT BALLS

## 4 TAMAGAWA MEASURES, ADELIC HEIGHT BALLS, AND THEIR VOLUMES



# REAL SEMI-SIMPLE LIE GROUPS

Let  $G$  be a semi-simple Lie group with trivial center,  $\mu$  a Haar measure on  $G$ .

Let  $\rho: G \rightarrow \mathrm{GL}(V)$  be a finite dimensional faithful representation of  $G$  in a real vector space  $V$ . Let  $\|\cdot\|$  be a norm on  $\mathrm{End}(V)$ .

For any  $T > 0$ , let  $B_T = \{g \in G; \|\rho(g)\| \leq T\}$  — compact in  $G$ .

## THEOREM (MAUCOURANT, 2004)

When  $T \rightarrow \infty$ :

- ① **volume estimate:**  $\mu(B_T) \sim cT^d \log(T)^e$  for some real number  $c$ , some rational number  $d$  and some integer  $e$ ;
- ② **convergence of measures:** there exists a measure  $\mu_\infty$  on  $\mathrm{End}(V)$  such that for any function  $f \in \mathcal{C}(\mathbb{P} \mathrm{End}(V))$ ,

$$\frac{1}{\mu(B_T)} \int_{B_T} f(\rho(g)) d\mu(g) \rightarrow \int_{\mathbb{P} \mathrm{End}(V)} f(g) d\mu_\infty(g).$$

# REAL SEMI-SIMPLE LIE GROUPS

## THEOREM (MAUCOURANT)

$$\mu(B_T) \sim cT^d \log(T)^e$$
$$\mu(B_T)^{-1} \int_{B_T} f(\rho(g)) d\mu(g) \rightarrow \int_{\mathbb{P}\mathrm{End}(V)} f(g) d\mu_\infty(g).$$

The numbers  $d$  and  $e$  are explicitly defined in terms of the relative root system of  $G$  and the weights of  $\rho$ .

One has  $0 \leq e \leq \mathrm{rank}_{\mathbb{R}}(G)$ .

The measure  $\mu_\infty$  is supported by a submanifold of  $\mathbb{P}\mathrm{End}(V)$  which is bi-invariant under  $G$ .

**Principle of proof:**  $K\alpha^+K$ -decomposition and integration formula.



# ADELIC SEMI-SIMPLE ALGEBRAIC GROUPS

Let  $G$  be a semi-simple algebraic group over  $\mathbb{Q}$ .

Let  $\rho: G \rightarrow \mathrm{GL}(V)$  be a faithful representation of  $G$  in a finite dimensional  $\mathbb{Q}$ -vector space  $V$  (with a unique highest weight).

For any  $p \in \{\text{prime numbers}\} \cup \{\infty\}$ , let  $\|\cdot\|_p$  be a  $p$ -adic norm on  $\mathrm{End}(V) \otimes \mathbb{Q}_p$ .

**Compatibility assumption:** there exists a basis  $(e_i)$  of  $V$  such that for almost all  $p$ : for any  $u \in \mathrm{End}(V) \otimes \mathbb{Q}_p$  with matrix representation  $U = (u_{i,j})$ , one has  $\|u\|_p = \max(|u_{i,j}|_p)$ .

Adeles:  $\mathbb{A} = \text{restricted product } \prod'_{\mathfrak{p}} \mathbb{Q}_p$ .

Then,  $G(\mathbb{A})$  is a locally compact group — restricted product  $\prod'_{\mathfrak{p}} G(\mathbb{Q}_p)$ .

For any  $T > 0$ , let  $B_T = \{g = (g_p) \in G(\mathbb{A}) ; \prod_p \|\rho(g_p)\|_p \leq T\}$  — this is a **compact set in  $G(\mathbb{A})$** .



# ADELIC SEMI-SIMPLE GROUPS

$$B_T = \{g = (g_p) \in G(\mathbb{A}) ; \prod_p \|\rho(g_p)\|_p \leq T\}$$

Fix a Haar measure  $\mu$  on  $\mathbf{G}(\mathbb{A})$ .

THEOREM (GORODNIK, MAUCOURANT, OH, 2007)

When  $T \rightarrow \infty$ :

- ① **volume estimate:**  $\mu(B_T) \sim cT^a \log(T)^b$  for some positive real number  $c$ , some rational number  $a$  and some non-negative integer  $b$ ;
- ② **convergence of measures:** for any function  $f \in \mathcal{C}(\mathbb{P}(\mathrm{End}(V) \otimes \mathbb{A}))$ ,

$$\frac{1}{\mu(B_T)} \int_{B_T} f(\rho(g)) d\mu(g) \rightarrow \int_{\mathbb{P}(\mathrm{End}(V) \otimes \mathbb{A})} f(g) d\mu_\infty(g).$$

# ADELIC SEMI-SIMPLE ALGEBRAIC GROUPS

THEOREM (GORODNIK, MAUCOURANT, OH)

$$\mu(B_T) \sim cT^a \log(T)^b$$

$$\frac{1}{\mu(B_T)} \int_{B_T} f(\rho(g)) d\mu(g) \rightarrow \int_{\mathbb{P}(\text{End}(V) \otimes \mathbb{A})} f(g) d\mu_\infty(g)$$

Again,  $a$  and  $b$  can be computed explicitly in terms of the weights of  $\rho$ , the root system of  $G$  and the action of  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  they possess.

The measure  $\mu_\infty$  is supported by  $X_\rho(\mathbb{A})$ , where  $X_\rho$  is the Zariski closure of  $\rho(G)$  in  $\mathbb{P}\text{End}(V)$  — **DE CONCINI–PROCESI’s wonderful compactification** of  $G$ .



# MOTIVATION/CONSEQUENCE OF THESE ESTIMATES

These **volume estimates**, resp. **convergence of measures** are one step in understanding the number, resp. the distribution, of

- points in  $\Gamma \cap B_T$ , where  $\Gamma$  is a lattice of the Lie group  $G$  — **lattice points in balls**;
- points in  $G(\mathbb{Q}) \cap B_T$  — **rational points of “bounded height”**.

When  $T \rightarrow \infty$ , and for adequate representations  $\rho$ , the obtained estimates are

- ①  $\#(\Gamma \cap B_T) \sim V(T)/\mu(G/\Gamma)$ ;
- ②  $\#(G(\mathbb{Q}) \cap B_T) \sim V(T)/\mu(G(\mathbb{A})/G(\mathbb{Q}))$  — with a deliberately ignored twist caused by automorphic characters.

Other actors of the play:

- ① Duke, Rudnick, Sarnak; Eskin, McMullen;
- ② Shalika, Tschinkel, Takloo-Bigash.



# GOAL AND MOTIVATION FOR THIS TALK

Generalize these results when the algebraic group is replaced by any affine variety:

- define limit measures; in the adelic setting, this will require to introduce convergence factors;
- define a notion of height so that  $p$ -adic, resp. adelic balls have finite volume;
- understand the volume of these height balls, or the measure-theoretical behaviour of these height balls, when  $T \rightarrow \infty$ .

## Motivation:

- conjectures by Batyrev, Manin and Peyre on the **distribution of rational points of bounded height**;
- investigation of analogous conjectures in the context of **integral points**



# CONJECTURES OF BATYREV, MANIN AND PEYRE

- 1 TWO VOLUME ESTIMATES FOR SEMI-SIMPLE GROUPS
- 2 CONJECTURES OF BATYREV, MANIN AND PEYRE
  - Rapid preview
  - Peyre's Tamagawa measure
- 3 VOLUMES AND DISTRIBUTION OF REAL OR  $p$ -ADIC HEIGHT BALLS
- 4 TAMAGAWA MEASURES, ADELIC HEIGHT BALLS, AND THEIR VOLUMES



# RATIONAL POINTS OF ALGEBRAIC VARIETIES

A basic problem in **diophantine geometry** consists in deciding whether **diophantine equations** have solutions or not, more generally, to tell as much as possible about the set of solutions.

From a **geometrical point of view**, describe the set of rational points of algebraic varieties defined over  $\mathbb{Q}$ , or the set of integral points of algebraic varieties over  $\mathbb{Z}$ .

We are interested in varieties whose rational points are dense for the Zariski topology. We thus have to sort them according to their “arithmetic complexity”, that is, their **height**.



# HEIGHTS

**Essential example:** a point  $P \in \mathbb{P}^n(\mathbb{Q})$ , with homogeneous coordinates  $[x_0 : \dots : x_n]$  coprime integers, has height  $H(P) = \max(|x_0|, \dots, |x_n|)$ .

**Finiteness property (NORTHCOTT):** for any  $B > 0$ , there are only finitely many points  $P \in \mathbb{P}^n(\mathbb{Q})$  such that  $H(P) \leq B$ .

**Question:** How many, when  $B \rightarrow \infty$ ?

**Answer (SCHANUEL):**  $\sim \frac{2^n}{\zeta(n+1)} B^{n+1}$ .

◀ Back

**Analytical tool:** the “height zeta function”, *i.e.*, the generating series

$$Z_{\mathbb{P}^n}(s) = \sum_{P \in \mathbb{P}^n(\mathbb{Q})} H(P)^{-s}.$$

Understand abscissa of convergence, meromorphic continuation, location of poles,...



# THE CONJECTURE OF BATYREV, MANIN, PEYRE

**Question:** What happens if one restricts to points lying in a subvariety  $X$  of  $\mathbb{P}^n$ ?

**Conjectural answer (MANIN):** If  $X$  is smooth, anticanonically embedded, it should be  $\approx B(\log B)^{t-1}$ , where  $t = \text{rank Pic}(X)$ , provided:

- $X(\mathbb{Q})$  is Zariski dense in  $X$ ;
- you allow to enlarge the ground field;
- you exclude from  $X$  some strict algebraic subvarieties.

**Refinement (PEYRE):** it might even be  $\sim cB(\log B)^{t-1}$ , with an arithmetical description of the constant  $c$ .



# THE CONJECTURE OF BATYREV, MANIN, PEYRE: EXAMPLES

Many, often non-trivial, examples:

- complete intersection of very large dimension (circle method);
- flag varieties (**LANGLANDS**'s theory of Eisenstein series);
- toric varieties; equivariant compactifications of vector spaces;
- wonderful compactifications of adjoint semi-simple groups;
- some Del Pezzo surfaces (using universal torsors)...

... but a counter-example (total space of the family of diagonal cubic surfaces in  $\mathbb{P}^3$ ).



# PEYRE'S CONSTANT — VOLUMES

In all cases where the conjectures have been established for the variety  $X$ , the constant  $c$  in front of the asymptotic expansion features three kinds of invariants:

- the position of the anticanonical class in the effective cone;
- the cardinality of a Galois cohomology group;
- the volume of (an adequate part of) the adelic space  $X(\mathbb{A})$  with respect to a suitable measure.



# PEYRE'S TAMAGAWA MEASURE

Assume  $K_X = \mathcal{O}_X(-d)$  for some integer  $d$ .

- for each prime  $p \leq \infty$ ,  $p$ -adic measure defined by suitable local gauge forms: since  $K_X \sim \mathcal{O}_X(-d)$ , one can take a meromorphic differential form  $\omega$  with  $\text{div}(\omega) = -dH_0 \cap X$  and set

$$\tau_p = |\omega|_p \left( \frac{|x_0|_p}{\max(|x_0|_p, \dots, |x_n|_p)} \right)^d$$

- find suitable convergence factors  $\lambda_p$  — Peyre takes  $\tau_p = L_p(1, \text{Pic}(\bar{X}))^{-1}$  ;
- define  $\tau$  as the absolutely convergent product  $L^*(1, \text{Pic}(\bar{X})) \prod_p (\lambda_p \tau_p)$ .



## EXAMPLE: PEYRE'S MEASURE FOR $\mathbb{P}^n$

For  $X = \mathbb{P}^n$ ,  $d = n + 1$  — Homogeneous coordinates  $[x_0 : \dots : x_n]$

On the (compact) chart  $x_0 = 1$  and  $|x_j| \leq 1$  for all  $j$ , one has

$$\tau_p = \frac{|dx_1 \cdots dx_n|_p}{\max(|x_0|_p, \dots, |x_n|_p)^d} = |dx_1|_p \cdots |dx_n|_p.$$

One has  $\tau_\infty(\mathbb{P}^n(\mathbb{R})) = (n+1)2^n$ .

For  $p$  prime,  $\tau_p$  is nothing but the “canonical measure” on  $\mathbb{P}^n(\mathbb{Q}_p) = \mathbb{P}^n(\mathbb{Z}_p)$ , hence:

$$\tau_p(\mathbb{P}^n(\mathbb{Q}_p)) = p^{-n} \# \mathbb{P}^n(\mathbb{F}_p) = 1 + p^{-1} + \cdots + p^{-n} = \frac{1 - p^{-n-1}}{1 - p^{-1}}.$$

Convergence factors: Let  $L_p(s, \text{Pic}(\overline{\mathbb{P}^n})) = (1 - p^{-s})^{-1}$  and set  $\lambda_p = L_p(1, \cdot)$ , hence  $\lambda_p \tau_p(\mathbb{P}^n(\mathbb{Q}_p)) = 1 - p^{-n-1}$ , and finally:

$$\tau(X(\mathbb{A})) = L^*(1, \text{Pic}(\overline{X})) \prod (\lambda_p \tau_p(\mathbb{P}^n(\mathbb{Q}_p))) = (n+1)2^n / \zeta(n+1).$$

This is essentially **SCHANUEL's constant!**

► Want to check?



# VOLUMES AND DISTRIBUTION OF REAL OR $p$ -ADIC HEIGHT BALLS

1 TWO VOLUME ESTIMATES FOR SEMI-SIMPLE GROUPS

2 CONJECTURES OF BATYREV, MANIN AND PEYRE

3 VOLUMES AND DISTRIBUTION OF REAL OR  $p$ -ADIC HEIGHT BALLS

- Measures
- Height balls
- An Igusa zeta function

4 TAMAGAWA MEASURES, ADELIC HEIGHT BALLS, AND THEIR VOLUMES



# MEASURES: GAUGE FORMS VS. METRICS

$F$ , local field with a fixed Haar measure.

Let  $X$  be a smooth projective variety over a local field  $F$ , purely of dimension  $n$ ,

$D$  divisor on  $X$ ,  $U = X \setminus |D|$ .

How to define measures on  $U$ ?

- ① a **gauge form**  $\omega \in K_X(U) = \Omega_X^n(U)$  defines a measure  $|\omega|$ : in local coordinates, write  $\omega = f(x) dx_1 \wedge \cdots \wedge dx_n$  and consider the measure  $|f(x)| dx_1 \dots dx_n$ . This is well-defined because of the formula for change of variables in multiple integrals;
- ② given a **metric on the canonical line bundle**  $K_X$ , one may take local forms  $\omega$  and define  $\tau_X$  by glueing the measures  $|\omega| / \|\omega\|$ ;
- ③ **metric on  $K_X(D)$ :** take local forms  $\omega$  and glue the measures  $|\omega| / \|\omega f_D\|$  to define  $\tau_{(X,D)}$ .



# MEASURES: EXAMPLE OF COMPACTIFICATIONS OF ALGEBRAIC GROUPS

Assume that  $X$  is a smooth equivariant compactification of an algebraic group  $G$ .

**Gauge forms.** — Pick  $\omega \in K_X(G)$  a (left-)invariant differential form; this is a gauge form on  $G$  and  $|\omega|$  is a Haar measure on  $G(F)$ .

**Metric on  $K_X$ .** — One has  $K_X \simeq \mathcal{O}_X(-D)$  with  $|D| = X \setminus G$ ; if  $K_X$  is metrized, then  $|\omega| / \|\omega\|$  defines a measure for which  $G(F)$  has finite volume.

**Metric on  $K_X(D)$ .** — Observe that  $\text{div}(\omega) = -D$  and  $\omega f_D$  is a global section of the trivial line bundle  $K_X(D)$ ; we may assume that  $\|\omega f_D\| = \text{cst} = 1$ . Then,  $\tau_{(X,D)} = \frac{|\omega|}{\|\omega f_D\|} = |\omega|$  is a Haar measure on  $G$ .



# HEIGHT BALLS

$L$  effective divisor with support  $|D|$ , metric on  $\mathcal{O}_X(L)$ ,  $f_L$  canonical section of  $\mathcal{O}_X(L)$ ;

for  $T > 0$ , the inequality  $\|f_L\| \geq 1/T$  defines a compact subset  $B_T$  in  $U(F)$  — “**height ball**”;

fix a metric on  $K_X(D)$ , gets a measure  $\tau_{(X,D)}$  on  $U(F)$ ,

volume of  $B_T$ :  $V(T) = \tau_{(X,D)}(B_T)$ .

## DEFINITION

Mellin transform:

$$Z(s) = \int_0^\infty t^{-s} dV(t) = \int_{U(F)} \|f_L\|^s \tau_{(X,D)}.$$



# IGUSA ZETA FUNCTION

## DEFINITION (MELLIN TRANSFORM)

$$Z(s) = \int_0^\infty t^{-s} dV(t) = \int_{U(F)} \|f_L\|^s \tau_{(X,D)}.$$

**Tauberian theory** relates analytic properties of  $Z(s)$  to the asymptotic behaviour of  $V(T)$ .

Then, the detailed asymptotic behaviour of  $V(T)$  can be used to study the convergence of the probability measures  $V(T)^{-1} \tau_{(X,D)}|_{B_T}$ .

**Simple remark:**  $Z(s)$  is a kind of Igusa zeta function; it should not be looked as in integral on  $U(F)$  but computed using the projective compactification  $X(F)$ .



# ANALYSIS OF THE IGUSA ZETA FUNCTION

**Geometric assumption:** over  $\overline{F}$ , the irreducible components  $D_\alpha$  of  $D$  are smooth, and intersect transversally.

**To simplify the exposition, I pretend here that the irreducible components of  $D$  over  $F$  are geometrically irreducible.**

Consider the corresponding stratification  $(D_A)$  of  $X$ , such that  $D_A^\circ$  is the open stratum consisting of points of  $X$  which belong exactly to the divisors  $D_\alpha$  for  $\alpha \in A$ .

By the transversality assumption,  $D_A$  is a smooth subvariety of  $X$  of codimension  $\#A$  (or empty).

Decomposition of divisors:  $D = \sum \rho_\alpha D_\alpha$ ,  $L = \sum \lambda_\alpha D_\alpha$ .



# LOCAL COMPUTATION “AT INFINITY”

We may use finitely many local charts on  $X(F)$  to study the integral  $Z(s)$ .  
The part “around” a point  $x \in D_A^\circ(F)$  can be computed as

$$\int \prod_{\alpha} \|f_{D_\alpha}\| (x)^{\lambda_\alpha s - \rho_\alpha} d\tau_X(x) = \int \prod_{\alpha \in A} |x_\alpha|^{\lambda_\alpha s - \rho_\alpha} \varphi(x; y; s) \prod_{\alpha} dx_\alpha dy.$$

Then, the analytic properties of  $Z(s)$  are completely clear and can be expressed in terms of the combinatorics of the stratification  $(D_A)$ .

For example, **abscissa of convergence** =

$$\max_{\substack{D_\alpha(F) \neq \emptyset \\ \lambda_\alpha > 0}} \frac{\rho_\alpha - 1}{\lambda_\alpha} \quad ;$$

**order of pole** = numbers of  $\alpha$  that achieve equality;

**leading coefficient** = sum of integrals over all stratas  $D_A$  of minimal dimension where  $A$  consists only of such  $\alpha$ s.



# TAMAGAWA MEASURES, ADELIC HEIGHT BALLS, AND THEIR VOLUMES

- 1 TWO VOLUME ESTIMATES FOR SEMI-SIMPLE GROUPS
- 2 CONJECTURES OF BATYREV, MANIN AND PEYRE
- 3 VOLUMES AND DISTRIBUTION OF REAL OR  $p$ -ADIC HEIGHT BALLS
- 4 TAMAGAWA MEASURES, ADELIC HEIGHT BALLS, AND THEIR VOLUMES
  - Definition of a Tamagawa measure
  - The adelic Igusa zeta function



# LOCAL MEASURES AND CONVERGENCE FACTORS

Let  $X$  be a smooth projective variety over  $\mathbb{Q}$ ,  
 $D$  effective divisor on  $X$ ,  $U = X \setminus |D|$ .

Fix an adelic metric on  $K_X(D)$ ; this defines measures  $\tau_{(X,D),p}$  on  $U(\mathbb{Q}_p)$  for all  $p$ .

To define a measure on  $U(\mathbb{A})$  from these  $\tau_p$ , one needs convergence factors  $\lambda_p$  such that the infinite product

$$\prod_p \lambda_p \tau_p(U(\mathbb{Z}_p))$$

converges absolutely.

## Examples:

- $X$  equivariant compactification of a semi-simple algebraic group  $G$ ,  $\tau_p$ =Haar measure: one may take  $\lambda_p = 1$ ;
- same, but  $G$  unipotent:  $\lambda_p = 1$ ;
- same, but  $G$  is a torus,  $\lambda_p = L_p(1, X^*(G_{\overline{\mathbb{Q}}}))$ ;
- if  $D = \emptyset$ ,  $\lambda_p = L_p(1, \text{Pic}(X_{\overline{\mathbb{Q}}}))^{-1}$  (**PEYRE**).



# A CHOICE OF CONVERGENCE FACTORS

Same notation:  $X, D, U, \tau_p$ .

**Geometric assumption:**  $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$ .

Two free  $\mathbb{Z}$ -modules with a  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -action:

- $\Gamma(U_{\overline{\mathbb{Q}}}, \mathcal{O}_X^*)/\overline{\mathbb{Q}}^*$ ;
- $\text{Pic}(U_{\overline{\mathbb{Q}}})/\text{torsion}$ .

Virtual  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -module:  $\text{EP}(U) = \Gamma(U_{\overline{\mathbb{Q}}}, \mathcal{O}_X^*)/\overline{\mathbb{Q}}^* - \text{Pic}(U_{\overline{\mathbb{Q}}})/\text{torsion}$ .

## THEOREM

*One may take  $\lambda_p = L_p(1, \text{EP}(U))$  for all  $p < \infty$ .*

## DEFINITION

Global measure on  $U(\mathbb{A})$ :

$$\tau_{(X,D)} = L^*(1, \text{EP}(U))^{-1} \prod_{p < \infty} (L_p(1, \text{EP}(U)) \tau_{(X,D),p}) \tau_{(X,D),\infty}.$$

# HEIGHT ON THE ADELIC SPACE $U(\mathbb{A})$

Let  $L$  be an effective divisor supported on  $|D|$

$f_L$  the canonical section of  $\mathcal{O}_X(L)$

Choosing an adelic metric on  $\mathcal{O}_X(L)$ , one get a **height function on the adelic space**  $U(\mathbb{A})$  defined by

$$H_L((x_p)) = \prod_{p \leq \infty} \|f_L(x_p)\|_p^{-1}.$$

## PROPOSITION

*The function  $H_L$  defines a continuous exhaustion of  $U(\mathbb{A})$ .*

**Height ball:** compact subset  $B_T = \{x \in U(\mathbb{A}) ; H_L(x) \leq T\}$ .

Volume and zeta function:

$$V(T) = \tau_{(X,D)}(B_T), \quad Z(s) = \int_0^\infty t^{-s} dV(t) = \int_{U(\mathbb{A})} H_L(x)^{-s} d\tau_{(X,D)}(x).$$



# PRODUCT OF $p$ -ADIC ZETA FUNCTIONS

Modulo absolute convergence, one has

$$Z(s) = L^*(1, \text{EP}(U))^{-1} Z_\infty(s) \prod_{p < \infty} (L_p(1, \text{EP}(U)) Z_p(s)),$$

where for  $p \leq \infty$ ,

$$Z_p(s) = \int_{U(\mathbb{Q}_p)} \|f_L(x)\|^s d\tau_{(X,D),p}(x)$$

is the  $p$ -adic Igusa zeta function described previously.

Recall the decomposition  $D = \sum \rho_\alpha D_\alpha$ ,  $L = \sum \lambda_\alpha D_\alpha$ , as well as the transversality assumption on the  $D_\alpha$ .

Then, choosing compatibly adelic metrics on  $\mathcal{O}(D_\alpha)$ , one has:

$$Z_p(s) = \int_{X(\mathbb{Q}_p)} \prod_\alpha \|f_{D_\alpha}\|^{s\lambda_\alpha - \rho_\alpha} d\tau_{X,p}(x).$$

The previous computation in charts shows that it converges absolutely for  $\Re(s) > \max((\rho_\alpha - 1)/\lambda_\alpha)$ . 

# DENEF'S FORMULA

For almost all  $p$ , one can give a precise formula for  $Z_p(s)$  in terms of the reduction mod.  $p$  of the whole situation. This is done by adapting the method used by **J. DENEF** to prove that the degrees of the local zeta functions are bounded when one makes the prime number  $p$  vary.

## PROPOSITION

*For  $p$  large enough, and for any complex number  $s$  such that  $\Re(s) > (\rho_\alpha - 1)/\lambda_\alpha$ , one has*

$$Z_p(s) = \sum_A p^{-\dim X} \#D_A^\circ(\mathbb{F}_p) \prod_{\alpha \in A} \frac{p-1}{p^{s\lambda_\alpha - \rho_\alpha + 1} - 1}.$$

This follows from the fact that the local computation in charts around  $x \in D_A^\circ$  can be done using étale coordinates  $((x_\alpha)_{\alpha \in A}, y)$  such that  $\|f_{D_\alpha}\| = |x_\alpha|$ , etc., and from the explicit computation:

$$\int_{p\mathbb{Z}_p} |x|^s dx = \sum_{n=1}^{\infty} \int_{p^n \mathbb{Z}_p \setminus p^{n+1} \mathbb{Z}_p} p^{-ns} dx = \sum_{n=1}^{\infty} p^{-ns-n} \left(1 - \frac{1}{p}\right) = p^{-1} \frac{p-1}{p^{s+1}-1}$$

# MEROMORPHIC CONTINUATION OF AN EULER PRODUCT

Let  $\sigma = \max(\rho_\alpha / \lambda_\alpha)$ , let  $A(L, D)$  be the set of  $\alpha$  where equality is achieved

One can deduce from Denef's formula that for  $\Re(s) > \sigma - \varepsilon$ ,

$$Z_p(s) = p^{-\dim X} \# U(\mathbb{F}_p) \prod_{\alpha \in A(L, D)} (1 + p^{-s\lambda_\alpha + \rho_\alpha - 1}) (1 + O(p^{-1-\varepsilon}))$$

hence

- ①  $\prod L_p(1, \text{EP}(U)) Z_p(s)$  converges absolutely for  $\Re(s) > \sigma$ ;
- ②  $\prod L_p(1, \text{EP}(U)) Z_p(s) \prod_{\alpha \in A(L, D)} (1 - p^{-s\lambda_\alpha + \rho_\alpha - 1})$  converges absolutely for  $\Re(s) > \sigma - \varepsilon$ .

Consequently, one obtains a meromorphic continuation of the form

$$Z(s) = L^*(1, \text{EP}(U))^{-1} \prod_p (L_p(1, \text{EP}(U)) Z_p(s)) = \varphi(s) \prod_{\alpha \in A(L, D)} \zeta(\lambda_\alpha(s - \sigma) + 1),$$

with

$$\varphi(1) = \prod_{\alpha \in A(L, D)} \zeta^*(1) \int_{X(\mathbb{A}_F)} \prod_{\alpha \notin A(L, D)} H_{D_\alpha}(x)^{\rho_\alpha - \sigma \lambda_\alpha} d\tau_X(x).$$



# CONCLUSION

Let  $E$  be the divisor  $\sigma L - D$ ; it is effective and its support is contained in  $|D|$ .

Let  $t = \#A(L, D)$ .

Some more calculation implies:

$$\lim_{s \rightarrow \sigma} Z(s)(s - \sigma)^t \prod_{\alpha \in A(L, D)} \lambda_\alpha = \int_{X(\mathbb{A})} H_E(x)^{-1} d\tau_X(x).$$

Using tauberian theorems, we deduce:

## THEOREM

When  $T \rightarrow \infty$ ,

- ① one has the asymptotic expansion  $V(T) = \tau_{(X, D)}(B_T) \sim B^\sigma (\log B)^{t-1} (\sigma(t-1)! \prod_{\alpha \in A(L, D)} \lambda_\alpha)^{-1} \int_{X(\mathbb{A})} H_E(x)^{-1} d\tau_X(x);$
- ② the probability measures  $V(T)^{-1} \tau_{(X, D)}|_{B_T}$  equidistribute to the unique probability measure proportional to  $H_E(x)^{-1} d\tau_X(x)$ .

The comparison of the geometric estimates we obtained with those of **MAUCOURANT *et al.*** is an exercice for specialists of wonderful compactifications of algebraic groups.

