

LINEAR INDEPENDENCE OF CLUSTER MONOMIALS FOR SKEW-SYMMETRIC CLUSTER ALGEBRAS

GIOVANNI CERULLI IRELLI, BERNHARD KELLER,
DANIEL LABARDINI-FRAGOSO, AND PIERRE-GUY PLAMONDON

Dedicated to Idun Reiten on the occasion of her seventieth birthday.

ABSTRACT. Fomin-Zelevinsky conjectured that in any cluster algebra, the cluster monomials are linearly independent and that the exchange graph is independent of the choice of coefficients. We confirm these conjectures for all skew-symmetric cluster algebras.

CONTENTS

1. Introduction	1
2. Recollections	2
2.1. Skew-symmetric cluster algebras of geometric type	2
2.2. Cluster categories	3
3. Linear independence of cluster monomials	4
4. Exchange graphs	7
5. Cluster algebras with arbitrary coefficients	10
References	11

1. INTRODUCTION

When S. Fomin and A. Zelevinsky invented cluster algebras more than ten years ago [16], one of their hopes was that the cluster monomials would be part of a ‘canonical’ basis. This naturally led them to the conjecture that cluster monomials should be linearly independent (Conjecture 4.16 of [17]). The conjecture has been checked for several classes, notably for finite type cluster algebras [7] [12], for skew-symmetric acyclic cluster algebras [19] and for cluster algebras associated with surfaces [15]. It was proved for all skew-symmetric cluster algebras of geometric type whose exchange matrix is of full rank in [14] and [26]. In this note, we show that the condition on the rank may be dropped, and that we need not restrict ourselves to the geometric type. In other words, we show that cluster monomials are linearly independent in any cluster algebra associated with a skew-symmetric matrix.

Date: March 7, 2012.

The fourth author was financially supported by the FQRNT.

Our proof is based on the additive categorification of cluster algebras using the cluster category [26] (closely related to that using decorated representations [14]) and on an idea from [8]. We show that the canonical cluster character yields a bijection from the set of isomorphism classes of reachable rigid objects onto the set of cluster monomials (Corollary 3.5). In this way, we treat the cluster algebras of geometric type, *i.e.* those associated with ice quivers. As a corollary of our proof, we show that for cluster algebras associated with ice quivers, the exchange graph does not depend on the frozen part of the quiver, which confirms Conjecture 4.3 of [18] for this class. This has been shown previously in [20] (see also the book [21]) for skew-symmetrizable cluster algebras whose defining matrix is of full rank. Once these results have been obtained for cluster algebras of geometric type, we show how we can extend them to arbitrary coefficients thanks to Fomin-Zelevinsky's separation formula.

We expect that our methods and results can be extended to the skew-symmetrizable cluster algebras categorified by L. Demonet in [11].

2. RECOLLECTIONS

2.1. Skew-symmetric cluster algebras of geometric type. We recall notions from [18]. Let \tilde{B} be an $n \times r$ matrix whose top $r \times r$ matrix B is skew-symmetric. We will work over the field $\mathbb{Q}(x_1, \dots, x_n)$; let \mathbb{P} be the tropical semifield generated by x_{r+1}, \dots, x_n .

Definition 2.1. A *seed* is a pair (\tilde{B}, \mathbf{u}) , where

- \tilde{B} is a matrix as above, and
- $\mathbf{u} = (u_1, \dots, u_n)$ is an ordered set whose elements form a free generating subset of the field $\mathbb{Q}(x_1, \dots, x_n)$.

Definition 2.2. Let $i \in \{1, \dots, r\}$. The *mutation* of the seed (\tilde{B}, \mathbf{u}) at vertex i is the seed $(\tilde{B}', \mathbf{u}')$, where

- $\tilde{B}' = (b'_{kj})$ is the $n \times r$ matrix defined by

$$b'_{kj} = \begin{cases} -b_{kj} & \text{if } k = i \text{ or } j = i, \\ b_{kj} + \frac{b_{ki}|b_{ij}| + |b_{ki}|b_{ij}}{2} & \text{otherwise,} \end{cases}$$

where $[z]_+ = \max(0, z)$ for any real number z .

- $\mathbf{u}' = (u'_1, \dots, u'_n)$, with $u'_j = u_j$ if $i \neq j$ and

$$u_i u'_i = \prod_{j=1}^n u_j^{[b_{ji}]_+} + \prod_{j=1}^n u_j^{[-b_{ji}]_+}.$$

Definition 2.3. Let \tilde{B} be a matrix as above.

- (1) The *clusters* are the sets \mathbf{u} contained in seeds (\tilde{B}', \mathbf{u}) obtained from the *initial seed* $(\tilde{B}, \mathbf{x} = (x_1, \dots, x_n))$ by iterated mutations.

- (2) The *cluster variables* are elements u_i of any cluster \mathbf{u} , with $i \in \{1, \dots, r\}$.
- (3) The *coefficients* are the elements x_{r+1}, \dots, x_n .
- (4) A *cluster monomial* is a product of cluster variables all belonging to the same cluster.
- (5) The (*skew-symmetric*) *cluster algebra (of geometric type)* $\mathcal{A}(\tilde{B})$ is the $\mathbb{Z}\mathbb{P}$ -subalgebra of $\mathbb{Q}(x_1, \dots, x_n)$ generated by all cluster variables.

Notice that one can associate an ice quiver (Q, F) to the matrix \tilde{B} in the following way:

- Q has n vertices, labelled by the rows of \tilde{B} ;
- the set F of frozen vertices is the set $\{r+1, \dots, n\}$;
- for any vertices i and j of Q , there are $[b_{ij}]_+$ arrows from i to j .

Conjecture 2.4 (Conjecture 7.2 of [18]). *The cluster monomials of any cluster algebra are linearly independent over $\mathbb{Z}\mathbb{P}$.*

2.2. Cluster categories. We now briefly recall results on the categorification of cluster algebras via triangulated categories. Recent surveys on the subject include [28], [29], [2] and [22].

Let (Q, W) be a quiver with a non-degenerate potential [13]. Generalizing the cluster category introduced in [3], C. Amiot [1] has defined the cluster category $\mathcal{C}_{Q,W}$ of (Q, W) ; it is a triangulated category with a rigid object $\Gamma = \Gamma_1 \oplus \dots \oplus \Gamma_n$ whose endomorphism algebra is isomorphic to the Jacobian algebra $J(Q, W)$.

As in [27, Definition 3.9], define \mathcal{D} to be the full subcategory of all objects X of $\mathcal{C}_{Q,W}$ such that

- there exists a triangle $T_1^X \rightarrow T_0^X \rightarrow X \rightarrow \Sigma T_1^X$, with T_0^X and T_1^X in $\text{add } \Gamma$;
- there exists a triangle $T_X^0 \rightarrow T_X^1 \rightarrow \Sigma X \rightarrow \Sigma T_X^0$, with T_X^0 and T_X^1 in $\text{add } \Gamma$; and
- $\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma X)$ is finite-dimensional.

Define the *index* [10] of an object X of \mathcal{D} as

$$\text{ind}_{\Gamma} X = [T_0^X] - [T_1^X] \in K_0(\text{add } \Gamma) \cong \mathbb{Z}^n.$$

An object X *admits an index* if there exists a triangle $T_1^X \rightarrow T_0^X \rightarrow X \rightarrow \Sigma T_1^X$, with T_0^X and T_1^X in $\text{add } \Gamma$ (so that the index of X is defined).

The *cluster character* [5] [25] is the map $CC : \text{Obj}(\mathcal{D}) \rightarrow \mathbb{Z}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ defined by

$$CC(X) = \mathbf{x}^{\text{ind}_{\Gamma} X} \sum_{\mathbf{e}} \chi\left(\text{Gr}_{\mathbf{e}}(\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma X))\right) \mathbf{x}^{B \cdot \mathbf{e}},$$

where

- B is the skew-symmetric matrix associated to Q ;

- $\text{Gr}_{\mathbf{e}}(\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma X))$ is the *quiver Grassmannian* (see Section 2.3 of [5]). It is a projective variety whose points correspond to submodules with dimension vector \mathbf{e} of the $\text{End}_{\mathcal{C}}(\Gamma)$ -module $\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma X)$;
- χ is the Euler–Poincaré characteristic.

A result of [23] implies the following proposition if $H^0\Gamma$ is finite-dimensional.

Proposition 2.5 ([27]). *The functor $F = \text{Hom}_{\mathcal{C}}(\Gamma, \Sigma(?))$ induces an equivalence of additive categories*

$$\mathcal{D}/(\Gamma) \xrightarrow{\sim} \text{mod } J(Q, W),$$

where $\text{mod } J(Q, W)$ is the category of finite-dimensional right modules over the Jacobian algebra $J(Q, W)$.

There is a notion of mutation of objects inside $\mathcal{C}_{Q,W}$, studied in [24]. A *reachable object* is an object obtained by iterated mutations from Γ . Any reachable object lies in the subcategory \mathcal{D} [27, Section 3.3].

The following theorem generalizes results obtained in [5] [7] [25].

Theorem 2.6 ([27]). *Let B be an $r \times r$ skew-symmetric matrix, and let Q be the associated quiver. Let W be a non-degenerate potential on Q . Then the cluster character CC induces a surjection*

$\{\text{isoclasses of indecomposable reachable objects of } \mathcal{D}\} \rightarrow \{\text{cluster variables of } \mathcal{A}(B)\}$,
which commutes with mutation on both sides.

In particular, if \tilde{B} is an $n \times r$ matrix whose upper $r \times r$ matrix is B , and if (\tilde{Q}, F) is the associated ice quiver, then the cluster variables of $\mathcal{A}(\tilde{B})$ are obtained from the cluster character CC by restricting mutation on the left hand side to non-frozen vertices.

3. LINEAR INDEPENDENCE OF CLUSTER MONOMIALS

Definition 3.1 (Definition 6.1 of [9]). Let B be a skew-symmetric matrix and \tilde{B} be an $n \times r$ integer matrix whose top $r \times r$ matrix is B .

- (1) Let $\mathbf{u} = (u_1, \dots, u_r)$ be a cluster of $\mathcal{A}(\tilde{B})$. A *proper Laurent monomial in \mathbf{u}* is a product of the form $u_1^{c_1} \cdots u_r^{c_r}$ where at least one of the c_i is negative.
- (2) The cluster algebra $\mathcal{A}(\tilde{B})$ has the *proper Laurent monomial property* if for any two clusters \mathbf{u} and \mathbf{u}' of $\mathcal{A}(\tilde{B})$, every monomial in \mathbf{u}' in which at least one element from $\mathbf{u}' \setminus \mathbf{u}$ appears with positive exponent is a $\mathbb{Z}\mathbb{P}$ -linear combination of proper Laurent monomials in \mathbf{u} .

Theorem 3.2 (Theorem 6.4 of [9]). *If $\mathcal{A}(\tilde{B})$ has the proper Laurent monomial property, then its cluster monomials are linearly independent over $\mathbb{Z}\mathbb{P}$.*

Using the key idea of Lemma 5.1 in [8] it was proved in [9, Theorem 6.2] that a cluster algebra $\mathcal{A}(\tilde{B})$ has the proper Laurent monomial property if there exists a potential on the quiver Q of \tilde{B} satisfying certain conditions.

These conditions allow the use of a homological interpretation of the E -invariant (see [14, Section 10]).

If (Q, W) is any quiver with a non-degenerate potential W , then the E -invariant has a similar interpretation in the cluster category $\mathcal{C}_{Q,W}$ (see [26, Proposition 4.15]). Using it, we prove the following result.

Let B be any skew-symmetric matrix and \tilde{B} be an $n \times r$ integer matrix whose top $r \times r$ matrix is B . Let (Q, F) be the ice quiver corresponding to \tilde{B} , and let W be a non-degenerate potential on Q .

Theorem 3.3. *If R is a rigid object of $\mathcal{D} \subset \mathcal{C}_{Q,W}$ which does not belong to $\text{add } \Gamma$ and such that $\text{Hom}_{\mathcal{C}}(\Gamma_i, \Sigma R) = 0$ for $i = r + 1, \dots, n$, then $CC(R)$ is a $\mathbb{Z}\mathbb{P}$ -linear combination of proper Laurent monomials in the initial cluster.*

PROOF. We follow the proof of [9, Theorem 6.2]. Letting $\text{ind}_{\Gamma} R = \sum_{i=1}^n g_i[\Gamma_i]$, we have that

$$\begin{aligned} CC(R) &= \mathbf{x}^{\text{ind}_{\Gamma} R} \sum_{\mathbf{e}} \chi\left(\text{Gr}_{\mathbf{e}}(\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma R))\right) \mathbf{x}^{\tilde{B} \cdot \mathbf{e}_0} \\ &= \sum_{\mathbf{e}} \chi\left(\text{Gr}_{\mathbf{e}}(\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma R))\right) \mathbf{x}_N^{\text{ind}_{\Gamma}^N R + B \cdot \mathbf{e}_0} \mathbf{x}_F^{\text{ind}_{\Gamma}^F R + B_c \cdot \mathbf{e}_0}, \end{aligned}$$

where B_c is the lower $(n-r) \times r$ submatrix of \tilde{B} , $\mathbf{e}_0 = (e_1, \dots, e_r)$, $\mathbf{x}_N = (x_1, \dots, x_r)$, $\mathbf{x}_F = (x_{r+1}, \dots, x_n)$, $\text{ind}_{\Gamma}^N R = \sum_{i=1}^r g_i[\Gamma_i]$ and $\text{ind}_{\Gamma}^F R = \sum_{i=r+1}^n g_i[\Gamma_i]$.

It suffices to show that, if \mathbf{e} is such that $\text{Gr}_{\mathbf{e}}(\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma R))$ is non-empty, then $\text{ind}_{\Gamma}^N R + B \cdot \mathbf{e}_0$ (the power of \mathbf{x}_N) has at least one negative entry.

Assume first that \mathbf{e} is non-zero. Since B is skew-symmetric, we have that $\mathbf{e}_0 \cdot B \cdot \mathbf{e}_0$ vanishes, and thus $\mathbf{e}_0 \cdot (\text{ind}_{\Gamma}^N R + B \cdot \mathbf{e}_0) = \mathbf{e}_0 \cdot \text{ind}_{\Gamma}^N R$. Thus it is sufficient to prove that $\mathbf{e}_0 \cdot \text{ind}_{\Gamma}^N R$ is negative. Since the last $(n-r)$ entries of \mathbf{e} are zero, we have that $\mathbf{e}_0 \cdot \text{ind}_{\Gamma}^N R = \mathbf{e} \cdot \text{ind}_{\Gamma} R$.

Let F be the functor $\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma(?))$. Let L be a submodule of FR with dimension vector \mathbf{e} , and let V be an object of \mathcal{D} such that $FV = L$. Then, by [26, Proposition 4.15] (or rather, by the second-to-last equality at the end of the proof)¹, we have that

$$\dim(\Sigma\Gamma)(V, \Sigma R) = \dim \text{Hom}_{J(Q,W)}(FV, FR) + \mathbf{e} \cdot \text{ind}_{\Gamma} R.$$

Since $FV = L$ is a submodule of FR and since the latter is non-zero, we have that $\dim \text{Hom}_{J(Q,W)}(FV, FR)$ is positive. Moreover, let $i : V \rightarrow R$ be a lift in $\mathcal{C}_{Q,W}$ of the inclusion of L in FR . Embed i in a triangle to get a diagram

$$\begin{array}{ccccc} U & \xrightarrow{u} & V & \xrightarrow{i} & R & \longrightarrow & \Sigma U \\ & & \downarrow f & \swarrow 0 & & & \\ & & \Sigma R & & & & \end{array}$$

¹There is a typo in this equality in [26]; the term $\text{ind}_{\Gamma} \Sigma Y$ should be read $\text{ind}_{\Gamma} Y$.

where the upper row is a triangle, and where u is in the ideal (Γ) , since $Fu = 0$ (Fi being injective). Now, if f is in $(\Sigma\Gamma)(V, \Sigma R)$, then the composition fu vanishes since Γ is rigid, and thus f factors through R . But R is also rigid; thus f is zero. Therefore $\dim(\Sigma\Gamma)(V, \Sigma R) = 0$, and $\mathbf{e} \cdot \text{ind}_\Gamma R$ is negative; this is what we wanted to prove.

Now, assume that \mathbf{e} is zero. Then we need to show that $\text{ind}_\Gamma^N R$ has a negative entry. Let $R = T_1^{a_1} \oplus \dots \oplus T_s^{a_s}$, with each T_i indecomposable. By hypothesis, one of the T_i , with $i \in \{1, \dots, r\}$, is not a direct factor of Γ ; thus $\text{ind}_\Gamma^N T_i$ has a negative entry (because rigid objects whose indices have only non-negative entries are in $\text{add } \Gamma$; this is a consequence of [26, Proposition 3.1]). Moreover, by [26, Theorem 3.7(1)], the indices of T_1, \dots, T_s are sign-coherent. Thus $\text{ind}_\Gamma^N R = \sum_{i=1}^s a_i \text{ind}_\Gamma^N T_i$ has a negative entry.

Another way of dealing with the case $\mathbf{e} = 0$ is by replacing V by R in the above equality, which becomes

$$\dim(\Sigma\Gamma)(R, \Sigma R) = \dim \text{Hom}_{J(Q,W)}(FR, FR) + \underline{\dim} FR \cdot \text{ind}_\Gamma R.$$

Since R is rigid, the left-hand side vanishes; since R is not in $\text{add } \Gamma$, FR is not zero. Thus we get

$$\underline{\dim} FR \cdot \text{ind}_\Gamma R = -\dim \text{Hom}_{J(Q,W)}(FR, FR) < 0,$$

and since the entries of $\underline{\dim} FR$ corresponding to frozen vertices vanish, this implies that $\text{ind}_\Gamma^N R$ has a negative entry. This finishes the proof. \square

Corollary 3.4. *If B is any skew-symmetric matrix and \tilde{B} is an $n \times r$ integer matrix whose top $r \times r$ matrix is B , then the cluster algebra $\mathcal{A}(\tilde{B})$ has the proper Laurent monomial property. In particular, the cluster monomials of any skew-symmetric cluster algebra of geometric type are linearly independent over $\mathbb{Z}\mathbb{P}$.*

PROOF. If the first statement of the theorem is true, then the independence of cluster monomials follows from [9, Theorem 6.4]. Let us prove that $\mathcal{A}(\tilde{B})$ has the proper Laurent monomial property.

Let Q be the quiver associated to \tilde{B} , and let W be a non-degenerate potential on Q . Consider the cluster category $\mathcal{C}_{Q,W}$. Let (u_1, \dots, u_r) be a cluster of $\mathcal{A}(\tilde{B})$, and let $z = u_1^{a_1} \dots u_r^{a_r}$ be a cluster monomial. By [27, Theorem 4.1], there exists a rigid object $T = T_1 \oplus \dots \oplus T_n$ of $\mathcal{C}_{Q,W}$ obtained from Γ by iterated mutations at vertices in $\{1, \dots, r\}$ and such that

$$z = CC(T_1^{a_1} \oplus \dots \oplus T_r^{a_r}).$$

Assume, moreover, that there is an index i such that u_i is not in the initial cluster and such that a_i is positive. Then $R = T_1^{a_1} \oplus \dots \oplus T_r^{a_r}$ satisfies the hypothesis of Theorem 3.3; thus z is a $\mathbb{Z}\mathbb{P}$ -linear combination of proper Laurent monomials in the initial cluster. This finishes the proof. \square

Corollary 3.5. *The surjection of Theorem 2.6 is a bijection.*

PROOF. Assume that R and R' are indecomposable reachable objects such that $CC(R) = CC(R')$. Let \mathbf{i} be a sequence of vertices of Q and j be a vertex of Q such that $\mu_{\mathbf{i}}(R)$ is isomorphic to Γ_j . Then we have that

$$x_j = CC(\Gamma_j) = CC(\mu_{\mathbf{i}}(R)) = \mu_{\mathbf{i}}(CC(R)) = \mu_{\mathbf{i}}(CC(R')) = CC(\mu_{\mathbf{i}}(R')).$$

Therefore, $CC(\mu_{\mathbf{i}}(R'))$ is not a linear combination of proper Laurent monomials in the initial variables. By Theorem 3.3, this implies that $\mu_{\mathbf{i}}(R')$ lies in $\text{add } \Gamma$. Since $CC(\mu_{\mathbf{i}}(R')) = x_j$, we must have that $\mu_{\mathbf{i}}(R')$ is isomorphic to Γ_j , and thus to $\mu_{\mathbf{i}}(R)$. Therefore, R and R' are isomorphic. \square

As noticed in [4, Theorem 4.1] for acyclic cluster algebras with trivial coefficients, the bijection of Corollary 3.5 allows us to prove that seeds are determined by their clusters, as conjectured in [17, Conjecture 4.14(2)]. This was also proved in [20] for cluster algebras of geometric type, and for cluster algebras whose matrix is of full rank (including the skew-symmetrizable case).

Corollary 3.6. *In any skew-symmetric cluster algebra, each seed is determined by its cluster.*

PROOF. We first deal with cluster algebras with trivial coefficients. Let (\mathbf{u}, Q) and (\mathbf{u}, Q') be two seeds with the same cluster. By Corollary 3.5, there is a unique reachable object T corresponding to \mathbf{u} in the cluster category; moreover, the ordinary quiver of $\text{End}_{\mathcal{C}}(T)$ has to be both Q and Q' . Thus the two seeds coincide. The result is proved for trivial coefficients.

If the coefficients are taken in an arbitrary semifield \mathbb{P} , then the morphism of semifields $\mathbb{P} \rightarrow \{1\}$ induces a morphism of cluster algebras $f : \mathcal{A} \rightarrow \mathcal{A}_{triv}$, where \mathcal{A}_{triv} is the cluster algebra defined from the same quiver as \mathcal{A} , but with trivial coefficients. Thus if two seeds (\mathbf{u}, Q) and (\mathbf{u}, Q') in \mathcal{A} have the same cluster, their images in \mathcal{A}_{triv} by f also have the same cluster, and thus Q and Q' are equal, since clusters determine their seeds when the coefficients are trivial. \square

4. EXCHANGE GRAPHS

Definition 4.1 (Definition 4.2 of [18]). The *exchange graph* of a cluster algebra $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$ is the n -regular connected graph whose vertices are the seeds of $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$ (up to simultaneous renumbering of rows, columns and variables) and whose edges connect the seeds related by a single mutation.

Conjecture 4.2 (Conjecture 4.3 of [18]). *The exchange graph of a cluster algebra $\mathcal{A}(\mathbf{x}, \mathbf{y}, B)$ only depends on the matrix B .*

Definition 4.3. Let (Q, W) be a non-degenerate quiver with potential (where the vertices $r + 1, \dots, n$ are frozen), and let $\mathcal{C}_{Q, W}$ be the associated cluster category.

- (1) The *exchange graph of Γ* is the r -ary graph whose vertices are iso-classes of objects reachable from Γ (by mutation at non-frozen vertices), and where two vertices are joined by an edge if the corresponding objects are related by a mutation.
- (2) The *exchange graph of indices* is the r -ary graph obtained from the exchange graph of Γ by replacing any vertex $T = T_1 \oplus \dots \oplus T_n$ by the n -tuple $(\text{ind}_\Gamma T_1, \dots, \text{ind}_\Gamma T_n)$ of elements of $K_0(\text{add } \Gamma) \cong \mathbb{Z}^n$.
- (3) The *exchange graph of reduced indices* is the graph obtained from the exchange graph of indices by replacing each n -tuple

$$(\text{ind}_\Gamma T_1, \dots, \text{ind}_\Gamma T_n)$$

by the r -tuple $(\text{ind}_\Gamma^N T_1, \dots, \text{ind}_\Gamma^N T_r)$, where $\text{ind}_\Gamma^N T_i$ is obtained from $\text{ind}_\Gamma T_i$ by forgetting the coefficients of $[\Gamma_{r+1}], \dots, [\Gamma_n]$.

The following corollary is a consequence of Corollary 3.5.

Corollary 4.4. *The exchange graphs of $\mathcal{A}(\tilde{B})$, of Γ , of indices and of reduced indices are isomorphic.*

PROOF. The fact that the exchange graphs of $\mathcal{A}(\tilde{B})$ and of Γ are isomorphic is a direct consequence of Corollary 3.5. The fact that the exchange graphs of Γ and of indices are isomorphic follows from the fact that rigid objects are determined by their index, by section 3.1 of [26], *cf.* also [10].

Let Γ_N be the direct sum of the Γ_j , $1 \leq j \leq r$, and Γ_F be the direct sum of the Γ_j , $r+1 \leq j \leq n$. To prove that the exchange graph of indices and of reduced indices are isomorphic, it is sufficient to show that if X and X' are indecomposable reachable objects (not in $\text{add } \Gamma_F$) such that $\text{ind}_\Gamma^N X = \text{ind}_\Gamma^N X'$, then they are isomorphic. Let

$$T_1 \rightarrow T_0 \oplus R \rightarrow X \rightarrow \Sigma T_1 \quad \text{and} \quad T_1 \rightarrow T_0 \oplus R' \rightarrow X' \rightarrow \Sigma T_1$$

be triangles, with $T_1, T_0 \in \text{add } \Gamma_N$ and $R, R' \in \text{add } \Gamma_F$. We can assume that T_1 has no direct factor in $\text{add } \Gamma_F$ since we have that $\text{Hom}_{\mathcal{C}}(X, \Sigma \Gamma_F)$ and $\text{Hom}_{\mathcal{C}}(X', \Sigma \Gamma_F)$ vanish (this is true because we only allow mutations at non-frozen vertices). But then $X \oplus R'$ and $X' \oplus R$ are rigid objects which have the same index; by [10], they are isomorphic. Since X and X' are indecomposable and not in $\text{add } \Gamma_F$, we get that they are isomorphic. \square

Lemma 4.5. *Let (Q, W) be a non-degenerate quiver with potential, and let Q^0 be the quiver obtained from Q by deleting the frozen vertices and arrows incident with them. Then the graph of reduced indices of (Q, W) only depends on Q^0 .*

PROOF. Note first that it follows from the definition of mutation of quivers that $\mu_i(Q)^0 = \mu_i(Q^0)$ for any non-frozen vertex i .

For any quiver R and any sequence $\mathbf{i} = (i_1, \dots, i_s)$ of non-frozen vertices, denote by $\mu_{\mathbf{i}}(R)$ the mutated quiver $\mu_{i_s} \cdots \mu_{i_2} \mu_{i_1}(R)$.

Let $\mathbf{i} = (i_1, \dots, i_s)$ be a sequence of non-frozen vertices of Q , and let $T = T_1 \oplus \dots \oplus T_n$ be the object obtained from Γ by iterated mutations at \mathbf{i} . The corresponding vertex in the exchange graph of reduced indices is $(\text{ind}_{\Gamma}^N T_1, \dots, \text{ind}_{\Gamma}^N T_r)$. We must prove that each entry only depends on Q^0 (and, implicitly, on the sequence \mathbf{i} of mutations).

Recall from [24] the functor $\Phi_i : \mathcal{C}_{\mu_i(Q,W)} \rightarrow \mathcal{C}_{Q,W}$ which sends the direct factor $(\Gamma_{\mu_i(Q,W)})_j$ of $\Gamma_{\mu_i(Q,W)}$ to Γ_j if $i \neq j$ and to a cone of the natural morphism

$$\Gamma_i \longrightarrow \bigoplus_{\alpha: i \rightarrow k} \Gamma_k$$

if $i = j$. Following [24], T is the image of $\Gamma_{\mu_i(Q,W)} \in \mathcal{C}_{\mu_i(Q,W)}$ by the composition of functors

$$\mathcal{C}_{\mu_i(Q,W)} \xrightarrow{\Phi_{i_s}} \dots \xrightarrow{\Phi_{i_2}} \mathcal{C}_{\mu_{i_1}(Q,W)} \xrightarrow{\Phi_{i_1}} \mathcal{C}_{Q,W}.$$

We will write $\Gamma' = \Gamma_{\mu_i(Q,W)}$ and $T = \Phi_{\mathbf{i}}(\Gamma')$.

The lemma will be a consequence of the following stronger statement: *If X is a rigid object of $\mathcal{C}_{\mu_i(Q,W)}$ which admits an index, then $\text{ind}_{\Gamma}^N \Phi_{\mathbf{i}}(X)$ only depends on Q^0 and on $\text{ind}_{\Gamma'}^N X$.*

We prove this statement by induction on s . For $s = 1$, we only perform one mutation at vertex $i = i_1$. Let $\text{ind}_{\Gamma} \Phi_i(X) = \sum_{j=1}^n g_j[\Gamma_j]$ and $\text{ind}_{\Gamma'} X = \sum_{j=1}^n g'_j[\Gamma'_j]$. Then the proof of [18, Conjecture 7.12] given in [26, Theorem 3.7 (iv)] gives us the formula

$$g_j = \begin{cases} -g'_i & \text{if } i = j, \\ g'_j + g'_i[b'_{ji}]_+ - b'_{ji} \min(g'_i, 0) & \text{if } i \neq j, \end{cases}$$

where $B' = (b'_{ij})$ is the matrix of $\mu_i(Q)$, and $[z]_+$ stands for $\max(z, 0)$ for any real number z . It follows from this formula that $\text{ind}_{\Gamma}^N \Phi_i(X) = \sum_{j=1}^r g_j[\Gamma_j]$ only depends on $\text{ind}_{\Gamma'}^N X$ and on $\mu_i(Q)^0$, and this last quiver only depends on Q^0 . The statement is shown for $s = 1$.

Assume it is proved for $s - 1$. Let $\mathbf{i} = (i_1, \dots, i_s)$. Then $\text{ind}_{\Gamma}^N \Phi_{\mathbf{i}}(X)$ only depends on Q^0 and on $\text{ind}_{\Gamma''}^N \Phi_{i_s}(X)$, where $\Gamma'' = \Gamma_{\mu_{i_{s-1}} \dots \mu_{i_1}(Q,W)}$. Moreover, $\text{ind}_{\Gamma''}^N \Phi_{i_s}(X)$ only depends on $\text{ind}_{\Gamma'}^N X$ and on $\mu_{i_{s-1}} \dots \mu_{i_1}(Q)^0$. This last quiver only depends on Q^0 . This proves the statement.

To finish the proof of the lemma, apply the statement to $X = \Gamma'_j$. Since $\Phi_{\mathbf{i}}(\Gamma'_j) = T_j$, and since Γ'_j is determined by its index, we get that $\text{ind}_{\Gamma}^N T_j$ only depends on Q_0 . This finishes the proof of the lemma. \square

Theorem 4.6. *Conjecture 4.2 is true for skew-symmetric cluster algebras of geometric type, that is, the exchange graph of $\mathcal{A}(\tilde{B})$ only depends on the top $r \times r$ matrix B .*

PROOF. By Corollary 4.4, the exchange graph of $\mathcal{A}(\tilde{B})$ is isomorphic to the exchange graph of reduced indices, which only depends on B by Lemma 4.5. \square

5. CLUSTER ALGEBRAS WITH ARBITRARY COEFFICIENTS

In this section, we extend Corollary 3.4 and Theorem 4.6 to the case of skew-symmetric cluster algebras with arbitrary coefficients. For the definition of cluster algebras with arbitrary coefficients, we refer the reader to [18, Section 2].

The linear independence of cluster monomials will follow from Corollary 3.4 and from the “separation of additions” of Fomin-Zelevinsky, which we recall.

Theorem 5.1 (Theorem 3.7 of [18]). *Let \mathcal{A} be a cluster algebra associated to the matrix B and with coefficients in an arbitrary semifield \mathbb{P} . Let t be a seed of \mathcal{A} , and let $x_{\ell,t}$ be a cluster variable of that seed. Then*

$$x_{\ell,t} = \frac{X_{\ell,t}|_{\mathcal{F}}(x_1, \dots, x_m; y_1, \dots, y_m)}{F_{\ell,t}|_{\mathbb{P}}(y_1, \dots, y_m)}.$$

In this theorem, $X_{\ell,t}$ and $F_{\ell,t}$ are rational functions which depend on the cluster algebra with principal coefficients (see [18, Definition 3.1]) and with defining matrix B . They are defined in [18, Definition 3.3].

Corollary 5.2. *If the cluster algebra with principal coefficients \mathcal{A}_0 associated to the matrix B has the proper Laurent monomial property, then any cluster algebra associated to B has the same property.*

PROOF. By definition, $X_{\ell,t}$ is an expression of a cluster variable with respect to the initial variables, inside the cluster algebra with principal coefficients. By assumption, this is an initial variable or a proper Laurent monomial in the initial variables. Specializing the coefficients to y_1, \dots, y_m and dividing by $F_{\ell,t}|_{\mathbb{P}}(y_1, \dots, y_m)$ does not change this property. Therefore any cluster variable $x_{\ell,t}$ is a proper Laurent monomial. \square

Corollary 5.3. *A skew-symmetric cluster algebra with arbitrary coefficients has the proper Laurent monomial property. In particular, its cluster monomials are linearly independent.*

PROOF. By Corollary 3.4, any skew-symmetric cluster algebra with principal coefficients has the proper Laurent monomial property (since cluster algebras with principal coefficients are of geometric type). Therefore, by Corollary 5.2, any skew-symmetric cluster algebra also has this property. Then Theorem 6.4 of [9] implies that cluster monomials in such a cluster algebra are linearly independent. Note that Theorem 6.4 of [9] is only stated for cluster algebras of geometric type; however, its proof adapts without change to arbitrary cluster algebras. \square

The proof that the exchange graph of a cluster algebra only depends on the defining matrix will follow from Theorem 4.6 and from the following theorem of Fomin-Zelevinsky.

Theorem 5.4 (Theorem 4.6 of [18]). *The exchange graph of an arbitrary cluster algebra is covered by the exchange graph of the cluster algebra with principal coefficients having the same defining matrix.*

Corollary 5.5. *The exchange graph of any skew-symmetric cluster algebra only depends on its defining matrix.*

PROOF. Let \mathcal{A} be any skew-symmetric cluster algebra, with defining matrix B . Let \mathcal{A}_0 and \mathcal{A}_{triv} be the cluster algebras with principal and trivial coefficients, respectively, associated to B . Let G , G_0 and G_{triv} be the exchange graphs of \mathcal{A} , \mathcal{A}_0 and \mathcal{A}_{triv} , respectively. By Theorem 5.4, G_0 is a covering of G . Moreover, G is a covering of G_{triv} . Since \mathcal{A}_0 and \mathcal{A}_{triv} are both of geometric type, Theorem 4.6 implies that the composed covering $G_0 \rightarrow G_{triv}$ is an isomorphism. Therefore, the coverings $G_0 \rightarrow G$ and $G \rightarrow G_{triv}$ are isomorphisms. This finishes the proof. \square

REFERENCES

1. Claire Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 6, 2525–2590.
2. Claire Amiot, *On generalized cluster categories*, ICRA XIV Proceedings, Eur. Math. Soc. (2011).
3. Aslak Bakke Buan, Robert Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), no. 2, 572–618.
4. Aslak Bakke Buan, Robert J. Marsh, Idun Reiten and Gordana Todorov, *Clusters and seeds in acyclic cluster algebras*, with an appendix coauthored in addition by P. Caldero and B. Keller, Proc. Amer. Math. Soc. **135** (2007), no. 10, 3049–3060 (electronic).
5. Philippe Caldero and Frédéric Chapoton, *Cluster algebras as Hall algebras of quiver representations*, Comment. Math. Helv. **81** (2006), no. 3, 595–616.
6. Philippe Caldero and Bernhard Keller, *From triangulated categories to cluster algebras. II*, Ann. Sci. École Norm. Sup. (4) **39** (2006), no. 6, 983–1009.
7. ———, *From triangulated categories to cluster algebras*, Invent. Math. **172** (2008), no. 1, 169–211.
8. Giovanni Cerulli Irelli, *Positivity in skew-symmetric cluster algebras of finite type*, arXiv:1102.3050 [math.RA], 2011.
9. Giovanni Cerulli Irelli and Daniel Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces, part III: Tagged triangulations and cluster monomials*, arXiv:1108.1774v3 [math.RT], 2011.
10. Raika Dehy and Bernhard Keller, *On the combinatorics of rigid objects in 2-Calabi-Yau categories*, Int. Math. Res. Not. IMRN (2008), no. 11, Art. ID rnn029, 17.
11. Laurent Demonet, *Mutations of group species with potentials and their representations. Applications to cluster algebras*, arXiv:1003.5078 [math.RT].
12. ———, *Categorification of skew-symmetrizable cluster algebras*, Algebras and Representation Theory **14** (2011), no. 6, 1087–1162.

13. Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky, *Quivers with potentials and their representations. I. Mutations*, *Selecta Math. (N.S.)* **14** (2008), no. 1, 59–119.
14. ———, *Quivers with potentials and their representations II: applications to cluster algebras*, *J. Amer. Math. Soc.* **23** (2010), no. 3, 749–790.
15. Sergey Fomin, Michael Shapiro, and Dylan Thurston, *Cluster algebras and triangulated surfaces. I. Cluster complexes*, *Acta Math.* **201** (2008), no. 1, 83–146.
16. Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. I. Foundations*, *J. Amer. Math. Soc.* **15** (2002), no. 2, 497–529 (electronic).
17. ———, *Cluster algebras: notes for the CDM-03 conference*, *Current developments in mathematics, 2003*, *Int. Press, Somerville, MA, 2003*, pp. 1–34.
18. ———, *Cluster algebras. IV. Coefficients*, *Compos. Math.* **143** (2007), no. 1, 112–164.
19. Christof Geiss, Bernard Leclerc and Jan Schröer, *Generic bases for cluster algebras and the Chamber Ansatz*, *J. of the Amer. Math. Soc.* **25** (2012), 21–76.
20. Michael Gekhtman, Michael Shapiro and Alek Vainshtein, *On the properties of the exchange graph of a cluster algebra*, *Math. Res. Lett.* **15** (2008), no. 2, 321–330.
21. ———, *Cluster Algebras and Poisson Geometry*, *Amer. Math. Soc., Math. Surveys and Monographs* **167**, 2010, 246 pages.
22. Bernhard Keller, *Cluster algebras and derived categories*, arXiv:1202.4161v3 [math.RT].
23. Bernhard Keller and Idun Reiten, *Cluster-tilted algebras are Gorenstein and stably Calabi-Yau*, *Adv. Math.* **211** (2007), no. 1, 123–151.
24. Bernhard Keller and Dong Yang, *Derived equivalences from mutations of quivers with potential*, *Advances in Mathematics* **226** (2011), 2118–2168.
25. Yann Palu, *Cluster characters for 2-Calabi-Yau triangulated categories*, *Ann. Inst. Fourier (Grenoble)* **58** (2008), no. 6, 2221–2248.
26. Pierre-Guy Plamondon, *Cluster algebras via cluster categories with infinite-dimensional morphism spaces*, *Compos. Math.* **147** (2011), 1921 – 1954.
27. ———, *Cluster characters for cluster categories with infinite-dimensional morphism spaces*, *Advances in Mathematics* **227** (2011), no. 1, 1 – 39.
28. Idun Reiten, *Tilting theory and cluster algebras*, arXiv: 1012.6014, preprint 2008, *Proc. Trieste Workshop*.
29. Idun Reiten, *Cluster categories*, *Proceedings of the ICM 2010, Hyderabad*.

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, 53115 BONN, GERMANY

UNIVERSITÉ PARIS DIDEROT - PARIS 7, UFR DE MATHÉMATIQUES, CASE 7012, BÂTIMENT CHEVALERET, 75205 PARIS CEDEX 13, FRANCE

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, 53115 BONN, GERMANY

LABORATOIRE LMNO, ÉQUIPE ALGÈBRE, GÉOMÉTRIE ET LOGIQUE, UNIVERSITÉ DE CAEN, F14032 CAEN CEDEX, FRANCE

E-mail address: cerulli@math.uni-bonn.de

E-mail address: keller@math.jussieu.fr

E-mail address: labardini@math.uni-bonn.de

E-mail address: pierre-guy.plamondon@unicaen.fr