

---



---

**Problem set 5: Hyperbolic surfaces**


---



---

**Exercise 1 (Unique geodesics).** The goal of this exercise is to prove that free homotopy classes of essential curves on hyperbolic surfaces contain unique geodesics.

- (a) Let us first state the Arzelà–Ascoli theorem. Recall that a map  $f : X \rightarrow Z$  between metric spaces  $X$  and  $Z$  is called  $L$ -Lipschitz, for some  $L > 0$  if

$$d_Z(f(x), f(y)) \leq L \cdot d_X(x, y)$$

for all  $x, y \in X$ . The Arzelà–Ascoli theorem now states:

**Theorem (Arzelà–Ascoli)** Let  $X$  be a metric space that has a countable dense subset and  $Z$  a compact metric space. Suppose  $\gamma_n : X \rightarrow Z$  is an  $L$ -Lipschitz map for all  $n \in \mathbb{N}$  and some fixed  $L > 0$ . Then there exists a subsequence  $(\gamma_{n_k})_{k \in \mathbb{N}}$  that converges uniformly on compact sets in  $X$  to an  $L$ -Lipschitz map  $\gamma : X \rightarrow Z$ .

Use this theorem to show that every non-trivial free homotopy class on a closed hyperbolic surface  $X$  contains at least one closed geodesic that minimizes the length in the homotopy class.

- (b) Show that this geodesic is unique. Hint: Suppose that there are two parallel geodesics and lift these to  $\mathbb{H}^2$ .

**Exercise 2 (The band model).**

- (a) Show that

$$\mathbb{B} = \left\{ z \in \mathbb{C}; |\operatorname{Im}(z)| < \frac{\pi}{2} \right\},$$

equipped with the metric

$$ds^2 = \frac{dx^2 + dy^2}{\cos^2(y)}$$

is isometric to  $\mathbb{H}^2$ . *Hint: this can be done without providing an isometry.*

- (b) Show that  $\mathbb{R} \subset \mathbb{B}$  is a geodesic.

**Exercise 3 (Twist curves).** Let  $\mathcal{P}$  be a pants decomposition of a surface  $S$ . Show that there exists collection of disjoint simple closed curves  $\Gamma$  so that for each pair of pants  $P$  in  $S \setminus \mathcal{P}$ ,  $\Gamma \cap \mathcal{P}$  consists of three arcs, each connecting a different pair of boundary components of  $P$ .

**Exercise 4 (Gauss–Bonnet for hyperbolic surfaces).**

(a) Suppose  $T$  is a hyperbolic triangle with angles  $\alpha, \beta, \gamma \geq 0$  at the vertices. Show that

$$\text{area}(T) = \pi - \alpha - \beta - \gamma$$

(b) Given a closed orientable hyperbolic surface  $X$  of genus  $g$ , equipped with a topological triangulation, we may straighten the edges to geodesic segments without moving the vertices. The result is still a triangulation (no intersections between edges will be created), this is a similar result to that of the first exercise that we will assume. Prove that

$$\text{area}(X) = 4\pi \cdot (g - 1),$$

without using the Gauss–Bonnet formula (of which this is a special case).