

THE FULL FAITHFULNESS CONJECTURES IN CHARACTERISTIC p

BRUNO KAHN

ABSTRACT. We present a triangulated version of the conjectures of Tate and Beilinson on algebraic cycles over a finite field. This sheds a new light on Lichtenbaum’s Weil-étale cohomology.

CONTENTS

Introduction	1
1. General overview	3
2. The Tate conjecture: a review	5
3. The Beilinson, Parshin and Friedlander conjectures	7
4. Motivic cohomology and l -adic cohomology	11
5. Categories of motives over a field	13
6. Categories of motives over a base	19
7. Effectivity and spectra	26
8. A proof of Theorem 3.2.1	30
9. Motivic reformulation of Friedlander’s conjecture	32
10. Duality and finite generation	40
11. Weil-étale reformulation	48
12. A positive case of the conjectures	52
Appendix A. A letter to T. Geisser	54
References	57

INTRODUCTION

It is generally understood that the “standard” conjectures on mixed motives predict that certain triangulated realisation functors should be conservative. The aim of this text is to explain that, at least in characteristic p , they predict much more: namely, that suitable triangulated realisation functors should be *fully faithful*.

The main result is the following. Let \mathbf{F} be a finite field, and let $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(\mathbf{F})$ be Voevodsky’s stable category of (unbounded, étale) motivic complexes.

Date: September 19, 2012.

It contains the category $\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(\mathbf{F})$ of effective motivic complexes as a full subcategory. Let l be a prime number different from 2 and $\mathrm{char} \mathbf{F}$. By work of Ayoub [4], there is a pair of adjoint functors:

$$\mathrm{DM}_{\acute{e}t}(\mathbf{F}) \underset{R_l}{\overset{\Omega_l}{\rightleftarrows}} \hat{D}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)$$

where the right hand side is Ekedahl's category of l -adic coefficients [22]. In particular, we have the object

$$\Gamma = \Omega_l(\mathbf{Z}_l) \in \mathrm{DM}_{\acute{e}t}(\mathbf{F}).$$

Theorem 1 (cf. Corollary 9.8.4). *The following conditions are equivalent:*

- (i) *The Tate conjecture (on the poles of the zeta function) and the Beilinson conjecture (on rational equivalence agreeing with numerical equivalence) hold for any smooth projective \mathbf{F} -variety.*
- (ii) $\Gamma \in \mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(\mathbf{F})$.

The full faithfulness statement announced above appears as another equivalent condition in Proposition 10.3.3 a); further equivalent conditions (finite generation of Hom groups) appear in Theorem 10.4.2. After the fact, see §11, these reformulations involve Weil étale cohomology. For a case when they hold in the triangulated context, see Theorem 12.2.1.

Curiously, the Beilinson conjecture and the Parshin conjecture (on vanishing of higher rational K -groups of smooth projective \mathbf{F} -varieties) are sufficient to imply the existence of a motivic t -structure on $\mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$, as well as semi-simplicity and independence of l for the \mathbf{Q}_l -adic realisations of objects of this category: see Proposition 10.5.1.

A problem is that there is no known analogue of this picture in characteristic 0 at the moment. While in characteristic p a single l -adic cohomology is sufficient to approach cycles modulo rational equivalence, it seems that in characteristic 0 one should consider the full array of realisation functors, plus their comparison isomorphisms. Even with this idea it does not seem obvious how to get a clean conjectural statement. In the light of §11, this might be of great interest to get the right definition of Weil-étale cohomology in characteristic 0.

This is a write-up of the talk I gave at the summer school on July 27, 2006. Much of the oral version was tentative because the suitable l -adic realisation functors were not constructed at the time. The final version is much more substantial than I had envisioned: this is both because of technical difficulties and because I tried to make the exposition as pedagogical as possible, in the spirit of the summer school. I hope the reader will bear with the first reason, and be satisfied with the second one.

I also hope that some readers will, like me, find the coherence and beauty of the picture below compelling reasons to believe in these conjectures.

I wish to thank Joseph Ayoub for a great number of exchanges while preparing this work, and the referee for a thorough reading which helped me improve the exposition.

Notation. k denotes a perfect field; we write $\mathrm{Sm}(k)$ for the category of smooth separated k -schemes of finite type. When k is finite we write \mathbf{F} instead of k and denote by $G \simeq \hat{Z}$ its absolute Galois group.

If \mathcal{C} is a category, we write $\mathcal{C}(X, Y)$, $\mathrm{Hom}_{\mathcal{C}}(X, Y)$ or $\mathrm{Hom}(X, Y)$ for the set of morphisms between two objects X, Y , according to notational convenience.

1. GENERAL OVERVIEW

This section gives a background to the sequel of the paper.

1.1. Triangulated categories of motives. As explained in André's book [1, Ch. 7], the classical conjectures of Hodge and Tate, and less classical ones of Grothendieck and Ogus, may be interpreted as requesting certain realisation functors on pure Grothendieck motives to be fully faithful. These conjectures concern algebraic cycles on smooth projective varieties modulo homological equivalence. On the other hand, both Bloch's answer to Mumford's nonrepresentability theorem for 0-cycles [10, Lect. 1] and Beilinson's approach to special values of L -functions [5, 7] led to conjectures on cycles modulo *rational* equivalence: the conjectures of Bloch-Beilinson and Murre (see Jannsen [45] for an exposition).

This development came parallel to another idea of Beilinson: in order to construct the (still conjectural) abelian category $\mathcal{M}(k)$ of mixed motives over a field k , one might start with the easier problem of constructing a triangulated category of motives, leaving for later the issue of finding a good t -structure on this category. Perhaps Beilinson had two main insights: first, the theory of perverse sheaves he had been developing with Bernstein, Deligne and Gabber [8] and second, his vanishing conjecture for Adams eigenspaces on algebraic K -groups (found independently by Soulé) which deals with an *a priori* obstruction to the existence of $\mathcal{M}(k)$.

The latter programme: constructing triangulated categories of motives, was successfully developed by Levine [53], Hanamura [31] and Voevodsky [78] independently. All three defined tensor triangulated categories of motives over k , by approaches similar in flavour but quite different in detail. It is now known that all these categories are equivalent, if $\mathrm{char} k = 0$ or if we take rational coefficients.¹ More precisely, the comparison between Levine's and Voevodsky's categories is due to Levine in characteristic 0 [53,

¹Gabber's recent refinement of de Jong's alteration theorem [36] now allows us to just invert the exponential characteristic for these theorems.

Part I, Ch. VI, 2.5.5] and to Ivorra in general [38], while the comparison between Hanamura’s and Voevodsky’s categories is due to Bondarko [12] and independently to Hanamura (unpublished).

These three constructions extend when replacing the field k by a rather general base S [53, 81, 32]². At this stage, the issue of Grothendieck’s six operations [7, 5.10 A] starts to make sense. In a talk at the ICTP in 2002, Voevodsky gave hints on how to carry this over in an abstract framework which would fit with his constructions, at least for the four functors f^* , f_* , $f_!$ and $f^!$. This programme was taken up by Ayoub in [2]; he added a great deal to Voevodsky’s outline, namely a study of the missing operations \otimes and $\underline{\text{Hom}}$ plus related issues like constructibility and Verdier duality, as well as an impressive theory of specialisation systems, a vast generalisation of the theory of nearby cycle functors.

It remained to see whether this abstract framework applied to categories of motives over a base, for example to the Voevodsky version $S \mapsto \text{DM}(S)$ constructed using relative cycles (“sheaves with transfers”). It did apply to a variant “without transfers” $S \mapsto \text{DA}(S)$ (as well as to Voevodsky’s motivating example: the Morel-Voevodsky stable \mathbf{A}^1 -homotopy categories $S \mapsto \text{SH}_{\mathbf{A}^1}(S)$): see [2, Ch. 4] for this. It did not apply directly to DM , however. This issue was solved to some extent by Cisinski and Déglise [15], who showed that the natural functor $\text{DA}_{\text{ét}}(S, \mathbf{Q}) \rightarrow \text{DM}(S, \mathbf{Q})$ is an equivalence of categories when S is a normal scheme, where $\text{DA}_{\text{ét}}$ is an étale variant of DA . All this will be explained in much more detail in §6.

1.2. Motivic conjectures and categories of motives. It is both a conceptual and a tactical issue to reformulate the conjectures alluded to at the beginning of §1.1 in this triangulated framework. The first necessary thing is to have triangulated realisation functors at hand. In Levine’s framework, many of them are constructed in his book [53, Part I, Ch. V]. In Voevodsky’s framework, with rational coefficients and over a field of characteristic 0, this was done by Huber using her triangulated category of mixed realisations as a target [33]. Over a separated Noetherian base, with integral coefficients and for l -adic cohomology, this was done by Ivorra [37].

Then came up the issue whether realisation functors commute with the six operations. The only context where the question made full sense was Ivorra’s. But there were three problems at the outset: Ivorra’s functors 1) are only defined on geometric motives, and 2) are contravariant. The third problem is that the formalism of six operations is not known to exist on $S \mapsto \text{DM}(S)$ in full generality, as explained above.

These issues were recently solved by Ayoub who constructed covariant l -adic realisation functors from $\text{DA}_{\text{ét}}(S)$ to Ekedahl’s l -adic categories $\hat{D}(S, \mathbf{Z}_l)$ [4]. He proved that they commute with the six operations and with the right choice of a specialisation system. More details are in §6.

²As far as I know, no comparison between these extensions has been attempted yet.

2. THE TATE CONJECTURE: A REVIEW

In this section, $\mathbf{F} = \mathbf{F}_q$ is a finite field with q elements. The main reference here is Tate's survey [76].

2.1. The zeta function and the Weil conjectures. Let X be a \mathbf{F} -scheme of finite type. It has a zeta function:

$$\zeta(X, s) = \exp \left(\sum_{n \geq 1} |X(\mathbf{F}_{q^n})| \frac{q^{-ns}}{n} \right) = \prod_{x \in X(0)} (1 - |\mathbf{F}(x)|^{-s})^{-1}$$

Weil conjectured that $\zeta(X, s) \in \mathbf{Q}(q^{-s})$ for any X : Dwork was first to prove it in [21]. A different proof, based on l -adic cohomology, was given by Grothendieck et al in [SGA5]. It provided the following extra property, also conjectured by Weil: if X is smooth projective, there is a functional equation of the form

$$\zeta(X, s) = AB^s \zeta(X, \dim X - s)$$

where A, B are constants.

Continue to assume X smooth projective and let $n \geq 0$. We write $\mathcal{Z}^n(X)$ for the group of cycles of codimension n on X and $A_{\text{num}}^n(X)$ for its quotient by numerical equivalence: $A_{\text{num}}^n(X)$ is finitely generated (see [1, 3.4.6] or [66, Th. 2.15]). The starting point of this work is the following

Conjecture 2.1.1 (Tate [74]). $\text{ord}_{s=n} \zeta(X, s) = -\text{rk } A_{\text{num}}^n(X)$.

2.2. Cohomological interpretation. The proofs of [SGA5] give the factorisation (another Weil conjecture)

$$\zeta(X, s) = \prod_{i=0}^{2d} P_i(q^{-s})^{(-1)^{i+1}}$$

with

$$P_i(t) = \det(1 - \varphi t \mid H_i^i(X))$$

$$\varphi = \text{Frobenius}, \quad H_l^i(X) := H_{\text{cont}}^i(\bar{X}, \mathbf{Q}_l) \quad (l \neq p).$$

Here $\bar{X} = X \otimes_{\mathbf{F}} \bar{\mathbf{F}}$, $H_{\text{cont}}^i(\bar{X}, \mathbf{Q}_l) := \varprojlim H_{\text{ét}}^i(\bar{X}, \mathbf{Z}/l^n) \otimes \mathbf{Q}_l$.

Finally the last Weil conjecture was proven by Deligne in [19]:

Theorem 2.2.1 (“Riemann Hypothesis over finite fields”). *For all i , $P_i(t) \in \mathbf{Z}[t]$ and its inverse roots have complex absolute values $= q^{i/2}$.*

This yields:

Theorem 2.2.2 (Milne [58, Prop. 8.2 and 8.4], Tate [76, Th. (2.9)]). *For any (X, n, l) , the following are equivalent:*

- (1) *Conjecture 2.1.1*;
- (2) $\text{rk } A_{\text{num}}^n(X) = \dim_{\mathbf{Q}_l} H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))^G$, $G = \text{Gal}(\bar{\mathbf{F}}/\mathbf{F})$;
- (3) (a) *the cycle map* $\mathcal{Z}^n(X) \otimes \mathbf{Q}_l \rightarrow H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))^G$ *is surjective*,
 (b) *homological and numerical equivalences agree on* $\mathcal{Z}^n(X) \otimes \mathbf{Q}$.

(3) (a) is called the *cohomological Tate conjecture*: it makes sense (and is conjectured to hold) over any finitely generated field.

Remark 2.2.3. Let us write S^n for the following condition: the composition

$$H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))^G \hookrightarrow H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n)) \twoheadrightarrow H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))_G$$

is bijective. Poincaré duality easily implies that $S^n \iff S^{d-n}$ if $d = \dim X$. Let us also write more precisely $(a)_n$ for (a) in Theorem 2.2.2 (3). Then, by [76, Th. (2.9)], the conditions of this theorem are also equivalent to

- (4) $(a)_n + (a)_{d-n} + S^n$.

Condition S^n is verified if X is an abelian variety, essentially because the Frobenius endomorphism of X is semi-simple as an element of the centre of $\text{End}^0(X)$ [48, lemme 1.9]. Thus, in this case, (1) $\iff (a)_n + (a)_{d-n}$ in Theorem 2.2.2.

Remark 2.2.4. Here are other known consequences of Conjecture 2.1.1 (see Tate [76] and Milne [59]):

- (i) Grothendieck’s standard conjecture B (by Theorem 2.2.2 (3) (b)).
- (ii) Semi-simplicity of Galois action on l -adic cohomology ([58, Rk. 8.6], [49]).
- (iii) Any (pure) homological motive M is a direct summand of $h(A_K)(n)$ for an abelian \mathbf{F} -variety A , K/\mathbf{F} a finite extension and $n \in \mathbf{Z}$ [59, Rk. 2.7]. We say that M is *of abelian type*.
- (iv) “Any mixed motive [over a finite field!] is pure” [59, Th. 2.49] (see also Proposition 10.5.1 below).

2.3. Known cases of the cohomological Tate conjecture. They are scarce: see [76, §5] for a 1994 state of the art.

The main case is $n = 1$, where the conjecture is known for abelian varieties over any finitely generated field k (Tate [75] if k is finite, Zarhin [83] if $\text{char } k > 0$, Faltings [23, 24] if $\text{char } k = 0$). This trivially extends to smooth projective varieties X whose homological motive is of abelian type in the sense of Remark 2.2.4 (iii): for simplicity, we shall then say that X is *homologically of abelian type*. For some other X ’s, see [76, Th. (5.6) and (5.8)].

One can then sometimes extend the cohomological Tate conjecture from codimension 1 to all codimensions. It was essentially observed by Soulé

that this is automatic if X is homologically of abelian type and of dimension ≤ 3 [69, Th. 3].³ In another direction, this will work if the algebra $H_{\text{cont}}^{2*}(X \otimes_K \bar{k}, \mathbf{Q}_l(*))^G$ is generated in cohomological degree 2: this happens when k is finite and X is a product of elliptic curves (Spiess [71]), and also for powers of certain simple abelian varieties (Lenstra, Zarhin). See [60], where Milne also proves the Tate conjecture for powers of certain abelian varieties for which the above generation condition is not satisfied.

3. THE BEILINSON, PARSHIN AND FRIEDLANDER CONJECTURES

3.1. The conjectures.

Conjecture 3.1.1 (Beilinson). *Let X/\mathbf{F} be smooth projective and let $n \geq 0$. Then rational and numerical equivalences coincide on $\mathcal{Z}^n(X) \otimes \mathbf{Q}$.*

This is the strongest possible conjecture: it implies that all adequate equivalence relations should coincide on $\mathcal{Z}^n(X) \otimes \mathbf{Q}$! (Compare [1, 3.2.2.1 and 3.2.7.2].)

Conjecture 3.1.2 (Parshin). *For any smooth projective \mathbf{F} -variety X , the algebraic K -group $K_i(X)$ is torsion for $i > 0$.*

This contains in particular the Beilinson-Soulé vanishing conjecture on eigenspaces for the Adams operations ([5], [70, p. 501, Conj.]).

Remark 3.1.3. Milne writes in [61, §1] that Tate already formulated conjecture 3.1.1 orally in the Woods Hole seminar of 1964. On the other hand, Conjecture 3.1.2 is attributed to Parshin by Jannsen in [43, Conj. 12.2], but is also formulated by Beilinson in [6, Conj. 8.3.3 b)]⁴. So the attributions of these conjectures are a bit blurred. For simplicity, I will keep their established names of “Tate conjecture”, “Beilinson conjecture” and “Parshin conjecture” here.

3.2. “Elementary” implications. We now start reviewing implications between these conjectures, in the same spirit as Theorem 2.2.2 and Remark 2.2.4.

Theorem 3.2.1 (Geisser [28]). *Conjecture 2.1.1 + Conjecture 3.1.1 \Rightarrow Conjecture 3.1.2.*

See Theorem 8.0.2 below for a more precise version.

Geisser’s proof is based on Jannsen’s semi-simplicity theorem [44] plus a Frobenius argument going back to Soulé [69]. The same is true for the

³Soulé’s proof works over a finite field, but the argument of [50, proof of Th. 82] extends this to any finitely generated field.

⁴Beilinson attributes it to Parshin in [5, Conj. 2.4.2.3].

proof of the following theorem, in which CH^d is the quotient of \mathcal{Z}^d by rational equivalence.

Theorem 3.2.2 (Kahn [48]). *Let X be a smooth projective \mathbf{F} -variety of dimension d . Assume that $\text{Ker}(CH^d(X \times X) \otimes \mathbf{Q} \rightarrow A_{\text{num}}^d(X \times X) \otimes \mathbf{Q})$ is a nilpotent ideal. Then Conjecture 2.1.1 for X and all $n \geq 0 \Rightarrow$ Conjecture 3.1.1 for X and all $n \geq 0$.*

The point is that the hypothesis of Theorem 3.2.2 is verified if the Chow motive of X is *finite dimensional* in the sense of Kimura and O’Sullivan (see [1, Ch. 12] or [39]). This will be the case if X is *of abelian type*, that is, if its motive verifies condition (iii) of Remark 2.2.4 for *rational equivalence* (*ibid.*). This gives many cases where one can apply Theorem 3.2.2, see §2.3.

3.3. From smooth projective to more general varieties. Putting together the cohomological Tate conjecture ((3) (a) in Theorem 2.2.2) and the Beilinson conjecture (Conjecture 3.1.1) gives the following

Conjecture 3.3.1. *For any smooth projective \mathbf{F} -variety X and any $n \geq 0$, the cycle class map*

$$CH^n(X) \otimes \mathbf{Q}_l \rightarrow H^{2n}(\bar{X}, \mathbf{Q}_l(n))^G$$

is bijective.

This statement continues to make sense when removing the word “projective”. It was conjectured to be true by Friedlander. Actually, Friedlander made a general conjecture involving the l -adic Chern character from algebraic K -theory, with targets of the form $H^i(\bar{X}, \mathbf{Q}_l(n))^G$ for all $i \in \mathbf{Z}$ (*ibid.*), namely:

Conjecture 3.3.2 (Friedlander, [6, 8.3.4 b])). *For any smooth \mathbf{F} -variety X , the Chern character*

$$ch_{n,i} : K_{2n-i}(X)_{\mathbf{Q}}^{(n)} \otimes \mathbf{Q}_l \rightarrow H^i(\bar{X}, \mathbf{Q}_l(n))^G$$

is an isomorphism, where the left hand side is the n -th Adams eigenspace on the algebraic K -theory of X .

For $i = 2n$ we get the Adams eigenspace of weight n on $K_0(X) \otimes \mathbf{Q}$ which may be identified with $CH^n(X) \otimes \mathbf{Q}$ thanks to the Grothendieck Riemann-Roch theorem, so Friedlander’s conjecture includes Conjecture 3.3.1 in the extended case of smooth varieties. For X smooth projective, it implies Parshin’s conjecture 3.1.2 in view of Theorem 2.2.1, as observed by Beilinson (*ibid.*). Geisser’s theorem 3.2.1 refines this observation.

Remarks 3.3.3.

- (1) I want to avoid entering the details of algebraic K -theory and the construction of l -adic Chern characters here, in order to concentrate on the already substantial case of motivic cohomology and l -adic cycle class maps.
- (2) Chern characters involve denominators, so that Friedlander's conjecture is intrinsically with rational coefficients. One could get some integrality by using higher Chern classes, but not completely avoid the denominators. On the other hand, as we shall see, a motivic version of this conjecture can be formulated integrally.

In his habilitation thesis [43, Th. 12.7 a)], Jannsen proved that Friedlander's conjecture follows from its restriction to smooth projective varieties, provided one adds a generalisation of Condition S^n from Remark 2.2.3 and assumes resolution of singularities. Jannsen's proof is a little awkward mainly because the formulation of Friedlander's conjecture involves fixed points under Galois action, which is not an exact functor: this is where the recourse to semi-simplicity (and the weight filtration) was essential.

Jannsen also gives an extension of these conjectures to singular varieties using K' -theory and (Borel-Moore) l -adic homology.

Let K be a finitely generated field of characteristic p , and let X be a smooth K -variety. Then X spreads as a smooth S -scheme of finite type \mathcal{X} for a suitable smooth model S of K over \mathbf{F}_p . Using this remark, one sees that Friedlander's conjecture potentially implies statements concerning K -varieties: this is one of its many interests. For example, one easily sees that it implies Conjecture 2.4.2.2 of [5] on rational K -groups of fields of positive characteristic, see [46, 8.28 (v) and 8.30] and [50, Th. 60 and 61].

3.4. Motivic reformulations. Since the appearance of [43], two major progresses have been made: 1) de Jong's theorem on alterations [41], and 2) definitions of motivic cohomology, together with motivic l -adic cycle class maps.

Motivic cohomology groups generalise Chow groups, and they agree rationally with Adams eigenspaces on algebraic K -theory. For smooth varieties there are mainly two versions of these groups: Bloch's higher Chow groups [9] and the Suslin-Voevodsky motivic cohomology groups. They are shown to be canonically (but not very directly) isomorphic over any field in [79].

In [46], I was interested in formulating a version of Conjecture 3.3.2 using motivic cohomology. I used the Suslin-Voevodsky version, but assumed resolution of singularities (over finite fields!) because it was built in Voevodsky's theory of triangulated motives at the time. To get around resolution, I changed course and used higher Chow groups in [47] and [48]. Here I want to give a version of Friedlander's conjecture 3.3.2 involving

realisation functors on Voevodsky's $DM(\mathbf{F})$, thus going back to the original viewpoint of using Suslin-Voevodsky motivic cohomology. It has several advantages:

- (1) While Bloch's theory avoids resolution of singularities, it relies on a subtle sophistication of Chow's moving lemma which renders the functoriality of his cycles complexes (as opposed to their homology groups) delicate. By contrast, the motivic complexes of Suslin and Voevodsky have a straightforward functoriality.
- (2) Much of the resolution of singularities needed in the basics of Voevodsky's theory has been removed. On the one hand, he proved very simply the cancellation theorem over any perfect field [82]. On the other hand, de Jong's alteration theorem turns out to be sufficient for most of the development, even integrally.
- (3) Some of the remaining original constructions of Voevodsky, which still require resolution of singularities, like the motives (with or without compact supports) of singular schemes, can now be bypassed by using the 4 operations., see §6.7.
- (4) The triangulated framework is much richer than the one involving smooth varieties: for example, it automatically yields conjectural statements for the cohomology and cohomology with compact supports of possibly singular varieties in the spirit of Jannsen, see (3).

3.5. A reading guide. Let me now outline the sequel of this paper. I tried to go step by step and by increasing degree of sophistication:

- (1) In §4, I introduce motivic cohomology à la Suslin-Voevodsky by using their concrete motivic complexes; the construction of the l -adic cycle class map is then very easy but insufficient.
- (2) In §5, I introduce Voevodsky's categories of motives over a field; the motivic cohomology of §4 may then be interpreted as Hom groups in these categories (Theorem 5.2.1 and 5.5.1).
- (3) In §6, I describe the situation over a base, and in particular outline the formalism of six operations. Here the story becomes more complicated as several versions of categories of motives come into play. The main point is that they often agree: see §6.5. An important outcome is the construction of motives (with or without compact supports) of singular varieties, with the right functoriality: see §6.7. The l -adic realisation functor is finally introduced in §6.8.
- (4) §7 is technical but essential to get Theorem 1 of the introduction.
- (5) In §8, I prove a more precise version of Geisser's Theorem 3.2.1. The main reason why I did this was to convince myself that there is no way to get the equivalence (ii) \iff (iv) of Corollary 9.8.4 " n by n ".

- (6) In §9, we get slowly towards the motivic formulation of Friedlander’s conjecture and the proof of Theorem 1: this is achieved in §9.8, see Proposition 9.8.2, Theorem 9.8.3 and Corollary 9.8.4. In particular, the conditions of Theorem 1 are equivalent to Friedlander’s conjecture plus a generalisation of Condition S^n of Remark 2.2.3. A simple lemma of Ayoub (Lemma 9.3.1) drastically simplifies an argument I had outlined in my 2006 talk.
- (7) In §10, I give two other reformulations of these conjectures: a) a realisation functor from a suitable category is fully faithful, b) the Hom groups of this category are finitely generated. See Proposition 10.3.3 and Theorem 10.4.2. This involves an apparently artificial construction, which is better explained in the next section.
- (8) In §11, I recall Lichtenbaum’s Weil-étale topology and inflict it to DM. This sheds a better light on the previous section, but these new categories have their shortcomings.

4. MOTIVIC COHOMOLOGY AND l -ADIC COHOMOLOGY

4.1. Motivic cohomology, Nisnevich and étale. Let us first recall Voevodsky’s groups of *finite correspondences* [78]: if X, Y are smooth k -schemes, we denote by $c(X, Y)$ the free abelian group with basis the closed integral subschemes $Z \subseteq X \times Y$ such that the projection $Z \rightarrow X$ is finite and surjective on some connected component of X . The beauty of finite correspondences is that they compose “on the nose”, that is, without having to mod out by any adequate equivalence relation. In particular, they are covariant in Y and contravariant in X .

If $(Y_1, y_1), (Y_2, y_2)$ are two pointed smooth k -schemes ($y_i \in Y_i(k)$), we define

$$c(X, (Y_1, y_1) \wedge (Y_2, y_2)) = \text{Coker}(c(X, Y_1 \times y_2) \oplus c(X, y_1 \times Y_2) \rightarrow c(X, Y_1 \times Y_2))$$

and more generally, we define $c(X, (Y_1, y_1) \wedge \cdots \wedge (Y_n, y_n))$ by a similar formula: this is a direct summand of $c(X, Y_1 \times \cdots \times Y_n)$.

We denote by Δ be the standard cosimplicial k -scheme, with

$$\Delta^n = \text{Spec } k[t_0, \dots, t_n] / (\sum t_i - 1).$$

Let now $X \in \text{Sm}(k)$, and let $n \geq 0$. The n -th *Suslin-Voevodsky complex* is the complex whose i -th term is

$$C^i(X, \mathbf{Z}(n)) = c(X \times \Delta^{n-i}, \mathbb{G}_m^{\wedge n})$$

where $\mathbb{G}_m^{\wedge n} = (\mathbb{G}_m, 1) \wedge \cdots \wedge (\mathbb{G}_m, 1)$, and the differentials are as usual.

Varying X , this defines a complex of presheaves (even sheaves) over the étale site on $\mathrm{Sm}(k)$, and *a fortiori* over the corresponding Nisnevich site. Let us denote this complex by $\underline{C}^*(\mathbf{Z}(n))$. We set

Definition 4.1.1. a) $H^i(X, \mathbf{Z}(n)) = \mathbb{H}_{\mathrm{Nis}}^i(X, \underline{C}^*(\mathbf{Z}(n)))$. This is *motivic cohomology*.

b) $H_{\mathrm{ét}}^i(X, \mathbf{Z}(n)) = \mathbb{H}_{\mathrm{ét}}^i(X, \underline{C}^*(\mathbf{Z}(n)))$. This is *étale motivic cohomology*.

The link between motivic cohomology and Chow groups is the following:

Proposition 4.1.2. *One has isomorphisms*

$$CH^n(X) \simeq H^{2n}(X, \mathbf{Z}(n)).$$

More generally, motivic cohomology groups (for the Nisnevich topology) are isomorphic to Bloch's higher Chow groups, with a suitable reindexing [79]. This also implies:

$$(4.1.1) \quad H^i(X, \mathbf{Z}(n)) = 0 \text{ for } i > 2n.$$

4.2. Continuous étale cohomology. We now assume that $k = \mathbf{F}$ is finite until the end of this section.

Let $X \in \mathrm{Sm}(\mathbf{F})$. We write $H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n))$ for Jannsen's *continuous étale cohomology* [42]: recall that if (A_n) is an inverse system of étale sheaves, $H_{\mathrm{cont}}^i(X, (A_n))$ is defined as the i -th derived functor of the composite functor

$$(A_n) \mapsto (\Gamma(X, A_n)) \mapsto \varprojlim \Gamma(X, A_n).$$

We also write $H_{\mathrm{cont}}^i(X, \mathbf{Q}_l(n)) := H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)) \otimes \mathbf{Q}$.

Since étale cohomology of X with finite coefficients is finite, this coincides with the naïve l -adic cohomology:

$$(4.2.1) \quad H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)) \xrightarrow{\sim} \varprojlim H_{\mathrm{ét}}^i(X, \mu_l^{\otimes n}).$$

The Hochschild-Serre spectral sequence yields short exact sequences:

$$(4.2.2) \quad 0 \rightarrow H_{\mathrm{cont}}^{i-1}(\bar{X}, \mathbf{Z}_l(n))_G \rightarrow H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)) \rightarrow H_{\mathrm{cont}}^i(\bar{X}, \mathbf{Z}_l(n))^G \rightarrow 0.$$

4.3. An l -adic lemma. Let

$$(4.3.1) \quad e \in H_{\mathrm{cont}}^1(\mathbf{F}, \hat{\mathbf{Z}}) \simeq \mathrm{Hom}_{\mathrm{cont}}(\hat{\mathbf{Z}}, \hat{\mathbf{Z}}) \simeq \hat{\mathbf{Z}}$$

be the canonical generator corresponding to the identity, where the first isomorphism is given by the arithmetic Frobenius of \mathbf{F} . Note that $e^2 = 0$ since $cd(\mathbf{F}) = 1$. The following generalises [58, Prop. 6.5] (the proof is the same as for [46, Prop. 6.5]):

Lemma 4.3.1. *Let $C \in \hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)$ be a perfect complex of l -adic sheaves over $\mathrm{Spec} \mathbf{F}$. Then the diagram*

$$(4.3.2) \quad \begin{array}{ccc} H_{\mathrm{cont}}^j(\mathbf{F}, C) & \xrightarrow{a} & H^j(C)^G \\ \downarrow \cdot c & & \searrow i \\ & & H^j(C) \\ & & \swarrow p \\ H_{\mathrm{cont}}^{j+1}(\mathbf{F}, C) & \xleftarrow{b} & H^j(C)_G \end{array}$$

commutes for any $j \in \mathbf{Z}$. Here i is the inclusion, p is the projection and a, b are the maps stemming from the hypercohomology spectral sequence for $H_{\mathrm{cont}}^*(\mathbf{F}, C)$. \square

4.4. Naïve cycle class maps. The rigidity theorem of Suslin-Voevodsky ([72, Th. 7.6] or [80, Th. 6.1]) yields canonical isomorphisms

$$(4.4.1) \quad H_{\mathrm{ét}}^i(X, \mathbf{Z}(n) \otimes \mathbf{Z}/l^\nu) \simeq H_{\mathrm{ét}}^i(X, \mu_{l^\nu}^{\otimes n})$$

which, in view of (4.2.1), immediately yield higher l -adic cycle class maps:

$$(4.4.2) \quad H_{\mathrm{ét}}^j(X, \mathbf{Z}(n)) \otimes \mathbf{Z}_l \rightarrow H_{\mathrm{cont}}^j(X, \mathbf{Z}_l(n)).$$

One has isomorphisms [9]

$$K_{2n-i}(X)_{\mathbf{Q}}^{(n)} \simeq H^j(X, \mathbf{Q}(n)) \xrightarrow{\sim} H_{\mathrm{ét}}^j(X, \mathbf{Q}(n)),$$

which convert the maps (4.4.2) into the Chern characters $ch_{n,i}$ of Conjecture 3.3.2. Thus, this construction already allows us to reformulate Conjecture 3.3.2 in terms of motivic cohomology. However it is too crude to provide a correct *integral* version of this conjecture (see §9.1 for a detailed discussion). For this, one needs a triangulated version of (4.4.2), namely a map

$$\mathbf{Z}(n)_{\mathrm{ét}} \otimes \mathbf{Z}_l \rightarrow R \varprojlim \mu_{l^\nu}^{\otimes n}$$

in a suitable triangulated category, which has to be modified. In [47], this was done in the derived category of étale sheaves on $Sm(\mathbf{F})$, using Bloch's cycle complexes instead of the Suslin-Voevodsky complexes. Since we are going to ultimately give a reformulation using $\mathrm{DM}_{\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})$, it is best to construct the above morphism directly in this category. This will be done in §9.6.

5. CATEGORIES OF MOTIVES OVER A FIELD

5.1. Categories of effective geometric motives. Let $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ be Voevodsky's tensor triangulated category of effective geometric motives [78]: we also refer to Levine's survey [54, Lect. 1 and 2] for a nice exposition. Then

any $X \in \text{Sm}(k)$ has a *motive* $M(X) \in \text{DM}_{\text{gm}}^{\text{eff}}(k)$. The functor $X \mapsto M(X)$ has the following properties:

Homotopy invariance: $M(X \times \mathbf{A}^1) \xrightarrow{\sim} M(X)$.

Mayer Vietoris: If $X = U \cup V$ with U, V open, then one has an exact triangle

$$M(U \cap V) \rightarrow M(U) \oplus M(V) \rightarrow M(X) \xrightarrow{+1}.$$

Projective bundle formula: Write $\mathbf{Z}(1)$ for the direct summand of $M(\mathbf{P}^1)[-2]$ defined by any rational point: this is the *Tate object*. For $n > 0$, define $\mathbf{Z}(n) = \mathbf{Z}(1)^{\otimes n}$: if $M \in \text{DM}_{\text{gm}}^{\text{eff}}(k)$, we write $M(n) := M \otimes \mathbf{Z}(n)$. Now let $E \rightarrow X$ be a vector bundle of rank r . Then

$$M(\mathbf{P}(E)) \simeq \bigoplus_{n=0}^{r-1} M(X)(n)[2n].$$

Gysin: Let $Z \subset X$ be a closed subscheme, smooth of codimension c , and let $U = X - Z$. Then one has an exact triangle

$$M(U) \rightarrow M(X) \rightarrow M(Z)(c)[2c] \xrightarrow{+1}.$$

Let us recall the construction of $\text{DM}_{\text{gm}}^{\text{eff}}(k)$ in two words: it parallels the construction of Chow motives to a large extent, but is based on smooth, not necessarily projective, varieties. We start from the category $\text{Cor}(k)$ of *finite correspondences* over k : its objects are smooth k -schemes of finite type and its morphisms are the groups $c(X, Y)$ of §4.1. The graph of a morphism is a finite correspondence, which yields a (covariant) functor $\text{Sm}(k) \rightarrow \text{Cor}(k)$.

Since $\text{Cor}(k)$ is an additive category, one may consider the homotopy category $K^b(\text{Cor}(k))$ of bounded chain complexes of objects of $\text{Cor}(k)$. Then $\text{DM}_{\text{gm}}^{\text{eff}}(k)$ is defined as the pseudo-abelian envelope of the Verdier localisation of $K^b(\text{Cor}(k))$ with respect to the “relations” corresponding to homotopy invariance and Mayer-Vietoris as above. This yields a string of functors:

$$\text{Sm}(k) \rightarrow \text{Cor}(k) \rightarrow K^b(\text{Cor}(k)) \rightarrow \text{DM}_{\text{gm}}^{\text{eff}}(k).$$

Their composition is the above functor M .

5.2. Categories of motivic complexes, Nisnevich topology. The relationship between motivic cohomology and $\text{DM}_{\text{gm}}^{\text{eff}}(k)$ is

Theorem 5.2.1 ([78]). *One has canonical isomorphisms*

$$H^i(X, \mathbf{Z}(n)) \simeq \text{Hom}_{\text{DM}_{\text{gm}}^{\text{eff}}(k)}(M(X), \mathbf{Z}(n)[i]).$$

This is proven by introducing a larger tensor triangulated category $\mathrm{DM}_-^{\mathrm{eff}}(k)$, constructing a fully faithful functor $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k) \rightarrow \mathrm{DM}_-^{\mathrm{eff}}(k)$ and proving Theorem 5.2.1 in $\mathrm{DM}_-^{\mathrm{eff}}(k)$.

In short, $\mathrm{DM}_-^{\mathrm{eff}}(k)$ is constructed out of Nisnevich sheaves with transfers. A *presheaf with transfers* \mathcal{F} is simply an additive contravariant functor from the category $\mathrm{Cor}(k)$ of the previous subsection to abelian groups; \mathcal{F} is a *Nisnevich sheaf with transfers* if, viewed as a functor on $\mathrm{Sm}(k)$, it is a sheaf for the Nisnevich topology. Write $\mathrm{NST}(k)$ for the abelian category of Nisnevich sheaves with transfers. Inside the bounded above derived category $D^-(\mathrm{NST}(k))$, Voevodsky considers the full subcategory consisting of those complexes whose homology sheaves are *homotopy invariant*: this is $\mathrm{DM}_-^{\mathrm{eff}}(k)$. The embedding of $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ into $\mathrm{DM}_-^{\mathrm{eff}}(k)$ may then be thought of as a Yoneda embedding: it is constructed using complexes $\underline{C}_*(\mathcal{F})$ generalising those described in §4.1.

The category $\mathrm{DM}_-^{\mathrm{eff}}(k)$ is constructed out of bounded above complexes of Nisnevich sheaves. It will be more convenient for us to work with the corresponding category $\mathrm{DM}^{\mathrm{eff}}(k)$ constructed in the same way but without boundedness conditions. In fact we have a commutative square of inclusions of tensor triangulated categories

$$(5.2.1) \quad \begin{array}{ccc} \mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k) & \xrightarrow{a} & \mathrm{DM}^{\mathrm{eff}}(k) \\ c \downarrow & & d \downarrow \\ \mathrm{DM}_{\mathrm{gm}}(k) & \xrightarrow{b} & \mathrm{DM}(k). \end{array}$$

The two bottom categories are obtained from the top categories by inverting the Tate object, but in rather different ways. On the left hand side, one defines $\mathrm{DM}_{\mathrm{gm}}(k)$ as the category whose objects are pairs (M, m) with $M \in \mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ and $m \in \mathbf{Z}$, and Hom sets are defined by

$$\mathrm{Hom}_{\mathrm{DM}_{\mathrm{gm}}(k)}((M, m), (N, n)) = \varinjlim_{r \geq -m, -n} \mathrm{Hom}_{\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)}(M(m+r), N(n+r))$$

cf. [78, p. 192]. The full faithfulness of a follows from [the proof of] [78, Th. 3.2.6] and the full faithfulness of c follows from [82]; both results use the perfectness of the field k .

On the other hand, if we try and define $\mathrm{DM}(k)$ from $\mathrm{DM}^{\mathrm{eff}}(k)$ in the same way, the category we get is too small: in particular it does not have infinite direct sums, while $\mathrm{DM}^{\mathrm{eff}}(k)$ does. The solution is to pass to the model-categorical level and imitate the construction of spectra in algebraic topology: namely one considers “ $\mathbf{Z}(1)$ -spectra” of the form

$$(C_n, e_n)_{n \in \mathbf{Z}}$$

where $C_n \in C(NST)$ and e_n is a morphism of complexes $C_n \otimes \mathbf{Z}(1) \rightarrow C_{n+1}$ for $\mathbf{Z}(1)$ as in §4.1. This is outlined by Morel in [62, §5.2], as an analogue to the construction given by Voevodsky in [77, §5] for the stable homotopy category of schemes. A detailed construction, over a base, is given by Cisinski and Déglise in [14, Ex. 7.15].

One can show that the functors b and d are also fully faithful. The categories $\mathrm{DM}^{\mathrm{eff}}(k)$ and $\mathrm{DM}(k)$ are closed, i.e., have internal Homs.

5.3. Variant. If A is a commutative ring, one may define similar categories of motives with coefficients in A ; we shall denote them by $\mathrm{DM}(k, A)$ and the decorated analogues. The most important cases for us will be $A = \mathbf{Q}$, $A = \mathbf{Z}/n$ and, later, $A = \mathbf{Z}_l$.

5.4. Remark on notation. I will never write $\mathrm{DM}(k, \mathbf{Z})$ for $\mathrm{DM}(k)$. There is a dilemma (or a trilemma) for writing the unit object of $\mathrm{DM}(k, A)$. A spontaneous choice would be A , but this leads to ambiguities. One ambiguity which would be disastrous here would be to confuse notationally the complex of sheaves $\mathbf{Z}(n) \otimes \mathbf{Z}_l$ and the l -adic sheaf $\mathbf{Z}_l(n)$.

Another choice which has been made elsewhere to avoid this problem is to use the neutral notation $\mathbf{1}$ for the unit object. This leads to notation like $\mathbf{1}(n)[i]$, which I find disgraceous.

For these reasons, I have decided to retain the notation \mathbf{Z} for the unit object of $\mathrm{DM}(k, A)$ even when A is another ring than \mathbf{Z} . Similarly, the notation $\mathbf{Z}(n)$ is retained in $\mathrm{DM}(k, A)$ for any A . This means that $\mathrm{DM}(k, A)(\mathbf{Z}, \mathbf{Z}) = A$ and $\mathrm{DM}(k, A)(M(X), \mathbf{Z}(n)[i]) = H^i(X, \mathbf{Z}(n)) \otimes A$ for $X \in \mathrm{Sm}(k)$.

This also applies to the variants of $\mathrm{DM}(k)$ to come.

On the other hand, I will sometimes write $H^i(X, A(n))$ for $H^i(X, \mathbf{Z}(n)) \otimes A$ when there is no risk of ambiguity, e.g. for $A = \mathbf{Q}$ or \mathbf{Z}/m (but not $A = \mathbf{Z}_l$).

I hope this will not create any confusion.

5.5. Categories of motivic complexes, étale topology. Let A be a commutative ring. There is an étale analogue to the inclusion $\mathrm{DM}^{\mathrm{eff}}(k, A) \hookrightarrow \mathrm{DM}(k, A)$, using the étale topology rather than the Nisnevich topology. This yields a naturally commutative diagram of tenros triangulated categories:

$$(5.5.1) \quad \begin{array}{ccc} \mathrm{DM}^{\mathrm{eff}}(k, A) & \xrightarrow{\alpha^*} & \mathrm{DM}_{\mathrm{\acute{e}t}}^{\mathrm{eff}}(k, A) \\ d \downarrow & & d_{\mathrm{\acute{e}t}} \downarrow \\ \mathrm{DM}(k, A) & \xrightarrow{\alpha^*} & \mathrm{DM}_{\mathrm{\acute{e}t}}(k, A) \end{array}$$

where α^* denotes the change of topology functor. The functor $d_{\mathrm{\acute{e}t}}$ is still fully faithful.

The basic results on the effective categories are identical to those stated in [78, Prop. 3.3.2 and 3.3.3] for their bounded versions:

Theorem 5.5.1. *Assume that the field k has finite cohomological dimension, and exponential characteristic p . Then:*

a) *For $A = \mathbf{Q}$, the functors α^* of (5.5.1) are equivalences of tensor triangulated categories.*

b) *for $n > 0$ prime to p , the natural functor*

$$\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k, \mathbf{Z}/n) \rightarrow D(k_{\acute{e}t}, \mathbf{Z}/n)$$

is an equivalence of triangulated categories, where the right hand side is the derived category of sheaves of \mathbf{Z}/n -modules on the small étale site of $\mathrm{Spec} k$ and the functor is induced by restriction to the small étale site.

c) *For $\nu \geq 1$, the categories $\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k, \mathbf{Z}/p^\nu)$ and $\mathrm{DM}_{\acute{e}t}(k, \mathbf{Z}/p^\nu)$ are 0.*

Part c) of Theorem 5.5.1 means that $\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k)$ and $\mathrm{DM}_{\acute{e}t}(k)$ are $\mathbf{Z}[1/p]$ -linear; the basic reason why this follows from homotopy invariance is the Artin-Schreier exact sequence (of étale shaves with transfers)

$$0 \rightarrow \mathbf{Z}/p \rightarrow \mathbb{G}_a \xrightarrow{F-1} \mathbb{G}_a \rightarrow 0.$$

One deduces a pendant to Theorem 5.2.1, plus an important comparison between motivic and étale motivic cohomology:

Theorem 5.5.2. a) *One has canonical isomorphisms*

$$H_{\acute{e}t}^i(X, \mathbf{Z}(n)) \otimes \mathbf{Z}[1/p] \simeq \mathrm{Hom}_{\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k)}(M(X), \mathbf{Z}(n)[i]).$$

b) *For any smooth X and any $n \geq 0$, $i \in \mathbf{Z}$, the natural map*

$$H^i(X, \mathbf{Z}(n)) \otimes \mathbf{Q} \rightarrow H_{\acute{e}t}^i(X, \mathbf{Z}(n)) \otimes \mathbf{Q}$$

is an isomorphism.

An upshot is that if we work with DM, we are bound to lose p -torsion information on étale motivic cohomology. Using the groups from Definition 4.1.1 b) retains this information, but we lose the power of the triangulated formalism.⁵

We shall also need:

Lemma 5.5.3. *Let $\mathrm{DM}_{\mathrm{gm},\acute{e}t}(k)$ be the full subcategory of $\mathrm{DM}_{\acute{e}t}(k)$ which is the pseudo-abelian envelope of the image of $\mathrm{DM}_{\mathrm{gm}}(k, \mathbf{Z}[1/p]) \rightarrow \mathrm{DM}_{\acute{e}t}(k)$. Then $\mathrm{DM}_{\mathrm{gm},\acute{e}t}(k)$ is a rigid subcategory of $\mathrm{DM}_{\acute{e}t}(k)$: every object is strongly dualisable.*

⁵To recover this power for the p -torsion, one would have to invent new triangulated categories which would accomodate non homotopy invariant phenomena: this is very much work in progress now.

(Recall that an object M of a unital symmetric monoidal category \mathcal{A} is *strongly dualisable* if there exists a triple (M^*, η, ϵ) with $M^* \in \mathcal{A}$ and $\eta : \mathbf{1} \rightarrow M \otimes M^*$, $\epsilon : M^* \otimes M \rightarrow \mathbf{1}$, verifying adjunction-like identities. Such a triple is then unique up to unique isomorphism; see [20].)

Proof. Since $\mathrm{DM}_{\mathrm{gm}}(k, \mathbf{Z}[1/p]) \rightarrow \mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(k)$ is a (strict) \otimes -functor, it suffices to show that $\mathrm{DM}_{\mathrm{gm}}(k, \mathbf{Z}[1/p])$ is rigid. But this triangulated category is generated by motives of smooth projective varieties, as follows from Gabber’s refinement of de Jong’s theorem, and these objects are strongly dualisable (cf. [34, App. B]). \square

5.6. Motivic complexes in DM. They are easy to understand in weight ≤ 1 :

Theorem 5.6.1 (see [78, Cor. 3.4.3]). *One has isomorphisms in $\mathrm{DM}^{\mathrm{eff}}(k)$*

$$\begin{aligned} \mathbf{Z}(0) &\simeq \mathbf{Z} \\ \mathbf{Z}(1) &\simeq \mathbb{G}_m[-1] \end{aligned}$$

where \mathbb{G}_m is viewed as a homotopy invariant Nisnevich sheaf with transfers.

These quasi-isomorphisms persist when passing to $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(k)$: we shall put an index $\acute{\mathrm{e}}\mathrm{t}$ for clarity. In particular, if m is invertible in k , multiplication by m gives a “Kummer” isomorphism

$$\mu_m \xrightarrow{\sim} \mathbf{Z}/m(1)_{\acute{\mathrm{e}}\mathrm{t}}$$

hence morphisms

$$(5.6.1) \quad \mu_m^{\otimes n} \rightarrow \mathbf{Z}/m(n)_{\acute{\mathrm{e}}\mathrm{t}}$$

in the category $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(k)$. The following reformulation of (4.4.1) is a special case of Theorem 5.5.1 b) (see also [80, Th. 6.1]):

Theorem 5.6.2. (5.6.1) is an isomorphism for any $n \geq 0$.

5.7. The l -adic realisation functor. Let us mention immediately that, for a prime number l invertible in k , there exists an l -adic realisation \otimes -functor

$$R_l : \mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(k) \rightarrow \hat{D}_{\acute{\mathrm{e}}\mathrm{t}}(k, \mathbf{Z}_l)$$

where the right hand side is the category defined by Ekedahl in [22]. Intuitively, its restriction to $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(k)$ is obtained as the “inverse limit” of the compositions

$$\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(k) \rightarrow \mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(k, \mathbf{Z}/l^n) \simeq D(k_{\acute{\mathrm{e}}\mathrm{t}}, \mathbf{Z}/l^n)$$

cf. Theorem 5.5.1 b). This functor commutes with infinite direct sums, which implies formally that it has a right adjoint Ω_l . When k is a finite field, the motivic reformulation of Friedlander’s conjecture will rely on this pair of adjoints and its relationship with the subcategory $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}^{\mathrm{eff}}(k)$.

For details on the definition of R_l , I refer to the next section.

6. CATEGORIES OF MOTIVES OVER A BASE

In this section, S is a Noetherian separated scheme.

6.1. Generalising Voevodsky’s definition over a field. This presents no basic difficulty: in short, one replaces the category $\text{Cor}(k)$ used in the previous section by a category $\text{Cor}(S)$ based on the theory of relative cycles developed by Suslin and Voevodsky in [73]. The description of $\text{Cor}(S)$ is outlined in [57, App. 1A], and in more detail in [82, §2]. The construction of a category $\text{DM}_-^{\text{eff}}(S)$ (even for S a simplicial scheme!) is then given in [81].

Ivorra’s exposition [40] develops this a bit further in §2, and introduces a corresponding category of geometric motives $\text{DM}_{\text{gm}}^{\text{eff}}(S)$ in §4: the only difference with the case of a field is that the Mayer-Vietoris relation is generalised to “Mayer-Vietoris for the Nisnevich topology”. Finally, Ivorra constructs a fully faithful functor $\text{DM}_{\text{gm}}^{\text{eff}}(S) \hookrightarrow \text{DM}_-^{\text{eff}}(S)$ in §5, just as in the case of a field.

Another exposition of the theory of finite correspondences, in the case of a regular base, is given by Déglise in [18]. He then gives a general exposition with Cisinski in [15, §8], that they use in *loc. cit.*, §10 to construct categories $\text{DM}_{\text{gm}}^{\text{eff}}(S)$ and $\text{DM}^{\text{eff}}(S)$ as above, but also stable versions $\text{DM}_{\text{gm}}(S)$ and $\text{DM}(S)$. (In [15], the index gm is replaced by c .) In this way, the picture of §§5.1 and 5.2 is extended to a general base. The only difference is that, in the diagram

$$\begin{array}{ccc} \text{DM}_{\text{gm}}^{\text{eff}}(S) & \xrightarrow{a} & \text{DM}^{\text{eff}}(S) \\ c \downarrow & & d \downarrow \\ \text{DM}_{\text{gm}}(S) & \xrightarrow{b} & \text{DM}(S) \end{array}$$

generalising (5.2.1), the horizontal functors are still fully faithful but it is not known whether the vertical functors are.

6.2. Where the buck stops. So far we seem to live in the best possible world, having fully extended Voevodsky’s theory over a base. Unfortunately, one runs into trouble in two situations:

- (1) When trying to apply to this construction the yoga of six operations of Voevodsky-Ayoub.
- (2) Similarly for realisation functors, most importantly the l -adic realisation functors.

In (1), the aim would be to endow $S \mapsto \mathrm{DM}(S)$ with the six functors of Grothendieck: $f^*, f_*, f_!, f^!, \otimes, \underline{\mathrm{Hom}}$ where $f : S' \rightarrow S$ is a morphism of Noetherian schemes (cf. [35, §2]). The strategy of Grothendieck, Verdier and Deligne for coherent and étale cohomology is awkward here, mainly because Voevodsky works with big sites rather than small sites. For this reason, Voevodsky proposed another axiomatic strategy which fits better in his context.

Roughly speaking, his framework is the following. One starts from associating to any (reasonable) scheme S a triangulated category $\mathbb{H}(S)$; one assumes that some of the operations are already constructed and verify certain axioms. From this input, one can then deduce the other operations and all their properties.

After Voevodsky sketched this programme for $f^*, f_*, f_!, f^!$, it was worked out by Ayoub in [2, Ch. 1] (see his 1.4.1 for a precise statement of the above).

Voevodsky had in mind primarily the stable homotopy categories of schemes $\mathrm{SH}(S)$ that he developed jointly with Morel [63, 77]. In [2, Ch. 4], Ayoub gives examples of theories $S \mapsto \mathbb{H}(S)$ which verify his axioms 1.4.1. They include SH , but not DM .

The problem with DM is that transfers make it difficult to verify the axiom of “locality”. This axiom (number 4 in [2, 1.4.1]) roughly requires that if Z is a closed subset of S with open complement U , one can obtain $\mathbb{H}(S)$ by glueing $\mathbb{H}(Z)$ and $\mathbb{H}(U)$.

The issue for (2): constructing l -adic realisation functors, is to extend the rigidity theorem of Suslin-Voevodsky over a base. For this, a strategy to reduce to the case of a perfect base field treated in [72] is obstructed by the same axiom of locality.

6.3. Where the buck restarts: motives without transfers. On the other hand, the examples of [2, Ch. 4] do include *motives without transfers*. If τ is a suitable topology, the category $\mathrm{DA}_\tau(S)$ is constructed as $\mathrm{DM}_\tau(S)$, except that one replaces the category $\mathrm{Cor}(S)$ by the simpler category $\mathrm{Sm}(S)$ (more correctly: $\mathbf{Z}[\mathrm{Sm}(S)]$) throughout. We shall use the established practice of dropping τ if it is the Nisnevich topology. For S the spectrum of a field k , this construction appears first in [62, §5.2] with the notation $\widetilde{\mathrm{DM}}(k)$.

That this theory satisfies locality is the contents of [2, Cor. 4.5.44]⁶, whose proof is inspired by that of [63, Th. 2.21]. An essentially self-contained construction of DA_τ is given in [4, §3].

⁶The notation of [2] may be confusing: compare [4, App. A].

As above, one may define this theory with coefficients in a commutative ring A ; there is a natural \otimes -functor

$$DA_\tau(S, A) \rightarrow DM_\tau(S, A).$$

In [15, §15.2], Cisinski and Déglise prove that this functor is an equivalence of categories when $A = \mathbf{Q}$ and $\tau = \text{ét}$ if S is excellent and geometrically unibranch. In [4, Th. B.1], Ayoub extends this to the case where any prime number is invertible either in A or on S , provided S is universally Japanese, normal and of finite étale cohomological dimension. (There is an extra minor condition, and 2 must be inverted unless the characteristic is 0.) In this way, the formalism of the 4 operations is largely repaired.

If S has characteristic $p > 0$, we may choose $A = \mathbf{Z}[1/2p]$. But the Artin-Schreier argument of Theorem 5.5.1 c) shows that

$$DM_{\text{ét}}(S) \rightarrow DM_{\text{ét}}(S, \mathbf{Z}[1/p])$$

is an equivalence of categories, and the same works for $DA_{\text{ét}}$. So the comparison theorem holds with coefficients $\mathbf{Z}[1/2]$ in this case. If S is a \mathbf{Q} -scheme, then it holds with integer coefficients.

6.4. A competitor: Beilinson motives. In [15], Cisinski and Déglise do prove the locality axiom outlined in §6.2 for DM when S and Z are smooth over a common base. They can relax this smoothness assumption provided one works with rational coefficients. To this aim, they introduce yet another theory: *Beilinson motives*.

Roughly speaking, Beilinson motives are modules over the K -theory spectrum, an object of SH which represents algebraic K -theory. They form a \mathbf{Q} -linear tensor triangulated category $DM_{\mathbb{B}}(S)$, defined for any reasonable S . The locality axiom for SH then implies rather formally the locality axiom for $DM_{\mathbb{B}}$.

They also construct an equivalence of \otimes -triangulated categories

$$DM_{\mathbb{B}}(S) \simeq DM(S, \mathbf{Q})$$

for any excellent and geometrically unibranch scheme S [15, Th. 8 and 9], which extends the locality axiom for $DM(-, \mathbf{Q})$ to these schemes.

Finally, they prove an equivalence

$$DM_{\mathbb{B}}(S) \simeq DA_{\text{ét}}(S, \mathbf{Q})$$

for any S [15, Th. 13], which connects with the results explained in §6.3.

6.5. Recapitulation. Since the above discussion may have confused the reader, let me summarise it:

- (1) There are three types of triangulated theories of motives on the market, associated to a reasonable scheme S : $DM_\tau(S, A)$, $DA_\tau(S, A)$, $DM_{\mathbb{B}}(S)$. Here τ denotes a suitable Grothendieck topology (usually Nisnevich or étale) and A denotes a commutative ring. The categories $DM_{\mathbb{B}}(S)$ are \mathbf{Q} -linear.

(This does not exhaust the triangulated theories of motives on the market: there are also the versions of Levine and Hanamura, see §1.1. Here I limit myself to those which are constructed in the spirit of Voevodsky and Morel.)

- (2) For any S , there is an equivalence of \otimes -categories $DM_{\mathbb{B}}(S) \simeq DA_{\text{ét}}(S, \mathbf{Q})$.
- (3) Suppose for simplicity that S is a k -scheme for a field k of exponential characteristic p . Then the natural \otimes -functor

$$DA_{\text{ét}}(S, \mathbf{Z}[1/2]) \rightarrow DM_{\text{ét}}(S, \mathbf{Z}[1/2])$$

is an equivalence of categories provided S is normal and universally Japanese. If $p = 1$, this holds with integer coefficients.

- (4) The assignment $S \mapsto DA_\tau(S, A)$ satisfies the formalism of the six operations of Voevodsky-Ayoub (for $\tau = \text{Nis}, \text{ét}$).

6.6. Some philosophy. It may be disturbing that one has to replace the beautiful categories of motives based on finite correspondences by apparently uglier categories in order to have a working theory. In fact, this is not so serious. What matters is that this theory coincides with $DM_{\text{ét}}(-, A)$ in important cases, especially when the base is a field. In this way, the connection with algebraic cycles is not lost. As for other bases, I like to think of some theory $S \mapsto \mathbb{H}(S)$ extending $k \mapsto DM_{\text{ét}}(k, A)$ as a black box, which is helpful for applying the six functors formalism but whose specific description for a general S is not so important. An example is given just below.

6.7. Application: motives of singular varieties. Let k be a field. In [78, §4.1], Voevodsky associates to any k -scheme X of finite type two objects $\underline{C}_*(X)$ and $\underline{C}_*^c(X)$ of $DM_{\text{ét}}^{\text{eff}}(k)$. These objects play the respective rôles of the motive and motive with compact supports of X . Unfortunately, the proof of their functoriality rests on [78, Th. 4.1.2], whose proof uses Hironaka's resolution of singularities in a central way, so is valid only in characteristic 0. Under this condition, the functorialities show that these objects belong to $DM_{\text{gm}}^{\text{eff}}(k)$, by reduction to the case of smooth projective schemes.

As I first learned from Joël Riou, the six operations allow us to define such objects in nonzero characteristic at least in the étale categories, by avoiding Hironaka resolution:

Definition 6.7.1. Let τ be the Nisnevich or the étale topology, and let $p_X : X \rightarrow \text{Spec } k$ be a k -scheme of finite type. We define:

- (1) Its motive: $M(X) = \underline{\text{Hom}}((p_X)_* \mathbf{Z}_X, \mathbf{Z}) \in \text{DA}_\tau(k)$.
- (2) Its motive with compact supports: $M_c(X) = \underline{\text{Hom}}((p_X)_! \mathbf{Z}_X, \mathbf{Z}) \in \text{DA}_\tau(k)$.

Proposition 6.7.2. *a) The motive is covariant for any morphism.
b) The motive with compact supports is covariant for projective morphisms and contravariant for étale morphisms.
c) Let X be a k -scheme of finite type, $i : Z \rightarrow X$ a closed immersion and $j : U \rightarrow X$ the complementary open immersion. Then the sequence*

$$M_c(Z) \xrightarrow{i_*} M_c(X) \xrightarrow{j^*} M_c(U)$$

defines an exact triangle in $\text{DA}_\tau(k)$.

d) Let

$$\begin{array}{ccc} Z' & \xrightarrow{i'} & X' \\ q \downarrow & & p \downarrow \\ Z & \xrightarrow{i} & X \end{array}$$

be an abstract alteration in the following sense: p is projective, i is a closed immersion, $Z' = p^{-1}(Z)$ and

$$r = p|_{X'-Z'} : X' - Y' \rightarrow X - Z$$

is an étale (finite) morphism of degree m . Then the map of cones

$$\text{cone}(M(Z') \rightarrow M(X')) \rightarrow \text{cone}(M(Z) \rightarrow M(X))$$

is split surjective either in $\text{DM}(k, \mathbf{Z}[1/m])$ or in $\text{DA}_{\text{ét}}(k, \mathbf{Z}[1/m])$ and is an isomorphism in $\text{DA}_\tau(k, \mathbf{Z}[1/m])$ if $m = 1$. In particular, if $m = 1$ we have an exact triangle

$$M(Z') \rightarrow M(X') \oplus M(Z) \rightarrow M(X) \xrightarrow{+1}.$$

e) If X is projective, $M_c(X) \simeq M(X)$.

f) If X is smooth, $M(X)$ is the classical motive of X . If we work in $\text{DM}(k)$ or $\text{DA}_{\text{ét}}(k, A)$, with 2 invertible in A , then

$$M_c(X) \simeq M(X)^*(-d)[-2d],$$

where $d = \dim X$.

g) Let $\mathcal{T} = \text{DA}_{\text{gm}, \text{ét}}^{\text{eff}}(k)$ or $\text{DM}_{\text{gm}}^{\text{eff}}(k, \mathbf{Z}[1/p])$, where p is the exponential characteristic of k . Then $M(X), M_c(X) \in \mathcal{T}$ for any X .

Proof. We shall prove the (a priori finer) dual functoriality to a) – f) for $M^*(X) := (p_X)_* \mathbf{Z}_X$ and $BM(X) := (p_X)! \mathbf{Z}_X$.

a) Let $f : X \rightarrow Y$. The isomorphism of functors

$$(p_X)_* \simeq (p_Y)_* \circ f_*$$

gives an isomorphism

$$M^*(X) \simeq (p_Y)_*(f_* \mathbf{Z}_X).$$

We get a morphism $f^* : M^*(Y) \rightarrow M^*(X)$ from the unit morphism $\mathbf{Z}_Y \rightarrow f_* f^* \mathbf{Z}_Y = f_* \mathbf{Z}_X$.

b) Same argument for the contravariance of BM , using the covariance of $f_!$ and the isomorphism $f_! \xrightarrow{\sim} f_*$ for f projective [2, Sch. 1.4.2]. Let now $f : X \rightarrow Y$ be an étale morphism. To define a covariant functoriality, we have to exhibit a canonical morphism $f_! \mathbf{Z}_X \rightarrow \mathbf{Z}_Y$ or equivalently $\mathbf{Z}_X \rightarrow f^! \mathbf{Z}_Y$. But since f is étale, we have $f^! = f^*$ (ibid.) and we take the identity morphism.

c) follows from applying $(p_X)_!$ to the exact triangle

$$j_! j^! \mathbf{Z}_X \rightarrow \mathbf{Z}_X \rightarrow i_* i^* \mathbf{Z}_X \xrightarrow{+1}$$

from [2, Lemma 1.4.6], noting that $i^* \mathbf{Z}_X = \mathbf{Z}_Z$, $j^! \mathbf{Z}_X = j^* \mathbf{Z}_X = \mathbf{Z}_U$ and $i_* = i_!$.

d) Let $j : X - Z \rightarrow X$, $j' : X' - Z' \rightarrow X'$ be the complementary open immersions to i, i' . Applying to $p_* \mathbf{Z}_{X'}$ the exact triangle of functors used in the proof of c), we get a commutative diagram of exact triangles

$$\begin{array}{ccccc} j_! j^! p_* \mathbf{Z}_{X'} & \longrightarrow & p_* \mathbf{Z}_{X'} & \longrightarrow & i_* i^* p_* \mathbf{Z}_{X'} \xrightarrow{+1} \\ \uparrow & & \uparrow & & \uparrow \\ j_! j^! \mathbf{Z}_X & \longrightarrow & \mathbf{Z}_X & \longrightarrow & i_* i^* \mathbf{Z}_X \xrightarrow{+1} . \end{array}$$

By projective base change [2, Sch. 1.4.2 5], this may be rewritten as

$$\begin{array}{ccccc} j_! r_* \mathbf{Z}_{X'-Z'} & \longrightarrow & p_* \mathbf{Z}_{X'} & \longrightarrow & i_* q_* \mathbf{Z}_{Z'} \xrightarrow{+1} \\ \uparrow & & \uparrow & & \uparrow \\ j_! \mathbf{Z}_{X-Z} & \longrightarrow & \mathbf{Z}_X & \longrightarrow & i_* \mathbf{Z}_Z \xrightarrow{+1} \end{array}$$

or

$$\begin{array}{ccccc} j_! r_* \mathbf{Z}_{X'-Z'} & \longrightarrow & p_* \mathbf{Z}_{X'} & \longrightarrow & p_* i'^* \mathbf{Z}_{Z'} \xrightarrow{+1} \\ \uparrow & & \uparrow & & \uparrow \\ j_! \mathbf{Z}_{X-Z} & \longrightarrow & \mathbf{Z}_X & \longrightarrow & i_* \mathbf{Z}_Z \xrightarrow{+1} . \end{array}$$

Consider the unit map $a : \mathbf{Z}_{X-Z} \rightarrow r_* \mathbf{Z}_{X'-Z'} = r_! \mathbf{Z}_{X'-Z'}$ and the counit map $b : r_! \mathbf{Z}_{X'-Z'} \rightarrow \mathbf{Z}_{X-Z}$. The composition ba is multiplication by m

in two cases: if we push the situation in $\mathrm{DM}(S)$ (because b is given by the transpose of the graph of r) or if $\tau = \acute{e}t$ (same proof as for [4, Lemma 2.3]). If $m = 1$, r is the identity in any situation. Applying $(p_X)_*$ to the above diagram, this concludes the proof of d).

e) follows from the isomorphism $(p_X)_* \xrightarrow{\sim} (p_X)!$ (when X is projective).

In f), the second statement follows from the isomorphism $(p_X)!\xrightarrow{\sim} (p_X)_\#(-d)[-2d]$, where $(p_X)_\#$ is the left adjoint of p_X^* , defined because p_X is smooth ([2, Sch. 1.4.2 3] plus [4, cor. 2.14]); recall that the isomorphism $(p_X)_\#\mathbf{Z}_X = M(X)$ is formal. The first statement follows from the isomorphism

$$\underline{\mathrm{Hom}}_k((p_X)_\#A, B) \xrightarrow{\sim} (p_X)_* \underline{\mathrm{Hom}}_X(A, p_X^*B)$$

from [2, Prop. 2.3.52], applied with $A = \mathbf{Z}_X, B = \mathbf{Z}$.

Finally, g) follows from the previous properties by dévissage using Gabber's refinement of de Jong's theorem [36], by reduction to the case of smooth projective varieties. \square

Remarks 6.7.3. 1) The condition “projective” may be relaxed to “proper” by arguments involving Chow's lemma⁷, cf. [SGA4, Exp. XVII, §7] and [15].

2) There are other useful formulas for $M(X)$ and $M^c(X)$:

$$M(X) = f_!f^!\mathbf{Z}, \quad M^c(X) = f_*f^!\mathbf{Z}$$

obtained by applying the Verdier duality of [2, Vol. I, p. 435, Th. 2.3.75].

3) It would remain to compare the motives $M(X)$ and $M_c(X)$ of Proposition 6.7.2 (or rather their images in $\mathrm{DM}_\tau(k)$) with those defined by Voevodsky. We leave this as a problem for the interested reader.

4) This provides definitions for motivic theories associated to a k -scheme of finite type X , in the style of [25, §9]:

Motivic cohomology:

$$H_\tau^i(X, \mathbf{Z}(n)) = \mathrm{DA}_\tau(k)(M(X), \mathbf{Z}(n)[i]).$$

Motivic cohomology with compact supports:

$$H_{\tau,c}^i(X, \mathbf{Z}(n)) = \mathrm{DA}_\tau(k)(M_c(X), \mathbf{Z}(n)[i]).$$

Motivic homology:

$$H_i^\tau(X, \mathbf{Z}(n)) = \mathrm{DA}_\tau(k)(\mathbf{Z}(n)[i], M(X)).$$

Borel-Moore motivic homology:

$$H_i^{\tau,c}(X, \mathbf{Z}(n)) = \mathrm{DA}_\tau(k)(\mathbf{Z}(n)[i], M_c(X)).$$

⁷plus a support property [15, 2.2.5], as pointed out by the referee.

It would remain to compare these (co)homology groups with those given by cdh cohomology as in [25, 78] or éh cohomology as in [29]: we leave this as another problem for the interested reader.

6.8. The l -adic realisation functor. The correct version of the target is the category defined by Ekedahl in [22]; technical details are spelled out in Ayoub’s article [4, §5]. He gets l -adic realisation functors:

Theorem 6.8.1. *Let S be a k -scheme of finite type and $l \neq 2, p$, where k is a finite field of characteristic p . Then there exist \otimes -triangulated functors*

$$(6.8.1) \quad R_l : \mathrm{DA}_{\acute{\mathrm{e}}\mathrm{t}}(S, \mathbf{Z}_l) \rightarrow \hat{D}_{\acute{\mathrm{e}}\mathrm{t}}(S, \mathbf{Z}_l)$$

sending the “motive” of a smooth S -scheme $f : X \rightarrow S$ to the dual of $Rf_*\mathbf{Z}_l$. We shall also write R_l for its restriction to $\mathrm{DA}_{\acute{\mathrm{e}}\mathrm{t}}(S)$.

These functors commute with the 6 operations of Grothendieck.

Proof. See [4, Prop. 5.8] for (6.8.1), *ibid.*, Th. 6.6 and Prop. 6.7 for the compatibility with the 6 operations and *ibid.*, Prop. 5.9 for the value of R_l on the motive of X . \square

6.9. The motivic version of the l -adic cycle class map. Assume that $S = \mathrm{Spec} k$ in Theorem 6.8.1. For a smooth k -scheme $f : X \rightarrow \mathrm{Spec} k$, R_l induces homomorphisms

$$(6.9.1) \quad \begin{aligned} H_{\acute{\mathrm{e}}\mathrm{t}}^i(X, \mathbf{Z}(n)) &= \mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(k)(M(X), \mathbf{Z}(n)[i]) \\ \xrightarrow{R_l} \hat{D}_{\acute{\mathrm{e}}\mathrm{t}}(k, \mathbf{Z}_l)(R_l M(X), R_l \mathbf{Z}(n)[i]) &= \hat{D}_{\acute{\mathrm{e}}\mathrm{t}}(k, \mathbf{Z}_l)(\mathbf{Z}_l, Rf_*\mathbf{Z}_l(n)[i]) \\ &= H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)). \end{aligned}$$

Here the equality on the first line follows from Theorem 5.5.2 a) and the full faithfulness of the functor $d_{\acute{\mathrm{e}}\mathrm{t}}$ in (5.5.1). The equality on the second line uses the fact that $Rf_*\mathbf{Z}_l$ is a strongly dualisable object: this is a formal consequence of the fact that R_l is a (strict) \otimes -functor and that $M(X)$ is strongly dualisable in $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(k)$ (Lemma 5.5.3).

One can check that (4.4.2) is induced by (6.9.1) when k is finite.

We get similar cycle class maps for the (co)homology theories considered in Remark 6.7.3 4).

However, as said in §4.4, these maps are insufficient to get a good reformulation of Friedlander’s conjecture: we shall have to wait until §9.6 to get the latter.

7. EFFECTIVITY AND SPECTRA

This section is very formal and would work in a wider generality; it will be used in the proof of Corollary 9.8.4.

7.1. Compacts, duals and Brown representability. In this subsection I review some now basic facts about “large” triangulated categories. First some terminology from [64]:

Definition 7.1.1. Let \mathcal{S} be a triangulated category.

- a) An object $X \in \mathcal{S}$ is *compact* if the functor $\mathcal{S}(X, -)$ commutes with representable direct sums.
- b) A subcategory of \mathcal{S} is *localising* if it is full, triangulated and stable under representable direct sums.
- c) A set of objects \mathcal{X} of \mathcal{S} is *dense in \mathcal{S}* if

$$\mathcal{X}^\perp := \{Y \in \mathcal{S} \mid \mathcal{S}(X, Y[i]) = 0 \forall X \in \mathcal{X} \forall i \in \mathbf{Z}\}$$

is reduced to 0.

- d) The category \mathcal{S} *satisfies TR5* if small direct sums are representable in \mathcal{S} , and define triangulated functors.
- e) The category \mathcal{S} is *compactly generated* if it satisfies TR5 and there exists a dense set of compact objects.

Clearly, the full subcategory $\mathcal{S}^c \subseteq \mathcal{S}$ consisting of compact objects is triangulated and *thick*, i.e., closed under direct summands.

The following is Neeman’s version of Brown’s representability theorem (a special case of [64, Prop. 8.4.2]):

Theorem 7.1.2. *Let \mathcal{S} be a compactly generated triangulated category. Let $H : \mathcal{S}^{op} \rightarrow \text{Ab}$ be a cohomological functor. Then H is representable if and only if it converts small direct sums into small products.*

Corollary 7.1.3. *Let \mathcal{S} be as in Theorem 7.1.2, and let f be a triangulated functor to another triangulated category \mathcal{T} . Then f has a right adjoint if and only if it commutes with small direct sums. This right adjoint is triangulated.*

To deduce Corollary 7.1.3 from Theorem 7.1.2 is left to the reader as an exercise.

We also note the following useful lemma of Ayoub:

Lemma 7.1.4 ([2, Lemma 2.1.28]). *Let $\mathcal{S}, \mathcal{T}, f$ be as in Corollary 7.1.3, and assume that f has a right adjoint g . Then g commutes with small direct sums if and only if f preserves compact objects.*

Finally, if \mathcal{S} is a tensor triangulated category [2, Def. 2.1.148], we have the notion of strongly dualisable object, see Lemma 5.5.3 and its proof.

Lemma 7.1.5. *Let \mathcal{S} be a tensor triangulated category, with unit object $\mathbf{1}$. Then strongly dualisable objects are compact if and only if $\mathbf{1}$ is compact.*

Proof. If $X, Y \in \mathcal{S}$ with X strongly dualisable, then

$$\mathcal{S}(X, Y) \simeq \mathcal{S}(\mathbf{1}, X^* \otimes Y)$$

where X^* is the dual of X . □

7.2. The hocolim construction. Let \mathcal{S} be a triangulated category in which countable direct sums are representable. In [11], Bökstedt and Neeman observed that one can perform the hocolim construction, up to non unique isomorphism: given a sequence

$$(7.2.1) \quad \dots \xrightarrow{f_{n-1}} X_n \xrightarrow{f_n} X_{n+1} \xrightarrow{f_{n+1}} \dots$$

in \mathcal{S} , define $\text{hocolim}(X_n, f_n)$ as a cone of the morphism

$$\bigoplus_{n \in \mathbf{Z}} X_n \xrightarrow{1-f_n} \bigoplus_{n \in \mathbf{Z}} X_n.$$

If Y is a compact object of \mathcal{S} , one easily constructs an isomorphism

$$(7.2.2) \quad \varinjlim \mathcal{S}(Y, X_n) \xrightarrow{\sim} \mathcal{S}(Y, \text{hocolim } X_n).$$

7.3. $\mathbf{Z}(1)$ -spectra. Let S be a Noetherian scheme and A be a commutative ring. Write $cd(S, A)$ for the étale cohomological dimension of S relatively to sheaves of A -modules. We need:

Proposition 7.3.1. *Suppose that $cd(S, A) < \infty$.*

a) *The category $\text{DA}_{\text{ét}}^{\text{eff}}(S, A)$ is compactly generated.*

b) *The “suspension spectrum” functor $i : \text{DA}_{\text{ét}}^{\text{eff}}(S, A) \rightarrow \text{DA}_{\text{ét}}(S, A)$ has a right adjoint $M \mapsto M^{\text{eff}}$.*

c) *The category $\text{DA}_{\text{ét}}^{\text{eff}}(S, A)$ is closed, with internal Hom given by*

$$\underline{\text{Hom}}^{\text{eff}}(M, N) = \underline{\text{Hom}}(i(M), i(N))^{\text{eff}}$$

where $\underline{\text{Hom}}$ is the internal Hom of $\text{DA}_{\text{ét}}(S, A)$.

Proof. a) If $cd(S, A) < \infty$, then $cd(X, A) < \infty$ for any X/S of finite type [SGA4, Exp. X]. For such X ,

$$\text{DA}_{\text{ét}}^{\text{eff}}(S, A)(M(X), C) \simeq H_{\text{ét}}^0(X, C)$$

for any $C \in \text{DA}_{\text{ét}}^{\text{eff}}(S, A)$ viewed as a complex of étale sheaves. To see that $M(X)$ is compact, one reduces to the cohomology of a single sheaf via hypercohomology spectral sequences, and then one uses that étale cohomology commutes with filtering direct limits of sheaves. Clearly, the set of $M(X)$ 's⁸ is dense in $\text{DA}_{\text{ét}}^{\text{eff}}(k, A)$. Axiom TR5 is also clear in terms of complexes of sheaves (this is where the unboundedness is important).

⁸more correctly, X running through a set of representatives of isomorphism classes of smooth S -schemes

b) follows from a) and Theorem 7.1.2, or directly from the construction of $\mathrm{DA}_{\acute{e}t}(S)$ as a category of spectra. To apply it, we use the fact that i commutes with small direct sums. c) follows formally. \square

In $\mathrm{DA}_{\acute{e}t}^{\mathrm{eff}}(S)$, we have the object $\mathbf{Z}(1)$ which may be defined as $\pi^*\mathbf{Z}(1)$ for $\pi : S \rightarrow \mathrm{Spec} k$, or alternately as a shifted direct summand of the motive of \mathbf{P}_S^1 . For an object $M \in \mathrm{DA}_{\acute{e}t}^{\mathrm{eff}}(S)$, we write as usual $M(1) := M \otimes \mathbf{Z}(1)$.

We may consider “ $\mathbf{Z}(1)$ -spectra” in $\mathrm{DA}_{\acute{e}t}^{\mathrm{eff}}(S)$: these are systems

$$(M_n, \varphi_n)_{n \in \mathbf{Z}}, \quad \varphi_n : M_n(1) \rightarrow M_{n+1}.$$

Of course, one could also consider $\mathbf{Z}(1)[i]$ -spectra for some $i \in \mathbf{Z}$: examples which may be relevant are $i = 1$ ($\mathbf{Z}(1)[1] \simeq \mathbb{G}_m[0]$) and $i = 2$ ($\mathbf{Z}(1)[2] \in \mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(k)$ is the image of the Lefschetz motive). This does not make any difference in practice.

Let $M \in \mathrm{DA}_{\acute{e}t}(S)$: we may associate to it the $\mathbf{Z}(1)$ -spectrum

$$M_n = (M(n))^{\mathrm{eff}}.$$

The transition morphisms

$$(7.3.1) \quad \varphi_n : M_n(1) \rightarrow M_{n+1}$$

are obtained by twisting once the counit morphisms $iM_n \rightarrow M(n)$ and using that i is a monoidal functor.

Lemma 7.3.2. *The morphisms (7.3.1) induce isomorphisms*

$$M_n \xrightarrow{\sim} \underline{\mathrm{Hom}}^{\mathrm{eff}}(\mathbf{Z}(1), M_{n+1}).$$

In other words, the spectrum (M_n) is an Ω -spectrum.

Proof. By an adjunction game, this follows from the fact that $\mathbf{Z}(1)$ is invertible in $\mathrm{DA}_{\acute{e}t}(S)$. \square

7.4. The case of a field. We now assume $S = \mathrm{Spec} k$, where k is a perfect field of finite étale cohomological dimension.

Let us apply the hocolim construction to $X_n = iM_n(-n) \in \mathrm{DA}_{\acute{e}t}(k)$: the transition morphisms $M_n(1) \rightarrow M_{n+1}$ induce morphisms $f_n : X_n \rightarrow X_{n+1}$. These morphisms are compatible with the untwisted counit morphisms $X_n \rightarrow M$, hence induce a morphism

$$(7.4.1) \quad \mathrm{hocolim} iM_n(-n) \rightarrow M.$$

Proposition 7.4.1. *Suppose that $S = \mathrm{Spec} k$. Then, for any $M \in \mathrm{DA}_{\acute{e}t}(k) = \mathrm{DM}_{\acute{e}t}(k)$, (7.4.1) is an isomorphism.*

(I don’t know if this is true over a general base; the proof below uses the full faithfulness of i , which relies on the cancellation theorem.)

Proof. Since $\mathrm{DM}_{\acute{e}t}(k)$ is generated by $\mathrm{DM}_{\mathrm{gm},\acute{e}t}(k)$, it is enough to check that, for any compact object $N \in \mathrm{DM}_{\mathrm{gm},\acute{e}t}(k)$, the morphism

$$\begin{aligned} \varinjlim_n \mathrm{DM}_{\acute{e}t}(k)(N, iM_n(-n)) &\xrightarrow{\sim} \mathrm{DM}_{\acute{e}t}(k)(N, \mathrm{hocolim}_n iM_n(-n)) \\ &\rightarrow \mathrm{DM}_{\acute{e}t}(k)(N, M) \end{aligned}$$

given by (7.2.2) and (7.4.1) is an isomorphism.

There exists $n_0 \gg 0$ such that $N(n) \in \mathrm{Im} i$ for $n \geq n_0$. For clarity, let us write $N(n) = iN_n$ for $n \geq n_0$. Since i is fully faithful, we get for $n \geq n_0$

$$\begin{aligned} \mathrm{DM}_{\acute{e}t}(k)(N, iM_n(-n)) &= \mathrm{DM}_{\acute{e}t}(k)(iN_n, iM_n) \\ &\xleftarrow{\sim} \mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k)(N_n, (M(n))^{\mathrm{eff}}) \simeq \mathrm{DM}_{\acute{e}t}(k)(iN_n, M(n)) \\ &= \mathrm{DM}_{\acute{e}t}(k)(N(n), M(n)) \xleftarrow{\sim} \mathrm{DM}_{\acute{e}t}(k)(N, M). \end{aligned}$$

We thus get an isomorphism

$$\begin{aligned} \mathrm{DM}_{\acute{e}t}(k)(N, M) &\xrightarrow{\sim} \varinjlim_{n \geq n_0} \mathrm{DM}_{\acute{e}t}(k)(N, iM_n(-n)) \\ &= \varinjlim_n \mathrm{DM}_{\acute{e}t}(k)(N, iM_n(-n)). \end{aligned}$$

One check that this isomorphism is inverse to the above map, which concludes the proof. \square

7.5. Ω -spectra and Σ -spectra. The following corollary is the *raison d'être* of this section:

Corollary 7.5.1. *For $M \in \mathrm{DM}_{\acute{e}t}(k)$, the following are equivalent:*

- (i) $M \in \mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k)$.
- (ii) For any $n \geq 0$, the natural morphism $M^{\mathrm{eff}}(n) \rightarrow (M(n))^{\mathrm{eff}}$ is an isomorphism. ($(M(n))^{\mathrm{eff}}$ is a “ Σ -spectrum”.)
- (iii) For any $n > 0$, $i(M(n))^{\mathrm{eff}}(-1) \in \mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k)$.

Proof. (i) \Rightarrow (ii) \Rightarrow (iii) is obvious; (ii) \Rightarrow (i) follows from Proposition 7.4.1 applied to M . Finally, assume (iii) true: thus we may write $(M(n))^{\mathrm{eff}} \simeq N(1)$ for some $N \in \mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(k)$. Using Lemma 7.3.2, we find

$$(M(n-1))^{\mathrm{eff}} \xrightarrow{\sim} \underline{\mathrm{Hom}}^{\mathrm{eff}}(\mathbf{Z}(1), (M(n))^{\mathrm{eff}}) \simeq \underline{\mathrm{Hom}}^{\mathrm{eff}}(\mathbf{Z}(1), N(1)) \xleftarrow{\sim} N$$

or $(M(n-1))^{\mathrm{eff}}(1) \xrightarrow{\sim} (M(n))^{\mathrm{eff}}$; by induction on n , we get (ii). \square

8. A PROOF OF THEOREM 3.2.1

From now on, \mathbf{F} is a finite field.

The aim of this section is to give the following effective version of Geisser’s theorem 3.2.1 [27, Th. 3.3]:

Theorem 8.0.2. *Let X be a smooth projective \mathbf{F} -variety of dimension d , and let $n \geq 0$. Assume:*

(1) Conjecture 3.1.1 holds for $X \times X$ in codimension d .

(2) The cohomological Tate conjecture holds for X in codimension n .

Then, for any $i \neq 2n$, $H^i(X, \mathbf{Q}(n)) = 0$ (“Parshin’s conjecture in weight n ”).

Proof. We shall use Theorem 5.2.1 as well as the functor

$$(8.0.1) \quad \Phi : \mathcal{M}_{\text{rat}}^{\text{eff}}(\mathbf{F}, \mathbf{Q}) \rightarrow \text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})$$

of [78, Prop. 2.1.4], where the left hand side is the category of effective Chow motives over \mathbf{F} . This shows that $\text{End}(h(X)) = CH^d(X \times X) \otimes \mathbf{Q}$ acts on the motivic cohomology of X , where $h(X)$ is the Chow motive of X . By assumption,

$$CH^d(X \times X) \otimes \mathbf{Q} \xrightarrow{\sim} A_{\text{num}}^d(X \times X, \mathbf{Q}).$$

By Jannsen [44], the right hand side is a semi-simple \mathbf{Q} -algebra. Thus we may decompose $h(X)$ as a direct sum

$$h(X) = \bigoplus_{\alpha \in A} S_{\alpha}$$

where the S_{α} are simple motives. Accordingly, we may decompose the motivic cohomology of X as

$$H^i(X, \mathbf{Q}(n)) = \bigoplus_{\alpha \in A} \text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})(\Phi(S_{\alpha}), \mathbf{Z}(n)[i]).$$

Let \mathbb{L} be the Lefschetz motive. If $S_{\alpha} = \mathbb{L}^n$, then

$$\begin{aligned} \text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})(\Phi(S_{\alpha}), \mathbf{Z}(n)[i]) &= \text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})(\mathbf{Z}(n)[2n], \mathbf{Z}(n)[i]) \\ &= 0 \text{ if } i \neq 2n. \end{aligned}$$

To conclude, it remains to see that $\text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})(\Phi(S_{\alpha}), \mathbf{Z}(n)[i]) = 0$ if $S_{\alpha} \not\cong \mathbb{L}^n$.

Let us first show that $\mathcal{M}_{\text{rat}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})(S_{\alpha}, \mathbb{L}^n) = 0$. This is obvious if $n \notin [0, d]$. If $n \in [0, d]$, then $A_{\text{num}}^n(X) \neq 0$ ⁹, which shows that \mathbb{L}^n appears as a direct summand of $h(X)$ and we conclude by semi-simplicity.

We now use that Frobenius defines an endomorphism of the identity functor of $\text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})$: this follows from the easy fact that it is natural with respect to the action of finite correspondences. In particular, for $M, N \in \text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})$, with Frobenius endomorphisms F_M, F_N , we have the equality $F_M^* = (F_N)_*$ on $\text{DM}_{\text{gm}}^{\text{eff}}(\mathbf{F}, \mathbf{Q})(M, N)$. Since the Frobenius endomorphism of \mathbb{L}^n equals q^n , we get:

⁹If L is an ample line bundle on X , then $\deg(L^d) > 0$ and *a fortiori* $0 \neq L^n \in A_{\text{num}}^n(X)$ for all $n \in [0, d]$...

Lemma 8.0.3. *Let F_α be the Frobenius endomorphism of S_α . Then F_α^* acts on $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(\mathbf{F}, \mathbf{Q})(\Phi(S_\alpha), \mathbf{Z}(n)[i])$ by multiplication by q^n . \square*

To conclude, it suffices to show that $F_\alpha \neq q^n$ in $D_\alpha = \mathrm{End}_{\mathcal{M}_{\mathrm{num}}^{\mathrm{eff}}(\mathbf{F}, \mathbf{Q})}(S_\alpha)$. (Then $F_\alpha^* - q^n$ will act both invertibly and by 0 on the D_α -vector space $\mathrm{DM}_{\mathrm{gm}}^{\mathrm{eff}}(\mathbf{F}, \mathbf{Q})(\Phi(S_\alpha), \mathbf{Z}(n)[i])$, and the latter will have to be 0.)

The cycle class map of Conjecture 3.3.1 splits into components:

$$\mathcal{M}_{\mathrm{rat}}^{\mathrm{eff}}(\mathbf{F}, \mathbf{Q})(S_\alpha, \mathbb{L}^n) \rightarrow H^{2n}(\bar{S}_\alpha, \mathbf{Q}_l(n))^G$$

with an obvious abuse of notation on the right. As we saw above, the left hand side is 0. By (2), the right hand side is therefore 0 as well. But this means that 1 is not an eigenvalue of the characteristic polynomial of F_α acting on $H^{2n}(\bar{S}_\alpha, \mathbf{Q}_l(n))$, which concludes the proof. \square

9. MOTIVIC REFORMULATION OF FRIEDLANDER'S CONJECTURE

9.1. An integral version? Getting a motivic version of Conjecture 3.3.2 with rational coefficients from (4.4.2) or (6.9.1) is easy: one considers the composition

(9.1.1)

$$\bar{c}_X^{n,i} : H_{\mathrm{ét}}^i(X, \mathbf{Z}(n)) \otimes \mathbf{Q}_l \rightarrow H_{\mathrm{cont}}^i(X, \mathbf{Q}_l(n)) \rightarrow H_{\mathrm{cont}}^i(\bar{X}, \mathbf{Q}_l(n))^G$$

and conjectures that it is an isomorphism. What about an integral version?

The first idea, to consider the composition

$$H_{\mathrm{ét}}^i(X, \mathbf{Z}(n)) \otimes \mathbf{Z}_l \rightarrow H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)) \rightarrow H_{\mathrm{cont}}^i(\bar{X}, \mathbf{Z}_l(n))^G$$

is tempting but looks unreasonable: a special case would imply a naïve integral version of the Tate conjecture, and it is well-known that the corresponding integral refinement of the Hodge conjecture has counterexamples, the first ones being due to Atiyah-Hirzebruch using Godeaux varieties.

Over \mathbf{F} , counterexamples to the surjectivity of the map

$$CH^n(X) \otimes \mathbf{Z}_l \rightarrow H_{\mathrm{cont}}^{2n}(\bar{X}, \mathbf{Z}_l(n))^G$$

also exist. In [61, Aside 1.4], Milne hints that the Atiyah-Hirzebruch examples also work here: this was fully justified in [17, Th. 2.1]. These counterexamples persist after any finite extension of \mathbf{F} . More recently, A. Pirutka has given an example of a smooth projective \mathbf{F} -variety X for which the map $CH^2(X) \rightarrow CH^2(\bar{X})^G$ is not surjective [65]: this provides a different kind of counterexample.

Of course, this rules out the even more naïve idea that (4.4.2) and (6.9.1) could be isomorphisms. In fact they cannot even be isomorphisms rationally, in view of the exact sequence (4.2.2). If X is smooth projective and $i = 2n + 1$, one gets a map

$$0 = H^{2n+1}(X, \mathbf{Z}(n)) \otimes \mathbf{Q}_l \rightarrow H_{\mathrm{cont}}^{2n+1}(X, \mathbf{Q}_l(n)) \xleftarrow{\sim} H_{\mathrm{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))^G.$$

Here we used (4.1.1) and Theorem 5.5.2 b) for the first vanishing, as well as (4.2.2) and Theorem 2.2.1, which gives the vanishing of $H^{2n+1}(\bar{X}, \mathbf{Q}_l(n))^G$. The equality

$$\dim_{\mathbf{Q}_l} H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))_G = \dim_{\mathbf{Q}_l} H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))^G$$

shows that $H_{\text{cont}}^{2n}(\bar{X}, \mathbf{Q}_l(n))_G$ is in general nonzero.

9.2. The complex \mathbf{Z}^c . The simplest example of the above is for $n = 0$ and $X = \text{Spec } \mathbf{F} = \text{Spec } \mathbf{F}_q$. Then

$$(9.2.1) \quad H_{\text{cont}}^j(\mathbf{F}_{q^r}, \mathbf{Z}_l(0)) = \begin{cases} \mathbf{Z}_l & \text{if } j = 0, 1 \\ 0 & \text{else} \end{cases}$$

while

$$H_{\text{ét}}^j(\mathbf{F}_{q^r}, \mathbf{Q}(0)) = \begin{cases} \mathbf{Q} & \text{if } j = 0 \\ 0 & \text{else.} \end{cases}$$

In (9.2.1), transition morphisms for passing from \mathbf{F}_q to \mathbf{F}_{q^r} are the identity on H^0 but are multiplication by r on H^1 . This gives a computation of the cohomology sheaves of $R\varprojlim(\mathbf{Z}_l)$, where $R\varprojlim : \hat{D}_{\text{ét}}(\mathbf{F}, \mathbf{Z}_l) \rightarrow D(\mathbf{F}_{\text{ét}}, \mathbf{Z}_l)$ is the total derived functor of \varprojlim :

$$\varprojlim^j \mathbf{Z}_l = \begin{cases} \mathbf{Z}_l & \text{if } j = 0 \\ \mathbf{Q}_l & \text{if } j = 1 \\ 0 & \text{else.} \end{cases}$$

In [46, §4], I computed the object $R\varprojlim \mathbf{Z}_l$ as follows: there exists an object $\mathbf{Z}^c \in D^b(\mathbf{F}_{\text{ét}}, \mathbf{Z})$, represented by an explicit complex of étale sheaves of length 1, such that

$$R\varprojlim \mathbf{Z}_l \simeq \mathbf{Z}^c \otimes \mathbf{Z}_l \text{ in } D(\mathbf{F}_{\text{ét}}, \mathbf{Z}_l)$$

for any $l \neq p$ (and even for $l = p$) [46, Def. 4.1 and Th. 4.6 b)].

Let us repeat the properties of \mathbf{Z}^c : its cohomology sheaves are \mathbf{Z} in degree 0 and \mathbf{Q} in degree 1; in particular

$$(9.2.2) \quad \mathbf{Z}/n[0] \xrightarrow{\sim} \mathbf{Z}^c \otimes^L \mathbf{Z}/n.$$

The extension class $\partial \in \text{Ext}_{\mathbf{F}_{\text{ét}}}^2(\mathbf{Q}, \mathbf{Z})$ corresponding to \mathbf{Z}^c is nonzero; it can be factored as follows in $D^b(\mathbf{F}_{\text{ét}}, \mathbf{Z})$:

$$(9.2.3) \quad \begin{array}{ccc} \mathbf{Q}[-1] & \xrightarrow{\partial} & \mathbf{Z}[1] \\ \pi \downarrow & & \uparrow \beta \\ \mathbf{Q}/\mathbf{Z}[-1] & \xrightarrow{e} & \mathbf{Q}/\mathbf{Z}. \end{array}$$

Here π is induced by the projection $\mathbf{Q} \rightarrow \mathbf{Q}/\mathbf{Z}$, β is the integral Bockstein and $\cdot e$ is cup-product with the canonical generator from (4.3.1).

In particular, $\partial \otimes \mathbf{Q} = 0$, yielding an isomorphism

$$(9.2.4) \quad \mathbf{Z}^c \otimes \mathbf{Q} \simeq \mathbf{Q}[0] \oplus \mathbf{Q}[-1].$$

Finally, we have

$$(9.2.5) \quad H_{\text{ét}}^j(\mathbf{F}, \mathbf{Z}^c) = \begin{cases} \mathbf{Z} & \text{if } j = 0, 1 \\ 0 & \text{else.} \end{cases}$$

In particular, (4.3.1) refines to a generator

$$(9.2.6) \quad e \in H_{\text{ét}}^1(\mathbf{F}, \mathbf{Z}^c).$$

See Appendix A for more details on \mathbf{Z}^c .

I used \mathbf{Z}^c in [46] and [47] to define a *modified cycle class map*:

$$(9.2.7) \quad H_{\text{ét}}^j(X, \mathbf{Z}(n) \overset{L}{\otimes} \mathbf{Z}^c) \otimes \mathbf{Z}_l \xrightarrow{\text{cl}_X^{n,j}} H_{\text{cont}}^j(X, \mathbf{Z}_l(n))$$

which serves for an integral version of Friedlander's conjecture. In the sequel, we shall recover this map more conceptually.

9.3. A lemma of Ayoub. Let \mathcal{M}, \mathcal{N} be two unital symmetric monoidal categories, and let $f : \mathcal{M} \rightarrow \mathcal{N}$ be a (strong, symmetric, unital) monoidal functor. Assume that f has a right adjoint g . Then the unit map

$$M \rightarrow gf(M)$$

enriches into a bi-natural transformation

$$(9.3.1) \quad M \otimes gf(M') \rightarrow gf(M \otimes M'), \quad M, M' \in \mathcal{M}$$

which can be described as follows: by adjunction, such a morphism corresponds to a morphism

$$f(M) \otimes fgf(M') \simeq f(M \otimes gf(M')) \rightarrow f(M \otimes M') \simeq f(M) \otimes f(M').$$

This morphism is simply induced by the counit map $fgf(M') \rightarrow f(M')$.

In particular, we may take $M' = \mathbf{1}$ and get a natural transformation

$$(9.3.2) \quad M \otimes gf(\mathbf{1}) \rightarrow gf(M).$$

We are interested in a sufficient condition for (9.3.2) to be an isomorphism. For this, note that we have another bi-natural transformation

$$(9.3.3) \quad M \otimes g(N) \rightarrow g(f(M) \otimes N), \quad M \in \mathcal{M}, N \in \mathcal{N}$$

which corresponds by adjunction to the map

$$f(M) \otimes fg(N) \simeq f(M \otimes g(N)) \rightarrow f(M) \otimes N$$

given by the counit at N . To recover (9.3.1) from (9.3.3), just take $N = f(M')$. The following “projection formula” is due to Joseph Ayoub [3, Lemme 2.8]:

Lemma 9.3.1. *(9.3.3) is an isomorphism if M is strongly dualisable.*

Proof. It is so simple that we don’t resist in reproducing it:

Given an object X of \mathcal{M} or \mathcal{N} , write t_X for the functor $X \otimes -$. Let C be a strong dual of M , i.e., there exist morphisms $C \otimes M \rightarrow \mathbf{1}$ and $\mathbf{1} \rightarrow M \otimes C$ which make t_C a left adjoint of t_M . As f is monoidal, symmetric and unital, one deduces that $f(C)$ is a strong dual of $f(M)$, hence that $t_{f(C)}$ is left adjoint to $t_{f(M)}$. On the other hand, there is an obvious isomorphism $t_{f(C)} \circ f \simeq f \circ t_C$. Using the adjunctions $(f; g)$, $(t_C; t_M)$ and $(t_{f(C)}; t_{f(M)})$, one deduces an isomorphism of functors $t_M \circ g \simeq g \circ t_{f(M)}$. It remains to see that this isomorphism coincides with (9.3.3). This verification is omitted. \square

Corollary 9.3.2. *If M is strongly dualisable, (9.3.2) is an isomorphism.* \square

Remark 9.3.3. Of course Lemma 9.3.1 and Corollary 9.3.2 can be extended: suppose that \mathcal{M}, \mathcal{N} are triangulated categories and f (hence g) is a triangulated functor. Suppose also that g commutes with small direct sums (compare Lemma 7.1.4). Then, given $N \in \mathcal{N}$, the full subcategory of \mathcal{M} consisting of those M for which (9.3.3) is an isomorphism is localising. In particular, if strongly dualisable objects are dense in \mathcal{M} , then (9.3.3) holds universally as soon as g commutes with small direct sums.

9.4. A stable theorem. We shall need:

Lemma 9.4.1. *The functor R_l of Theorem 6.8.1 has a right adjoint Ω_l .*

Proof. This is true for both versions: with coefficients \mathbf{Z} or \mathbf{Z}_l . Once again this follows from Corollary 7.1.3 (note that R_l commutes with small direct sums). \square

Remark 9.4.2. The unit object \mathbf{Z}_l of $\hat{D}_{\text{ét}}(S, \mathbf{Z}_l)$ is not compact. This implies that Ω_l does not commute with small direct sums (see lemma 7.1.4).

We now apply Corollary 9.3.2 and Remark 9.3.3 to the adjunction (R_l, Ω_l) of Lemma 9.4.1; we get:

Theorem 9.4.3. *For any strongly dualisable object M of $\text{DA}_{\text{ét}}(S)$, the map*

$$M \otimes \Omega_l(\mathbf{Z}_l) \rightarrow \Omega_l R_l(M)$$

is an isomorphism.

If $S = \text{Spec } \mathbf{F}$ and $l \neq 2$, this applies to all objects of $\text{DM}_{\text{gm}, \text{ét}}(\mathbf{F})$.

Corollary 9.4.4. *Let $\Gamma = \Omega_l(\mathbf{Z}_l)$. Let $M, N \in \mathrm{DA}_{\acute{e}t}(S)$ and suppose that N is strongly dualisable. Then there is a natural isomorphism*

$$\mathrm{DA}_{\acute{e}t}(S)(M, N \otimes \Gamma) \xrightarrow{\sim} \hat{D}_{\acute{e}t}(S, \mathbf{Z}_l)(R_l(M), R_l(N)).$$

In particular, assume that $S = \mathrm{Spec} \mathbf{F}$ and $l \neq 2$; taking $M = M(X)$ for a smooth \mathbf{F} -scheme X and $N = \mathbf{Z}(n)[i]$, we get an isomorphism

$$\mathrm{DM}_{\acute{e}t}(\mathbf{F})(M(X), \Gamma(n)[i]) \xrightarrow{\sim} H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)).$$

Lemma 9.4.5. *Let $f : S \rightarrow \mathrm{Spec} \mathbf{F}$ be a smooth \mathbf{F} -scheme of finite type. Then there is a canonical isomorphism $\Omega_l^S(\mathbf{Z}_l) \simeq f^* \Omega_l^{\mathrm{Spec} \mathbf{F}} \mathbf{Z}_l$.*

Proof. Since f is smooth, f^* has a left adjoint f_{\sharp} which commutes with R_l [4]. This implies formally that f^* commutes with Ω_l . Since f^* also commutes with R_l (ibid.), the conclusion follows. \square

9.5. Towards the integral Friedlander conjecture. Theorem 9.4.3 and its corollary look fantastic, but they are in some sense a *trompe-l'œil*: all the hard l -adic information is concentrated in the object $\Gamma = \Omega_l(\mathbf{Z}_l)$. To explain this, let us evaluate this object against the motive of some smooth variety detwisted: for simplicity we assume $S = \mathrm{Spec} \mathbf{F}$. Thus, if $M = M(X)(-n)[-i]$ with X a smooth \mathbf{F} -scheme and $n, i \in \mathbf{Z}$, we have

$$\begin{aligned} \mathrm{DM}_{\acute{e}t}(\mathbf{F})(M(X)(-n)[-i], \Omega_l(\mathbf{Z}_l)) &= \hat{D}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(R_l(M(X)(-n)[-i]), \mathbf{Z}_l) \\ &= \hat{D}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(\mathbf{Z}_l, Rf_* \mathbf{Z}_l(n)[i]) = H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n)) \end{aligned}$$

where $f : X \rightarrow \mathrm{Spec} \mathbf{F}$ is the structural morphism.

Nevertheless, Corollary 9.4.4 gives the flavour of the integral Friedlander conjecture, to come. In order to get something interesting, we need to go to the *effective* subcategory $\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(\mathbf{F})$.

9.6. The modified motivic cycle map. From Lemma 9.4.1 and Proposition 7.3.1, we deduce that the composite functor

$$\mathrm{DA}_{\acute{e}t}^{\mathrm{eff}}(S) \xrightarrow{i} \mathrm{DA}_{\acute{e}t}(S) \xrightarrow{R_l} \hat{D}_{\acute{e}t}(S, \mathbf{Z}_l)$$

has the right adjoint

$$\Omega_l^{\mathrm{eff}} : \hat{D}_{\acute{e}t}(S, \mathbf{Z}_l) \xrightarrow{\Omega_l} \mathrm{DA}_{\acute{e}t}(S) \xrightarrow{-\mathrm{eff}} \mathrm{DA}_{\acute{e}t}^{\mathrm{eff}}(S).$$

Using (9.3.2), we get a morphism for any $M \in \mathrm{DA}_{\acute{e}t}^{\mathrm{eff}}(S)$:

$$(9.6.1) \quad M \otimes \Gamma^{\mathrm{eff}} \rightarrow \Omega_l^{\mathrm{eff}} R_l(M)$$

where $\Gamma = \Omega_l(\mathbf{Z}_l)$, as in Corollary 9.4.4. The most important case is $M = \mathbf{Z}(n)$ for $n \geq 0$, which we now record:

$$(9.6.2) \quad \Gamma^{\mathrm{eff}}(n) \rightarrow \Omega_l^{\mathrm{eff}} \mathbf{Z}_l(n).$$

The right hand side of (9.6.2) is denoted by $\mathbf{Z}_l(n)^c$ in [46] and [47]. It computes continuous étale cohomology with coefficients in $\mathbf{Z}_l(n)$.

Definition 9.6.1. We call (9.6.2) the *modified motivic cycle map*.

9.7. The object Γ^{eff} . From now on, we assume that $S = \text{Spec } \mathbf{F}$ and $l \neq 2$.¹⁰ Of course, Corollary 9.3.2 tells us that (9.6.1) is an isomorphism when M is strongly dualisable in $\text{DM}_{\text{ét}}^{\text{eff}}(\mathbf{F})$, but there are very few such objects: we only have the objects of $d_{\leq 0} \text{DM}_{\text{gm,ét}}^{\text{eff}}(\mathbf{F})$ (see below). The motive $\mathbf{Z}(n)$, for $n > 0$, is certainly not of this kind. Nevertheless we are in a better situation than in §9.5, in that it is possible to compute Γ^{eff} :

Proposition 9.7.1. *We have a canonical isomorphism*

$$(9.7.1) \quad \mathbf{Z}^c \otimes \mathbf{Z}_l \xrightarrow{\sim} \Gamma^{\text{eff}}$$

where \mathbf{Z}^c is the object of $D^b(\mathbf{F}_{\text{ét}}, \mathbf{Z})$ recalled in §9.2. In particular, $\Gamma^{\text{eff}} \otimes \mathbf{Q} \simeq \mathbf{Q}_l[0] \oplus \mathbf{Q}_l[-1]$.

Proof. This corresponds to [46, Th. 6.3 and Cor. 6.4]. Let me repeat the proof, which is a simple application of Theorem 2.2.1. We first construct the map (9.7.1). For $n \geq 0$, let $d_{\leq n} \text{DM}_{\text{ét}}^{\text{eff}}(\mathbf{F})$ be the localising subcategory of $\text{DM}_{\text{ét}}^{\text{eff}}(\mathbf{F})$ generated by motives of varieties of dimension $\leq n$ [78, §3.4]. The inclusion functor $i_n : d_{\leq n} \text{DM}_{\text{ét}}^{\text{eff}}(\mathbf{F}) \rightarrow \text{DM}_{\text{ét}}^{\text{eff}}(\mathbf{F})$ has a right adjoint ρ_n . Suppose $n = 0$: by §9.2 we have

$$\rho_0 \Gamma^{\text{eff}} \simeq \mathbf{Z}^c \otimes \mathbf{Z}_l$$

and we get the map by adjunction.

It is clear that

$$\Gamma^{\text{eff}} \otimes \mathbf{Z}/l \simeq \Omega_l^{\text{eff}}(\mathbf{Z}/l) = \mathbf{Z}/l.$$

Combining this with (9.2.2), we find that the cone of (9.7.1) is uniquely divisible. Hence it suffices to show that (9.7.1) $\otimes \mathbf{Q}$ is an isomorphism.

Let X be a smooth projective \mathbf{F} -variety, and let $X \rightarrow \pi_0(X)$ be the Stein factorisation of the structural morphism $X \rightarrow \text{Spec } \mathbf{F}$: $\pi_0(X)$ is the “scheme of constants” of X . For $i \in \mathbf{Z}$, we have a commutative diagram

$$\begin{array}{ccc} H_{\text{ét}}^i(\pi_0(X), \mathbf{Z}^c \otimes \mathbf{Q}_l) & \xrightarrow{a} & H_{\text{ét}}^i(X, \mathbf{Z}^c \otimes \mathbf{Q}_l) \\ c \downarrow & & d \downarrow \\ H_{\text{cont}}^i(\pi_0(X), \mathbf{Q}_l) & \xrightarrow{b} & H_{\text{cont}}^i(X, \mathbf{Q}_l) \end{array}$$

where the vertical maps are induced by (9.7.1) $\otimes \mathbf{Q}$.

In this diagram, c is an isomorphism by the result of §9.2. The map a is an isomorphism for trivial reasons. The Riemann hypothesis over finite fields and (4.2.2) imply that b is an isomorphism. Hence so is d .

¹⁰In order to work with $\text{DM}_{\text{ét}}(\mathbf{F})$ rather than $\text{DA}_{\text{ét}}(\mathbf{F})$.

Thus $(9.7.1) \otimes \mathbf{Q}$ induces an isomorphism when evaluated against the motive $M(X)[-i]$ for any smooth projective X and any $i \in \mathbf{Z}$. Since these objects generate $\mathrm{DM}_{\acute{e}t}^{\mathrm{eff}}(\mathbf{F}, \mathbf{Q})$, $(9.7.1) \otimes \mathbf{Q}$ is an isomorphism by Yoneda's lemma. \square

The terminology of Definition 9.6.1 is now justified by

Corollary 9.7.2. *Evaluating (9.6.2) against the motive of a smooth variety X yields a map*

$$H_{\acute{e}t}^j(X, \mathbf{Z}(n) \otimes^L \mathbf{Z}^c) \otimes \mathbf{Z}_l \xrightarrow{\mathrm{cl}_X^{n,j}} H_{\mathrm{cont}}^j(X, \mathbf{Z}_l(n))$$

of the form (9.2.7). \square

Remark 9.7.3. Let us record the other computations of [46, Cor. 6.10 and 6.12], which also rely on the Riemann hypothesis:

- (1) For $n \in \mathbf{Z}$, the cohomology sheaves of $\Omega_l^{\mathrm{eff}} \mathbf{Z}_l(n) \otimes \mathbf{Q}$ are concentrated in $[n, 2n + 1]$; in particular, $\Omega_l^{\mathrm{eff}} \mathbf{Z}_l(n) \otimes \mathbf{Q} = 0$ for $n < 0$.
- (2) For $0 \leq m < n$, $\rho_m(\Omega_l^{\mathrm{eff}} \mathbf{Z}_l(n) \otimes \mathbf{Q}) = 0$ (see proof of Proposition 9.7.1 for the definition of ρ_m).

9.8. The motivic Friedlander conjecture. We start with:

Lemma 9.8.1. *For any $M \in \mathrm{DM}_{\acute{e}t}(\mathbf{F})$ and $\nu \geq 1$, $(9.6.1) \otimes \mathbf{Z}/l^\nu$ is an isomorphism (hence so is $(9.6.2) \otimes \mathbf{Z}/l^\nu$).*

Proof. Theorem 5.5.1 b) implies that $R_l^{\mathrm{eff}} = R_l \circ i$ and Ω_l^{eff} become quasi-inverse equivalence of categories when we pass to \mathbf{Z}/l^ν coefficients. \square

Proposition 9.8.2. *Let X be a smooth \mathbf{F} -variety and $n \in \mathbf{Z}$. Then the following are equivalent:*

- (i) *The map $\mathrm{cl}_X^{n,j}$ of Corollary 9.7.2 is an isomorphism for any $j \in \mathbf{Z}$.*
- (ii) *Same as (i), replacing $\mathrm{cl}_X^{n,j}$ by $\mathrm{cl}_X^{n,j} \otimes \mathbf{Q}$.*
- (iii) *The map $\bar{\mathrm{cl}}_X^{n,j}$ of (9.1.1) is bijective for any $j \in \mathbf{Z}$ (Friedlander's conjecture), and the composition as in Remark 2.2.3*

$$\rho_X^{n,j} : H_{\mathrm{cont}}^j(\bar{X}, \mathbf{Q}_l(n))^G \hookrightarrow H_{\mathrm{cont}}^j(\bar{X}, \mathbf{Q}_l(n)) \twoheadrightarrow H_{\mathrm{cont}}^j(\bar{X}, \mathbf{Q}_l(n))_G$$

is bijective for any $j \in \mathbf{Z}$.

Proof. This is an elaboration of the proof of [47, Prop. 3.9]:

The equivalence (i) \iff (ii) follows from Lemma 9.8.1. We now prove (ii) \iff (iii).

Using (4.2.2), (9.2.4) and the commutative diagram (4.3.2) applied to $C = Rf_* \mathbf{Q}_l(n)$ for $f : X \rightarrow \mathrm{Spec} \mathbf{F}$ the structural morphism, one checks

that $\mathrm{cl}_X^{n,j} \otimes \mathbf{Q}$ fits in a diagram of short exact sequences

$$\begin{array}{ccccccc} 0 \rightarrow & H^{j-1}(X, \mathbf{Q}(n))_{\mathbf{Q}_l} & \rightarrow & H^j(X, \mathbf{Q}^c(n))_{\mathbf{Q}_l} & \rightarrow & H^j(X, \mathbf{Q}(n))_{\mathbf{Q}_l} & \rightarrow 0 \\ & f^{j-1} \downarrow & & \mathrm{cl}_X^{n,j} \otimes \mathbf{Q} \downarrow & & \bar{\mathrm{cl}}_X^{n,j} \downarrow & \\ 0 \rightarrow & H_{\mathrm{cont}}^{j-1}(\bar{X}, \mathbf{Q}_l(n))_G & \rightarrow & H_{\mathrm{cont}}^j(X, \mathbf{Q}_l(n)) & \rightarrow & H_{\mathrm{cont}}^j(\bar{X}, \mathbf{Q}_l(n))_G & \rightarrow 0 \end{array}$$

where $-\mathbf{Q}_l := - \otimes_{\mathbf{Q}} \mathbf{Q}_l$, $\mathbf{Q}^c(n) := \mathbf{Z}(n) \otimes^L \mathbf{Z}^c \otimes \mathbf{Q}$. Moreover, $f^j = \rho_X^{n,j} \circ \bar{\mathrm{cl}}_X^{n,j}$.

If $\mathrm{cl}_X^{n,j} \otimes \mathbf{Q}$ is bijective, we get the surjectivity of $\bar{\mathrm{cl}}_X^{n,j}$ and the injectivity of f^{j-1} . So if $\mathrm{cl}_X^{n,j+1} \otimes \mathbf{Q}$ is also bijective, then $\bar{\mathrm{cl}}_X^{n,j}$ and $\rho_X^{n,j}$ are bijective.

Conversely, if $\bar{\mathrm{cl}}_X^{n,j}$, $\rho_X^{n,j-1}$ and $\bar{\mathrm{cl}}_X^{n,j-1}$ are bijective, then $\mathrm{cl}_X^{n,j} \otimes \mathbf{Q}$ is bijective. \square

Theorem 9.8.3. *For X smooth projective and $n \geq 0$, consider the following conditions:*

$A(X, n)$: *Conjectures 2.1.1 and 3.1.1 hold for (X, n) .*

$B(X, n)$: *$\mathrm{cl}_X^{n,j}$ is an isomorphism for all $j \in \mathbf{Z}$.*

Let $d = \dim X$. Then

$$B(X, n) + B(X, d - n) \Rightarrow A(X, n) + A(X, d - n)$$

and

$$A(X \times X, d) + A(X, n) \Rightarrow B(X, n).$$

In particular, the validity of $B(X^m, n)$ for all (m, n) is independent of l .

Proof. This is a DM version of the equivalence between conjectures 3.1 and 3.2 of [47]. Let me recall the proof:

1) By Proposition 9.8.2 and Theorem 2.2.1, $B(X, n)$ is equivalent to the following three conditions put together:

- (i) $H^i(X, \mathbf{Q}(n)) = 0$ for $i \neq 2n$.
- (ii) Conjecture 3.3.1 holds for (X, n) .
- (iii) Condition S^n of Remark 2.2.3 holds for X .

2) By Theorem 2.2.2 and Remark 2.2.3, (ii) + (iii) for (X, n) and $(X, d - n) \iff A(X, n) + A(X, d - n)$, where $d = \dim X$. In particular we get the first implication in Theorem 9.8.3.

3) It remains to see that $A(X \times X, d) + A(X, n) \Rightarrow$ (i): this follows from Theorem 8.0.2. \square

Corollary 9.8.4. *a) The following are equivalent:*

- (i) *Conjectures 2.1.1 and 3.1.1.*

- (ii) $\mathrm{cl}_X^{n,j}$ is an isomorphism for all smooth projective X and any $n \geq 0$, $j \in \mathbf{Z}$.
- (iii) $\mathrm{cl}_X^{n,j}$ is an isomorphism for all smooth X and any $n \geq 0$, $j \in \mathbf{Z}$.
- (iv) (9.6.2) is an isomorphism for any $n \geq 0$.
- (v) For any $n > 0$, $(\Omega_l^{\mathrm{eff}} \mathbf{Z}_l(n))(-1) \in \mathrm{DM}_{\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})$.
- (vi) $\Gamma \in \mathrm{DM}_{\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})$; equivalently, $\Gamma^{\mathrm{eff}} \xrightarrow{\sim} \Gamma$.
- (vii) Same as (vi), replacing $\mathrm{DM}_{\mathrm{ét}}(\mathbf{F})$ by $\mathrm{DM}_{\mathrm{ét}}(\mathbf{F}, \mathbf{Z}_l)$.

b) If this holds, then (9.6.1) is an isomorphism for any $M \in \mathrm{DM}_{\mathrm{gm},\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})$. In particular, for such M and for any $N \in \mathrm{DM}_{\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})$, there is an isomorphism

$$\mathrm{DM}_{\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})(N, M \otimes \mathbf{Z}^c) \otimes \mathbf{Z}_l \xrightarrow{\sim} \hat{D}_{\mathrm{ét}}(\mathbf{F}, \mathbf{Z}_l)(R_l(N), R_l(M)).$$

Proof. a) (i) \iff (ii): this follows from Theorem 9.8.3.

(ii) \iff (iv): this is formal, since $\mathrm{DM}_{\mathrm{gm},\mathrm{ét}}^{\mathrm{eff}}(\mathbf{F})$ is generated by the $M(X)$ for X smooth projective.¹¹

(iv) \Rightarrow (iii) \Rightarrow (ii) is trivial.

(iv) \iff (v) \iff (vi): this follows from Corollary 7.5.1.

(vi) \iff (vii) is formal. (Note that the right adjoint to $\mathrm{DM}_{\mathrm{ét}}(\mathbf{F}) \rightarrow \mathrm{DM}_{\mathrm{ét}}(\mathbf{F}, \mathbf{Z}_l)$ is just given by restriction of scalars.)

b) Applying the functor $-\mathrm{eff}$ to the isomorphism of Theorem 9.4.3, we get an isomorphism

$$(M \otimes \Gamma)^{\mathrm{eff}} \xrightarrow{\sim} \Omega_l^{\mathrm{eff}} R_l(M).$$

If (vi) holds, the obvious isomorphism $(M \otimes \Gamma^{\mathrm{eff}})^{\mathrm{eff}} \xrightarrow{\sim} M \otimes \Gamma^{\mathrm{eff}}$ yields an isomorphism $M \otimes \Gamma^{\mathrm{eff}} \xrightarrow{\sim} \Omega_l^{\mathrm{eff}} R_l(M)$, that one checks is (9.6.1). \square

The next corollary follows from Lemma 9.4.5:

Corollary 9.8.5. *Suppose that the equivalent conditions of Corollary 9.8.4 hold, and let S be a smooth \mathbf{F} -scheme of finite type. Then there is a canonical isomorphism*

$$M \otimes \mathbf{Z}_l \otimes \mathbf{Z}^c \xrightarrow{\sim} \Omega_l R_l(M)$$

for any strongly dualisable $M \in \mathrm{DA}_{\mathrm{ét}}(S)$. \square

10. DUALITY AND FINITE GENERATION

Motivated by the last isomorphism of Corollary 9.8.4, we define here a new category $\mathrm{DA}_W(\mathbf{F})$ out of $\mathrm{DA}_{\mathrm{ét}}(\mathbf{F})$ by using the product $\mu : \mathbf{Z}^c \otimes^L \mathbf{Z}^c \rightarrow \mathbf{Z}^c$ from Appendix A. We shall then prove that the equivalent statements

¹¹One may avoid the use of Gabber's theorem here by using the equivalence (i) \iff (ii) of Proposition 9.8.2 and working in $\mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$: then only de Jong's theorem is needed.

of corollary 9.8.4 are also equivalent to the finite generation (as $\mathbf{Z}[1/p]$ -modules) of the Hom groups for the constructible objects of $\mathrm{DA}_W(\mathbf{F})$.

10.1. l -adic duality. Let $\hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)$ be the full subcategory of $\hat{D}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)$ consisting of perfect complexes: it is equivalent to the derived category of bounded complexes of finitely generated $\mathbf{Z}_l[[G]]$ -modules. A form of Verdier duality is the following (cf. [68, Ch. I, Ann. 2]):

Proposition 10.1.1. *The “global sections” functor $R\Gamma : \hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l) \rightarrow D^b(\mathbf{Z}_l)$ has a right adjoint Π , and $\Pi(X) \simeq X[1]$ where $X \in D^b(\mathbf{Z}_l)$ is viewed as a complex of constant sheaves in $\hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)$. \square*

Corollary 10.1.2. *Let $C \in \hat{D}_c^b := \hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)$. Then*

a) *The composition*

$$\mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, C) \times \mathrm{Hom}_{\hat{D}_c^b}(C, \mathbf{Z}_l[1]) \rightarrow \mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, \mathbf{Z}_l[1]) = \mathbf{Z}_l$$

is a perfect duality modulo torsion.

b) *The compositions*

$$\mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, C) \times \mathrm{Hom}_{\hat{D}_c^b}(C, \mathbf{Z}/l^\nu[1]) \rightarrow \mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, \mathbf{Z}/l^\nu[1]) = \mathbf{Z}/l^\nu$$

induce a perfect duality of finite groups

$$\mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, C)_{\mathrm{tors}} \times \mathrm{Hom}_{\hat{D}_c^b}(C, \mathbf{Z}_l[2])_{\mathrm{tors}} \rightarrow \mathbf{Q}_l/\mathbf{Z}_l.$$

Proof. Proposition 10.1.1 yields an isomorphism

$$R\Gamma \underline{\mathrm{Hom}}_{\mathbf{F}}(C, \mathbf{Z}_l[1]) \xrightarrow{\sim} \underline{\mathrm{Hom}}(R\Gamma(C), \mathbf{Z}_l)$$

where $\underline{\mathrm{Hom}}_{\mathbf{F}}$ denotes the internal Hom of $\hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)$. Taking cohomology, we get

$$\mathrm{Hom}_{\hat{D}_c^b}(C, \mathbf{Z}_l[q+1]) \xrightarrow{\sim} H^q(\underline{\mathrm{Hom}}(R\Gamma(C), \mathbf{Z}_l))$$

and we conclude with the well-known exact sequences

$$\begin{aligned} 0 \rightarrow \mathrm{Ext}_{\mathbf{Z}_l}^1(\mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, C[1-q]), \mathbf{Z}_l) &\rightarrow H^q(\underline{\mathrm{Hom}}(R\Gamma(C), \mathbf{Z}_l)) \\ &\rightarrow \mathrm{Hom}_{\mathbf{Z}_l}(\mathrm{Hom}_{\hat{D}_c^b}(\mathbf{Z}_l, C[-q]), \mathbf{Z}_l) \rightarrow 0 \end{aligned}$$

for $q = 0, -1$. (We leave it to the reader to verify that the corresponding isomorphisms are really induced by the pairings in a) and b). \square)

10.2. An abstract construction. Let \mathcal{M} be a unital, symmetric monoidal category and let Γ be a commutative monoid in \mathcal{M} [56, Ch. VII, §3]. We shall construct a new (unital, symmetric) monoidal category $\mathcal{M}(\Gamma)$ and a pair of adjoint functors

$$\mathcal{M} \underset{\gamma_*}{\overset{\gamma^*}{\rightleftarrows}} \mathcal{M}(\Gamma)$$

with γ^* symmetric monoidal.

For this, we observe that the endofunctor $M \mapsto M \otimes \Gamma$ has the structure of a monad [56, Ch. VI, §1]; we define $\mathcal{M}(\Gamma)$ as the Kleisli category associated to this monad (ibid., §5, Th. 1). Recall its definition:

- The objects of $\mathcal{M}(\Gamma)$ are the objects of \mathcal{M} .
- Given two objects M, N ,

$$\mathcal{M}(\Gamma)(M, N) = \mathcal{M}(M, N \otimes \Gamma).$$

- Composition is defined by using the multiplication of Γ .

This construction comes with an adjunction (γ^*, γ_*) as required. Let us describe these two functors:

- γ^* is the identity on objects and acts on morphisms via the unit of Γ .
- γ_* sends an object M to $M \otimes \Gamma$ and acts on morphisms via the multiplication of Γ .

The only thing which is not in Mac Lane's book is the symmetric monoidal structure on $\mathcal{M}(\Gamma)$ and γ^* : this will be left to the reader.

10.3. An artificial category. Recall from Appendix A that the object \mathbf{Z}^c of $\mathrm{DM}_{\acute{e}t}(\mathbf{F})$ is provided with a multiplication $\mu : \mathbf{Z}^c \otimes^L \mathbf{Z}^c \rightarrow \mathbf{Z}^c$; together with the map $\eta : \mathbf{Z}[0] \rightarrow \mathbf{Z}^c$, one checks that \mathbf{Z}^c has the structure of a commutative monoid. This will be better justified in §11.

Definition 10.3.1. Let $f : S \rightarrow \mathrm{Spec} \mathbf{F}$ be a smooth scheme of finite type, and let A be a commutative ring.

a) We write $\mathrm{DA}_W(S, A)$ for the category $\mathrm{DA}_{\acute{e}t}(S, A)(f^*\mathbf{Z}^c)$, with the notation of the previous subsection.

b) If $S = \mathrm{Spec} \mathbf{F}$, we write $\mathrm{DA}_W(\mathbf{F}, A)$ for $\mathrm{DA}_W(\mathrm{Spec} \mathbf{F}, A)$ and define $\mathrm{DA}_W^{\mathrm{eff}}(\mathbf{F})$, $\mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F})$ and $\mathrm{DA}_{\mathrm{gm}, W}^{\mathrm{eff}}(\mathbf{F})$ as the corresponding thick subcategories.

In particular, we have a symmetric monoidal functor $\gamma^* : \mathrm{DA}_{\acute{e}t}(S, A) \rightarrow \mathrm{DA}_W(S, A)$ with right adjoint γ_* , and

Lemma 10.3.2. *There is a unique symmetric monoidal functor*

$$R_l^W : \mathrm{DA}_W(S, \mathbf{Z}_l) \rightarrow \hat{D}_{\acute{e}t}(S, \mathbf{Z}_l)$$

such that $R_l = R_l^W \circ \gamma^*$.

Proof. The definition of R_l^W on objects is forced since γ^* is the identity on objects; we need to define $R_l^W(f)$ for $f \in \mathrm{DA}_W(S, \mathbf{Z}_l)(M, N)$. Thus f is a morphism from M to $N \otimes f^*\mathbf{Z}^c$ in $\mathrm{DA}_{\acute{e}t}(S, \mathbf{Z}_l)$. We then get

$$R_l(f) : R_l(M) \rightarrow R_l(N \otimes f^*\mathbf{Z}^c) \simeq R_l(N) \otimes f^*R_l(\mathbf{Z}^c).$$

But $R_l(\eta) : \mathbf{Z}_l[0] \rightarrow R_l(\mathbf{Z}^c)$ is an isomorphism, in view of the cohomology sheaves of \mathbf{Z}^c ; this gives the desired morphism. \square

One interest of the categories $\mathrm{DA}_W(S, A)$ is that they allow us to reformulate the equivalent conditions of Corollary 9.8.4 in a way which justifies the title of this paper:

Proposition 10.3.3. *a) The conditions of Corollary 9.8.4 a) are also equivalent to the following: the restriction of R_l^W to $\mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F}, \mathbf{Z}_l)$ is fully faithful.*

b) If they are verified, then the same is true for the restriction of R_l^W to strongly dualisable objects of $\mathrm{DA}_W(S, \mathbf{Z}_l)$, for any $S \in \mathrm{Sm}(\mathbf{F})$.

Proof. a) By part b) of Corollary 9.8.4, the conditions of its part a) imply the said full faithfulness. Conversely, if we specialise to motives $M(X)$ and $\mathbf{Z}(n)[i]$ for X smooth and $n \geq 0$, $i \in \mathbf{Z}$, we get back Condition (iii) of Corollary 9.8.4 a).

b) This follows similarly from Corollary 9.8.5. \square

Recall the generator $e \in H_{\acute{\mathrm{e}}\mathrm{t}}^1(\mathbf{F}, \mathbf{Z}^c)$ of (9.2.5). We may view it as a map $e \in \mathrm{DA}_W(\mathbf{F})(\mathbf{Z}, \mathbf{Z}[1])$; it is easy to see that $e^2 = 0$, for example by using its compatibility with the generator of (4.3.1). Therefore, for any $M, N \in \mathrm{DA}_W(\mathbf{F})$, we have a complex of abelian groups

$$(10.3.1) \quad \dots \xrightarrow{\cdot e} \mathrm{DA}_W(S)(M, N[j]) \xrightarrow{\cdot e} \mathrm{DA}_W(S)(M, N[j+1]) \xrightarrow{\cdot e} \dots$$

Lemma 10.3.4. *The complex (10.3.1) has torsion cohomology groups.*

Proof. It is equivalent to show that the complex

$$\dots \xrightarrow{\cdot e} \mathrm{DA}_{\acute{\mathrm{e}}\mathrm{t}}(S, \mathbf{Q})(M, N \otimes \mathbf{Q}^c[j]) \xrightarrow{\cdot e} \mathrm{DA}_{\acute{\mathrm{e}}\mathrm{t}}(S, \mathbf{Q})(M, N \otimes \mathbf{Q}^c[j+1]) \xrightarrow{\cdot e} \dots$$

is acyclic. Inspection shows that the map

$$e : \mathbf{Q}[0] \oplus \mathbf{Q}[-1] \simeq \mathbf{Q}^c \rightarrow \mathbf{Q}^c[1] \simeq \mathbf{Q}[1] \oplus \mathbf{Q}[0]$$

inducing the differentials is the identity on $\mathbf{Q}[0]$ and 0 on the other summands. The claim then follows from a simple computation. \square

10.4. Two pairings. We now imitate §10.1. Let $M \in \mathrm{DA}_W(\mathbf{F})$. Composition of morphisms defines a pairing

$$(10.4.1) \quad \mathrm{Hom}(\mathbf{Z}, M)/\mathrm{tors} \otimes \mathrm{Hom}(M, \mathbf{Z}[1])/\mathrm{tors} \rightarrow \mathrm{Hom}(\mathbf{Z}, \mathbf{Z}[1]) = \mathbf{Z}[1/p].$$

Here the value of the target group comes from (9.2.5) (recall that $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(\mathbf{F})$, hence $\mathrm{DA}_W(\mathbf{F})$, is $\mathbf{Z}[1/p]$ -linear).

Thanks now to (9.2.2), we have a canonical isomorphism

$$\mathrm{Hom}(\mathbf{Z}, \mathbf{Q}/\mathbf{Z}[1]) \simeq (\mathbf{Q}/\mathbf{Z})'$$

(given by the choice of the arithmetic Frobenius of \mathbf{F}). Here we wrote $(\mathbf{Q}/\mathbf{Z})'$ for $\bigoplus_{l \neq p} \mathbf{Q}_l/\mathbf{Z}_l$. Thus we get a pairing

$$(10.4.2) \quad \mathrm{Hom}(\mathbf{Z}, M) \otimes \mathrm{Hom}(M, \mathbf{Q}/\mathbf{Z}[1]) \rightarrow (\mathbf{Q}/\mathbf{Z})'$$

The tautological exact sequence

$$0 \rightarrow \mathrm{Hom}(M, \mathbf{Z}[1]) \otimes \mathbf{Q}/\mathbf{Z} \rightarrow \mathrm{Hom}(M, \mathbf{Q}/\mathbf{Z}[1]) \\ \rightarrow \mathrm{Hom}(M, \mathbf{Z}[2])_{\mathrm{tors}} \rightarrow 0$$

shows that (10.4.2) induces a pairing

$$(10.4.3) \quad \mathrm{Hom}(\mathbf{Z}, M)_{\mathrm{tors}} \otimes \mathrm{Hom}(M, \mathbf{Z}[2])_{\mathrm{tors}} \rightarrow (\mathbf{Q}/\mathbf{Z})'$$

Lemma 10.4.1. *Let R be a commutative ring and $A \times B \rightarrow R$ be a pairing of flat R -modules.*

- a) *Suppose that the pairing is non-degenerate after $\otimes R_l$ for some prime ideal l of R , where R_l is the completion of R at l . Then it is non-degenerate.*
- b) *Assume that R is a Noetherian domain with quotient field K . If moreover $A \otimes_R K$ or $B \otimes_R K$ is a finite-dimensional K -vector space, then A and B are finitely generated.*
- c) *In the case of b), suppose moreover that the pairing is perfect after $\otimes R_l$ for every l . Then it is perfect.*

Proof. a) and b) are exactly the contents of [48, Lemma 3.9]: we refer to loc. cit. for the proof. The latter involves the composition

$$A_l \xrightarrow{a} \mathrm{Hom}_R(B, R)_l \xrightarrow{b} \mathrm{Hom}_{R_l}(B_l, R_l)$$

where $A_l = A \otimes_R R_l$, etc. Since B is finitely generated and R_l is flat over R [13, II.3.4, Th. 3 (iii)], b is injective (ibid., I.2.10, Prop. 11), thus if ba is bijective so is a . If this is true for all l , then $A \rightarrow \mathrm{Hom}(B, R)$ is bijective by the same proof as [13, II.3.3, Th. 1]. This proves c). \square

Theorem 10.4.2. *The conditions of Corollary 9.8.4 are also equivalent to the following:¹²*

- (i) $\mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z})$ is a finitely generated $\mathbf{Z}[1/2p]$ -module for any $M \in \mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F}, \mathbf{Z}[1/2])$.
- (ii) $\mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, N)$ is a finitely generated $\mathbf{Z}[1/2p]$ -module for any $M, N \in \mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F}, \mathbf{Z}[1/2])$.
- (iii) For some prime $l \neq 2, p$, $\mathrm{DA}_W(\mathbf{F}, \mathbf{Z}_l)(M, N)$ is a finitely generated \mathbf{Z}_l -module for any $M, N \in \mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F}, \mathbf{Z}_l)$.

These conditions imply:

¹²For the reason why 2 is inverted, see Theorem 6.8.1.

(*) For any $M \in \mathrm{DA}_{\mathrm{gm},W}(\mathbf{F}, \mathbf{Z}[1/2])$, (10.4.1) is a perfect pairing of finitely generated free $\mathbf{Z}[1/2p]$ -modules and (10.4.3) is a perfect pairing of finite groups.

Proof. Let (**) the condition of Proposition 10.3.3 a): we show (**) \Rightarrow (i) + (*) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (**).

(**) \Rightarrow (i) + (*): we first observe that $R_l(M) \in \hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)$ for any $l \neq p$. Indeed, this is true for $M = M(X)$, $f : X \rightarrow \mathrm{Spec} \mathbf{F}$ smooth by Deligne's finiteness theorem for l -adic cohomology (recall that $R_l(M(X)) = (Rf_* \mathbf{Z}_l)^*$). Let $A = \mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z})$. Clearly, (**) implies that $A \otimes \mathbf{Z}_l$ is finitely generated over \mathbf{Z}_l for any $l \neq p$. Further, Corollary 10.1.2 implies the conclusion of (*) after $\otimes \mathbf{Z}_l$. Lemma 10.4.1 then implies that A/A_{tors} is a finitely generated free $\mathbf{Z}[1/p]$ -module, that (10.4.3) is perfect and that $A\{l\}$ is finite for all $l \neq p$.

To get (i), we observe that the full subcategory of $\mathrm{DA}_{\mathrm{gm},\acute{e}t}(\mathbf{F}, \mathbf{Z}[1/2])$ formed of those M such that $\mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z}[i])$ is finitely generated for all $i \in \mathbf{Z}$ is triangulated and thick (stable under direct summands). Thus we may reduce to $M = M(X)(-n)$ with X smooth projective and $n \geq 0$. In this case, it is known that

$$\hat{D}_c^b(\mathbf{F}, \mathbf{Z}_l)(R_l(M(X)(-n)), \mathbf{Z}_l[i]) = H_{\mathrm{cont}}^i(X, \mathbf{Z}_l(n))$$

is torsion-free for almost all l [16, Th. 2]. Hence $\mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z}[i])$ has finite torsion, hence is finitely generated, for any i . This also finishes to prove (*).

(i) \Rightarrow (ii): this is clear from the formula

$$\mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, N) \simeq \mathrm{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])(N^* \otimes M, \mathbf{Z}).$$

(ii) \Rightarrow (iii): this is obvious.

(iii) \Rightarrow (**): let $M, N \in \mathrm{DA}_{\mathrm{gm},W}(\mathbf{F}, \mathbf{Z}_l)$. By hypothesis, we have a homomorphism of finitely generated \mathbf{Z}_l -modules:

$$(10.4.4) \quad \mathrm{DA}_W(\mathbf{F}, \mathbf{Z}_l)(M, N) \rightarrow D_c(\mathbf{F}, \mathbf{Z}_l)(R_l(M), R_l(N)).$$

To work in a triangulated category, we place ourselves in $\mathrm{DM}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)$ and rewrite the above as

$$\mathrm{DM}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(M, N \otimes \mathbf{Z}^c) \rightarrow \mathrm{DM}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(M, \Omega_l R_l(N)).$$

Let $C(N)$ be a cone of the map $N \otimes \mathbf{Z}^c \rightarrow \Omega_l R_l N$ in $D_{\acute{e}t}(\mathbf{F})$. By the same argument as in the proof of Lemma 9.8.1, $C(N) = 0$ if N is torsion, which implies that $C(N)$ is uniquely divisible in general. Then the exact sequence

$$\begin{aligned} \mathrm{DM}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(M, \Omega_l R_l(N)) &\rightarrow \mathrm{DM}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(M, C(N)) \\ &\rightarrow \mathrm{DM}_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)(M, N \otimes \mathbf{Z}^c[1]) \end{aligned}$$

shows that the \mathbf{Q}_l -vector space $\mathrm{DM}_{\acute{\mathrm{e}}\mathrm{t}}(\mathbf{F}, \mathbf{Z}_l)(M, C(N))$ is an extension of two finitely generated \mathbf{Z}_l -modules, hence is 0. Applying this to $N[-1]$ instead of N , we get the bijectivity of (10.4.4). \square

Remark 10.4.3. To control the torsion in Hom groups, we used [16, Th. 2] which rests ultimately on a theorem of Gabber on the torsion in the l -adic cohomology of smooth projective varieties [26]. This ingredient is not in the spirit of the rest of the paper: is there a way to avoid it?

Corollary 10.4.4. *Suppose that the conditions of Corollary 9.8.4 a) hold. Then, for any $M, N \in \mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F}, \mathbf{Z}[1/2])$, the complex (10.3.1) is bounded and has finite cohomology groups.*

Proof. It follows from Lemma 10.3.4 and Theorem 10.4.2 that, under the conditions of Corollary 9.8.4 a), the cohomology groups of (10.3.1) are finite. It remains to see its boundedness. In terms of $\mathrm{DA}_{\acute{\mathrm{e}}\mathrm{t}}(\mathbf{F}, \mathbf{Z}[1/2])$, amounts to the vanishing of $\mathrm{DA}_{\acute{\mathrm{e}}\mathrm{t}}(\mathbf{F}, \mathbf{Z}[1/2])(M, N \otimes \mathbf{Z}^c[j])$ for $|j| \gg 0$. By duality, we may reduce to $N = \mathbf{Z}$. The vanishing is stable under cones and direct summands, so it suffices to check it on motives of the form $M = M(X)(-n)$, $X \in \mathrm{Sm}(\mathbf{F})$. This amounts to the vanishing of the cohomology groups $H_{\acute{\mathrm{e}}\mathrm{t}}^i(X, \mathbf{Z}^c(n))$ for $|j| \gg 0$. In view of the cohomology sheaves of \mathbf{Z}^c , this reduces to the same vanishing for $H_{\acute{\mathrm{e}}\mathrm{t}}^j(X, \mathbf{Z}(n))$; since X has finite étale cohomological dimension, this in turn reduces in view of (5.6.1) to the same vanishing for $H_{\acute{\mathrm{e}}\mathrm{t}}^j(X, \mathbf{Q}(n))$. This group is 0 for $j > 2n$ by (4.1.1) and Theorem 5.5.2 b). To get the boundedness below we may even reduce to X smooth projective, and then $H_{\acute{\mathrm{e}}\mathrm{t}}^j(X, \mathbf{Q}(n)) = 0$ for $j < 2n$ by the assumption and Theorem 8.0.2. \square

In [51], we shall use Corollary 10.4.4 to give a conjectural expression for the special values of the zeta function of M , generalising Lichtenbaum's conjecture up to a power of p (cf. [50, Th. 72]).

10.5. Rational coefficients.

Proposition 10.5.1. *The following conditions are equivalent:*

- (i) *The Beilinson and Parshin conjectures hold for any smooth projective \mathbf{F} -variety.*
- (ii) *For any $M \in \mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$, the composition pairing*

$$\mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})(\mathbf{Z}, M) \otimes \mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})(M, \mathbf{Z}) \rightarrow \mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})(\mathbf{Z}, \mathbf{Z}) = \mathbf{Q}$$

is non-degenerate.

If this is the case,

- a) *The Hom groups in $\mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$ are finite-dimensional \mathbf{Q} -vector spaces.*
- b) *The functor Φ of (8.0.1) induces an equivalence of categories*

$$D^b(\mathcal{M}_{\mathrm{num}}(\mathbf{F}, \mathbf{Q})) \xrightarrow{\sim} \mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q}).$$

c) There is a unique t -structure on $\mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$ such that $S[-w]$ is in the heart for any $S \in \mathcal{M}_{\mathrm{num}}(\mathbf{F}, \mathbf{Q})$ of pure weight w .

d) For any $M \in \mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$, the action of Frobenius on $R_l(M)$ is semi-simple and its characteristic polynomial is independent of l .

Proof. The full category of $\mathrm{DM}_{\mathrm{gm}}(\mathbf{F}, \mathbf{Q})$ consisting of those M such that the hypothesis of (ii) is satisfied for $M[i]$ for any $i \in \mathbf{Z}$ is thick and triangulated. Hence (ii) is equivalent to the same statement restricted to $M = M(X)(-n)[-i]$, X smooth projective, $n \geq 0$, $i \in \mathbf{Z}$. Then the equivalence between (i) and (ii) (for this specific X) is an observation essentially due to Jakob Scholbach:

In this case, the said pairing is of the form

$$H_i(X, \mathbf{Q}(n)) \times H^i(X, \mathbf{Q}(n)) \rightarrow \mathbf{Q}.$$

By Poincaré duality, the left group may be rewritten $H^{2d-i}(X, \mathbf{Q}(d-n))$.

Suppose $i = 2n$. Then we find the intersection pairing between rational Chow groups. To say that it is nondegenerate is equivalent to say that rational and numerical equivalences agree.

Suppose $i \neq 2n$. Then either $i > 2n$ or $2d - i > 2(d - n)$. So one of the two groups in the pairing is 0, and (ii) is equivalent to the vanishing of the other, which is Parshin's conjecture.

a) follows from (i) by internal duality and reduction to smooth projective varieties. For b), see the proof of [50, Th. 56]. By [44], $\mathcal{M}_{\mathrm{num}}(\mathbf{F}, \mathbf{Q})$ is semi-simple and by [52], any simple $S \in \mathcal{M}_{\mathrm{num}}(\mathbf{F}, \mathbf{Q})$ has a weight; then c) follows readily from b). Finally, d) follows from b) and Theorem 2.2.1, since M is a direct sum of shifted simple motives. \square

Remarks 10.5.2. 1) Proposition 10.5.1 d) may be applied in particular to the motives $M(X)$ and $M_c(X)$ of Definition 6.7.1, for X of finite type over \mathbf{F} .

2) If we consider $\mathrm{DA}_W(\mathbf{F}, \mathbf{Q})$, the isomorphism stemming from (9.2.4)

$$\mathrm{DA}_W(\mathbf{F}, \mathbf{Q})(M, N) \simeq \mathrm{DM}(\mathbf{F}, \mathbf{Q})(M, N) \oplus \mathrm{DM}(\mathbf{F}, \mathbf{Q})(M, N[-1])$$

easily shows that (10.4.1) boils down to the pairing of Proposition 10.5.1 (ii).

3) Let $l \neq p$. Using Theorem 5.5.1 b) and the duality of Proposition 10.1.1, it is easy to see that one has a perfect pairing for any $M \in \mathrm{DA}_{\mathrm{gm}, W}(\mathbf{F}, \mathbf{Z}/l)$:

$$\mathrm{Hom}(\mathbf{Z}, M) \times \mathrm{Hom}(M, \mathbf{Z}[1]) \rightarrow \mathrm{Hom}(\mathbf{Z}, \mathbf{Z}[1]) = \mathbf{Z}/l.$$

This suggests that the conditions of Proposition 10.5.1 might be equivalent to those of Theorem 10.4.2. Unfortunately a proof of this fails, as we shall see in Remark 11.6.2.

11. WEIL-ÉTALE REFORMULATION

11.1. Lichtenbaum's Weil-étale topology and cohomology.

Definition 11.1.1 (Lichtenbaum [55]). a) Let X/\mathbf{F} be a scheme of finite type, and let $\bar{X} = X \otimes_{\mathbf{F}} \bar{\mathbf{F}}$. A *Weil-étale sheaf* on X is

- a sheaf \mathcal{F} on \bar{X} ;
- a map $\varphi_{\mathcal{F}} : \varphi^* \mathcal{F} \rightarrow \mathcal{F}$, φ the Frobenius of \mathbf{F} .

Weil-étale sheaves form the *Weil-étale topos* of X . There is an underlying site X_W described in [55, §2].

b) If \mathcal{F} is an abelian Weil-étale sheaf, one defines

$$H_W^0(X, \mathcal{F}) := H_{\text{ét}}^0(\bar{X}, \mathcal{F})^{\varphi_{\mathcal{F}}}.$$

c) Weil-étale cohomology is defined as the derived functors of H_W^0 .

The definition of the Weil-étale cohomology yields short exact sequences

$$0 \rightarrow H_{\text{ét}}^{i-1}(\bar{X}, \mathcal{F})_{\varphi} \rightarrow H_W^i(X, \mathcal{F}) \rightarrow H_{\text{ét}}^i(\bar{X}, \mathcal{F})^{\varphi} \rightarrow 0$$

which show that Weil-étale cohomology commutes with filtering direct limits of sheaves and has finite cohomological dimension.

There is a morphism γ from the Weil-étale topos to the étale topos (forget that Frobenius acts continuously), hence adjoint functors: $Ab(X_{\text{ét}}) \xrightleftharpoons[\gamma_*]{\gamma^*} Ab(X_W)$.

11.2. **Two basic results.** They are due to Geisser [28, Th. 5.1 and 5.3]:

Theorem 11.2.1. a) For any $C \in D(X_{\text{ét}})$, the map

$$R\gamma_* \mathbf{Z} \otimes^L C \rightarrow R\gamma_* \gamma^* C$$

of (9.3.2) is an isomorphism.

b) $R\gamma_* \gamma^* \mathbf{Z} \simeq \mathbf{Z}^c$.

Note that b) is the promised better justification of the monoid structure on \mathbf{Z}^c (see beginning of §10.3)!

Since $D(X_{\text{ét}})$ is generated by strongly dualisable objects¹³ and γ^* preserves compact objects, a) is in fact a formal consequence of Corollary 9.3.2 and Remark 9.3.3. One will find another proof of b) in Appendix A.

The description of \mathbf{Z}^c given in §9.2 then gives:

Corollary 11.2.2. a) One has long exact sequences

$$\dots \rightarrow H_{\text{ét}}^i(X, C) \rightarrow H_W^i(X, \gamma^* C) \rightarrow H_{\text{ét}}^{i-1}(X, C) \otimes \mathbf{Q} \xrightarrow{\partial} H_{\text{ét}}^i(X, C) \rightarrow \dots$$

¹³namely, locally constant constructible sheaves

for C a complex of étale sheaves on X , where ∂ is described by (9.2.3).

b) If C is torsion, $H_{\text{ét}}^i(X, C) \xrightarrow{\sim} H_W^i(X, \gamma^*C)$.

c) If C is a complex of sheaves of \mathbf{Q} -vector spaces,

$$H_W^i(X, \gamma^*C) \simeq H_{\text{ét}}^i(X, C) \oplus H_{\text{ét}}^{i-1}(X, C).$$

11.3. Inflicting the Weil-étale topology on DA. One can faithfully follow the template of [4, §3] and get triangulated categories $\widetilde{\text{DA}}_W(S, A)$ for S a \mathbf{F} -scheme of finite type and A a commutative ring of coefficients. To do this we should define a Weil-étale site $\text{Sm}_W(S)$. We can do it by simply mimicking Lichtenbaum's definition in [55]:

We first define a “big Weil-étale” site $\text{Sch}_W(\mathbf{F})$. Objects of the underlying category are $\overline{\mathbf{F}}$ -schemes of finite type. Let \overline{S} be such a scheme; for $n \in \mathbf{Z}$ we write $\overline{S}^{(n)} = (\varphi^n)^*\overline{S}$; there is a canonical \overline{F} -morphism $\overline{S}^{(n)} \rightarrow \overline{S}$ (iterated relative Frobenius). If $\overline{S}, \overline{T} \in \text{Sch}_W(\mathbf{F})$ with \overline{S} connected, a morphism $\overline{S} \rightarrow \overline{T}$ is a \overline{F} -morphism $\overline{S} \rightarrow \overline{T}^{(n)}$ for some $n \in \mathbf{Z}$; for \overline{S} arbitrary a morphism from \overline{S} to \overline{T} is a collection of such morphisms over the connected components of \overline{S} . Finally, coverings are étale coverings.

If S/\mathbf{F} is of finite type, we define $\text{Sm}_W(S)$ as the full subsite of $\text{Sch}_W(\mathbf{F})$ whose objects are smooth \overline{S} -schemes.

One still has a pair of adjoint functors

$$\text{DA}_{\text{ét}}(S, A) \underset{\gamma_*}{\overset{\gamma^*}{\rightleftarrows}} \widetilde{\text{DA}}_W(S, A).$$

Here are some good features of this construction:

- Lemma 11.3.1.** a) γ^* is fully faithful if $A = \mathbf{Z}/l^\nu$.
b) Theorem 11.2.1 a) remains valid here.
c) The functor γ^* factors canonically as a composition

$$\text{DA}_{\text{ét}}(S, A) \xrightarrow{\gamma^*} \text{DA}_W(S, A) \xrightarrow{\iota} \widetilde{\text{DA}}_W(S, A)$$

where $\text{DA}_W(S, A)$ is the category from Definition 10.3.1 a) and ι is fully faithful. Moreover, the functor γ_* of §10.3 is the restriction to $\text{DA}_W(S, A)$ of the present functor γ_* .

d) If $A = \mathbf{Z}/l^\nu$, $\text{DA}_W(S, A)$ is a triangulated subcategory of $\widetilde{\text{DA}}_W(S, A)$.

Proof. a) follows from Corollary 11.2.2 b). b) is formal by localisation from Theorem 11.2.1 a). c) follows formally from b) and Theorem 11.2.1 b). Finally, d) follows easily from a). \square

Remark 11.3.2. Unfortunately $\text{DA}_W(S, A)$ is not a triangulated subcategory of $\widetilde{\text{DA}}_W(S, A)$ in general, e.g. for $A = \mathbf{Z}$ or \mathbf{Q} . Indeed, consider a fibre F of the map $e \in \widetilde{\text{DA}}_W(\mathbf{F})(\mathbf{Z}, \mathbf{Z}[1])$ coming from (9.2.5) (see above (10.3.1)). Then F is represented by a Weil-étale sheaf (still denoted by

F), extension of \mathbf{Z} by itself. (One can check that the Frobenius φ acts on $F \simeq \mathbf{Z} \oplus \mathbf{Z}$ by

$$\varphi(m, n) = (m, m + n),$$

compare [46, (4.1)] and Appendix A, but we won't need this here.)

I claim that F is not in the essential image of γ^* . It is enough to show this for its image $F_{\mathbf{Q}}$ in $\widetilde{\overline{DA}}_W(\mathbf{F}, \mathbf{Q})$. If $F_{\mathbf{Q}}$ were of the form γ^*C for $C \in \text{DA}_{\text{ét}}(\mathbf{F}, \mathbf{Q})$, we would have

$$\gamma_*F_{\mathbf{Q}} \simeq \gamma_*\gamma^*C \simeq C \otimes \gamma_*\mathbf{Q} \simeq C \otimes \mathbf{Q}^c \simeq C \oplus C[1].$$

But it is easy to compute $\gamma_*F_{\mathbf{Q}}$ as the fibre of γ_*e : one finds

$$\gamma_*F_{\mathbf{Q}} \simeq \mathbf{Q} \oplus \mathbf{Q}[1].$$

One would then have $C \oplus C[1] \simeq \mathbf{Q} \oplus \mathbf{Q}[1]$ in $\text{DA}_{\text{ét}}(\mathbf{F}, \mathbf{Q})$. Taking cohomology sheaves, this easily gives $C \simeq \mathbf{Q}[0]$. Hence a contradiction, since then $F_{\mathbf{Q}} = \gamma^*C = \mathbf{Q}[0]$ would be of \mathbf{Q} -rank 1.

11.4. The Weil-étale l -adic realisation functor. Remark 11.3.2 leads to

Definition 11.4.1. We write $\overline{DA}_W(S, A)$ for the localising subcategory of $\widetilde{\overline{DA}}_W(S, A)$ generated by $DA_W(S, A) = \gamma^*\text{DA}_{\text{ét}}(S, A)$. It is a \otimes -triangulated subcategory.

Lemma 11.4.2. *If $A = \mathbf{Z}/l^n$, $\overline{DA}_W(S, A) = DA_W(S, A) \simeq \text{DA}_{\text{ét}}(S, A)$.*

Proof. This follows formally from Lemma 11.3.1 a). \square

Proposition 11.4.3. *Let $l \neq 2$. The functor R_l^W of Lemma 10.3.2 extends to $\overline{DA}_W(S)$, and has a right adjoint Ω_l^W . Similarly with coefficients \mathbf{Z}_l .*

Proof. It suffices to do it with \mathbf{Z}_l coefficients. This follows immediately from Ayoub's construction of R_l [4, Def. 5.6]: it is the composition of two functors

$$\text{DA}_{\text{ét}}(S, \mathbf{Z}_l) \rightarrow \text{DA}_{\text{ét}}(S, \mathbf{Z}/l^\bullet) \rightarrow D_{\text{ét}}(S, \mathbf{Z}/l^\bullet)$$

where $\text{DA}_{\text{ét}}(S, \mathbf{Z}/l^\bullet)$ is the “profinite” version of $\text{DA}_{\text{ét}}(S, \mathbf{Z}_l)$ described in [4, §5]. There is an obvious functor

$$\text{DA}_{\text{ét}}(S, \mathbf{Z}/l^\bullet) \xrightarrow{\gamma^*} \overline{DA}_W(S, \mathbf{Z}/l^\bullet)$$

where the right category is defined similarly, and γ^* has a right adjoint γ_* . These adjoints are clearly compatible with the projections

$$\text{DA}_{\text{ét}}(S, \mathbf{Z}/l^\bullet) \xrightarrow{s_{\text{ét}}^*} \text{DA}_{\text{ét}}(S, \mathbf{Z}/l^s), \quad \overline{DA}_W(S, \mathbf{Z}/l^\bullet) \xrightarrow{s_W^*} \overline{DA}_W(S, \mathbf{Z}/l^s).$$

By [4, Lemma 5.3], the $(s_{\text{ét}}^*)$ form a conservative set of functors, and the same proof shows that the same holds for the (s_W^*) . From this and Lemma 11.4.2, one deduces that the unit and counit of the adjunction (γ^*, γ_*) are

isomorphisms, hence γ^* and γ_* are quasi-inverse equivalences of categories. The functor R_l^W is then defined as the composition

$$\overline{DA}_W(S, \mathbf{Z}_l) \rightarrow \overline{DA}_W(S, \mathbf{Z}/l^\bullet) \xrightarrow{\gamma^*} DA_{\acute{e}t}(S, \mathbf{Z}/l^\bullet) \xrightarrow{R_l} D_{\acute{e}t}(S, \mathbf{Z}/l^\bullet).$$

It remains to show that the image of R_l^W is contained in $\hat{D}_{\acute{e}t}(S, \mathbf{Z}_l)$. This is a localising subcategory of $D_{\acute{e}t}(S, \mathbf{Z}/l^\bullet)$; since it contains the image of R_l and since $\gamma^* DA_{\acute{e}t}(S, \mathbf{Z}_l)$ generates $\overline{DA}_W(S, \mathbf{Z}_l)$, $\hat{D}_{\acute{e}t}(S, \mathbf{Z}_l)$ also contains the image of R_l^W . The existence of Ω_l^W follows from Theorem 7.1.2. \square

On the other hand, there are too many objects in $\widetilde{DA}_W(S, A)$:

- (1) Even if $A = \mathbf{Z}/l^\nu$, γ^* is not essentially surjective. For example, the free $\mathbf{Z}/l^\nu[\varphi, \varphi^{-1}]$ -module of rank 1 defines an object of $\widetilde{DA}_W(\mathbf{F}, \mathbf{Z}/l^\nu)$ which is not in the essential image of γ^* . In particular, it is not clear whether R_l^W extends to $\widetilde{DA}_W(\mathbf{F})$.
- (2) $DA_W(\mathbf{F}, \mathbf{Q})$ is not dense in $\overline{DA}_W(\mathbf{F}, \mathbf{Q})$. For example, the sheaf $\mathbf{Q}\langle n \rangle$ where Frobenius acts by multiplication by q^n has no nonzero Weil-étale cohomology if $n \neq 0$ (this example was given by Geisser). Hence it is right orthogonal to $DA_{\text{gm}, W}(\mathbf{F}, \mathbf{Q})$, hence to $DA_W(\mathbf{F}, \mathbf{Q})$.

11.5. A reformulation of Proposition 10.3.3. The following statement is a bit more elegant:

Proposition 11.5.1. *The conditions of Corollary 9.8.4 a) are also equivalent to the following: the unit map*

$$\mathbf{Z}_l \rightarrow \Omega_l^W R_l^W \mathbf{Z}_l$$

of the adjunction in Proposition 11.4.3 is an isomorphism.

Proof. Using Theorem 11.2.1 a) and the fact that $\gamma^* DA_{\acute{e}t}(\mathbf{F}, \mathbf{Z}_l)$ generates $\overline{DA}_W(\mathbf{F}, \mathbf{Z}_l)$, this is formally equivalent to Corollary 9.8.4 a) (vii). \square

11.6. A reformulation of Theorem 10.4.2. Consider the composite functor

$$\begin{aligned} D(\mathbf{Z}[1/2p]) &\simeq d_{\leq 0} DA(\mathbf{F}, \mathbf{Z}[1/2p]) \rightarrow DA_{\acute{e}t}(\mathbf{F}, \mathbf{Z}[1/2]) \\ &\rightarrow \overline{DA}_W(\mathbf{F}, \mathbf{Z}[1/2]). \end{aligned}$$

It commutes with representable direct sums and preserves compact objects, hence has a right adjoint $R\Gamma$ which in turn has a right adjoint Π .

The cohomology groups of $R\Gamma(\mathbf{Z})$ are given by (9.2.5); in particular, we have a canonical map $R\Gamma(\mathbf{Z}) \rightarrow \mathbf{Z}[1]$, which yields a map in $\overline{DA}_W(\mathbf{F}, \mathbf{Z}[1/2])$

$$(11.6.1) \quad \mathbf{Z}[-1] \xrightarrow{\theta} \Pi(\mathbf{Z}).$$

Theorem 11.6.1. *The conditions of Corollary 9.8.4 a) are equivalent to the invertibility of (11.6.1).*

Proof. Let $M \in \overline{\text{DA}}_W(\mathbf{F}, \mathbf{Z}[1/2])$. Evaluating (11.6.1) against M produces a map

$$\overline{\text{DA}}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z}[1]) \xrightarrow{\theta_M} D(\mathbf{Z}[1/2p])(R\Gamma(M), \mathbf{Z}).$$

The right hand side fits in an exact sequence

$$\begin{aligned} 0 \rightarrow \text{Ext}^1(\overline{\text{DA}}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z}[1]), \mathbf{Z}[1/p]) &\rightarrow D(\mathbf{Z}[1/2p])(R\Gamma(M), \mathbf{Z}) \\ &\rightarrow \text{Hom}(\overline{\text{DA}}_W(\mathbf{F}, \mathbf{Z}[1/2])(M, \mathbf{Z}), \mathbf{Z}[1/p]) \rightarrow 0 \end{aligned}$$

One checks that θ_M induces the pairings (10.4.1) and (10.4.3), and that its bijectivity is equivalent to their perfectness (compare proof of Corollary 10.1.2). \square

Remark 11.6.2. As in §10.5, one easily sees that, in (11.6.1), $\theta \otimes \mathbf{Z}/m$ is an isomorphism for any $m \neq 0$. Hence θ is an isomorphism if and only if $\theta \otimes \mathbf{Q}$ is an isomorphism. The problem to go further is that the map

$$\Pi(\mathbf{Z}) \otimes \mathbf{Q} \rightarrow \Pi(\mathbf{Q})$$

is not *a priori* an isomorphism. It will be, provided Π commutes with homotopy colimits, which happens if and only if $R\Gamma$ preserves compact objects (Lemma 7.1.4), in other words, if $\text{Hom}(M, \mathbf{Z}[i])$ is finitely generated and 0 for almost all i , for any $M \in \text{DA}_W(\mathbf{F}) \dots$

The isomorphisms $\text{Tor}(\mathbf{Z}/m, \mathbf{Z}/m) \simeq \mathbf{Z}/m$ provide a pairing in $D(\mathbf{Z})$

$$\mathbf{Q}/\mathbf{Z} \times \mathbf{Q}/\mathbf{Z} \rightarrow \mathbf{Q}/\mathbf{Z}[1].$$

This pairing is perfect after $\otimes \mathbf{Q}$ (in $D(\mathbf{Q})$) and after $\overset{L}{\otimes} \mathbf{Z}/m$ (in $D(\mathbf{Z}/m)$) for any m . But it is clearly not perfect, since $\text{Hom}(\mathbf{Q}/\mathbf{Z}, \mathbf{Q}/\mathbf{Z})$ is not a torsion group.

12. A POSITIVE CASE OF THE CONJECTURES

12.1. **Motives of weight ≤ 1 .** Write

$$w_{\leq 1} \text{DM}_{\text{gm}, \text{ét}}^{\text{eff}}(\mathbf{F})$$

for the thick subcategory of $\text{DM}_{\text{gm}, \text{ét}}^{\text{eff}}(\mathbf{F})$ generated by the $M(X)$ with X smooth of dimension 0 and the $\Phi(h_1(C))$ where C is a smooth projective curve and Φ is the integral version of the functor in (8.0.1). (Note that C always has a zero-cycle of degree 1 by Weil's theorem, hence an integral Chow-Künneth decomposition.)

12.2. **A positive result.** The aim of this section is to prove:

Theorem 12.2.1. *If $M \in w_{\leq 1} \text{DM}_{\text{gm}, \text{ét}}^{\text{eff}}(\mathbf{F})$, (9.6.1) is an isomorphism.*

Proof. By Lemma 9.8.1, it suffices to prove this after tensoring with \mathbf{Q} . For simplicity, we drop Φ from the notation: we reduce to the case $M = M(Y)$, Y smooth (projective) of dimension 0, or $M = h_1(C)$, C a smooth projective curve. The case of an Artin motive is trivial and left to the reader. As we shall see, that of h_1 of a curve follows essentially from Theorem 2.2.1 plus Tate's theorem on homomorphisms of abelian varieties [76].

It is equivalent to show that the morphism of Corollary 9.8.4 b) is an isomorphism for any N when $M = h_1(C)$ and after tensoring with \mathbf{Q} . We reduce as usual to $N = M(X)[-i]$, X smooth projective and $i \in \mathbf{Z}$.

Observe that the l -adic realisation of $h_1(C)$ is $T_l(J)[1]$, where J is the Jacobian of C . Thus we want to show that the homomorphism

$$(12.2.1) \quad \text{DM}_{\text{gm}, \text{ét}}^{\text{eff}}(k)(M(X), \Gamma^{\text{eff}} \otimes h_1(C)[i-1]) \otimes \mathbf{Q} \rightarrow H_{\text{cont}}^i(X, V_l(J))$$

is an isomorphism for all $i \in \mathbf{Z}$.

By duality, the left hand side of (12.2.1) is isomorphic to

$$\text{DM}_{\text{gm}}^{\text{ét}, \text{eff}}(k)(M(X) \otimes h_1(C), \Gamma^{\text{eff}}(1)[i+1]) \otimes \mathbf{Q}$$

which is a direct summand of

$$H_{\text{ét}}^{i+1}(X \times C, \Gamma^{\text{eff}}(1)) \otimes \mathbf{Q} \simeq H_{\text{ét}}^{i+1}(X \times C, \mathbf{Z}(1)) \otimes \mathbf{Q}_l \oplus H_{\text{ét}}^{i+2}(X \times C, \mathbf{Z}(1)) \otimes \mathbf{Q}_l$$

cf. (9.2.4) and (9.7.1). We have

$$H_{\text{ét}}^i(X \times C, \mathbf{Z}(1)) \otimes \mathbf{Q}_l = \begin{cases} \text{Pic}(X \times C) \otimes \mathbf{Q}_l & \text{if } i = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Using the isomorphism (theorem of the cube + finiteness of $\text{Pic}^0(X \times C)$)

$$\text{Pic}(X \times C) \otimes \mathbf{Q}_l \simeq \text{NS}(X) \otimes \mathbf{Q}_l \oplus \text{NS}(C) \otimes \mathbf{Q}_l \oplus \text{Hom}(\text{Alb}(X), J) \otimes \mathbf{Q}_l$$

we see that the direct summand $\text{DM}^{\text{ét}}(k)(M(X) \otimes h_1(C), \mathbf{Z}(1)[2]) \otimes \mathbf{Q}_l$ of $H_{\text{ét}}^2(X \times C, \mathbf{Z}(1)) \otimes \mathbf{Q}_l$ gets identified with $\text{Hom}(\text{Alb}(X), J) \otimes \mathbf{Q}_l$.

A weight computation shows that the right hand side of (12.2.1) is

$$H_{\text{cont}}^i(X, V_l(J)) = \begin{cases} H_{\text{cont}}^1(\bar{X}, V_l(J))^G & \text{si } i = 1 \\ H_{\text{cont}}^1(\bar{X}, V_l(J))_G & \text{si } i = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, the action of G on $H_{\text{cont}}^1(\bar{X}, V_l(J)) \simeq \text{Hom}(V_l(\text{Alb}(X)), V_l(J))$ is semi-simple, hence the composition

$$H_{\text{cont}}^1(\bar{X}, V_l(J))^G \rightarrow H_{\text{cont}}^1(\bar{X}, V_l(J)) \rightarrow H_{\text{cont}}^1(\bar{X}, V_l(J))_G$$

is an isomorphism; by Lemma 4.3.1, this isomorphism may be identified with cup-product by $e \in H_{\text{cont}}^1(k, \mathbf{Q}_l)$:

$$H_{\text{cont}}^1(X, V_l(J)) \xrightarrow{e} H_{\text{cont}}^2(X, V_l(J)).$$

Poincaré duality now realises $H_{\text{cont}}^i(X, V_l(J))$ as a direct summand of $H_{\text{cont}}^{i+1}(X \times C, \mathbf{Q}_l(1))$.

In (12.2.1), the two sides are therefore 0 for $i \neq 1, 2$; for $i = 1, 2$ they fit in a commutative diagram

$$\begin{array}{ccc} \text{Hom}(\text{Alb}(X), J) \otimes \mathbf{Q}_l & \xrightarrow{u} & \text{Hom}(V_l(\text{Alb}(X)), V_l(J))^G \\ \parallel & & \downarrow \wr \\ \text{Hom}(\text{Alb}(X), J) \otimes \mathbf{Q}_l & \xrightarrow{v} & \text{Hom}(V_l(\text{Alb}(X)), V_l(J))_G. \end{array}$$

By Tate's theorem [76], u is an isomorphism. The diagram shows that so is v , hence (12.2.1) is indeed an isomorphism for any $i \in \mathbf{Z}$. \square

APPENDIX A. A LETTER TO T. GEISSER

Paris, May 2, 2004.

Dear Thomas,

To start with, let Γ be a group, A a Γ -module and $e \in H^1(\Gamma, A)$ a class. To e corresponds an extension of Γ -modules

$$0 \rightarrow A \rightarrow A' \rightarrow \mathbf{Z} \rightarrow 0.$$

It is an exercise of homological algebra to check that A' may be described¹⁴ as $\mathbf{Z} \times A$ (the underlying abelian group) with Γ -action given by

$$a(r, s) = (r, re(a) + as).$$

This can be used as follows. Given a complex of Γ -modules C , cup-product by e

$$\cdot e : \mathbb{H}^i(\Gamma, C) \rightarrow \mathbb{H}^{i+1}(\Gamma, A \overset{L}{\otimes} C)$$

may be described, up to sign, as the boundary morphism ∂_C of the exact triangle

$$A \overset{L}{\otimes} C \rightarrow A' \overset{L}{\otimes} C \rightarrow C \xrightarrow{\partial_C} A \overset{L}{\otimes} C[1].$$

(This is another exercise of homological algebra: observe that $e = \partial_{\mathbf{Z}}(1)$ and compare [67, Ch. VIII, §3, Prop. 5]).

All this can be [made] topological. In particular, for $\Gamma = \hat{\mathbf{Z}}$, $A = \hat{\mathbf{Z}}$ and $e = e$ we get a corresponding $A' = \hat{M}$ such that $\hat{M} \otimes \mathbf{Q}/\mathbf{Z} = \tilde{M}$ (notation from [46, §4]).

¹⁴Up to sign, as in all this letter.

This first justifies the claim in loc. cit., Prop. 4.4 that the bottom row in the diagram is cup-product by e . Second, it gives another way to look at M : it is the push-out of the extension

$$(A.0.2) \quad 0 \rightarrow \hat{\mathbf{Z}} \otimes \mathbf{Q} \rightarrow \hat{M} \otimes \mathbf{Q} \rightarrow \mathbf{Q} \rightarrow 0$$

by the projection $\hat{\mathbf{Z}} \otimes \mathbf{Q} \rightarrow \hat{\mathbf{Z}} \otimes \mathbf{Q}/\mathbf{Z} = \mathbf{Q}/\mathbf{Z}$. Viewed from this side, we may recover $\gamma : \mathbf{Q} \rightarrow M$ as the composite $\mathbf{Q} \rightarrow \hat{\mathbf{Z}} \otimes \mathbf{Q} \rightarrow \mathbf{Q}/\mathbf{Z} \rightarrow M$.

Another thing you asked me is to explain the proof of [46, Cor. 4.8]. First, by Theorem 4.6, the map

$$\mathbb{H}^1(\mathbf{F}_p, \mathbf{Z}^c) \otimes \mathbf{Z}_l = \mathbb{H}^1(\mathbf{F}_p, \mathbf{Z}^c \otimes \mathbf{Z}_l) \rightarrow \mathbb{H}^1(\mathbf{F}_p, \mathbf{Z}_l(0)^c) = H_{\text{cont}}^1(\mathbf{F}_p, \mathbf{Z}_l)$$

is an isomorphism and therefore sends a generator e_0 of $\mathbb{H}^1(\mathbf{F}_p, \mathbf{Z}^c)$ to ue , where u is an l -adic unit. To prove that $u = 1$, it suffices to go mod l^ν for all ν and to prove that the induced map

$$\mathbf{Z} = \mathbb{H}^1(\mathbf{F}_p, \mathbf{Z}^c) \rightarrow H^1(\mathbf{F}_p, \mathbf{Z}/l^\nu) = \mathbf{Z}/l^\nu$$

is the projection. For this, observe that by definition of \tilde{M} and M , we have a short exact sequence

$$(A.0.3) \quad 0 \rightarrow \mathbf{Z} \rightarrow M \rightarrow \tilde{M} \rightarrow 0$$

and that the map $\mathbf{Z} \rightarrow M$ splits the exact sequence of [46, Lemma 4.3]. This implies that $e_0 \in H^1(\hat{\mathbf{Z}}, \mathbf{Z}^c)$ is the image of the generator of $H^0(\hat{\mathbf{Z}}, \mathbf{Z}) = H^1(\hat{\mathbf{Z}}, \mathbf{Z}[-1])$ under the map of complexes

$$(A.0.4) \quad \mathbf{Z}[-1] \rightarrow \mathbf{Z}^c$$

defined by (A.0.3). Another way to formulate this is that the morphism defined by e_0 in the derived category is realised by (A.0.4). To prove the claimed compatibility, we now have to identify the cone of (A.0.4) tensored with \mathbf{Z}/l^ν with $\hat{M} \otimes^L \mathbf{Z}/l^\nu$. This cone is $[\mathbf{Q} \rightarrow \tilde{M}]$; we have a string of quasi-isomorphisms

$$[\mathbf{Q} \rightarrow \tilde{M}] \otimes \mathbf{Z}/l^\nu \xrightarrow{\sim} \tilde{M} \otimes^L \mathbf{Z}/l^\nu[-1] \simeq \hat{M} \otimes^L \mathbf{Q}/\mathbf{Z} \otimes^L \mathbf{Z}/l^\nu[-1] \simeq \hat{M} \otimes^L \mathbf{Z}/l^\nu.$$

Then for the second claim of Corollary 4.8. First note that the statement is nonsense. In fact, the splitting $\mathbf{Q}^c \simeq \mathbf{Q} \oplus \mathbf{Q}[-1]$ of Corollary 4.5 defines a “projection onto H^0 ” $\mathbf{Q}^c \rightarrow \mathbf{Q}$ and an “inclusion of H^1 ” $\mathbf{Q}[-1] \rightarrow \mathbf{Q}^c$. (Note that the only endomorphism of $\mathbf{Q} \oplus \mathbf{Q}[-1]$ that induces the identity on cohomology is the identity because $\text{Ext}_{\mathbf{Z}}^1(\mathbf{Q}, \mathbf{Q}) = 0$, so these projection and inclusion are unambiguous.) The correct statement is now that the corresponding composition

$$\mathbf{Q}^c \rightarrow \mathbf{Q} \rightarrow \mathbf{Q}^c[1]$$

is given by cup product by e .

In order to make sense of this claim, I should define a product

$$\mathbf{Q}^c \otimes^L \mathbf{Q}^c \rightarrow \mathbf{Q}^c$$

matching the product

$$\mathbf{Q}_l(0)^c \otimes^L \mathbf{Q}_l(0)^c \rightarrow \mathbf{Q}_l(0)^c$$

which corresponds to the ‘‘continuous’’ product $\mathbf{Q}_l \otimes \mathbf{Q}_l \rightarrow \mathbf{Q}_l$. There is a stupid way to do this, and then one can use the isomorphism of Theorem 4.6 b) tensored with \mathbf{Q} and the obvious description of cup-product by e on $\mathbf{Q}_l(0)^c$: this is ugly but it works, and rationally this is all there is to it. Nevertheless let me do all this integrally as it might be useful. We have a short exact sequence of topological modules over $\hat{\mathbf{Z}}$:

$$0 \rightarrow \hat{\mathbf{Z}} \rightarrow \hat{M} \otimes \mathbf{Q} \rightarrow M \rightarrow 0$$

hence a flat version of \mathbf{Z}^c is given by the length 1 complex

$$(A.0.5) \quad [\mathbf{Q} \oplus \hat{\mathbf{Z}} \rightarrow \hat{M} \otimes \mathbf{Q}]$$

(note that, as already remarked, the map $\gamma : \mathbf{Q} \rightarrow M$ lifts canonically to $\hat{M} \otimes \mathbf{Q}$ via (A.0.2).) This gives us a version of $\mathbf{Z}^c \otimes^L \mathbf{Z}^c$:

$$[\mathbf{Q} \oplus (\mathbf{Q} \otimes \hat{\mathbf{Z}})^2 \oplus \hat{\mathbf{Z}} \rightarrow (\hat{M} \otimes \mathbf{Q})^2 \rightarrow (\hat{M} \otimes \mathbf{Q})^{\otimes 2}].$$

We define the product $\mathbf{Z}^c \otimes^L \mathbf{Z}^c \rightarrow \mathbf{Z}^c$ by the cochain map which is the projection on $\mathbf{Q} \oplus \hat{\mathbf{Z}}$ in degree 0 and the sum in degree 1. With this definition, the diagrams

$$\begin{array}{ccc} \mathbf{Z}^c \otimes^L \mathbf{Z}^c & \longrightarrow & \mathbf{Z}^c \\ \downarrow & & \downarrow \\ \mathbf{Z}/l^\nu \otimes^L \mathbf{Z}/l^\nu & \longrightarrow & \mathbf{Z}/l^\nu \end{array}$$

obviously commute coherently (for solid versions of the maps obtained by naïvely tensoring the flat version of \mathbf{Z}^c by \mathbf{Z}/l^ν), which proves the compatibility with the product $\mathbf{Z}_l(0)^c \otimes^L \mathbf{Z}_l(0)^c \rightarrow \mathbf{Z}_l(0)^c$. If we now tensor with \mathbf{Q} , we find that on $\mathbf{Q}^c \simeq \mathbf{Q} \oplus \mathbf{Q}[-1]$, this product is given by the identity $\mathbf{Q} \rightarrow \mathbf{Q}$ in degree 0 and the sum $\mathbf{Q} \oplus \mathbf{Q} \rightarrow \mathbf{Q}$ in degree 1. From this and the description (A.0.4) of the first map of the composition

$$\mathbf{Q}^c[-1] \xrightarrow{\cdot e_0} \mathbf{Q}^c \otimes^L \mathbf{Q}^c \rightarrow \mathbf{Q}^c$$

you easily deduce the second claim of [46, Cor. 4.8].

You might also be interested in the following construction of an isomorphism $\mathbf{Z}^c \xrightarrow{\sim} R\gamma_* \mathbf{Z}$. By adjunction I need to give a map $\varphi : \gamma^* \mathbf{Z}^c \rightarrow \mathbf{Z}$.

Now to describe $\gamma^*\mathbf{Z}^c$, I may use a discrete version of the topological $\hat{\mathbf{Z}}$ -module \hat{M} , say \bar{M} :

$$0 \rightarrow \mathbf{Z} \rightarrow \bar{M} \rightarrow \mathbf{Z} \rightarrow 0.$$

Then a flat version of $\gamma^*\mathbf{Z}^c$ is given by

$$\begin{aligned} \mathbf{Q} \oplus \mathbf{Z} &\rightarrow \bar{M} \otimes \mathbf{Q} \\ (r, s) &\mapsto (0, s - r) \end{aligned}$$

(the discrete version of (A.0.5)), and there is an obvious \mathbf{Z} -equivariant map φ from this complex to $\mathbf{Z}[0]$ ($(r, s) \mapsto s$). You will easily check by considering cones that the diagram

$$\begin{array}{ccc} \mathbf{Z} & \xlongequal{\quad} & \mathbf{Z} \\ \gamma^*e_0 \downarrow & & e_1 \downarrow \\ \gamma^*\mathbf{Z}^c[1] & \xrightarrow{\varphi} & \mathbf{Z}[1] \end{array}$$

commutes, where e_1 is the generator of $H^1(\mathbf{Z}, \mathbf{Z}) = Ext_{\mathbf{Z}[\mathbf{Z}]}^1(\mathbf{Z}, \mathbf{Z})$. By adjunction it follows that the diagram

$$\begin{array}{ccc} \mathbf{Z} & \xlongequal{\quad} & \mathbf{Z} \\ e_0 \downarrow & & \tilde{e}_1 \downarrow \\ \mathbf{Z}^c[1] & \xrightarrow{\tilde{\varphi}} & R\gamma_*\mathbf{Z}[1] \end{array}$$

commutes, where $\tilde{\varphi}$ and \tilde{e}_1 denote the maps corresponding to φ and e_1 under adjunction. This implies that $H^1(\hat{\mathbf{Z}}, \mathbf{Z}^c) \xrightarrow{\sim} H^1(\hat{\mathbf{Z}}, R\gamma_*\mathbf{Z})$ under $\tilde{\varphi}$; passing to the open subgroups of $\hat{\mathbf{Z}}$ we get that $\tilde{\varphi}$ induces an isomorphism on the \mathcal{H}^1 s. Since it also obviously induces an isomorphism on the \mathcal{H}^0 s, this proves that it is a quasi-isomorphism.

I hope this is useful.

Best regards,

Bruno

REFERENCES

- [1] Y. André Une introduction aux motifs (motifs purs, motifs mixtes, périodes), Panoramas et Synthèses **17**, SMF, 2004.
- [2] J. Ayoub Les six opérations de Grothendieck et le formalisme des cycles évanescents dans le monde motivique, Astérisque **314**, **315**, SMF, 2007.
- [3] J. Ayoub *L'algèbre de Hopf et le groupe de Galois motivique d'un corps de caractéristique nulle, I*, preprint, 2010.
- [4] J. Ayoub *La réalisation étale et les opérations de Grothendieck*, preprint, 2010.
- [5] A. Beilinson *Higher regulators and values of L-functions*, J. Math. Sciences **30** (1985), 2036–2070. Translated from Itogi Nauki i Tekhniki, Ser. Sovr. Problemy Mat. **24**, (1984), 181–238.
- [6] A. Beilinson *Notes on absolute Hodge cohomology*, Contemp. Math. **55** (I), AMS, 1986, 35–68.

- [7] A. Beilinson *Height pairings between algebraic cycles*, Lect. Notes in Math. **1289**, Springer, 1987, 1–25 or Contemp. Math. **67**, Amer. Math. Soc., 1987, 1–24.
- [8] A. Beilinson, J. Bernstein, P. Deligne *Faisceaux pervers*, Astérisque **100** (1982), 5–171.
- [9] S. Bloch *Algebraic cycles and higher K-theory*, Adv. Math. **61** (1986), 267–304.
- [10] S. Bloch *Lectures on algebraic cycles*, second edition, New Math. Monographs **16**, Cambridge Univ. Press, 2010.
- [11] M. Bökstedt, A. Neeman *Homotopy limits in triangulated categories*, Compositio Math. **86** (1993), 209–234.
- [12] M. Bondarko *Differential graded motives: weight complex, weight filtrations and spectral sequences for realizations; Voevodsky versus Hanamura*, J. Inst. Math. Jussieu **8** (2009), 39–97.
- [13] N. Bourbaki, *Algèbre commutative*, Masson, 1984.
- [14] D.-C. Cisinski, F. Déglise *Local and stable homological algebra in Grothendieck abelian categories*, Homology, Homotopy Appl. **11** (2009), 219–260.
- [15] D.-C. Cisinski, F. Déglise *Triangulated categories of motives*, preprint, 2009.
- [16] J.-L. Colliot-Thélène, J.-J. Sansuc et C. Soulé, *Torsion dans le groupe de Chow de codimension deux*, Duke Math. J. **50** (1983), 763–801.
- [17] J.-L. Colliot-Thélène, T. Szamuely *Autour de la conjecture de Tate à coefficients \mathbf{Z}_ℓ pour les variétés sur les corps finis*, in The geometry of algebraic cycles, Clay Math. Proc. **9**, Amer. Math. Soc., 2010, 83–98.
- [18] F. Déglise *Finite correspondences and transfers over a regular base*, in Algebraic cycles and motives (J. Nagel, C. Peters, eds.), Vol. 1, LMS Lect. Notes Series **343**, Cambridge Univ. Press, 2007, 138–205.
- [19] P. Deligne *La conjecture de Weil, I*, Publ. Math. IHÉS **43** (1974), 5–77.
- [20] A. Dold, D. Puppe *Duality, trace and transfer*, Proc. Int. Conf. in Geometric Topology (Warsaw, 1978), PWN, Warsaw, 1980, 81–102.
- [21] B. Dwork *On the rationality of the zeta function of an algebraic variety*, Amer. J. Math. **82** (1960), 631–648.
- [22] T. Ekedahl *On the adic formalism*, The Grothendieck Festschrift, Vol. II, Birkhäuser, 1990.
- [23] G. Faltings *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983), 349–366.
- [24] G. Faltings, G. Wüstholz, *Rational points*, Vieweg, 1984.
- [25] E. Friedlander, V. Voevodsky *Bivariant cycle cohomology*, in Cycles, transfers and motivic cohomology theories, Ann. of Math. Studies **143**, Princeton Univ. Press, 2000, 137–187.
- [26] O. Gabber *Sur la torsion dans la cohomologie l -adique d'une variété*, C. R. Acad. Sci. Paris **297** (1983), 179–182.
- [27] T. Geisser *Tate's conjecture, algebraic cycles and rational K-theory in characteristic p* , K-Theory **13** (1998), 109–122.
- [28] T. Geisser *Weil-étale cohomology over finite fields*, Math. Ann. **330** (2004), 665–692.
- [29] T. Geisser *Arithmetic cohomology over finite fields and special values of ζ -functions*, Duke Math. J. **133** (2006), 27–57.
- [30] T. Geisser, M. Levine *The Bloch-Kato conjecture and a theorem of Suslin-Voevodsky*, J. Reine Angew. Math. **530** (2001), 55–103.

- [31] M. Hanamura *Mixed motives and algebraic cycles*, I: Math. Res. Lett. **2** (1995), 811–821, II: Invent. Math. **158** (2004), 105–179, III: Math. Res. Lett. **6** (1999), 61–82.
- [32] M. Hanamura *Quasi-DG categories and mixed motivic sheaves*, preprint, 2011.
- [33] A. Huber *Realization of Voevodsky’s motives*, J. Algebraic Geom. **9** (2000), 755–799; Corrigendum: *ibid.*, **13** (2004), 195–207.
- [34] A. Huber, B. Kahn *The slice filtration and mixed Tate motives*, Compositio Math. **142** (2006), 907–936.
- [35] L. Illusie *Catégories dérivées et dualité: travaux de J.-L. Verdier*, Enseign. Math. **36** (1990), 369–391.
- [36] L. Illusie *On Gabbers uniformization theorems: outline and applications to étale cohomology*, Conf. on Ramification and vanishing cycles, Univ. Tokyo, Sept. 11, 2007, <http://www.math.u-psud.fr/~illusie/IllusieTokyo.pdf>.
- [37] F. Ivorra *Réalisation l -adique des motifs triangulés géométriques*, I: Doc. Math. **12** (2007), 607–671; II: Math. Z. **265** (2010), 221–247.
- [38] F. Ivorra *Levine’s motivic comparison theorem revisited*, J. Reine Angew. Math. **617** (2008), 67–107.
- [39] F. Ivorra *Finite dimensional motives and applications following S.-I. Kimura, P. O’Sullivan and others*, Panoramas et synthèses **29**, SMF, 2009.
- [40] F. Ivorra *Triangulated motives over Noetherian separated schemes*, preprint, 2010.
- [41] A. J. de Jong *Smoothness, semi-stability and alterations*, Publ. Math. IHÉS **83** (1996), 51–93.
- [42] U. Jannsen *Continuous étale cohomology*, Math. Ann. **280** (1988), 207–245.
- [43] U. Jannsen, *Mixed motives and algebraic K -theory*, Lect. Notes in Math. **1400**, Springer, 1989.
- [44] U. Jannsen *Motives, numerical equivalence, and semi-simplicity*, Invent. Math. **107** (1992), 447–452.
- [45] U. Jannsen *Motivic sheaves and filtrations on Chow groups*, Proc. Sympos. Pure Math. **55** (1), Amer. Math. Soc., 1994, 245–302.
- [46] B. Kahn *A sheaf-theoretic reformulation of the Tate conjecture*, preprint, 1998, [arXiv:math/9801017](https://arxiv.org/abs/math/9801017).
- [47] B. Kahn *The Geisser-Levine method revisited and algebraic cycles over a finite field*, Math. Ann. **324** (2002), 581–617.
- [48] B. Kahn *Équivalence rationnelle et numérique sur certaines variétés de type abélien sur un corps fini*, Ann. Sci. Éc. Norm. Sup. **36** (2003), 977–1002.
- [49] B. Kahn *On the semi-simplicity of Galois actions*, Rend. Sem. Mat. Univ. Padova **112** (2004), 97–102.
- [50] B. Kahn *Algebraic K -theory, algebraic cycles and arithmetic geometry*, Handbook of K -theory, Springer, 2005.
- [51] B. Kahn *Zeta and L -functions of triangulated motives*, in preparation.
- [52] N. Katz, W. Messing *Some consequences of the Riemann hypothesis for varieties over finite fields*, Invent. Math. **23** (1974), 73–77.
- [53] M. Levine *Mixed motives*, Math. Surveys and Monographs **57**, American Mathematical Society, 1998.
- [54] M. Levine *Six lectures on motives*, Panoramas et synthèses **29**, SMF, 2009.

- [55] S. Lichtenbaum *The Weil-étale topology on schemes over finite fields*, Compos. Math. **141** (2005), 689–702.
- [56] S. Mac Lane *Categories for the working mathematician* (2nd ed.), Springer, 1998.
- [57] C. Mazza, V. Voevodsky, C. Weibel, *Lecture notes on motivic cohomology*, Clay Mathematics Monographs **2**, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2006.
- [58] J. Milne *Values of zeta functions of varieties over finite fields*, Amer. J. Math. **108** (1986), 297–360.
- [59] J. Milne *Motives over finite fields*, in *Motives* (Jannsen, Kleiman, Serre, eds), Proc. Symp. Pure Math. **55** (I), AMS, 1994.
- [60] J.S. Milne *The Tate conjecture for certain abelian varieties over finite fields*, Acta Arithm. **100** (2001), 135–166.
- [61] J. Milne *The Tate conjecture over finite fields*, AIM talk, 2007, available on www.jmilne.org.
- [62] F. Morel *On the motivic π_0 of the sphere spectrum*, in *Axiomatic, enriched and motivic homotopy theory* (J.P.C. Greenlees, ed.), NATO Science Series II, **131**, Kluwer, 2004, 219–260.
- [63] F. Morel, V. Voevodsky *\mathbf{A}^1 -homotopy theory of schemes*, Publ. Math. IHÉS **90** (1999), 45–143.
- [64] A. Neeman *Triangulated categories*, Ann. Math. Studies **148**, Princeton University Press, 2001.
- [65] A. Pirutka *Sur le groupe de Chow de codimension deux des variétés sur les corps finis*, to appear in *Algebra and Number Theory*.
- [66] J. Riou *Realization functors*, Panoramas et synthèses **29**, SMF, 2009.
- [67] J.-P. Serre *Corps locaux* (2nd ed.), Hermann, 1968.
- [68] J.-P. Serre *Cohomologie galoisienne* (5th ed.), Lect. Notes in Math. **5**, Springer, 1994.
- [69] C. Soulé *Groupes de Chow et K -théorie des variétés sur les corps finis*, Math. Ann. **268** (1984), 317–345.
- [70] C. Soulé *Opérations en K -théorie algébrique*, Canad. J. Math. **37** (1985), 488–550.
- [71] M. Spiess *Proof of the Tate conjecture for products of elliptic curves over finite fields*, Math. Ann. **314** (1999), 285–290.
- [72] A. Suslin, V. Voevodsky *Singular homology of abstract algebraic varieties*, Invent. Math. **123** (1996), 61–94.
- [73] A. Suslin, V. Voevodsky *Relative cycles and Chow sheaves*, in *Cycles, transfers and motivic cohomology theories*, Ann. of Math. Studies **143**, Princeton Univ. Press, 2000, 10–86.
- [74] J. Tate *Algebraic cycles and poles of zeta functions*, *Arithmetical Algebraic Geometry* (Proc. Conf. Purdue Univ., 1963), Harper & Row, 1965, 93–110.
- [75] J. Tate *Endomorphisms of abelian varieties over finite fields*, Invent. Math. **2** (1966), 134–144.
- [76] J. Tate *Conjectures on algebraic cycles in ℓ -adic cohomology*, in *Motives* (Jannsen, Kleiman, Serre, eds), Proc. Symp. Pure Math. **55** (I), AMS, 1994, 71–83.
- [77] V. Voevodsky *\mathbf{A}^1 -homotopy theory*, Proc. Intern. Congress of Math., Vol. I (Berlin, 1998), Doc. Math. 1998, Extra Vol. I, 579–604.

- [78] V. Voevodsky *Triangulated categories of sheaves over a field*, in *Cycles, transfers and motivic cohomology theories*, Ann. of Math. Studies **143**, Princeton Univ. Press, 2000, 188–238.
- [79] V. Voevodsky *Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic*, IMRN **2002**, 351–355.
- [80] V. Voevodsky *Motivic cohomology with $\mathbf{Z}/2$ coefficients*, Publ. Math. IHÉS **98** (2003), 59–104.
- [81] V. Voevodsky *Motives over simplicial schemes*, J. K-Theory **5** (2010), 1–38.
- [82] V. Voevodsky *Cancellation theorem*, Doc. Math. **2010**, Extra volume: Andrei A. Suslin sixtieth birthday, 671–685.
- [83] Y. G. Zarhin *Isogenies of abelian varieties over fields of finite characteristics*, Math. USSR-Sbornik **24** (1974), 451–461.
- [SGA4] Séminaire de géométrie algébrique du Bois-Marie: Théorie des topos et cohomologie étale des schémas, Vol. 3, Lect. Notes in Math. **305**, Springer, 1973.
- [SGA5] Séminaire de géométrie algébrique du Bois-Marie: Cohomologie l -adique et fonctions L , Lect. Notes in Math. **589**, Springer, 1977.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, CASE 247, 4 PLACE JUSSIEU, 75252
PARIS CEDEX 05, FRANCE

E-mail address: kahn@math.jussieu.fr