

Liftings of Reduction Maps for Quaternion Algebras

Christophe Cornut and Dimitar Jetchev

ABSTRACT

We construct liftings of reduction maps from CM points to supersingular points for general quaternion algebras and use these liftings to establish a precise correspondence between CM points on indefinite quaternion algebras with a given conductor and CM points on certain corresponding totally definite quaternion algebras.

1. Introduction

Let F be a totally real number field and let B be a quaternion algebra over F . Let $G = \text{Res}_{F/\mathbb{Q}} B^\times$ be the algebraic group over \mathbb{Q} associated to B^\times . Let \mathbb{A}_f be the finite adèles of \mathbb{Q} .

For an open subgroup $H \subset G(\mathbb{A}_f)$ that is compact modulo the center let

$$\mathcal{X}(G, H) = G(\mathbb{Q}) \backslash G(\mathbb{A}_f) / H.$$

This is a finite set. In the case when $H = \widehat{R}^\times$ for some \mathcal{O}_F -order R in B , the map $\widehat{b} \mapsto (\widehat{b} \cdot \widehat{R}) \cap B$ identifies $\mathcal{X}(G, H)$ with the set of B^\times -homothety classes of locally principal fractional right R -ideals in B . When B is not totally definite, the reduced norm $\text{nr}: B \rightarrow F$ induces a bijection

$$\mathcal{X}(G, H) \xrightarrow{\cong} \text{nr}(B^\times) \backslash \widehat{F}^\times / \text{nr}(H)$$

by the strong approximation theorem [Vig80, p.81]. Moreover, the norm theorem [Vig80, Thm.III.4.1] implies that $\text{nr}(B^\times)$ is precisely the subgroup of elements $\lambda \in F^\times$ such that $\lambda_v > 0$ for all places $v \mid \infty$ of F where B is not split, i.e., where $B_v \not\cong M_2(F_v)$. On the other hand, if B is totally definite, then $\mathcal{X}(G, H)$ is a genuinely non-commutative object, which shows up quite often in low dimensional arithmetic geometry (see, e.g., [Gro87] and [Rib90]).

Next, let K be a totally imaginary quadratic extension of F that embeds into B . Fix an embedding $K \hookrightarrow B$. If $T = \text{Res}_{K/\mathbb{Q}} K^\times$ is the associated rational torus, we get a corresponding embedding $T \hookrightarrow G$. Define

$$\mathcal{CM}(G, H) = T(\mathbb{Q}) \backslash G(\mathbb{A}_f) / H.$$

This is now an infinite set with an obvious projection map

$$\pi: \mathcal{CM}(G, H) \rightarrow \mathcal{X}(G, H) \tag{1}$$

In addition, it is equipped with a left action of $T(\mathbb{A}_f)$ with finite orbits. Using Artin's reciprocity map $\text{rec}_K: T(\mathbb{A}_f) = \widehat{K}^\times \rightarrow \text{Gal}_K^{ab}$ whose kernel equals the closure of K^\times in \widehat{K}^\times , we can also view this action as a continuous action of Gal_K^{ab} . We shall thus refer to it as the *Galois action* on $\mathcal{CM}(G, H)$.

As suggested by the notation, such sets are most frequently occurring as sets of complex multiplication points, or special points, on certain Shimura varieties. Assume for instance that B is split at a single place $v \mid \infty$ of F , say $v = v_\iota$ corresponding to an embedding $\iota: F \rightarrow \mathbb{R}$. Let X be the

Shimura curve of level H attached to B . It is an algebraic curve over the reflex field ιF , and

$$\mathcal{CM}(G, H) \simeq G(\mathbb{Q}) \backslash (G(\mathbb{A}_f)/H \times G(\mathbb{Q}) \cdot \tau) \subset X(\mathbb{C}) = G(\mathbb{Q}) \backslash (G(\mathbb{A}_f)/H \times (\mathbb{C} - \mathbb{R}))$$

is the set of special points with complex multiplication by K in X . In this formula, we have fixed an isomorphism $B \otimes_{F, \iota} \mathbb{R} \simeq M_2(\mathbb{R})$ to define the action of $G(\mathbb{Q})$ on $\mathbb{C} - \mathbb{R}$, and τ is the unique point on the upper half-plane whose stabilizer is $K^\times = T(\mathbb{Q})$ under this isomorphism. These special points are defined over the maximal abelian extension of ιK in \mathbb{C} , and the above Galois action is the natural one (this follows from the definition of Shimura varieties). In this setting, the projection map is almost the projection to the set of connected components of X , namely

$$\mathcal{CM}(G, H) \hookrightarrow X(\mathbb{C}) \rightarrow \pi_0(X(\mathbb{C})) = G(\mathbb{Q}) \backslash (G(\mathbb{A}_f)/H \times \{\pm 1\}) = G(\mathbb{Q})^+ \backslash G(\mathbb{A}_f)/H.$$

Here $G(\mathbb{Q})$ acts on $\{\pm 1\}$ by the sign of $\iota \circ \det: G(\mathbb{Q}) \rightarrow \mathbb{R}^\times$, and $G(\mathbb{Q})^+$ is the kernel of this action.

Reduction maps. There are similar but more interesting maps to consider. For instance, let v be a finite place of F such that $B_v \simeq M_2(F_v)$. Then our Shimura curve has a natural model over the corresponding local ring, the special fiber of which contains a distinguished finite set of supersingular points and this finite set of points may be identified with a set $\mathcal{X}(G', H')$ as above, where G' is the algebraic group over \mathbb{Q} associated to the totally definite quaternion algebra B' over F which is obtained from B by changing the invariants at v_i and v (see, e.g., [DR73] and [KM85] for $B = M_2(\mathbb{Q})$, and [Car86] for the remaining cases). If we choose an extension of v to the maximal abelian extension of ιK in \mathbb{C} then each CM point extends to the corresponding local ring. Moreover, if v does not split in K , these extended points reduce to supersingular points on the special fiber and one obtains a reduction map

$$\text{red}: \mathcal{CM}(G, H) \rightarrow \mathcal{X}(G', H') \tag{2}$$

that is described in [Cor02] and Section 4 if $B = M_2(\mathbb{Q})$ and in [CV05] for the remaining cases. On the other hand, if B is ramified at v , there is a similar description for the reduction map with values in the set of irreducible components of the special fiber of a suitable model of X via Ribet's theory of bimodules. We refer to [Mol10b] for a survey of such maps, and to [Mol10a] for applications to the determination of explicit equations for certain hyperelliptic Shimura curves.

Given such a reduction map, it is expected that the Galois orbits in $\mathcal{CM}(G, H)$ tend to be equidistributed among the finitely many fibers of the reduction map red . Such equidistribution results are already known in various cases [Mic04], [CV05], [HM06], [JK10], [Mol10b] and were crucial in the proof of Mazur's non-vanishing conjecture by the first author and Vatsal [Cor02], [Vat02], [CV07].

Our contributions. We propose a simple strategy to reduce the study of the arithmetically interesting reduction maps (2) to that of the more straightforward projections (1). In all cases, there is indeed a natural Gal_K -equivariant map

$$\theta: \mathcal{CM}(G, H) \rightarrow \mathcal{CM}(G', H')$$

such that $\text{red} = \pi \circ \theta$. Thus, for any Gal_K -orbit Γ in $\mathcal{CM}(G, H)$ and any point $s \in \mathcal{X}(G', H')$, we obtain a κ -to-1 surjective map

$$\theta: \Gamma \cap \text{red}^{-1}(s) \rightarrow \Gamma' \cap \pi^{-1}(s)$$

where $\Gamma' = \theta(\Gamma)$ is a Gal_K -orbit in $\mathcal{CM}(G', H')$ and $\kappa = \kappa(\Gamma) = |\Gamma| / |\Gamma'|$.

This paper essentially implements this strategy when $H = \widehat{R}^\times$ for some Eichler \mathcal{O}_F -order R in B . The algebraic description of θ (and also of $\text{red} = \pi \circ \theta$) is given in Section 3.2 in a more general setting following the conventions of [CV05]. The size of the Galois orbits (and thus also the constant

κ above) is controlled by a simple invariant that is defined in Section 3.3 together with a refinement: these are the coarse and fine conductors \mathfrak{c}_g and \mathfrak{c}_f , respectively. The number of Galois orbits with a prescribed fine conductor is also given there using elementary local computations that are carried on in Section 2. Our main result, Theorem 3.1, then describes the restriction of θ to the fibers of the fine conductor map. If we furthermore restrict θ to a given fiber of red and π , we obtain an explicit correspondence between certain sets of CM points on two distinct quaternion algebras, a special case of which is used in [JK10] and thoroughly explained in the final section. Another case of our main theorem is used in [Mol10a] and [Mol10b] to compute explicit equations for Shimura curves with no cusps.

2. Review of quadratic orders

Only for this particular section, \mathcal{O}_F will be a Dedekind domain with fraction field F . Let K be a semi-simple commutative F -algebra of dimension 2, i.e., $K \simeq F \times F$ or K is a quadratic field extension of F . Let \mathcal{O}_K be the integral closure of \mathcal{O}_F in K . The map which sends \mathfrak{c} to $\mathcal{O}_{\mathfrak{c}} = \mathcal{O}_F + \mathfrak{c}\mathcal{O}_K$ is a bijection from the set of all non-zero ideals $\mathfrak{c} \subset \mathcal{O}_F$ onto the set of all \mathcal{O}_F -orders in K . It is well-known that all such orders are *Gorenstein* rings. We refer to \mathfrak{c} as the conductor of $\mathcal{O} = \mathcal{O}_{\mathfrak{c}}$.

2.1 Quadratic lattices

Fix a free, rank one K -module V . Let \mathcal{L} be the set of all full \mathcal{O}_F -lattices in V . The *conductor* of a lattice $\Lambda \in \mathcal{L}$ is the conductor $c(\Lambda)$ of the \mathcal{O}_F -order $\mathcal{O}(\Lambda) = \{\lambda \in K : \lambda\Lambda \subset \Lambda\} = \mathcal{O}_{c(\Lambda)}$. It follows from [Bas63, Prop.7.2] that Λ is a projective rank one $\mathcal{O}_{c(\Lambda)}$ -module. Let $[\Lambda]$ be its isomorphism class in the Picard group $\text{Pic}(\mathcal{O}_{c(\Lambda)})$. Since any two \mathcal{O}_F -lattices in V are K^\times -homothetic precisely when they have the same conductor and define the same class in the relevant Picard group, the map $\Lambda \mapsto [\Lambda]$ induces a bijection

$$K^\times \backslash \mathcal{L} \simeq \bigsqcup_{\mathfrak{c}} \text{Pic}(\mathcal{O}_{\mathfrak{c}}). \quad (3)$$

When \mathcal{O}_F is a local ring with maximal ideal \mathfrak{p}_F , $\text{Pic}(\mathcal{O}_{\mathfrak{p}_F^n}) = \{1\}$ for all n , so the above bijection becomes

$$K^\times \backslash \mathcal{L} \simeq \mathbb{N}, \quad K^\times \Lambda \mapsto n(\Lambda), \quad (4)$$

where $n(\Lambda)$ is the positive integer for which $c(\Lambda) = \mathfrak{p}_F^{n(\Lambda)}$.

2.2 The action of K^\times on $\mathcal{L} \times \mathcal{L}$

Assume that \mathcal{O}_F is a discrete valuation ring. We will describe the orbits of K^\times acting diagonally on $\mathcal{L} \times \mathcal{L}$.

A K^\times -invariant of \mathcal{L}^2 . There is an invariant $K^\times \backslash \mathcal{L}^2 \rightarrow (K^\times \backslash \mathcal{L})^2 \simeq \mathbb{N}^2$, given by

$$x = (\Lambda', \Lambda'') \mapsto \underline{n}(x) = (n(\Lambda'), n(\Lambda'')). \quad (5)$$

There is another invariant $K^\times \backslash \mathcal{L}^2 \rightarrow \text{GL}_F(V) \backslash \mathcal{L}^2 \simeq \mathfrak{S}_2 \backslash \mathbb{Z}^2$ that describes the relative position of two lattices. It maps $x = (\Lambda', \Lambda'')$ to the unique pair of integers $\text{inv}(x) = \{i_1, i_2\} \in \mathfrak{S}_2 \backslash \mathbb{Z}^2$ for which there exists an F -basis (e_1, e_2) of V such that

$$\Lambda' = \mathcal{O}_F e_1 \oplus \mathcal{O}_F e_2 \quad \text{and} \quad \Lambda'' = \mathfrak{p}_F^{i_1} e_1 \oplus \mathfrak{p}_F^{i_2} e_2. \quad (6)$$

The distance function. This latter invariant is related to the distance function on the set of vertices $\mathcal{V} = F^\times \backslash \mathcal{L}$ of the Bruhat-Tits tree of $\text{PGL}_F(V)$: if v' and v'' are the images of Λ' and Λ'' in \mathcal{V} , then $\text{dist}(v', v'') = |i_1 - i_2|$. Since $K^\times \backslash \mathcal{L} \cong \mathbb{N}$, also $K^\times \backslash \mathcal{V} \cong \mathbb{N}$. In other words, the function n

on \mathcal{L} descends to a function $n: \mathcal{V} \rightarrow \mathbb{N}$ whose fibers $\mathcal{V}(k) = \{v \in \mathcal{V}: n(v) = k\}$ are precisely the K^\times -orbits in \mathcal{V} . We claim that also $\mathcal{V}(k) = \{v \in \mathcal{V}: \text{dist}(v, \mathcal{V}(0)) = k\}$ for all $k \in \mathbb{N}$. To prove this, first note that $\mathcal{V}(0)$ is a convex subset of \mathcal{V} , namely a single vertex if K is an unramified extension of F , a pair of adjacent vertices if K is a ramified extension of F , and a line in the building (i.e. an appartement) if $K \cong F \times F$.

Now, let v be any vertex in $\mathcal{V}(k)$. Then v is represented by $\Lambda = \mathcal{O}_{\mathfrak{p}_F^k} e$ for some K -basis e of V . For $0 \leq j \leq k$, let $v_j \in \mathcal{V}$ be the F^\times -homothety class of $\Lambda_j = \mathcal{O}_{\mathfrak{p}_F^j} e \in \mathcal{L}$. Then $v_j \in \mathcal{V}(j)$ and $(v_k, v_{k-1}, \dots, v_0)$ is a path of length k from $v = v_k$ to the vertex v_0 of $\mathcal{V}(0)$. Therefore $\text{dist}(v, \mathcal{V}(0)) = j$ for some $0 \leq j \leq k$, and the convexity of $\mathcal{V}(0)$ implies that the j th term in our path, namely v_{k-j} , should already be in $\mathcal{V}(0)$. Therefore $k - j = 0$ and $\text{dist}(v, \mathcal{V}(0)) = k$.

Counting the K^\times -orbits. Finally, fix $(n', n'') \in \mathbb{N} \times \mathbb{N}$, $(i_1, i_2) \in \mathbb{Z} \times \mathbb{Z}$ and let $\delta = |i_1 - i_2|$. It follows from the above considerations that the projection $\mathcal{L} \rightarrow \mathcal{V}$ induces a bijection between

$$\mathcal{L}(n', n''; i_1, i_2) = \{x \in K^\times \backslash \mathcal{L}^2 \text{ s.t. } \underline{n}(x) = (n', n'') \text{ and } \text{inv}(x) = \{i_1, i_2\}\}$$

and the set of K^\times -orbits of pairs $(v', v'') \in \mathcal{V} \times \mathcal{V}$ such that

$$\text{dist}(v', \mathcal{V}(0)) = n', \quad \text{dist}(v'', \mathcal{V}(0)) = n'' \quad \text{and} \quad \text{dist}(v', v'') = \delta.$$

If for instance $n' \geq n''$, the choice of a vertex $v' \in \mathcal{V}(n')$ identifies the latter set of K^\times -orbits with the set of all vertices $v'' \in \mathcal{V}(n'')$ at distance δ from v' . Using then the above description of $\mathcal{V}(0)$, it is a simple combinatorial exercise to prove the following:

LEMMA 2.1. *If $\mathcal{O}_F/\mathfrak{p}_F$ is finite of order q , then $\mathcal{L}(n', n''; i_1, i_2)$ is finite of order*

$$N(n', n'', \delta) = |\mathcal{L}(n', n''; i_1, i_2)| \quad \text{with } \delta = |i_1 - i_2|.$$

Moreover $N(n', n'', \delta) = 0$ unless one of the following conditions holds:

i) $\delta = |n' - n''| + 2r$ for some $0 \leq r < \min(n', n'')$. Then

$$N(n', n'', \delta) = \begin{cases} 1 & \text{if } r = 0, \\ (q-1)q^{r-1} & \text{if } r > 0 \end{cases}$$

ii) K is an unramified extension of F and $\delta = n' + n''$. Then

$$N(n', n'', \delta) = q^{\min(n', n'')}.$$

iii) K is a ramified extension of F and $\delta = n' + n'' + s$ with $s \in \{0, 1\}$. Then

$$N(n', n'', \delta) = \begin{cases} q^{\min(n', n'')} & \text{if } s = 1 \text{ or } \min(n', n'') = 0, \\ (q-1)q^{\min(n', n'')-1} & \text{if } s = 0 < \min(n', n''). \end{cases}$$

iv) $K \simeq F \times F$ and $\delta = n' + n'' + s$ with $s \in \mathbb{N}$. Then

$$N(n', n'', \delta) = \begin{cases} 1 & \text{if } \min(n', n'') = 0 = s, \\ 2 & \text{if } \min(n', n'') = 0 < s, \\ (q-2)q^{\min(n', n'')-1} & \text{if } \min(n', n'') > 0 = s, \\ 2(q-1)q^{\min(n', n'')-1} & \text{if } \min(n', n'', s) > 0. \end{cases}$$

3. Adelic constructions of the lifting and the reduction maps

3.1 Notation and preliminaries

We now switch back to the notation in the introduction, where F is a totally real number field, B is a quaternion algebra over F , K is a totally imaginary quadratic extension of F and $\iota: K \hookrightarrow B$ is

an embedding. We denote by $\text{Ram}_f B$ the set of all finite places of F where B does not split. Let S be a finite set of finite places of F that satisfy the following hypotheses:

- H.1.** B is unramified at every $v \in S$, i.e., $\text{Ram}_f B \cap S = \emptyset$,
- H.2.** $|S| + |\text{Ram}_f(B)| + [F : \mathbb{Q}]$ is even,
- H.3.** Every $v \in S$ is either inert or ramified in K .

Hypotheses H.1. and H.2. imply that there exists a unique (up to isomorphism) totally definite quaternion algebra B_S over F ramified exactly at the finite places $\text{Ram}_f(B) \cup S$. Hypothesis H.3. implies that K embeds into B_S . We fix once and for all such an embedding $\iota_S : K \hookrightarrow B_S$. We let

$$G = \text{Res}_{F/\mathbb{Q}} B^\times, \quad G_S = \text{Res}_{F/\mathbb{Q}} B_S^\times \quad \text{and} \quad T = \text{Res}_{F/\mathbb{Q}} K^\times.$$

3.2 Adelic construction of the map θ_S

Following Cornut and Vatsal (see [CV05, §2.1]), we consider the following locally compact and totally discontinuous groups:

$$\left\{ \begin{array}{l} T(\mathbb{A}_f) = (K \otimes \mathbb{A}_f)^\times = \prod' K_v^\times \\ G(\mathbb{A}_f) = (B \otimes \mathbb{A}_f)^\times = \prod' B_v^\times \\ G_S(\mathbb{A}_f) = (B_S \otimes \mathbb{A}_f)^\times = \prod' B_{S,v}^\times \end{array} \right. \quad \text{and} \quad G(S) = \left(\prod'_{v \notin S} B_{S,v}^\times \right) \times \prod_{v \in S} F_v^\times.$$

The restricted products on the left are the usual ones defined by the arbitrary choice of an integral structure on the algebraic groups, whereas the restricted product on the right is induced by that of $G_S(\mathbb{A}_f)$. These topological groups together fit in a commutative diagram

$$\begin{array}{ccc} & G(\mathbb{A}_f) & \\ \iota \nearrow & & \searrow \phi_S \\ T(\mathbb{A}_f) & & G(S) \\ \iota_S \searrow & & \nearrow \pi_S \\ & G_S(\mathbb{A}_f) & \end{array}$$

where ι and ι_S are the continuous embeddings induced by their algebraic counterparts, where π_S is the continuous, open and surjective morphism

$$G_S(\mathbb{A}_f) = \prod'_{v \notin S} B_{S,v}^\times \times \prod_{v \in S} B_{S,v}^\times \twoheadrightarrow \prod'_{v \notin S} B_{S,v}^\times \times \prod_{v \in S} F_v^\times = G(S)$$

which is induced by the identity on $B_{S,v}$ for $v \notin S$ and by the reduced norm $\text{nr}_{S,v} : B_{S,v}^\times \rightarrow F_v^\times$ for $v \in S$. Finally, the continuous, open and surjective morphism

$$\phi_S : G(\mathbb{A}_f) = \prod'_{v \notin S} B_v^\times \times \prod_{v \in S} B_v^\times \twoheadrightarrow \prod'_{v \notin S} B_{S,v}^\times \times \prod_{v \in S} F_v^\times = G(S)$$

is again induced by the reduced norm $\text{nr}_v : B_v^\times \rightarrow F_v^\times$ for $v \in S$, and by well-chosen isomorphisms $\theta_{S,v} : B_v \xrightarrow{\cong} B_{S,v}$ for $v \notin S$ whose choice is carefully explained in [CV05, §2.1.3].

Topological properties and Galois action. We use the above commutative diagram to let $T(\mathbb{A}_f)$ act on $G(\mathbb{A}_f)$, $G_S(\mathbb{A}_f)$ and $G(S)$ by multiplication on the left. Let $\overline{T(\mathbb{Q})}$ be the closure of $T(\mathbb{Q})$ in $T(\mathbb{A}_f)$, and recall that class field theory yields an isomorphism of topological groups

$$\text{Art}_K : T(\mathbb{A}_f) / \overline{T(\mathbb{Q})} \xrightarrow{\cong} \text{Gal}_K^{ab}.$$

We thus obtain continuous Gal_K^{ab} -equivariant maps of topological Gal_K^{ab} -sets

$$\overline{T(\mathbb{Q})} \backslash G(\mathbb{A}_f) \xrightarrow{\phi_S} \overline{T(\mathbb{Q})} \backslash G(S) \xleftarrow{\pi_S} \overline{T(\mathbb{Q})} \backslash G_S(\mathbb{A}_f).$$

These maps are also respectively equivariant for the right actions of $G(\mathbb{A}_f)$ and $G_S(\mathbb{A}_f)$. For a compact open subgroup H of $G(\mathbb{A}_f)$, define $H(S) = \phi_S(H)$ and $H_S = \pi_S^{-1}(H(S))$. We now have Gal_K^{ab} -equivariant maps of discrete Gal_K^{ab} -sets

$$\overline{T(\mathbb{Q})} \backslash G(\mathbb{A}_f) / H \xrightarrow{\phi_S} \overline{T(\mathbb{Q})} \backslash G(S) / H(S) \xleftarrow{\pi_S} \overline{T(\mathbb{Q})} \backslash G_S(\mathbb{A}_f) / H_S$$

and the above π_S is a bijection by construction of H_S . Since

$$\mathcal{CM}(G, H) = T(\mathbb{Q}) \backslash G(\mathbb{A}_f) / H = \overline{T(\mathbb{Q})} \backslash G(\mathbb{A}_f) / H$$

and similarly for G_S , we obtain a Gal_K^{ab} -equivariant map of discrete Gal_K^{ab} -sets

$$\theta_S = \pi_S^{-1} \circ \phi_S : \mathcal{CM}(G, H) \rightarrow \mathcal{CM}(G_S, H_S).$$

If $H = \prod_v H_v$ in $G(\mathbb{A}_f) = \prod'_v B_v^\times$, then $H_S = \prod_v H_{S,v}$ in $G_S(\mathbb{A}_f) = \prod'_v B_{S,v}^\times$ with

$$H_{S,v} = \phi_v(H_v) \text{ for } v \notin S \quad \text{and} \quad H_{S,v} = \text{nr}_{S,v}^{-1}(\text{nr}_v(H_v)) \text{ for } v \in S.$$

In this case, the map induced by θ_S on the Gal_K^{ab} -orbit spaces,

$$\bar{\theta}_S : \text{Gal}_K^{ab} \backslash \mathcal{CM}(G, H) \rightarrow \text{Gal}_K^{ab} \backslash \mathcal{CM}(G_S, H_S) \quad (7)$$

has a purely local description, namely

$$\bar{\theta}_S = (\bar{\theta}_{S,v}) : \prod'_v K_v^\times \backslash B_v^\times / H_v \rightarrow \prod'_v K_v^\times \backslash B_{S,v}^\times / H_{S,v} \quad (8)$$

where $\bar{\theta}_{S,v}$ is the bijection induced by $\theta_{S,v} : B_v \xrightarrow{\simeq} B_{S,v}$ for $v \notin S$, and equals

$$K_v^\times \backslash B_v^\times / H_v \xrightarrow{\text{nr}_v} \text{nr}_v(K_v^\times) \backslash F_v^\times / \text{nr}_v(H_v) \xrightarrow{\text{nr}_{S,v}^{-1}} K_v^\times \backslash B_{S,v}^\times / H_{S,v}$$

for $v \in S$.

The construction of the map θ_S also gives us a reduction map

$$\text{red}_S : \mathcal{CM}(G, H) \rightarrow \mathcal{X}(G_S, H_S) \quad (9)$$

defined by $\text{red}_S := \pi \circ \theta_S$, where $\pi : \mathcal{CM}(G_S, H_S) \rightarrow \mathcal{X}(G_S, H_S)$ is the natural projection map.

3.3 Fine and coarse conductors

We now specialize the above constructions to the case $H = \widehat{R}^\times$ for some Eichler \mathcal{O}_F -order R in B that will be fixed throughout. The level of R is a nonzero integral ideal of \mathcal{O}_F that we denote by \mathfrak{n} . We also choose two maximal \mathcal{O}_F -orders R' and R'' in B such that $R = R' \cap R''$. Then $H_S = \widehat{R}_S^\times$ where R_S is an Eichler \mathcal{O}_F -order in B_S whose level \mathfrak{n}_S is the prime-to- S -part of \mathfrak{n} . We also obtain two maximal \mathcal{O}_F -orders R'_S and R''_S in B_S such that $R_S = R'_S \cap R''_S$. For a finite place v of F , $R_{S,v} = \theta_{S,v}(R_v)$ if $v \notin S$ while $R_{S,v}$ is the unique maximal order of $B_{S,v}$ if $v \in S$. Explicitly,

$$R_S = \left\{ b \in B_S : \forall v \notin S, \theta_{S,v}^{-1}(b) \in R_v \text{ and } \forall v \in S, \text{nr}_{S,v}(b) \in \mathcal{O}_{F,v} \right\}.$$

We have a similar explicit description for R'_S and R''_S .

Definitions. For any finite set of places T of F , let $\mathcal{I}(T)$ be the monoid of integral ideals of \mathcal{O}_F that are coprime to the places of T . The *fine conductor* is a Gal_K^{ab} -invariant map

$$\mathfrak{c}_f : \mathcal{CM}(G, H) \rightarrow \mathcal{I}(\text{Ram}_f B) \times \mathcal{I}(\text{Ram}_f B) \quad (10)$$

that is defined as follows: given a CM point $x = [g]$ in $\mathcal{CM}(G, H) = T(\mathbb{Q}) \backslash G(\mathbb{A}_f) / H$, consider the images x' and x'' of x in $T(\mathbb{Q}) \backslash G(\mathbb{A}_f) / \widehat{R}'^\times$ and $T(\mathbb{Q}) \backslash G(\mathbb{A}_f) / \widehat{R}''^\times$, respectively. The intersection $K \cap g \widehat{R}' g^{-1}$ is an \mathcal{O}_F -order in K that depends only on the Gal_K^{ab} -orbit of x and whose conductor

REDUCTION MAPS

$\mathfrak{c}(x')$ is an \mathcal{O}_F -ideal that is prime to $\text{Ram}_f B$. Similarly, we obtain an integral ideal $\mathfrak{c}(x'') \subset \mathcal{O}_F$ for R'' . One then defines $\mathfrak{c}_f(x) := (\mathfrak{c}(x'), \mathfrak{c}(x''))$. The *coarse conductor map*

$$\mathfrak{c}_g: \mathcal{CM}(G, H) \rightarrow \mathcal{I}(\text{Ram}_f B) \quad (11)$$

is defined as $\mathfrak{c}_g(x) = \mathfrak{c}(x') \cap \mathfrak{c}(x'') \subset \mathcal{O}_F$. The stabilizer of x in $T(\mathbb{A}_f) = \widehat{K}^\times$ is then

$$\text{Stab}_{T(\mathbb{A}_f)}(x) = K^\times \cdot \left(\widehat{K}^\times \cap g \widehat{R}^\times g^{-1} \right) = K^\times \widehat{\mathcal{O}_{\mathfrak{c}_g(x)}}^\times,$$

and the field $K(x)$ that is fixed by the stabilizer of x in Gal_K^{ab} is the ring class field $K[\mathfrak{c}_g(x)]$ of conductor $\mathfrak{c}_g(x)$, i.e., the abelian extension of K fixed by $\text{rec}_K(K^\times \widehat{\mathcal{O}_{\mathfrak{c}_g(x)}}^\times)$.

Similarly, we define the fine and coarse conductors

$$\mathfrak{c}_{S,f}: \mathcal{CM}(G_S, H_S) \rightarrow \mathcal{I}(\text{Ram}_f B_S)^2 \quad \text{and} \quad \mathfrak{c}_{S,g}: \mathcal{CM}(G_S, H_S) \rightarrow \mathcal{I}(\text{Ram}_f B_S).$$

Note that if $y = \theta_S(x)$ then $\mathfrak{c}(y)$ is the prime-to- S part of $\mathfrak{c}(x)$. We thus have a commutative diagram

$$\begin{array}{ccccc} \mathcal{CM}(G, H) & \twoheadrightarrow & \text{Gal}_K^{\text{ab}} \backslash \mathcal{CM}(G, H) & \xrightarrow{\mathfrak{c}_f} & \mathcal{I}(\text{Ram}_f B)^2 \\ \downarrow \theta_S & & \downarrow \bar{\theta}_S & & \downarrow ()_S \\ \mathcal{CM}(G_S, H_S) & \twoheadrightarrow & \text{Gal}_K^{\text{ab}} \backslash \mathcal{CM}(G_S, H_S) & \xrightarrow{\mathfrak{c}_{S,f}} & \mathcal{I}(\text{Ram}_f B_S)^2, \end{array} \quad (12)$$

where the map $()_S$ sends $I \in \mathcal{I}(\text{Ram}_f B)$ to its prime-to- S part $I_S \in \mathcal{I}(\text{Ram}_f B_S)$.

Local analysis. The right-hand square of this diagram can be analyzed by purely local means: it is the restricted product over all finite primes v of F of one of the following diagrams:

Case 1: $v \notin \text{Ram}_f B \cup S$.

$$\begin{array}{ccc} K_v^\times \backslash B_v^\times / R_v^\times & \xrightarrow{n_v} & \mathbb{N} \times \mathbb{N} \\ \downarrow \theta_{S,v} & & \downarrow \text{id} \\ K_v^\times \backslash B_{S,v}^\times / R_{S,v}^\times & \xrightarrow{n_{S,v}} & \mathbb{N} \times \mathbb{N} \end{array}$$

Case 2: $v \in S$.

$$\begin{array}{ccc} K_v^\times \backslash B_v^\times / R_v^\times & \xrightarrow{n_v} & \mathbb{N} \times \mathbb{N} \\ \downarrow \theta_{S,v} & & \downarrow 0 \\ K_v^\times \backslash B_{S,v}^\times / R_{S,v}^\times & \longrightarrow & 0 \end{array}$$

Case 3: $v \in \text{Ram}_f B$.

$$\begin{array}{ccc} K_v^\times \backslash B_v^\times / R_v^\times & \xrightarrow{n_v} & 0 \\ \downarrow \theta_{S,v} & & \downarrow 0 \\ K_v^\times \backslash B_{S,v}^\times / R_{S,v}^\times & \longrightarrow & 0 \end{array}$$

Here, for $\star \in \{\emptyset, S\}$, the map $n_{\star,v}$ sends $K_v^\times g_v R_{\star,v}^\times$ to the pair of integers (n', n'') such that $K_v \cap g_v R'_{\star,v} g_v^{-1}$ (resp. $K_v \cap g_v R''_{\star,v} g_v^{-1}$) is the order of conductor $\mathfrak{p}_v^{n'}$ (resp. $\mathfrak{p}_v^{n''}$) in \mathcal{O}_{K_v} , where \mathfrak{p}_v is the maximal ideal in \mathcal{O}_{F_v} .

These maps are related to the one defined in (5) as follows: fix a simple left $B_{\star,v}$ -module $V_{\star,v}$, let \mathcal{L}_v be the set of all \mathcal{O}_{F_v} -lattices in $V_{\star,v}$ and fix two lattices Λ'_v and Λ''_v in \mathcal{L}_v that are fixed

by $(R'_{\star,v})^\times$ and $(R''_{\star,v})^\times$, respectively. Thus, if $(i_1, i_2) = \text{inv}(\Lambda'_v, \Lambda''_v)$ as defined in (6) then we have $|i_1 - i_2| = v(\mathfrak{n})$, $R'_{\star,v}$ is the stabilizer of $(\Lambda'_v, \Lambda''_v) \in \mathcal{L}_v \times \mathcal{L}_v$ and $B_{\star,v}^\times$ acts transitively on the set $\mathcal{L}_v(i_1, i_2)$ of pairs of lattices (Λ', Λ'') with $\text{inv}(\Lambda', \Lambda'') = (i_1, i_2)$. Therefore,

$$K_v^\times \backslash B_{\star,v}^\times / R_{\star,v}^\times \simeq K_v^\times \backslash \mathcal{L}_v(i_1, i_2) \quad (13)$$

is contained in $K_v^\times \backslash \mathcal{L}_v \times \mathcal{L}_v$. The map \underline{n}_v that we have defined above on the former set is equal to the restriction of the map defined by (5) on the latter. In particular, the fiber of \underline{n}_v over any $(n', n'') \in \mathbb{N} \times \mathbb{N}$ is finite of order $N_v(n', n'', v(\mathfrak{n}))$ as defined in Lemma 2.1 with $q = N(v)$ the order of the residue field of F at v .

3.4 A sign invariant

In the three diagrams above, the vertical maps are surjective. In Cases 1 and 3, the restriction of the first vertical map to compatible fibers of the two horizontal maps is still obviously surjective. But in Case 2, it may be that such a restriction fails to be surjective. Since for $v \in S$,

$$K_v^\times \backslash B_{S,v}^\times / R_{S,v}^\times \simeq \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if } K_v/F_v \text{ is unramified} \\ \{0\} & \text{otherwise,} \end{cases}$$

this occurs precisely when $v \in S$ is inert in K . Let thus S' be the set of all places v in S that are inert in K . For any such $v \in S'$, the valuation of the reduced norm induces a projection

$$K_v^\times \backslash B_v^\times / R_v^\times \twoheadrightarrow NK_v^\times \backslash F_v^\times / \text{nr}(R_v^\times) \cong \mathbb{Z}/2\mathbb{Z},$$

as well as a bijection $K_v^\times \backslash B_{S,v}^\times / R_{S,v}^\times \cong \mathbb{Z}/2\mathbb{Z}$. We thus obtain maps

$$\varphi_{S'}: \mathcal{CM}(G, H) \rightarrow \text{Gal}_K^{\text{ab}} \backslash \mathcal{CM}(G, H) \rightarrow \prod_{v \in S'} K_v^\times \backslash B_v^\times / R_v^\times \rightarrow (\mathbb{Z}/2\mathbb{Z})^{|S'|}$$

and

$$\psi_{S'}: \mathcal{CM}(G_S, H_S) \rightarrow \text{Gal}_K^{\text{ab}} \backslash \mathcal{CM}(G_S, H_S) \rightarrow \prod_{v \in S'} K_v^\times \backslash B_{S,v}^\times / R_{S,v}^\times \cong (\mathbb{Z}/2\mathbb{Z})^{|S'|}.$$

By construction, $\varphi_{S'} = \psi_{S'} \circ \theta_S$.

The map $\varphi_{S'}$ admits a description in terms of lattices. Under the identification (13), the element $K_v^\times g_v R_v^\times$ corresponds to the pair of lattices $(g_v \Lambda'_v, g_v \Lambda''_v)$. Let y' and y'' be the images of Λ'_v and Λ''_v in the Bruhat–Tits tree $\mathcal{V}_v = F_v^\times \backslash \mathcal{L}_v$ and recall from Section 2 that for all $k \in \mathbb{N}$,

$$\mathcal{V}_v(k) = \{y \in \mathcal{V}_v: n(y) = k\} = \{y \in \mathcal{V}_v: \text{dist}(y, y_0) = k\},$$

where $\mathcal{V}_v(0) = \{y_0\}$ since v is inert in K . Therefore,

$$n(g_v y') - n(y') = \text{dist}(g_v y', y_0) - \text{dist}(y', y_0) \equiv \text{dist}(g_v y', y') \equiv v(\det(g_v)) \pmod{2}$$

and similarly for y'' . Note that the same argument also shows that

$$n(y') - n(y'') \equiv v(\mathfrak{n}) \pmod{2}.$$

The map $\varphi_{S'}$ can thus be computed as follows: for $x \in \mathcal{CM}(G, H)$ with $\mathfrak{c}_f(x) = (\mathfrak{c}', \mathfrak{c}'')$,

$$\varphi_{S'}(x) = (v(\mathfrak{c}') - n(\Lambda'_v) \pmod{2})_{v \in S'} = (v(\mathfrak{c}'') - n(\Lambda''_v) \pmod{2})_{v \in S'} \in (\mathbb{Z}/2\mathbb{Z})^{|S'|}. \quad (14)$$

In particular, $\varphi_{S'}$ factors through the fine conductor map \mathfrak{c}_f .

3.5 Statement of the main theorem

Fix $(\mathfrak{c}', \mathfrak{c}'') \in \mathcal{I}(\text{Ram}_f B)$ such that $v(\mathfrak{c}') - v(\mathfrak{c}'') \equiv v(\mathfrak{n}) \pmod{2}$ for every $v \in S'$. Let $e_{S'}(\mathfrak{c}', \mathfrak{c}'')$ be the element of $(\mathbb{Z}/2\mathbb{Z})^{|S'|}$ defined by the right-hand side of (14). Let $\mathfrak{c} = \mathfrak{c}' \cap \mathfrak{c}''$ and let

$$\kappa = |\text{Gal}(K[\mathfrak{c}]/K[\mathfrak{c}_S])| \prod_{v \in S} N_v(v(\mathfrak{c}'), v(\mathfrak{c}''), v(\mathfrak{n})),$$

where $N_v(n', n'', \delta)$ is the function defined in Lemma 2.1 with $q = N(v)$ being the order of the residue field of F at v .

THEOREM 3.1. *By restriction, the map θ_S induces a surjective κ -to-1 correspondence*

$$\theta_S: \mathfrak{c}_f^{-1}(\mathfrak{c}', \mathfrak{c}'') \rightarrow \mathfrak{c}_{S,f}^{-1}(\mathfrak{c}'_S, \mathfrak{c}''_S) \cap \psi_{S'}^{-1}(e_{S'}(\mathfrak{c}', \mathfrak{c}'')).$$

By restricting to a subspace of the target space, we immediately obtain the following:

COROLLARY 3.2. *For any point $s \in \mathcal{X}(G_S, H_S)$, the map θ_S induces a κ -to-1 correspondence*

$$\theta_S: \text{red}_S(s)^{-1} \cap \mathfrak{c}_f^{-1}(\mathfrak{c}', \mathfrak{c}'') \rightarrow \pi^{-1}(s) \cap \mathfrak{c}_{S,f}^{-1}(\mathfrak{c}'_S, \mathfrak{c}''_S) \cap \psi_{S'}^{-1}(e_{S'}(\mathfrak{c}', \mathfrak{c}'')). \quad (15)$$

3.6 Computation of the fiber for the definite algebra B_S

Before proving the theorem, we shall explain how to compute the right-hand side of the correspondence (15). Fix $s \in \mathcal{X}(G_S, H_S)$ and some $g \in G_S(\mathbb{A}_f)$ above s . Then $b \mapsto b^{-1}g$ induces a bijection

$$R_{S,g}^\times \backslash G_S(\mathbb{Q})/T(\mathbb{Q}) \xrightarrow{\sim} \pi^{-1}(s),$$

where $R_{S,g} = g\widehat{R}_S g^{-1} \cap B_S$ is an Eichler order of level \mathfrak{n}_S in B_S . On the other hand, the map $b \mapsto \text{ad}(b) \circ \iota_S$ induces a bijection

$$G_S(\mathbb{Q})/T(\mathbb{Q}) \xrightarrow{\sim} \text{Hom}_{F\text{-alg}}(K, B_S)$$

that is $G_S(\mathbb{Q}) = B_S^\times$ -equivariant for the natural left actions on both sides. Combining these two identifications, we obtain

$$\pi^{-1}(s) \cong R_{S,g}^\times \backslash \text{Hom}_{F\text{-alg}}(K, B_S).$$

Let also $R'_{S,g} = g\widehat{R}'_S g^{-1} \cap B_S$ and $R''_{S,g} = g\widehat{R}''_S g^{-1} \cap B_S$. These are maximal orders in B_S and $R_{S,g} = R'_{S,g} \cap R''_{S,g}$. Under these identifications,

- The restriction to $\pi^{-1}(s)$ of the fine conductor map $\mathfrak{c}_{S,f}: \mathcal{CM}(G_S, H_S) \rightarrow \mathcal{I}(\text{Ram}_f B_S)^2$ is induced by the $R_{S,g}^\times$ -invariant map

$$\mathfrak{c}_{S,f}^g: \text{Hom}_{F\text{-alg}}(K, B_S) \rightarrow \mathcal{I}(\text{Ram}_f B_S)^2$$

whose fiber over $(\mathfrak{c}'_S, \mathfrak{c}''_S)$ consists of the embeddings $j: K \rightarrow B_S$ such that $j^{-1}(R'_{S,g}) = \mathcal{O}_{\mathfrak{c}'_S}$ and $j^{-1}(R''_{S,g}) = \mathcal{O}_{\mathfrak{c}''_S}$, so that $j^{-1}(R_{S,g}) = \mathcal{O}_{\mathfrak{c}_S}$ with $\mathfrak{c}_S = \mathfrak{c}'_S \cap \mathfrak{c}''_S$.

- The restriction to $\pi^{-1}(s)$ of the sign invariant $\psi_{S'}: \mathcal{CM}(G_S, H_S) \rightarrow (\mathbb{Z}/2\mathbb{Z})^{|S'|}$ is induced by the $R_{S,g}^\times$ -invariant map

$$\psi_{S'}^g: \text{Hom}_{F\text{-alg}}(K, B_S) \xrightarrow{\text{sp}} \prod_{v \in S'} \text{Hom}_{\mathbb{F}(v)}(\mathbb{K}(v), \mathbb{B}_S(v)) \xrightarrow{\sim} (\mathbb{Z}/2\mathbb{Z})^{|S'|},$$

where $\mathbb{F}(v)$, $\mathbb{K}(v)$ and $\mathbb{B}_S(v)$ are the residue fields of the maximal \mathcal{O}_{F_v} -orders in F_v , K_v and $B_{S,v}$, respectively and sp is the natural specialization map. Moreover, the last bijection is the unique isomorphism of $(\mathbb{Z}/2\mathbb{Z})^{|S'|}$ -torsors that maps $\text{sp}(\iota_S)$ to $\psi_{S'}(s) = (v(\det g_v) \pmod{2})_{v \in S'}$.

Remark 1. For a coarse conductor $\mathfrak{c} \in \mathcal{I}(\text{Ram}_f B)$ that is prime to the level \mathfrak{n} , we may replace the fine conductor maps by the coarse ones in the above statements since

$$\mathfrak{c}_f(\mathfrak{c}, \mathfrak{c}) = \mathfrak{c}_g^{-1}(\mathfrak{c}) \quad \text{and} \quad \mathfrak{c}_{S,f}^{-1}(\mathfrak{c}_S, \mathfrak{c}_S) = \mathfrak{c}_{S,g}^{-1}(\mathfrak{c}_S).$$

In this situation, the right-hand side of Corollary 3.2 counts the number of $R_{S,g}^\times$ -conjugacy classes of optimal embeddings $\mathcal{O}_{\mathfrak{c}_S} \hookrightarrow R_S$ that induce a given collection of isomorphisms $\mathbb{K}(v) \rightarrow \mathbb{B}_S(v)$ between the corresponding residue field at $v \in S'$. If, in addition, S' consists of a single prime $S' = \{\ell\}$, one can remove this last condition by identifying the embeddings that are conjugated by the non-trivial automorphism of $\text{Gal}(K/F) \cong \text{Gal}(\mathbb{K}(v)/\mathbb{F}(v))$.

Remark 2. For $F = \mathbb{Q}$, one can interpret the right-hand side of (15) in terms of primitive representations of integers by ternary quadratic forms (see [Gro87, pp.172–173] and [JK10, Prop.4.2] for details).

3.7 Proof of Theorem 3.1

To prove the theorem, let $(\mathfrak{c}'_S, \mathfrak{c}''_S) \in \mathcal{I}(\text{Ram}_f B_S)^2$ be the prime-to- S parts of $(\mathfrak{c}', \mathfrak{c}'') \in \mathcal{I}(\text{Ram}_f B)^2$. Consider the diagram

$$\begin{array}{ccc} \mathfrak{c}_f^{-1}(\mathfrak{c}', \mathfrak{c}'') & \longrightarrow & \text{Gal}_K^{ab} \setminus \mathfrak{c}_f^{-1}(\mathfrak{c}', \mathfrak{c}'') \\ \downarrow \theta_S^{(1)} & & \downarrow \theta_S^{(2)} \\ \mathfrak{c}_{S,f}^{-1}(\mathfrak{c}'_S, \mathfrak{c}''_S) & \longrightarrow & \text{Gal}_K^{ab} \setminus \mathfrak{c}_{S,f}^{-1}(\mathfrak{c}'_S, \mathfrak{c}''_S) \end{array}$$

obtained from the first square of (12) by restriction to the relevant fibers of \mathfrak{c}_f and $\mathfrak{c}_{S,f}$. Note that the vertical maps may fail to be surjective, but the local analysis of Section 2.2 shows that

LEMMA 3.3. *The map $\theta_S^{(2)}$ maps $\text{Gal}_K^{ab} \setminus \mathfrak{c}_f^{-1}(\mathfrak{c}', \mathfrak{c}'')$ onto $\text{Gal}_K^{ab} \setminus \left(\mathfrak{c}_{S,f}^{-1}(\mathfrak{c}'_S, \mathfrak{c}''_S) \cap \psi_{S'}^{-1}(e_{S'}(\mathfrak{c}', \mathfrak{c}'')) \right)$ and moreover, it is $k^{(2)}$ -to-1, where*

$$k^{(2)} = \prod_{v \in S} N_v(v(\mathfrak{c}'), v(\mathfrak{c}''), v(\mathfrak{n})).$$

The same local analysis allows us to compute the number of Galois orbits of CM points with a prescribed fine conductor for each of the algebras B and B_S :

LEMMA 3.4. (i) *The number of Gal_K^{ab} -orbits in $\mathcal{CM}(G, H)$ with fine conductor $(\mathfrak{c}', \mathfrak{c}'') \in \mathcal{I}(\text{Ram}_f B)^2$ is finite and equal to*

$$N_B(\mathfrak{c}', \mathfrak{c}'', \mathfrak{n}) = 2^{\#\{v \in \text{Ram}_f B, v \text{ inert in } K\}} \prod_{v \notin \text{Ram}_f B} N_v(v(\mathfrak{c}'), v(\mathfrak{c}''), v(\mathfrak{n})),$$

where the product is taken over all finite primes. The number of CM points in each of these orbits is equal to

$$h(\mathfrak{c}) = |\text{Gal}(K[\mathfrak{c}]/K)| \quad \text{where } \mathfrak{c} = \mathfrak{c}' \cap \mathfrak{c}''.$$

(ii) *The number of Gal_K^{ab} -orbits in $\mathcal{CM}(G_S, H_S)$ with fine conductor $(\mathfrak{c}'_S, \mathfrak{c}''_S) \in \mathcal{I}(\text{Ram}_f B_S)^2$ is finite and equal to*

$$N_{B_S}(\mathfrak{c}'_S, \mathfrak{c}''_S, \mathfrak{n}) = 2^{\#\{v \in \text{Ram}_f B_S, v \text{ inert in } K\}} \prod_{v \notin \text{Ram}_f B_S} N_v(v(\mathfrak{c}'_S), v(\mathfrak{c}''_S), v(\mathfrak{n})),$$

where the product is taken over all finite primes. The number of CM points in each of these orbits is equal to

$$h(\mathfrak{c}_S) = |\text{Gal}(K[\mathfrak{c}_S]/K)| \quad \text{where } \mathfrak{c}_S = \mathfrak{c}'_S \cap \mathfrak{c}''_S.$$

Theorem 3.1 now follows easily from Lemma 3.3.

4. Application to Galois orbits of Heegner points on modular curves

4.1 The modular curve $Y_0(N)$

Let $N \geq 1$ be an integer and let S be a scheme over $\text{Spec } \mathbb{Z} \left[\frac{1}{N} \right]$. An enhanced elliptic curve (E, C) over S is an elliptic curve E over S together with a closed subgroup C that is locally isomorphic to the constant group scheme $(\mathbb{Z}/N\mathbb{Z})_S$ for the étale topology on S . The modular curve $Y = Y_0(N)$ is a smooth affine curve over $\text{Spec } \mathbb{Z} \left[\frac{1}{N} \right]$ that coarsely represents the contravariant functor mapping S to the set of isomorphism classes of enhanced elliptic curves over S . Here, we say that two enhanced elliptic curves (E_1, C_1) and (E_2, C_2) over S are isomorphic if there is an isomorphism $E_1 \xrightarrow{\sim} E_2$ of elliptic curves over S that maps C_1 to C_2 . We denote by $[E, C]$ the S -valued point of Y defined by an enhanced elliptic curve (E, C) over S . For $S = \text{Spec } \mathbb{C}$, we have the usual isomorphism

$$\Gamma_0(N) \backslash \mathfrak{h} \xrightarrow{\sim} Y(\mathbb{C}) \quad \tau \mapsto [E_\tau, C_\tau] \quad (16)$$

where \mathfrak{h} is the upper half-plane, $\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) : c \equiv 0 \pmod{N} \right\}$ acts on \mathfrak{h} by $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \tau = \frac{a\tau + b}{c\tau + d}$, and $E_\tau = \mathbb{C} / \langle 1, \tau \rangle$ with $C_\tau = \langle N^{-1}, \tau \rangle / \langle 1, \tau \rangle$.

4.2 Isogeny classes

Fix an elliptic curve \mathcal{E} over a base S . Let $\mathcal{I}(\mathcal{E}_*)$ be the contravariant functor that assigns to an S -scheme T the set $\mathcal{I}(\mathcal{E}_T)$ of isomorphism classes of triples (E, C, ϕ) where (E, C) is an enhanced elliptic curve over T and where ϕ is an invertible element of $\text{Hom}_T(E/C, \mathcal{E}_T) \otimes \mathbb{Q}$. An isomorphism between two such triples (E_1, C_1, ϕ_1) and (E_2, C_2, ϕ_2) is an isomorphism of enhanced elliptic curves $\theta: (E_1, C_1) \rightarrow (E_2, C_2)$ such that $\phi_2 \circ \theta = \phi_1$, where $\theta: E_1/C_1 \rightarrow E_2/C_2$ is the induced isomorphism. The group $\text{Aut}_T^0(\mathcal{E}_T)$ of invertible elements in $\text{End}_T^0(\mathcal{E}_T) = \text{End}_T(\mathcal{E}_T) \otimes \mathbb{Q}$ acts on $\mathcal{I}(\mathcal{E}_T)$ by

$$\sigma \cdot (E, C, \phi) = (E, C, \sigma \circ \phi).$$

If $s \in S$ is a geometric point, the map $(E, C, \phi) \mapsto [E, C]$ yields a bijection

$$\text{Aut}_s^0(\mathcal{E}_s) \backslash \mathcal{I}(\mathcal{E}_s) \simeq Y(\mathcal{E}_s) = \{x \in Y(s) \mid x = [E, C] \text{ s.t. } \text{Hom}_s^0(E, \mathcal{E}_s) \neq 0\} \subset Y(s).$$

4.3 Lattices

For a geometric point s of S and a prime number p let $T_p(\mathcal{E}_s)$ be the p -adic Tate module of \mathcal{E}_s if $\text{char}(s) \neq p$ and the covariant Dieudonné crystal of the ℓ -divisible group $\mathcal{E}_s[\ell^\infty]$ if $p = \ell = \text{char}(s)$. Let also $\widehat{T}(\mathcal{E}_s) = \prod_p T_p(\mathcal{E}_s)$ and $\widehat{V}(\mathcal{E}_s) = \widehat{T}(\mathcal{E}_s) \otimes \mathbb{Q}$. If $\text{char}(s) = 0$, a lattice in $\widehat{V}(\mathcal{E}_s)$ is any $\widehat{\mathbb{Z}}$ -submodule that is commensurable with $\widehat{T}(\mathcal{E}_s)$. If $\text{char}(s) = \ell$, a lattice in $\widehat{V}(\mathcal{E}_s) = \widehat{V}^{(\ell)}(\mathcal{E}_s) \times V_\ell(\mathcal{E}_s)$ is a submodule of the form $\widehat{T}^{(\ell)} \times T_\ell$ where $\widehat{T}^{(\ell)}$ is a $\widehat{\mathbb{Z}}^{(\ell)}$ -submodule of $\widehat{V}^{(\ell)}(\mathcal{E}_s)$ that is commensurable with $\widehat{T}^{(\ell)}(\mathcal{E}_s)$, and where T_ℓ is a subcrystal of $V_\ell(\mathcal{E}_s)$. Here,

$$\widehat{\mathbb{Z}}^{(\ell)} = \prod_{p \neq \ell} \mathbb{Z}_p \quad \widehat{V}^{(\ell)}(\mathcal{E}_s) = \prod_{p \neq \ell} V_p(\mathcal{E}_s) \quad \widehat{T}^{(\ell)}(\mathcal{E}_s) = \prod_{p \neq \ell} T_p(\mathcal{E}_s).$$

Let $\mathcal{L}(\mathcal{E}_s)$ be the set of pairs of lattices $(\widehat{T}_1, \widehat{T}_2)$ in $\widehat{V}(\mathcal{E}_s)$ such that $\widehat{T}_1 \subset \widehat{T}_2$ and $\widehat{T}_2/\widehat{T}_1 \simeq \mathbb{Z}/N\mathbb{Z}$ (thus if $\text{char}(s) = \ell$, \widehat{T}_1 and \widehat{T}_2 share the same ℓ -component, since $\ell \nmid N$). For $(E, C, \phi) \in \mathcal{I}(\mathcal{E}_s)$, we have morphisms

$$\widehat{T}(E) \xrightarrow{\text{can}} \widehat{T}(E/C) \subset \widehat{V}(E/C) \xrightarrow{\phi} \widehat{V}(\mathcal{E}_s)$$

and the resulting map

$$(E, C, \phi) \mapsto \left(\phi \circ \text{can} \left(\widehat{T}(E) \right), \phi \left(\widehat{T}(E/C) \right) \right)$$

yields a bijection $\mathcal{I}(\mathcal{E}_s) \simeq \mathcal{L}(\mathcal{E}, s)$. Thus also

$$Y(\mathcal{E}_s) \simeq \text{Aut}_s^0(\mathcal{E}_s) \backslash \mathcal{L}(\mathcal{E}, s).$$

4.4 CM points as special points

Let K be a quadratic imaginary field and let $K \hookrightarrow \mathbb{C}$ be a fixed embedding. An elliptic curve E over a field F is said to have complex multiplication by K if $\text{End}_F^0(E)$ is isomorphic to K . If F is a subfield of \mathbb{C} , we normalize the isomorphism $K \simeq \text{End}_F^0(E)$ by requiring that K acts on the tangent space $\text{Lie } E(\mathbb{C})$ through our fixed embedding $K \hookrightarrow \mathbb{C}$. The conductor of E is the unique positive integer $c(E)$ such that $\text{End}_F(E) \simeq \mathbb{Z} + c(E)\mathcal{O}_K$ inside $\text{End}_F^0(E) \simeq K$. A complex point $x \in Y(\mathbb{C})$ is said to have complex multiplication by K if $x = [E, C]$ for some elliptic curve E over \mathbb{C} with complex multiplication by K . The elliptic curve E/C then also has complex multiplication by K . The fine and coarse conductors of x are respectively equal to

$$\mathfrak{c}_f(x) = (c(E), c(E/C)) \in \mathbb{N} \times \mathbb{N} \quad \text{and} \quad \mathfrak{c}_g(x) = \text{lcm}(c(E), c(E/C)) \in \mathbb{N}.$$

We denote by \mathcal{CM}_K the subset of $Y(\mathbb{C})$ thus defined and refer to its elements as CM points. Note that the bijection (16) restricts to

$$\Gamma_0(N) \backslash (\mathfrak{h} \cap K) \xrightarrow{\cong} \mathcal{CM}_K \quad \tau \mapsto [E_\tau, C_\tau]. \quad (17)$$

Let $\tau \in \mathfrak{h} \cap K$ satisfy the quadratic equation $A\tau^2 + B\tau + C = 0$ where $A > 0$, $A, B, C \in \mathbb{Z}$ and $(A, B, C) = 1$. One can calculate the fine conductor of $[E_\tau, C_\tau]$ as follows: if $\Delta(\tau) = B^2 - 4AC$ is the discriminant modular function and if $D < 0$ is the fundamental discriminant of K then (see, e.g., [Cox89, §7])

$$\mathfrak{c}_f[E_\tau, C_\tau] = \left(\sqrt{\left| \frac{\Delta(\tau)}{D} \right|}, \sqrt{\left| \frac{\Delta(N\tau)}{D} \right|} \right).$$

The point $[E_\tau, C_\tau]$ is a Heegner point if and only if $\Delta(\tau) = \Delta(N\tau)$. It is not hard to show that the latter is equivalent to $A \mid N$ and $(A/N, B, NC) = 1$.

4.5 CM points as isogeny class

All elliptic curves over \mathbb{C} with complex multiplication by K are isogenous. Fix one such curve \mathbb{E} . Then (notation as above)

$$\mathcal{CM}_K = Y(\mathbb{E}) \xleftarrow{\cong} K^\times \backslash \mathcal{I}(\mathbb{E}) \xrightarrow{\cong} K^\times \backslash \mathcal{L}(\mathbb{E}, \mathbb{C}). \quad (18)$$

The fine conductor corresponds to

$$\mathfrak{c}_f(\widehat{T}_1, \widehat{T}_2) = \left(c(\widehat{T}_1), c(\widehat{T}_2) \right)$$

where for a lattice \widehat{T} in $\widehat{V}(\mathbb{E})$, $c(\widehat{T})$ is the conductor of the quadratic order

$$\mathcal{O}_{c(\widehat{T})} = \{s \in K : s\widehat{T} \subseteq \widehat{T}\}.$$

The $\mathcal{O}_{c(\widehat{T})}$ -module \widehat{T} is thus free of rank one.

4.6 The Galois action on CM points

Let K^{ab} be the maximal abelian extension of K inside \mathbb{C} , and let $\text{Art}_K: \widehat{K}^\times \rightarrow \text{Gal}(K^{ab}/K)$ be the reciprocal of the usual Artin reciprocity. In other words, Art_K sends uniformizers to geometric Frobenii [Mil06, p.90]. The main theorem of complex multiplication then says (see [Mil07, Thm.3.10] or [Mil06, Thm.9.10]):

For any $\sigma \in \text{Gal}(\mathbb{C}/K)$ and any $s \in \widehat{K}^\times$ such that $\sigma|_{K^{ab}} = \text{Art}_K(s)$ in $\text{Gal}(K^{ab}/K)$, there exists a unique isogeny $\lambda: \mathbb{E} \rightarrow \sigma\mathbb{E}$ such that for all $y \in \widehat{V}(\mathbb{E})$, $\lambda(sy) = \sigma y$ in $\widehat{V}(\sigma\mathbb{E})$.

Suppose now that $x = [E, C]$ belongs to \mathcal{CM}_K , corresponding to a pair of lattices $(\widehat{T}_1, \widehat{T}_2)$ in $\mathcal{L}(\mathbb{E}, \mathbb{C})$ obtained from (E, C) by choosing a non-zero $\phi: E/C \rightarrow \mathbb{E}$. Fix σ, s and λ as above. Then $\sigma x = [\sigma E, \sigma C]$ belongs to \mathcal{CM}_K , and it corresponds to the pair of lattices $(s\widehat{T}_1, s\widehat{T}_2)$ for the choice

$$\phi' = \lambda^{-1} \circ \sigma\phi: \sigma E/\sigma C \rightarrow \mathbb{E}.$$

The bijection (18) therefore maps the action of $\sigma \in \text{Gal}(\mathbb{C}/K)$ on \mathcal{CM}_K to left multiplication by $s \in \widehat{K}^\times$ on $\mathcal{L}(\mathbb{E}, \mathbb{C})$. It follows that the field of definition of $x \in \mathcal{CM}_K$ is equal to the ring class field $K[\mathfrak{c}_g(x)]$ specified by the coarse conductor $\mathfrak{c}_g(x)$ of x . In fact, it is well-known that the corresponding point $x \in Y(K[\mathfrak{c}_g(x)])$ can be represented by an enhanced elliptic curve (E, C) over $K[\mathfrak{c}_g(x)]$.

4.7 The good reduction of CM points

Let $\overline{\mathbb{Q}}$ be the algebraic closure of \mathbb{Q} inside \mathbb{C} . Let $\ell \nmid N$ be a prime. Fix an embedding $\iota_\ell: \overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_\ell$. Note that ι_ℓ determines a valuation ring $\overline{\mathcal{O}} \subset \overline{\mathbb{Q}}$ whose residue field $\overline{\mathbb{F}}_\ell$ is an algebraic closure of \mathbb{F}_ℓ and a place $\bar{\lambda}$ of $\overline{\mathbb{Q}}$ over ℓ . Let $K[\infty] \subset K^{ab}$ be the union of all ring class fields $K[c]$. For each c (including $c = \infty$), let λ_c be the place of $K[c]$ below $\bar{\lambda}$. Let $\mathcal{O}[c]$ be the valuation ring of λ_c inside $K[c]$ and let $\mathbb{F}[c]$ be its residue field.

There are various equivalent approaches to the reduction theory of CM points at λ :

1. *Reduction via properness.* Let $X_{/\text{Spec } \mathbb{Z}[1/N]}$ be the smooth and proper compactification of Y constructed by Deligne–Rapoport [DR73] when N is prime and by Katz–Mazur [KM85] in general. One gets a reduction map

$$\text{red}: Y(K[c]) \hookrightarrow X(K[c]) \simeq X(\mathcal{O}[c]) \xrightarrow{\text{red}_{\lambda_c}} X(\mathbb{F}[c]),$$

where the bijection $X(K[c]) \simeq X(\mathcal{O}[c])$ follows from the valuative criterion of properness.

2. *Reduction via Néron models.* For a point $x \in \mathcal{CM}_K$ with coarse conductor $c = \mathfrak{c}_g(x)$ write $x = [E, C]$ for some enhanced elliptic curve (E, C) over $K[c]$ with complex multiplication by K . Suppose first that E has good reduction at λ_c . Then (E, C) extends to an enhanced elliptic curve $(\mathcal{E}, \mathcal{C})$ over $\mathcal{O}[c]$ where $\mathcal{E}_{/\mathcal{O}[c]}$ is the Néron model of E . The special fiber of the latter gives a point $\text{red}(x) = [\mathcal{E}_{\mathbb{F}[c]}, \mathcal{C}_{\mathbb{F}[c]}]$ in $Y(\mathbb{F}[c])$. If E does not have good reduction at λ_c , we know by [ST68, Thm.7] that E acquires good reduction at $\bar{\lambda}$ (and indeed everywhere) after a suitable cyclic extension of $K[c]$. We may thus define $\text{red}(x)$ as the point corresponding to the special fiber of the Néron model of the base change of (E, C) to such an extension. This yields a well-defined reduction map

$$\text{red}: \mathcal{CM}_K \rightarrow Y(\overline{\mathbb{F}}_\ell).$$

The above two constructions give the same map on \mathcal{CM}_K with values in $X(\overline{\mathbb{F}}_\ell)$. Since $Y(\mathbb{F}[\infty]) = X(\mathbb{F}[\infty]) \cap Y(\overline{\mathbb{F}}_\ell)$ in $X(\overline{\mathbb{F}}_\ell)$, we have a well-defined map

$$\text{red}: \mathcal{CM}_K \rightarrow Y(\mathbb{F}[\infty]). \tag{19}$$

For a more explicit construction of this map, we can also choose an elliptic curve \mathbb{E} over $K[\infty]$ with complex multiplication by K and good reduction at λ_∞ . Such a curve exists by the elementary theory of complex multiplication and by [ST68, Cor.1]. We then reduce \mathcal{CM}_K as the isogeny class $Y(\mathbb{E})$, as we shall now explain.

3. *Reduction maps for isogeny classes.* Let $S = \text{Spec } \overline{\mathcal{O}}$ with geometric points $g = \text{Spec } \overline{\mathbb{Q}}$ and $s = \text{Spec } \overline{\mathbb{F}}_\ell$. Let $\mathcal{E}_{/S}$ be an elliptic curve. We will eventually take \mathcal{E} to be the Néron model of

an elliptic curve \mathbb{E} with complex multiplication by K , but in what follows we will not make this assumption. The theory of Néron models implies that restriction from S to g yields bijections

$$\mathcal{I}(\mathcal{E}) \xrightarrow{\simeq} \mathcal{I}(\mathcal{E}_g) \quad \text{and} \quad \text{End}_S(\mathcal{E}) \xrightarrow{\simeq} \text{End}_g(\mathcal{E}_g).$$

Restriction from S to s on the other hand gives

$$\mathcal{I}(\mathcal{E}) \rightarrow \mathcal{I}(\mathcal{E}_s) \quad \text{and} \quad \text{End}_S(\mathcal{E}) \hookrightarrow \text{End}_s(\mathcal{E}_s).$$

We get reduction maps $\text{red}: \mathcal{L}(\mathcal{E}, g) \rightarrow \mathcal{L}(\mathcal{E}, s)$ (from left to right):

$$\begin{array}{ccccc} \mathcal{L}(\mathcal{E}, g) & \xleftarrow{\simeq} & \mathcal{I}(\mathcal{E}) & \rightarrow & \mathcal{L}(\mathcal{E}, s) \\ \uparrow \simeq & & \parallel & & \uparrow \simeq \\ \mathcal{I}(\mathcal{E}_g) & \xleftarrow{\simeq} & \mathcal{I}(\mathcal{E}) & \rightarrow & \mathcal{I}(\mathcal{E}_s) \\ \downarrow & & \downarrow & & \downarrow \\ Y(\mathcal{E}_g) = \text{Aut}_g^0(\mathcal{E}_g) \backslash \mathcal{I}(\mathcal{E}_g) & \xleftarrow{\simeq} & \text{Aut}_S^0(\mathcal{E}) \backslash \mathcal{I}(\mathcal{E}) & \rightarrow & \text{Aut}_s^0(\mathcal{E}_s) \backslash \mathcal{I}(\mathcal{E}_s) = Y(\mathcal{E}_s) \end{array}$$

The map $\text{red}: \mathcal{L}(\mathcal{E}, g) \rightarrow \mathcal{L}(\mathcal{E}, s)$ is induced by the natural isomorphism $\widehat{T}^{(\ell)}(\mathcal{E}_g) \xrightarrow{\simeq} \widehat{T}^{(\ell)}(\mathcal{E}_s)$ between the Tate modules away from ℓ , together with a map

$$\text{red}_\ell: \mathcal{L}_\ell(\mathcal{E}, g) \rightarrow \mathcal{L}_\ell(\mathcal{E}, s) \tag{20}$$

from the set $\mathcal{L}_\ell(\mathcal{E}, g)$ of \mathbb{Z}_ℓ -lattices in $V_\ell(\mathcal{E}_g)$ to the set $\mathcal{L}_\ell(\mathcal{E}, s)$ of crystals in $V_\ell(\mathcal{E}_s)$. For $X, Y \in \mathcal{L}_\ell(\mathcal{E}, s)$, we set

$$[X : Y] = \text{length}(X/X \cap Y) - \text{length}(Y/X \cap Y)$$

where the length function is relative to the \mathbb{Z}_ℓ -module structure if $u = g$ and to the W -module structure if $u = s$ (here, $W := W(\overline{\mathbb{F}}_\ell)$ is the ring of Witt vectors of $\overline{\mathbb{F}}_\ell$). Then for all $X \in \mathcal{L}_\ell(\mathcal{E}, g)$,

$$[T_\ell(\mathcal{E}_g) : X] = [T_\ell(\mathcal{E}_s) : \text{red}_\ell(X)] \tag{21}$$

Indeed, we may choose a triple $(E, C, \phi) \in \mathcal{I}(\mathcal{E})$ such that

$$X = \phi(T_\ell((E/C)_g)) \quad \text{and} \quad \text{red}_\ell(X) = \phi(T_\ell((E/C)_s))$$

and then both sides of (21) are equal to the exponent of ℓ in the degree of ϕ .

4.8 The supersingular case

If \mathcal{E}_s is a supersingular elliptic curve, then

$$(T_\ell(\mathcal{E}_s), F, V) \simeq (W^2, \pi_\ell \sigma, \pi_\ell \sigma^{-1}) \quad \text{with} \quad \pi_\ell = \begin{pmatrix} 0 & 1 \\ p & 0 \end{pmatrix} \in M_2(W)$$

where σ is the Frobenius automorphism of W . We thus obtain

$$\mathcal{L}_\ell(\mathcal{E}_s) \simeq \{\pi_\ell^i W^2 \subset \mathcal{K}^2 : i \in \mathbb{Z}\}$$

where \mathcal{K} is the fraction field of W . In particular, the map (20) is uniquely determined by (21). Moreover, the endomorphism ring

$$\text{End}(T_\ell(\mathcal{E}_s), F, V) \simeq \{x \in M_2(W) : \pi \sigma(x) \pi^{-1} = x\}$$

is the maximal \mathbb{Z}_ℓ -order \mathcal{R} of a non-split quaternion algebra $\mathcal{B} = \mathcal{R} \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$ over \mathbb{Q}_ℓ and $\mathcal{B}^\times \simeq \mathcal{R}^\times \times \pi_\ell^{\mathbb{Z}}$ acts transitively on $\mathcal{L}_\ell(\mathcal{E}, z)$.

Remark 3. The reduction map (20) is more difficult to analyze when \mathcal{E}_s is an ordinary elliptic curve, especially when \mathcal{E}_g does *not* have complex multiplication.

4.9 Matrices

We choose a subgroup T' of $T = H_1(\mathcal{E}(\mathbb{C}), \mathbb{Z})$ such that $T/T' \simeq \mathbb{Z}/N\mathbb{Z}$. We put $V = T \otimes \mathbb{Q}$, $B = \text{End}_{\mathbb{Q}}(V)$ and

$$R = \{x \in B : xT' \subset T' \text{ and } xT \subset T\},$$

an Eichler order of level N in $B \simeq M_2(\mathbb{Q})$. This gives rise to identifications

$$\widehat{T}(\mathcal{E}_g) \simeq \widehat{T}(\mathcal{E}, \mathbb{C}) \simeq \widehat{T}, \quad \widehat{V}(\mathcal{E}_g) \simeq \widehat{V}(\mathcal{E}, \mathbb{C}) \simeq \widehat{V} \quad \text{and} \quad \widehat{B}^\times / \widehat{R}^\times \simeq \mathcal{L}(\mathcal{E}, g),$$

where the last map sends b to $(b\widehat{T}', b\widehat{T})$. Suppose that \mathcal{E}_s is supersingular and let $B_{\{\ell\}} = \text{End}_s^0(\mathcal{E}_s)$, a definite quaternion algebra over \mathbb{Q} with $\text{Ram}_f(B_{\{\ell\}}) = \{\ell\}$. For $p \neq \ell$, the left action of $B_{\{\ell\}, p}$ on $V_p(\mathcal{E}_s) \simeq V_p(\mathcal{E}, g) \simeq V_g$ yields an isomorphism $\theta_p : B_p \simeq B_{\{\ell\}, p}$, and the left action of $B_{\{\ell\}, \ell}$ on $V_\ell(\mathcal{E}_s)$ yields an isomorphism $\theta_\ell : \mathcal{B} \simeq B_{\{\ell\}, \ell}$ with \mathcal{B} as above. Put $\text{red}(\widehat{T}', \widehat{T}) = (\widehat{T}'_s, \widehat{T}_s)$ and

$$R_{\{\ell\}} = \left\{ x \in B_{\{\ell\}} : x\widehat{T}'_s \subset \widehat{T}'_s \text{ and } x\widehat{T}_s \subset \widehat{T}_s \right\}.$$

Thus $\theta_p(R_p) = R_{\{\ell\}, p}$ for all $p \neq \ell$, and $\theta_\ell(\mathcal{R}) = R_{\{\ell\}, \ell}$ with \mathcal{R} as above. The map $b \mapsto (b\widehat{T}'_s, b\widehat{T}_s)$ yields an identification $\widehat{B}_{\{\ell\}}^\times / \widehat{R}_{\{\ell\}}^\times \simeq \mathcal{L}(\mathcal{E}, s)$, and the reduction map

$$\widehat{B}^\times / \widehat{R}^\times \simeq \mathcal{L}(\mathcal{E}, g) \xrightarrow{\text{red}} \mathcal{L}(\mathcal{E}, s) \simeq \widehat{B}_{\{\ell\}}^\times / \widehat{R}_{\{\ell\}}^\times$$

sends $\widehat{bR}^\times = \prod b_p R_p^\times$ to $\widehat{b'R}_{\{\ell\}}^\times = \prod b'_p R_{\{\ell\}, p}^\times$ where $b'_p = \theta_p(b_p)$ for $p \neq \ell$ and $b'_\ell = \theta_\ell(\pi_\ell)^{v_\ell}$ for $v_\ell = \text{ord}_\ell(\text{nr}(b_\ell))$. We finally obtain a reduction map

$$\text{Aut}_g^0(\mathcal{E}_g) \backslash \widehat{B}^\times / \widehat{R}^\times \simeq Y(\mathcal{E}_g) \xrightarrow{\text{red}} Y(\mathcal{E}_s) \simeq B_{\{\ell\}}^\times \backslash \widehat{B}_{\{\ell\}}^\times / \widehat{R}_{\{\ell\}}^\times$$

where $\text{End}_g^0(\mathcal{E}_g)$ embeds in B through its action on $V = H_1(\mathcal{E}_g(\mathbb{C}), \mathbb{Q})$.

4.10 The supersingular reduction of CM points

We now assume that ℓ is inert in K , and let \mathcal{E} be the Néron model over $\overline{\mathcal{O}}$ of an elliptic curve with complex multiplication by K , i.e. $\text{End}_g^0(\mathcal{E}_g) = K$. Then \mathcal{E}_s is indeed supersingular, $Y(\mathcal{E}_g) = \mathcal{CM}_K$ and $Y^{ss}(\mathcal{E}_s) = Y^{ss}(\overline{\mathbb{F}}_\ell)$, the set of supersingular points in $Y(\overline{\mathbb{F}}_\ell)$. We have now identified the geometric reduction map

$$\text{red} : \mathcal{CM}_K \rightarrow Y^{ss}(\overline{\mathbb{F}}_\ell) \tag{22}$$

with the adelic reduction map which we had previously considered.

Remark 4. The surjectivity of (22) implies that $Y^{ss}(\overline{\mathbb{F}}_\ell) \subset Y(\mathbb{F}[\infty])$. On the other hand, class field theory shows that $\mathbb{F}[\infty] \simeq \mathbb{F}_{\ell^2}$. We thus retrieve the well-known fact that $Y^{ss}(\overline{\mathbb{F}}_\ell) \subset Y(\mathbb{F}_{\ell^2})$.

The main correspondence between Heegner points and optimal embeddings. Let now c be a positive integer satisfying $(c, \ell N) = 1$ and let \mathcal{O}_c be the corresponding order in K . As a consequence of the above identifications, our Corollary 3.2 implies the following:

COROLLARY 4.1. *Let $s \in Y^{ss}(\overline{\mathbb{F}}_\ell)$ be a supersingular point. Choose $[\widetilde{E}, \widetilde{C}] = s$ and define*

$$R_s = \text{End}(\widetilde{E}, \widetilde{C}) = \left\{ \alpha \in \text{End}(\widetilde{E}) : \alpha(\widetilde{C}) \subset \widetilde{C} \right\}.$$

There is a one-to-one correspondence between the following two sets:

$$\left\{ \begin{array}{l} \text{Points } x \in \mathcal{CM}_K \text{ on } Y \\ \text{of conductor } c \text{ reducing to } s \end{array} \right\} \iff \left\{ \begin{array}{l} R_s^\times - \text{conjugacy classes of} \\ \text{conjugate pairs of} \\ \text{optimal embeddings } \mathcal{O}_c \hookrightarrow R_s \end{array} \right\}.$$

Remark 5. This was precisely the correspondence needed in [JK10] to translate the equidistribution question for Heegner points to a question about optimal embeddings. It is shown in [JK10, §4.1] that the latter relates to counting primitive representations of integers by ternary quadratic forms. For $c = 1$, the above correspondence is known as Deuring lifting theorem (see [Deu41]) and has been subsequently refined (as a correspondence) by Gross and Zagier [GZ85, Prop.2.7].

Remark 6. The left-to-right map in Corollary 4.1 is rather natural. Let (E, C) be an enhanced elliptic curve over $\overline{\mathbb{Q}}$ with complex multiplication by K and coarse conductor c . Extend (E, C) to an enhanced elliptic curve $(\mathcal{E}, \mathcal{C})$ over $\overline{\mathcal{O}}$ and suppose that $(\mathcal{E}_s, \mathcal{C}_s) \simeq (\tilde{E}, \tilde{C})$. A choice of isomorphisms $(\mathcal{E}_s, \mathcal{C}_s) \xrightarrow{\simeq} (\tilde{E}, \tilde{C})$ and $K \xrightarrow{\simeq} \text{End}^0(\mathcal{E}, \mathcal{C})$ yields an embedding $\mathcal{O}_c \hookrightarrow R_s$, and the resulting R_s^\times -conjugacy class of pairs of conjugate embeddings does not depend upon these two choices. Since $\ell \nmid c$, it is still fairly straightforward to verify that the embeddings thus obtained are optimal. What is not obvious is that this construction gives a one-to-one correspondence. This is precisely what we establish in our Theorem 3.1 in a greater generality for quaternion algebras.

ACKNOWLEDGEMENTS

We would like to thank Dick Gross, David Jao, Ben Kane, Philippe Michel, Ken Ribet and Nicolas Templier for helpful discussions.

REFERENCES

- Bas63 H. Bass, *On the ubiquity of Gorenstein rings*, Math. Z. (1963), no. 82, 8–28.
- Car86 H. Carayol, *Sur la mauvaise réduction des courbes de Shimura*, Compositio Math. **59** (1986), no. 2, 151–230.
- Cor02 C. Cornut, *Mazur’s conjecture on higher Heegner points*, Invent. Math. **148** (2002), no. 3, 495–523.
- Cox89 D. Cox, *Primes of the form $x^2 + ny^2$* , Wiley, New York, NY, 1989.
- CV05 C. Cornut and V. Vatsal, *CM points and quaternion algebras*, Doc. Math. **10** (2005), 263–309 (electronic).
- CV07 ———, *Nontriviality of Rankin-Selberg L -functions and CM points*, L -functions and Galois representations, London Math. Soc. Lecture Note Ser., vol. 320, Cambridge Univ. Press, Cambridge, 2007, pp. 121–186.
- Deu41 M. Deuring, *Die Typen der Multiplikatorenringe elliptischer Funktionenkörpern*, Abh. Math. Sem. Hansischen Univ., vol. 14, 1941, pp. 197–272.
- DR73 P. Deligne and M. Rapoport, *Les schémas de modules de courbes elliptiques*, Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972) (Berlin), Springer, 1973, pp. 143–316. Lecture Notes in Math., Vol. 349.
- Gro87 B. H. Gross, *Heights and the special values of L -series*, Number theory (Montreal, Que., 1985), CMS Conf. Proc., vol. 7, Amer. Math. Soc., Providence, RI, 1987, pp. 115–187.
- GZ85 B. Gross and D. Zagier, *On singular moduli*, J. Reine Angew. Math. **355** (1985), 191–220.
- HM06 G. Harcos and P. Michel, *The subconvexity problem for Rankin-Selberg L -functions and equidistribution of Heegner points. II*, Invent. Math. **163** (2006), no. 3, 581–655.
- JK10 D. Jetchev and B. Kane, *Equidistribution of Heegner points and ternary quadratic forms*, Math. Annalen (2010), to appear.
- KM85 N. M. Katz and B. Mazur, *Arithmetic moduli of elliptic curves*, Princeton University Press, Princeton, N.J., 1985.
- Mic04 P. Michel, *The subconvexity problem for Rankin-Selberg L -functions and equidistribution of Heegner points*, Ann. of Math. (2) **160** (2004), no. 1, 185–236.
- Mil06 J. Milne, *Complex multiplication*, <http://www.jmilne.org/math/CourseNotes/CM.pdf> (2006).

REDUCTION MAPS

- Mil07 ———, *The fundamental theorem of complex multiplication*, <http://www.jmilne.org/math/articles/2007c.pdf> (2007).
- Mol10a S. Molina, *Equations of hyperelliptic Shimura curves*, preprint (2010).
- Mol10b ———, *Ribet bimodules and the specialization of Heegner points*, preprint (2010).
- Rib90 K. A. Ribet, *On modular representations of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ arising from modular forms*, *Invent. Math.* **100** (1990), no. 2, 431–476.
- ST68 J.-P. Serre and J. T. Tate, *Good reduction of abelian varieties*, *Ann. of Math. (2)* **88** (1968), 492–517.
- Vat02 V. Vatsal, *Uniform distribution of Heegner points*, *Invent. Math.* **148** (2002), no. 1, 1–46.
- Vig80 M.-F. Vignéras, *Arithmétique des algèbres de quaternions*, *Lecture Notes in Mathematics*, vol. 800, Springer, Berlin, 1980.

Christophe Cornut cornut@math.jussieu.fr

Institut de Mathématiques de Jussieu, 175 rue du Chevaleret, 75013 Paris, France

Dimitar Jetchev dimitar.jetchev@epfl.ch

Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland