

On the Chow ring of very general abelian varieties and a question of Pirola

Claire Voisin*

Abstract

We prove that for a very general abelian variety of dimension ≥ 4 , a divisor $D \in \mathrm{CH}^1(A)$ that satisfies $D^2 = 0$ in $\mathrm{CH}^2(A)$ is of torsion. The same result is also established for a very general Jacobian in genus 4. We use then the second statement in order to prove a conjecture of Pirola, which states that any rational section of the Kummer fibration $K = J/\pm \mathrm{Id} \rightarrow \mathcal{M}_4$, where $J \rightarrow \mathcal{M}_4$ is the Jacobian fibration, must be a multiple of the Griffiths-Pirola section given by the difference of the two trigonal divisors.

1 Introduction

Let A be an abelian variety over $\mathbb{C}$. The study of the Chow ring $\mathrm{CH}^*(A)$ with $\mathbb{Q}$ -coefficients has been initiated by Bloch [5] and Beauville [3] who discovered beautiful structures like the Beauville decomposition [3], [4] and the Fourier-Mukai transform [16] induced by the abelian group structure of A on one hand, and the isogeny between A and its dual on the other hand. In [23], we started the study of more basic questions about the ring structure of $\mathrm{CH}^*(A)$. The paper [23] focused on the Pontryagin powers of 0-cycles, but via the Fourier-Mukai transform, this can be reformulated as questions concerning the ordinary ring structure of $\mathrm{CH}^*(A)$ (see [23, Lemma 1.7]). The basic question is then the following: given r , what can be said about the set of divisors $D \in \mathrm{CH}^1(A)_{\mathrm{hom}}$ such that $D^r = 0$ in $\mathrm{CH}^r(A)$? If $r \geq 2$, this set is a countable union of closed algebraic subsets of $\mathrm{Pic}^0(A)$ which contains the torsion points of $\mathrm{Pic}^0(A)$. In *loc. cit.*, it was proved that for a very general abelian variety of dimension ≥ 3 , the set of divisors $D \in \mathrm{Pic}^0(A)$ such that $D^2 = 0$ in $\mathrm{CH}^2(A)$ has dimension 0. In the papers [9], the locus of abelian varieties for which this statement fails is studied in a very precise way : Away from decomposable abelian varieties, this locus has dimension $\leq 2g - 1$ (so in particular, one recovers the fact that this locus is not the whole of $\mathcal{A}_g$ when $g \geq 3$).

Note that one can ask similar questions for symmetric products of curves, or for their Jacobians. Let C be a smooth projective curve and let $j : C \hookrightarrow J(C)$ be an embedding. The pull-back map

$$j^* : \mathrm{Pic}^0(J(C)) \rightarrow \mathrm{Pic}^0(C)$$

is an isomorphism, which does not depend on the choice of j . Note that we also have the isomorphism $\mathrm{Pic}^0(C) \cong \mathrm{Pic}^0(C^{(2)})$, and we can consider, for any

$$D \in \mathrm{Pic}^0(C) \cong \mathrm{Pic}^0(C^{(2)}) \cong \mathrm{Pic}^0(J(C))$$

the codimension 2 cycles $D^2 \in \mathrm{CH}^2(J(C))$ and $D^2 \in \mathrm{CH}^2(C^{(2)})$. It turns out that the vanishing of these two squares are equivalent:

Lemma 1.1. *One has $D^2 = 0$ in $\mathrm{CH}^2(J(C))$ if and only if $D^2 = 0$ in $\mathrm{CH}^2(C^{(2)})$.*

*The author is supported by the ERC Synergy Grant HyperK (Grant agreement No. 854361).

Proof. Clearly the first vanishing implies the second one by restriction from $J(C)$ to $C^{(2)}$. In the other direction, assume that D^2 vanishes in $\mathrm{CH}_0(C^{(2)})$. Then by Bloch's formula [5, Lemma 1.1(ii)], one deduces that $s^{*2} = 0$ in $\mathrm{CH}_0(J(C))$ where $s := D \cdot C \in \mathrm{CH}_0(J(C))$ and $*$ is the Pontryagin product. Denoting by F the Fourier-Mukai transform of $J(C)$ (see [3]) it follows that

$$F(D \cdot C) \cdot F(D \cdot C) = 0 \text{ in } \mathrm{CH}^*(J(C)) \quad (1)$$

since F exchanges the Pontryagin product and the intersection product. Equation (1) then implies $D^2 = 0$ in $\mathrm{CH}^2(J(C))$ because $F(D \cdot C) = D + \Gamma$ where Γ is a sum of cycles of codimension ≥ 2 . $\square$

Our first main result in the present paper is the following

Theorem 1.2. (i) *Let A be a very general abelian variety of dimension at least 4. Then, if $D \in \mathrm{CH}^1(A)_{\mathrm{hom}} = \mathrm{Pic}^0(A)$, and $D^2 = 0$ in $\mathrm{CH}^2(A)$, D is of torsion.*

(ii) *Let C be a very general curve of genus 4. Then if $D \in \mathrm{Pic}^0(C) = \mathrm{Pic}^0(C^{(2)})$ and $D^2 = 0$ in $\mathrm{CH}^2(C^{(2)})$ (or equivalently by Lemma 1.1, $D^2 = 0$ in $\mathrm{CH}^2(J(C))$), D is of torsion.*

Using [23, Lemma 1.7], one can also rephrase Theorem 1.2 as follows, using the Pontryagin product of 0-cycles on A . Denote by 0_A the origin of an abelian variety A (it is needed if we allude to the group structure of A , which appears in the definition of the Pontryagin product).

Corollary 1.3. *For a very general abelian variety A of dimension $g \geq 4$, the set of points $x \in A$ such that $(\{x\} - \{0_A\})^{*2} = 0$ in $\mathrm{CH}_0(A)$ consists of the torsion points of A . The same statement holds if A is a very general Jacobian in genus 4.*

Corollary 1.4. *If A is a very general abelian variety of dimension $g \geq 4$ or the Jacobian of a very general curve of genus 4, the orbit $O_{2\{0_A\}} \subset A^{(2)}$ for rational equivalence on A of the 0-cycle $2\{0_A\} \in A^{(2)}$, that is, the set of $z \in A^{(2)}$ such that z is rationally equivalent to $2\{0_A\}$ in A , consists only of (effective, degree 2) 0-cycles supported on torsion points of A .*

Proof. By [23, Proposition 1.9], we know that a rational equivalence relation

$$\{x_1\} + \dots + \{x_k\} = k\{0_A\} \text{ in } \mathrm{CH}_0(A) \quad (2)$$

implies that $(\{x_i\} - \{0_A\})^{*k} = 0$ in $\mathrm{CH}_0(A)$ for $i = 1, \dots, k$. Thus, for $k = 2$, when A is very general, (2) implies that each x_i is a torsion point of A by Corollary 1.3. $\square$

Remark 1.5. It is likely that Theorem 1.2(ii) also holds for any genus $g \geq 4$. The extra work needed to make the same proof work in higher genus is to prove Proposition 3.1(ii) in higher genus.

Remark 1.6. The situation is unclear in genus 3, but for abelian surfaces, there are uncountably many points $x \in A$ such that $(\{x\} - \{0_A\})^{*2} = 0$ in $\mathrm{CH}_0(A)$ (in particular there are nontorsion points). Indeed, let $A = J(C)$ for a genus 2 curve $C \subset A$ and assume that C passes through 0_A , and that the corresponding point $x_0 \in C$ is a Weierstrass point. Then, by the same argument as above, any $x \in C \subset A$ has the property that $(\{x\} - \{0_A\})^{*2} = 0$ in $\mathrm{CH}_0(A)$, since, for some $y \in C$, $\{x\} + \{y\}$ is rationally equivalent to $2x_0$ in C , hence to $2\{0_A\}$ in A .

Remark 1.7. It is likely that the method used to prove Theorem 1.2 also allows to study higher powers. One may even conjecture that if $g \geq 2k$, and A is a very general abelian variety of dimension g , the only divisors $D \in \mathrm{CH}^1(A)$ such that $D^k = 0$ in $\mathrm{CH}^k(A)$ are the torsion divisors.

Our motivation to prove Theorem 1.2 was an intriguing and seemingly unrelated question of Pirola related to the Franchetta conjecture for generic curves. The Franchetta conjecture (now a theorem, see [15]) says that if C is the generic curve of genus g , then $\text{Pic}(C)$ is the cyclic group generated by K_C . If one considers a general curve of genus 4 over $\mathbb{C}$, it is naturally embedded in a 2-dimensional quadric in its canonical embedding, hence it has two g_3^1 's, say D_1, D_2 , coming from the restrictions of the two rulings of the quadric to C . The difference $D_1 - D_2 \in \text{Pic}^0(C)$ is a divisor of degree 0 on C , nontrivial in general, which is canonically defined up to sign, so for the generic curve of genus 4, which is defined over $\mathbb{C}(\mathcal{M}_4)$, it provides an element of $\text{Pic}^0(C_L)$ defined over a degree 2-field extension L of the function field $\mathbb{C}(\mathcal{M}_4)$. Equivalently, it provides a relative divisor on the universal curve $\tilde{C} \rightarrow \tilde{\mathcal{M}}_4$ for some degree 2 cover $\tilde{\mathcal{M}}_4 \rightarrow \mathcal{M}_4$.

We will call the corresponding section γ_P of $\text{Pic}^0(\tilde{C}/\tilde{\mathcal{M}}_4)$ the Griffiths-Pirola normal function. It has been studied indeed in [13] by Griffiths, who explained how to recover the general curve C of genus 4 from the infinitesimal invariant of this normal function at $[C] \in \mathcal{M}_4$ (see also [20] and Section 2). We can also see γ_P as a rational section of the Kummer fibration $K = J/\pm \text{Id} \rightarrow \mathcal{M}_4$. The following result answers a question of Pirola.

Theorem 1.8. *Let $C \rightarrow \mathcal{M}_4$ be the universal curve of genus 4, and $J \rightarrow \mathcal{M}_4$ the associated Jacobian fibration. Then all rational sections of the fibration $K = J/\pm \text{Id} \rightarrow \mathcal{M}_4$ are obtained by taking multiples of the Griffiths-Pirola normal function γ_P .*

This result suggests the following conjecture (proposed by Pirola and Farkas).

Conjecture 1.9. *For $g \geq 5$, the only rational section of the Kummer fibration $K = J/\pm \text{Id} \rightarrow \mathcal{M}_g$ is the zero section.*

The proof of Theorem 1.2 uses the theory of Griffiths infinitesimal invariants $\delta\nu$ of normal functions ν [13], and also its extension to higher codimensional cycles introduced in [24], where rational equivalence of 0-cycles of very general surfaces in $\mathbb{P}^3$ was studied (see Section 2). For the proof of Theorem 1.2, one has to analyze the square map for infinitesimal invariants. For a normal function ν defined on a generically finite cover of $\mathcal{A}_g$, $g \geq 4$, with the property that $(\delta\nu)^2 = 0$, this formal analysis provides us with distinguished representative of the cohomology class $[\nu]$ given by a closed and decomposable algebraic section of the bundle $\mathcal{H}^1 \otimes \Omega_B$. Results of Ax-Shanuel type (see [6], [1], [2]) then imply that such a form must be 0, from which the conclusion that ν is of torsion follows in a rather classical way.

The proof of Theorem 1.8 uses another ingredient. The main difficulty in Pirola's question, which makes it rather different from Franchetta's conjecture, is the fact that the rational sections γ that we want to describe correspond to sections of the Jacobian fibration over a double cover of $\mathcal{M}_4$ that is not specified. Our starting point is the fact that, the divisor $D_\gamma(t) \in \text{Pic}^0(C_t) = \text{Pic}^0(C_t^{(2)})$ being defined up to sign, the codimension 2 cycle $D_\gamma(t)^2 \in \text{CH}^2(C_t^{(2)})$ is defined over the function field of $\mathcal{M}_4$. One easily deduces from this that it is a multiple of $D_{\gamma_P}(t)^2$ (which is also proportional to the Green-Griffiths 0-cycle of [10], see [20]). Our proof consists then in analyzing, using again the theory of infinitesimal invariants, the rational equivalence relation

$$D_\gamma^2 = \lambda D_{\gamma_P}^2 \text{ in } \text{CH}^2(C_t^{(2)}).$$

Thanks. *This work was inspired by lectures of Gian Pietro Pirola at the workshop "Topics in Hodge theory" at the university of Ashoka. I thank the organizers of this nice event. I also thank Greg Baldi for the proof of Proposition 3.16, and Bruno Klingler and Gavril Farkas for their interest and constructive comments.*

2 A review of infinitesimal invariants

Infinitesimal invariants of normal functions were first introduced by Griffiths (see [13]). Assume that we have an abelian fibration $\pi : J \rightarrow M$. Let D be a divisor on J , which is

homologous to 0 along the fibers of π . There is an associated section ν_D of the sheaf $\mathcal{J}^\vee$ of holomorphic sections of the dual abelian fibration $J^\vee := \text{Pic}^0(J/M) \rightarrow M$, such that $\nu_D(m) := D|_{J_m}$. By uniformization, this sheaf fits into an exact sequence

$$0 \rightarrow H_{\mathbb{Z}}^1 \oplus \mathcal{H}^{1,0} \rightarrow \mathcal{H}^1 \rightarrow \mathcal{J}^\vee \rightarrow 0, \quad (3)$$

where $H_{\mathbb{Z}}^1 := R^1\pi_*\mathbb{Z}$, $\mathcal{H}^1 := H_{\mathbb{Z}}^1 \otimes \mathcal{O}_M$, and $\mathcal{H}^{1,0} \subset \mathcal{H}^1$ are the Hodge bundles for the weight 1 variation of Hodge structure on $H_{\mathbb{Z}}^1$.

The Griffiths infinitesimal invariant $\delta\nu_D(m)$ at the point $m \in M$ is defined as follows: using the exact sequence (3), one can choose local (for the Euclidean topology) lifts

$$\tilde{\nu} \in \Gamma(\mathcal{H}^1)$$

of ν_D . Such local lifts are unique up to the addition of sections $\nu_{\mathbb{Z}}$ of $H_{\mathbb{Z}}^1$ and $\nu^{1,0}$ of $\mathcal{H}^{1,0}$. It follows that $\nabla\tilde{\nu} \in \Gamma(\mathcal{H}^1 \otimes \Omega_M)$ is defined up to adding a section $\nabla\nu^{1,0} \in \Gamma(\mathcal{H}^1 \otimes \Omega_M)$, and thus the $(0,1)$ -projection $(\nabla\tilde{\nu})^{0,1} \in \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M)$ of $\nabla\tilde{\nu}$ is defined modulo $\overline{\nabla}\mathcal{H}^{1,0}$, where

$$\overline{\nabla} : \mathcal{H}^{1,0} \rightarrow \mathcal{H}^{0,1} \otimes \Omega$$

is the infinitesimal variation of Hodge structures of our family. As $\overline{\nabla}$ is $\mathcal{O}_M$ -linear, the quantity

$$\delta\nu_D(m) := (\nabla\tilde{\nu})_m^{0,1} \in (\mathcal{H}_m^{0,1} \otimes \Omega_{M,m}) / \overline{\nabla}(\mathcal{H}_m^{1,0}) \quad (4)$$

is thus well defined at any point $m \in M$. It computes part of the derivative of the normal function ν_D and depends only on the data of ν_D at first order at m . We will use the following

Lemma 2.1. *Let $\pi : J \rightarrow M$ be a family of abelian varieties of dimension g with corresponding sheaf $\mathcal{J}^\vee$ as above and let $\nu \in \Gamma(M, \mathcal{J}^\vee)$ be a normal function. Assume that ν has trivial infinitesimal invariant at the general point of M , and that*

(i) *The Griffiths differential $\overline{\nabla} : \mathcal{H}^{1,0} \rightarrow \mathcal{H}^{0,1} \otimes \Omega_M$ is injective (that is, pointwise injective at the general point of M).*

(ii) *The second Griffiths differential $\overline{\nabla}_1 : \mathcal{H}^{1,0} \otimes \Omega_M \rightarrow \mathcal{H}^{0,1} \otimes \Omega_M^2$ is injective (that is, pointwise injective at the general point of M).*

(iii) *The monodromy group acting on the degree 1 cohomology of the fibers of the family $J \rightarrow M$ has finite index in $\text{Sp}(2g)$.*

Then ν is a torsion section of $\mathcal{J}^\vee$.

In (ii), the map $\overline{\nabla}_1$ is defined by the formula $\overline{\nabla}_1(\omega \otimes \alpha) = \overline{\nabla}\omega \wedge \alpha$. Note that Condition (ii) implies Condition (i), but we will need the two injectivity statements in the proof.

Remark 2.2. Condition (ii) is actually essential. Consider the case of a 1-dimensional basis M and assume that (i) is satisfied. Then $\overline{\nabla} : \mathcal{H}^{1,0} \rightarrow \mathcal{H}^{0,1} \otimes \Omega_M$ is an isomorphism at the general point of M . Thus the space of infinitesimal invariants is trivial in this case, so any normal function has trivial infinitesimal invariant at the general point, while of course there always exist nontorsion normal functions, maybe after making a base change.

Proof of Lemma 2.1. The argument is classical and appears in the proof of [26, Theorem 7.22] in a slightly different geometric context. We reproduce it here for completeness. Note that it suffices to prove the conclusion after restriction to a dense Zariski open set of M , which does not change assumption (iii). Thus we can assume that $\delta\nu = 0$ at any point of M . This implies that any local lift $\tilde{\nu}_0$ of ν satisfies $(\nabla\tilde{\nu}_0)^{0,1} = (\nabla\eta)^{0,1}$ in $\Gamma(\mathcal{H}^{0,1})$ for some local section $\eta \in \Gamma(\mathcal{H}^{1,0})$, so that the local lift $\tilde{\nu} := \tilde{\nu}_0 - \eta$ of ν has the property that

$$\nabla\tilde{\nu} \in \Gamma(\mathcal{H}^{1,0} \otimes \Omega_M). \quad (5)$$

We observe now that, since $\nabla \circ \nabla = 0$, we have

$$\nabla(\nabla\tilde{\nu}) = 0. \quad (6)$$

Then by taking the $(0, 1)$ -part in (6) and using (5), we get that $\bar{\nabla}_1(\nabla\tilde{\nu}) = 0$. Thus condition (ii) implies that we have in fact

$$\nabla\tilde{\nu} = 0, \quad (7)$$

that is, $\tilde{\nu}$ is a horizontal section of $\mathcal{H}^1$. Any other local horizontal local lift $\tilde{\nu}'$ of ν is of the form

$$\tilde{\nu}' = \tilde{\nu} + \eta_{\mathbb{Z}} + \eta^{1,0}$$

for some local sections $\eta_{\mathbb{Z}} \in \Gamma(H_{\mathbb{Z}}^1)$ and $\eta^{1,0} \in \Gamma(\mathcal{H}^{1,0})$. We then have

$$\nabla(\eta^{1,0}) = 0, \quad (8)$$

as $\tilde{\nu}', \tilde{\nu}$ and $\eta_{\mathbb{Z}}$ are horizontal. By Condition (i), (8) implies that η_F vanishes identically. It follows that the horizontal local lifts $\tilde{\nu}$ of ν are unique up to sections of $H_{\mathbb{Z}}^1$. We thus proved that the global section $\nu \in \Gamma(M, \mathcal{J})$ is induced by a global section

$$\bar{\nu} \in \Gamma(M, H_C^1/H_{\mathbb{Z}}^1).$$

If $\rho : \pi_1(M, m_0) \rightarrow \text{Aut}(H^1(J_{m_0}, \mathbb{Z}))$ denotes the monodromy representation of our family, such a global section $\bar{\nu}$ gives an element of $H^1(J_{m_0}, \mathbb{C}/\mathbb{Z})$ which is invariant under ρ , or equivalently, an element $\tilde{\nu}_{m_0} \in H^1(J_{m_0}, \mathbb{C})$ satisfying $\rho_{\gamma}(\tilde{\nu}_{m_0}) - \tilde{\nu}_{m_0} \in H^1(J_{m_0}, \mathbb{Z})$ for any $\gamma \in \pi_1(M, m_0)$. As the image of the monodromy representation contains a finite index subgroup of $\text{Sp}(H^1(J_{m_0}, \mathbb{Z}))$, this last condition is satisfied only when $\tilde{\nu}_{m_0} \in H^1(J_{m_0}, \mathbb{Q})$, hence $\bar{\nu}$, and a fortiori ν , is of torsion. $\square$

In [22], [27], the following cohomological interpretation of $\delta\nu$ is given. Writing the exact sequence

$$0 \rightarrow \pi^*\Omega_M \rightarrow \Omega_J \rightarrow \Omega_{J/M} \rightarrow 0, \quad (9)$$

restricting it to the fiber J_m , and using the fact that the connecting map of the associated long exact sequence is the map $\bar{\nabla}$ (a fundamental result due to Griffiths, see [26, Section 5.1.2]), one gets a canonical isomorphism

$$H^1(J_m, \Omega_{J|J_m})^0 \cong (\mathcal{H}_m^{0,1} \otimes \Omega_{M,m})/\bar{\nabla}(\mathcal{H}_m^{1,0}),$$

where on the left $H^1(J_m, \Omega_{J|J_m})^0 \subset H^1(J_m, \Omega_{J|J_m})$ is defined as the kernel of the restriction map

$$H^1(J_m, \Omega_{J|J_m}) \rightarrow H^1(J_m, \Omega_{J_m}).$$

Theorem 2.3. [22] *Let D be a divisor on J which is homologous to 0 along the fibers of π , and ν_D be the associated normal function. One has*

$$\delta\nu_D(m) = [D]_{|J_m}^{1,1} \text{ in } H^1(J_m, \Omega_{J|J_m})^0, \quad (10)$$

where $[D]^{1,1} \in H^1(J, \Omega_J)$ is the Dolbeault cycle class of D , and $[D]_{|J_m}^{1,1}$ is its restriction to J_m .

This result allowed us in [24], [27] to define more generally infinitesimal invariants

$$\delta Z(m) \in H^k(Y_m, \Omega_{Y|Y_m}^k)$$

at any point $m \in M$, for higher codimensional cycles

$$Z \subset Y$$

in a smooth variety Y admitting a smooth fibration $\pi : Y \rightarrow M$ over a base M . These infinitesimal invariants are obtained from the Dolbeault class $[Z]^{k,k} \in H^k(Y, \Omega_Y^k)$, which provides, by restriction to the fibers Y_m , the invariant

$$\delta Z(m) := [Z]_{|Y_m}^{k,k} \in H^k(Y_m, \Omega_{Y|Y_m}^k).$$

In [26, Section 5.2.1], the filtration L on $H^k(Y_m, \Omega_{Y|Y_m}^k)$ induced by the filtration

$$L^p \Omega_Y^k := \pi^* \Omega_M^p \wedge \Omega_Y^{k-p}$$

on Ω_Y^k and its restriction to $\Omega_{Y|Y_m}^k$ is introduced. Note that $Gr_L^p(\Omega_Y^k) \cong \pi^* \Omega_M^p \otimes \Omega_{Y/M}^{k-p}$. This filtration allows to introduce successive infinitesimal invariants living in the graded pieces of $Gr_L^p H^k(Y_m, \Omega_{Y|Y_m}^k)$. The following result is proved in [26, Proposition 5.9] concerning the corresponding spectral sequences

$$\begin{aligned} E_r^{p,q} &\Rightarrow H^{p+q}(Y_m, \Omega_{Y|Y_m}^k), \\ E_r^{p,q} &\Rightarrow R^{p+q} \pi_*(\Omega_Y^k). \end{aligned} \quad (11)$$

Proposition 2.4. *The E_1 -terms of (11)*

$$\dots E_1^{p-1, l-p} \xrightarrow{d_1} E_1^{p, l-p} \xrightarrow{d_1} E_1^{p+1, l-p} \dots$$

identify respectively with the Griffiths complexes

$$\begin{aligned} \dots H^{k-p+1, l-1}(Y_m) \otimes \Omega_{M,m}^{p-1} \xrightarrow{\bar{\nabla}_{\bar{\gamma}}^{-1}} H^{k-p, l}(Y_m) \otimes \Omega_{M,m}^p \xrightarrow{\bar{\nabla}_{\bar{\gamma}}^p} H^{k-p-1, l+1}(Y_m) \otimes \Omega_{M,m}^{p+1} \dots, \\ \dots \mathcal{H}^{k-p+1, l-1} \otimes \Omega_M^{p-1} \xrightarrow{\bar{\nabla}_{\bar{\gamma}}^{-1}} \mathcal{H}^{k-p, l} \otimes \Omega_M^p \xrightarrow{\bar{\nabla}_{\bar{\gamma}}^p} \mathcal{H}^{k-p-1, l+1} \otimes \Omega_M^{p+1} \dots, \end{aligned}$$

where in the second formula $\mathcal{H}^{p,q} := R^q \pi_* \Omega_{Y/M}^p$ and we set $\bar{\nabla}_p(\alpha \otimes \omega) = \bar{\nabla}(\alpha) \wedge \omega$ for any $\alpha \in H^{p,q}(Y_m)$ and $\omega \in \Omega_{M,m}$.

It the paper [24] (see also [27]) where infinitesimal invariants are introduced using this spectral sequence, it degenerates at E_2 for degree reasons, because the infinitesimal variation of Hodge structures considered are that of hypersurfaces, so are supported in only one degree. We could not find a reference for its degeneracy at E_2 in general, so we include here a proof for completeness.

Proposition 2.5. *Let $\pi : Y \rightarrow M$ be a smooth projective morphism, where M is smooth and quasi-projective. Then the spectral sequences (11) degenerate at E_2 .*

Remark 2.6. In this paper, we are going to study infinitesimal invariants for cycles on families $\pi : J \rightarrow M$ of abelian varieties. In this case, proving the degeneracy at E_1 of the spectral sequences (11) is even easier. Indeed, for any integer $n > 1$, let $\mu_n : J \rightarrow J$ be the multiplication by n map. It is a morphism of fibrations over M , and the pull-back morphisms

$$\mu_n^* : \mu_n^* \Omega_J^k \rightarrow \Omega_J^k$$

are compatible with the filtration L , which provide filtered isomorphisms

$$\begin{aligned} \mu_n^* : H^k(J_m, \Omega_{J|J_m}^k) &\rightarrow H^k(J_m, \Omega_{J|J_m}^k) \\ \mu_n^* : R^k \pi_* \Omega_J^k &\rightarrow R^k \pi_* \Omega_J^k. \end{aligned}$$

The morphisms μ_n^* induce morphisms of spectral sequences (11) for J , and by Proposition 2.4, these morphisms act as $n^{k-p+l} Id$ on $E_1^{p, l-p}$. It thus follows that they also act as $n^{k-p+l} Id$ on $E_r^{p, l-p}$, for any $r \geq 1$. But then, the differentials

$$d_r : E_r^{p, l-p} \rightarrow E_r^{p+r, l-p-r+1}$$

vanish for $r \geq 2$ since they are compatible with the μ_n -action and μ_n^* acts by different powers of n on both sides for $r \geq 2$.

Proof of Proposition 2.5. The reasoning is similar to the one presented above, except that we need a substitute for the μ_n^* . This is done using the relative Künneth projectors constructed as follows. Let d be the relative dimension of π . Consider $Y_2 := Y \times_M Y \xrightarrow{\pi'} M$, with projections $\pi_1, \pi_2 : Y_2 \rightarrow Y$, which are both smooth projective morphisms of relative dimension d . The diagonal $\Delta_Y \subset Y_2$ has a cohomology class

$$\delta_Y \in H^{2d}(Y_2, \mathbb{Q})$$

which is algebraic hence extends to a Hodge class on a given projective compactification $\bar{Y}_2$ of Y_2 . It induces a relative action

$$\begin{aligned} \delta_{Y,*} : R^j \pi_{1*} \mathbb{Q} &\rightarrow R^j \pi_{2*} \mathbb{Q}, \\ \delta_{Y,*}(\alpha) &= \pi_{2*}(\pi_1^* \alpha \cup \delta_Y) \end{aligned}$$

which is nothing but the identity. The class δ_Y induces a global section of

$$R^{2d} \pi'_* \mathbb{Q} = \oplus_{j+i=2d} R^j \pi_* \mathbb{Q} \otimes R^i \pi_* \mathbb{Q} = \oplus_i \text{Hom}(R^i \pi_* \mathbb{Q}, R^i \pi_* \mathbb{Q}). \quad (12)$$

Using (12), we can define the global section $\delta_{Y,i}$ of the local system $R^{2d} \pi'_* \mathbb{Q}$ as the one which acts as the identity on $R^i \pi_* \mathbb{Q}$ and by 0 on $R^j \pi_* \mathbb{Q}$ for $j \neq i$. The classes $\delta_{Y,i}(m)$ are Hodge classes at any point $m \in M$ which are not known to be algebraic but by Deligne's global invariant cycles theorem [11], see also [26, Theorem 4.24], there exists for any i a Hodge class on $\bar{Y}_2$, such that its restriction δ_i to Y_2 induces the section $\delta_{Y,i} \in H^0(M, R^{2d} \pi'_* \mathbb{Q})$. The class $\delta_i \in H^{2d}(Y_2, \mathbb{Q})$ (which is what we call a relative Künneth projector) is thus constructed so that

$$\delta_{i*}(\alpha) := \pi_{2*}(\pi_1^* \alpha \cup \delta_i) : R^j \pi_* \mathbb{Q} \rightarrow R^j \pi_* \mathbb{Q} \quad (13)$$

is the identity for $j = i$ and 0 otherwise. As δ_i is the restriction of a Hodge class on $\bar{Y}_2$, there is a Dolbeault version

$$\delta_i^{d,d} \in H^d(Y_2, \Omega_{Y_2}^d)$$

of δ_i . Let us choose a Dolbeault representative $\tilde{\delta}_i^{d,d}$ of $\delta_i^{d,d}$, given by a closed (d, d) -form on Y_2 .

Lemma 2.7. (i) For each k , the $\bar{\partial}$ -closed forms $\tilde{\delta}_i^{d,d}$ act on the push-forward to M of the Dolbeault complex $(\mathcal{A}^{0,*}(\Omega_Y^k), \bar{\partial}_Y)$ of Ω_Y^k , whose cohomology computes $R^* \pi_* \Omega_Y^k$. This action is compatible with the filtration L on Ω_Y^k .

(ii) The induced action of $\tilde{\delta}_i^{d,d}$ on $E_1^{p,l-p} = \mathcal{H}^{k-p,l} \otimes \Omega_M^p$ is by the identity if $k-p+l = i$ and by 0 otherwise.

(iii) Similar statement at any point $m \in M$.

Proof. The action of $\tilde{\delta}_i^{d,d}$ on the relative Dolbeault complex of Ω_Y is given by

$$\tilde{\delta}_{i*}(\alpha) = \pi_{2*}(\pi_1^* \alpha \cup \tilde{\delta}_i), \quad (14)$$

where $\pi_{2*} : A^{p,q}(Y_2/M) \rightarrow A^{p-d,q-d}(Y/M)$ is given by integration along the fibers. As the three operations π_1^* , $\cup \tilde{\delta}_i$ and π_{2*} commute with the Dolbeault operator $\bar{\partial}$ and preserve the L -filtration, so does $\tilde{\delta}_{i*}$. The $\tilde{\delta}_{i*}$ thus provide morphisms of filtered complexes quasi-isomorphic to Ω_Y^k , hence morphisms of the corresponding spectral sequences. This proves (i). Concerning (ii), the action of $\tilde{\delta}_{i*}$ on $E_1^{p,l-p} = \mathcal{H}^{k-p,l} \otimes \Omega_M^p$ is compatible with its action of $\mathcal{H}^{k-p,l}$ (and is the identity on Ω_M^p), and the latter is also compatible with the action of $\tilde{\delta}_{i*}$ on $R^{k-p+l} \pi_* \mathbb{C}$ given by the same formula as (14), which makes sense since we chose $\tilde{\delta}_{i*}$ to be a closed form. This latter action is the action of δ_{i*} given by (13), hence it vanishes if $k-p+l \neq i$, and it is the identity if $k-p+l = i$, which proves (ii). $\square$

By Lemma 2.7(i), the form $\sum_i \tilde{\delta}_i$ acts on the L -filtered relative Dolbeault complex of Ω_Y^k , hence on the whole spectral sequences (11). Furthermore by Lemma 2.7(ii), it acts by the identity on the E_1 -term, hence on all E_r -terms. In order to prove that $d_r = 0$ for $r \geq 2$, it thus suffices to prove that $d_r \circ \tilde{\delta}_{i*} = 0$ for all $r \geq 2$. This is now obvious for degree reason by Lemma 2.7(ii). $\square$

A generalization of Mumford's theorem (see [24], [25]) is as follows.

Theorem 2.8. *Let $\pi : Y \rightarrow M$ be a smooth projective morphism, and let $Z \in \text{CH}^k(Y)$ satisfy the condition that for any fiber $m \in M$, $Z|_{Y_m} = 0$ in $\text{CH}^k(Y_m)$. Then, for a general point $m \in M$, $\delta Z(m) = 0$ in $H^k(Y_m, \Omega_{Y|Y_m}^k)$.*

Indeed, under these assumptions, a multiple NZ is rationally equivalent to 0 on Y_U , for some dense Zariski open set $U \subset M$ (see [8], [25]). A fortiori, one has $[Z]_{Y_U}^{k,k} = 0$ in $H^k(Y_U, \Omega_{Y_U}^k)$ and so $\delta Z(m) = 0$ for any $m \in U$, by restriction to Y_m . Note that the vanishing of $\delta Z(m)$ may not hold everywhere, due to the fact that $R^k \pi_* \Omega_Y^k$ is not torsion free in general.

The proofs of Theorems 1.2 and 1.8 rely on Theorem 2.8 applied to the case of codimension 2 cycles on an abelian fibration $J \rightarrow M$. More precisely, the cycles Z that we are going to consider are of the form $Z = D^2$, for some divisor D on J . The compatibility of the cycle class with cup-product and intersection product gives the equality

$$[Z]^{2,2} = ([D]^{1,1})^2 \text{ in } H^2(J, \Omega_J^2). \quad (15)$$

Restricting this equalities along the fibers of π , one gets that δZ is computed explicitly as the square of δD , although the square map

$$\text{Sym}^2 H^1(J_m, \Omega_{J|J_m}) \rightarrow H^2(J_m, \Omega_{J|J_m}^2)$$

is not so easy to analyze. This analysis was started in [20] and our results rely on the analysis made in *loc. cit.*. The paper [20] also computes the infinitesimal invariant for the Griffiths-Pirola normal function γ_P (defined over a double cover $\widetilde{\mathcal{M}}_4$ of $\mathcal{M}_4$), and for its square. Let $\tilde{\mathcal{C}} \rightarrow \widetilde{\mathcal{M}}_4$ be the pull-back of the universal curve of genus 4 as in the introduction, and let $\tilde{\mathcal{C}}^{(2)}/\widetilde{\mathcal{M}}_4 \rightarrow \widetilde{\mathcal{M}}_4$ be its relative second symmetric product. The normal function γ_P provides as well a divisor D_{γ_P} on $\tilde{\mathcal{C}}^{(2)}$, which is cohomologous to 0 along fibers.

The following results proved in [20] will be used in Section 4. Theorem 2.9 strengthens an earlier result by Griffiths [13], and fully computes the infinitesimal invariant $\delta\gamma_{P,[C]}$ for a general curve C of genus 4. Let C be a general curve of genus 4. Let

$$V := H^{1,0}(C), \quad V_2 := H^0(C, K_C^{\otimes 2}) = \text{Sym}^2 V/q,$$

where $q \in \text{Sym}^2 V$ is nondegenerate and defines the unique quadric Q containing C in its canonical embedding. The space of infinitesimal invariants at $[C]$ for normal functions $\nu \in \Gamma(\mathcal{J}^\vee)$ defined on a neighborhood of $[C] \in \mathcal{M}_4$ is

$$I_{1,V} = \text{Hom}(V, V_2)/V$$

where the inclusion $V \hookrightarrow \text{Hom}(V, V_2)$ maps $v \in V$ to the multiplication map by v . The space $I_{1,V}$ is a representation of the group $\text{Aut}(V, q)$, and, as such, splits naturally as the direct sum of $H^0(Q, \mathcal{O}_Q(3))$ and $H^0(Q, T_Q(1))$.

Theorem 2.9. *[20, Theorem 1.2] The infinitesimal invariant $\delta\gamma_P$ at $[C]$ (which is well-defined up to sign) belongs to $H^0(Q, \mathcal{O}_Q(3)) \subset I_{1,V}$. It is a nonzero multiple of the cubic equation defining $C \subset Q$.*

Theorem 2.10. *[20, Theorem 1.3] One has $(\delta\gamma_P)^2 = \delta(D_{\gamma_P}^2) \neq 0$ in $H^2(J_m, \Omega_{J|J_m}^2)$ at a general point $m \in \widetilde{\mathcal{M}}_4$, hence for a very general point $m \in \widetilde{\mathcal{M}}_4$, one has $D_{\gamma_P}^2 \neq 0$ in $\text{CH}_0(C^{(2)})$.*

The second statement in Theorem 2.10 follows from the first by Theorem 2.8.

Remark 2.11. Theorem 2.10 is also implied by Theorem 1.2(ii). However the proof in [20] is much more direct, because it directly proves by an explicit computation the nonvanishing of the infinitesimal invariant $\delta(D_{\gamma_P}^2)$, hence uses none of the ingredients of Section 3, apart of course for the general theory of infinitesimal invariants.

Remark 2.12. The spaces of infinitesimal invariants for codimension 2 cycles on $J(C)$ and on $C^{(2)}$ coincide. So the (non)vanishing properties of the infinitesimal invariants for $C^{(2)}$ and for $J(C)$ are equivalent, which is coherent with Lemma 1.1. However, it is important to also consider in the course of the paper cycles on $C^{(2)}$ because the Chow group of codimension 2 cycles on $C_\eta^{(2)}$, where C_η is the generic curve of genus 4 is well understood, see for example [21].

3 Proof of Theorem 1.2

3.1 The equation $\overline{M}_2(\overline{\phi}) = 0$

Let V be a g -dimensional vector space, with $g \geq 4$. We denote by W_2 the vector space $\text{Sym}^2 V$. If $g = 4$, given a nondegenerate quadratic form q on V , we denote by V_2 the vector space $\text{Sym}^2 V/q$. If A is a principally polarized abelian variety of dimension g , and $V = H^{1,0}(A)$, W_2 is the cotangent space at $[A]$ to the moduli stack $\mathcal{A}_g$ (or to the base of the Kuranishi family) of polarized deformations of A . If $A = J(C)$, where C is a non-hyperelliptic curve of genus 4, and q is the unique quadratic form vanishing on C in its canonical embedding, q is nondegenerate if C is general, and $V_2 = H^0(C, 2K_C)$ is the cotangent space at $[C]$ to the moduli stack $\mathcal{M}_4$ of deformations of C .

Let

$$I_{1,W} := \text{Hom}(V, W_2)/V, \text{ resp. } I_{1,V} := \text{Hom}(V, V_2)/V, \quad (16)$$

where, in both cases, V is naturally embedded in $\text{Hom}(V, W_2)$, resp. $\text{Hom}(V, V_2)$, via the multiplication maps. Let $J_g \rightarrow \mathcal{A}_g$ be the universal principally polarized abelian variety of dimension g and let U be an (analytic or étale) open set of $\mathcal{A}_g$. Let D be a divisor on $J_{g,U}$, which is homologous to 0 on fibers of π . As we saw in Section 2, using the exact sequence

$$0 \rightarrow \Omega_{U,m} \otimes \mathcal{O}_{J_{g,m}} \rightarrow \Omega_{J_g|J_{g,m}} \rightarrow \Omega_{J_{g,m}} \rightarrow 0 \quad (17)$$

at any point $m \in U$, the infinitesimal invariant $\delta D(m) = \delta \nu_{D,m}$ at $m \in U$ belongs to

$$H^1(J_{g,m}, \Omega_{J_g|J_{g,m}})^0 = (H^{0,1}(J_{g,m}) \otimes \Omega_{U,m})/\overline{\nabla}(H^{1,0}(J_{g,m})),$$

and the right hand side is naturally isomorphic to $I_{1,W}$, with $V = H^{1,0}(J_{g,m})$.

Similarly, if $\pi : J \rightarrow \mathcal{M}_4$ is the universal Jacobian fibration of the universal family of curves $\mathcal{C} \rightarrow \mathcal{M}_4$ of genus 4, and V_2 is as above, the space

$$I_{1,V} \cong H^1(\Omega_{J|J_m})^0 \cong H^1(\Omega_{\mathcal{C}|C})^0,$$

where $C \cong \mathcal{C}_m$, is the space of infinitesimal invariants at $m = [C]$ for normal functions $\nu_D \in \Gamma(\mathcal{J}^\vee)$ defined on open (étale or analytic) sets of $\mathcal{M}_4$.

For any $\phi, \psi \in V^* \otimes W_2$, we denote by $\phi \cdot \psi \in \text{Sym}^2(V^* \otimes W_2)$ the image of $\phi \otimes \psi \in (V^* \otimes W_2)^{\otimes 2}$. Similarly, recalling that V embeds naturally in $\text{Hom}(V, W_2) = V^* \otimes W_2$, we denote by $V \cdot (V^* \otimes W_2) \subset \text{Sym}^2(V^* \otimes W_2)$ the subspace generated by elements $\phi \cdot \psi$, where $\phi \in V$. The quotient

$$\text{Sym}^2(V^* \otimes W_2)/V \cdot (V^* \otimes W_2)$$

is naturally isomorphic to $\text{Sym}^2((V^* \otimes W_2)/V) = \text{Sym}^2 I_{1,W}$.

The 2×2 -minors map

$$V^* \otimes W_2 \ni \phi \mapsto M_{2,W}(\phi) := \wedge^2 \phi \in \bigwedge^2 V^* \otimes \bigwedge^2 W_2$$

extends by linearity to a morphism

$$M_{2,W} : \text{Sym}^2(V^* \otimes W_2) \rightarrow \bigwedge^2 V^* \otimes \bigwedge^2 W_2.$$

Let

$$I_{2,W} := (\bigwedge^2 V^* \otimes \bigwedge^2 W_2) / M_{2,W}(V \cdot (V^* \otimes W_2)). \quad (18)$$

We denote by

$$\overline{M}_{2,W} : \text{Sym}^2((V^* \otimes W_2)/V) = \text{Sym}^2 I_{1,W} \rightarrow I_{2,W} \quad (19)$$

the natural factorization.

In the second situation, where $\dim V = 4$ and we work with $V_2 = W_2/q$, we will use the similarly defined notation $M_{2,V} : \text{Sym}^2(V^* \otimes V_2) \rightarrow \bigwedge^2 V^* \otimes \bigwedge^2 V_2$, and

$$I_{2,V} := (\bigwedge^2 V^* \otimes \bigwedge^2 V_2) / M_{2,V}(V \cdot (V^* \otimes V_2)). \quad (20)$$

As before, we also have the induced morphism

$$\overline{M}_{2,V} : \text{Sym}^2((V^* \otimes V_2)/V) = \text{Sym}^2 I_{1,V} \rightarrow I_{2,V}. \quad (21)$$

Coming back to the first situation, we have as explained in Section 2 a filtration $L^\bullet$ on $H^2(\Omega_{J_g|J_{g,m}}^2)$. Given a divisor D homologous to 0 on fibers of π on J_g , the infinitesimal invariant

$$\delta(D^2) = (\delta D)^2$$

clearly belongs to the deepest part

$$L^2 H^2(\Omega_{J_g|J_{g,m}}^2) \subset H^2(\Omega_{J_g|J_{g,m}}^2)$$

for this filtration. This deepest part is computed using Propositions 2.4 and 2.5 as

$$L^2 H^2(\Omega_{J_g|J_{g,m}}^2) \cong (H^{0,2}(J_{g,m}) \otimes \Omega_{U,m}^2) / \overline{\nabla}_1(H^{1,1}(J_{g,m}) \otimes \Omega_{U,m}), \quad (22)$$

where $\overline{\nabla}_1 : H^{1,1}(J_{g,m}) \otimes \Omega_{U,m} \rightarrow H^{0,2}(J_{g,m}) \otimes \Omega_{U,m}$ is given by

$$\overline{\nabla}_1(\omega \otimes \alpha) = \overline{\nabla}(\omega) \wedge \alpha.$$

Comparing (22) and (18), one gets (see [20] for more detail) that the space $L^2 H^2(\Omega_{J_g|J_{g,m}}^2)$ is naturally isomorphic to the space $I_{2,W}$. It is also proved in [20] that the linear map

$$\text{Sym}^2 H^1(\Omega_{J_g|J_{g,m}})^0 \rightarrow L^2 H^2(\Omega_{J_g|J_{g,m}}^2)$$

induced by the square map

$$H^1(\Omega_{J_g|J_{g,m}})^0 = L^1 H^1(\Omega_{J_g|J_{g,m}}) \ni \delta D \mapsto (\delta D)^2 \in L^2 H^2(\Omega_{J_g|J_{g,m}}^2)$$

is nothing but the product $\overline{M}_{2,W}$ of (19). Similarly, in the second situation where we work over open sets U of $\mathcal{M}_4$, with tangent space V_2 , and $J \rightarrow \mathcal{M}_4$ is the universal Jacobian, the

space $L^2 H^2(\Omega_{J|J_m}^2)$ is naturally isomorphic to the space $I_{2,V}$ of (20) and it is proved in [20, Proposition 5.1] that the linear map

$$\mathrm{Sym}^2 H^1(\Omega_{J|J_m})^0 \rightarrow L^2 H^2(\Omega_{J|J_m}^2)$$

induced by the square map

$$H^1(\Omega_{J|J_m})^0 = L^1 H^1(\Omega_{J|J_m}) \ni \delta D \mapsto (\delta D)^2 \in L^2 H^2(\Omega_{J|J_m}^2)$$

is nothing but the product $\overline{M}_{2,V}$ of (21).

Proposition 3.1. *The notation being as above, the following hold:*

(i) *Assume $g \geq 4$. If $\overline{\phi} \in I_{1,W} = (V^* \otimes W_2)/V$ has the property that*

$$\overline{M}_{2,W}(\overline{\phi}) = 0 \text{ in } (\bigwedge^2 V^* \otimes \bigwedge^2 W_2)/M_2(V \cdot (V^* \otimes W_2)) = I_{2,W},$$

there exists a unique $\phi \in V^ \otimes W_2$ mapping to $\overline{\phi}$, such that*

$$M_2(\phi) = 0 \text{ in } \bigwedge^2 V^* \otimes \bigwedge^2 W_2,$$

that is, ϕ has rank ≤ 1 .

(ii) *Assume $g = 4$. If $\overline{\phi} \in I_{1,V} = (V^* \otimes V_2)/V$ has the property that*

$$\overline{M}_{2,V}(\overline{\phi}) = 0 \text{ in } (\bigwedge^2 V^* \otimes \bigwedge^2 V_2)/M_2(V \cdot (V^* \otimes V_2)) = I_{2,V},$$

there exists a unique $\phi \in V^ \otimes V_2$ mapping to $\overline{\phi}$, such that*

$$M_2(\phi) = 0 \text{ in } \bigwedge^2 V^* \otimes \bigwedge^2 V_2,$$

that is, ϕ has rank ≤ 1 .

In both cases, the uniqueness follows from the assumption $g \geq 4$ (in fact $g \geq 3$ would suffice here). Indeed, assume there are two lifts $\phi, \phi' \in V^* \otimes W_2$, resp. $\phi, \phi' \in V^* \otimes V_2$ of $\overline{\phi} \in I_{1,W}$ (resp. $\overline{\phi} \in I_{1,V}$) that have rank ≤ 1 . Then the difference $\phi - \phi'$ has rank ≤ 2 , and belongs to $V \subset \mathrm{Hom}(V, W_2)$ (resp. to $V \subset \mathrm{Hom}(V, V_2)$). This contradicts the fact that, for any $0 \neq v \in V$, the map $\mu_v : V \rightarrow W_2$, (resp. $\mu_v : V \rightarrow V_2$) of multiplication by v is injective, hence has rank g (resp. rank 4).

The unicity being proved, we concentrate below on the existence of the rank 1 lifts.

Proof of Proposition 3.1(i). For any 2-dimensional vector subspace $S \subset V$, let $E := V/S$. We have two restriction maps $V^* \rightarrow S^*$ and $W_2 \rightarrow \mathrm{Sym}^2 E$, which together induce

$$r_S : V^* \otimes W_2 \rightarrow S^* \otimes \mathrm{Sym}^2 E.$$

We use the following observation made in [20, Lemma 6.1].

Lemma 3.2. *For any $v \in V$, with induced morphism*

$$\mu_v \in \mathrm{Hom}(V, W_2) = V^* \otimes W_2$$

of multiplication by v , we have $r_S(\mu_v) = 0$ in $S^ \otimes \mathrm{Sym}^2 E$.*

From the restriction maps $V^* \rightarrow S^*$ and $W_2 \rightarrow \text{Sym}^2 E$, we also deduce a restriction morphism

$$r'_S : \bigwedge^2 V^* \otimes \bigwedge^2 W_2 \rightarrow \bigwedge^2 S^* \otimes \bigwedge^2 (\text{Sym}^2 E),$$

which is compatible with r_S and the 2×2 -minors maps on both sides. We then get from Lemma 3.2 that r'_S vanishes on $M_{2,W}(V \cdot (V^* \otimes W_2))$. Hence, using formula (18) defining $I_{2,W}$, we deduce that, for any vector subspace $S \subset V$ of dimension 2, there is a linear map

$$r_{S,2} : I_{2,W} \rightarrow \text{Hom}(\bigwedge^2 S, \bigwedge^2 (\text{Sym}^2 E)) \quad (23)$$

obtained by factorization of r'_S , such that, for any $\bar{\phi} \in V^* \otimes W_2/V$, we have

$$r_{S,2}(\bar{M}_2(\bar{\phi})) = M_{2,S}(r_S(\bar{\phi})), \quad (24)$$

where the map $M_{2,S}$ on the right is the 2×2 -minor map for $\text{Hom}(S, \text{Sym}^2 E)$.

Let $G = G(2, V)$ be the Grassmannian of 2-dimensional vector subspaces of V . Let $\mathcal{S} \subset V \otimes \mathcal{O}_G$ be the tautological rank 2 vector subbundle, and $\mathcal{E} := V \otimes \mathcal{O}_G / \mathcal{S}$ the rank $g - 2$ tautological quotient bundle. Lemma 3.2 provides a linear map

$$r_1 : (V^* \otimes W_2)/V \rightarrow H^0(G, \text{Hom}(\mathcal{S}, \text{Sym}^2 \mathcal{E})). \quad (25)$$

We also have the 2×2 -minors quadratic map

$$\begin{aligned} m_2 : \text{Hom}(\mathcal{S}, \text{Sym}^2 \mathcal{E}) &\rightarrow \text{Hom}(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E})) \\ m_2(\psi) &:= \wedge^2 \psi. \end{aligned} \quad (26)$$

We will also denote by m_2 the map induced by (26) at the level of global sections. By (23), we get a linear map

$$r_2 : I_{2,W} \rightarrow H^0(G, \text{Hom}(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E}))) \quad (27)$$

such that, for any $\bar{\phi} \in (V^* \otimes W_2)/V$, we have as in (24)

$$r_2(\bar{M}_2(\bar{\phi})) = m_2(r_1(\bar{\phi})) \text{ in } H^0(G, \text{Hom}(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E}))). \quad (28)$$

We now prove two lemmas.

Lemma 3.3. *For $g \geq 4$, the morphism r_1 of (25) is injective.*

Proof. Let $\phi \in V^* \otimes W_2$ such that $r_1(\bar{\phi}) = 0$ in $H^0(G, \text{Hom}(\mathcal{S}, \text{Sym}^2 \mathcal{E}))$. We note that, given a quotient $E = V/S$, the kernel of the natural map $\text{Sym}^2 V \rightarrow \text{Sym}^2 E$ maps naturally to $S \otimes E$, with kernel isomorphic to $\text{Sym}^2 S$. By our assumption, ϕ thus induces a section

$$s \in H^0(G, \text{Hom}(\mathcal{S}, \mathcal{S} \otimes \mathcal{E})).$$

We now write the exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow V \otimes \mathcal{O}_G \rightarrow \mathcal{E} \rightarrow 0$$

on G , tensor it by $\mathcal{E}nd(\mathcal{S})$, and get by taking sections the long exact sequence

$$\dots \rightarrow H^0(G, \mathcal{E}nd(\mathcal{S}) \otimes V) \rightarrow H^0(G, \text{Hom}(\mathcal{S}, \mathcal{S} \otimes \mathcal{E})) \rightarrow H^1(G, \mathcal{E}nd(\mathcal{S}) \otimes \mathcal{S}) \rightarrow \dots$$

The vanishing of $H^1(G, \mathcal{E}nd(\mathcal{S}) \otimes \mathcal{S})$ and the fact that $H^0(G, \mathcal{E}nd(\mathcal{S})) = \mathbb{C}\text{Id}$ (thanks to the assumption $g \geq 4$) thus imply that s comes from an element v of V . Thus the image of $\phi - \mu_v$ in $H^0(G, \text{Hom}(\mathcal{S}, \text{Sym}^2 V \otimes \mathcal{O}_G))$ belongs to $H^0(G, \text{Hom}(\mathcal{S}, \text{Sym}^2 \mathcal{S}))$, which is 0 because $g \geq 3$. This implies that $\phi = \mu_v$. $\square$

Lemma 3.4. *If $0 \neq \alpha \in H^0(G, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E}))$ satisfies*

$$m_2(\alpha) = 0 \text{ in } H^0(G, \mathcal{H}om(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E}))), \quad (29)$$

then there exists a rank 1 element $\psi \in V^ \otimes W_2$ whose image in $H^0(G, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E}))$ equals α .*

Proof. The condition $m_2(\alpha) = 0$ means that α has everywhere rank ≤ 1 . Thus, if $\alpha \neq 0$, there is a line bundle $\mathcal{L}$ on G and two nonzero morphisms

$$\beta : \mathcal{S} \rightarrow \mathcal{L}, \quad i : \mathcal{L} \rightarrow \text{Sym}^2 \mathcal{E}, \quad (30)$$

such that

$$\alpha = i \circ \beta : \mathcal{S} \rightarrow \text{Sym}^2 \mathcal{E}. \quad (31)$$

The line bundle $\mathcal{L} \in \text{Pic}(G)$ is a power of the Plücker line bundle. It cannot be strictly positive, because the existence of the morphism $i \neq 0$ shows that $H^0(G, \text{Sym}^2 \mathcal{E} \otimes \mathcal{L}^{-1}) \neq 0$. The line bundle $\mathcal{L}$ cannot be strictly negative because $H^0(G, \mathcal{S}^* \otimes \mathcal{L}) \neq 0$. It follows that $\mathcal{L} = \mathcal{O}_G$. Using (30) with $\mathcal{L}$ trivial, (31) gives that the morphism $\alpha \in H^0(G, \mathcal{S}^* \otimes \text{Sym}^2 \mathcal{E})$ can be written as $u \otimes v$, where $u \in H^0(G, \mathcal{S}^*) = V$, and $v \in H^0(G, \text{Sym}^2 \mathcal{E}) = \text{Sym}^2 H^0(G, \mathcal{E}) = W_2$. Thus Lemma 3.4 is proved. $\square$

Using formula (24), Lemma 3.4 shows that for $\bar{\phi} \in V^* \otimes W_2 / V$, the condition $\bar{M}_{2,W}(\bar{\phi}) = 0$ implies that there exists a rank 1 element $\psi \in V^* \otimes W_2$ whose image $\bar{\psi} \in (V^* \otimes W_2) / V$ satisfies $r_1(\bar{\phi} - \bar{\psi}) = 0$. Lemma 3.3 then gives that $\bar{\phi} = \bar{\psi}$, which proves Proposition 3.1(i). $\square$

Proof of Proposition 3.1(ii). The proof is very similar. We now work with $V_2 = \text{Sym}^2 V / q = W_2 / q$, and with the space of infinitesimal invariants

$$I_{2,V} := (\bigwedge^2 V^* \otimes \bigwedge^2 V_2) / M_{2,V}(V \cdot (V^* \otimes V_2)). \quad (32)$$

We now study the linear map

$$\bar{M}_{2,V} : \text{Sym}^2((V^* \otimes V_2) / V) \rightarrow I_{2,V}$$

of (21), obtained by factorization of the 2×2 -minor map

$$M_{2,V} : \text{Sym}^2(V^* \otimes V_2) \rightarrow \bigwedge^2 V^* \otimes \bigwedge^2 V_2.$$

The other adjustments that we have to make are the following: using Lemma 3.2, we now get the following analogue

$$r'_1 : (V^* \otimes V_2) / V \rightarrow H^0(G, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E} / q)) \quad (33)$$

of r_1 , where we use the notation q for its natural image in $\text{Sym}^2 \mathcal{E}$ under restriction. The coherent sheaf $\text{Sym}^2 \mathcal{E} / q$ is not locally free anymore on G , but it is locally free on the open set $G^0 \subset G$ parameterizing lines not contained in $\{q = 0\}$. We recall that we have now $g = 4$ so the rank of $\mathcal{E}$ is 2. In particular, the codimension of $G \setminus G^0$ in G is 3. On G^0 , the 2×2 -minors map m_2 of (26) is now replaced with the quadratic map

$$\begin{aligned} m'_2 : \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E} / q) &\rightarrow \mathcal{H}om(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E} / q)), \\ m'_2(\psi) &:= \wedge^2 \psi. \end{aligned} \quad (34)$$

(We will use the same notation m'_2 for the induced map at the level of global sections). We also get a linear map

$$r'_2 : I_{2,V} \rightarrow H^0(G^0, \mathcal{H}om(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E}/q))) \quad (35)$$

such that, for any $\bar{\phi} \in (V^* \otimes V_2)/V$,

$$r'_2(\bar{M}_{2,V}(\bar{\phi})) = m'_2(r'_1(\bar{\phi})) \text{ in } H^0(G^0, \mathcal{H}om(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E}/q))). \quad (36)$$

We now prove two lemmas:

Lemma 3.5. *The morphism r'_1 of (33) is injective.*

Proof. Let $\phi \in V^* \otimes W_2$ be such that its image $\bar{\phi} \in I_{1,V}$ satisfies $r'_1(\bar{\phi}) = 0$. Denoting by ψ the image of ϕ in $(V^* \otimes W_2)/V$, this implies that the morphism

$$r_1(\psi) \in H^0(G, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E}))$$

has its image contained in $q\mathcal{O}_G \subset \text{Sym}^2 \mathcal{E}$, hence has everywhere rank ≤ 1 . By Lemma 3.4, there exists a rank 1 element $w \in V^* \otimes W_2$ such that its image in $H^0(G, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E}))$ equals $r_1(\psi)$. One has $w = v \otimes q'$ for some element $q' \in S^2 V$, and in fact q' must be proportional to q since it becomes proportional to q in $\text{Sym}^2 E$, for a general quotient E of V of dimension 2. Hence one concludes by Lemma 3.3 that, modulo $V \subset V^* \otimes W_2$, ψ belongs to $V^* \otimes q \subset V^* \otimes W_2$. So modulo q , ψ belongs to $V \subset V^* \otimes V_2$ and the lemma is proved. $\square$

Lemma 3.6. *If $\alpha \in H^0(G^0, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E}/q))$ satisfies*

$$m'_2(\alpha) = 0 \text{ in } H^0(G^0, \mathcal{H}om(\bigwedge^2 \mathcal{S}, \bigwedge^2 (\text{Sym}^2 \mathcal{E}/q))), \quad (37)$$

then there exists a rank 1 element $\psi \in V^ \otimes V_2$ whose image in $H^0(G^0, \mathcal{H}om(\mathcal{S}, \text{Sym}^2 \mathcal{E}/q))$ equals α .*

Proof. Our assumption is that $\alpha : \mathcal{S} \rightarrow \text{Sym}^2 \mathcal{E}/q$ has rank ≤ 1 everywhere, hence, for some line bundle $\mathcal{L}$ on G , it factors as $i \circ \beta$, where

$$\beta : \mathcal{S} \rightarrow \mathcal{L}$$

is defined on G^0 and $i : \mathcal{L} \rightarrow \text{Sym}^2 \mathcal{E}/q$ is an injective morphism. As in the proof of Lemma 3.4, we discuss the possible values of $\mathcal{L}$. It cannot be strictly negative since

$$H^0(G^0, \mathcal{S}^* \otimes \mathcal{L}) = H^0(G, \mathcal{S}^* \otimes \mathcal{L})$$

is nonzero, because it contains β . Next, $H^0(G^0, \mathcal{L}^{-1} \otimes (\text{Sym}^2 \mathcal{E}/q)) \neq 0$ because it contains i . It follows that

$$H^0(G^0, \mathcal{L}^{-1} \otimes \text{Sym}^2 \mathcal{E}) \neq 0. \quad (38)$$

Indeed, $H^1(G^0, \mathcal{L}^{-1}) = 0$, because $\text{codim}(G \setminus G^0 \subset G) \geq 3$, so the claim follows from the exact sequence

$$0 \rightarrow \mathcal{L}^{-1} \rightarrow \mathcal{L}^{-1} \otimes \text{Sym}^2 \mathcal{E} \rightarrow \mathcal{L}^{-1} \otimes (\text{Sym}^2 \mathcal{E}/q) \rightarrow 0$$

on G^0 . As

$$H^0(G^0, \mathcal{L}^{-1} \otimes \text{Sym}^2 \mathcal{E}) = H^0(G, \mathcal{L}^{-1} \otimes \text{Sym}^2 \mathcal{E}),$$

we conclude from (38) that $H^0(G, \mathcal{L}^{-1} \otimes \text{Sym}^2 \mathcal{E}) \neq 0$ and thus $\mathcal{L}$ cannot be strictly positive. Thus the line bundle $\mathcal{L}$ is trivial, and we conclude exactly as in the proof of Lemma 3.4, using now the facts that $H^0(G, \mathcal{S}^*) = V^*$ and $H^0(G^0, \text{Sym}^2 \mathcal{E}/q) = W_2/q = V_2$. $\square$

The proof of Proposition 3.1(ii) follows from these two lemmas exactly as before. $\square$

3.2 Proof of Theorem 1.2

We recall that Theorem 1.2 is the following statement.

Theorem 3.7. (i) *Let A be a very general abelian variety of dimension at least 4. Then, if $D \in \text{CH}^1(A)_{\text{hom}} = \text{Pic}^0(A)$ satisfies $D^2 = 0$ in $\text{CH}^2(A)$, D is of torsion in $\text{Pic}^0(A)$.*

(ii) *Let C be a very general curve of genus 4. Then if $D \in \text{Pic}^0(C) = \text{Pic}^0(J(C)) = \text{CH}^1(J(C))$ satisfies $D^2 = 0$ in $\text{CH}^2(J(C))$, D is of torsion in $\text{Pic}^0(C)$.*

The proofs of Theorem 3.7(i) and Theorem 3.7(ii) are very similar, with some technical differences that are essentially already taken care in Proposition 3.1(i) and (ii), so we give them in parallel. When a different argument is needed for (i) and (ii), we will refer to case (i) and case (ii).

In case (i), we consider the moduli space $\mathcal{A}_g$ of principally polarized abelian varieties, and consider the universal abelian variety $J_g \rightarrow \mathcal{A}_g$ with associated (isomorphic) dual fibration

$$\widehat{J}_g := \text{Pic}^0(J/\mathcal{A}_g) \rightarrow \mathcal{A}_g.$$

In case (ii), we consider the moduli space $\mathcal{M}_4$ of genus 4 curves, the universal Jacobian $J \rightarrow \mathcal{M}_4$ and its dual $\widehat{J} \rightarrow \mathcal{M}_4$.

We start recalling the following general fact (see [25]).

Lemma 3.8. *For any abelian fibration $J \rightarrow B$, the set of points (b, d) , $d \in \widehat{J}_b = \text{Pic}^0(J_b) = \text{CH}^1(J_b)_{\text{hom}}$, such that $d^2 = 0$ in $\text{CH}^2(J_b)$, is a countable union of closed algebraic subsets of $\widehat{J}$.*

This set contains the torsion points in the fibers, by the Bloch-Roitman theorem on torsion codimension 2 algebraic cycles (see [7], [18]), because the cycles d^2 above have trivial Abel-Jacobi invariant. In case (i), it is also known for $g \geq 3$ (see [23]) that any irreducible algebraic component of this set dominating $\mathcal{A}_g$ is generically finite over it. We fix such a dominating component M and denote respectively by $J_{g,M} \rightarrow M$, $\widehat{J}_{g,M} \rightarrow M$ the fibered products $J_g \times_{\mathcal{A}_g} M$, $\widehat{J}_g \times_{\mathcal{A}_g} M$. By construction, there is a section $M \rightarrow \widehat{J}_{g,M}$, which (possibly after finite base change) gives a divisor $D \in \text{CH}^1(J_{g,M})$ which is cohomologous to 0 along the fibers $J_{g,m}$ and satisfies for any $m \in M$

$$D_m^2 = 0 \text{ in } \text{CH}^2(J_{g,m}). \quad (39)$$

The contents of Theorem 3.7(i) is that, under assumption (39), D is of torsion along the fibers $J_{g,m}$. In case (ii), we can make exactly the same construction, and we get this time a generically finite cover $M \rightarrow \mathcal{M}_4$, and a section $M \rightarrow \widehat{J}_M := \widehat{J} \times_{\mathcal{M}_4} M$, inducing a divisor D of $J_M := J \times_{\mathcal{M}_4} M$, which is homologous to zero on the fibers J_m and satisfies, for any $m \in M$

$$D_m^2 = 0 \text{ in } \text{CH}^2(J_m). \quad (40)$$

Proof of Theorem 1.2. With the same notations as above, the cycle $D^2 \in \text{CH}^2(J_{g,M})$, being rationally equivalent to 0 in the fibers of $J_M \rightarrow M$, has trivial infinitesimal invariant (see Theorem 2.8), that is:

$$\delta D^2(m) = 0 \text{ in } H^2(\Omega_{J_{g,M}|J_{g,m}}^2) \quad (41)$$

for a general $m \in M$. By compatibility of the cycle class with the intersection and cup-products, we have

$$\delta D^2(m) = (\delta D(m))^2 \text{ in } H^2(\Omega_{J_{g,M}|J_{g,m}}^2). \quad (42)$$

In case (i), as discussed in the previous section, the infinitesimal invariant $\delta D(m) \in I_{1,W}$ thus satisfies $\overline{M}_{2,W}(\delta D(m)) = 0$ in $I_{2,W}$. According to Proposition 3.1(i), this implies that the infinitesimal invariant

$$\delta D(m) \in (V^* \otimes W_2)/V$$

at a general point $m \in M$ has a unique lift

$$\widetilde{\delta D(m)} \in V^* \otimes W_2$$

which is of rank ≤ 1 . Note that the case where the rank is 0 at the general point is already treated in Lemma 2.1. When the generic rank is 1, we can restrict to the dense Zariski open set of M where this rank is exactly 1. In this case, we have:

Claim 3.9. *The normal function $\nu_D \in \Gamma(\mathcal{J}_g^\vee)$ admits local lifts $\tilde{\nu} \in \Gamma(\mathcal{H}^1)$ having the property that, at any point $m \in M$*

$$(\nabla \tilde{\nu})_m^{0,1} \in H^{0,1}(J_{g,m}) \otimes \Omega_{M,m} = V^* \otimes W_2 \quad (43)$$

is a rank 1 tensor in $V^* \otimes W_2$.

In (43), $(\nabla \tilde{\nu})_m^{0,1}$ denotes the $(0,1)$ -part of $\nabla \tilde{\nu}_m \in H^1(J_{g,m}, \mathbb{C}) \otimes \Omega_{M,m}$.

Proof of Claim 3.9. We choose a first local lift $\tilde{\nu}_0$ of ν_D . By definition, $\delta \nu_{D,m} = \delta D(m)$ is obtained by considering $(\nabla \tilde{\nu}_0)_m^{0,1} \in H^{0,1}(J_{g,m}) \otimes \Omega_{M,m}$ modulo $\overline{\nabla} H^{1,0}(J_{g,m})$. Thus we have

$$(\nabla \tilde{\nu}_0)^{0,1} = \widetilde{\delta D} + \overline{\nabla} \eta \text{ in } \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M), \quad (44)$$

where η is a section of $\mathcal{H}^{1,0}$ and $\widetilde{\delta D}$ is the section of $\mathcal{H}^{0,1} \otimes \Omega_M$ defined by $\widetilde{\delta D}(m) := \widetilde{\delta D(m)}$. By (44), the lift $\tilde{\nu} := \tilde{\nu}_0 - \eta$ then satisfies

$$(\nabla \tilde{\nu})^{0,1} = \widetilde{\delta D} \text{ in } \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M), \quad (45)$$

where $\widetilde{\delta D}$ is a rank 1 section of $\mathcal{H}^{0,1} \otimes \Omega_M$. $\square$

In case (ii), the infinitesimal invariant $\delta D \in I_{1,V}$ satisfies $\overline{M}_{2,V}(\delta D) = 0$ in $I_{2,V}$ and, using Proposition 3.1(ii) and arguing exactly as above, we get, after shrinking M , that the normal function $\nu_D \in \Gamma(\mathcal{J}^\vee)$ admits local lifts $\tilde{\nu} \in \Gamma(\mathcal{H}^1)$ having the property that, at any point $m \in M$

$$(\nabla \tilde{\nu})^{0,1} \in H^{0,1}(J_m) \otimes \Omega_{M,m} = V^* \otimes V_2 \quad (46)$$

is a rank 1 tensor in $V^* \otimes V_2$.

We will prove below the following result.

Proposition 3.10. *Let M be a smooth quasi-projective variety over $\mathbb{C}$ and let $J \rightarrow M$ be a family of abelian varieties, which is either the universal abelian variety of dimension $g \geq 4$ over M , for some generically finite morphism $M \rightarrow \mathcal{A}_g$ (case (i)), or the universal Jacobian over M , for some generically finite morphism $M \rightarrow \mathcal{M}_4$ (case (ii)). Let $D \in \text{Pic}(J)$ be homologous to 0 on fibers. Assume that the associated normal function ν_D has the property that it admits local lifts $\tilde{\nu} \in \Gamma(\mathcal{H}^1)$ such that the $(0,1)$ -part*

$$(\nabla \tilde{\nu})^{0,1} \in \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M)$$

of $\nabla \tilde{\nu}$ has rank ≤ 1 at any point of M . Then ν_D admits local lifts $\tilde{\nu} \in \Gamma(\mathcal{H}^1)$ which are horizontal, that is, $\nabla \tilde{\nu} = 0$.

Proposition 3.10 and Lemma 2.1 then imply that ν_D is of torsion, which concludes the proof of Theorem 3.7. $\square$

Proof of Proposition 3.10. We argue by contradiction and assume furthermore that over a dense Zariski open set of M , ν_D admits local lifts $\tilde{\nu} \in \Gamma(\mathcal{H}^1)$ such that the $(0,1)$ -part

$$(\nabla\tilde{\nu})^{0,1} \in \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M)$$

of $\nabla\tilde{\nu}$ has rank exactly 1.

Lemma 3.11. *The following hold.*

(i) *The normal function ν_D admits local lifts $\tilde{\nu} \in \Gamma(\mathcal{H}^1)$ with the property that*

$$\nabla\tilde{\nu} = \beta \otimes \alpha \text{ in } \Gamma(\mathcal{H}^1 \otimes \Omega_M), \quad (47)$$

where $\beta \in \Gamma(\mathcal{H}^1)$, $\alpha \in \Gamma(\Omega_M)$.

(ii) *Furthermore, the 1-form α defines an integrable distribution.*

Remark 3.12. Our assumption is that, near a general point of M , we can choose local lifts $\tilde{\nu}_0$ satisfying the property that the $(0,1)$ -part $(\nabla\tilde{\nu}_0)^{0,1} \in \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M)$ has rank 1, hence is locally decomposable as in (47). Statement (i) is that we can achieve this local decomposability for the full differential $\nabla\tilde{\nu}$ for an adequate choice of $\tilde{\nu}$.

Proof of Lemma 3.11. By assumption, near a general point of M , we can choose local lifts $\tilde{\nu}_0$ such that

$$\nabla\tilde{\nu}_0 = \beta_0 \otimes \alpha + \beta', \quad (48)$$

Where $\beta_0 \in \Gamma(\mathcal{H}^1)$, α is a holomorphic 1-form, and $\beta' \in \Gamma(\mathcal{H}^{1,0} \otimes \Omega_M)$ is a local section. Differentiating (48) and using $\nabla(\nabla\tilde{\nu}_0) = 0$, we get the local equality

$$\nabla\beta' = -\nabla\beta_0 \wedge \alpha - \beta_0 \otimes d\alpha \text{ in } \Gamma(\mathcal{H}^1 \otimes \Omega_M^2). \quad (49)$$

Projecting (49) in $\Gamma(\mathcal{H}^{0,1} \otimes \Omega_M^2)$ gives

$$\overline{\nabla}_1(\beta') = -(\nabla\beta_0)^{0,1} \wedge \alpha - \beta_0^{0,1} \otimes d\alpha \text{ in } \Gamma(\mathcal{H}^{0,1} \otimes \Omega_M^2), \quad (50)$$

where we recall that

$$\overline{\nabla}_1 : \mathcal{H}^{1,0} \otimes \Omega_M \rightarrow \mathcal{H}^{0,1} \otimes \Omega_M^2 \quad (51)$$

is induced by $\overline{\nabla} : \mathcal{H}^{1,0} \rightarrow \mathcal{H}^{0,1} \otimes \Omega_M$ and is defined by $\overline{\nabla}_1(\eta \otimes \gamma) = \overline{\nabla}\eta \wedge \gamma$. We now make the following

Claim 3.13. *Near a general point of M , Equation (50) implies that $\beta' = \beta'' \otimes \alpha$ for some local section β'' of $\mathcal{H}^{1,0}$.*

Assuming the claim, (48) now becomes

$$\nabla\tilde{\nu}_0 = (\beta_0 + \beta'') \otimes \alpha \text{ in } \Gamma(\mathcal{H}^1 \otimes \Omega_M), \quad (52)$$

which proves Lemma 3.11(i), with $\beta = \beta_0 + \beta''$. Finally we observe that by differentiating (47), we get

$$0 = \beta \otimes d\alpha + \nabla\beta \wedge \alpha \text{ in } \Gamma(\mathcal{H}^1 \otimes \Omega_M^2), \quad (53)$$

As $\beta \neq 0$, this implies that $\alpha \wedge d\alpha = 0$, so that the distribution defined by α is integrable. This proves Lemma 3.11(ii) assuming Claim 3.13. $\square$

Proof of Claim 3.13. We recall that M is a generically finite cover of either $\mathcal{A}_g$, with $g \geq 4$, (Case (i)) or $\mathcal{M}_4$ (case (ii)). We restrict to the dense Zariski open set of M where M is étale over $\mathcal{A}_g$ or $\mathcal{M}_4$. In case (i), keeping the same notation as before, so that in particular $V = H^{1,0}(A)$ for $[A] \in \mathcal{A}_g$, the map $\bar{\nabla}$ at the point $[A]$ is the multiplication map

$$\mu_W : V \rightarrow \text{Hom}(V, W_2),$$

with $W_2 := \text{Sym}^2 V$. In case (ii), with the notation $V = H^{1,0}(C)$ for $[C] \in \mathcal{M}_4$ a general curve, the map $\bar{\nabla}$ at the point $[C]$ is the multiplication map

$$\mu_V : V \rightarrow \text{Hom}(V, V_2),$$

with $V_2 = H^0(C, 2K_C) = W_2/q$, the quadratic form q being nondegenerate.

In case (i), let $\mu_{W,1} : V \otimes W_2 \rightarrow V^* \otimes \bigwedge^2 W_2$ be defined by

$$\mu_{W,1}(\omega) := \mu_W(\beta) \wedge \gamma \in V^* \otimes \bigwedge^2 W_2$$

for $\omega = \beta \otimes \gamma \in V \otimes W_2$ (this is the map $\bar{\nabla}_1$ of (51)). We have to show that for any element $\omega \in V \otimes W_2$ such that

$$\mu_{W,1}(\omega) = v_0^* \otimes \alpha_2 + \omega' \wedge \alpha, \quad (54)$$

for some $v_0^* \in V^*$, $\alpha_2 \in \bigwedge^2 W_2$, $\omega' \in V^* \otimes W_2$, and $\alpha \in W_2$, one has $\omega = v \otimes \alpha$ for some $v \in V$. Let

$$\bar{W}_2 := W_2 / \langle \alpha \rangle.$$

Let $V_0 \subset V$ be the hyperplane defined by v_0^* . Equation (54) is equivalent to the vanishing of $\mu_{W,1}(\omega) \in V^* \otimes \bigwedge^2 W_2$ in the quotient $\text{Hom}(V_0, \bigwedge^2 \bar{W}_2)$. Denoting by $\bar{\omega} \in V \otimes \bar{W}_2$ the projection of $\omega \in V \otimes W_2$, we have to prove that $\bar{\omega} = 0$.

We write $\bar{\omega} = \sum_{i=1}^r v_i \otimes w_i$, with w_i independent in $\bar{W}_2$, and v_i independent in V , for some $r \leq \dim V$. Equation (54) tells that, for any $u \in V_0$,

$$\sum_{i=1}^r uv_i \wedge w_i = 0 \text{ in } \bigwedge^2 \bar{W}_2. \quad (55)$$

As the w_i are independent in $\bar{W}_2$, (55) implies that

$$\overline{uv_i} \in \langle w_i, i \geq 1 \rangle \subset \bar{W}_2,$$

where $\overline{uv_i}$ denotes the projection of uv_i in $\bar{W}_2$. Equivalently, choosing lifts $\tilde{w}_i \in W_2$ of w_i , we have

$$uv_i \in \langle \alpha, \tilde{w}_i, i \geq 1 \rangle \subset W_2, \quad \forall u \in V_0. \quad (56)$$

Condition (56) says in particular that the image of $V_0 \otimes \langle v_i \rangle$ in $W_2 = \text{Sym}^2 V$ is contained in the space $\langle \alpha, \tilde{w}_i, i \geq 1 \rangle$ which has dimension $r + 1$. As $\dim V_0 = g - 1$ and $g \geq 4$, we easily get a contradiction unless $r = 0$ (see below for the precise argument).

In case (ii), we have similarly to show that for any element $\omega \in V \otimes V_2$ such that

$$\mu_{V,1}(\omega) = v_0^* \otimes \alpha_2 + \omega' \wedge \alpha, \quad (57)$$

for some $v_0^* \in V^*$, $\alpha_2 \in \bigwedge^2 V_2$, $\omega' \in V^* \otimes V_2$, and $\alpha \in V_2$, one has $\omega = v \otimes \alpha$ for some $v \in V$. We argue as before, write $\bar{V}_2 := V_2 / \alpha$, and

$$\bar{\omega} = \sum_{i=1}^r v_i \otimes w_i \text{ in } V \otimes \bar{V}_2,$$

with r minimal. The vanishing of $\mu_{V,1}(\omega)$ in $V_0^* \otimes \bigwedge^2(V_2/\alpha)$ then gives as above that, choosing lifts $\tilde{w}_i \in V_2$ of $w_i \in V_2/\alpha$,

$$uv_i \in \langle \alpha, \tilde{w}_i, i \geq 1 \rangle \subset V_2, \forall u \in V_0.$$

We then see that it suffices to prove the following statement:

(*) *Let V be of vector space of dimension 4 and $V_2 = \text{Sym}^2 V/q$, for some nondegenerate quadratic form q . Let $V_0 \subset V$ be a hyperplane, and let*

$$H_r \subset V, H_{r+1} \subset V_2$$

be vector subspaces of respective dimensions $r, r+1$. Then, if

$$V_0 \cdot H_r \subset H_{r+1},$$

one has $r = 0$.

Indeed, we apply statement (*) to $H_r = \langle v_i \rangle \subset V$, $H_{r+1} = \langle \tilde{w}_i, \alpha \rangle \subset V_2$, where the $\tilde{w}_i$ are lifts of w_i in V_2 . Then we conclude that $r = 0$.

We finally prove (*): as the multiplication map $V \otimes V \rightarrow V_2$ does not vanish on nonzero decomposable tensors, the Hopf Lemma says that $\dim(V_0 \cdot H_r) \geq g + r - 2$ if $r > 0$, hence, if $g \geq 4$, we get

$$\dim(V_0 \cdot H_r) \geq r + 2.$$

Thus we must have $r = 0$. □

Coming back to the proof of Proposition 3.10, we now observe the following facts.

Lemma 3.14. *Notation and assumptions being as in Lemma 3.11, we have*

(i) *The local lift $\tilde{\nu}$ of Lemma 3.11 is unique modulo horizontal sections of $\mathcal{H}^1$. It follows that $\nabla \tilde{\nu} = \beta \otimes \alpha$ is a global section of $\mathcal{H}^1 \otimes \Omega_M$. In particular, the holomorphic distribution induced by α is globally defined, at least on the Zariski open set of M where $\delta\nu \neq 0$.*

(ii) *Up to passing to a dense Zariski open set of M if necessary, one has locally defined holomorphic functions f on M such that*

$$d(f\alpha) = 0, \nabla(f^{-1}\beta) = \beta' \otimes \alpha, \tag{58}$$

for some local section β' of $\mathcal{H}^1$.

Proof. Any other local lift $\tilde{\nu}'$ is of the form $\tilde{\nu}' = \tilde{\nu} + \gamma_{\mathbb{Z}} + \gamma^{1,0}$, where $\gamma_{\mathbb{Z}}$ is a section of $H_{\mathbb{Z}}^1$ and $\gamma^{1,0}$ is a local section of $\mathcal{H}^{1,0}$. We then have

$$(\nabla \tilde{\nu}')^{0,1} = (\nabla \tilde{\nu})^{0,1} + \overline{\nabla} \gamma^{1,0}.$$

If $\tilde{\nu}'$, $\tilde{\nu}$ both satisfy the property that $(\nabla \tilde{\nu}')^{0,1}$, $(\nabla \tilde{\nu})^{0,1}$ are rank 1 elements of $\mathcal{H}^{0,1} \otimes \Omega_M$, the uniqueness in Proposition 3.1 implies that $\overline{\nabla} \gamma^{1,0} = 0$, hence that $\gamma^{1,0} = 0$ by injectivity of $\overline{\nabla}$. This proves (i).

By Lemma 3.11(ii), the distribution determined by α is integrable, which means that, on the Zariski open set where it is well-defined and nonzero, α can be locally (in the Euclidean topology) multiplied by an adequate holomorphic function f in such a way that $d(f\alpha) = 0$. Equation (53) then gives

$$0 = \nabla(f^{-1}\beta) \wedge f\alpha \text{ in } \Gamma(\mathcal{H}^1 \otimes \Omega_M^2), \tag{59}$$

which tells that $\nabla(f^{-1}\beta) = \beta' \otimes \alpha$ for some local section $\beta' \in \Gamma(\mathcal{H}^1)$, proving (ii). □

We next prove the following result:

Lemma 3.15. *The global holomorphic section $\nabla \tilde{\nu} = \beta \otimes \alpha$ of $\mathcal{H}^1 \otimes \Omega_M$ constructed in Lemma 3.14(i) is algebraic.*

Proof. We note first that we could have used, instead of the exact sequence (3) defining $\mathcal{J}^\vee$, the following exact sequence

$$0 \rightarrow \mathcal{H}^{1,0} \rightarrow \mathcal{H}^1/H_{\mathbb{Z}}^1 \rightarrow \mathcal{J}^\vee \rightarrow 0. \quad (60)$$

Then the local lifts $\tilde{\nu}$ of ν are local sections of $\mathcal{H}^1/H_{\mathbb{Z}}^1$ defined modulo local sections of $\mathcal{H}^{1,0}$, and we can use the Gauss-Manin connection

$$\nabla : \mathcal{H}^1/H_{\mathbb{Z}}^1 \rightarrow \mathcal{H}^1 \otimes \Omega_M. \quad (61)$$

Next we observe that the exact sequence (60) has an algebraic version, in which $\mathcal{J}^\vee$ is replaced with the sheaf of algebraic sections of $\text{Pic}^0(J_M/M) \rightarrow M$, $\mathcal{H}^{1,0} = R^0\pi_*\Omega_{J/M}$ is the algebraic version of the Hodge bundle, constructed using algebraic relative differentials, and the algebraic version of the term $\mathcal{H}^1/H_{\mathbb{Z}}^1$ in the middle is the sheaf $\mathcal{P}$ of algebraic sections of the abelian group scheme over M parameterizing line bundles on fibers of $\pi : J_M \rightarrow M$ equipped with an (integrable) algebraic connection. Via the Riemann-Hilbert correspondence, the complex points of this group scheme over $t \in M$ are representations of $\pi_1(J_t)$ with values in $\mathbb{C}^*$, that is, elements of

$$H^1(J_t, \mathbb{C}^*) = H^1(J_t, \mathbb{C})/H^1(J_t, \mathbb{Z}).$$

(Note however that the later viewpoint is transcendental.) The connection (61) is algebraically defined on the algebraic group scheme $\mathcal{P}$: if we have a line bundle with algebraic (integrable) connection (L, ∇_L) on a smooth projective variety X and a first order deformation $X_\epsilon \rightarrow \Delta_\epsilon$ of X , where $\Delta_\epsilon := \text{Spec}(\mathbb{C}[\epsilon]/\epsilon^2)$, then there exists a unique pair $(L_\epsilon, \nabla_{L_\epsilon})$ consisting of a line bundle with connection on X_ϵ , restricting to (L, ∇_L) on the central fiber. If $P_1(L)$ is the bundle of 1-jets of L , fitting in an extension

$$0 \rightarrow \Omega_X(L) \rightarrow P_1(L) \rightarrow L \rightarrow 0,$$

and $T_L := P_1(L)^* \otimes L$, there is an exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow T_L \rightarrow T_X \rightarrow 0, \quad (62)$$

and the connection ∇ gives a splitting of this exact sequence, and thus a canonical splitting

$$H^1(X, T_L) \cong H^1(X, \mathcal{O}_X) \bigoplus H^1(X, T_X). \quad (63)$$

The vector space $H^1(X, T_L)$ parameterizes the first order deformations of the pair (X, L) and the natural map $H^1(X, T_L) \rightarrow H^1(X, T_X)$ induced by (62) is the forgetful map. The splitting (63) induces a section $H^1(X, T_X) \rightarrow H^1(X, T_L)$ which provides the desired first order extension. These constructions are classical (see [14], [17]) and can be also seen as a particular case of Simpson's work [19] on de Rham moduli spaces.

Having this algebraic version of the exact sequence (3), we argue as before, except that our local lifts $\tilde{\nu}$ of the algebraic section ν_D of $\mathcal{J}^\vee$ are now algebraic sections of $\mathcal{P}$ which are local for the Zariski topology. The existence and uniqueness of these local lifts $\tilde{\nu}$ satisfying property (47) are proved as before. It follows that $\nabla \tilde{\nu}$ is algebraic. $\square$

Combining Lemmas 3.11, 3.14 and 3.15 together with Proposition 3.16 stated and proved below, we conclude that the local lifts $\tilde{\nu}$ of Lemma 3.11 are in fact horizontal, hence Proposition 3.10 is now fully proved. $\square$

Proposition 3.16. *Let M be smooth quasi-projective and $J_M \rightarrow M$ be a family of abelian varieties, which is either the universal abelian variety of dimension $g \geq 2$ over M for some generically finite morphism $M \rightarrow \mathcal{A}_g$, or the universal Jacobian over M for some generically finite morphism $M \rightarrow \mathcal{M}_g$, with $g \geq 2$. Let $\omega \in \Gamma(\mathcal{H}^1 \otimes \Omega_M)$ be an algebraic section which is locally decomposable, i.e. locally of the form $\beta \otimes \alpha$, and annihilated by ∇ . Then $\omega = 0$.*

Proof. We assume that $\omega \neq 0$ and get a contradiction. Let $\mathcal{H}^1$ be the Hodge bundle with Gauss-Manin connection ∇ . This is an algebraic vector bundle on M . Let $p : \mathfrak{F} \rightarrow M$ be the symplectic frame bundle of $\mathcal{H}^1$. This is a principal $\mathrm{Sp}(2g)$ -bundle which is equipped with a flat connection ∇ . The leaves for this connection are complex submanifolds of $\mathfrak{F}$ corresponding to the locally constant frames. The locally decomposable algebraic section $\omega = \beta \otimes \alpha$ provides a line subbundle $\mathcal{L} \subset \mathcal{H}^1$, hence an algebraic subvariety

$$\mathfrak{F}_{\mathcal{L}} \subset \mathfrak{F}$$

consisting of frames $e_1, \dots, e_{2g}$ with e_1 belonging to $\mathcal{L}$ (that is, proportional to β). The intersection of $\mathfrak{F}_{\mathcal{L}}$ with the leaf $\mathfrak{F}_e$ passing through a point $e = (e_1, \dots, e_{2g}) \in \mathfrak{F}_{m_0}$, is described over M by the condition that $e_1(m)$ is proportional to $\beta(m)$ in the fiber $\mathcal{H}_m^1 := H^1(J_m, \mathbb{C})$. By Lemma 3.14(ii), there exist an integrable codimension 1 algebraic distribution α on M and local holomorphic functions f on M such that $f\beta$ is horizontal along the leaves of the distribution α . Hence the intersection $\mathfrak{F}_e \cap \mathfrak{F}_{\mathcal{L}}$ contains the inverse image under p of the leaf of α passing through $m_0 \in M$. In particular, it has codimension ≤ 1 in $\mathfrak{F}_e$, while $\mathfrak{F}_{\mathcal{L}}$ has codimension $2g - 1$ in $\mathfrak{F}$. It follows that the intersection $\mathfrak{F}_e \cap \mathfrak{F}_{\mathcal{L}}$ is not transverse if $g \geq 2$. As in both considered cases, the Galois group of the connection is Zariski dense in the symplectic group, we can apply [6, Theorem A]. This tells us that the images $p(\mathfrak{F}_e \cap \mathfrak{F}_{\mathcal{L}})$ in M are contained in (proper) ∇ -special algebraic varieties (on which the Galois group of the restricted connection is not anymore Zariski dense). As the images $p(\mathfrak{F}_e \cap \mathfrak{F}_{\mathcal{L}})$ contain open sets of the leaves, which are of codimension 1 in M , we conclude that the leaves of α are algebraic, that is, the distribution α is algebraically integrable. It follows that the distribution α is, at the general point of M , the relative tangent bundle of a rational map $M \dashrightarrow D$, where D is a curve, and in particular it is defined by a closed rational 1-form on M . We then know by Lemma 3.14(ii) that the class β conveniently normalized satisfies $\nabla\beta = 0$ along any leaf $M_{\alpha,t}$ of α . By Deligne's global invariant cycles theorem, the set of classes globally invariant along the leaf $M_{\alpha,t}$ (which we know to be algebraic) generates at any point $m \in M_{\alpha,t}$ a constant Hodge substructure of $H^1(J_{M,m}, \mathbb{Q})$. As, by varying t , m can be taken very general in M , any Hodge substructure of $H^1(J_{M,m}, \mathbb{Q})$ is 0 or equals $H^1(J_{M,m}, \mathbb{Q})$. The first case would provide a contradiction with $\omega \neq 0$. The second case would say that the moduli map for the family $J_M \rightarrow M$ is constant along the leaves, hence factors through a curve, which is not the case in our two geometric situations. $\square$

Remark 3.17. Bruno Klingler suggested the following alternative argument replacing the use of the invariant cycles theorem at the end of the proof given above. Once one knows that the distribution α is algebraically integrable, one gets that the (Zariski closure of the) image of the monodromy representation restricted to the general leaf of α is a proper and normal subgroup of the full algebraic monodromy group, which is $\mathrm{Sp}(2g)$. As this image is a proper subgroup because the leaves are ∇ -special, it follows that the image is trivial. We are thus in the second case above.

4 Proof of Theorem 1.8

This section is devoted to the proof of Theorem 1.8. We start with the following statement, proved in [20]. For a smooth curve C of genus 4, let $\gamma_{P,C} \in \mathrm{CH}^1(C)$ be the Griffiths-Pirola divisor (which is defined up to sign). We can see also $\gamma_{P,C}$ as an element of $\mathrm{CH}^1(C^{(2)})_{\mathrm{hom}}$ using the isomorphism $\mathrm{Pic}^0(C^{(2)}) \cong \mathrm{Pic}^0(C)$.

Theorem 4.1. [20, Theorem 1.4] *Let $\pi : \mathcal{C} \rightarrow B$ be the universal family of smooth $(2, 3)$ -complete intersections in $\mathbb{P}^3$, and let $\pi_2 : \mathcal{C}^{(2)/B} \rightarrow B$ be the corresponding family of symmetric products. Let $C_\eta, C_\eta^{(2)}$ be the respective generic fibers of π, π_2 . Then $\mathrm{CH}_0(C_\eta^{(2)})_{\mathrm{hom}} \otimes \mathbb{Q}$ is 1-dimensional, generated over $\mathbb{Q}$ by $\gamma_{P,\eta}^2$.*

More precisely, Verni proves in [20] that $\gamma_{P,\eta}^2 \in \text{CH}_0(C_\eta^{(2)})_{\text{hom}} \otimes \mathbb{Q}$ is different from 0 (see Theorem 2.10), while earlier arguments appearing in [21] (see also [12]) show that $\text{CH}_0(C_\eta^{(2)})_{\text{hom}} \otimes \mathbb{Q}$ is at most 1-dimensional over $\mathbb{Q}$, generated by the Green-Griffiths 0-cycle of [10]. Note that, due to the sign ambiguity, $\gamma_{P,\eta}$ is not defined as an element of $\text{CH}^1(C_\eta^{(2)})_{\text{hom}}$, but only as an element of $\text{CH}^1(C_{\eta'}^{(2)})_{\text{hom}}$ for a degree 2 field extension of $\mathbb{C}(\mathcal{M}_4)$. However, its square $\gamma_{P,\eta'}^2$ descends to an element $\gamma_{P,\eta}^2$ of $\text{CH}_0(C_\eta^{(2)})$, although maybe only with rational coefficients (see below for the precise argument).

Let $\pi : \mathcal{C} \rightarrow \mathcal{M}_4$ be the universal curve of genus 4 and let $J \rightarrow \mathcal{M}_4$ be the corresponding Jacobian fibration. Assume there is a rational section $\gamma : \mathcal{M}_4 \dashrightarrow K$ of the associated Kummer fibration

$$K := J / \pm \text{Id}.$$

As the quotient map $J \rightarrow K$ is a double cover, one gets by base changing under γ a double cover

$$\mathcal{M}'_4 := \mathcal{M}_4 \times_K J.$$

The double cover $\mathcal{M}'_4$ also maps to J , via a map that we denote by γ' . Let

$$\mathcal{C}' := \mathcal{C} \times_{\mathcal{M}_4} \mathcal{M}'_4$$

and let $J' = J \times_{\mathcal{M}_4} \mathcal{M}'_4$ be the Jacobian fibration of the family $\mathcal{C}' \rightarrow \mathcal{M}'_4$ of genus 4 curves. The rational map γ' provides a rational section of $\pi' : J' \rightarrow \mathcal{M}'_4$ that we also denote by γ' . By construction, the variety $\mathcal{M}'_4$ has an involution i which makes γ' compatible with the involution $-\text{Id}$ on J , that is,

$$\gamma' \circ i = -\gamma'. \quad (64)$$

The section γ' gives (maybe with rational coefficients, and after replacing $\mathcal{M}'_4$ by a dense Zariski open set) a divisor $D_{\gamma'}$ on $\mathcal{C}'$ and also on the induced relative second symmetric product $\mathcal{C}'^{(2)}/\mathcal{M}'_4$. Hence we get a codimension 2-cycle

$$D_{\gamma'}^2 \in \text{CH}^2(\mathcal{C}'^{(2)}/\mathcal{M}'_4).$$

Let $q : \mathcal{C}'^{(2)}/\mathcal{M}'_4 \rightarrow \mathcal{C}^{(2)}/\mathcal{M}_4$ be the natural double cover. Equation (64) tells that, over the same Zariski open set of $\mathcal{M}'_4$, the divisor $D_{\gamma'} \in \text{CH}^1(\mathcal{C}'^{(2)}/\mathcal{M}'_4)$ associated to γ' is anti-invariant under the involution q . It follows that, over the same Zariski open set,

$$D_{\gamma'}^2 \in \text{CH}^2(\mathcal{C}'^{(2)}/\mathcal{M}'_4)$$

is invariant under q , hence the cycle $D_{\gamma'}^2$ comes (with rational coefficients) from a codimension 2-cycle on $\mathcal{C}^{(2)}/\mathcal{M}_4$. More precisely, possibly after shrinking $\mathcal{M}'_4$ and $\mathcal{M}_4$, one has by q -invariance

$$D_{\gamma'}^2 = 2q^* q_* D_{\gamma'}^2 \text{ in } \text{CH}^2(\mathcal{C}'^{(2)}/\mathcal{M}'_4).$$

Applying Theorem 4.1 to $q_* D_{\gamma'}^2$, one gets the following first step in the proof of Theorem 1.8.

Proposition 4.2. *Let $\mathcal{C} \rightarrow \mathcal{M}_4$ be the universal curve of genus 4, and $J \rightarrow \mathcal{M}_4$ the associated Jacobian fibration. Then, if γ is a rational section of the fibration $K = J / \pm \text{Id} \rightarrow \mathcal{M}_4$, there exists a rational number λ such that, for a general point $b \in \mathcal{M}_4$,*

$$D_{\gamma,b}^2 = \lambda D_{\gamma_P,b}^2 \text{ in } \text{CH}_0(C_b^{(2)}). \quad (65)$$

Remark 4.3. The descent to $J \rightarrow \mathcal{M}_4$ has been necessary only to apply Proposition 4.1 in order to get (65). From now on, we will work on a degree 4 cover $J'' \rightarrow \mathcal{M}_4''$ of $J \rightarrow \mathcal{M}_4$, where both divisors $D_{\gamma_P} \in \text{CH}^1(J'')$ and $D_{\gamma'} \in \text{CH}^1(J'')$ are defined.

We now apply the theory of infinitesimal invariants described in Section 2. From Proposition 4.2 and Lemma 1.1, we get that, shrinking $\mathcal{M}_4''$ to a dense Zariski open set if necessary, the cycle

$$D_{\gamma'}^2 - \lambda D_{\gamma_P}^2 \in \text{CH}^2(J'')_{\mathbb{Q}}$$

is trivial, hence also its infinitesimal invariant

$$\begin{aligned} \delta D_{\gamma'}^2(b) - \lambda \delta D_{\gamma_P}^2(b) &= ([D_{\gamma'}^2]^{2,2} - \lambda [D_{\gamma_P}^2]^{2,2})|_{J''} \\ &= ([D_{\gamma'}]_{|J''}^{1,1})^2 - \lambda ([D_{\gamma_P}]_{|J''}^{1,1})^2 \text{ in } H^2(\Omega_{J''|J''}^2) \end{aligned}$$

at the general point $b \in \mathcal{M}_4''$. The map $\mathcal{M}_4'' \rightarrow \mathcal{M}_4$ is étale at the general point of $\mathcal{M}_4''$, so the infinitesimal invariants (see Section 2)

$$[D_{\gamma'}]_{|J''}^{1,1} = \delta\gamma'(b) \in H^1(\Omega_{J''|J''}),$$

$$[D_{\gamma_P}]_{|J''}^{1,1} = \delta\gamma_P(b) \in H^1(\Omega_{J''|J''})$$

and their squares are as in the previous section, in case (ii). Using the explicit description (see (16), (20)) of the spaces $I_{1,V}$ and $I_{2,V}$ of infinitesimal invariants given in Section 3.1, and the notation $\overline{M}_{2,V}$ of (21) for the square map, one concludes the following.

Corollary 4.4. *With the assumptions and notation of Proposition 4.2, the infinitesimal invariant $\delta\gamma'(b) \in I_{1,V}$ of the section γ' at a general point $b \in \mathcal{M}_4'$ satisfies*

$$\overline{M}_{2,V}(\delta\gamma'(b)) = \lambda \overline{M}_{2,V}(\delta\gamma_P(b)) \text{ in } I_{2,V}. \quad (66)$$

4.1 The equation $\overline{M}_{2,V}(\overline{\phi}) = \lambda \overline{M}_{2,V}(\delta\gamma_P)$

Let C be a generic curve of genus 4, $V := H^0(C, K_C)$ and let $q \in \text{Sym}^2 V$ be the unique (nondegenerate) quadratic form vanishing in $H^0(C, 2K_C)$. According to Section 2, the Griffiths-Pirola normal function γ_P (defined up to sign on $\mathcal{M}_4$), has an infinitesimal invariant

$$\delta\gamma_{P,C} \in I_{1,V} \cong (V^* \otimes V_2)/V \quad (67)$$

at $[C] \in \mathcal{M}_4$ (which is well-defined up to sign).

Proposition 4.5. *Let C be a general curve of genus 4 and let $\delta\gamma_{P,C} \in I_{1,V}$ be the infinitesimal invariant (67). Then, if an element $\overline{\phi} \in I_{1,V}$ satisfies*

$$\overline{M}_{2,V}(\overline{\phi}) = \lambda \overline{M}_{2,V}(\delta\gamma_{P,C}) \text{ in } I_{2,V},$$

one has either

$$\overline{M}_{2,V}(\overline{\phi}) = 0 \text{ and } \lambda = 0 \quad (68)$$

or

$$\overline{\phi} = \sqrt{\lambda} \delta\gamma_{P,C} \quad (69)$$

for some square root $\sqrt{\lambda}$ of λ .

Remark 4.6. The meaning of Proposition 4.5 is that the quadratic rational map

$$\overline{M}_{2,V} : \mathbb{P}(I_{1,V}) \dashrightarrow \mathbb{P}(I_{2,V})$$

is, away from its indeterminacy locus, 1-to-1 on its image over the points of the form $\overline{M}_{2,V}(\delta\gamma_{P,C})$, with C general.

The proof will use several lemmas. We are going to use for this proof the lines contained in the quadric $Q = \{q = 0\}$. There are two such families of lines parameterized by $P_1 \cong \mathbb{P}^1$, $P_2 \cong \mathbb{P}^1$. For each line $\Delta \subset Q$, there is a corresponding vector subspace $S_\Delta \subset V$ of dimension 2, and a quotient E_Δ of V of dimension 2, with

$$S_\Delta := I_\Delta(1), E_\Delta = H^0(\Delta, \mathcal{O}_\Delta(1)).$$

We thus get two vector bundles $\mathcal{S}_i, \mathcal{E}_i$ on P_i , $i = 1, 2$, which are restricted from the Grassmannian $G(2, V)$. As we are considering lines contained in Q , on which $q = 0$, the construction in the previous section gives for $i = 1, 2$ linear maps

$$\alpha_i : I_{1,V} \rightarrow H^0(P_i, \mathcal{H}om(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i)) \quad (70)$$

and linear maps

$$\beta_i : I_{2,V} \rightarrow H^0(P_i, \mathcal{H}om(\bigwedge^2 \mathcal{S}_i, \bigwedge^2 (\text{Sym}^2 \mathcal{E}_i))), \quad (71)$$

such that the following formula holds for any $\bar{\phi} \in I_{1,V}$:

$$\beta_i(\overline{M}_{2,V}(\bar{\phi})) = \wedge^2(\alpha_i(\bar{\phi})) \text{ in } H^0(P_i, \mathcal{H}om(\bigwedge^2 \mathcal{S}_i, \bigwedge^2 (\text{Sym}^2 \mathcal{E}_i))). \quad (72)$$

We now have the following result.

Lemma 4.7. *If C is a general curve of genus 4, both morphisms*

$$\alpha_i(\delta\gamma_{P,C}) \in H^0(P_i, \mathcal{H}om(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i)), \quad i = 1, 2,$$

have rank 2 everywhere on P_i .

Proof. Let $C \subset Q \subset \mathbb{P}^3$ be canonically embedded and let $t \in H^0(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(3))$ be a cubic equation defining C inside Q . We know by Theorem 2.9 (see also [20, Section 4.2] for more detail) that $\delta\gamma_{P,C} \in I_{1,V}$ is given by the restrictions to Q of the partial derivatives of t , that is, the image $(\partial t)|_Q$ of t in $V \otimes V_2$, seen as an element of $V^* \otimes V_2$ thanks to the isomorphism $V \cong V^*$ given by q . Indeed, if we change t to $t + aq$, for some $a \in H^0(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(1))$, then $(\partial t)|_Q$ is changed to

$$(\partial(t + aq))|_Q = (\partial t)|_Q + a(\partial q)|_Q,$$

hence we get the same element in $I_{1,V} = (V^* \otimes V_2)/V$. Hence,

$$t \mapsto \partial t|_Q \text{ mod. } V$$

provides a well defined $O(Q)$ -equivariant inclusion of $H^0(Q, \mathcal{O}_Q(3))$ in $I_{1,V} = (V^* \otimes V_2)/V$, which is in fact unique.

If $\Delta \subset Q$ is a member of the i -th ruling, then via the isomorphism $V \cong V^*$ given by q , the ideal $I_\Delta(1)$ is mapped to the space of sections $H^0(\Delta, T_\Delta(-1))$. It follows that, for any point $[\Delta] \in P_i$,

$$\alpha_i(\delta\gamma_{P,C})|_{[\Delta]} \in \text{Hom}(I_\Delta(1), H^0(\Delta, \mathcal{O}_\Delta(2)))$$

identifies with the morphism

$$H^0(\Delta, T_\Delta(-1)) \rightarrow H^0(\Delta, \mathcal{O}_\Delta(2))$$

given by the partial derivatives of $t|_\Delta$. The lemma then follows from the fact that the set of cubic polynomials on Δ with one vanishing partial derivative is a closed algebraic subset of codimension 2 in $H^0(\Delta, \mathcal{O}_\Delta(3))$. $\square$

We will also need the following

Lemma 4.8. *The restriction map*

$$(\alpha_1, \alpha_2) : I_{1,V} \rightarrow H^0(P_1, \mathcal{H}om(\mathcal{S}_1, \text{Sym}^2 \mathcal{E}_1)) \bigoplus H^0(P_2, \mathcal{H}om(\mathcal{S}_2, \text{Sym}^2 \mathcal{E}_2))$$

is injective, and its cokernel is isomorphic to $H^0(Q, \mathcal{O}_Q(3))$.

Before proving the lemma, let us explain how both spaces $H^0(P_i, \mathcal{H}om(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i))$ map naturally to $H^0(Q, \mathcal{O}_Q(3))$. Let

$$\pi_i : Q \rightarrow P_i$$

be the natural morphisms (note that $Q \cong P_1 \times P_2$ via (π_1, π_2)). We have

$$H^0(P_i, \mathcal{H}om(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i)) = H^0(Q, \pi_i^* \mathcal{S}_i^* \otimes \mathcal{O}_Q(2)). \quad (73)$$

We now note that we have an injection

$$j_i : \mathcal{O}_Q(-1) \hookrightarrow \pi_i^* \mathcal{S}_i \quad (74)$$

given by the Gauss map of Q , which produces at each point $x \in Q$ a linear form vanishing at x and along the two lines in Q passing through x . Combining (73) and (74) gives us the desired maps

$$\eta_i : H^0(P_i, \mathcal{H}om(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i)) \rightarrow H^0(Q, \mathcal{O}_Q(3)). \quad (75)$$

As we have the obvious equality of morphisms

$$j'_1 \circ j_1 = j'_2 \circ j_2 : \mathcal{O}_Q(-1) \rightarrow V^* \otimes \mathcal{O}_Q,$$

where $j'_i : \pi_i^* \mathcal{S}_i \hookrightarrow V^* \otimes \mathcal{O}_Q$, $i = 1, 2$, are the natural inclusions, one gets that, for any $\bar{\phi} \in I_{1,V}$,

$$\eta_1(\alpha_1(\bar{\phi})) = \eta_2(\alpha_2(\bar{\phi})) \text{ in } H^0(Q, \mathcal{O}_Q(3)). \quad (76)$$

Proof of Lemma 4.8. Let $\phi \in \text{Hom}(V, V_2)$ with image $\bar{\phi} \in I_{1,V}$ and assume that $\alpha_i(\bar{\phi}) = 0$ in $H^0(P_i, \mathcal{H}om(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i))$ for $i = 1, 2$. Then, defining the vector bundle $\mathcal{K}$ on Q as the kernel of the evaluation map $\text{ev} : V \otimes \mathcal{O}_Q \rightarrow \mathcal{O}_Q(1)$, the image of ϕ in $H^0(Q, V^* \otimes \mathcal{O}_Q(2)) = V^* \otimes V_2$ vanishes in $H^0(Q, \mathcal{K}^*(2))$, since by our assumption, at any point x of Q , it vanishes in $\text{Hom}(I_\Delta, \mathcal{O}_x(2))$ and in $\text{Hom}(I_{\Delta'}, \mathcal{O}_x(2))$, where Δ and Δ' are the two lines contained in Q and passing through x , so that $I_x(1) = I_\Delta(1) \oplus I_{\Delta'}(1)$. Dualizing the exact sequence

$$0 \rightarrow \mathcal{K} \rightarrow V \otimes \mathcal{O}_Q \rightarrow \mathcal{O}_Q(1) \rightarrow 0,$$

we get that the image of ϕ in $H^0(Q, V^* \otimes \mathcal{O}_Q(2))$ comes from an element ϕ' of

$$H^0(Q, \mathcal{O}_Q(-1) \otimes \mathcal{O}_Q(2)) = H^0(Q, \mathcal{O}_Q(1)) = V.$$

Thus $\phi - \phi'$ vanishes in $H^0(Q, V^* \otimes \mathcal{O}_Q(2))$, or equivalently $\phi - \phi' = 0$ in $V^* \otimes V_2$. As $\phi' \in V$, it follows that $\bar{\phi} = 0$ in $I_{1,V} = (V^* \otimes V_2)/V$, which proves the injectivity statement. The other statement is then proved by a dimension computation. Indeed, it is clear that

$$\eta_1 - \eta_2 : H^0(P_1, \mathcal{H}om(\mathcal{S}_1, \text{Sym}^2 \mathcal{E}_1)) \bigoplus H^0(P_2, \mathcal{H}om(\mathcal{S}_2, \text{Sym}^2 \mathcal{E}_2)) \rightarrow H^0(Q, \mathcal{O}_Q(3)) \quad (77)$$

is surjective. The right hand side in (77) has dimension 16 and the left hand side has dimension 48. By (76), the kernel of $\eta_1 - \eta_2$ contains $\text{Im}(\alpha_1, \alpha_2)$ which, as just proved, has dimension equal to $\dim((V^* \otimes V_2)/V) = 32$. Hence the kernel of $\eta_1 - \eta_2$ equals $\text{Im}(\alpha_1, \alpha_2)$. $\square$

Proof of Proposition 4.5. Let C be a general curve of genus 4, and let $\bar{\phi} \in I_{1,V}$ be such that

$$\overline{M}_{2,V}(\bar{\phi}) = \lambda \overline{M}_{2,V}(\delta\gamma_{P,C}) \text{ in } I_{2,V} \quad (78)$$

for some coefficient λ . If $\lambda = 0$, then (68) is satisfied, so we just have to study the case where $\lambda \neq 0$. We now use (72) and apply it to both $\bar{\phi}$ and $\delta\gamma_P$. We thus get from (78) that for $i = 1, 2$,

$$\wedge^2(\alpha_i(\bar{\phi})) = \lambda \wedge^2(\alpha_i(\delta\gamma_{P,C})) \text{ in } H^0(P_i, \text{Hom}(\bigwedge^2 \mathcal{S}_i, \bigwedge^2(\text{Sym}^2 \mathcal{E}_i))). \quad (79)$$

By Lemma 4.7, $\alpha_i(\delta\gamma_{P,C}) \in H^0(P_i, \text{Hom}(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i))$ has rank 2 everywhere, hence by (79), the same is true for the morphism $\alpha_i(\bar{\phi})$. Furthermore (79) implies that the images of the two pointwise injective morphisms $\alpha_i(\bar{\phi}) \in H^0(P_i, \text{Hom}(\mathcal{S}_i, \text{Sym}^2 \mathcal{E}_i))$ and $\alpha_i(\delta\gamma_P)$ are equal. Thus we have, for $i = 1, 2$, a factorization

$$\alpha_i(\bar{\phi}) = \sqrt{\lambda} \alpha_i(\delta\gamma_{P,C}) \circ \psi_i, \quad (80)$$

for some determinant 1 automorphism ψ_i of the vector bundle $\mathcal{S}_i$ on P_i . The vector bundle $\mathcal{S}_1$ on $P_1 \cong \mathbb{P}^1$ is isomorphic to $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$, hence its automorphisms group is isomorphic to $\text{GL}(2)$. The automorphism ψ_1 thus acts on $Q \cong P_1 \times P_2$ via its action on P_2 , and this action is linearized on $\mathcal{O}_Q(1)$, via the action of ψ_1 on $H^0(P_1, \mathcal{S}_1^*) = V$. One can argue similarly with P_1 and P_2 exchanged. Using the maps η_i from (75), we get from the two equations (80) the equalities

$$\begin{aligned} \eta_1(\alpha_1(\bar{\phi})) &= \sqrt{\lambda} \psi_1^*(\eta_1(\alpha_1(\delta\gamma_{P,C}))) \text{ in } H^0(Q, \mathcal{O}_Q(3)), \\ \eta_2(\alpha_2(\bar{\phi})) &= \sqrt{\lambda} \psi_2^*(\eta_2(\alpha_2(\delta\gamma_{P,C}))) \text{ in } H^0(Q, \mathcal{O}_Q(3)). \end{aligned} \quad (81)$$

We now use (76), which tells that

$$\begin{aligned} \eta_1(\alpha_1(\bar{\phi})) &= \eta_2(\alpha_2(\bar{\phi})) \text{ in } H^0(Q, \mathcal{O}_Q(3)), \\ \eta_1(\alpha_1(\delta\gamma_{P,C})) &= \eta_2(\alpha_2(\delta\gamma_{P,C})) \text{ in } H^0(Q, \mathcal{O}_Q(3)). \end{aligned}$$

Combining these equalities with (81), we conclude that the cubic

$$t := \eta_2(\alpha_2(\delta\gamma_{P,C})) \in H^0(Q, \mathcal{O}_Q(3))$$

satisfies

$$\psi_1^* t = \psi_2^* t,$$

which means that t is invariant under the $\mathcal{O}(1)$ -linearized automorphism (ψ_1, ψ_2^{-1}) of $Q \cong P_1 \times P_2$. As the cubic polynomial t is generic (it gives the equation of the curve C by Theorem 2.9), it is not fixed by a nontrivial automorphism of Q . Hence we conclude that $\psi_1 = \text{Id}_{P_1}$, $\psi_2 = \text{Id}_{P_2}$, and so (80) becomes $\alpha_i(\bar{\phi}) = \sqrt{\lambda} \alpha_i(\delta\gamma_{P,C})$ for $i = 1, 2$. Thus we get by Lemma 4.8

$$\bar{\phi} = \sqrt{\lambda} \delta\gamma_{P,C}$$

and (69) is satisfied, which concludes the proof. $\square$

4.2 Proof of Theorem 1.8

We complete in this section the proof of Theorem 1.8. We first establish a slightly weaker statement.

Theorem 4.9. *Let $C \rightarrow \mathcal{M}_4$ be the universal curve of genus 4, and $J \rightarrow \mathcal{M}_4$ the associated Jacobian fibration. Then all rational sections of the fibration $K = J/\pm \text{Id} \rightarrow \mathcal{M}_4$ are given by rational multiples of the Griffiths-Pirola normal function γ_P .*

The meaning of this statement is that, any rational section γ satisfies a relation

$$N\gamma = N'\gamma_P \quad (82)$$

as rational sections of K , for some integers N, N' with $N \neq 0$.

Proof of Theorem 4.9. By Proposition 4.2 and its Corollary 4.4, given a rational section γ of $K \rightarrow \mathcal{M}_4$, the infinitesimal invariant $\delta\gamma'_C \in I_{1,V}$ at a general point $[C]$ of $\mathcal{M}_4''$ of the corresponding normal function γ' defined on an adequate cover $\mathcal{M}_4''$ of $\mathcal{M}_4$, (on which the normal function $\gamma_P \in \Gamma(\mathcal{J}^\vee)$ is also defined), satisfies the hypotheses of Proposition 4.5 for some rational number λ for which (65) holds. It follows that either (68) or (69) is satisfied by $\bar{\phi} := \delta\gamma'_C$. If (68) is satisfied, then the coefficient λ vanishes, hence by (65) the divisor $D_\gamma \in \text{Pic}^0(C) = \text{Pic}^0(C^{(2)}) = \text{CH}^1(C^{(2)})$ satisfies

$$D_\gamma^2 = 0 \text{ in } \text{CH}_0(C^{(2)}). \quad (83)$$

By Theorem 1.2(ii), it follows that D_γ is a torsion element of $\text{Pic}^0(C)$ for very general C , hence by a Baire countability argument, there exists an integer $N \neq 0$ such that (82) is satisfied for any C , with $N' = 0$.

It remains to study what happens if (69) is satisfied. In this case, $\lambda \neq 0$ and the infinitesimal invariants $\delta\gamma', \delta\gamma_P \in \Gamma((\mathcal{H}^{0,1} \otimes \Omega_{\mathcal{M}_4'}/\text{Im } \bar{\nabla}))$ of γ' and γ_P respectively, satisfy the relation

$$\delta\gamma' = \sqrt{\lambda} \delta\gamma_P. \quad (84)$$

Recall that these infinitesimal invariants are computed locally in the Euclidean topology by choosing local lifts $\tilde{\gamma}', \tilde{\gamma}_P$ in $\Gamma(\mathcal{H}^1)$, and defining

$$\begin{aligned} \delta\gamma' &:= (\nabla\tilde{\gamma}')^{0,1} \bmod \bar{\nabla}\mathcal{H}^{1,0}, \\ \delta\gamma_P &:= (\nabla\tilde{\gamma}_P)^{0,1} \bmod \bar{\nabla}\mathcal{H}^{1,0}. \end{aligned}$$

The local lifts are defined up to the addition of local sections of $\mathcal{H}^{1,0}$ and $H_{\mathbb{Z}}^1$, making the definition above independent of the local lift. From this description and (84), we conclude that there exist local lifts $\tilde{\gamma}'$, resp. $\tilde{\gamma}_P$ of γ' , resp. γ_P , satisfying

$$(\nabla(\tilde{\gamma}' - \sqrt{\lambda}\tilde{\gamma}_P))^{0,1} = 0 \text{ in } \Gamma(\mathcal{H}^{0,1} \otimes \Omega_{\mathcal{M}_4'}). \quad (85)$$

Arguing as in the proof of Lemma 2.1, we find that the local lifts $\tilde{\gamma}' - \sqrt{\lambda}\tilde{\gamma}_P$ satisfying (85) are in fact horizontal, and unique modulo sections of the local system $H_{\mathbb{Q}(\sqrt{\lambda})}^1$. (Due to the coefficient $\sqrt{\lambda}$ in (85), we need to enlarge the coefficients of the local system here.) This is equivalent to saying that there exists a global section

$$\nu \in \Gamma(\mathcal{M}_4'', H_{\mathbb{C}}^1/H_{\mathbb{Q}(\sqrt{\lambda})}^1)$$

whose image in $\Gamma(\mathcal{M}_4'', \mathcal{H}^{0,1}/H_{\mathbb{Q}(\sqrt{\lambda})}^1)$ equals $\gamma' - \sqrt{\lambda}\gamma_P$. Arguing as in the proof of Lemma 2.1, we get that $\nu = 0$, so that

$$\gamma' - \sqrt{\lambda}\gamma_P = 0 \text{ in } \Gamma(\mathcal{M}_4'', \mathcal{H}^{0,1}/H_{\mathbb{Q}(\sqrt{\lambda})}^1). \quad (86)$$

To conclude the proof of Theorem 4.9, we just need to prove that $\sqrt{\lambda}$ is a rational number. This is done as follows: given a family $J \rightarrow B$ of abelian varieties and a normal function $\nu \in \Gamma(B, \mathcal{J}^\vee) = \Gamma(B, \mathcal{H}^{0,1}/H_{\mathbb{Z}}^1)$, the class

$$[\nu] \in H^1(B, H_{\mathbb{Z}}^1)$$

of ν , introduced by Griffiths, is obtained by writing the exact sequence (60)

$$0 \rightarrow H_{\mathbb{Z}}^1 \rightarrow \mathcal{H}^{0,1} \rightarrow \mathcal{J}^\vee \rightarrow 0$$

and by applying to ν the associated long exact sequence.

If J is the Jacobian fibration of a family of curves $\mathcal{C} \rightarrow B$ and $\nu = \nu_D$ comes from a relative divisor $D \in \text{Pic}^0(\mathcal{C}/B)$, this class is nothing but the class $[D]$ of the divisor D , at least with $\mathbb{Q}$ -coefficients, for which we have

$$H^1(B, H_{\mathbb{Q}}^1) \subset H^2(\mathcal{C}, \mathbb{Q})/H^2(B, \mathbb{Q}).$$

Coming back to our normal function γ_P for the family $\mathcal{C}'' \rightarrow \mathcal{M}_4''$, the class

$$[\gamma_P] \in H^2(\mathcal{C}'', \mathbb{Q})/H^2(\mathcal{M}_4'', \mathbb{Q})$$

does not vanish. Indeed, this would imply otherwise by a standard argument that the normal function γ_P is trivial. The condition $\gamma' - \sqrt{\lambda}\gamma_P = 0$ of (86) implies that the two cohomology classes $[\gamma']$ and $[\gamma_P]$ satisfy

$$[\gamma'] - \sqrt{\lambda}[\gamma_P] = 0 \text{ in } H^2(\mathcal{C}'', \mathbb{Q}(\sqrt{\lambda}))/H^2(\mathcal{M}_4'', \mathbb{Q}(\sqrt{\lambda})).$$

As $[\gamma']$ and $[\gamma_P]$ are rational, that is, belong to $H^2(\mathcal{C}'', \mathbb{Q})/H^2(\mathcal{M}_4'', \mathbb{Q})$, and $[\gamma_P] \neq 0$, we get that $\sqrt{\lambda}$ is rational and the proof of Theorem 4.9 is finished. $\square$

The proof of Theorem 1.8 is then concluded by the following two lemmas.

Lemma 4.10. *Assumptions and notation being as before, the rational number λ of (84) is an integer.*

Proof. Let γ be a rational section of $K \rightarrow \mathcal{M}_4$. We restrict γ over a rational curve in $\mathcal{M}_4$ constructed as follows: let $Q \subset \mathbb{P}^3$ be a smooth quadric and let $T_0, T_\infty \in H^0(Q, \mathcal{O}_Q(3))$ generate a Lefschetz pencil $\mathbb{P}^1 = \langle T_0, T_\infty \rangle$. The smooth fibers $C_t := \{T_t = 0\}$ form a 1-parameter family of genus 4 curves, hence we get a curve $\mathbb{P}^1 = B \supseteq B^0 \subset \mathcal{M}_4$. The Lefschetz pencil provides a morphism

$$\pi_T : \tilde{Q} \rightarrow B,$$

where $\tilde{Q}$ is the quadric Q blown-up along its base locus, and we get a family

$$\pi_T^0 : \tilde{Q}^0 \rightarrow B^0, \tag{87}$$

by restricting π_T over the regular locus B^0 of π_T .

We observe now that on the family (87), the Griffiths-Pirola normal function is well-defined (there is no need to pass to a double cover), by ordering the two rulings δ_1, δ_2 of Q , and by defining

$$\gamma_P(t) = (\delta_1 - \delta_2)|_{\tilde{Q}_t^0} \in \text{Pic}^0(\tilde{Q}_t^0)$$

for $t \in B^0$. We claim that the section γ restricted to B^0 also lifts to a section of the Jacobian fibration $J_{\pi_T} \rightarrow B^0$. Indeed, the section γ restricted to B^0 lifts in any case to a section γ' of the pulled-back Jacobian fibration $J'_{\pi_T} \rightarrow B'^0$, for some double cover $B'^0 \rightarrow B^0$. On any irreducible component of this double cover, we have by Theorem 4.9 that

$$\gamma' = \lambda\gamma_P.$$

On the other hand, we have by construction of the double cover B'^0 :

$$\gamma' \circ i = -\gamma',$$

where i is the involution of B'^0 over B^0 , while $\gamma_P \circ i = \gamma_P$. Assuming the section γ' is not of torsion (equivalently is nonzero by Lemma 4.11 below), this is possible only if the double cover B'^0 has two components, so that γ' is well defined on each component isomorphic to

B^0 . This proves the claim. The two sections γ_P and γ' of the Jacobian fibration $J_{\pi_T} \rightarrow B^0$ have cohomology classes

$$[\gamma_P], [\gamma'] \in H^2(\tilde{Q}^0, \mathbb{Z})$$

which satisfy the relation

$$[\gamma'] = \lambda[\gamma_P], \in H^2(\tilde{Q}^0, \mathbb{Q}).$$

That λ is an integer then follows from the fact that the class $[\gamma_P] \in H^2(\tilde{Q}_T^0, \mathbb{Z})$ is nonzero and primitive (that is, not divisible by an integer > 1) in $H^2(\tilde{Q}_T^0, \mathbb{Z})$. $\square$

Lemma 4.11. *Any torsion section of the Kummer fibration $K \rightarrow \mathcal{M}_4$ is the zero section.*

Proof. A torsion section of $K \rightarrow \mathcal{M}_4$ is given by local sections of the local system $H_{\mathbb{Z}/N}^1$ on $\mathcal{M}_4$, that glue up to sign. Given a point $t \in \mathcal{M}_4$, this would provide equivalently an element γ_t of $H^1(\mathcal{C}_t, \mathbb{Z}/N\mathbb{Z})$ such that for any $\eta \in \pi_1(\mathcal{M}_4, t)$,

$$\rho(\eta)(\gamma_t) = \pm \gamma_t. \quad (88)$$

As the monodromy representation ρ with integral coefficients has for image the full symplectic group of $H^1(\mathcal{C}_t, \mathbb{Z})$ equipped with the intersection form, condition (88) implies that $\gamma_t = 0$. $\square$

References

- [1] B. Bakker, J. Tsimerman. Functional transcendence of periods and the geometric André-Grothendieck period conjecture. *Forum Math. Sigma* 13 (2025), Paper No. e97.
- [2] G. Baldi, D. Urbanik. Effective atypical intersections and applications to orbit closures, arXiv:2406.16628.
- [3] A. Beauville. Sur l'anneau de Chow d'une variété abélienne. *Math. Ann.* 273 (1986), no. 4, 647-651.
- [4] A. Beauville. Quelques remarques sur la transformation de Fourier dans l'anneau de Chow d'une variété abélienne. *Algebraic geometry (Tokyo/Kyoto, 1982)*, 238-260, *Lecture Notes in Math.*, 1016, Springer, Berlin, (1983).
- [5] S. Bloch. Some elementary theorems about algebraic cycles on Abelian varieties. *Invent. Math.* 37 (1976), no. 3, 215-228.
- [6] D. Blázquez-Sanz, G. Casale, J. Freitag, J. Nagloo. A differential approach to Ax-Schanuel, I. Arxiv:2102.03384v4, to appear in the *Annals of Mathematics*.
- [7] S. Bloch. Torsion algebraic cycles and a theorem of Roitman. *Compositio Math.* 39 (1979), no. 1, 107-127.
- [8] S. Bloch, V. Srinivas. Remarks on correspondences and algebraic cycles, *Amer. J. of Math.* 105 (1983) 1235-1253.
- [9] E. Colombo, J. C. Naranjo, G. P. Pirola. On the dimension of Voisin sets in the moduli space of abelian varieties. *Math. Ann.* 381 (2021), no. 1-2, 91-104.
- [10] M. Green, Ph. Griffiths. An interesting 0-cycle. *Duke Math. J.* 119 (2003), no. 2, 261-313.
- [11] P. Deligne. Théorie de Hodge. II. (French) *Inst. Hautes Études Sci. Publ. Math.* No. 40 (1971), 5-57.

- [12] L. Fu, R. Laterveer, Ch. Vial. The generalized Franchetta conjecture for some hyper-Kähler varieties. (With an appendix by the authors and Mingmin Shen.) *J. Math. Pures Appl.* (9) 130 (2019), 1-35.
- [13] Ph. Griffiths. Infinitesimal variations of Hodge structure. III. Determinantal varieties and the infinitesimal invariant of normal functions. *Compositio Math.* 50 (1983), no. 2-3, 267-324.
- [14] B. Mazur, W. Messing. *Universal extensions and one dimensional crystalline cohomology*. Lecture Notes in Mathematics, Vol. 370. Springer-Verlag, Berlin-New York, (1974).
- [15] N. Mestrano. Conjecture de Franchetta forte. *Invent. Math.* 87 (1987), no. 2, 365-376.
- [16] S. Mukai. Duality between $D(X)$ and $D(\widehat{X})$ with its application to Picard sheaves. *Nagoya Math. J.* 81 (1981), 153-175.
- [17] N. Nitsure. Deformation theory for vector bundles. in *Moduli spaces and vector bundles*, 128–164, London Math. Soc. Lecture Note Ser., 359, Cambridge Univ. Press, Cambridge, (2009).
- [18] A. A. Rojtmán. The torsion of the group of 0-cycles modulo rational equivalence. *Ann. of Math.* (2) 111 (1980), no. 3, 553-569.
- [19] C. Simpson. Moduli of representations of the fundamental group of a smooth projective variety. II. *Inst. Hautes Études Sci. Publ. Math.* No. 80 (1994), 5-79 (1995).
- [20] M. Verni. On infinitesimal invariants of genus 4 curves and the Griffiths-Pirola cycle, [arXiv:2607.12793](https://arxiv.org/abs/2607.12793).
- [21] C. Voisin. The generalized Hodge and Bloch conjectures are equivalent for general complete intersections, *Annales scientifiques de l'ENS* 46, fascicule 3 (2013), 449-475.
- [22] C. Voisin. Une remarque sur l'invariant infinitésimal des fonctions normales. *C. R. Acad. Sci. Paris Sér. I Math.* 307 (1988), no. 4, 157-160.
- [23] C. Voisin. Chow ring and gonality of general abelian varieties. *Ann. H. Lebesgue* 1 (2018), 313-332.
- [24] C. Voisin. Variations de structure de Hodge et zéro-cycles sur les surfaces générales. *Math. Ann.* 299 (1994), no. 1, 77-103.
- [25] C. Voisin. *Chow rings, decomposition of the diagonal, and the topology of families*. Annals of Mathematics Studies, 187. Princeton University Press, Princeton, NJ, 2014. viii+163 pp. ISBN: 978-0-691-16051-1; 0-691-16051-1.
- [26] C. Voisin. *Hodge theory and complex algebraic geometry*. II. Translated from the French by Leila Schneps. Cambridge Studies in Advanced Mathematics, 77. Cambridge University Press, Cambridge, (2003).
- [27] C. Voisin. Variations of Hodge structure and algebraic cycles. *Proceedings of the International Congress of Mathematicians*, Vol. 1, 2 (Zürich, 1994), 706-715, Birkhäuser, Basel, 1995.