

Infinitesimal invariants for cycles modulo algebraic equivalence and 1-cycles on Jacobians

Claire Voisin
CNRS and École Polytechnique

Abstract

We construct an infinitesimal invariant for cycles in a family with cohomology class in the total space lying in a given level of the Leray filtration. This infinitesimal invariant detects cycles modulo algebraic equivalence in the fibers. We apply this construction to the Ikeda family, which gives optimal results for the Beauville decomposition of the 1-cycle of a very general plane curve in its Jacobian.

0 Introduction

This paper is devoted first of all to the construction of infinitesimal invariants for (families of) cycles modulo algebraic equivalence, similar to the infinitesimal invariants used in [31] for the study of families of cycles modulo rational equivalence, and secondly to a geometric application of these infinitesimal invariants to the study of the length of the Beauville decomposition of the 1-cycle of a curve in its Jacobian.

Infinitesimal invariants appear in the following situation: Let $\pi : \mathcal{Y} \rightarrow B$ be a smooth projective morphism, and let \mathcal{Z} be a codimension n cycle in \mathcal{Y} . We assume that the cohomology class $[\mathcal{Z}] \in H^{2n}(\mathcal{Y}, \mathbb{Q})$ belongs to the s -th level of the Leray filtration on $H^{2n}(\mathcal{Y}, \mathbb{Q})$. Then we produce for any point $b \in B$ an infinitesimal invariant $\delta[\mathcal{Z}]_{b,alg}$ which depends only on the first order neighborhood $\mathcal{Z}_{b,2} := \mathcal{Z}|_{\mathcal{Y}_{b,2}}$ of the restricted cycle $\mathcal{Z}_b := \mathcal{Z}|_{\mathcal{Y}_b}$ of the fibre \mathcal{Y}_b (hence the name “infinitesimal invariant”). More precisely, \mathcal{Z} has a Dolbeault cohomology class $[\mathcal{Z}]^{n,n} \in H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$ and this infinitesimal invariant will depend only on the restricted class

$$[\mathcal{Z}]_{|\mathcal{Y}_b}^{n,n} \in H^n(\mathcal{Y}_b, \Omega_{\mathcal{Y}_b}^n).$$

This infinitesimal invariant has the property that it allows to decide whether the restricted cycle \mathcal{Z}_b is algebraically equivalent to 0 at the very general point $b \in B$. More precisely, we will prove the following result:

Theorem 0.1 *In the situation above, assuming $s \geq 2$ and $[\mathcal{Z}] \in L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$, there is an infinitesimal invariant $\delta[\mathcal{Z}]_{alg}$ which is a section of the sheaf*

$$\frac{Ker \bar{\nabla}_{s,n-s,n}}{\Omega_B \wedge Ker \bar{\nabla}_{s-1,n-s,n}},$$

with restriction

$$\delta[\mathcal{Z}]_{b,alg} \in \frac{Ker \bar{\nabla}_{s,n-s,n,b}}{\Omega_{B,b} \wedge Ker \bar{\nabla}_{s-1,n-s,n,b}},$$

which is 0 at a general point $b \in B$ if for general $b \in B$, \mathcal{Z}_b is algebraically equivalent to 0 up to torsion in the fiber \mathcal{Y}_b .

The maps

$$\bar{\nabla}_{s,n-s,n} : \Omega_B^s \otimes \mathcal{H}^{n-s,n} \rightarrow \Omega_B^{s+1} \otimes \mathcal{H}^{n-s-1,n+1}$$

are induced by the infinitesimal variation of Hodge structure on the cohomology of the fibers of π ; they are defined using the Gauss-Manin connection and the Hodge filtration on Hodge bundles associated to the family $\mathcal{Y} \rightarrow B$ (cf. [17], [33, 5.1.2]).

Since the set of points $b \in B$ such that a multiple of the restricted cycle \mathcal{Z}_b is algebraically equivalent to 0 is a countable union of closed algebraic subsets of B , we can rephrase Theorem 0.1 as follows:

Corollary 0.2 *There is a Zariski open set $U \subset B$ such that, if $\delta[\mathcal{Z}]_{b,alg} \neq 0$ for some $b \in U$, then $[\mathcal{Z}]_{b'}$ is not algebraically equivalent to 0 up to torsion for very general $b' \in B$.*

In fact we can take for U the Zariski open set of B where the sheaf $\frac{\text{Ker } \bar{\nabla}_{s,n-s,n}}{\Omega_{B,b} \wedge \text{Ker } \bar{\nabla}_{s-1,n-s,n}}$ is locally free; the nonvanishing of $\delta[\mathcal{Z}]_{b,alg}$ for some $b \in U$ then guarantees the nonvanishing of $\delta[\mathcal{Z}]_{b,alg}$ for general $b \in B$.

The main difference between the present construction and the one in [31] lies in the fact that the infinitesimal invariant introduced here detects cycles modulo algebraic equivalence, while in our previous work only cycles modulo rational equivalence (a much bigger group) could be detected. There is a serious difficulty here, since Lemma 1.2, which says that up to shrinking the base B , the cycle \mathcal{Z} as above is cohomologous to 0 modulo torsion on the total space \mathcal{Y} if the restricted cycles \mathcal{Z}_t are rationally equivalent to 0, obviously does not hold when replacing rational equivalence by algebraic equivalence.

Our infinitesimal invariants are not present, at least explicitly, in the work of Nori [23] and Fakhruddin [15], but they are in fact hidden in the infinitesimal variations of Hodge structures arguments used in their computations of the Leray filtration (we refer to [30], [33] for the relation between infinitesimal invariants and Leray filtration). The work of Ikeda [19] involves similar infinitesimal invariants, which lie however in a bigger cohomology group, and allow to detect the nontriviality of cycles modulo a relation which is much finer than algebraic equivalence. His computations have been extended by Pirola and Rizzi in [26].

Our main application concerns the length of the Beauville decomposition of the canonical 1-cycle of a generic Jacobian. To avoid heavy notation, we denote by $\text{CH}^i(Y)$ the Chow groups of a variety Y with rational coefficients. Recall that Beauville defined in [3] a canonical decomposition on the Chow groups $\text{CH}^i(A)$ of an abelian variety, as a direct sum of eigenspaces for the actions of the homotheties:

$$\text{CH}^i(A) = \bigoplus_s \text{CH}^i(A)_s,$$

where

$$z \in \text{CH}^i(A)_s \Leftrightarrow \mu_m^* z = m^{2i-s} z, \forall m \in \mathbb{Z}.$$

Here for $m \in \mathbb{Z}$ we denote by μ_m the multiplication by m on A . This decomposition works as well for cycles modulo algebraic equivalence. It should be understood as a canonical splitting of the conjectural Bloch-Beilinson filtration, which is supposed

to be a decreasing filtration $F^l \text{CH}^i(A)$ on $\text{CH}^i(A)$, whose graded pieces $Gr_F^l \text{CH}^i(A)$ are governed by the Hodge structures on the cohomology groups $H^{2i-l}(A, \mathbb{Q})$, and which finishes, more precisely $F^{i+1} \text{CH}^i(A) = 0$.

It is not known however if the filtration defined by

$$F^l \text{CH}^i(A) = \bigoplus_{s \geq l} \text{CH}^i(A)_s \quad (0.1)$$

satisfies the axioms of a Bloch-Beilinson filtration, and in particular if

$$F^1 \text{CH}^i(A) = \text{CH}^i(A)_{\text{hom}}.$$

Here the inclusion $F^1 \text{CH}^i(A) \subset \text{CH}^i(A)_{\text{hom}}$ is easy to see : for $s > 0$, $\text{CH}^i(A)_s$ is made of cycles homologous to 0, because μ_n^* acts as multiplication by n^{2i} on $H^{2i}(A, \mathbb{Q})$, but the reverse inclusion, which is equivalent to saying that $\text{CH}^i(A)_s = 0$ for $s < 0$ and the cycle class map is injective on $\text{CH}^i(A)_0$, is unknown.

In this paper, we consider the induced decomposition on the groups $\text{CH}^i(A)/\text{alg}$ of cycles with rational coefficients modulo algebraic equivalence. As mentioned above, for $s > 0$, $\text{CH}^i(A)_s \subset \text{CH}^i(A)_{\text{hom}}$, so that the s -th piece $\text{CH}^i(A)_s/\text{alg}$ is in fact contained in the Griffiths group

$$\text{CH}^i(A)_{\text{hom}}/\text{alg} = \text{Griff}^i(A) \otimes \mathbb{Q}.$$

We will avoid using the notation $\text{Griff}^i(A)_s$ since it is used by Ikeda with a different meaning (see below). In the case where $A = JC$ is the Jacobian of a curve of genus g , there is a natural 1-cycle (codimension $g - 1$ cycle) Z on JC defined as the image of C under the Abel map, with respect to any chosen point or 0-cycle of degree 1 on C . Being defined up to translation, it is well-defined modulo algebraic equivalence. We consider the Beauville decomposition

$$Z = \sum_s Z_s$$

of this cycle, with $Z_s \in \text{CH}^{g-1}(JC)_s/\text{alg}$. The study of the components Z_s has a long history. The first result is the one by Ceresa [7], saying that Z_1 is non zero in $\text{CH}^{g-1}(JC)_1/\text{alg}$ for $g \geq 3$ and C very general in modulus. Ceresa uses the Griffiths criterion [16] saying that if a cycle is algebraically equivalent to 0, its Abel-Jacobi invariant belongs to the maximal abelian subvariety of the intermediate Jacobian, together with a degeneration argument for the study of this Abel-Jacobi invariant. Using Griffiths' infinitesimal invariant [17], an infinitesimal argument can be applied as well (cf. [8]) to prove the nontriviality of the primitive (or transcendental) part of the normal function given by the Abel-Jacobi image of the cycle Z_1 .

In the opposite direction, there is the following vanishing result due to Colombo and Van Geemen [9]:

Theorem 0.3 *Assume a smooth curve C has a g_d^1 , that is a morphism of degree d to \mathbb{P}^1 . Then*

$$Z_s = 0 \text{ in } \text{CH}^{g-1}(JC)_s/\text{alg}, \forall s \geq d - 1.$$

Further results concerning the so-called ‘‘tautological ring’’ introduced by Beauville in [4] and generated under Pontryagin product by the Z_s have been obtained by Herbaut [18], Van der Geer-Kouvidakis [29], Polishchuk and Moonen [25], [21]. The

results of Herbaut (reproved by Van der Geer-Kouvidakis) generalize the Colombo-Van Geemen result to the case where the curve admits a g_d^r , $r \geq 2$. This however does not give more vanishing for the individual components Z_s , but rather polynomial relations between them in the tautological ring (here the ring structure considered is given by Pontryagin product of cycles). The results by Moonen and Polishchuk are devoted to relations in the tautological ring of a general curve, and the paper [4] by Beauville describes a set of generators for this ring.

In the paper [9], Colombo and Van Geemen asked for the optimality of their result, for a very general curve. (As noticed by Ben Moonen, it is not expected that their result is optimal for any curve, because for curves defined over a number field, the Beilinson conjecture [5] predicts

$$Z_s = 0, \forall s \geq 2$$

while the general curve defined over a number field and of genus ≥ 5 has gonality ≥ 4 .)

Concretely, the gonality $\text{gon}(C)$ of a general curve C of genus g , that is the minimal degree of a g_d^1 on C is computed by Brill-Noether theory (cf. [1]) which gives:

$$g = 2k + 1 \text{ or } g = 2k + 2 \Rightarrow \text{gon}(C) = k + 2 \text{ for generic } C.$$

Thus the question asked by Colombo and Van Geemen can be stated in the following form :

Conjecture 0.4 *Let C be a very general curve of genus $g \geq 2k + 1$. Then*

$$Z_k \neq 0 \text{ in } CH^{g-1}(JC)_k/\text{alg}.$$

Note that by a degeneration argument as in Ceresa's paper [7], we can reduce to the case where $g = 2k + 1$. Some results have been obtained in this direction. First of all Fakhruddin [15] shows the nonvanishing of Z_2 for C a general curve of genus ≥ 11 (the expected bound being $g \geq 5$). In a different direction, Ikeda obtains a nonvanishing result for an analogue of the cycles Z_s , $s \leq d - 2$ when the curve is a very general plane curve of degree d (thus of gonality $d - 1$). Colombo-van Geemen's Theorem 0.3 says that for such a curve, the cycle Z_{d-2} is zero in $CH^{g-1}(JC)_{d-2}/\text{alg}$. This last result is also possibly optimal since by Herbaut's above mentioned results in [18], we know that for plane curves, there are extra polynomial relations between the various Z_k , but a priori no further vanishing of a given component Z_k is expected except for the vanishing of Z_{d-2} given by Theorem 0.3. This leads to the following conjecture:

Conjecture 0.5 *Let C be a very general plane curve of degree d (so $g = g(C)$ is given by $2g - 2 = d(d - 3)$). Then the cycle Z_{d-3} is $\neq 0$ in $CH^{g-1}(JC)_{d-3}/\text{alg}$.*

Ikeda addresses this conjecture but not for cycles modulo algebraic equivalence. The quotient of the group of cycles considered by Ikeda is an a priori much bigger group than the group of cycles modulo algebraic equivalence. These groups called "higher Griffiths groups" were defined by Shuji Saito [27] and involve his construction of a Bloch-Beilinson filtration $F_S^i CH^k(Y)$ [28]. The "higher Griffiths groups" of Saito are then defined as the quotient

$$\frac{F_S^i CH^k(Y)}{F_S^{i+1} CH^k(Y) + \langle \Gamma_* F_S^i CH_0(Z) \rangle},$$

where Γ runs over the set of correspondences of codimension k between any smooth projective variety Z and Y . Of course when $i = 1$, the subgroup $\langle \Gamma_* F_S^1 \text{CH}_0(Z) \rangle$ consists exactly of cycles algebraically equivalent to 0, but for $i \geq 1$, the group $\langle \Gamma_* F_S^i \text{CH}_0(Z) \rangle \subset F_S^i \text{CH}^k(Y)$ may be very small, and different from $F_S^i \text{CH}^k(Y) \cap \text{CH}^k(Y)_{\text{alg}}$. We will make a comparison between the Ikeda infinitesimal invariants and ours in section 3, and we will see that, due to the vanishing of canonical syzygies for generic curves [35], for the case of a general curve of genus $2k+1$, his infinitesimal invariants for the cycle Z_s , $s \leq k$, highly depend on the choice of embedding of C into its Jacobian (this embedding, hence also the cycles Z_s , is defined up to translation).

Our main application of the infinitesimal invariant constructed in Section 1 is the proof of conjecture 0.5.

Theorem 0.6 *Let C be a very general plane curve of degree d . Then*

$$Z_{d-3} \neq 0 \text{ in } \text{CH}^{g-1}(JC)_{d-3}/\text{alg}.$$

Theorem 0.6 implies in particular the following result, which is certainly not optimal in view of Conjecture 0.4, but answers partially the question asked in [4]:

Corollary 0.7 *Let C be a very general curve of genus $g > \frac{k(k+3)}{2}$. Then the cycle Z satisfies $Z_k \neq 0$ in $\text{CH}^{g-1}(JC)_k/\text{alg}$.*

Proof. Indeed, we first reduce by degeneration the problem to the case where $g = \frac{k(k+3)}{2} + 1$; then we specialize C to a very general plane curve of degree $k+3$ and apply Theorem 0.6. Since the specialized cycle Z_k is not algebraically equivalent to 0, the same is true at the very general point. ■

For $k = 2$, the above corollary says that the very general curve of genus $g > 5$ has $Z_2 \neq 0$ in $\text{CH}^{g-1}(JC)_2/\text{alg}$. The bound expected from Colombo-Van Geemen conjecture 0.4 is $g \geq 5$, and as already mentioned, Fakhruddin [15] proved the result in the range $g \geq 11$.

In order to prove Theorem 0.6, we compute the infinitesimal invariant of Theorem 0.1 on the Ikeda family of plane curves of degree d (cf. [19]). This family has a very special variation of Hodge structure, which makes the explicit computation quite easy.

1 Infinitesimal invariants for cycles modulo algebraic equivalence

1.1 Infinitesimal invariants for cycles modulo rational equivalence

In this subsection, we recall for the convenience of the reader the construction of infinitesimal invariants for families of cycles modulo rational equivalence. We will explain in next subsection 1.2 the modifications needed to get infinitesimal invariants for families of cycles modulo algebraic equivalence.

Let $\pi : \mathcal{Y} \rightarrow B$ be a smooth projective morphism, where B is smooth quasi-projective, and $\mathcal{Z} \subset \mathcal{Y}$ be a cycle of codimension n . Following Nori [23], consider the cycle class

$$[\mathcal{Z}]^{n,n} \in H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$$

with restriction at $t \in B$:

$$[\mathcal{Z}]_t^{n,n} \in H^n(\mathcal{Y}_t, \Omega_{\mathcal{Y}_t}^n).$$

We have the following lemma, which will be useful later on. It is not completely immediate because we are not working on a smooth projective variety, but only on a quasiprojective one.

Lemma 1.1 *The class $[\mathcal{Z}]^{n,n}$ is determined by the Betti cohomology class $[\mathcal{Z}] \in H^{2n}(\mathcal{Y}, \mathbb{Q})$.*

Proof. The class $[\mathcal{Z}]$ belongs to the pure weight $2n$ part $W_{\text{pure}} H^{2n}(\mathcal{Y}, \mathbb{Q})$ of the Deligne mixed Hodge structure on $H^{2n}(\mathcal{Y}, \mathbb{Q})$, which is defined as the image of the restriction map

$$H^{2n}(\overline{\mathcal{Y}}, \mathbb{Q}) \rightarrow H^{2n}(\mathcal{Y}, \mathbb{Q})$$

for any smooth projective completion $\overline{\mathcal{Y}}$ of \mathcal{Y} . It is a Hodge class in $W_{\text{pure}} H^{2n}(\mathcal{Y}, \mathbb{Q})$, which means that it comes from a Hodge class α on $\overline{\mathcal{Y}}$. This Hodge class can now be seen via the Hodge decomposition as a class $\alpha^{n,n} \in H^n(\overline{\mathcal{Y}}, \Omega_{\overline{\mathcal{Y}}}^n)$, which restricts to a class in $H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. If we change α to α' , the difference $\alpha - \alpha'$ is a Hodge class in $\text{Ker}(H^{2n}(\overline{\mathcal{Y}}, \mathbb{Q}) \rightarrow H^{2n}(\mathcal{Y}, \mathbb{Q}))$. It is known (cf. [34, Proof of Lemma 2.22] or [24, Corollary 7.3]) that if a Hodge class β of degree $2n$ on a smooth projective compactification $\overline{\mathcal{Y}}$ of \mathcal{Y} vanishes on \mathcal{Y} , then there exists a Hodge class β' of degree $2n - 2$ on a desingularization \overline{D} of $D := \overline{\mathcal{Y}} \setminus \mathcal{Y}$ such that $j_* \beta' = \beta$, where $j : \overline{D} \rightarrow \overline{\mathcal{Y}}$ is the natural morphism. It follows that $j_* \beta'^{m-1, n-1} = \beta^{n,n}$, with $\beta'^{m-1, n-1} \in H^{n-1}(\overline{D}, \Omega_{\overline{D}}^{n-1})$ and thus $\beta^{n,n}$ vanishes in $H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. Applying this to $\beta = \alpha - \alpha'$, we conclude that $\alpha^{n,n} = \alpha'^{n,n}$ in $H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. \blacksquare

The class $[\mathcal{Z}]^{n,n}$ was again considered and studied in [31], in the case of a family of 0-cycles in a family of surfaces. Its usefulness for the study of the algebraic cycle \mathcal{Z} comes from the following easy fact (see [33, Theorem 10.19]) :

Lemma 1.2 *If for any $t \in B$, the restricted cycle $\mathcal{Z}_t := \mathcal{Z}|_{\mathcal{Y}_t}$ is rationally equivalent to 0, then there is a dense Zariski open set $U \subset B$ such that \mathcal{Z}_U is of torsion in $CH^n(\mathcal{Y}_U)$, hence in particular*

$$[\mathcal{Z}]^{n,n} = 0 \text{ in } H^n(\mathcal{Y}_U, \Omega_{\mathcal{Y}_U}^n), \quad [\mathcal{Z}]_t^{n,n} = 0 \text{ in } H^n(\mathcal{Y}_t, \Omega_{\mathcal{Y}_t}^n), \quad t \in U.$$

In [31], [30] (see also [33, 5.2.2]), it was furthermore explained how to study the nonvanishing of the class $[\mathcal{Z}]^{n,n}$ by considering the ‘‘Leray filtration’’ on $\Omega_{\mathcal{Y}}^n$:

$$L^s \Omega_{\mathcal{Y}}^n := \pi^* \Omega_B^s \wedge \Omega_{\mathcal{Y}}^{n-s}. \quad (1.2)$$

The associated graded is

$$Gr_L^s \Omega_{\mathcal{Y}}^n = \pi^* \Omega_B^s \otimes \Omega_{\mathcal{Y}/B}^{n-s}.$$

It thus follows that there is a spectral sequence abutting to $R^n \pi_* \Omega_{\mathcal{Y}}^n$:

$$E_1^{s,q}(\Omega_{\mathcal{Y}}^n, L) = \Omega_B^s \otimes R^{s+q} \pi_* \Omega_{\mathcal{Y}/B}^{n-s} \Rightarrow R^{s+q} \pi_* \Omega_{\mathcal{Y}}^n. \quad (1.3)$$

Of course, if B is affine, we have

$$H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n) = H^0(B, R^n \pi_* \Omega_{\mathcal{Y}}^n)$$

and thus the vanishing of $[\mathcal{Z}]^{n,n}$ is equivalent to the vanishing of $[\mathcal{Z}]_t^{n,n}$ for any $t \in B$.

One verifies (see [33, 5.1]), as a consequence of the Katz-Oda description of the Gauss-Manin connection, that the first differentials d_1 are the Griffiths $\overline{\nabla}$ -maps induced by Gauss-Manin connection and Griffiths transversality :

$$d_1 = \overline{\nabla}_{s,n-s,s+q} : \Omega_B^s \otimes \mathcal{H}^{n-s,s+q} \rightarrow \Omega_B^{s+1} \otimes \mathcal{H}^{n-s-1,s+q+1}, \quad (1.4)$$

where the bundles $\mathcal{H}^{p,q} = R^q \pi_* \Omega_{\mathcal{Y}/B}^p$ are the Hodge bundles with fiber at $t \in B$ the spaces $H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}_t}^p)$.

This spectral sequence degenerates at E_2 (see [19], [14]), as it is the case for the topological Leray spectral sequence, and by the same argument as in [12].

Assume now that our cycle \mathcal{Z} has its class in $L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$, which is equivalent, if B is affine and small enough, to the fact that $[\mathcal{Z}]_t^{n,n}$ belongs to the subspace $L^s H^n(\mathcal{Y}_t, \Omega_{\mathcal{Y}_t}^n)$ for any $t \in B$, since we then have the base change property for the sheaf $R^n \pi_* \Omega_{\mathcal{Y}}^n$ endowed with the L -filtration, and the formula

$$L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n) = \Gamma(B, L^s R^n \pi_* \Omega_{\mathcal{Y}}^n).$$

Then it has an ‘‘infinitesimal invariant’’ (see [30], [33, 5.2.2])

$$\begin{aligned} \delta[\mathcal{Z}] \in \Gamma(B, E_\infty^{s,n-s}(\Omega_{\mathcal{Y}}^n, L)) &= \Gamma(B, E_2^{s,n-s}(\Omega_{\mathcal{Y}}^n, L)) \\ &= \Gamma(B, \text{Ker } \overline{\nabla}_{s,n-s,n} / \text{Im } \overline{\nabla}_{s-1,n-s+1,n-1}) \end{aligned} \quad (1.5)$$

where

$$\overline{\nabla}_{s,n-s,n} : \Omega_B^s \otimes \mathcal{H}^{n-s,n} \rightarrow \Omega_B^{s+1} \otimes \mathcal{H}^{n-s-1,n+1}$$

and

$$\overline{\nabla}_{s-1,n-s+1,n-1} : \Omega_B^{s-1} \otimes \mathcal{H}^{n-s+1,n-1} \rightarrow \Omega_B^s \otimes \mathcal{H}^{n-s,n},$$

are the $\overline{\nabla}$ -maps of (1.4).

Let $U \subset B$ be the dense Zariski open set where the coherent sheaf $E_\infty^{s,n-s,n}$ is locally free. The nonvanishing of the restricted infinitesimal invariant

$$\delta[\mathcal{Z}]_t \in \text{Ker } \overline{\nabla}_{s,n-s,n,t} / \text{Im } \overline{\nabla}_{s-1,n-s+1,n-1,t}$$

at some point $t \in U$ implies that the class $[\mathcal{Z}]^{n,n}$ is nonzero on any dense Zariski open subset of U , and thus by Lemma 1.2, the cycle $\mathcal{Z}_{t'}$ is not rationally equivalent to 0 at a very general point $t' \in B$.

Let us comment on the assumption made on $\mathcal{Z}^{n,n}$, namely that it belongs to $L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$, and relate it to the assumption made in Theorem 0.1 that $[\mathcal{Z}] \in L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$.

Proposition 1.3 *Let \mathcal{Z} be a codimension n cycle on \mathcal{Y} such that the Betti cohomology class $[\mathcal{Z}]$ belongs to $L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$. Then, shrinking the base B if necessary, the Dolbeault cycle class $[\mathcal{Z}]^{n,n}$ belongs to $L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$.*

Proof. We use the fact, which is a consequence of the global invariant cycles theorem of Deligne (see [10], [33, 4.3.3]), that there exist Hodge classes δ_i of degree $2d$, $d := \dim \mathcal{Y}/B$, on $\mathcal{Y} \times_B \mathcal{Y}$, (where by Hodge classes, we mean Hodge classes in the pure part $W_{\text{pure}} H^{2d}(\mathcal{Y} \times_B \mathcal{Y}, \mathbb{Q})$) satisfying the following properties: The classes

δ_i act as Künneth projectors, which means the following : The induced morphisms $\delta_{i*} : R\pi_*\mathbb{Q} \rightarrow R\pi_*\mathbb{Q}$ factors as

$$R\pi_*\mathbb{Q} \rightarrow R^i\pi_*\mathbb{Q}[-i] \rightarrow R\pi_*\mathbb{Q},$$

where both maps induce the identity on cohomology of degree i (and 0 in the other degrees). Furthermore, the sum $\sum_i \delta_i$ is equal to the class of the relative diagonal.

When the fibres \mathcal{Y}_t have the property, as it is the case for an abelian fibration, that the Künneth component of their diagonal are algebraic, thus providing cycles $\Delta_{i,t}$ on $\mathcal{Y}_t \times \mathcal{Y}_t$ with classes the i -th Künneth projectors $\delta_{i,t} \in H^{2d}(\mathcal{Y}_t \times \mathcal{Y}_t, \mathbb{Q})$, we can construct the δ_i 's over a generically finite cover B' of B , by spreading out the $\Delta_{i,t}$ (which needs to make a base change) and taking the cohomology class δ_i of the spread-out cycles Δ_i .

We now let the δ_i act on the classes $[\mathcal{Z}]$ and $[\mathcal{Z}]^{n,n}$. For the second action, we use the fact that δ_i is a Hodge class on $\mathcal{Y} \times_B \mathcal{Y}$ hence has a natural image in $H^d(\mathcal{Y} \times_B \mathcal{Y}, \Omega_{\mathcal{Y} \times_B \mathcal{Y}}^d)$. As we have $\sum_i \delta_i = [\Delta_{\mathcal{Y}/B}]$, we get

$$\sum_i \delta_{i*} = Id : H^{2n}(\mathcal{Y}, \mathbb{Q}) \rightarrow H^{2n}(\mathcal{Y}, \mathbb{Q}). \quad (1.6)$$

Note also that the actions δ_{i*} are compatible with the Leray filtration on $H^{2n}(\mathcal{Y}, \mathbb{Q})$ and the filtration L on $H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. Looking at the way the δ_i operate on the spectral sequence associated to the filtration L on $\Omega_{\mathcal{Y}}^n$, one finds that

$$\delta_{i*}[\mathcal{Z}]^{n,n} \in L^{2n-i}H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n).$$

We write now using (1.6)

$$[\mathcal{Z}] = \sum_i \delta_{i*}[\mathcal{Z}].$$

By Lemma 1.1, we get that if the Hodge class $\delta_{i*}[\mathcal{Z}]$ vanishes in $H^{2n}(\mathcal{Y}, \mathbb{Q})$, then its Dolbeault counterpart $\delta_{i*}[\mathcal{Z}]^{n,n}$ vanishes in $H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$.

The assumption that $[\mathcal{Z}] \in L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$ is equivalent to the fact that $\delta_{i*}[\mathcal{Z}] = 0$ for $i > 2n - s$, because $\text{Im } \delta_{i*} : H^{2n}(\mathcal{Y}, \mathbb{Q}) \rightarrow H^{2n}(\mathcal{Y}, \mathbb{Q})$ is a direct summand contained in $L^{2n-i}H^{2n}(\mathcal{Y}, \mathbb{Q})$ and isomorphic to $Gr_L^{2n-i}H^{2n}(\mathcal{Y}, \mathbb{Q})$. But then $\delta_{i*}[\mathcal{Z}]^{n,n}$ vanishes for $i > 2n - s$, that is,

$$[\mathcal{Z}]^{n,n} = \sum_i \delta_{i*}[\mathcal{Z}]^{n,n} = \sum_{i \leq 2n-s} \delta_{i*}[\mathcal{Z}]^{n,n} \in \sum_{s \leq 2n-i} L^{2n-i}H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n) = L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n). \quad \blacksquare$$

The name ‘‘infinitesimal invariant’’ comes from the fact, proved in [32], that for $s = 1$, the infinitesimal invariant is the Griffiths infinitesimal invariant (cf. [17]) obtained by differentiating the normal function associated to \mathcal{Z} .

1.2 Infinitesimal invariants for cycles modulo algebraic equivalence

Our main result in this section is the construction of an infinitesimal invariant for cycles *modulo algebraic equivalence* and the proof of Theorem 0.1. Let $\pi : \mathcal{Y} \rightarrow B$ be a smooth projective morphism and $\mathcal{Z} \subset \mathcal{Y}$ be a cycle of codimension n . Note that

Lemma 1.2 is not true for cycles modulo algebraic equivalence. However, we have the following Proposition 1.4 which will allow us to extract from the $\delta[\mathcal{Z}]$ defined in previous section an infinitesimal invariant associated to cycles modulo algebraic equivalence.

We will use the fact that the maps

$$\bar{\nabla}_{s,n-s,n} : \Omega_B^s \otimes \mathcal{H}^{n-s,n} \rightarrow \Omega_B^{s+1} \otimes \mathcal{H}^{n-s-1,n+1}$$

satisfy the relation:

$$\bar{\nabla}_{s,n-s,n}(\alpha \wedge \sigma) = -\alpha \wedge \bar{\nabla}_{s-1,n-s,n}(\sigma) \quad (1.7)$$

for α a section of Ω_B , σ a section of $\Omega_B^{s-1} \otimes \mathcal{H}^{n-s,n}$. This is proved as follows : for α, σ as above, let $\tilde{\sigma}$ be a section of $F^{n-s}\mathcal{H}^{2n-s}$ which lifts σ . Then by definition,

$$\bar{\nabla}_{s,n-s,n}(\alpha \wedge \sigma) = \nabla(\alpha \wedge \tilde{\sigma}) \text{ mod. } \Omega_B^{s+1} \otimes F^n \mathcal{H}^{2n-s}.$$

By Leibniz rule, this is also equal to $-\alpha \wedge \nabla \tilde{\sigma} + d\alpha \wedge \tilde{\sigma}$ mod. $\Omega_B^{s+1} \otimes F^n \mathcal{H}^{2n-s}$. As $d\alpha \wedge \tilde{\sigma}$ is a section of $\Omega_B^{s+1} \otimes F^n \mathcal{H}^{2n-s}$, we get

$$\bar{\nabla}_{s,n-s,n}(\alpha \wedge \sigma) = -\alpha \wedge \nabla \tilde{\sigma} \text{ mod. } \Omega_B^{s+1} \otimes F^n \mathcal{H}^{2n-s}.$$

It follows from (1.7) that

$$\Omega_B \wedge \text{Ker } \bar{\nabla}_{s-1,n-s,n} \subset \text{Ker } \bar{\nabla}_{s,n-s,n}, \quad (1.8)$$

as implicitly stated in Theorem 0.1. We have now:

Proposition 1.4 *Assume that for any $t \in B$, (a multiple of) \mathcal{Z}_t is algebraically equivalent to 0 and furthermore $[\mathcal{Z}]^{n,n} \in L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. Then there is a Zariski open set $U \subset B$ such that over U , the infinitesimal invariant $\delta[\mathcal{Z}]^{n,n}$ belongs to the image of $\Omega_B \wedge \text{Ker } \bar{\nabla}_{s-1,n-s,n}$ in*

$$\frac{\text{Ker } \bar{\nabla}_{s,n-s,n}}{\text{Im } \bar{\nabla}_{s-1,n-s+1,n-1}}.$$

The following lemma is useful as it explains the simple form that the space of infinitesimal invariants in Theorem 0.1 takes.

Lemma 1.5 *For $s \geq 2$, the space $\Omega_B \wedge \text{Ker } \bar{\nabla}_{s-1,n-s,n} \subset \text{Ker } \bar{\nabla}_{s,n-s,n}$ contains $\text{Im } \bar{\nabla}_{s-1,n-s+1,n-1}$.*

It follows that the quotient

$$\frac{\text{Ker } \bar{\nabla}_{s-1,n-s,n}}{\text{Im } \bar{\nabla}_{s-1,n-s+1,n-1} + \Omega_B \wedge \text{Ker } \bar{\nabla}_{s-1,n-s,n}}$$

is equal to

$$\frac{\text{Ker } \bar{\nabla}_{s-1,n-s,n}}{\Omega_B \wedge \text{Ker } \bar{\nabla}_{s-1,n-s,n}}.$$

Proof. Indeed, for $s \geq 2$, we have $s-1 \geq 1$ and thus for a section η of $\Omega_B^{s-1} \otimes \mathcal{H}^{n-s+1,n-1}$, we can write locally $\eta = \sum_i \alpha_i \wedge \eta_i$, where α_i are sections of Ω_B and η_i are sections of $\Omega_B^{s-2} \otimes \mathcal{H}^{n-s+1,n-1}$. By (1.7),

$$\bar{\nabla}_{s-1,n-s+1,n-1}(\eta) = - \sum_i \alpha_i \wedge \bar{\nabla}_{s-2,n-s+1,n-1}(\eta_i).$$

This proves the result since $\bar{\nabla}_{s-2,n-s+1,n-1}(\eta_i) \in \text{Ker } \bar{\nabla}_{s-1,n-s,n}$. ■

The proof of Proposition 1.4 is postponed to the end of the section. We first show how it implies Theorem 0.1.

Proof of Theorem 0.1. Starting from a cycle \mathcal{Z} with class $[\mathcal{Z}] \in L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$, we know by Proposition 1.3 that up to shrinking B if necessary, the class $[\mathcal{Z}]^{n,n}$ belongs to $L^s H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. We thus have the infinitesimal invariant $\delta[\mathcal{Z}]$ and its restrictions $\delta[\mathcal{Z}]_t \in Gr_L^s H^n(\mathcal{Y}_t, \Omega_{\mathcal{Y}_t}^n)$ for any point $t \in B$. Let U be the dense Zariski open subset of B where the sheaf

$$\frac{\text{Ker } \overline{\nabla}_{s,n-s,n}}{\Omega_B \wedge \text{Ker } \overline{\nabla}_{s-1,n-s,n} + \text{Im } \overline{\nabla}_{s-1,n-s-1,n-1}}$$

is locally free. Then the nonvanishing of the infinitesimal invariant $\delta[\mathcal{Z}]_t \in \frac{\text{Ker } \overline{\nabla}_{s,n-s,n,t}}{\text{Im } \overline{\nabla}_{s-1,n-s-1,n-1,t}}$ modulo the image of $\Omega_{B,t} \otimes \text{Ker } \overline{\nabla}_{s-1,n-s,n,t}$ under the wedge product map implies that for any dense Zariski open subset $U' \subset U$, $\delta[\mathcal{Z}]_{|U'} \in \Gamma(U', \frac{\text{Ker } \overline{\nabla}_{s,n-s,n}}{\text{Im } \overline{\nabla}_{s-1,n-s-1,n-1}})$ does not vanish modulo

$$\text{Im} \left[\Gamma(U', \Omega_{U'} \wedge \text{Ker } \overline{\nabla}_{s-1,n-s,n}) \rightarrow \Gamma(U', \frac{\text{Ker } \overline{\nabla}_{s,n-s,n}}{\text{Im } \overline{\nabla}_{s-1,n-s-1,n-1}}) \right],$$

hence it follows from Proposition 1.4 that the cycle \mathcal{Z}_t is not algebraically equivalent to 0 for very general $t \in U$.

For $s \geq 2$, we can apply Lemma 1.5. Denoting then by $\delta[\mathcal{Z}]_{alg}$ (resp. $\delta[\mathcal{Z}]_{t,alg}$) the image of $\delta[\mathcal{Z}]$ in

$$\Gamma(U, \frac{\text{Ker } \overline{\nabla}_{s,n-s,n}}{\Omega_U \wedge \text{Ker } \overline{\nabla}_{s-1,n-s,n}}),$$

(resp. $\delta[\mathcal{Z}]_t$ in $\frac{\text{Ker } \overline{\nabla}_{s,n-s,n,t}}{\Omega_{B,t} \wedge \text{Ker } \overline{\nabla}_{s-1,n-s,n,t}}$), we constructed the desired infinitesimal invariant for cycles modulo algebraic equivalence satisfying the conclusion of Theorem 0.1 and Corollary 0.2. \blacksquare

Let us first make a few remarks.

Remark 1.6 In [23], Nori uses the cycle class $[\mathcal{Z}]^{n,n}$ to conclude that a certain cycle is not algebraically equivalent to 0 at a general point $t \in B$, assuming its cohomology class \mathcal{Z} does not vanish on the total space. However, he does not introduce the infinitesimal invariant above (although the infinitesimal computation is hidden in his arguments), for the following reason. In his situation, the kernel of the map

$$\overline{\nabla}_{s-1,n-s,n} : \Omega_B^{s-1} \otimes \mathcal{H}^{n-s,n} \rightarrow \Omega_B^s \otimes \mathcal{H}^{n-s-1,n+1}$$

is equal to the image of the map

$$\overline{\nabla}_{s-2,n-s+1,n-1} : \Omega_B^{s-2} \otimes \mathcal{H}^{n-s+1,n-1} \rightarrow \Omega_B^{s-1} \otimes \mathcal{H}^{n-s,n},$$

and it thus follows that

$$\Omega_B \wedge \text{Ker } \overline{\nabla}_{s-1,n-s,n} \subset \text{Im} (\overline{\nabla}_{s-1,n-s+1,n-1} : \Omega_B^{s-1} \otimes \mathcal{H}^{n-s+1,n-1} \rightarrow \Omega_B^s \otimes \mathcal{H}^{n-s,n}).$$

This is why Nori can use the previously defined invariant $\delta[\mathcal{Z}]$ (see Section 1.1) to deal with cycles modulo algebraic equivalence. The same remark applies to the paper of Fakhruddin who uses similar methods applied to general abelian varieties.

Our second remark concerns the consistency of the existence of the above defined invariant and the conjecture made in [36]. We first recall what is this conjecture: Here we can consider for the Bloch-Beilinson filtration on Chow groups with rational coefficients any of the filtrations mentioned in the introduction, for example the one defined by Shuji Saito in [28], or, in the case of an abelian variety, the one which is induced by the Beauville decomposition as in (0.1). Let us denote by F such a filtration. These filtrations are conjectured to satisfy the crucial property:

$$F^{k+1}\mathrm{CH}^k(Y)_{\mathbb{Q}} = 0.$$

These filtrations also induce similar filtrations on Chow groups modulo algebraic equivalence.

We made in [36] (and explored the consequences of) the following conjecture (which generalizes Nori's conjecture [23] on CH^2):

Conjecture 1.7 *We have*

$$F^k \mathrm{CH}^k(Y)/\mathrm{alg} = 0.$$

We want to observe now the following fact, which is consistent with the above conjecture

Lemma 1.8 *Let $\mathcal{Z} \subset \mathcal{Y} \rightarrow B$ be a cycle of codimension n , such that $[\mathcal{Z}] \in L^n H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$. Then for any $t \in B$, the infinitesimal invariant*

$$\delta[\mathcal{Z}]_{t,\mathrm{alg}} \in \frac{\mathrm{Ker} \bar{\nabla}_{n,0,n,t}}{\Omega_{B,t} \wedge \mathrm{Ker} \bar{\nabla}_{n-1,0,n,t}}$$

vanishes.

Proof. This is obvious because the group we are looking at is the following (assuming $n \geq 2$, see Lemma 1.5):

$$\frac{\mathrm{Ker} (\bar{\nabla}_{n,0,n} : \Omega_{B,t}^n \otimes H^{0,n}(\mathcal{Y}_t) \rightarrow \Omega_{B,t}^{n+1} \otimes H^{-1,n+1}(\mathcal{Y}_t))}{\Omega_{B,t} \wedge \mathrm{Ker} (\bar{\nabla}_{n-1,0,n} : \Omega_{B,t}^{n-1} \otimes H^{0,n}(\mathcal{Y}_t) \rightarrow \Omega_{B,t}^n \otimes H^{-1,n+1}(\mathcal{Y}_t))}.$$

Hence, because $H^{-1,n+1}(\mathcal{Y}_t) = 0$, the $\bar{\nabla}$ -maps vanish in this range, and our group is in fact

$$\frac{\Omega_{B,t}^n \otimes H^{0,n}(\mathcal{Y}_t)}{\Omega_{B,t} \wedge \Omega_{B,t}^{n-1} \otimes H^{0,n}(\mathcal{Y}_t)}$$

which is 0 because $n > 0$. ■

Our last remark concerns the assumption that $[\mathcal{Z}] \in L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$ and will be useful for our main application. There are two cases where the Leray level of an algebraic cycle in a family of varieties is easy to compute:

1) The family \mathcal{Y} is a family of smooth ample hypersurfaces or complete intersections in projective space (or more generally, any ambient variety X whose rational cohomology is made of classes of algebraic cycles). Then the hard Lefschetz theorem in this case says that the interesting part of the cohomology of \mathcal{Y}_t is supported in degree $m := \dim \mathcal{Y}_t$. Given any cycle \mathcal{Z} of codimension $n < \dim \mathcal{Y}_t/2$ in \mathcal{Y} , we can write (at least with rational coefficients)

$$\mathcal{Z} = \mathcal{Z}' + \Gamma$$

where Γ is the restriction of a cycle coming from X and $\mathcal{Z}'_{|\mathcal{Y}_t}$ is cohomologous to 0. Looking more precisely at the shape of the Leray spectral sequence of $\mathcal{Y} \rightarrow B$ and applying the Lefschetz hyperplane section theorem, one finds that on an adequate dense Zariski open set $U \subset B$, the Dolbeault cohomology class of $\mathcal{Z}'_{|\mathcal{Y}_U}$ belongs to $L^{2n-d}H^{2n}(\mathcal{Y}_U, \mathbb{Q})$, where $d = \dim \mathcal{Y}_t$.

2) The second case where the Leray level is easy to compute is the case of a cycle in a family of abelian varieties $\mathcal{A} \rightarrow B$. In this case we have the following fact:

Lemma 1.9 *If the cycle \mathcal{Z} satisfies the condition that*

$$Z_t \in CH^n(\mathcal{A}_t)_s$$

for any $t \in B$, then there is a dense Zariski open set $U \subset B$ such that $[\mathcal{Z}]_U \in L^s H^{2n}(\mathcal{A}_U, \mathbb{Q})$.

Proof. Indeed, under our assumptions, Lemma 1.2 and the relative Beauville decomposition (see [13]) show that there exists a dense Zariski open set $U \in B$ such that for any $k \in \mathbb{Z}^*$,

$$\mu_k^* \mathcal{Z}_U = k^{2n-s} \mathcal{Z} \text{ in } CH^n(\mathcal{A}_U). \quad (1.9)$$

It thus follows that the class $[\mathcal{Z}] \in H^{2n}(\mathcal{A}_U, \mathbb{Q})$ satisfies for any k :

$$\mu_k^* [\mathcal{Z}] = k^{2n-s} [\mathcal{Z}] \text{ in } H^{2n}(\mathcal{A}_U, \mathbb{Q}). \quad (1.10)$$

But there is a canonical Deligne decomposition

$$R\pi_* \mathbb{Q} = \bigoplus R^i \pi_* \mathbb{Q}[-i] \quad (1.11)$$

such that for any i ,

$$\mu_k^* \circ \pi_i = k^i \pi_i : R\pi_* \mathbb{Q} \rightarrow R\pi_* \mathbb{Q},$$

where the π_i are the projectors associated to the decomposition (1.11) (cf. [13]). It follows that

$$L^s H^{2n}(\mathcal{A}_U, \mathbb{Q}) = \bigoplus_{i \geq s} H^{2n}(\mathcal{A}_U, \mathbb{Q})_i, \quad (1.12)$$

where

$$H^{2n}(\mathcal{A}_U, \mathbb{Q})_i = \{\alpha \in H^{2n}(\mathcal{A}_U, \mathbb{Q}), \mu_k^* [\mathcal{Z}] = k^{2n-i} [\mathcal{Z}] \text{ in } H^{2n}(\mathcal{A}_U, \mathbb{Q}) \text{ for any } k\}.$$

By (1.10) and (1.12), $[\mathcal{Z}]_U \in L^s H^{2n}(\mathcal{A}_U, \mathbb{Q})$. ■

Proof of Proposition 1.4. We first prove the result in the case which will be useful for us, namely when $\pi : \mathcal{Y} \rightarrow B$ is a family of abelian varieties. The proof is easier in this case because, as the proof of lemma 1.9 shows, not only the spectral sequence abutting to $R^n \pi_*(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$ degenerates at E_2 but there is a natural decomposition

$$R^n \pi_* \Omega_{\mathcal{Y}}^n = \bigoplus_s (R^n \pi_* \Omega_{\mathcal{Y}}^n)_s, \quad (1.13)$$

where

$$(R^n \pi_* \Omega_{\mathcal{Y}}^n)_s = \{\alpha \in R^n \pi_* \Omega_{\mathcal{Y}}^n, \mu_k^* \alpha = k^{2n-s} \alpha, \forall k \in \mathbb{Z}\}$$

and the decomposition (1.13) is a splitting of the filtration induced by the filtration L on $\Omega_{\mathcal{Y}}$. We will explain at the end of the proof how to modify the argument to make it work in general.

So assume $\mathcal{Y} \rightarrow B$ is a family of abelian varieties over a smooth affine variety B , and consider a codimension n cycle \mathcal{Z} on \mathcal{Y} . Assume that for any $t \in B$, \mathcal{Z}_t is algebraically equivalent to 0, and furthermore $[\mathcal{Z}] \in L^s H^{2n}(\mathcal{Y}, \mathbb{Q})$. Using the relative Beauville decomposition of Deninger-Murre [13], we can replace \mathcal{Z} by \mathcal{Z}_s , which does not change the class $\delta[\mathcal{Z}] \in Gr_s^L H^n(\mathcal{Y}, \Omega_{\mathcal{Y}}^n)$ and it still satisfies the assumption that its restrictions to the fibers is algebraically equivalent to 0. So we assume from now on that $\mathcal{Z} = \mathcal{Z}_s$.

The definition of algebraic equivalence together with an elementary argument involving countability of Chow varieties and uncountability of \mathbb{C} implies that over a Zariski dense open subset U of B , there exist

1. A family of smooth nonnecessarily connected curves $\rho : \mathcal{C} \rightarrow U$;
2. A divisor \mathcal{D} of \mathcal{C} which is homologous to 0 on the fibers of ρ ;
3. A codimension n correspondence $\Gamma \subset \mathcal{C} \times_U \mathcal{Y}_U$ with \mathbb{Q} -coefficients,

such that

$$\mathcal{Z} = \Gamma_*(\mathcal{D}) \text{ in } \text{CH}^n(\mathcal{Y}_U). \quad (1.14)$$

Here we recall that, denoting by p_1, p_2 the two proper smooth projections from $\mathcal{C} \times_U \mathcal{Y}_U$ to \mathcal{C} and \mathcal{Y}_U respectively,

$$\Gamma_*(\mathcal{D}) := p_{2*}(p_1^* \mathcal{D} \cdot \Gamma).$$

The equality of cycles (1.14) provides now, using the natural functoriality properties of the Dolbeault cycle class, the corresponding equality of Dolbeault cohomology classes:

$$[\mathcal{Z}_U]^{n,n} = p_{2*}([\Gamma]^{n,n} \cdot p_1^*[\mathcal{D}]^{1,1}) \text{ in } H^n(\mathcal{Y}_U, \Omega_{\mathcal{Y}_U}^n). \quad (1.15)$$

We use now again the relative Beauville decomposition for cycles in the family $\mathcal{C} \times_U \mathcal{Y} \rightarrow \mathcal{C}$. The cycle Γ decomposes as

$$\Gamma = \sum_p \Gamma_p,$$

with $\mu_k^* \Gamma_p = k^{2n-p} \Gamma_p$ and accordingly, the class

$$[\Gamma]^{n,n} \in H^n(\mathcal{C} \times_U \mathcal{Y}_U, \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)$$

decomposes then as

$$[\Gamma]^{n,n} = \bigoplus_p [\Gamma_p]^{n,n},$$

where $[\Gamma_p]^{n,n} \in H^n(\mathcal{C} \times_U \mathcal{Y}_U, \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)$ satisfies the condition

$$\mu_k^* [\Gamma_p]^{n,n} = k^{2n-p} [\Gamma_p]^{n,n}, \forall k \in \mathbb{Z}^*. \quad (1.16)$$

Recall that $[\mathcal{Z}]_U^{n,n}$ belongs to $H^n(\mathcal{Y}_U, \Omega_{\mathcal{Y}_U}^n)_s$. Observing that the μ_k^* maps are compatible with the pushforward map p_{2*} , in the sense that

$$p_{2*} \circ \mu_k^* = \mu_k^* \circ p_{2*}, \quad (1.17)$$

we get from (1.15), (1.16) and (1.17) the following equality:

$$[\mathcal{Z}_U]^{n,n} = p_{2*}([\Gamma_s]^{n,n} \cdot p_1^*[\mathcal{D}]^{1,1}) \text{ in } H^n(\mathcal{Y}_U, \Omega_{\mathcal{Y}_U}^n)_s \subset L^s H^n(\mathcal{Y}_U, \Omega_{\mathcal{Y}_U}^n), \quad (1.18)$$

and projecting everything modulo L^{s+1} , we get :

$$\delta[\mathcal{Z}] = p_{2*}(\delta[\Gamma_s] \cdot p_1^* \delta[\mathcal{D}]) \text{ in } Gr_L^s H^n(\mathcal{Y}_U, \Omega_{\mathcal{Y}_U}^n) = \Gamma(U, E_2^{s, n-s}(\Omega_{\mathcal{Y}_U}^n, L)). \quad (1.19)$$

Here $[\mathcal{D}]^{1,1} \in H^1(\mathcal{C}, \Omega_{\mathcal{C}})$, and as \mathcal{D} is cohomologous to 0 on the fibers \mathcal{C}_t , $t \in U$, we have $[\mathcal{D}] \in L^1 H^1(\mathcal{C}, \Omega_{\mathcal{C}})$, so

$$\delta[\mathcal{D}] \in \Gamma(U, \Omega_U \otimes \mathcal{H}_{\mathcal{C}}^{0,1} / \overline{\nabla}_{\mathcal{C}} \mathcal{H}_{\mathcal{C}}^{1,0}), \quad (1.20)$$

where the bundles $\mathcal{H}_{\mathcal{C}}^{0,1} = R^1 \rho_* \mathcal{O}_{\mathcal{C}}$, $\mathcal{H}_{\mathcal{C}}^{1,0} = R^0 \rho_* \Omega_{\mathcal{C}/U}$ are the Hodge bundles of the family $\rho : \mathcal{C} \rightarrow U$ and $\nabla_{\mathcal{C}}$, $\overline{\nabla}_{\mathcal{C}}$ denote the corresponding Gauss-Manin connection and $\overline{\nabla}$ -map. (To avoid confusion, we will also write below $\overline{\nabla}_{\mathcal{Y}}$, $\mathcal{H}_{\mathcal{Y}}^{p,q}$ for the infinitesimal variation of Hodge structure of the family \mathcal{Y} or \mathcal{Y}_U .) We will denote below by $\widetilde{\delta}[\mathcal{D}]$ any lift of $\delta[\mathcal{D}]$ in $\Omega_U \otimes \mathcal{H}_{\mathcal{C}}^{0,1}$.

We now examine

$$\gamma := \delta[\Gamma_s] \in H^n(\mathcal{C} \times_U \mathcal{Y}_U, \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s = H^0(U, (R^n \pi'_* \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s),$$

where $\pi' := \pi \circ p_2 : \mathcal{C} \times_U \mathcal{Y}_U \rightarrow U$, and as before

$$(R^n \pi'_* \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s = \{\alpha \in R^n \pi'_* \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n, \mu_k^* \alpha = k^{2n-s} \alpha\}. \quad (1.21)$$

We recall that the class γ is a section of

$$(R^n \pi'_* \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s,$$

with $\pi' = \rho \circ p_1 = \pi \circ p_2 : \mathcal{C} \times_U \mathcal{Y}_U \rightarrow U$. Considering the exact sequence

$$0 \rightarrow R^1 \rho_*(R^{n-1} p_{1*} \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n) \rightarrow R^n \pi'_* \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n \rightarrow \rho_*(R^n p_{1*} \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n) \rightarrow 0,$$

and taking the direct summand (where μ_k^* acts by multiplication by k^{2n-s}), we get:

$$0 \rightarrow R^1 \rho_*(R^{n-1} p_{1*} \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s \rightarrow (R^n \pi'_* \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s \rightarrow \rho_*(R^n p_{1*} \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s \rightarrow 0.$$

As $\Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n$ projects naturally to $p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^{n-1}$, we get a morphism

$$\rho_*(R^n p_{1*} \Omega_{\mathcal{C} \times_U \mathcal{Y}_U}^n)_s \rightarrow \mathcal{H}_{\mathcal{C}}^{1,0} \otimes (R^n \pi'_* \Omega_{\mathcal{Y}_U}^{n-1})_{s-1},$$

where the subscript $s-1$ instead of s on the right follows from formula (1.21) and the shift in total degree.

As was explained in Section 1.1, $(R^n \pi_* \Omega_{\mathcal{Y}_U}^{n-1})_{s-1}$ is computed as

$$\frac{\text{Ker}(\bar{\nabla}_{\mathcal{Y},s-1,n-s,n} : \Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n} \rightarrow \Omega_U^s \otimes \mathcal{H}_{\mathcal{Y}}^{n-s-1,n+1})}{\text{Im}(\bar{\nabla}_{\mathcal{Y},s-2,n-s+1,n-1} : \Omega_U^{s-2} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s+1,n-1} \rightarrow \Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n})}.$$

Combining these various morphisms and identifications, we find that γ provides a class

$$\gamma_1 \in \Gamma(U, \mathcal{H}_{\mathcal{C}}^{1,0} \otimes \frac{\text{Ker} \bar{\nabla}_{\mathcal{Y},s-1,n-s,n}}{\text{Im} \bar{\nabla}_{\mathcal{Y},s-2,n-s+1,n-1}}).$$

Below, we will denote by $\tilde{\gamma}_1$ any lift of γ_1 in $\Gamma(U, \mathcal{H}_{\mathcal{C}}^{1,0} \otimes \text{Ker} \bar{\nabla}_{\mathcal{Y},s-1,n-s,n}) \subset \Gamma(U, \mathcal{H}_{\mathcal{C}}^{1,0} \otimes \Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n})$.

The proof of Proposition 1.4 in the case of a family of abelian varieties is then an immediate consequence of the following Lemma 1.10. Indeed, it says in particular that, under the assumptions of Proposition 1.4, $\delta[\mathcal{Z}] \in \frac{\text{Ker} \bar{\nabla}_{\mathcal{Y},s,n-s,n}}{\text{Im} \bar{\nabla}_{\mathcal{Y},s-1,n-s+1,n-1}}$ belongs to the image, under the wedge product map, of the sheaf $\Omega_U \otimes \frac{\text{Ker} \bar{\nabla}_{\mathcal{Y},s-1,n-s,n}}{\text{Im} \bar{\nabla}_{\mathcal{Y},s-2,n-s+1,n-1}}$.

Lemma 1.10 *With the same assumptions and notation as above, we have*

$$\delta[\mathcal{Z}] = w(\langle \delta[\widetilde{\mathcal{D}}], \tilde{\gamma}_1 \rangle) \text{ modulo } \text{Im} \bar{\nabla}_{\mathcal{Y},s-1,n-s+1,n-1}, \quad (1.22)$$

where the brackets mean that we use the duality between $\mathcal{H}_{\mathcal{C}}^{0,1}$ and $\mathcal{H}_{\mathcal{C}}^{1,0}$ to get a contraction map

$$\langle, \rangle : (\Omega_U \otimes \mathcal{H}_{\mathcal{C}}^{0,1}) \otimes (\mathcal{H}_{\mathcal{C}}^{1,0} \otimes \Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n}) \rightarrow \Omega_U \otimes \Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n},$$

and the map w is induced by the wedge product map $\Omega_U \otimes \Omega_U^{s-1} \rightarrow \Omega_U^s$.

Before giving the proof of (1.22), let us explain why the projection of

$$w(\langle \delta[\widetilde{\mathcal{D}}], \tilde{\gamma}_1 \rangle) \in \Gamma(U, \text{Ker} \bar{\nabla}_{\mathcal{Y},s,n-s,n}) \subset \Gamma(U, \Omega_U^s \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n})$$

in the quotient $\frac{\text{Ker} \bar{\nabla}_{\mathcal{Y},s,n-s,n}}{\text{Im} \bar{\nabla}_{\mathcal{Y},s-1,n-s+1,n-1}}$ does not depend on the choice of the lifts $\delta[\widetilde{\mathcal{D}}]$, $\tilde{\gamma}_1$.

If we change $\tilde{\gamma}_1$ by a $\bar{\nabla}_{\mathcal{Y}}$ -exact section of $\Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{C}}^{1,0} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s,n}$, that is $(\tilde{\gamma}_1)' = \tilde{\gamma}_1 + \bar{\nabla}_{\mathcal{Y},s-2,n-s+1,n-1}(\gamma')$ for a section γ' of $\Omega_U^{s-2} \otimes \mathcal{H}_{\mathcal{C}}^{1,0} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s+1,n-1}$, then we clearly have:

$$w(\langle \delta[\widetilde{\mathcal{D}}], (\tilde{\gamma}_1)' \rangle) = w(\langle \delta[\widetilde{\mathcal{D}}], \tilde{\gamma}_1 \rangle) + \bar{\nabla}_{\mathcal{Y},s-1,n-s+1,n-1}(w(\langle \delta[\widetilde{\mathcal{D}}], \gamma' \rangle)),$$

and the difference is $\bar{\nabla}_{\mathcal{Y},s-1,n-s+1,n-1}$ -exact. On the other hand, if we change $\delta[\widetilde{\mathcal{D}}]$ to

$$(\delta[\widetilde{\mathcal{D}}])' = \delta[\widetilde{\mathcal{D}}] + \bar{\nabla}_{\mathcal{C}}(\alpha)$$

where α is a section of $\mathcal{H}_{\mathcal{C}}^{1,0}$, then we get

$$\begin{aligned} w(\langle (\delta[\widetilde{\mathcal{D}}])', \tilde{\gamma}_1 \rangle) &= w(\langle \delta[\widetilde{\mathcal{D}}], \tilde{\gamma}_1 \rangle) + w(\langle \bar{\nabla}_{\mathcal{C}}(\alpha), \tilde{\gamma}_1 \rangle) \\ &= w(\langle \delta[\widetilde{\mathcal{D}}], \tilde{\gamma}_1 \rangle) - w(\langle \alpha, \bar{\nabla}_{\mathcal{C}}(\tilde{\gamma}_1) \rangle). \end{aligned}$$

On the other hand, the class $\overline{\nabla}_{\mathcal{C}}(\tilde{\gamma}_1)$, hence also the contraction $-w(\langle \alpha, \overline{\nabla}_{\mathcal{C}}(\tilde{\gamma}_1) \rangle)$, is $\overline{\nabla}_{\mathcal{Y}}$ -exact for the following reason: Recall that $\tilde{\gamma}_1$ is the projection of a $\overline{\nabla}_{\mathcal{C} \times_B \mathcal{Y}}$ -closed section $\tilde{\gamma}$ of $(R^n \pi'_* \Omega_{\mathcal{C} \times_B \mathcal{Y}/B}^n)_s$. This section has a Künneth decomposition $\tilde{\gamma} = \tilde{\gamma}_1 + \tilde{\gamma}_2 + \dots$ with $\tilde{\gamma}_2 \in \Gamma(U, \Omega_U^{s-1} \otimes \mathcal{H}_{\mathcal{C}}^{0,1} \otimes \mathcal{H}_{\mathcal{Y}}^{n-s+1, n-1})$. The condition $\overline{\nabla}_{\mathcal{C} \times \mathcal{Y}}(\tilde{\gamma}) = 0$ implies then

$$\overline{\nabla}_{\mathcal{C}}(\tilde{\gamma}_1) = -\overline{\nabla}_{\mathcal{Y}}(\tilde{\gamma}_2).$$

Proof of Lemma 1.10. We apply formula (1.18). We observe that the map

$$p_{2*} : R^{n+1} \pi'_* \Omega_{\mathcal{C} \times \mathcal{Y}}^{n+1} \rightarrow R^n \pi_* \Omega_{\mathcal{Y}}^n$$

is obtained by composing the maps I and g , where

$$g : R^{n+1} \pi'_* \Omega_{\mathcal{C} \times \mathcal{Y}}^{n+1} \rightarrow R^{n+1} \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^n)$$

is induced by the morphism

$$\Omega_{\mathcal{C} \times \mathcal{Y}}^{n+1} \rightarrow p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^n$$

coming from the exact sequence:

$$0 \rightarrow p_2^* \Omega_{\mathcal{Y}} \rightarrow \Omega_{\mathcal{C} \times \mathcal{Y}} \rightarrow p_1^* \Omega_{\mathcal{C}/U} \rightarrow 0$$

and

$$I : R^{n+1} \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^n) \rightarrow R^n \pi_* \Omega_{\mathcal{Y}}^n$$

is the integration over the fibre of p_2 , also obtained by writing $\pi' = \pi \circ p_2$ and sending $R^{n+1} \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^n)$ to

$$R^n \pi_*(R^1 p_{2*}(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^n)) = R^n \pi_*(\Omega_{\mathcal{Y}}^n)$$

by the corresponding Leray spectral sequence.

We apply the morphism g to the class $p_1^*[\mathcal{D}]^{1,1} \cdot [\Gamma_s]^{n,n}$. As $[\mathcal{D}]^{1,1} \in H^1(\mathcal{C}, \Omega_{\mathcal{C}})$ vanishes in $H^0(U, R^1 \rho_* \Omega_{\mathcal{C}/U})$, we find that

$$g(p_1^*[\mathcal{D}]^{1,1} \cdot [\Gamma_s]^{n,n}) = \delta[\widetilde{\mathcal{D}}] \cdot g(\gamma) \text{ in } \Gamma(U, R^{n+1} \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^n)), \quad (1.23)$$

where $g(\gamma)$ is the image of $\gamma \in \Gamma(U, R^n \pi'_* \Omega_{\mathcal{C} \times \mathcal{Y}}^n)$ in $\Gamma(U, R^n \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* \Omega_{\mathcal{Y}}^{n-1}))$ and $\delta[\widetilde{\mathcal{D}}]$ is any lift of $[\mathcal{D}]$ in $H^1(\mathcal{C}, \rho^* \Omega_U) = \Gamma(U, \mathcal{H}_{\mathcal{C}}^{0,1} \otimes \Omega_U)$.

We now observe that for $g(\gamma) \in \Gamma(U, R^n \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* L^{s-1} \Omega_{\mathcal{Y}}^{n-1}))$, and $\delta[\widetilde{\mathcal{D}}] \in H^1(\mathcal{C}, \rho^* \Omega_U)$ we have

$$I(\delta[\widetilde{\mathcal{D}}] \cdot g(\gamma)) = w(\langle \delta[\mathcal{D}], \gamma_1 \rangle) \text{ in } \Gamma(U, R^n \pi_* \Omega_{\mathcal{Y}}^n / L^{s+1} R^n \pi_* \Omega_{\mathcal{Y}}^n). \quad (1.24)$$

In order to prove (1.24), we choose a class $\mu \in H^1(\Omega_{\mathcal{C}/U})$ which does not vanish in $H^0(U, R^1 \rho_* \Omega_{\mathcal{C}/U})$ (for example the first Chern class of a relatively ample line bundle on \mathcal{C}). The class $g(\gamma) \in \Gamma(U, R^n \pi'_*(p_1^* \Omega_{\mathcal{C}/U} \otimes p_2^* L^{s-1} \Omega_{\mathcal{Y}}^{n-1}))$ can be written as

$$g(\gamma) = p_1^* \mu \cdot p_2^* \eta + p_2^* \gamma_1,$$

with $\eta \in \Gamma(U, R^{n-1}\pi_*(L^{s-1}\Omega_{\mathcal{Y}_U}^{n-1}))$ and $\gamma_1 \in \Gamma(U, \mathcal{H}_C^{1,0} \otimes R^n\pi_*(L^{s-1}\Omega_{\mathcal{Y}_U}^{n-1}))$. We then find that

$$I(\widetilde{\delta[\mathcal{D}]} \cdot p_1^*\mu \cdot p_2^*\eta) = \rho_*(\widetilde{\delta[\mathcal{D}]} \cdot \mu) \cdot \eta$$

which is 0 since we assumed U affine so $\widetilde{\delta[\mathcal{D}]} \cdot \mu = 0$ in $H^2(\mathcal{C}, \Omega_{\mathcal{C}/U})$. On the other hand, it is clear that for

$$\widetilde{\delta[\mathcal{D}]} \in \Gamma(U, \Omega_U) \otimes H^1(\mathcal{O}_C) = \Gamma(U, \Omega_U \otimes \mathcal{H}_C^{0,1})$$

we have

$$I(\widetilde{\delta[\mathcal{D}]} \cdot p_2^*\gamma_1) = w(\langle \widetilde{\delta[\mathcal{D}]} \cdot \gamma_1 \rangle)$$

because for any $u \in H^1(\mathcal{C}, \mathcal{O}_C) = \Gamma(U, R^1\rho_*\mathcal{O}_C)$ and $\omega \in \Gamma(U, \mathcal{H}_C^{1,0})$, with cup-product $u \cdot \omega \in H^1(\mathcal{C}, \Omega_{\mathcal{C}/U})$, we have $I(u \cdot \omega) = \langle u, \omega \rangle$.

Combining (1.23) and (1.24) clearly proves the desired formula (1.22). ■

We now turn to a general family $\mathcal{Y} \rightarrow B$. The proof works in fact exactly in the same way, except that we have to replace the natural splitting of the Leray filtration on $H^n(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^n)$ given by the character decomposition by another one, which is not canonical, and depends only on the choice of a polarization on the family $\mathcal{Y} \rightarrow B$, that is a line bundle \mathcal{L} on \mathcal{Y} which is ample on the fibers \mathcal{Y}_t .

Proposition 1.11 *Assume the base B is smooth quasi-projective. Given a relative polarization \mathcal{L} on $\mathcal{Y} \xrightarrow{\pi} B$, there is a canonically induced splitting of the filtration on $H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p)$ or of the sheaves $R^q\pi_*\Omega_{\mathcal{Y}}^p$ induced by the filtration L on $\Omega_{\mathcal{Y}|\mathcal{Y}_t}^p$. This splitting is functorial with respect to pull-back maps.*

The proof of proposition 1.11 is given below.

Choosing the pulled-back line bundle $p_2^*\mathcal{L}$ on the pulled-back family $\mathcal{C} \times_U \mathcal{Y}$, we get a similar decomposition of $H^n(\mathcal{C}_t \times \mathcal{Y}_t, \Omega_{\mathcal{C} \times_U \mathcal{Y}|\mathcal{C}_t \times \mathcal{Y}_t}^n)$ which splits the filtration L on this space, relative to the projection $\mathcal{C} \times_U \mathcal{Y} \rightarrow \mathcal{C}$.

The decompositions are compatible with pull-back and push-forward maps, and the rest of the argument works the same way as before. This concludes the proof of Proposition 1.4. ■

Proof of proposition 1.11. We could do it at hand, but it is quicker to see this as a consequence of Deligne's canonical splitting result in [11]. Deligne proves that the choice of a relative polarization on $\pi : \mathcal{Y} \rightarrow B$ induces a canonical quasi-isomorphism

$$R\pi_*\mathbb{Q} \cong \bigoplus_i R^i\pi_*\mathbb{Q}.$$

It follows that there are canonical cohomology classes $\tilde{\pi}_k \in H^{2d}(\mathcal{Y} \times_B \mathcal{Y}, \mathbb{Q})$, $d = \dim \mathcal{Y}_t$ whose images in $H^0(B, R^{2d}(\pi, \pi)_*\mathbb{Q})$ are the Künneth projectors π_k :

$$\begin{aligned} R^{2d}(\pi, \pi)_*\mathbb{Q} &= \bigoplus_{i+j=2d} R^i\pi_*\mathbb{Q} \otimes R^j\pi_*\mathbb{Q} = \\ &\bigoplus_j \text{Hom}(R^j\pi_*\mathbb{Q}, R^j\pi_*\mathbb{Q}) \ni \pi_j := \text{Id}_{R^j\pi_*\mathbb{Q}}. \end{aligned}$$

From the construction of [11], it is clear that these classes $\tilde{\pi}_k \in H^{2d}(\mathcal{Y} \times_B \mathcal{Y}, \mathbb{Q})$ are Hodge classes, which means that they extend to any smooth compactification of

$\mathcal{Y} \times_B \mathcal{Y}$ and are Hodge classes there. It then follows that these classes have Dolbeault counterparts:

$$\tilde{\pi}_k^{d,d} \in H^d(\mathcal{Y} \times_B \mathcal{Y}, \Omega_{\mathcal{Y} \times_B \mathcal{Y}}^d).$$

We can now use the $\tilde{\pi}_k^{d,d}$ as acting on the relative Dolbeault cohomology of \mathcal{Y} . Namely, denoting by pr_1, pr_2 the projections from $\mathcal{Y} \times_B \mathcal{Y}$ to B , we have maps:

$$pr_{2*} \circ (\tilde{\pi}_k) \cup \circ pr_1^* : R^q \pi_* \Omega_{\mathcal{Y}}^p \rightarrow R^q \pi_* \Omega_{\mathcal{Y}}^p.$$

They similarly act on the fibers at t :

$$pr_{2*} \circ (\tilde{\pi}_k) \cup \circ pr_1^* : H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p) \rightarrow H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p).$$

These maps are compatible with the L -filtration and induce a morphism of spectral sequences associated to the filtration L . On the other hand, their action on $E_1^{s,q-s} = \Omega_{B,t}^s \otimes H^{q-s}(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p)$ is induced by π_k , hence it is equal to 0 for $p+q-s \neq k$, and the identity if $p+q-s = k$, and similarly for $E_2^{s,q-s} = E_\infty^{s,q-s}$. It follows that we have induced maps

$$pr_{2*} \circ (\tilde{\pi}_{p+q-s}) \cup \circ pr_1^* : L^s H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p) \rightarrow L^s H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p)$$

which have the property that they induce after passing to the graded Gr_L the zero map on $E_2^{s',q-s'}$ for $s' > s$, and the identity map on $E_2^{s,q-s} = E_\infty^{s,q-s}$. But then a sufficiently high power of these maps vanishes on $L^{s+1} H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p)$ and thus factors through $E_\infty^{s,q-s}$ to give a retraction

$$E_\infty^{s,q-s} \rightarrow L^s H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p),$$

hence a canonical splitting

$$L^s H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p) = L^{s+1} H^q(\mathcal{Y}_t, \Omega_{\mathcal{Y}|\mathcal{Y}_t}^p) \oplus E_\infty^{s,q-s}, \quad E_\infty^{s,q-s} = E_2^{s,q-s}.$$

■

2 1-cycles modulo algebraic equivalence in Jacobians

2.1 Constructing test elements in the dual space

We consider now a family $\mathcal{C} \rightarrow B$ of curves of genus g , and the associated Jacobian fibration $\pi : \mathcal{J} \rightarrow B$. We choose an embedding $\mathcal{C} \subset \mathcal{J}$, which provides a codimension $g-1$ cycle $\mathcal{Z} \in \text{CH}^{g-1}(\mathcal{J})/\text{alg}$ and we want to study the cycles $\mathcal{Z}_s \in \text{CH}^{g-1}(\mathcal{J})_s/\text{alg}$. Our goal is to exhibit such families with a non trivial infinitesimal invariant

$$\delta[\mathcal{Z}_s]_{\text{alg}} \in I_{s,g-1-s,g-1} := \Gamma\left(B, \frac{\text{Ker}(\bar{\nabla}_{s,g-1-s,g-1})}{\Omega_B \wedge \text{Ker}(\bar{\nabla}_{s-1,g-s-1,g-1})}\right)$$

for $s \geq 2$. Here $\bar{\nabla}$ is the $\bar{\nabla}$ -map for the Hodge bundles of the family $\mathcal{J} \rightarrow B$. The infinitesimal variation of Hodge structure on the cohomology of the fibres of the family $(\mathcal{J}_t)_{t \in B}$ of Jacobians is deduced from the one of the family of curves $(\mathcal{C}_t)_{t \in B}$,

in the obvious way, namely it coincides with the latter in degree 1, and with its k -th exterior power for higher k .

The map

$$\bar{\nabla}_{s,p,q} : \Omega_B^s \otimes \mathcal{H}^{p,q} \rightarrow \Omega_B^{s+1} \otimes \mathcal{H}^{p-1,q+1}$$

has for transposed map

$${}^t\bar{\nabla}_{s+1,g-p+1,g-q-1} : \bigwedge^{s+1} T_B \otimes \mathcal{H}^{g-p+1,g-q-1} \rightarrow \bigwedge^s T_B \otimes \mathcal{H}^{g-p,g-q}.$$

This map is given by the formula

$$\begin{aligned} & {}^t\bar{\nabla}_{s+1,g-p+1,g-q-1}(u_1 \wedge \dots \wedge u_{s+1} \otimes \omega_1 \wedge \dots \wedge \omega_{g-p+1} \otimes \eta_1 \wedge \dots \wedge \eta_{g-q-1}) \\ = & \sum_{1 \leq i \leq s+1, 1 \leq j \leq g-p+1} (-1)^{i+j} u_1 \wedge \dots \wedge \hat{u}_i \wedge \dots \wedge u_{s+1} \otimes \omega_1 \wedge \dots \wedge \hat{\omega}_j \wedge \dots \wedge \omega_{g-p+1} \\ & \otimes {}^t\bar{\nabla}_{1,1,0}(u_i \otimes \omega_j) \wedge \eta_1 \wedge \dots \wedge \eta_{g-q-1}. \end{aligned}$$

Fix a point $b \in B$. The dual of the space of infinitesimal invariants

$$\frac{\text{Ker}(\bar{\nabla}_{s,g-1-s,g-1,b} : \Omega_{B,b}^s \otimes \mathcal{H}_b^{g-1-s,g-1} \rightarrow \Omega_{B,b}^{s+1} \otimes \mathcal{H}_b^{g-2-s,g})}{\Omega_{B,b} \wedge \text{Ker}(\bar{\nabla}_{s-1,g-s-1,g-1,b} : \Omega_{B,b}^{s-1} \otimes \mathcal{H}_b^{g-1-s,g-1} \rightarrow \Omega_{B,b}^s \otimes \mathcal{H}_b^{g-2-s,g})}$$

is the quotient of the space

$$I_{s,g-1-s,g-1}^* := \text{Ker}\left(\bigwedge^s T_{B,b} \otimes \mathcal{H}_b^{s+1,1} \rightarrow T_{B,b} \otimes \frac{\bigwedge^{s-1} T_{B,b} \otimes \mathcal{H}_b^{s+1,1}}{\text{Im } {}^t\bar{\nabla}_{s,s+2,0,b}}\right) \quad (2.25)$$

by the subspace $\text{Im}({}^t\bar{\nabla}_{s+1,s+2,0,b} : \bigwedge^{s+1} T_{B,b} \otimes \mathcal{H}_b^{s+2,0} \rightarrow \bigwedge^s T_{B,b} \otimes \mathcal{H}_b^{s+1,1})$, where $\mathcal{H}_b^{p,q} = \bigwedge^p H^0(C, K_C) \otimes \bigwedge^q H^1(C, \mathcal{O}_C)$. Let us describe the simplest possible elements in the space $I_{s,g-1-s,g-1}^*$:

Lemma 2.1 *Assume we have*

1. a s -dimensional subspace $W = \langle u_1, \dots, u_s \rangle \subset T_{B,b}$,
2. a $s+1$ -dimensional subspace $K = \langle \omega_1, \dots, \omega_{s+1} \rangle \subset \mathcal{H}_b^{1,0}$ and a s -dimensional subspace $K_1 = \langle \mu_1, \dots, \mu_s \rangle \subset \mathcal{H}_b^{1,0}$,
3. an element $\eta \in \mathcal{H}_b^{0,1}$,

such that

1. For any $i \in \{1, \dots, s\}$ and $j \in \{1, \dots, s+1\}$, ${}^t\bar{\nabla}(u_i \otimes \omega_j) = 0$ in $\mathcal{H}_b^{0,1}$.
2. For any $i \in \{1, \dots, s\}$ and $j \in \{1, \dots, s\}$, ${}^t\bar{\nabla}(u_i \otimes \mu_j) = \lambda_{ij}\eta$ in $\mathcal{H}_b^{0,1}$, where the (s, s) -matrix (λ_{ij}) is invertible.

Then $w := u_1 \wedge \dots \wedge u_s \otimes \omega_1 \wedge \dots \wedge \omega_{s+1} \otimes \eta$ belongs to the subspace $I_{s,g-1-s,g-1}^*$ of (2.25).

Proof. Indeed, we may assume that $\lambda_{ij} = \delta_{ij}$, changing the basis u_i if necessary. Then the image of $u_1 \wedge \dots \wedge u_s \otimes \omega_1 \wedge \dots \wedge \omega_{s+1} \otimes \eta$ in $T_{B,b} \otimes \bigwedge^{s-1} T_{B,b} \otimes \mathcal{H}^{s+1,1}$ is equal to $\sum_i (-1)^i u_i \otimes u_1 \wedge \dots \wedge \hat{u}_i \wedge \dots \wedge u_s \otimes \omega_1 \wedge \dots \wedge \omega_{s+1} \otimes \eta$. On the other hand, we have for any i :

$$\begin{aligned} & u_1 \wedge \dots \wedge \hat{u}_i \wedge \dots \wedge u_s \otimes \omega_1 \wedge \dots \wedge \omega_{s+1} \otimes \eta \\ &= (-1)^i {}^t \bar{\nabla}_{s,s+2,0,b} (u_1 \wedge \dots \wedge u_s \otimes \mu_i \wedge \omega_1 \wedge \dots \wedge \omega_{s+1}). \end{aligned}$$

Thus $u_1 \wedge \dots \wedge u_s \otimes \omega_1 \wedge \dots \wedge \omega_{s+1} \otimes \eta \in \text{Ker} \left(\bigwedge^s T_{B,b} \otimes \mathcal{H}_b^{s+1,1} \rightarrow T_{B,b} \otimes \frac{\bigwedge^{s-1} T_{B,b} \otimes \mathcal{H}_b^{s+1,1}}{\text{Im } {}^t \bar{\nabla}_{s,s+2,0,b}} \right)$. \blacksquare

2.2 The Ikeda family

The Ikeda family [19] is simply the family of plane curves which are cyclic covers of \mathbb{P}^1 of degree d ramified along a degree d divisor. In other words, the general equation f of such a plane curve takes the form (in homogeneous coordinates Y, X_0, X_1 on \mathbb{P}^2):

$$f(Y, X_0, X_1) = Y^d - f_0(X_0, X_1)$$

for some homogeneous polynomial f_0 of degree d . This family is parameterized by the quotient $U/PGL(2)$ where $U \subset \mathbb{P}(H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(d)))$ is the open set parameterizing reduced divisors, hence smooth plane curves.

The tangent space to this quotient at a general point f_0 is the quotient

$$R_{f_0}^d = S_X^d / J_{f_0}^d,$$

where $S_X^k := H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(k))$ and $J_{f_0}^k \subset S^k$ is the degree k part of the Jacobian ideal of f_0 , generated by the partial derivatives $\frac{\partial f_0}{\partial X_i}$.

We consider the universal family $\mathcal{C} \rightarrow B$, where $B \subset U$ is any slice for the action of $PGL(2)$ and the corresponding infinitesimal variation of Hodge structure at a point $b \in B$ corresponding to an equation f as above. Let us exhibit for any such f data as in Lemma 2.1 with $s = d - 3$.

We recall from [6] (see also [33, 6.2.1]) that the infinitesimal variation of Hodge structure of an hypersurface defined by a homogeneous equation f is governed by the product in the Jacobian ring R_f of f , quotient of the polynomial ring by the ideal generated by the partial derivatives of f . In the case of a curve C of degree d in \mathbb{P}^2 defined by an equation f , we get

$$H^{1,0}(C) \cong R_f^{d-3}, \quad H^{0,1}(C) \cong R_f^{2d-3}$$

and the infinitesimal variation of Hodge structure restricted to the family of plane deformations of C identifies

$${}^t \bar{\nabla}_{1,1,0} : T_{B,b} \otimes H^{1,0}(C) \rightarrow H^{0,1}(C)$$

to the multiplication map

$$R_f^d \otimes R_f^{d-3} \rightarrow R_f^{2d-3},$$

where B is a slice for the $PGL(3)$ -action and $b \in B$ is the parameter for C .

Let now $f = Y^d - f_0(X_0, X_1)$, which is homogeneous of degree d in the three variables Y, X_0, X_1 . Note that $\frac{\partial f}{\partial X_i} = -\frac{\partial f_0}{\partial X_i}$ for $i = 0, 1$, so that R_f is a module over R_{f_0} .

Consider the following subspaces:

$$W := R_{f_0}^d \subset R_f^d, K := R_{f_0}^{d-3} = S_X^{d-3} \quad (2.26)$$

and $K_1 := YR_{f_0}^{d-4} = YS_X^{d-4}$. Note that $\dim W = d - 3$, $\dim K = d - 2$, $\dim K_1 = d - 3$. We have

Lemma 2.2 (i) For any $u \in W, \omega \in K$, we have ${}^t\bar{\nabla}(u \otimes \omega) = 0$.

(ii) There is an element $\eta \in H^{0,1}(C)$, such that the map $W \otimes K_1 \rightarrow H^{0,1}(C)$, $u \otimes \omega \mapsto {}^t\bar{\nabla}(u \otimes \omega)$ takes the form

$${}^t\bar{\nabla}(u \otimes \omega) = \lambda(u, \omega)\eta,$$

where the bilinear form λ gives a perfect pairing between W and K_1 .

Proof. (i) This follows from the fact that $R_{f_0}^{2d-3} = 0$.

(ii) We choose for η the generator of the 1-dimensional vector space $YR_{f_0}^{2d-4}$, which means geometrically (cf. [6]) that

$$\eta = \text{Res}_C \frac{P(X_0, X_1)Y\Omega}{f^2} \text{ in } H^{0,1}(C), \quad (2.27)$$

with $\Omega = X_0dX_1 \wedge dY - X_1dX_0 \wedge dY + YdX_0 \wedge dX_1 \in H^0(\mathbb{P}^2, K_{\mathbb{P}^2}(3))$ and P generic of degree $2d - 4$ in the variables X_0, X_1 .

Then (ii) follows from Macaulay's theorem (cf. [33, 6.2.2]), which says that $R_{f_0}^{2d-4} = \mathbb{C}$ and the multiplication map $R_{f_0}^d \otimes R_{f_0}^{d-4} \rightarrow R_{f_0}^{2d-4}$ is a perfect pairing. ■

We get now the following corollary. Here we consider the Ikeda family $\mathcal{C} \rightarrow B$ (defined over a slice of the quotient map $\mathbb{P}(S_X^d) \rightarrow \mathbb{P}(S_X^d)/PGl(2)$), the associated Jacobian fibration $\pi : \mathcal{J} \rightarrow B$, an embedding $\mathcal{C} \rightarrow \mathcal{J}$ giving rise to a codimension $g - 1$ cycle \mathcal{Z} of \mathcal{J} and its (relative) Beauville component \mathcal{Z}_{d-3} .

Corollary 2.3 Choose a basis u_1, \dots, u_{d-3} of W and a basis $\omega_1, \dots, \omega_{d-2}$ of K . Let η be a generator of $YR_{f_0}^{2d-4} = \mathbb{C} \subset H^1(C, \mathcal{O}_C)$, where the curve C is defined by $Y^d = f_0(X_0, X_1)$. Then

$$w := u_1 \wedge \dots \wedge u_{d-3} \otimes \omega_1 \wedge \dots \wedge \omega_{d-2} \otimes \eta \quad (2.28)$$

belongs to the space $I_{d-3, g-d+2, g-1}^*$, $g = \frac{d(d-3)}{2} + 1$.

Proof. We only have to observe that the space $W \subset H^1(C, T_C)$ is exactly the tangent space to the Ikeda family, so that w belongs to $\bigwedge^{d-3} T_{B,b} \otimes H^{d-2,1}(JC)$. Applying Lemmas 2.1 and Lemma 2.2, we also get that the image of w in $T_{B,b} \otimes \bigwedge^{d-4} T_{B,b} \otimes H^{d-2,1}(JC)$ vanishes in the quotient

$$T_{B,b} \otimes \frac{\bigwedge^{d-4} T_{B,b} \otimes H^{d-2,1}(JC)}{\text{Im } {}^t\bar{\nabla}_{d-3, d-1, 0}},$$

which proves the result. ■

2.3 End of the proof of Theorem 0.6

This section is devoted to the proof of the following result:

Proposition 2.4 *Let $f_0 \in S_X^d$ define a smooth divisor in \mathbb{P}^1 . Let C be the curve defined by $Y^d - f_0(X_0, X_1)$ and let $w \in I_{d-3, g-d+2, g-1}^*$ be the element constructed in Corollary 2.3. Then, the pairing*

$$\langle \delta[\mathcal{Z}_{d-3}]_{b,alg}, w \rangle$$

is nonzero, where $b \in B$ is the point of the Ikeda family $\mathcal{C} \rightarrow B$ parameterizing C .

Note that Proposition 2.4 implies Theorem 0.6. Indeed, it implies that for the Ikeda family $\mathcal{C} \rightarrow B$, the infinitesimal invariant $\delta[\mathcal{Z}_{d-3}]_{b,alg}$ of the corresponding codimension $g-1$ cycle \mathcal{Z} of $\mathcal{J} \rightarrow B$ is nonzero at the general point of B , which implies by Theorem 0.1 that for the very general curve C parameterized by B (hence for the very general plane curve), the Beauville component Z_{d-3} is non zero in $\text{CH}_1(JC)_{d-3}/alg$.

The proof of Proposition 2.4 will use the following description of the pairing $\langle \delta[\mathcal{Z}_{d-3}]_{b,alg}, w \rangle$. Let $\mathcal{C} \subset \mathcal{J} \rightarrow B$ be a family of curves of genus g and let $w \in I_{s, g-1-s, g-1}^*$ be an element constructed as in Lemma 2.1. Assume for simplicity that $\dim B = s$ so that $W = T_{B,b}$, where the point b parameterizes the curve $C := \mathcal{C}_b$. Then the pairing

$$\langle \delta[\mathcal{Z}_s]_{b,alg}, w \rangle$$

is computed as follows : Since all the form $\omega_j \in H^0(C, K_C)$ satisfy ${}^t\bar{\nabla}(u_i \otimes \omega_j) = 0$ in $H^1(C, \mathcal{O}_C)$, they lift to sections of

$$H^0(C, \Omega_{\mathcal{C}|C}) = H^0(C, \Omega_{\mathcal{J}|C}) = H^0(J, \Omega_{\mathcal{J}|J}),$$

where $J = JC = \mathcal{J}_b$. For any k , the multiplication μ_k induces an endomorphism μ_k^* of $H^0(J, \Omega_{\mathcal{J}|J})$ which is compatible with the restriction map

$$H^0(J, \Omega_{\mathcal{J}|J}) \rightarrow H^0(J, \Omega_J),$$

the action of μ_k^* on the right hand term being the homothety of factor k . It follows that there is a unique lift $\tilde{\omega}_j$ of ω_j in $H^0(J, \Omega_{\mathcal{J}|J})$ satisfying the property

$$\mu_k^*(\tilde{\omega}_j) = k\tilde{\omega}_j \text{ for any } k.$$

The $d-2$ -form

$$\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{s+1} \in H^0(C, \Omega_{\mathcal{J}|C}^{s+1})$$

restricts to a form $\mu \in H^0(C, \Omega_{\mathcal{C}|C}^{s+1})$. There is an isomorphism

$$\Omega_{\mathcal{C}|C}^{s+1} \cong K_C$$

which becomes canonical once $\bigwedge^s W$ is trivialized and we trivialize $\bigwedge^s W$ using the multivector $u_1 \wedge \dots \wedge u_s$. We thus get from w an element $\mu \cdot \eta \in H^1(C, K_C)$.

Lemma 2.5 *We have*

$$\langle \delta[\mathcal{Z}_s]_{b,alg}, w \rangle = \int_C \mu \cdot \eta.$$

Proof. In a general situation $\mathcal{C} \hookrightarrow \mathcal{J} \rightarrow B$, the class $[\mathcal{Z}]_b^{g-1, g-1} \in H^{g-1}(JC, \Omega_{\mathcal{J}|JC}^{g-1})$ is the Dolbeault cohomology class of $\mathcal{C} \subset \mathcal{J}$ and the class $[\mathcal{Z}]_{s, alg, b}$ is its component lying in the space

$$H^{g-1}(JC, \Omega_{\mathcal{J}|JC}^{g-1})_s := \{\alpha \in H^{g-1}(JC, \Omega_{\mathcal{J}|JC}^{g-1}), \mu_k^* \alpha = k^{2g-2-s}, \forall k\}.$$

For any class β in the dual space

$$H^1(JC, \Omega_{\mathcal{J}|JC}^{N+1}), \quad N := \dim B$$

we have

$$\langle [\mathcal{C}]_b^{g-1, g-1}, \beta \rangle = \int_{\mathcal{C}} \beta|_{\mathcal{C}}, \quad (2.29)$$

where $\beta|_{\mathcal{C}}$ is the image of β in $H^1(C, \Omega_{\mathcal{C}|C}^{N+1})$ which is identified to $H^1(C, K_C)$ via a trivialization of $\bigwedge^N \Omega_{B, b}$. Coming back to our situation above, so $N = s$, the statement thus follows from the following facts:

- 1) The choice of lift $\tilde{\omega}_j \in H^0(\Omega_{\mathcal{J}|JC})_1$ gives the unique lift

$$\tilde{w} := \tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{s+1} \otimes \eta \in H^1(JC, \Omega_{\mathcal{J}^{s+1}|JC})_{s+2}$$

of $w \in Gr_L^{s+2} H^1(JC, \Omega_{\mathcal{J}|JC}^{s+1})$.

- 2) Writing $[\mathcal{Z}]^{g-1, g-1} = \sum_i [\mathcal{Z}_i]^{g-1, g-1}$, then we have

$$\langle [\mathcal{Z}]_b^{g-1, g-1}, \tilde{w} \rangle = \langle [\mathcal{Z}_s]_b^{g-1, g-1}, \tilde{w} \rangle, \quad (2.30)$$

since the coupling is multiplied by k^{2g} under μ_k^* , which implies that all the other terms $\langle [\mathcal{Z}_{s'}]^{g-1, g-1}, \tilde{w} \rangle$ vanish.

The term on the right in (2.30) gives a coupling $\langle \delta[\mathcal{Z}_s]_b^{g-1, g-1}, w \rangle$ which, by the fact that w is in the kernel $I_{d-3, g-d+2, g-1}^*$ of (2.25), is equal to the desired coupling $\langle \delta[\mathcal{Z}_s]_{b, alg}^{g-1, g-1}, w \rangle$. On the other hand, the term on the left is computed by formula (2.29). The equality (2.30) thus proves Lemma 2.5. \blacksquare

Proof of Proposition 2.4. We follow the description given in Lemma 2.5. As ${}^t\bar{\nabla}(u_i \otimes \omega_j) = 0$ for all i, j , each ω_j has a canonical lift $\tilde{\omega}_j \in H^0(J, \Omega_{\mathcal{J}|J})$. This lift is determined by the condition that $\mu_k^* \tilde{\omega}_j = k \tilde{\omega}_j$ for all k .

We observe that the cyclic group $G := \mathbb{Z}/d\mathbb{Z}$ with generator g identified to a d -th primitive root of unity ζ acts over B on the families \mathcal{C} and \mathcal{J} , in a compatible way if we embed \mathcal{C} in \mathcal{J} using one of the fixed points of the action of G on the curves. We claim that the forms $\tilde{\omega}_j$ satisfy

$$g^* \tilde{\omega}_j = \zeta \tilde{\omega}_j. \quad (2.31)$$

To see this, we write the exact sequence

$$0 \rightarrow \Omega_{B, b} \otimes \mathcal{O}_{\mathcal{C}} \rightarrow \Omega_{\mathcal{C}|B} \rightarrow \Omega_{\mathcal{C}} \rightarrow 0,$$

which induces the exact sequence

$$0 \rightarrow \Omega_{B, f} \rightarrow H^0(C, \Omega_{\mathcal{J}|C}) \rightarrow K \rightarrow 0 \quad (2.32)$$

with $K \subset H^{1,0}(C)$ defined as in (2.26). This follows indeed from Macaulay's theorem, which implies that $K = \{\alpha \in H^0(\Omega_C), \bar{\nabla}(\alpha) = 0 \text{ in } \Omega_{B,b} \otimes H^1(C, \mathcal{O}_C)\}$. Then the claim follows from the fact that the ω_j , which are the images of $\tilde{\omega}_j$ in $H^0(J, \Omega_{\mathcal{J}/B|J}) = H^0(J, \Omega_J) = H^{1,0}(C)$, satisfy the property

$$g^*\omega_j = \zeta\omega_j,$$

and from the fact that the two actions μ_k^* and g^* on the space

$$H^0(J, \Omega_{\mathcal{J}|J}) = H^0(C, \Omega_{\mathcal{J}|C}) = H^0(C, \Omega_{C|C}),$$

commute, both acting trivially on the space on the left in (2.32), and as a homothety of factor k , resp. ζ on the space on the right in (2.32).

As an immediate consequence of (2.31), we find that

$$g^*(\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{d-2}) = \zeta^{d-2} \tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{d-2} \text{ in } H^0(\bigwedge^{d-2} \Omega_{\mathcal{J}|C}),$$

hence a fortiori the restricted form μ which is the image of $\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{d-2}$ in $H^0(\bigwedge^{d-2} \Omega_{C|C}) = H^{1,0}(C)$ satisfies

$$g^*\mu = \zeta^{d-2}\mu.$$

Next we observe that the form $\eta \in H^{0,1}(C)$ defined in (2.27) satisfies

$$g^*\eta = \zeta^2\eta$$

as follows immediately from the formula

$$\eta = \text{Res}_C \frac{PY\Omega}{f^2},$$

with $g^*Y = \zeta Y$, $g^*\Omega = \zeta\Omega$, $g^*f = f$, $g^*P = P$.

The two spaces $H^{1,0}(C)^{\zeta^{d-2}}$ and $H^{0,1}(C)^{\zeta^2}$ where g^* acts respectively by multiplication by ζ^{d-2} and ζ^2 are dual and 1-dimensional and the second space is generated by η . It follows that the pairing $\langle \delta[\mathcal{Z}_{d-3}]_{b,alg}, w \rangle$, which by Lemma 2.5 is computed as the pairing $\langle \mu, \eta \rangle$ between $\mu \in H^{1,0}(C)$ and $\eta \in H^{0,1}(C)$, vanishes if and only if μ vanishes identically.

It thus suffices to prove that μ is nonzero. This is done as follows: Recall that the bundle $\Omega_{C|C}$ has rank $d-2$ and possesses the $d-2$ sections $\tilde{\omega}_j$. Assume that μ vanishes in $H^0(C, \bigwedge^{d-2} \Omega_{C|C}) = H^0(C, K_C)$. Then these sections generate a subbundle $N \subset \Omega_{C|C}$ of rank $\leq d-3$, and it follows that if $x \in C$ is a general point, there is a section $\tilde{\omega}_x \in \langle \tilde{\omega}_1, \dots, \tilde{\omega}_{d-2} \rangle$ vanishing at x . As all the sections $\tilde{\omega}_j$ satisfy $g^*\tilde{\omega}_j = \zeta\tilde{\omega}_j$, it follows that $\tilde{\omega}_x$ also vanishes at $gx, g^2x, \dots, g^{d-1}x$, so that $\tilde{\omega}_x$ is in fact a section of $\Omega_{C|C}(-1)$. It is easy to check however using the multiplication in the Jacobian ring R_f that $H^0(C, \Omega_{C|C}(-1)) = 0$. Indeed, we have the exact sequence

$$0 \rightarrow \Omega_{B,b} \otimes \mathcal{O}_C(-1) \rightarrow \Omega_{C|C}(-1) \rightarrow K_C(-1) \rightarrow 0$$

and the induced map

$$H^0(C, K_C(-1)) \rightarrow \Omega_{B,b} \otimes H^1(C, \mathcal{O}_C(-1)) = \text{Hom}(R_{f_0}^d, R_f^{2d-4})$$

is given by multiplication in the Jacobian ring of f . Such a section would provide a nonzero element of

$$R_f^{d-4} = \bigoplus_{i=0}^{d-1} Y^i R_{f_0}^{d-4-i}$$

which is annihilated by multiplication by all elements of $W = R_{f_0}^d$, and this does not exist by Macaulay's Theorem. ■

3 Comparison with Ikeda's invariants

As already mentioned in the introduction, Ikeda [19] proves the nonvanishing of the cycle Z_{d-3} for the very general curve in the Ikeda family of plane curves of degree d , modulo a certain subgroup of $\text{CH}^{g-1}(JC)$. Here the cycle Z_{d-3} is obtained by embedding of the curve C in JC using a general point $p \in JC$.

The infinitesimal invariant used by Ikeda is the projection of

$$\delta[\mathcal{Z}_{d-3,b}] \in \frac{\text{Ker } \bar{\nabla}_{d-3,g-d+2,g-1,b} \subset \Omega_{B,b}^{d-3} \otimes H^{g-d+2,g-1}(JC)}{\text{Im } \bar{\nabla}_{d-4,g-d+3,g-2}}$$

modulo the subspace

$$\Omega_{B,b}^{d-3} \otimes \text{Ker } \bar{\nabla}_{0,g-d+2,g-1,b} \subset \text{Ker } \bar{\nabla}_{d-3,g-d+2,g-1,b}. \quad (3.33)$$

In contrast, our infinitesimal invariant $\delta[\mathcal{Z}_{d-3,alg,b}]$ is the projection of

$$\delta[\mathcal{Z}_{d-3,b}] \in \frac{\text{Ker } \bar{\nabla}_{d-3,g-d+2,g-1,b} \subset \Omega_{B,b}^{d-3} \otimes H^{g-d+2,g-1}(JC)}{\text{Im } \bar{\nabla}_{d-4,g-d+3,g-2}}$$

modulo the subspace

$$\Omega_{B,b} \wedge \text{Ker } \bar{\nabla}_{d-4,g-d+2,g-1,b} \subset \text{Ker } \bar{\nabla}_{d-3,g-d+2,g-1,b} \quad (3.34)$$

which contains (3.33) by formula (1.7).

If we consider (instead of the family of plane curves) the universal family of curves $\mathcal{C} \rightarrow B$ of genus g , and replace $d-3$ by any integer s such that $g \geq 2s+1$, (this is the range we are interested in by Conjecture 0.4) then at the general point $b \in B$, the subspace (3.33) can be in fact explicitly computed, namely:

Proposition 3.1 *Let $C = C_b$ be a general curve of genus $g \geq 2s+1$. Then the kernel*

$$\text{Ker}(\bar{\nabla}_{0,g-s-1,g-1,b} : H^{g-s-1,g-1}(JC) \rightarrow \Omega_{B,b} \otimes H^{g-s-2,g}) \quad (3.35)$$

is equal to the image

$$\text{Im}([C]_* : H^{g-s,g}(JC) \rightarrow H^{g-s-1,g-1}(JC)),$$

where $[C]_*$ is the Pontryagin product with the class of C in $H^{g-1,g-1}(JC)$.

Notice that $\text{Im}([C]* : H^{g-s,g}(JC) \rightarrow H^{g-s-1,g-1}(JC))$ is also equal to the non-primitive part of the cohomology

$$\text{Im}(\theta^{g-s-1} : H^{0,s}(JC) \rightarrow H^{g-s-1,g-1}(JC)).$$

Proof of Proposition 3.1. The proposition above is in fact a geometric translation of our result in [35], namely the vanishing of the Koszul cohomology group $K_{k,1}(C, K_C)$ for a general curve of genus g in the range $g \leq 2k + 1$. This vanishing says that in this range, the kernel of the Koszul differential

$$\text{Ker}\left(\bigwedge^k H^0(C, K_C) \otimes H^0(C, K_C) \rightarrow \bigwedge^{k-1} H^0(C, K_C) \otimes H^0(C, 2K_C)\right) \quad (3.36)$$

is equal to the image of the natural map (also a Koszul differential)

$$\bigwedge^{k+1} H^0(C, K_C) \rightarrow \bigwedge^k H^0(C, K_C) \otimes H^0(C, K_C).$$

If we now identify $H^0(C, K_C)$ with $\bigwedge^{g-1} H^{0,1}(C) = \bigwedge^{g-1} H^{0,1}(C)$, the Koszul differential (3.36) identifies to the map $\bar{\nabla}_{0,k,g-1}$, $k = g - s - 1$ of (3.35) because, as is well known, the multiplication map

$$H^0(C, K_C) \otimes H^0(C, K_C) \rightarrow H^0(C, 2K_C)$$

seen in the form

$$H^0(C, K_C) \rightarrow \text{Hom}(H^0(C, K_C), H^0(C, 2K_C)) = H^{0,1}(C) \otimes H^0(C, 2K_C)$$

identifies to the map $\bar{\nabla}_{0,1,0}$.

For $g \geq 2s + 1$, we have $g \leq 2(g - s - 1) + 1$. It thus follows from the above that the kernel of $\bar{\nabla}_{0,g-s-1,g-1}$ is equal to the image of $\bigwedge^{g-s} H^0(C, K_C)$. We finally observe that the natural map

$$\bigwedge^{g-s} H^0(C, K_C) \rightarrow \bigwedge^{g-s-1} H^0(C, K_C) \otimes H^0(C, K_C)$$

identifies to the map $[C]* : H^{g-s,g}(JC) \rightarrow H^{g-s-1,g-1}(JC)$ given by Pontryagin product with the class of C .

We refer to [2, 8.1.4] for a more detailed version of this Hodge theoretic interpretation of the vanishing of canonical syzygies. \blacksquare

Proposition 3.1 shows that the Ikeda subspace (3.33) is too small to make the Ikeda infinitesimal invariant of the cycles $\mathcal{Z}_s \in \text{CH}^{g-1}(\mathcal{J})$ for $g \geq 2s + 1$ invariant under relative translation of the cycle. Indeed, choose a degree 1 codimension g cycle Γ in \mathcal{J} , and consider the cycle

$$\mathcal{Z}_\Gamma := \mathcal{Z} * \Gamma,$$

where $*$ is the relative Pontryagin product over B for cycles in \mathcal{J} . Then

$$\mathcal{Z}_\Gamma - \mathcal{Z} = \mathcal{Z} * (\Gamma - 0_B),$$

where 0_B is the zero section of \mathcal{J} , and since the 0-cycles $\Gamma_b - 0_b \in \text{CH}_0(\mathcal{J}_b)$ are of degree 0, they are algebraically equivalent to 0, and so are the cycles $\mathcal{Z}_{\Gamma,b} - \mathcal{Z}_b$.

If we now introduce the relative Beauville decomposition

$$\Gamma = \sum_i \Gamma_i, \Gamma_0 = 0_B$$

of Γ , we find that

$$\mathcal{Z}_{\Gamma,s} = \sum_{i \leq s} \mathcal{Z}_{s-i} * \Gamma_i.$$

In this sum, the first term is \mathcal{Z}_s and the last term is $\mathcal{Z}_0 * \Gamma_s$, and its restriction to \mathcal{J}_b over the point $b \in B$ belongs by definition to the subgroup $Z_0 F^s \text{CH}_{g-1}(\mathcal{J}_b)$. So its infinitesimal invariant belongs by Ikeda's results to the Ikeda subspace (3.33). The other terms however have infinitesimal invariants which can be shown, using Proposition 3.1 and Ceresa's Theorem, not to belong to the Ikeda subspace (3.33) for $g \geq 2s + 1$, (while of course they belong to the subspace (3.34) by Proposition 1.4, since $\mathcal{Z}_{s-i} * \Gamma_i$ is algebraically equivalent to 0 for $i > 0$).

References

- [1] E. Arbarello, M. Cornalba, P. Griffiths, J. Harris. Geometry of algebraic curves. Vol. I. Grundlehren der Mathematischen Wissenschaften 267. Springer-Verlag, New York, 1985.
- [2] M. Aprodu, J. Nagel. *Koszul cohomology and algebraic geometry*, University Lecture Series, 52. American Mathematical Society, Providence, RI, (2010).
- [3] A. Beauville. Sur l'anneau de Chow d'une variété abélienne. Math. Ann. 273 (1986), no. 4, 647–651.
- [4] A. Beauville. Algebraic cycles on Jacobian varieties, Compositio Math. 140 (2004), 683–688.
- [5] A. Beilinson. Height pairing between algebraic cycles. *K-theory, arithmetic and geometry* (Moscow, 1984–1986), 1–25, Lecture Notes in Math., 1289, Springer, Berlin, 1987.
- [6] J. Carlson, P. Griffiths. Infinitesimal variations of Hodge structure and the global Torelli theorem, in *Géométrie algébrique, Angers 1980*, (Ed. A. Beauville), Sijthoff-Noordhoff, 51–76.
- [7] G. Ceresa. C is not algebraically equivalent to C^- in its Jacobian. Ann. of Math. (2) 117 (1983), no. 2, 285–291.
- [8] A. Collino, G. P. Pirola. The Griffiths infinitesimal invariant for a curve in its Jacobian. Duke Math. J. 78 (1995), no. 1, 59–88.
- [9] E. Colombo, B. Van Geemen. Note on curves in a Jacobian. Compositio Math. 88 (1993), no. 3, 333–353.
- [10] P. Deligne. Théorie de Hodge II, Inst. Hautes Études Sci. Publ. Math. **40** (1971), 5–57.

- [11] P. Deligne. Décompositions dans la catégorie dérivée. *Motives* (Seattle, WA, 1991), 115–128, Proc. Sympos. Pure Math., 55, Part 1, Amer. Math. Soc., Providence, RI, 1994.
- [12] P. Deligne. Théorème de Lefschetz et critères de dégénérescence de suites spectrales, *Inst. Hautes Études Sci. Publ. Math.* No. 35 (1968) 259–278.
- [13] Ch. Deninger, J. Murre. Motivic decomposition of abelian schemes and the Fourier transform. *J. Reine Angew. Math.* 422 (1991), 201–219.
- [14] H. Esnault, K. Paranjape. Remarks on absolute de Rham and absolute Hodge cycles. *C. R. Acad. Sci. Paris Sér. I Math.* 319 (1994), no. 1, 67–72.
- [15] N. Fakhruddin, Algebraic cycles on generic Abelian varieties, *Compositio Math.* 100 (1996), 101–119.
- [16] Ph. Griffiths. On the periods of certain rational integrals. I, II. *Ann. of Math.* (2) 90 (1969), 460–495; *ibid.* (2) 90 1969 496–541.
- [17] Ph. Griffiths. Infinitesimal variations of Hodge structure. III. Determinantal varieties and the infinitesimal invariant of normal functions. *Compositio Math.* 50 (1983), no. 2-3, 267–324.
- [18] F. Herbaut. Algebraic cycles on the Jacobian of a curve with a linear system of given dimension. *Compos. Math.* 143 (2007), no. 4, 883–899.
- [19] A. Ikeda. Algebraic cycles and infinitesimal invariants on Jacobian varieties, *J. Algebraic Geom.* 12 (2003), 573–603.
- [20] G. Marini. Tautological cycles on Jacobian varieties. *Collect. Math.* 59 (2008), no. 2, 167–190.
- [21] B. Moonen. Relations between tautological cycles on Jacobians, *Comment. Math. Helv.* 84 (2009), no. 3, 471–502.
- [22] J. P. Murre. On the motive of an algebraic surface. *J. Reine Angew. Math.* 409 (1990), 190–204.
- [23] M. Nori. Algebraic cycles and Hodge-theoretic connectivity. *Invent. Math.* 111 (1993), no. 2, 349–373.
- [24] C. Peters, J. Steenbrink. *Mixed Hodge structures*, *Ergebnisse der Mathematik und ihrer Grenzgebiete.* 3, 52. Springer-Verlag, Berlin, 2008.
- [25] A. Polishchuk. Algebraic cycles on the relative symmetric powers and on the relative Jacobian of a family of curves. I. *Selecta Math. (N.S.)* 13 (2007), no. 3, 531–569.
- [26] G. P. Pirola, C. Rizzi. Infinitesimal invariant and vector bundles. *Nagoya Math. J.* 186 (2007), 95–118.
- [27] S. Saito. Higher normal functions and Griffiths groups, *J. Algebraic Geom.* 11 (2002), 161–201.

- [28] S. Saito. Motives and filtration on Chow groups, *Invent. Math.* 125 (1996), 149–196.
- [29] G. Van der Geer, A. Kouvidakis. Cycle relations on Jacobian varieties. With an appendix by Don Zagier, *Compositio Math.* 143 (2007), 900–908.
- [30] C. Voisin. Variations of Hodge structure and algebraic cycles, *Proceedings of the international congress of mathematicians, Zürich, (1994)*
- [31] C. Voisin. Variations de structure de Hodge et zéro-cycles sur les surfaces générales, *Mathematische Annalen* 299, 77-103 (1994).
- [32] C. Voisin. Une remarque sur l’invariant infinitésimal des fonctions normales, *C.R.A.S. Paris, t.307, Série 1*, 157-160 (1988).
- [33] C. Voisin. *Hodge Theory and complex algebraic geometry II*, Cambridge University Press 2003.
- [34] C. Voisin. *Chow rings, decomposition of the diagonal and the topology of families*, *Annals of Mathematics Studies, Number 187*, Princeton University Press (2014).
- [35] C. Voisin. Green’s canonical syzygy conjecture for generic curves of odd genus. *Compos. Math.* 141 (2005), no. 5, 1163–1190.
- [36] C. Voisin. Remarks on filtrations on Chow groups and the Bloch conjecture. *Ann. Mat. Pura Appl. (4)* 183 (2004), no. 3, 421–438.