

A geometric application of Nori's connectivity theorem

Claire Voisin

Institut de mathématiques de Jussieu, CNRS, UMR 7586

0 Introduction

Our purpose in this paper is to contribute to the study of rational maps from r -dimensional varieties to general hypersurfaces in projective space (cf [4], [18], [10], [3]). In the last section, we shall eventually extend this to the study of correspondences instead of rational maps. The problem we consider is the following : *given a family $\mathcal{Y} \rightarrow \mathcal{S}$ of r -dimensional smooth projective varieties, when a general hypersurface X of degree d in projective space \mathbb{P}^{n+1} is swept out by images of rational maps from one member of this family to X ?*

Our approach to this problem is Hodge theoretic. Unlike [4], [18], [10], [3], the result has nothing to do with the canonical bundle of the varieties Y_t , $t \in \mathcal{S}$. Instead, our answer will depend only on the dimension of the moduli space \mathcal{S} . Roughly speaking, the idea is as follows : assume that $\dim \mathcal{S}$ is small, but the general X is covered by images of rational maps from one member of this family to X ; then there is a universal dominating rational map

$$\begin{array}{ccc} \Phi : \mathcal{K} & \dashrightarrow & \mathcal{X}_U \\ \downarrow \pi & & \downarrow \\ \mathcal{B} & \rightarrow & U \end{array} ,$$

where $\pi : \mathcal{K} \rightarrow \mathcal{B}$ is induced by the family $\mathcal{Y} \rightarrow \mathcal{S}$, and $\mathcal{X} \rightarrow U$ is the universal family of hypersurfaces of degree d in \mathbb{P}^{n+1} . Here U is the open set of $\mathbb{P}(H^0(\mathbb{P}^{n+1}, \mathcal{O}_{\mathbb{P}^{n+1}}(d)))$ parameterizing smooth hypersurfaces.

If we fix a point $s \in \mathcal{S}$, we have now \mathcal{B}_s , which is the fiber of the map $\mathcal{B} \rightarrow \mathcal{S}$, and $\mathcal{X}_{\mathcal{B}_s}$, which is the fibered product

$$\mathcal{B}_s \times_U \mathcal{X}.$$

The Φ above restricts to

$$\Phi_s : Y_s \times \mathcal{B}_s \rightarrow \mathcal{X}_{\mathcal{B}_s}.$$

The point now is that if $\dim \mathcal{S}$ is small, then the corank of the map $\mathcal{B}_s \rightarrow U$ is small, so that an adaptation of Nori's connectivity theorem [16] will show that the cohomology groups of \mathcal{X}_s , modulo the cohomology of $\mathbb{P}^{n+1} \times U_s$, are

0 or have small Hodge level. Studying the cohomology class of the graph of Φ in

$$H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q})$$

via the Leray spectral sequence of the map

$$\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}} \rightarrow \mathcal{S},$$

with fiber $Y_s \times \mathcal{X}_{\mathcal{B}_s}$, we will deduce a contradiction with the fact that our initial Φ was dominating.

Let us now state our precise results :

Definition 1 *Let $\mathcal{Y} \rightarrow \mathcal{S}$ be a family of r -dimensional smooth projective varieties. We say that a n -dimensional variety X is rationally swept out by varieties parametrized by \mathcal{S} , if there exist a quasiprojective variety B of dimension $n - r$, a family $K \rightarrow B$ which is the pull-back of the family \mathcal{Y} via a morphism $\psi : B \rightarrow \mathcal{S}$, and a dominant rational map*

$$\phi : K \dashrightarrow X,$$

(which is necessarily generically finite on the generic fiber K_b since $\dim K = \dim X$).

Remark 1 *One can show that this property is equivalent to the fact that the union of the images of generically finite rational maps from a variety Y_t , $t \in \mathcal{S}$, to X contains a Zariski open set of X .*

With this definition, we show :

Theorem 1 *Fix an integer $1 \leq r \leq n$. Let $\gamma = \frac{r-1}{2}$, r odd, or $\gamma = \frac{r}{2}$, r even, that is γ is the round-up of $\frac{r-1}{2}$. Let $\mathcal{Y} \rightarrow \mathcal{S}$, $\dim \mathcal{S} = C$, be a family of r -dimensional smooth projective varieties. Then the general hypersurface of degree d in \mathbb{P}^{n+1} is not rationally swept out by varieties parameterized by \mathcal{S} if*

$$(d+1)r \geq 2n + C + 2, \quad (\gamma+1)d \geq 2n - r + 1 + C. \quad (0.1)$$

(Note that except for $r = 1$, the second inequality implies the first.)

Remark 2 *One could of course prove a similar statement for sufficiently ample hypersurfaces in any smooth variety. In the case of projective space, the estimates on d are sharp, and allow applications to the Calabi-Yau case, which is never considered in the papers quoted above (see section 3).*

So let us consider the case of Calabi-Yau hypersurfaces. Because of the Lang conjecture that we shall recall below, we will be mainly interested into the case where the family $\mathcal{Y} \rightarrow \mathcal{S}$ is a family of abelian varieties. Our theorem 1 implies in this case :

Theorem 2 *Let X be a general Calabi-Yau hypersurface in projective space \mathbb{P}^{n+1} , that is $d = n + 2$. Then X is not rationally swept out by r -dimensional abelian varieties, for any $r \geq 2$.*

In the paper [14], Lang formulates a number of conjectures concerning smooth projective complex varieties X . One of them is that the analytic closure of the union of the images of holomorphic maps from \mathbb{C} to X is equal to the union of the images of non constant rational maps from an abelian variety to X . Another one is that this locus is equal to X itself if and only if X is not of general type.

Next, by a standard countability argument for Chow varieties (similar to the one given in the beginning of section 2), we see that, according to these conjectures, if X is not of general type, there should exist a quasiprojective variety B , a family $K \rightarrow B$ of abelian varieties, and a dominating rational map

$$\phi : K \dashrightarrow X,$$

which is non constant on the generic fiber K_b , $b \in B$.

Let us now consider the case where X is a Calabi-Yau variety, that is K_X is trivial. We claim that if a map ϕ as above exists, then we may assume that $\phi|_{K_b}$ is generically finite, for generic $b \in B$. Indeed, because $H^0(X, K_X) \neq 0$, for generic $b \in B$, the image $\phi(K_b)$ has effective canonical bundle, in the sense that any desingularization Z_b of it has effective canonical bundle, as follows from adjunction formula and the fact that the $\phi(K_b)$ cover X . Now it is immediate to prove that any dominant rational map

$$K_b \dashrightarrow Z_b,$$

where K_b is an abelian variety and Z_b has effective canonical bundle, factors through the quotient map $K_b \rightarrow K'_b$, where K'_b is an abelian variety which is a quotient of K_b , and has the same dimension as Z_b . Replacing the family of abelian varieties $(K_b)_{b \in B}$, by the family $(K'_b)_{b \in B}$ gives the desired ϕ' .

In other words, Lang's conjecture asserts in particular that a Calabi-Yau variety should be rationally swept out by r -dimensional abelian varieties, for some $r \geq 1$. Hence, if Lang's conjecture is true, such an X should be swept out by elliptic curves.

On the other hand, the following is proved in [5], lecture 22 :

Proposition 1 *If a general Calabi-Yau hypersurface of dimension ≥ 2 is rationally swept-out by elliptic curves, then it contains a divisor which is uniruled.*

In dimension 3, this proposition combined with Theorem 2 shows that Lang's conjecture and Clemens' conjecture (see *loc. cit.*) on the finiteness of rational curves of fixed degree in a general quintic threefold, contradict.

In the case of hypersurfaces of general type, inequality (0.1) can be applied to give a non trivial estimate on the minimal genus of covering families of

curves, but the estimate is not sharp and could be obtained directly by geometry. What is interesting however is that looking more precisely at the proof of Theorem 1, we shall see that the result concerns in fact only the Hodge structure on $H^n(X)_{prim}$ and not the effective geometry of X . In fact we get as well :

Theorem 3 *Let X be a general hypersurface of degree $d \geq 2n - 2 + 3g$, $g \geq 2$ or $d \geq 2n + 2$, $g = 1$, in \mathbb{P}^{n+1} . Then there exists no non-zero morphism of Hodge structures*

$$H^n(X, \mathbb{Q})_{prim} \rightarrow H^n(Y, \mathbb{Q}),$$

where Y is rationally swept out by curves of genus g .

Combining this statement with the higher dimensional generalization of Mumford's theorem on 0-cycles on surfaces [15], this implies in particular that for $d \geq 2n + 2$, X general, there exists no correspondence $\Gamma \in CH^n(Y \times X)$ inducing a surjective map

$$\Gamma_* : CH_0(Y)_0 \rightarrow CH_0(X)_0,$$

where Y admits an elliptic fibration. Similarly, if $g \geq 2$ and $d \geq 2n - 2 + 3g$, there exists no such correspondence $\Gamma \in CH^n(Y \times X)$ where Y admits a fibration whose generic fiber is a genus g curve. One may wonder whether these statements are true for any such hypersurface or only for the general one.

The paper is organized as follows : in section 1, we recall briefly the proof of Nori's connectivity theorem for hypersurfaces in projective space, in order to prove a variant of it concerning families of hypersurfaces parameterized by subvarieties of the moduli space which are of small codimension. This will show us that for any family of hypersurfaces parameterized by a subvariety of the moduli space which is of small codimension, the Hodge level of the cohomology groups of the total space of the family is small.

The next section is devoted to the proof (by contradiction) of Theorem 1.

In section 3, we prove the applications of this result described above.

Acknowledgements. I would like to thank J. Harris who started me thinking to these problems and to Herb Clemens for very interesting discussions and comments.

This work has been essentially done at the University La Sapienza, and I would like to thanks the organizers of the trimester "Moduli spaces, Lie Theory, interactions with Physics", for the excellent working conditions and atmosphere I found there.

1 Nori's connectivity theorem for hypersurfaces

This section is devoted to an intermediate result (Theorem 4) toward the proof of Theorem 1. This result is a variant of Nori's connectivity theorem [16], or more precisely of its explicit version for hypersurfaces in projective space (cf [19], 8.1, [20]). In [1], [17], a sharper study of similar explicit bounds can be found.

Nori's theorem concerns the cohomology of locally complete families of sufficiently ample complete intersections in any smooth complex projective variety, parameterized by a smooth quasiprojective basis. The explicit version in case of hypersurfaces in projective space makes precise how ample the hypersurfaces must be.

Here, we consider a variant of this explicit version, where instead of locally complete families, we consider families of hypersurfaces whose moduli are of small codimension in the moduli space of all hypersurfaces.

We consider hypersurfaces of degree d in \mathbb{P}^{n+1} , and we assume that $d \geq n + 2$. Fix an integer r such that $1 \leq r \leq n$, and let γ be the round-up of $\frac{r-1}{2}$. Denote by $U \subset H^0(\mathcal{O}_{\mathbb{P}^{n+1}}(d))$ the open set parametrizing smooth hypersurfaces. Let $\rho : \mathcal{M} \rightarrow U$ be a morphism, where \mathcal{M} is smooth quasiprojective. We assume that $\text{Corank } \rho$ is constant equal to C . We also assume for simplicity that $\text{Im } \rho$ is stable under the action of $Gl(n+2)$. Let \mathcal{X}_U be the universal hypersurface parameterized by U and

$$\mathcal{X}_{\mathcal{M}} := \mathcal{X}_U \times_U \mathcal{M}.$$

Let

$$j : \mathcal{X}_{\mathcal{M}} \hookrightarrow \mathcal{M} \times \mathbb{P}^{n+1}$$

be the natural embedding. $\mathcal{X}_{\mathcal{M}}$ is a smooth quasi-projective variety, hence its cohomology groups carry mixed Hodge structures with associated Hodge filtration $F^i H^k(\mathcal{X}_{\mathcal{M}}, \mathbb{C})$.

Theorem 4 *i) Assume that*

$$(d+1)r \geq 2n + C + 2. \tag{1.2}$$

Then, the restriction map

$$j^* : F^n H^{2n-r}(\mathcal{M} \times \mathbb{P}^{n+1}, \mathbb{C}) \rightarrow F^n H^{2n-r}(\mathcal{X}_{\mathcal{M}}, \mathbb{C})$$

is surjective.

ii) If

$$(\gamma+1)d \geq 2n + 1 - r + C, \tag{1.3}$$

then for any $i \geq 1$, the restriction map

$$j^* : H^{2n-r-i}(\mathcal{M} \times \mathbb{P}^{n+1}, \mathbb{C}) \rightarrow H^{2n-r-i}(\mathcal{X}_{\mathcal{M}}, \mathbb{C})$$

is surjective.

The proof mimics the proof given in [19], 8.1 (which is nothing but the original proof of Nori, in the case of hypersurfaces of projective space). For the reader who knows already the arguments given there, let us just say that the proof in [19], 8.1 reduces first to showing that complexes built from the infinitesimal variations of Hodge structures (IVHS) on the primitive part of the cohomology of hypersurfaces are exact in a certain range. Via Carlson-Griffiths description of these IVHS, using the Griffiths description of the Hodge structure of an hypersurface via residues, these complexes are pointwise identified to Koszul complexes of the Jacobian ring of the considered hypersurface. The exactness of these Koszul complexes then uses a theorem due to M. Green on the Koszul cohomology of projective space. Our variant is then obtained by applying Green's refined theorem, which concerns Koszul cohomology of projective space, with respect to a non-complete base point free linear system of small codimension.

We try now to go a little more into the details, without reproducing the whole proof in [19], 8.1. We refer to *loc. cit.* for complete proofs of the steps sketched below.

Proof of Theorem 4.

i) One first reduces the assertion, as in [16], to proving that under the assumption (1.2), the restriction map

$$j^* : H^l(\Omega_{\mathcal{M} \times \mathbb{P}^{n+1}}^k) \rightarrow H^l(\Omega_{\mathcal{X}_{\mathcal{M}}}^k)$$

is bijective, for $l \leq n - r$, $k + l \leq 2n - r$. This step uses the mixed Hodge structure on relative cohomology and the Frölicher spectral sequence : the relative cohomology of the pair $(\mathcal{M} \times \mathbb{P}^{n+1}, \mathcal{X}_{\mathcal{M}})$ is equal to the hypercohomology of the complex

$$\Omega_{\mathcal{M} \times \mathbb{P}^{n+1}, \mathcal{X}_{\mathcal{M}}} := \text{Ker}(j^* : \Omega_{\mathcal{M} \times \mathbb{P}^{n+1}} \rightarrow \Omega_{\mathcal{X}_{\mathcal{M}}}).$$

The Frölicher spectral sequence considered here is the spectral sequence associated to the naïve filtration on $\Omega_{\mathcal{M} \times \mathbb{P}^{n+1}, \mathcal{X}_{\mathcal{M}}}$.

Denote respectively by π_X , $\pi_{\mathbb{P}}$ the natural maps

$$\mathcal{X}_{\mathcal{M}} \rightarrow \mathcal{M}, \mathcal{M} \times \mathbb{P}^{n+1} \rightarrow \mathcal{M}.$$

A Leray spectral sequence argument shows that it suffices to prove that under the assumption (1.2) one has :

$$\textit{The restriction map } j^* : R^l \pi_{\mathbb{P}*}(\Omega_{\mathcal{M} \times \mathbb{P}^{d+1}}^k) \rightarrow R^l \pi_{X*}(\Omega_{\mathcal{X}_{\mathcal{M}}}^k) \textit{ is bijective (1.4)}$$

for $l \leq n - r$, $k + l \leq 2n - r$.

Let

$$\mathcal{H}_{prim}^n, \mathcal{H}_{prim}^{p,q}, p + q = n,$$

be the Hodge bundles associated to the variation of Hodge structure on the primitive cohomology of the family $\pi_X : \mathcal{X}_{\mathcal{M}} \rightarrow \mathcal{M}$. The infinitesimal variation

of Hodge structure on the primitive cohomology of the fibers of π_X is described by maps

$$\bar{\nabla} : \mathcal{H}_{prim}^{p,q} \rightarrow \mathcal{H}_{prim}^{p-1,q+1} \otimes \Omega_{\mathcal{M}}, \quad (1.5)$$

and they can be extended in the obvious way to produce a complex :

$$\dots \mathcal{H}_{prim}^{p+1,q-1} \otimes \Omega_{\mathcal{M}}^{s-1} \xrightarrow{\bar{\nabla}} \mathcal{H}_{prim}^{p,q} \otimes \Omega_{\mathcal{M}}^s \xrightarrow{\bar{\nabla}} \mathcal{H}_{prim}^{p-1,q+1} \otimes \Omega_{\mathcal{M}}^{s+1} \dots \quad (1.6)$$

One shows, using the filtration of $\Omega_{\mathcal{X}_{\mathcal{M}}}^k$ by the subbundles $\pi_X^* \Omega_B^s \wedge \Omega_{\mathcal{X}_{\mathcal{M}}}^{k-s}$ and the associated spectral sequence (cf [19], 5.2.1), that (1.4) is equivalent to the following

The sequence (1.6) is exact at the middle for $q \leq n - r$, $p + s + q \leq 2n - r$.

Note that since $p + q = n$, the last inequality reduces to $s \leq n - r$.

It is convenient to dualize (1.6) using Serre duality, which gives :

$$\mathcal{H}_{prim}^{q+1,p-1} \otimes \bigwedge^{s+1} T_{\mathcal{M}} \xrightarrow{t\bar{\nabla}} \mathcal{H}_{prim}^{q,p} \otimes \bigwedge^s T_{\mathcal{M}} \xrightarrow{t\bar{\nabla}} \mathcal{H}_{prim}^{q-1,p+1} \otimes \bigwedge^{s-1} T_{\mathcal{M}}. \quad (1.7)$$

We finally use Griffiths, Griffiths-Carlson description of the IVHS of hypersurfaces ([13], [2]) to describe the complex (1.7) at the point $f \in \mathcal{M}$ as follows. We have the map $\rho_* : T_{\mathcal{M},f} \rightarrow T_{U,f} = S^d$, where S is the polynomial ring in $n + 2$ variables. Next the residue map provides isomorphisms

$$R_f^{-n-2+d(p+1)} \cong H_{prim}^{q,p}(X_f),$$

where $R_f := S/J_f$ is the Jacobian ideal of f , and R_f^k denotes its degree k component. The map $\bar{\nabla}$ in (1.5) identifies then, up to a coefficient, to the map given by ρ_* and multiplication :

$$R_f^{-n-2+d(p+1)} \rightarrow \text{Hom}(T_{\mathcal{M},f}, R_f^{-n-2+d(p+2)}).$$

It follows from this that the sequence (1.7) identifies to the following piece of the Koszul complex of the Jacobian ring R_f with respect to the action of $T_{\mathcal{M},f}$ on it by multiplication:

$$R_f^{-n-2+dp} \otimes \bigwedge^{s+1} T_{\mathcal{M},f} \xrightarrow{\delta} R_f^{-n-2+d(p+1)} \otimes \bigwedge^s T_{\mathcal{M},f} \xrightarrow{\delta} R_f^{-n-2+d(p+2)} \otimes \bigwedge^{s-1} T_{\mathcal{M},f} \quad (1.8)$$

Now, by assumption, if W is the image of ρ_* , $W \subset S^d$ is a base-point free linear system, because it contains the jacobian ideal J_f^d , and it satisfies $\text{codim } W = C$.

One verifies that it suffices to check exactness at the middle of the exact sequences (1.8) in the considered range, with $T_{\mathcal{M},f}$ replaced with W . This last fact is then a consequence of the following theorem due to M. Green :

Theorem 5 [12] *Let $W \subset S^d$ be a base-point free linear system. Then the following sequence, where the differentials are the Koszul differentials*

$$S^{-n-2+dp} \otimes \bigwedge^{s+1} W \xrightarrow{\delta} S^{-n-2+d(p+1)} \otimes \bigwedge^s W \xrightarrow{\delta} S^{-n-2+d(p+2)} \otimes \bigwedge^{s-1} W \quad (1.9)$$

is exact for $-n - 2 + dp \geq s + \text{codim } W$.

Using the fact that the Jacobian ideal is generated by a regular sequence in degree $d - 1$, which provides a very simple resolution of R_f by graded free S -modules, one then shows that the same is true when S^i is replaced with R_f^i in (1.9), at least if $-n - 2 + d(p + 1) \geq d - 1$.

We now conclude the proof of i). We have just proved that (1.6) is exact at the middle if

$$-n - 2 + dp \geq s + C, \quad -n - 2 + d(p + 1) \geq d - 1.$$

Since we assumed $d \geq n + 2$, the second inequality is satisfied when $p \geq 1$. Next, if $q \leq n - r$, $s \leq n - r$, we have

$$p \geq r \geq 1, \quad s \leq n - r.$$

Hence, the exactness of (1.6) in the range $q \leq n - r$, $s \leq n - r$ will follow from the inequality

$$-n - 2 + dr \geq n - r + C,$$

that is (1.2).

ii) The proof is entirely similar, and we just sketch it in order to see where the numerical assumption is used. Following [16], we first observe that, in order to get the surjectivity of the restriction map :

$$j^* : H^{2n-r-i}(\mathcal{M} \times \mathbb{P}^{n+1}, \mathbb{C}) \rightarrow H^{2n-r-i}(\mathcal{X}_{\mathcal{M}}, \mathbb{C}),$$

it suffices to show the surjectivity of the restriction map :

$$j^* : F^{n-r+\gamma_i} H^{2n-r-i}(\mathcal{M} \times \mathbb{P}^{n+1}, \mathbb{C}) \rightarrow F^{n-r+\gamma_i} H^{2n-r-i}(\mathcal{X}_{\mathcal{M}}, \mathbb{C}), \quad (1.10)$$

where γ_i is the round-up of $\frac{r-i}{2}$.

Indeed, let K be the cokernel

$$H^{2n-r-i}(\mathcal{X}_{\mathcal{M}}, \mathbb{Q}) / j^* H^{2n-r-i}(\mathcal{M} \times \mathbb{P}^{n+1}, \mathbb{Q}).$$

It has a quotient mixed Hodge structure whose weights are $\geq 2n - r - i$, because $\mathcal{X}_{\mathcal{M}}$ is smooth (cf [8]). The surjectivity of

$$j^* : F^{n-r+\gamma_i} H^{2n-r-i}(\mathcal{M} \times \mathbb{P}^{n+1}, \mathbb{C}) \rightarrow F^{n-r+\gamma_i} H^{2n-r-i}(\mathcal{X}_{\mathcal{M}}, \mathbb{C}),$$

then says that $F^{n-r+\gamma_i}K_{\mathbb{C}} = 0$. But the round-up of $\frac{2n-r-i}{2}$ is $n-r+\gamma_i$. Applying Hodge symmetry to the weight graded pieces of K (which carry pure Hodge structures of weight $\geq 2n-r-i$), we see that

$$F^{n-r+\gamma_i}K_{\mathbb{C}} = 0 \Rightarrow K = 0.$$

As in the previous proof, we reduce now the proof of the surjectivity of (1.10) to showing :

$$\text{The restriction map } j^* : R^l \pi_{\mathbb{P}^*}(\Omega_{\mathcal{M} \times \mathbb{P}^{n+1}}^k) \rightarrow R^l \pi_{X*}(\Omega_{\mathcal{X}_{\mathcal{M}}}^k) \text{ is bijective} \quad (1.11)$$

for $l \leq n-i-\gamma_i$, $k+l \leq 2n-r-i$.

Expressing as in the previous proof the cohomology groups above with the help of the IVHS on the primitive cohomology of the fibers of π_X , this is reduced to proving :

The sequence (1.6) is exact at the middle for $q \leq n-i-\gamma_i$, $p+q = n$, $p+s+q \leq 2n-r-i$.

Using the Carlson-Griffiths theory, we are now reduced to prove :

the following sequence :

$$R_f^{-n-2+dp} \otimes \bigwedge^{s+1} T_{\mathcal{M},f} \xrightarrow{\delta} R_f^{-n-2+d(p+1)} \otimes \bigwedge^s T_{\mathcal{M},f} \xrightarrow{\delta} R_f^{-n-2+d(p+2)} \otimes \bigwedge^{s-1} T_{\mathcal{M},f} \quad (1.12)$$

is exact for $p \geq \gamma_i + i$, $s \leq n-r-i$.

As in the previous proof, we now apply Theorem 5 and conclude that the last statement is true if

$$-n-2+d(\gamma_i+i) \geq C+n-r-i. \quad (1.13)$$

Now it is clear that the γ_i+i are increasing with i , while the $C+n-r-i$ are decreasing with i . Hence it suffices to have (1.13) satisfied for $i=1$, which is exactly inequality (1.3). \blacksquare

Denoting by $H^{2n-r}(\mathcal{X}_{\mathcal{M}})_{prim}$ the quotient

$$H^{2n-r}(\mathcal{X}_{\mathcal{M}})_{prim} / j^*(H^{2n-r}(\mathcal{M} \times \mathbb{P}^{n+1})),$$

we shall only be interested with the pure part

$$W_{2n-r}H^{2n-r}(\mathcal{X}_{\mathcal{M}})_{prim},$$

which is the part of the cohomology which comes from any smooth projective compactification of $\mathcal{X}_{\mathcal{M}}$ [8]. It carries a pure Hodge structure of weight $2n-r$.

Corollary 1 *Under the assumptions of Theorem 4, the Hodge structure on $W_{2n-r}H^{2n-r}(\mathcal{X}_{\mathcal{M}})_{prim}$ is of Hodge level $\leq r-2$.*

Proof. Recall that the Hodge level of a pure Hodge structure H , $H_{\mathbb{C}} = \oplus H^{p,q}$ is

$$\text{Max} \{p - q, H^{p,q} \neq 0\}.$$

Since we know that $F^n W_{2n-r} H^{2n-r}(\mathcal{X}_{\mathcal{M}})_{\text{prim}} = 0$, we have

$$H^{p,q}(W_{2n-r} H^{2n-r}(\mathcal{X}_{\mathcal{M}})_{\text{prim}}) = 0, \text{ for } p \geq n.$$

Since $H^{p,q} = 0$ for $p + q \neq 2n - r$, it follows that the Hodge level is $\leq n - 1 - (2n - r - (n - 1)) = r - 2$. \blacksquare

2 Proof of Theorem 1

We prove Theorem 1 by contradiction. So assume the assumptions (0.1) are satisfied, but that the general X of degree d in \mathbb{P}^{n+1} is rationally swept-out by varieties from the family \mathcal{Y} .

Using Chow varieties, or relative Hilbert schemes, we see that there exist countably many quasi-projective varieties \mathcal{B}_k parameterizing triples (t, f, ϕ_s) , where $t \in \mathcal{S}$, $f \in U$, and ϕ_s is a rational map $\phi_s : Y_s \dashrightarrow X_f$ which is generically finite onto its image. Indeed, such a ϕ_s can be identified to its graph, which is a r -dimensional subvariety of the product $Y_s \times X_f$, which has to be of degree 1 over the first factor.

For fixed generic f , our assumption implies that the images of such ϕ_s fill-in X_f .

For each k , let

$$\pi_k : \mathcal{K}_k \rightarrow \mathcal{B}_k$$

be the family which is pulled-back via the first projection $\mathcal{B}_k \rightarrow \mathcal{S}$ from the family $\mathcal{Y} \rightarrow \mathcal{S}$. By definition of \mathcal{B}_k as a Hilbert scheme, there is the universal rational map

$$\Phi_k : \mathcal{K}_k \dashrightarrow \mathcal{X}_U,$$

where, as in the previous section, \mathcal{X}_U is the universal hypersurface parameterized by U . Note that Φ_k makes the following diagram commutative :

$$\begin{array}{ccc} \Phi_k : & \mathcal{K}_k & \dashrightarrow & \mathcal{X}_U \\ & \downarrow \pi_k & & \downarrow \\ pr_2 : & \mathcal{B}_k & \rightarrow & U \end{array} \quad . \quad (2.14)$$

Our assumption implies that the union of the images of Φ_k contains a subset of \mathcal{X}_U which is the complementary set of a countable union of proper closed algebraic subsets. By Baire, it follows that for some k_0 , Φ_{k_0} is dominating. In the sequel, we shall use the notations $\pi, \mathcal{B}, \mathcal{K}, \Phi$, for $\pi_{k_0}, \mathcal{B}_{k_0}, \mathcal{K}_{k_0}, \Phi_{k_0}$ respectively.

We shall denote by B_f the (generic) fiber of the second projection $\mathcal{B} \rightarrow U$ and $\pi_f : \mathcal{K}_f \rightarrow B_f$ the induced family. By taking desingularizations, we may assume that \mathcal{B} is smooth, and since by assumption the map π is smooth, \mathcal{K}

is smooth too. Since f is generic, B_f and K_f are then also smooth. Finally, we may, up to replacing \mathcal{B} by a closed subvariety, assume that the restriction $\Phi_f : K_f \dashrightarrow X_f$ of Φ to K_f is generically finite and dominating. (This is because the restriction of Φ to the generic fibers of π is generically finite.) In particular, $\dim B_f = n - r$.

We fix now a generic point $t \in \text{Im } pr_1 \subset \mathcal{S}$. We let

$$\mathcal{B}_t := pr_1^{-1}(t), \mathcal{K}_t := \pi^{-1}(\mathcal{B}_t).$$

We observe that \mathcal{K}_t is naturally isomorphic to $Y_t \times \mathcal{B}_t$. Furthermore, since

$$\dim \text{Im } pr_1 \leq \dim \mathcal{S} \leq C,$$

we have $\text{codim}(\mathcal{B}_t \subset \mathcal{B}) \leq C$.

The second projection $pr_2 : \mathcal{B} \rightarrow U$ is dominating, so can be assumed to be submersive, after shrinking \mathcal{B} . It follows that the corank of $pr_2|_{\mathcal{B}_t}$ is $\leq C$, and after shrinking \mathcal{B}_t , we may assume it is constant and $\leq C$ on \mathcal{B}_t .

Let $\mathcal{X}_{\mathcal{B}_t} := \mathcal{X}_U \times_U \mathcal{B}_t$. The map $\Phi : \mathcal{K} \dashrightarrow \mathcal{X}_U$ lifts to a map

$$\tilde{\Phi} : \mathcal{K} \dashrightarrow \mathcal{X}_{\mathcal{B}_t},$$

because of the commutativity of the diagram (2.14). $\tilde{\Phi}$ takes the value ϕ_s over the point (t, f, ϕ_s) of \mathcal{B} .

This map restricts in turn to a map

$$\tilde{\Phi}_t : \mathcal{K}_t \dashrightarrow \mathcal{X}_{\mathcal{B}_t},$$

which is over \mathcal{B}_t .

The graph of $\tilde{\Phi}_t$ provides then a codimension n subvariety in the fibred product

$$\mathcal{K}_t \times_{\mathcal{B}_t} \mathcal{X}_{\mathcal{B}_t} \cong Y_t \times \mathcal{X}_{\mathcal{B}_t}.$$

The desired contradiction will come from the following :

Proposition 2 *Let $\gamma_t \in H^{2n}(Y_t \times \mathcal{X}_{\mathcal{B}_t}, \mathbb{Q})$ be the rational cohomology class of the graph of $\tilde{\Phi}_t$. Then γ_t induces a morphism of mixed Hodge structures*

$$\gamma_{t*} : H^r(Y_t, \mathbb{Q}) \rightarrow H^{2n-r}(\mathcal{X}_{\mathcal{B}_t}, \mathbb{Q})_{\text{prim}}, \quad (2.15)$$

whose image has Hodge level r .

Remark 3 *The mixed Hodge structure on $H^r(Y_t, \mathbb{Q})$ is pure as is the mixed Hodge structure on $\text{Im } \gamma_{t*}$. Hence it makes sense to speak of the Hodge level of the image.*

Remark 4 *The fact that $\text{Im } \gamma_{t*}$ has Hodge level r is equivalent to the fact that γ_{t*} does not vanish identically on $H^{r,0}(Y_t)$, or more generally on the subspace $H^r(Y_t, \mathbb{Q})_{tr}$ introduced later on in the proof.*

Admitting the proposition, the proof concludes as follows : we know that the map $\mathcal{B}_t \rightarrow U$ is of constant corank $\leq C$, and by assumption, the inequality (1.2) needed in Corollary 1 is satisfied, so Corollary 1 holds. On the other hand Proposition 2 clearly contradicts Corollary 1. \blacksquare

Proof of Proposition 2. The graph of $\tilde{\Phi}$ lies in $\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}$ and is of codimension n there. Let $\gamma \in H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q})$ be its rational cohomology class. Of course

$$\gamma_t = \gamma|_{\mathcal{X}_{\mathcal{B}_t}}.$$

Let $\tilde{\pi} : \mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}} \rightarrow \mathcal{X}_{\mathcal{B}}$ be the second projection. Let \mathcal{L} be a relatively ample line bundle on $\mathcal{Y} \rightarrow \mathcal{S}$, and let $\tilde{\mathcal{L}}$ be its pull-back to $\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}$ via the natural composite map

$$\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}} \rightarrow \mathcal{K} \rightarrow \mathcal{Y}.$$

Then $\tilde{\mathcal{L}}$ is relatively ample w.r.t. $\tilde{\pi}$.

The hard Lefschetz theorem provides isomorphisms

$$c_1(\mathcal{L})^i \cup : R^{r-i} \tilde{\pi}_* \mathbb{Q} \cong R^{r+i} \tilde{\pi}_* \mathbb{Q}, \quad i \geq 0,$$

and correspondingly isomorphisms :

$$c_1(\mathcal{L})^i \cup : H^{2n-r-i}(\mathcal{X}_{\mathcal{B}}, R^{r-i} \tilde{\pi}_* \mathbb{Q}) \cong H^{2n-r-i}(\mathcal{X}_{\mathcal{B}}, R^{r+i} \tilde{\pi}_* \mathbb{Q}), \quad i \geq 0.$$

These isomorphisms are isomorphisms of mixed Hodge structures. Furthermore, by Deligne's theorem [7], the Leray spectral sequence of $\tilde{\pi}$ degenerates at E_2 , so that the above groups are graded pieces of

$$H^{2n-2i}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q}), \quad H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q})$$

respectively. It follows that we can write

$$\gamma = \gamma' + \sum_{r \geq i > 0} c_1(\tilde{\mathcal{L}})^i \cup \gamma_i, \quad (2.16)$$

where $\gamma' \in H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q})$ lies in the r -th level of the Leray filtration relative to $\tilde{\pi}$, and $\gamma_i \in H^{2n-2i}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q})$. By strictness of the morphisms of mixed Hodge structures with respect to both filtrations, we may even assume that γ' and γ_i are Hodge classes, that is lie respectively in

$$W_{2n} H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q}) \cap F^n H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q}),$$

$$W_{2n-2i} H^{2n-2i}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q}) \cap F^{n-i} H^{2n-2i}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q}).$$

Restrict now everything to

$$\mathcal{K}_t \times_{\mathcal{B}_t} \mathcal{X}_{\mathcal{B}_t} \cong Y_t \times \mathcal{X}_{\mathcal{B}_t}. \quad (2.17)$$

Since $\tilde{\mathcal{L}}$ is the pull-back of a line bundle on \mathcal{Y} , its restriction $\tilde{\mathcal{L}}_t$ to $\mathcal{K}_t \times_{\mathcal{B}_t} \mathcal{X}_{\mathcal{B}_t}$ is of the form $pr_1^* L$, for some line bundle L on Y_t , where pr_1 is defined using the isomorphism (2.17). It follows that each

$$\gamma_{i,t} := c_1(\tilde{\mathcal{L}})^i \cup \gamma_i|_{\mathcal{K}_t \times_{\mathcal{B}_t} \mathcal{X}_{\mathcal{B}_t}}$$

induces a morphism of mixed Hodge structures

$$H^r(Y_t, \mathbb{Q}) \rightarrow H^{2n-r}(\mathcal{X}_{B_t}, \mathbb{Q}),$$

which factors through the map $c_1(L)^i \cup : H^r(Y_t, \mathbb{Q}) \rightarrow H^{r+2i}(Y_t, \mathbb{Q})$. So, since $\dim Y_t = r$, each such morphism has an image which is of Hodge level $< r$ for $i > 0$.

We observe now the following : γ' lies in the r -th level of the Leray filtration of $H^{2n}(\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}, \mathbb{Q})$ relative to $\tilde{\pi}$, hence, using the degeneracy at E_2 of this Leray spectral sequence, it projects to an element γ'' in

$$H^{2n-r}(\mathcal{X}_{\mathcal{B}}, R^r \tilde{\pi}_* \mathbb{Q}),$$

which is the $E_2^{r, n-r} = E_{\infty}^{r, n-r}$ term of the Leray spectral sequence of $\tilde{\pi}$.

Denote by ψ the composite map

$$\mathcal{X}_{\mathcal{B}} \rightarrow \mathcal{B} \rightarrow \mathcal{S}.$$

Observe that the family

$$\tilde{\pi} : \mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}} \rightarrow \mathcal{X}_{\mathcal{B}}$$

is pulled-back via ψ from the family $\mathcal{Y} \rightarrow \mathcal{S}$. It follows that the local system is also pulled-back :

$$R^r \tilde{\pi}_* \mathbb{Q} = \psi^{-1} H^r,$$

for some local system H^r of \mathbb{Q} -vector spaces on \mathcal{S} . Hence the E_2 terms of the Leray spectral sequence relative to ψ and to the sheaf $R^r \tilde{\pi}_* \mathbb{Q}$ on $\mathcal{X}_{\mathcal{B}}$ are

$$H^i(\mathcal{S}, R^{2n-r-i} \psi_* \mathbb{Q} \otimes_{\mathbb{Q}} H^r).$$

From now on we assume that the map $pr_1 : \mathcal{B} \rightarrow \mathcal{S}$ is surjective, and topologically locally trivial, which we can do, replacing \mathcal{S} by a Zariski open set in $Im pr_1$. The local system H^r contains a sub-local system which via Poincaré duality is also canonically a direct summand, whose fiber at the general point of \mathcal{S} is the maximal sub-Hodge structure of $H^r(Y_t, \mathbb{Q})$ which is not of maximal Hodge level. Let us denote this sub-local system by $H_{<r}^r$, and the quotient by H_{tr}^r . H_{tr}^r can also be defined as the minimal sub-variation of Hodge structure of H^r whose corresponding $(r, 0)$ -part contains the whole bundle $\mathcal{H}^{r,0}$ with fiber $H^{r,0}(Y_t)$ at $t \in \mathcal{S}$. So we have an orthogonal decomposition :

$$H^r = H_{<r}^r \oplus H_{tr}^r.$$

Let $R^r \tilde{\pi}_* \mathbb{Q}_{tr} := \psi^{-1} H_{tr}^r$. With these notations, we have the component γ''_{tr} of γ'' , which lies in

$$H^{2n-r}(\mathcal{X}_{\mathcal{B}}, R^r \tilde{\pi}_* \mathbb{Q}_{tr}),$$

and as above, the spectral sequence abutting to this last group has E_2 -term

$$H^i(\mathcal{S}, R^{2n-r-i} \psi_* \mathbb{Q} \otimes_{\mathbb{Q}} H_{tr}^r). \quad (2.18)$$

We apply now Theorem 4, ii). The fibers of the maps ψ are the $\mathcal{X}_{\mathcal{B}_t}$ hence they satisfy the vanishing

$$H^{2n-r-i}(\mathcal{X}_{\mathcal{B}_t}, \mathbb{Q})_{prim} = 0.$$

Formula (2.18) allows then to conclude that for $i > 0$, we have

$$H^i(\mathcal{S}, R^{2n-r-i}\psi_*\mathbb{Q}_{prim} \otimes_{\mathbb{Q}} H_{tr}^r) = 0.$$

Hence we conclude that up to classes coming from

$$H^{2n-r}(\mathbb{P}^{n+1} \times \mathcal{B}, R^r(Id \times \pi)_*\mathbb{Q}_{tr}),$$

the class γ''_{tr} is determined by its 0-th Leray component, namely

$$\gamma''' \in H^0(\mathcal{S}, R^{2n-r}\psi_*\mathbb{Q}_{prim} \otimes_{\mathbb{Q}} H_{tr}^r).$$

Note that the value of γ''' at the general point $t \in \mathcal{S}$ is, (using Poincaré duality on Y_t ,) nothing but the restriction of the map γ'_{t*} or γ_{t*} of (2.15) to the transcendental part $H^r(Y_t, \mathbb{Q})_{tr}$.

Let us now conclude the proof by contradiction : assume the γ'_{t*} has an image which is of Hodge level $< r$ for general t . Then equivalently, its restriction to $H^r(Y_t, \mathbb{Q})_{tr}$ vanishes. This means that $\gamma''' = 0$, and by the reasoning above, this implies that $\gamma''_{tr} = 0$ modulo classes coming from $H^{2n-r}(\mathbb{P}^{n+1} \times \mathcal{B}, R^r(Id \times \pi)_*\mathbb{Q}_{tr})$.

We now observe that $\mathcal{X}_{\mathcal{B}}$ contains $B_f \times X_f$, \mathcal{K} contains K_f , and that correspondingly $\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}$ contains $K_f \times X_f$. The restriction of $\tilde{\Phi}$ to K_f is the map $(\pi_f, \Phi_f) : K_f \dashrightarrow B_f \times X_f$ and so the graph of $\tilde{\Phi}$ in $\mathcal{K} \times_{\mathcal{B}} \mathcal{X}_{\mathcal{B}}$ restricts to the graph of Φ_f in $K_f \times X_f$. The class γ restricts to the class γ_f of this graph.

Denote by $\gamma'_f, \gamma''_f, \gamma''_{f,tr}$ the restrictions of the classes $\gamma', \gamma'', \gamma''_{tr}$ respectively to $K_f \times X_f$. The vanishing $\gamma''_{tr} = 0$ implies the vanishing $\gamma''_{f,tr} = 0$. Retracing through the definitions, this implies that the map

$$\gamma_f^* : H^n(X_f, \mathbb{Q})_{prim} \rightarrow H^n(K_f, \mathbb{Q}),$$

where the subscript “prim” here means “the orthogonal part to $H^*(\mathbb{P}^{n+1}, \mathbb{Q})|_{X_f}$ ”, decomposes as

$$\gamma_f^* = \sum_{i>0} l^i \cup \gamma_{i,f}^* + \gamma_f'^*,$$

where l is the first Chern class of the line bundle on K_f induced by \mathcal{L} via the map $K_f \rightarrow \mathcal{Y}$. Here the $\gamma_{i,f}, \gamma'$ are the restriction to $K_f \times X_f$ of the classes γ_i, γ' of (2.16). The map $\gamma_f'^*$ takes value in the r -th level of $H^n(K_f, \mathbb{Q})$ with respect to the Leray filtration relative to the map $\pi_f : K_f \rightarrow B_f$, and our conclusion is that its projection in

$$Hom(H^n(X_f, \mathbb{Q})_{prim}, H^{n-r}(B_f, R^r\pi_{f*}\mathbb{Q}))$$

lies in $Hom(H^n(X_f, \mathbb{Q})_{prim}, H^{n-r}(B_f, R^r\pi_{f*}\mathbb{Q})_{<r})$.

The contradiction now comes from the fact that since Φ_f is a dominating rational map, and the Hodge structure on $H^n(X_f, \mathbb{Q})_{prim}$ has Hodge level n , (because $d \geq n + 2$), the associated map γ_f^* has an image which is of Hodge level n .

On the other hand, the maps of the form $l^i \cup \gamma_{i,f}^*$ have images of Hodge level $< n$, as do the maps $\gamma_{i,f}^*$. Next, the only Leray piece of $H^n(K_f, \mathbb{Q})$ whose weight n part is of Hodge level n is $H^{n-r}(R^r \pi_{f*} \mathbb{Q})$, because $\dim B_f = n - r$, and inside it $H^{n-r}(R^r \pi_{f*} \mathbb{Q}_{<r})$ has Hodge level $< n$. ■

3 Rational maps from abelian varieties to Calabi-Yau hypersurfaces and other applications

Proof of Theorem 2. We want to show that for $r \geq 2$, the general Calabi-Yau hypersurfaces, that is hypersurfaces of degree $d = n + 2$ in \mathbb{P}^{n+1} are not rationally swept-out by r -dimensional abelian varieties. We apply Theorem 1 : the family $\mathcal{Y} \rightarrow \mathcal{S}$ we consider is any locally universal family of abelian varieties. The moduli space of r -dimensional abelian varieties with given polarization type is of dimension $\frac{r(r+1)}{2}$. Hence the inequalities needed in Theorem 1 become in this case :

$$(n + 3)r \geq 2n + \frac{r(r + 1)}{2} + 2, (\gamma + 1)(n + 2) \geq 2n - r + 1 + \frac{r(r + 1)}{2}.$$

It is not hard to check that this is satisfied for $2 \leq r \leq n$. ■

When $r = 1$, inequality (0.1) is never satisfied so that our argument definitely does not apply to the study of elliptic curves in Calabi-Yau hypersurfaces. In fact we could adapt our proof of Theorem 2 to work as well for Calabi-Yau hypersurfaces in a product of projective spaces. On the other hand, certain generic Calabi-Yau hypersurfaces in a product of projective spaces are swept out by elliptic curves, eg the hypersurface of bidegree (3, 3) in $\mathbb{P}^2 \times \mathbb{P}^2$. This shows that for $r = 1$, a different argument has to be found. Note however that Theorem 2 has the following corollary, which says that Lang's and Clemens' conjecture contradict :

Corollary 2 *If Lang's conjecture is true, any Calabi-Yau hypersurface X of dimension ≥ 2 has a divisor which is uniruled.*

Proof. Indeed, if Lang's conjecture is true, then by Theorem 2, the general X is rationally swept out by elliptic curves. One then applies the following result, which is proved in the three dimensional case in [5], lecture 22 (the proof however applies to any dimension ≥ 2) :

Proposition 3 *If a general Calabi-Yau hypersurface is rationally swept-out by elliptic curves, then it contains a divisor which is rationally swept-out by rational curves.*

■

For completeness, we sketch here the argument for the proof of Proposition 3. The first step is to prove :

Lemma 1 *If X is a general Calabi-Yau hypersurface of dimension ≥ 2 , X is not rationally swept out by elliptic curves of fixed modulus.*

Proof. Indeed, fixing otherwise the modulus of the elliptic curve, we would get, for at least one elliptic curve E , an hypersurface \mathcal{M}_E in the moduli space \mathcal{M} of X consisting of X_f 's which are rationally dominated by some $E \times B$. For such an X_f , there must be an inclusion of rational Hodge structures induced by the dominant rational map $\phi : E \times B \dashrightarrow X_f$:

$$\phi^* : H^n(X_f)_{prim} \hookrightarrow H^1(E) \otimes H^{n-1}(B), \quad (3.19)$$

because the Hodge structure on $H^n(X)_{prim}$ is simple for general X .

If we now let f vary in \mathcal{M}_E , only B deforms with f , not E , and it follows that the infinitesimal variation of Hodge structure on $H^1(E) \otimes H^{n-1}(B)$

$$\bar{\nabla} : H^{p,q}(H^1(E) \otimes H^{n-1}(B)) \rightarrow H^{p-1,q+1}(H^1(E) \otimes H^{n-1}(B)) \otimes \Omega_{\mathcal{M}_E}$$

has the following form at the point $f \in \mathcal{M}_E$:

$$\bar{\nabla}(\alpha \otimes \beta) = \alpha \otimes \bar{\nabla}_B(\beta)$$

for $\alpha \in H^{r,s}(E)$, $\beta \in H^{p-r,q-s}(B)$, where $\bar{\nabla}_B$ is the infinitesimal variation of Hodge structure on $H^{n-1}(B)$. Hence the Yukawa couplings of the IVHS on $H^1(E) \otimes H^{n-1}(B)$, that is the iterations of $\bar{\nabla}$, have the following property:

$\forall \eta \in H^{n,0}(H^1(E) \otimes H^{n-1}(B))$, the map

$$\bar{\nabla}^n(\eta) : S^n T_{\mathcal{M}_E, f} \rightarrow H^{0,n}(H^1(E) \otimes H^{n-1}(B))$$

vanishes.

If there is along \mathcal{M}_E an injective morphism of Hodge structures (3.19), it follows that the same property is true for the IVHS of the family of X_f 's parameterized by \mathcal{M}_E , namely the Yukawa couplings of X_f vanish on the hyperplane $K := T_{\mathcal{M}_E, f} \subset S^{n+2}$. The Carlson-Griffiths theory [13], [2] shows easily that this is not the case. Indeed, these Yukawa couplings identify to the multiplication map

$$S^n(S^{n+2}) \rightarrow R_f^{n(n+2)}.$$

Assume they vanish on K . Since K is a hyperplane in S^{n+2} , the subspace

$$K' := [K : S^1] \subset S^{n+1}$$

has codimension $\leq n+2$. It is without base-point, since $T_{\mathcal{M}_E, f}$ contains J_f . It follows then from [11], that $S^{n+3}K' = S^{2n+4}$. But K^2 contains $S^1K' \cdot K = K' \cdot S^1K = K' \cdot S^{n+3} = S^{2n+4}$. Hence $K^2 = S^{2n+4}$ and similarly $K^n = S^{n(n+2)}$ contradicting the fact that $K^n \subset J_f^{n(n+2)}$. \blacksquare

The second step is then following

Lemma 2 *Assume a variety is rationally swept-out by elliptic curves, but not by elliptic curves with constant modulus. Then it admits a divisor which is rationally swept out by rational curves.*

Proof. By assumption there is a diagram

$$\begin{array}{ccc} \widetilde{\mathcal{K}} & \xrightarrow{\phi} & X \\ \pi \downarrow & & \\ \widetilde{B} & & \end{array},$$

where we may assume that \widetilde{K} and \widetilde{B} are smooth, projective, where $\widetilde{\mathcal{K}}$ is a smooth projective model of the family $\mathcal{K} \rightarrow B$ on which ϕ is defined, and that the map $j : \widetilde{B} \rightarrow \mathbb{P}^1$ is defined and non constant. Now, since ϕ is generically finite, for generic $t \in \mathbb{P}^1$, the divisor $\widetilde{\mathcal{K}}_t := (j \circ \pi)^{-1}(t)$ must be sent by ϕ onto a divisor of X , and it follows that for any t the image by ϕ of the divisor $\widetilde{\mathcal{K}}_t := (j \circ \pi)^{-1}(t)$ must contain a divisor of X . Taking $t = \infty$, and noting that any component of $\widetilde{\mathcal{K}}_\infty$ has a normalization which is uniruled, gives the result. ■

Proof of Theorem 3. We simply note for this that in the proof of Theorem 1, we used the dominating rational map

$$\phi : K \dashrightarrow X$$

only to deduce that there is a corresponding inclusion

$$\phi_f^* : H^n(X_f)_{prim} \rightarrow H^n(K_f)$$

of Hodge structures. The Chow argument we used would work equally with graphs of rational maps replaced with cycles in the product

$$Y_t \times X.$$

Hence we conclude that everywhere in the paper, we could replace “dominating rational maps” by “cycles in $CH^d(K \times X)$ inducing a non-zero morphism of Hodge structures”

$$H^n(X)_{prim} \rightarrow H^n(K).$$

(Note that such a non-zero morphism should be in fact injective because the Hodge structure on $H^n(X)_{prim}$ is simple for generic X .)

Using the fact that the moduli space of curves of genus $g \geq 1$ has dimension $3g - 3$, $g \geq 2$ or 1 , $g = 1$, we see that theorem 3 is then a consequence of theorem 1, where we replace dominating rational maps with correspondences inducing a non zero morphism of Hodge structure. ■

References

- [1] M. Asakura, S. Saito. Noether-Lefschetz locus for Beilinson-Hodge cycles on open complete intersections, preprint 2003.
- [2] J. Carlson, Ph. Griffiths. Infinitesimal variations of Hodge structures and the global Torelli problem, in *Journées de géométrie algébrique*, (A. Beauville Ed.), Sijthoff-Nordhoff, (1980), 51-76.
- [3] L. Chiantini, A.-F. Lopez, Z. Ran. Subvarieties of generic hypersurfaces in any variety, *Math. Proc. Cambr. Phil. Soc.* 2002.
- [4] H. Clemens. Curves in generic hypersurfaces, *Ann. Sci. École Norm. Sup.* 19 (1986), 629-636.
- [5] H. Clemens, J. Kollár, S. Mori. *Higher dimensional complex geometry*, Astérisque 166, SMF (1988).
- [6] H. Clemens, Z. Ran. Twisted genus bounds for subvarieties of generic hypersurfaces, to appear *Amer. J. Math.*
- [7] P. Deligne. Théorèmes de Lefschetz et critères de dégénérescence de suites spectrales, *Publ. math. IHES* 35 (1968) 107-126.
- [8] P. Deligne. Théorie de Hodge II, *Publ. Math. IHES* 40 (1971), 5-57.
- [9] P. Deligne. La conjecture de Weil pour les surfaces $K3$, *Inventiones Math.* 15, 206-226 (1972).
- [10] L. Ein. Subvarieties of generic complete intersections, *Invent. Math.* 94 (1988), 163-169.
- [11] M. Green. Restrictions of linear series to hyperplanes, and some results of Macaulay and Gotzmann, in *Algebraic curves and projective geometry*, (E. Ballico, C. Ciliberto Eds), *Lecture Notes in Mathematics* 1389, Springer-Verlag 1989, 76-86.
- [12] M. Green. A new proof of the explicit Noether-Lefschetz theorem, *J. Diff. geom.*
- [13] Ph. Griffiths. Periods of certain rational integrals, I, II, *Ann. of Math.* 90 (1969) 460-541.
- [14] S. Lang. Hyperbolic and diophantine Analysis, *Bulletin of the American Math. Soc.* Vol. 14, 2, (1986), 159-205.
- [15] D. Mumford. Rational equivalence of zero-cycles on surfaces, *J. Math. Kyoto Univ.* 9 (1968), 195-204.

- [16] M. Nori. Algebraic cycles and Hodge theoretic connectivity, *Invent. Math.* 111 (1993) 349-373.
- [17] A. Otwinowska. Asymptotic bounds for Nori's connectivity theorem, preprint 2002.
- [18] C. Voisin. On a conjecture of Clemens on rational curves on hypersurfaces, *Journal of Differential Geometry* 44 (1996) 200-214 (+ correction *J. Differential Geometry* 49 (1998), 601-611.
- [19] C. Voisin. *Hodge Theory and Complex Algebraic Geometry II*, Cambridge University Press (2003).
- [20] C. Voisin. Nori's connectivity theorem and higher Chow groups, *Journal of the Inst. Math. Jussieu* (2002) 1 (2), 307-329.