

Torsion cohomology classes and algebraic cycles on complex projective manifolds

C. Soulé, C. Voisin

Let X be a smooth complex projective manifold and $H^n(X, \mathbb{Z})$ its singular cohomology group of degree n , with integral coefficients. Given a torsion class $\alpha \in H^{2k}(X, \mathbb{Z})$, can we say that this class α is algebraic?

This is true when $k = 1$, and, apparently, Hodge thought that this would always be the case [10]. However, Atiyah and Hirzebruch found counterexamples to Hodge's assertion [2]. This is why the Hodge conjecture is now formulated for rational cohomology classes only. Recently, Totaro gave a new interpretation of the Atiyah-Hirzebruch counterexamples, in terms of the complex cobordism ring of X [19].

When looking at these examples, we noticed that the order of the non-algebraic torsion class α can be divisible only by primes at most equal to the complex dimension of X (Th. 1). On the other hand, a construction of Kollár [12] provides examples of manifolds X as above with a class $\alpha \in H^4(X, \mathbb{Z})$ which is not algebraic, while a non-zero high multiple of α is algebraic. Inspired by this construction, we give examples of a non-algebraic p -torsion class in $H^6(X, \mathbb{Z})$, with $\dim_{\mathbb{C}}(X) = 5$ and with p any prime bigger than 3 (Th. 3). In particular, these classes could not be studied by the (topological) methods of [2] and [19].

In [19], Totaro gives examples of non trivial torsion classes in the Griffiths group of homologically trivial algebraic cycles modulo those algebraically equivalent to zero. Such examples were also constructed by Schoen [15], but Totaro's construction provides non trivial torsion cycles annihilated by the Abel-Jacobi map, while Schoen uses the Abel-Jacobi invariant to conclude as in Griffiths [9] that his cycles are not algebraically equivalent to 0. Note that for codimension 2 cycles, the Abel-Jacobi map is known to be injective on torsion cycles homologous to 0 [8]. Theorem 4, again inspired by Kollár's argument, gives new examples of such algebraic cycles, which furthermore cannot be detected by Totaro's method (nor indeed by any locally constant invariant). More generally, we get non trivial algebraic cycles in any level of the Hiroshi Saito filtration on Chow groups [14] (Theorem 5).

On the positive side, Bloch made the beautiful remark [4] that a conjecture of his and Kato on the Milnor K -theory of fields [6] implies that any torsion class in $H^n(X, \mathbb{Z})$, $n > 0$, is supported in codimension one. We note in Theorem 7 that it implies also that the image of this class by the Atiyah-Hirzebruch differentials are supported in codimension two.

Sections 1 and 5 (resp. 2, 3, 4) are due to the first (resp. second) author. This work started at the Hodge's Centennial Conference in Edinburgh, 2003. Both authors thank the organizers of that meeting for their invitation. We also thank P. Guillot and B. Totaro for useful comments, and J. Kollár for allowing us to reproduce his arguments in section 2.

1 On the counterexamples of Atiyah and Hirzebruch

Let X be a smooth projective complex manifold, $H^n(X, \mathbb{Z})$ its codimension n singular cohomology with integral coefficients, and $\text{CH}^k(X)$ the group of codimension k algebraic cycles on X modulo rational equivalence. Any codimension k cycle Z on X defines a cohomology class $[Z] \in H^{2k}(X, \mathbb{Z})$, the image of which in $H^{2k}(X, \mathbb{C})$ has Hodge type (k, k) . Hodge asked

whether, conversely, any class $\alpha \in H^{2k}(X, \mathbb{Z})$ with image $\alpha_{\mathbb{C}} \in H^{2k}(X, \mathbb{C})$ of type (k, k) is of the form $\alpha = [Z]$ for some cycle Z on X ([10], end of § 2). In particular, when α is a *torsion class*, i.e. $\alpha_{\mathbb{C}} = 0$, is it algebraic? This is true when $k = 1$, but Atiyah and Hirzebruch found counterexamples when $k \geq 2$ [2]. In these examples, to prove that $\alpha \in H^{2k}(X, \mathbb{Z})$ is not algebraic, they use the following criterion. Consider the spectral sequence of generalized cohomology, with

$$E_2^{st}(X) = \begin{cases} H^s(X, \mathbb{Z}) & \text{if } t \text{ is even} \\ 0 & \text{if } t \text{ is odd,} \end{cases}$$

which converges to the topological K -groups $K_{\text{top}}^{s+t}(X)$. If a class $\alpha \in H^{2k}(X, \mathbb{Z})$ is algebraic, its image by every differential d^r , $r \geq 2$, in that spectral sequence must vanish ([2], Th. 6.1).

Recently, Totaro revisited these examples of Atiyah and Hirzebruch. Let $MU^*(X)$ be the complex cobordism graded ring of X , and $MU^*(X) \otimes_{MU^*} \mathbb{Z}$ its tensor product with \mathbb{Z} over $MU^* = MU^*(\text{point})$ (which maps to $\mathbb{Z} = H^0(\text{point}, \mathbb{Z})$). Totaro proved that the cycle map

$$\text{CH}^k(X) \rightarrow H^{2k}(X, \mathbb{Z})$$

is the composite of two maps

$$\text{CH}^k(X) \rightarrow (MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k} \rightarrow H^{2k}(X, \mathbb{Z}). \quad (1.1)$$

Therefore, a torsion class which is not in the image of $(MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k}$ cannot be algebraic.

Our first remark is that, when one uses the Atiyah-Hirzebruch criterion or Totaro's factorization of the cycle map, the examples of non-algebraic torsion classes one gets must be of small order compared to the dimension of X .

Theorem 1 *Let p be a prime integer and $\alpha \in H^{2k}(X, \mathbb{Z})$ a cohomology class such that $p\alpha = 0$. Assume that p is bigger than the complex dimension of X . Then, for every $r \geq 2$, $d^r(\alpha) = 0$. Furthermore, α lies in the image of*

$$\phi^k : (MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k} \rightarrow H^{2k}(X, \mathbb{Z}). \quad (1.2)$$

Proof. The first assertion is rather standard. For any $q > 1$, the q -th Adams operation ψ^q acts upon the Atiyah-Hirzebruch spectral sequence $E_r^{st}(X)$. When $t = -2i$, its action on $E_2^{st}(X)$ is the multiplication by q^i . Since ψ^q commutes with d^r , we get

$$(q^{k+r_0} - q^k) d^r(\alpha) = 0$$

when $r = 2r_0 + 1$ (d^r is zero when r is even). For the prime number p to divide $q^{k+r_0} - q^k$ for every $q > 1$, it is necessary that $p - 1$ divides r_0 . On the other hand, since $p\alpha = 0$, we must have $k > 0$ and $d^r(\alpha)$ is not of top degree. Therefore $r \leq 2 \dim_{\mathbb{C}}(X) - 2$, hence $r_0 < \dim_{\mathbb{C}}(X)$. Since $p > \dim_{\mathbb{C}}(X)$, we get a contradiction.

Now consider the edge homomorphism

$$e^k : K_{\text{top}}^0(X) \rightarrow H^{2k}(X, \mathbb{Z})$$

in the Atiyah-Hirzebruch spectral sequence. Since $d^r(\alpha) = 0$ for every $r \geq 2$, we have

$$\alpha = e^k(\xi) \quad (1.3)$$

for some virtual bundle $\xi \in K_{\text{top}}^0(X)$. We want to show that α lies in the image of the map

$$\phi^k : (MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k} \rightarrow H^{2k}(X, \mathbb{Z}).$$

First notice that, for every $i \geq 0$, the Chern class $c_i(\xi) \in H^{2i}(X, \mathbb{Z})$ is in the image of ϕ^i . Indeed we have

$$c_i(\xi) = \phi^i(\text{cf}_{\alpha}(\xi) \otimes 1), \quad (1.4)$$

where $cf_\alpha(\xi) \in MU^{2i}(X)$ is the i -th Conner-Floyd class of ξ , with $\alpha = (i, 0, 0, \dots)$ ([1], Th. 4.1). To check (1.4), by the splitting principle, we can assume that $i = 1$, in which case it follows from the definitions (loc. cit.).

Since ϕ^* is a ring homomorphism, any polynomial in the Chern classes $c_i(\xi)$ lies in its image. Let $N_k(c_1, \dots, c_k)$ be the k -th Newton polynomial. We claim that

$$k! e^k(\xi) = N_k(c_1(\xi), \dots, c_k(\xi)). \quad (1.5)$$

To check (1.5), since ξ is the pull-back of a vector bundle on a Grassmannian, we can assume that $H^*(X, \mathbb{Z})$ is torsion free. Then it is enough to prove that the image of $e^k(\xi)$ in the rational cohomology $H^{2k}(X, \mathbb{Q})$ is

$$ch_k(\xi) = N_k(c_1(\xi), \dots, c_k(\xi))/k!.$$

But this identity follows from the fact that the Chern character

$$ch : K_{\text{top}}^*(-) \rightarrow H^*(-, \mathbb{Q})$$

is a morphism of extraordinary cohomology theories ([2], § 2).

Since $p\alpha = 0$ with $p > \dim_{\mathbb{C}}(X) \geq k$, and since $k! e^k(\xi)$ lies in the image of ϕ^k by (1.4), we deduce from (1.3) and (1.5) that α is also in the image of ϕ^k .

2 An argument due to Kollár

We start this section by describing a method due to Kollár [12], which produces examples of smooth projective complex varieties X , together with an even degree integral cohomology class α , which is not algebraic, that is, which is not the cohomology class of an algebraic cycle of X , while a non-zero multiple of α is algebraic. This is another sort of counterexample to the Hodge conjecture over the integers, since the class α is of course a Hodge class, the other known examples being that of torsion classes [3] that we shall revisit in section 3.

The examples are as follows : consider a smooth hypersurface $X \subset \mathbb{P}^{n+1}$ of degree D . For $l < n$ the Lefschetz theorem on hyperplane sections says that the restriction map

$$H^l(\mathbb{P}^{n+1}, \mathbb{Z}) \rightarrow H^l(X, \mathbb{Z})$$

is an isomorphism. Since the left-hand side is isomorphic to $\mathbb{Z}H^k$ for $l = 2k < n$, where H is the cohomology class of a hyperplane, and 0 otherwise, we conclude by Poincaré duality on X that for $2k > n$, we have $H^{2k}(X, \mathbb{Z}) = \mathbb{Z}\alpha$, where α is determined by the condition $\langle \alpha, h^{n-k} \rangle = 1$, with the notation $h = H|_X = c_1(\mathcal{O}_X(1))$. Note that the class $D\alpha$ is equal to h^k , (both have intersection number D with h^{n-k}), hence is algebraic.

In the sequel, we consider for simplicity the case where $n = 3$, $k = 2$. Then $D\alpha$ is the class of a plane section of X .

Theorem 2 (Kollár, [12]) *Assume that for some integer p coprime to 6, p^3 divides D . Then for general X , any curve $C \subset X$ has degree divisible by p . Hence the class α is not algebraic.*

Recall that “general” means that the defining equation for X has to be chosen away from a specified union of countably many Zariski closed proper subsets of the parameter space.

Proof. Let $D = p^3s$, and let $Y \subset \mathbb{P}^4$ be a degree s smooth hypersurface. Let ϕ_0, \dots, ϕ_4 be sections of $\mathcal{O}_{\mathbb{P}^4}(p)$ without common zeroes. They provide a map

$$\phi : Y \rightarrow \mathbb{P}^4,$$

which for a generic choice of the ϕ_i 's satisfies the following properties :

1. ϕ is generically of degree 1 onto its image, which is a hypersurface $X_0 \subset \mathbb{P}^4$ of degree $p^3 s = D$.
2. ϕ is two-to-one generically over a surface in X_0 , three-to-one generically over a curve in X_0 , at most finitely many points of X_0 have more than 3 preimages, and no point has more than 4 preimages.

Let \mathbb{P}^N be the projective space of all polynomials of degree D on \mathbb{P}^4 , and let $\mathcal{X} \rightarrow \mathbb{P}^N$ be the universal hypersurface. Introduce the relative Hilbert schemes (cf [11])

$$\mathcal{H}_\nu \rightarrow \mathbb{P}^N,$$

parameterizing pairs $\{(Z, X), Z \subset X\}$, where Z is a 1-dimensional subscheme with Hilbert polynomial ν . The Hilbert polynomials ν encode the degree and genus of the considered subschemes, hence there are only countably many of them. The important points are the following :

- The morphism $\rho_\nu : \mathcal{H}_\nu \rightarrow \mathbb{P}^N$ is projective.
- There exists a universal subscheme

$$\mathcal{Z}_\nu \subset \mathcal{H}_\nu \times_{\mathbb{P}^N} \mathcal{X}$$

which is flat over \mathcal{H}_ν .

Let U be the set

$$\mathbb{P}^N \setminus \bigcup_{\nu \in I} \rho_\nu(\mathcal{H}_\nu),$$

where the set I is the set of Hilbert polynomials ν for which the map ρ_ν is not dominating. Let now $X \subset \mathbb{P}^4$ be a smooth hypersurface which is parameterized by a point $x \in U$ (so X is general). Let $C \subset X$ be a curve. The reduced structure on C makes C into a subscheme of X , which is parameterized by a point $c_x \in \mathcal{H}_\nu$ over x , for some ν . By definition of U , since $x = \rho_\nu(c_x)$, the map ρ_ν has to be dominating, hence surjective. Hence it follows that there is a point $c_0 \in \mathcal{H}_\nu$ over the point x_0 parameterizing the hypersurface X_0 . The fiber Z_0 of the universal subscheme \mathcal{H}_ν over c_0 provides a subscheme $Z_0 \subset X_0$, which by flatness has the same degree as C . Let z_0 be the associated cycle of X_0 . Recall the normalization map

$$\phi : Y \rightarrow X_0.$$

By property 2 above, there exists a 1-cycle \tilde{z}_0 in Y such that $\phi_*(\tilde{z}_0) = 6z_0$. It follows that

$$6 \deg z_0 = \deg \phi_*(\tilde{z}_0).$$

On the other hand, the right-hand side is equal to the degree of the line bundle $\phi^* \mathcal{O}_{X_0}(1)$ computed on the cycle \tilde{z}_0 . Since $\phi^* \mathcal{O}_{X_0}(1)$ is equal to $\mathcal{O}_Y(p)$ (recall that $Y \subset \mathbb{P}^4$ was a hypersurface of degree s), it follows that this degree is divisible by p . Hence we found that $6 \deg C$ is divisible by p , and since p is coprime to 6, it follows that $\deg C$ is also divisible by p . ■

Remark 1 In contrast, one can show that there exists a countable union of proper algebraic subsets, which is dense in the parameter space \mathbb{P}^N , parameterizing hypersurfaces X for which the class α is algebraic. It suffices for this to prove that the set of surfaces of degree D carrying an algebraic class $\lambda \in H^2(S, \mathbb{Z}) \cap H^{1,1}(S)$ satisfying the property that $\langle \lambda, c_1(\mathcal{O}_S(1)) \rangle$ is coprime to D , is dense in the space of all surfaces of degree D in \mathbb{P}^3 . Indeed, for any X containing such a surface, the class α is algebraic on X .

Now this fact follows from the density criterion for the Noether-Lefschetz locus explained in [21] 5.3.4, and from the fact that rational classes $\lambda \in H^2(S, \mathbb{Q})$ such that a multiple $b\lambda$ is integral, and satisfies $\langle b\lambda, c_1(\mathcal{O}_S(1)) \rangle = a$ with a coprime to D , are dense in $H^2(S, \mathbb{Q})$.

To conclude this section, we note that Kollár's construction works only for high degree hypersurfaces, and indeed hypersurfaces X of degree $\leq 2n - 1$ in \mathbb{P}^{n+1} contain lines, whose class is equal to the positive generator of $H^{2n-2}(X, \mathbb{Z})$. So the following question might still have a positive answer :

Let X be a Fano variety or more generally a rationally connected variety of dimension n . Is the Hodge conjecture true for the integral cohomology classes of degree $2n - 2$ on X ?

Note that all such classes are of type $(n-1, n-1)$, since $H^2(X, \mathcal{O}_X) = 0 = H^{n-2}(X, K_X)^*$. Since the rational Hodge conjecture is known to be true for degree $2n - 2$ classes, it is always true that a multiple of such a class is algebraic. Note also that the question has a negative answer for classes of degree $2n - 2k$, $n - 3 \geq k \geq 2$, at least in the rationally connected case. Indeed, it suffices to start with one of Kollár's examples $X \subset \mathbb{P}^{n+1} \subset \mathbb{P}^{n+l}$, $l \geq 2$, and to blow-up X in \mathbb{P}^{n+l} . Since X has a degree $2n - 2$ integral class which is of Hodge type $(n - 1, n - 1)$, but is not algebraic, the resulting variety Y , which is $n + l$ -dimensional and rationally connected, has degree $2n + 2s$, $l - 2 \geq s \geq 0$ integral classes which are of Hodge type $(n + s, n + s)$, but are not algebraic.

One reason to ask this question is the fact that there is the following criterion for rationality :

Lemma 1 *Let X be a variety which is birationally equivalent to \mathbb{P}^n . Then any integral class of degree $2n - 2$ on X is algebraic. Furthermore, the Hodge conjecture is true for degree 4 integral Hodge classes on X .*

Proof. In both cases, there is a variety Y which admits a morphism of degree 1, $\phi : Y \rightarrow X$, and is obtained from \mathbb{P}^n by a sequence of blow-ups along smooth centers. Using the ϕ_* map, one concludes that if the statement is true for Y , it is true for X . Since the statement is true for \mathbb{P}^n , it suffices then to show that if the statement is true for a smooth projective variety Z , it is true for the blowing-up Z_W of Z along a smooth center W . But the supplementary classes of degree $2n - 2$ on Z_W are generated by classes of curves contracted by the blow-down map, hence they are algebraic. The supplementary integral Hodge classes of degree 4 on Z_W come from integral Hodge classes of degree 2 or 0 on W , hence they are also algebraic. ■

3 Application to torsion classes

As said in the first section, examples of even degree torsion cohomology classes which are not algebraic were first found by Atiyah and Hirzebruch [3]. They exhibited topological obstructions for a torsion class to be the cohomology class of an algebraic cycle. These obstructions were reinterpreted by Totaro [19], who stated the following criterion for algebraicity :

() For a degree $2k$ class to be algebraic, it has to be in the image of the map ϕ^k of (1.2).*

This leads to the construction of even degree torsion classes which are not algebraic. On the other hand, as shown by Theorem 1, these examples must be of large dimension, and the criterion above cannot be applied to provide for any prime p , examples of p -torsion classes of a given degree on projective manifolds of given dimension, which are not algebraic.

In this section we apply Kollár's argument to construct, for any prime $p \geq 5$, examples of p -torsion cohomology classes of degree 6 on smooth projective varieties X of dimension 5, which are not algebraic.

Furthermore, the fact that these classes are not algebraic cannot be detected by topological arguments. Indeed, any obstruction to algebraicity which is locally constant on the parameter space of X must vanish on these classes, as we show that they become algebraic on a dense subset of the parameter space of X . In particular, these classes are in the image of the map ϕ^k , which shows that criterion (*) is not sufficient.

The construction is as follows : let $p \geq 5$ be a prime integer. Let S be a surface which admits a copy of $\mathbb{Z}/p\mathbb{Z}$ as a direct summand in $H^2(S, \mathbb{Z})$. Such a surface can be constructed by a Godeaux type construction : namely one can take for S the quotient of a

degree p smooth surface Σ in \mathbb{P}^3 , defined by an equation invariant under g , where g acts on homogeneous coordinates by

$$g^*X_i = \zeta^i X_i, i = 0, \dots, 3.$$

Here ζ is a p -th root of unity, so that g has order p . As Σ is simply connected, one sees easily that the torsion of $H^2(S, \mathbb{Z})$ is isomorphic to $\mathbb{Z}/p\mathbb{Z}$, generated by $c := c_1(\mathcal{L})$, where \mathcal{L} is any of the p -torsion line bundles on S corresponding to a non trivial character of $\mathbb{Z}/p\mathbb{Z}$. Note that the class c is not divisible by p , since the torsion of $H^2(S, \mathbb{Z})$ is isomorphic to $\mathbb{Z}/p\mathbb{Z}$.

We now consider a hypersurface $X \subset \mathbb{P}^4$ of degree p^3 . Recall from the previous section that $H^4(X, \mathbb{Z})$ is generated by α , where α satisfies the condition

$$\langle \alpha, h \rangle = 1.$$

The class $\gamma := pr_1^*c \cup pr_2^*\alpha$ is a degree 6 cohomology class on $S \times X$, which is of p -torsion. Observe that, according to remark 1, this class is algebraic for a dense set of parameters for X . Indeed, since $c = c_1(\mathcal{L})$ is algebraic, that is $c = [D]$ for some divisor D of S , once α is algebraic, say $\alpha = [Z]$, $Z \in \mathcal{Z}^2(X)$, we also have

$$\gamma = [pr_1^*D \cap pr_2^*Z].$$

We now have :

Theorem 3 *For general X , the class γ is not algebraic. More precisely, for X general in modulus, and for any surface $T \subset S \times X$, the Künneth component $[T]^{2,4}$ of $[T]$ which lies in $H^2(S, \mathbb{Z}) \otimes H^4(X, \mathbb{Z})$ is of the form $t \otimes \alpha$, where t is divisible by p .*

(Note that the Künneth decomposition is well defined for $S \times X$, because the cohomology of X has no torsion.)

Proof. For X, T as in the theorem, let the Künneth component $[T]^{2,4}$ of $[T]$ be of the form $t \otimes \alpha$. If j is the natural inclusion map of X into \mathbb{P}^4 , it follows that the Künneth component of $[(Id, j)(T)]$ which is of type $(2, 6)$ is equal to $t \otimes j_*\alpha$. Since the class $j_*\alpha$ generates $H^6(\mathbb{P}^4, \mathbb{Z})$, the statement is equivalent to the fact that $[(Id, j)(T)]^{2,6}$ is divisible by p .

We now apply the argument of section 2, except that instead of considering the relative Hilbert schemes of 1-dimensional subschemes in hypersurfaces X , we consider the relative Hilbert schemes of 2-dimensional subschemes in the products $S \times X$.

This provides us with a set U of parameters, which is the complementary set of a countable union of proper algebraic subsets in \mathbb{P}^N , and has the property that any surface $T \subset S \times X$ admits a flat specialization $T_0 \subset S \times X_0$, for any specialization of X .

We will take for X_0 the image of a generic morphism

$$\phi : \mathbb{P}^3 \rightarrow \mathbb{P}^4$$

given by a base-point free linear system of polynomials degree p .

Now, by property 2 (see section 2) of the morphism ϕ , there exists a 2-cycle \tilde{t}_0 in $S \times \mathbb{P}^3$, such that, denoting by t_0 the cycle associated to the subscheme T_0 , we have the following equality of cycles in $S \times X_0$:

$$12t_0 = (Id, \phi_0)_*\tilde{t}_0,$$

where ϕ_0 is the map ϕ viewed as a map from \mathbb{P}^3 to X_0 . It follows that we also have the equality of 2-cycles in $S \times \mathbb{P}^4$:

$$12(Id, j_0)_*t_0 = (Id, \phi)_*\tilde{t}_0,$$

where j_0 is the inclusion of X_0 in \mathbb{P}^4 . This equality translates into an equality between cycle classes :

$$12[(Id, j_0)_*(t_0)] = [(Id, \phi)_*\tilde{t}_0],$$

and between their Künneth components of type (2, 6) :

$$12[(Id, j_0)_*t_0]^{2,6} = [(Id, \phi)_*\tilde{t}_0]^{2,6}. \quad (3.6)$$

Now note that since T_0 is a flat specialization of T , we have $[(I, j_0)(T_0)] = [(Id, j)(T)]$ and hence

$$[(Id, j_0)_*t_0]^{2,6} = [(Id, j)(T)]^{2,6} = t \otimes j_*\alpha.$$

So the left-hand side in (3.6) is equal to $12t \otimes j_*\alpha$. On the other hand, the right-hand side in (3.6) is equal to $(Id, \phi)_*([\tilde{t}_0]^{2,4})$, where $[\tilde{t}_0]^{2,4}$ is the Künneth component of type (2, 4) of the class $[\tilde{t}_0]$. Writing

$$[\tilde{t}_0]^{2,4} = t' \otimes \beta,$$

where β is the positive generator of $H^4(\mathbb{P}^3, \mathbb{Z})$, we have now

$$(Id, \phi)_*([\tilde{t}_0]^{2,4}) = t' \otimes \phi_*\beta.$$

But since ϕ is given by polynomials of degree p , the class $\phi_*\beta$ is equal to p times the positive generator of $H^6(\mathbb{P}^4, \mathbb{Z})$. Hence $(Id, \phi)_*([\tilde{t}_0]^{2,4})$ is divisible by p , and since p is coprime to 12, so is $t \otimes j_*\alpha$. ■

4 Application to torsion cycles

We apply in this section Kollár's degeneration argument to construct interesting torsion algebraic cycles on smooth projective complex varieties.

In [19], Totaro constructed examples of cycles which are homologous to 0 and annihilated by the Abel-Jacobi map, but are not algebraically equivalent to 0. In fact they are not divisible, while it is well-known that the groups $CH^k(X)_{alg}$ of algebraically equivalent to 0 cycles are divisible. Totaro plays on the factorization of the map

$$CH^k(X) \rightarrow (MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k}$$

of (1.1) through algebraic equivalence. Denoting by

$$Griff^k(X) = \mathcal{Z}^k(X)_{hom} / \mathcal{Z}^k(X)_{alg}$$

the quotient of the group of cycles homologous to 0 by its subgroup $\mathcal{Z}^k(X)_{alg}$, this provides an invariant

$$Griff^k(X) \rightarrow (MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k}$$

with value in the kernel of the map $\phi^k : (MU^*(X) \otimes_{MU^*} \mathbb{Z})^{2k} \rightarrow H^{2k}(X, \mathbb{Z})$ of (1.2).

These are the invariants used by Totaro to detect non trivial torsion elements in the Griffiths group. The cycles constructed there turn out to be also annihilated by the Abel-Jacobi map.

The construction we give here will provide torsion cycles annihilated by the Deligne cycle class (i.e. cohomology class and Abel-Jacobi map) and also by the Totaro invariants, but which are non divisible, hence non trivial modulo algebraic equivalence. A variant of this construction (Theorem 5) also shows that one can construct such a cycle in any level of the Hiroshi Saito filtration on the groups CH^* (cf [14]).

We shall also construct non trivial torsion cycles, which are algebraically equivalent to 0 and annihilated by the Deligne cycle class map.

The interesting point is that in all three cases, the cycles cannot be detected by any locally constant invariant associated to a torsion cycle $Z \in CH^*(X)_{tors}$ in a smooth complex projective variety X . By this, we mean an invariant which takes value in a locally constant sheaf on the parameter space for X . Indeed, our construction provides torsion cycles on smooth varieties which have parameters, and we shall see that for some special value of the parameter, our cycles become rationally equivalent to 0. Hence any locally constant invariant attached to them is 0. In the case of cycles modulo algebraic equivalence, this is to our knowledge the first example of this phenomenon. Recall that, in contrast, for fixed X , torsion cycles are known to be discrete, by the following lemma (cf [13]):

Lemma 2 (Roitman) *If X is projective, W is smooth and connected, and $\Gamma \in CH^k(W \times X)$ satisfies the property that for any $w \in W$, the cycle $\Gamma_*w \in CH^k(X)$ is of torsion, then Γ_*w is in fact constant. Hence, if it vanishes at some point, it vanishes for any w .*

Our examples show that the last statement is not true if X is allowed to deform, even staying smooth.

Note finally that, while Bloch-Esnault's [5] and Totaro's non divisible cycles are defined over number fields, ours might well not be, since we have to restrict to the general point of a parameter space, which might exclude all the points defined over some number field.

The general idea of the construction is as follows : X will be again a hypersurface of degree p^3 in \mathbb{P}^4 , but we assume that X contains a curve C of degree p . We arrange things in such a way that the curve C is not divisible by p in $CH^2(X)$, as in the previous section. This suggests that the cycle

$$pr_1^*c \cdot pr_2^*C$$

on $S \times X$, where the notations are as in section 3), is non trivial, which is indeed what we prove.

Let us now give the detailed construction : Fix a prime integer $p \geq 5$. We will consider the following morphism

$$\phi : \mathbb{P}^3 \rightarrow \mathbb{P}^4,$$

$$\phi(x_0, \dots, x_3) = (x_0^p, \dots, x_3^p, f(x_0, \dots, x_3)), \quad (4.7)$$

where f is a degree p generic homogeneous polynomial. So $\phi(\mathbb{P}^3) =: X_0$ is a hypersurface of \mathbb{P}^4 of degree p^3 . For any line $l \subset \mathbb{P}^3$, the image $\phi(l)$ is a curve C_0 of degree p , and if the line is conveniently chosen, this curve is smooth, hence is contained in a smooth hypersurface X_∞ of degree p^3 .

Consider the pencil $(X_t)_{t \in \mathbb{P}^1}$ generated by X_0 and X_∞ . For any $t \in \mathbb{P}^1$, let $C_t \subset X_t$ be the curve C_0 , viewed as a curve in X_t . Note that for smooth X_t , the cohomology class of C_t is divisible by p : it is equal to $p\alpha$ where α was introduced in the previous section. It follows that the Deligne cohomology invariant $[C_t]_D$ of C_t is also divisible by p , since there is the exact sequence

$$0 \rightarrow J^3(X_t) \rightarrow H_D^4(X_t, \mathbb{Z}(2)) \rightarrow H^4(X_t, \mathbb{Z}) \rightarrow 0,$$

and the intermediate Jacobian $J^3(X_t)$ is a divisible group.

Let now W be a smooth projective variety and \mathcal{L} a p -torsion line bundle on W . In applications, W will be either a surface as in the previous section, and \mathcal{L} will not be topologically trivial, with $c_1(\mathcal{L})$ not divisible by p in $H^2(S, \mathbb{Z})$, or a curve of genus > 0 , and \mathcal{L} will be topologically trivial, but not trivial. Let us denote $c = cl(\mathcal{L}) \in CH^1(W)$. Since \mathcal{L} is of p -torsion, so is the Deligne cohomology invariant $[c]_D \in H_D^2(W, \mathbb{Z}(1))$.

The cycle we will consider is

$$c \times C_t := pr_1^*c \cdot pr_2^*C_t \in CH^3(W \times X_t).$$

Note that this cycle has vanishing Deligne invariant, since

$$[c \times C_t]_D = pr_1^*[c]_D \cdot_D pr_2^*[C_t]_D,$$

where \cdot_D is the product in Deligne cohomology, and the left factor is p -torsion while the right factor is divisible by p . This cycle is of p -torsion, since c is, and it is algebraically equivalent to 0 when c is, for example in the case where W is a curve.

We have now the following :

Theorem 4 1) For a general point $t \in \mathbb{P}^1$, the p -torsion cycle $c \times C_t \in CH^3(W \times X_t)$ is non 0.

2) If c is not divisible by p in $CH^1(W)$, so is $c \times C_t$ in $CH^3(W \times X_t)$ for general t , hence in particular $c \times C_t$ is non trivial in $Griff^3(W \times X_t)$.

Proof. Let

$$\mathcal{X} := \{(x, t) \in \mathbb{P}^4 \times \mathbb{P}^1 / x \in X_t\}.$$

\mathcal{X} contains the 2-cycle $\Gamma := C_0 \times \mathbb{P}^1$, and $W \times \mathcal{X}$ contains the cycle

$$c \times \Gamma := pr_1^* c \cdot pr_2^* \Gamma.$$

Each $W \times X_t \xrightarrow{j_t} W \times \mathcal{X}$ is the fiber of the natural composite map

$$W \times \mathcal{X} \xrightarrow{pr_2} \mathcal{X} \xrightarrow{f} \mathbb{P}^1,$$

where f is the map to \mathbb{P}^1 given by the pencil. The restrictions $j_t^* : CH_i(W \times \mathcal{X}) \rightarrow CH_{i-1}(W \times X_t)$ are well defined even if $W \times \mathcal{X}$ is singular, because j_t is the inclusion of a Cartier divisor. We have $j_t^*(c \times \Gamma) = c \times C_t$ in $CH_i(W \times X_t)$, where $i = \dim W$.

Lemma 3 1) Assume that for a general complex point $t \in \mathbb{P}^1$, the restriction $j_t^*(c \times \Gamma)$ is equal to 0 in $CH_i(W \times X_t)$. Then for any $t \in \mathbb{P}^1$, the restriction $j_t^*(c \times \Gamma)$ is equal to 0 in $CH_i(W \times X_t)$.

2) Similarly, if for a general complex point $t \in \mathbb{P}^1$, the restriction $j_t^*(c \times \Gamma)$ is equal to 0 in $CH_i(W \times X_t) \otimes \mathbb{Z}/p\mathbb{Z}$, then the same is true for any $t \in \mathbb{P}^1$.

Proof. Indeed, the assumption in 1) implies that there exist a smooth projective curve D and a finite morphism $r : D \rightarrow \mathbb{P}^1$, such that denoting by

$$\mathcal{X}_D \xrightarrow{\tilde{f}} D, \mathcal{X}_D \xrightarrow{\tilde{r}} \mathcal{X}$$

the fibered product $\mathcal{X} \times_{\mathbb{P}^1} D$, the cycle

$$(Id, \tilde{r})^*(c \times \Gamma) \in CH_{i+1}(W \times \mathcal{X}_D)$$

vanishes on some dense open set $W \times \mathcal{X}_U$, where $\mathcal{X}_U := \tilde{f}^{-1}(U)$ for some dense Zariski open set $U \subset D$.

Similarly in case 2), the cycle

$$(Id, \tilde{r})^*(c \times \Gamma) \in CH_{i+1}(W \times \mathcal{X}_D)$$

will vanish modulo p on some dense open set $W \times \mathcal{X}_U$.

This implies that the cycle $(Id, \tilde{r})^*(c \times \Gamma)$ is supported on fibers of the map $\tilde{f} \circ pr_2 : W \times \mathcal{X}_D \rightarrow D$, that is

$$(Id, \tilde{r})^*(c \times \Gamma) = \sum_d \tilde{j}_d^* \gamma_d \text{ in } CH_{i+1}(W \times \mathcal{X}_D), \quad (4.8)$$

where \tilde{j}_d is the inclusion of $W \times \tilde{f}^{-1}(d) \cong W \times f^{-1}(r(d))$ in $W \times \mathcal{X}_D$, and the sum on the right is finite. Similarly in case 2), we will get the same equation as in (4.8), but modulo p .

For any $t \in \mathbb{P}^1$, let $t' \in D$ be such that $r(t') = t$. Then we have

$$\tilde{j}_{t'}^*((Id, \tilde{r})^*(c \times \Gamma)) = j_t^*(c \times \Gamma) \text{ in } CH_i(W \times X_t). \quad (4.9)$$

But for any $d \in D$, the cycle

$$\tilde{j}_{t'}^*(\tilde{j}_{d*}(\gamma_d))$$

vanishes in $CH_i(W \times X_t)$ since either $t' \neq d$ and the two fibers do not meet, or $t' = d$, and then $\tilde{j}_{t'}^* \circ \tilde{j}_{d*}$ is by definition the intersection with the class of the Cartier divisor $cl(\mathcal{O}_{W \times X_D}(W \times X_{t'})|_{W \times X_{t'}})$, which is trivial.

Combining this with (4.8) and (4.9), we have shown that $j_t^*(c \times \Gamma) = 0$ in $CH_i(W \times X_t)$ in case 1) and $j_t^*(c \times \Gamma) = 0$ in $CH_i(W \times X_t) \otimes \mathbb{Z}/p\mathbb{Z}$ in case 2). ■

We now conclude the proof of Theorem 4 by contradiction. Assuming the conclusion of 1) or 2) in the theorem is wrong, we conclude using Lemma 3 that for $t = 0$, the cycle $c \times C_0 \in CH_i(W \times X)$ is trivial (resp. is trivial mod. p in case 2)). This means that we can write

$$c \times C_0 = \sum_i \tau_{i*} \text{div } \phi_i \text{ in } \mathcal{Z}_i(W \times X_0), \quad (4.10)$$

where W_i is normal irreducible of dimension $i + 1$, ϕ_i is a non zero rational function on W_i , and $\tau_i : W_i \rightarrow W \times X$ is proper. In case 2), this equality will be true in $\mathcal{Z}_i(W \times X_0) \text{ mod } p$.

We observe now that the map $\phi : \mathbb{P}^3 \rightarrow X_0$ has the following property :

Lemma 4 *For any irreducible closed algebraic subset $Z \subset \mathbb{P}^3$, the restriction $\phi|_Z$ has generic degree 1 onto its image.*

Proof. Indeed, if Z is as above, and ϕ has degree ≥ 2 on Z , then looking at the first 4 coordinates (x_0^p, \dots, x_3^p) of ϕ , one concludes that Z has to be invariant under a group $\mathbb{Z}/p\mathbb{Z}$ acting non trivially on Z , with generator g acting on coordinates by

$$g^*(X_i) = \lambda_i X_i,$$

with $\lambda_i^p = 1$. Furthermore, we must have, looking at the last coordinate of ϕ : $g^*f = f$ on Z . Thus Z has to be contained in the zero locus of

$$f - g^*f, f - (g^2)^*f, \dots, f - (g^{p-1})^*f.$$

But as f is generic, and $p \geq 5$, this zero locus consists of the fixed points of g , for any such g . Thus g acts trivially on Z , which is a contradiction. ■

It follows that the $\tau_i : W_i \rightarrow W \times X_0$ lift to $\tilde{\tau}_i : W_i \rightarrow W \times \mathbb{P}^3$, so that equation (4.10) provides

$$c \times l = \sum_i \tilde{\tau}_{i*} \text{div } \phi_i + z \text{ in } \mathcal{Z}_i(W \times \mathbb{P}^3), \quad (4.11)$$

where we recall that l is the line such that $\phi(l) = C_0$ and where z is a cycle which satisfies :

$$(Id, \phi)_* z = 0 \text{ in } \mathcal{Z}_i(W \times X_0). \quad (4.12)$$

In case 2), this equation becomes :

$$c \times l = \sum_i \tilde{\tau}_{i*} \text{div } \phi_i + z + pz' \text{ in } \mathcal{Z}_i(W \times \mathbb{P}^3), \quad (4.13)$$

for some cycle z' , where z satisfies property (4.12). Let $H = cl(\mathcal{O}_{\mathbb{P}^3}(1)) \in CH^1(\mathbb{P}^3)$. We have a map

$$q : CH_i(W \times \mathbb{P}^3) \rightarrow CH_{i-1}(W) = CH^1(W),$$

defined by

$$q(\gamma) = pr_{1*}(\gamma \cdot pr_2^* H).$$

We observe that, because $\langle l, H \rangle = 1$, we have

$$q(c \times l) = c.$$

Applying q to the right-hand sides in equations (4.11) and (4.13) gives now

$$c = q(z) \text{ in } CH^1(W), \text{ resp. } c = q(z) \text{ in } CH^1(W) \otimes \mathbb{Z}/p\mathbb{Z} \text{ in case 2),} \quad (4.14)$$

where z satisfies property (4.12). We have now the following lemma :

Lemma 5 *If $z \in \mathcal{Z}_i(W \times \mathbb{P}^3)$ satisfies property (4.12), then $z = 0$ in $CH_i(W \times \mathbb{P}^3)$.*

Proof. The kernel of the map $(Id, \phi)_*$ defined on cycles is generated by cycles $z_1 - z_2$, where z_i 's are effective and $(Id, \phi)_*(z_1) = (Id, \phi)_*(z_2)$. By Lemma 4, we can even assume that the z_i are irreducible and $(Id, \phi)(z_1) = (Id, \phi)(z_2)$; looking at the form of the map ϕ given in (4.7), we see that this is equivalent to the following :

There exists $\lambda = (\lambda_0, \dots, \lambda_3)$, where λ_i are p -th roots of unity, acting on \mathbb{P}^3 by $\lambda.(x_0, \dots, x_3) = (\lambda_0 x_0, \dots, \lambda_3 x_3)$, such that on z_1 the equality $f(\lambda.(x_0, \dots, x_3)) = f(x_0, \dots, x_3)$ is satisfied, and we have

$$z_2 = (Id, \lambda.)(z_1).$$

But the map $(Id, \lambda.)$ acts as the identity on $CH(W \times \mathbb{P}^3)$, because its graph is rationally equivalent to the graph of the identity. Hence z_2 is rationally equivalent to z_1 . ■

It follows from this lemma that equation (4.14) becomes in fact :

$$c = 0 \text{ in } CH^1(W), \text{ resp. } c = 0 \text{ in } CH^1(W) \otimes \mathbb{Z}/p\mathbb{Z} \text{ in case 2),} \quad (4.15)$$

which contradicts the fact that $c \neq 0$, resp. $c \neq 0 \pmod{p}$ in case 2). This concludes the proof of the theorem. ■

Theorem 4, applied to the case where \mathcal{L} is topologically trivial, hence c is algebraically equivalent to 0, provides non trivial torsion cycles algebraically equivalent to 0 and annihilated by the Abel-Jacobi map. Since cycles algebraically equivalent to 0 are images via correspondences of 0-cycles homologous to 0, this is to be put in contrast with Roitman's theorem [13], which says that the Albanese map is injective on the torsion part of the groups CH_0 .

As explained below, these cycles can in fact be specialized to 0 on a smooth fiber of a flat family of varieties, so that more generally any invariant of locally constant type has to vanish on them. Note that Chad Schoen [16] also constructed independently non zero torsion cycles, algebraically equivalent to 0, with trivial specialization.

Next, in the case where $W = S$ is a surface and \mathcal{L} is as in section 3, so that $c_1(\mathcal{L})$ is a non divisible class in $H^2(S, \mathbb{Z})$, Theorem 4 shows that for general t , the cycle $c \times C_t$ is a torsion cycle which is not algebraically equivalent to 0.

On the other hand, we can arrange things so that for some flat deformation of the pair (C_t, X_t) to a pair (C'_t, X'_t) , with X'_t smooth, $c \times C'_t$ becomes rationally equivalent to 0 on $W \times X'_t$.

Indeed, in the above construction, we may assume that the initial curve C_0 is a (singular) plane curve of degree p . Such a plane curve deforms in a flat way to a multiple line Z of multiplicity p in a plane, and one can then construct a deformation of the pair (C_t, X_t) to a pair (Z, X'_t) . One verifies that it is possible to do so with a smooth X'_t . The cycle z associated to Z is divisible by p , and it follows that $c \times z = 0$ in $CH^3(W \times X'_t)$.

It follows in particular from this that the cycle $c \times C_t$ has vanishing associated Totaro's invariant, since these are locally constant under deformation of the pair $(X, \text{torsion-cycle})$

on X), and more generally any locally constant invariant of a torsion cycle must vanish on it.

To conclude this section, let us note that the same proof can be used to construct p -torsion cycles which are not trivial modulo algebraic equivalence, (in fact non divisible) and which are in the k -th level of the Hiroshi Saito filtration F_{HSaito}^l on Chow groups, where k can be taken arbitrarily large. Recall that this filtration is smaller than any existing Bloch-Beilinson filtration, and is defined as follows : $F_{HSaito}^l CH^m(X)$ is the subgroup of $CH^m(X)$ which is generated via correspondences by products $z_1 \cdot \dots \cdot z_l$ of l cycles homologous to 0.

Namely, with the same notations as above, consider the case where $W = S$ and c is not divisible by p . Consider the cycle

$$c^k \times C_t^k = (c \times C_t)^k = \Pi_i pr_i^*(c \times C_t) \in CH^{3k}(S^k \times X_t^k).$$

It is a p -torsion cycle which lies in $F_{HSaito}^k CH^{3k}(S^k \times X^k)$, since each factor $c \times C_t$ is homologous to 0. Now we have

Theorem 5 *For general t , the cycle $c^k \times C_t^k$ is non 0 in $CH^{3k}(S^k \times X^k) \otimes \mathbb{Z}/p\mathbb{Z}$, hence in particular it is not algebraically equivalent to 0.*

Proof. We introduce as before the whole family

$$\mathcal{X}_{/\mathbb{P}^1}^k := \mathcal{X} \times_{\mathbb{P}^1} \dots \times_{\mathbb{P}^1} \mathcal{X}.$$

On $S^k \times \mathcal{X}_{/\mathbb{P}^1}^k$, we have the cycle

$$c^k \times \Gamma_k,$$

where Γ was introduced at the beginning of the proof of Theorem 4, and $\Gamma_k := \Pi_l pr_i^* \Gamma \in CH^{2k}(\mathcal{X}_{/\mathbb{P}^1}^k)$.

The cycle $c^k \times \Gamma_k$ restricts to $c^k \times C_t^k$ on $S^k \times X_t^k$, and hence, applying Lemma 3, 2), we conclude that it suffices to show that the restriction of $c^k \times \Gamma_k$ to the fiber $S^k \times X_0^k$ over 0 is non trivial modulo p .

But this restriction is equal to $c^k \times C_0^k$, where C_0 is the image of a line l in \mathbb{P}^3 via the map ϕ . We have now an analogue of Lemma 5 which works for the map $\phi^k : (\mathbb{P}^3)^k \rightarrow X_0^k$, and allows to conclude that it suffices to show that the cycle $c^k \times l^k$ is not divisible by p in $CH^{3k}(S^k \times (\mathbb{P}^3)^k)$. But the cycle c^k is not divisible by p in S^k , because its cohomology class is not divisible by p , and it follows by applying the map

$$CH^{3k}(S^k \times (\mathbb{P}^3)^k) \rightarrow CH^k(S^k),$$

$$\gamma \mapsto pr_{1*}(\gamma \cdot pr_2^* H^{\boxtimes k}),$$

that $c^k \times l^k$ is not divisible by p in $CH^{3k}(S^k \times (\mathbb{P}^3)^k)$. ■

One may wonder whether a similar construction might allow to construct examples of non zero torsion cycles in $F_{HSaito}^{n+1} CH^n(X)$ for some n and some smooth complex projective variety X . (Note that it is conjectured by Bloch and Beilinson that $F_{HSaito}^{n+1} CH^n(X)_{\mathbb{Q}} = 0$.)

5 Consequences of the Bloch-Kato conjecture

When F is a field and $n \geq 0$ an integer, we denote by $K_n^M(F)$ the n -th Milnor K -group of F . If a prime p is invertible in F there is a symbol map

$$K_n^M(F)/p \rightarrow H^n(F, \mu_p^{\otimes n})$$

from the Milnor K -theory of F modulo p to the Galois cohomology of F with coefficients in the n -th power of the Galois group of p -th roots of unity. We shall say that the *Bloch-Kato conjecture is true at the prime p* when this symbol map is an isomorphism for every $n \geq 0$

and any F of characteristic different from p [6]. Voevodsky proved that the Bloch-Kato conjecture is true at the prime 2 [20]. He and Rost are close to the proof of the Bloch-Kato conjecture at every prime.

Bloch noticed the following striking consequence of the Bloch-Kato conjecture. Let X be a smooth quasi-projective complex manifold, and $\alpha \in H^n(X, \mathbb{Z})$ an integral cohomology class. We say that α is supported in codimension q when there exists a Zariski closed subset $Y \subset X$ of codimension q such that the restriction of α to $X - Y$ vanishes.

Theorem 6 (Bloch [4], end of Lecture 5) *Assume that the Bloch-Kato conjecture is true at the prime p and that $\alpha \in H^n(X, \mathbb{Z})$ is killed by p . Then α is supported in codimension one.*

The Bloch-Kato conjecture gives also some information on the Atiyah-Hirzebruch spectral sequence computing the topological K -theory (this fact was first noticed for function fields by Thomason).

Theorem 7 *Under the same assumption as in Theorem 6, the image of α by every differential d^r , $r \geq 2$, is supported in codimension two.*

Proof. We consider the Atiyah-Hirzebruch spectral sequence with \mathbb{Z}/p coefficients:

$$E_2^{st}(X; \mathbb{Z}/p) = \begin{cases} H^s(X, \mathbb{Z}/p) & \text{when } t \text{ is even} \\ 0 & \text{when } t \text{ is odd,} \end{cases}$$

which converges to $K_{\text{top}}^{s+t}(X, \mathbb{Z}/p)$. If the Bloch-Kato conjecture is true at p , it is known [18] that the algebraic K -theory of X with \mathbb{Z}/p coefficients is the abutment of a spectral sequence

$$'E_2^{st}(X, \mathbb{Z}/p) = \begin{cases} H_{\text{Zar}}^s(X, \tau_{\leq i} R\varepsilon_* \mu_p^{\otimes i}) & \text{if } t = -2i \leq 0 \\ 0 & \text{otherwise,} \end{cases}$$

where $\varepsilon : X_{\text{ét}} \rightarrow X_{\text{Zar}}$ is the natural morphism from the big étale site of X to its big Zariski site, and $\tau_{\leq i}$ is the good truncation. There is a morphism of spectral sequences

$$'E_r^{st}(X, \mathbb{Z}/p) \rightarrow E_r^{st}(X, \mathbb{Z}/p).$$

Consider the differential

$$d^r : E_r^{n, -2n}(X, \mathbb{Z}/p) \rightarrow E_r^{n+r, -2n-r+1}(X, \mathbb{Z}/p).$$

The group

$$'E_2^{n, -2n}(X, \mathbb{Z}/p) = H_{\text{Zar}}^n(X, \tau_{\leq n} R\varepsilon_* \mu_p^{\otimes n}) = H_{\text{Zar}}^n(X, R\varepsilon_* \mu_p^{\otimes n}) = H_{\text{ét}}^n(X, \mu_p^{\otimes n})$$

maps isomorphically to

$$E_r^{n, -2n}(X, \mathbb{Z}/p) = H^n(X, \mathbb{Z}/p).$$

On the other hand, if $i = n + \frac{r-1}{2}$, the spectral sequence computing the hypercohomology of $\tau_{\leq i} R\varepsilon_* \mu_p^{\otimes i}$ is

$$''E_2^{st} = \begin{cases} H_{\text{Zar}}^s(X, R^t \varepsilon_* \mu_p^{\otimes i}) & \text{when } 0 \leq t \leq i \\ 0 & \text{otherwise,} \end{cases}$$

with abutment $H_{\text{Zar}}^{s+t}(X, \tau_{\leq i} R\varepsilon_* \mu_p^{\otimes i})$. When $s+t = n+r$ we find $''E_2^{st} = 0$ unless $n+r-s \leq i = n + \frac{r-1}{2}$, i.e.

$$r \leq 2s - 1.$$

According to Bloch-Ogus [7], any class in $''E_2^{st}$ is supported in codimension s . Therefore, any class in $H_{\text{Zar}}^{n+r}(X, \tau_{\leq i} R\varepsilon_* \mu_p^{\otimes i})$ is supported in codimension $\frac{r+1}{2} \geq 2$.

Therefore, given $x \in H^n(X, \mathbb{Z}/p) \simeq 'E_2^{n, -2n}(X, \mathbb{Z}/p)$, we can find a Zariski closed subset $Y \subset X$, $\text{codim}(Y) \geq 2$, such that, for every $r \geq 2$, the restriction of $'d^r(x) \in 'E_2^{n+r, -2n-r+1}(X, \mathbb{Z}/p)$ to $X - Y$ vanishes. Therefore the same is true for $d^r(x)$.

If $\alpha \in H^n(X, \mathbb{Z})$ is such that $p\alpha = 0$, it lies in the image of the Bockstein homomorphism

$$\beta : H^{n-1}(X, \mathbb{Z}/p) \rightarrow H^n(X, \mathbb{Z}).$$

Therefore $\alpha = \beta(x)$, and $d^r(\alpha) = \beta(d^r(x))$ is supported in codimension two.

References

- [1] ADAMS, J.F. *Stable homotopy and generalised homology*, Chicago Lectures in Mathematics, Part I: S.P. Novikov's work on operations on complex cobordism, Chicago - London: The University of Chicago Press. X, 373 p.
- [2] ATIYAH, M.F. AND HIRZEBRUCH, F. Vector bundles and homogeneous spaces, *Proc. Sympos. Pure Math.* **3**, 7-38 (1961).
- [3] ATIYAH, M.F. AND HIRZEBRUCH, F. Analytic cycles on complex manifolds, *Topology* **1**, 25-45 (1962).
- [4] BLOCH, S. *Lectures on algebraic cycles*, Duke University Mathematics Series. IV. Durham, North Carolina: Duke University, Mathematics Department.
- [5] BLOCH, S. AND ESNAULT, H. The coniveau filtration and non-divisibility for algebraic cycles, *Math. Annalen* **304**, 303-314 (1996).
- [6] BLOCH, S. AND KATO, K. p-adic étale cohomology, *Publ. Math., Inst. Hautes Étud. Sci.* **63**, 107-152 (1986).
- [7] BLOCH, S. AND OGUS, A. Gersten's conjecture and the homology of schemes, *Ann. Sci. Éc. Norm. Supér.*, IV. Sér. **7**, 181-201 (1974).
- [8] COLLIOT-THÉLÈNE, J.-L. Cycles algébriques de torsion et K-théorie algébrique, (CIME lecture notes) in *Arithmetic Algebraic Geometry* (Trento 1991), LNM 1553, Springer-Verlag Berlin (1993).
- [9] GRIFFITHS, P. On the periods of certain rational integrals, I, II, *Ann. Math.* **90** (1969), 460-541.
- [10] HODGE, W.V.D. The topological invariants of algebraic varieties, *Proc. Intern. Congr. Math.* (Cambridge, Mass., Aug. 30-Sept. 6, 1950) **1**, 182-192 (1952).
- [11] KOLLÁR, J. *Rational curves on algebraic varieties*, Ergebnisse der Math. und ihrer Grenzgebiete, Springer-Verlag (1996).
- [12] KOLLÁR, J. Lemma p. 134 in *Classification of irregular varieties*, edited by E. Ballico, F. Catanese, C. Ciliberto, Lecture Notes in Math. 1515, Springer (1990).
- [13] ROITMAN, A. The torsion of the group of zero-cycles modulo rational equivalence, *Ann. Math.* **111** (1980) 553-569.
- [14] SAITO, H. Generalization of Abel's theorem and some finiteness properties of 0-cycles on surfaces, *Compo. Math.* **84** (1992), 289-332.
- [15] SCHOEN, C. Some examples of torsion in the Griffiths group, *Math. Annalen* **293** (1992), 651-679.
- [16] SCHOEN, C. Specialization of the torsion subgroup of the Chow group, preprint 2004.
- [17] SCHOEN, C. On certain exterior product maps of Chow groups, *Math. Res. Letters* **7** (2000), 177-194.
- [18] SUSLIN, A. AND VÆVODSKY, V. Bloch-Kato conjecture and motivic cohomology with finite coefficients, (English), Gordon, B. Brent (ed.) et al., *The arithmetic and geometry of algebraic cycles*, Proceedings of the NATO Advanced Study Institute, Banff, Canada, June 7-19, 1998. Vol. 1. Dordrecht: Kluwer Academic Publishers. NATO ASI Ser., Ser. C, Math. Phys. Sci. **548**, 117-189 (2000).

- [19] TOTARO, B. Torsion algebraic cycles and complex cobordism, *Journal of the American Mathematical Society* **10**, 467-493, (1997).
- [20] VOEVODSKY, V. Motivic cohomology with $\mathbf{Z}/2$ coefficients, *Pub. Math. IHES* **98**, 59-104 (2003).
- [21] VOISIN, C. *Hodge Theory and Complex Algebraic Geometry II*, Cambridge University Press (2003).

C.Soulé: CNRS and IHES, 35 Route de Chartres, 91440 Bures sur Yvette. soule@ihes.fr

C.Voisin: CNRS, UMR 7586, Institut de Mathématiques de Jussieu, 175 rue du Chevaleret, 75013 Paris. voisin@math.jussieu.fr