

ISOMORPHISMS AMONG QUANTUM GROTHENDIECK RINGS AND PROPAGATION OF POSITIVITY

RYO FUJITA, DAVID HERNANDEZ, SE-JIN OH, AND HIRONORI OYA

ABSTRACT. Let $(\mathfrak{g}, \mathfrak{g})$ be a pair of complex finite-dimensional simple Lie algebras whose Dynkin diagrams are related by (un)folding, with \mathfrak{g} being of simply-laced type. We construct a collection of ring isomorphisms between the quantum Grothendieck rings of monoidal categories $\mathcal{C}_{\mathfrak{g}}$ and $\mathcal{C}_{\mathfrak{g}}$ of finite-dimensional representations over the quantum loop algebras of \mathfrak{g} and \mathfrak{g} respectively. As a consequence, we solve long-standing problems : the positivity of the analogs of Kazhdan-Lusztig polynomials and the positivity of the structure constants of the quantum Grothendieck rings for any non-simply-laced \mathfrak{g} . In addition, comparing our isomorphisms with the categorical relations arising from the generalized quantum affine Schur-Weyl dualities, we prove the analog of Kazhdan-Lusztig conjecture (formulated in [17]) for simple modules in remarkable monoidal subcategories of $\mathcal{C}_{\mathfrak{g}}$ for any non-simply-laced \mathfrak{g} , and for any simple finite-dimensional modules in $\mathcal{C}_{\mathfrak{g}}$ for \mathfrak{g} of type B_n . In the course of the proof we obtain and combine several new ingredients. In particular we establish a quantum analog of T -systems, and also we generalize the isomorphisms of [24, 26] to all \mathfrak{g} in a unified way, that is isomorphisms between subalgebras of the quantum group of \mathfrak{g} and subalgebras of the quantum Grothendieck ring of $\mathcal{C}_{\mathfrak{g}}$.

CONTENTS

1. Introduction	2
2. Quantum loop algebras and their finite-dimensional representations	9
3. Quantum Grothendieck rings	13
4. Combinatorics of Q-data	19
5. Monoidal subcategories associated with Q-data	26
6. Quantum T -systems	31
7. Quantized coordinate algebras	35
8. The HLO isomorphisms	41
9. R -matrices and commutation relations	48
10. Isomorphisms among quantum Grothendieck rings	53
11. Further results in type B	58
Appendix A. Quantum unipotent minors	62
Appendix B. Quantum cluster algebras	64
References	65

Date: December 14, 2021.

2010 Mathematics Subject Classification. 17B37, 20G42, 81R50, 17B10, 17B67.

1. INTRODUCTION

1.1. For a complex finite-dimensional simple Lie algebra \mathfrak{g} , the associated quantum loop algebra $U_q(L\mathfrak{g})$ serves as a natural quantum affinization of \mathfrak{g} . It can be obtained as a subquotient of the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ corresponding to the level zero representations and hence inherits a structure of Hopf algebra over $\overline{\mathbb{Q}(q)}$. In particular, finite-dimensional representations over $U_q(L\mathfrak{g})$ form a rigid monoidal category $\mathcal{C}_{\mathfrak{g}}$, whose structure is quite intricate. For example, in contrast to the classical finite-dimensional representation theory of \mathfrak{g} , it is neither semisimple as an abelian category nor braided as a monoidal category. Since the naissance of quantum groups in the middle of the 80s in the context of solvable lattice models in physics, the category $\mathcal{C}_{\mathfrak{g}}$ has been intensively studied from various perspectives.

From the representation-theoretic point of view, many fundamental results were obtained in the 90s. Chari-Pressley [5, 6] gave a complete classification of simple representations in $\mathcal{C}_{\mathfrak{g}}$ in terms of the Drinfel'd polynomials. After that, motivated by the theory of deformed \mathcal{W} -algebras, Frenkel-Reshetikhin [10] introduced the notion of q -characters of representations in $\mathcal{C}_{\mathfrak{g}}$ as a natural quantum loop analog of the usual characters for \mathfrak{g} . Their basic properties were studied in a subsequent work by Frenkel-Mukhin [9]. These results particularly imply that the Grothendieck ring $K(\mathcal{C}_{\mathfrak{g}})$ is a commutative polynomial ring generated by the classes of *fundamental modules*, whose q -characters are computable by Frenkel-Mukhin's combinatorial algorithm. These fundamental modules also have crystal bases [33] (up to a spectral parameter shift). Moreover, every simple module $L(m)$ in $\mathcal{C}_{\mathfrak{g}}$ can be obtained as a head of a unique ordered tensor product of fundamental modules, called *the standard module* $M(m)$. Here m belongs to an ordered set \mathcal{M} of *dominant monomials* which serves as a parameter set for simple modules in $\mathcal{C}_{\mathfrak{g}}$ (which is comparable with the set of Drinfel'd polynomials).

In this paper, we are particularly interested in the next step : the fundamental problem to determine the dimensions and q -characters of simple modules. One possible approach is to compute the q -characters of simple modules from those of standard modules with a *Kazhdan-Lusztig type algorithm* explained below.

1.2. When \mathfrak{g} is simply-laced (i.e., of type ADE), a remarkable Kazhdan-Lusztig type algorithm was established by Nakajima based on the geometry of quiver varieties [45, 47]. This algorithm computes the Jordan-Hölder multiplicity $P_{m,m'}$ of the simple module $L(m')$ occurring in the standard module $M(m)$. Since we have

$$[M(m)] = [L(m)] + \sum_{m' \in \mathcal{M}: m' < m} P_{m,m'} [L(m')]$$

in the Grothendieck ring $K(\mathcal{C}_{\mathfrak{g}})$, this algorithm enables us to compute all the simple q -characters in principle. It returns the multiplicity $P_{m,m'}$ as the specialization at $t = 1$ of a certain positive polynomial $P_{m,m'}(t) \in t\mathbb{Z}_{\geq 0}[t]$, which can be seen as an analog of Kazhdan-Lusztig polynomial. These polynomials $P_{m,m'}(t)$ are constructed from the *quantum Grothendieck ring* $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ which is a t -deformation of the Grothendieck ring $K(\mathcal{C}_{\mathfrak{g}})$ [47, 55]. Here we emphasize that the quiver varieties play an essential role to guarantee the validity of the algorithm $P_{m,m'}(1) = P_{m,m'}$ and the positivity of $P_{m,m'}(t)$.

When \mathfrak{g} is non-simply-laced, the above theory is not applicable for the absence of a fully developed theory of quiver varieties. However, one can still formulate a conjectural Kazhdan-Lusztig type algorithm for general \mathfrak{g} as it is possible to construct the quantum Grothendieck ring $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ in a purely algebraic way. This is what the second named author established in [18, 17]. In this algebraic setting, a certain quantum torus \mathcal{Y}_t is obtained from a different construction involving formal power series of elements in the quantum affine algebra which can be seen as vertex operators. Then, mimicking [47, 55], the quantum Grothendieck ring $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ is defined inside \mathcal{Y}_t as a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra whose specialization at $t = 1$ is identical to $K(\mathcal{C}_{\mathfrak{g}})$ via the q -character map (the purely algebraic proof of the existence of $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ in [17] works in non simply-laced types as well). By the same arguments as in [47], $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ has a standard $\mathbb{Z}[t^{\pm 1/2}]$ -basis $\{E_t(m)\}_{m \in \mathcal{M}}$, whose members are called *the (q, t) -characters of standard modules*. Under the specialization $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}}) \rightarrow K(\mathcal{C}_{\mathfrak{g}})$, the element $E_t(m)$ corresponds to the class $[M(m)]$ and hence the name. As an analog of the Kazhdan-Lusztig basis, we can construct a canonical $\mathbb{Z}[t^{\pm 1/2}]$ -basis $\{L_t(m)\}_{m \in \mathcal{M}}$ of $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ characterized by the invariance under a standard bar-involution and the unitriangular property

$$E_t(m) = L_t(m) + \sum_{m' \in \mathcal{M}: m' < m} P_{m, m'}(t) L_t(m')$$

for some $P_{m, m'}(t) \in t\mathbb{Z}[t]$. The polynomials $P_{m, m'}(t)$ are nothing but the desired analogs of Kazhdan-Lusztig polynomials.

1.3. With the above construction, the following conjectures naturally arise.

Conjecture 1.1 (Analog of Kazhdan-Lusztig conjecture, [17, Conjecture 7.3]). *Under the specialization $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}}) \rightarrow K(\mathcal{C}_{\mathfrak{g}})$, the element $L_t(m)$ corresponds to the class $[L(m)]$ for any $m \in \mathcal{M}$. In particular, we have $P_{m, m'}(1) = P_{m, m'}$ for any $m, m' \in \mathcal{M}$.*

Conjecture 1.2 (Positivities in the quantum Grothendieck ring).

- (P1) *The polynomial $P_{m, m'}(t)$ has non-negative coefficients, i.e., $P_{m, m'}(t) \in t\mathbb{Z}_{\geq 0}[t]$.*
- (P2) *The structure constants of $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ with respect to the canonical basis $\{L_t(m)\}_{m \in \mathcal{M}}$ have non-negative coefficients.*
- (P3) *In the quantum torus \mathcal{Y}_t , the elements $L_t(m)$ have non-negative coefficients.*

Motivated by Conjecture 1.1, we refer to the element $L_t(m)$ as *the (q, t) -character of the simple module $L(m)$* . Since they can be computed algorithmically as in the usual Kazhdan-Lusztig theory, Conjecture 1.1 enables us to compute all the simple q -characters algorithmically once it is verified. Conjecture 1.2 (P1) was checked in several small examples and expected generally in [18]. As mentioned before, when \mathfrak{g} is simply-laced, all of these conjectures are known to be true by [47, 55] using the geometry of quiver varieties. On the other hand, they are still unsolved for non-simply-laced \mathfrak{g} .

In this paper, using the new ingredients that we explain below, we improve this situation with the following results.

Theorem 1.3 (= Corollary 10.7 + Corollary 11.4 + Theorem 11.7).

- (1) *For general \mathfrak{g} , Conjecture 1.2 (P1) and (P2) hold.*
- (2) *If \mathfrak{g} is of type B, Conjecture 1.1 and Conjecture 1.2 (P3) hold as well.*

Our proof of Theorem 1.3 relies in part on the isomorphisms established in this paper between quantum Grothendieck rings for non-simply-laced quantum loop algebras and their unfolded simply-laced ones.

Note that in their recent study of the non-symmetric quantized Coulomb branches [48, 44], Nakajima and Weekes established a parametrization of simple modules in non-simply-laced types in terms of simple modules in simply-laced types and proved that q -characters of simple modules of non-symmetric Kac-Moody algebras are obtained from those of unfolded symmetric ones by imposing a certain modulus condition. In particular, they solved the character problem for simple modules by a different approach. Although they also used a connection between non-symmetric and symmetric quantum affine algebras, it seems to be different from the one we used in this paper.

1.4. In what follows, we outline our proof of the above main results. First of all, we notice that the description of the (quantum) Grothendieck ring of the whole category $\mathcal{C}_{\mathfrak{g}}$ can be reduced to that of a certain rigid monoidal subcategory $\mathcal{C}_{\mathbb{Z},\mathfrak{g}} \subset \mathcal{C}_{\mathfrak{g}}$ introduced in [23] (see Section 2.6). Roughly speaking, $\mathcal{C}_{\mathbb{Z},\mathfrak{g}}$ is a Serre subcategory of $\mathcal{C}_{\mathfrak{g}}$ generated by the simple modules whose Drinfel'd polynomials have their roots at some integral powers of q only. From now on, we focus on this skeleton subcategory $\mathcal{C}_{\mathbb{Z},\mathfrak{g}}$.¹

Now we assume that \mathfrak{g} is of non-simply-laced type. Then we choose another simple Lie algebra \mathfrak{g} of simply-laced type whose Dynkin diagram is obtained by *unfolding* the Dynkin diagram of \mathfrak{g} . In other words, we fix a pair $(\mathfrak{g}, \mathfrak{g})$ of simple Lie algebras whose Dynkin types are either (B_n, A_{2n-1}) , (C_n, D_{n+1}) , (F_4, E_6) or (G_2, D_4) . To prove the positivities (P1) and (P2) for non-simply-laced \mathfrak{g} , we establish the following.

Theorem 1.4 (\doteq Theorem 10.6). *There exists an isomorphism of $\mathbb{Z}[t^{\pm 1/2}]$ -algebras*

$$(1.1) \quad \mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}}) \simeq \mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$$

which extends an isomorphism (1.3) explained below and which induces a bijection between the (q, t) -characters of simple modules.

Once we establish an isomorphism (1.1), it propagates the known positivity in $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ to the desired positivity in $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})$. We note that such an isomorphism is not unique. In fact, we construct a collection of such isomorphisms (1.1) labeled by certain combinatorial objects called *Q-data*, which we explain later in 1.6 (they lead to the same positivity results). The isomorphisms (1.3) concern subcategories and already depend on the choice of Q-data (see Theorem 1.5 below). Also we note that our isomorphisms are not deduced from the results in [48, 44].

The specialization at $t = 1$ of the isomorphism (1.1) yields an isomorphism between the usual Grothendieck rings

$$K(\mathcal{C}_{\mathbb{Z},\mathfrak{g}}) \simeq K(\mathcal{C}_{\mathfrak{g}}).$$

We note that it is still non-trivial. For example, it does not respect the classes of fundamental modules nor of standard modules. We conjecture that it induces a bijection between the classes of simple modules. In this paper, we prove this conjecture when \mathfrak{g}

¹In the main body of this paper, we only consider the quantum Grothendieck ring of the subcategory $\mathcal{C}_{\mathbb{Z},\mathfrak{g}} \subset \mathcal{C}_{\mathfrak{g}}$ from the beginning.

is of type B with a specific choice of Q-data. Then Conjecture 1.1 for type B is obtained as a corollary. See 1.10.

1.5. Our main step to construct an isomorphism (1.1) is to compare a presentation of the *localized* quantum Grothendieck ring²

$$\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})_{loc} := \mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}}) \otimes_{\mathbb{Z}[t^{\pm 1/2}]} \mathbb{Q}(t^{1/2})$$

of $\mathcal{C}_{\mathbb{Z},\mathfrak{g}}$ with that of $\mathcal{C}_{\mathbb{Z},\mathfrak{g}}$.

A presentation of $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})_{loc}$ for simply-laced \mathfrak{g} was previously obtained by B. Leclerc and the second named author in [24], which revealed that $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})_{loc}$ is isomorphic to the derived Hall algebra of the category $\text{Rep } Q$ of representations of a Dynkin quiver Q of the same type as \mathfrak{g} . Under this isomorphism, the cohomological degree shift [1] in the derived category $D^b(\text{Rep } Q)$ corresponds to the left dual functor \mathcal{D} in the rigid monoidal category $\mathcal{C}_{\mathbb{Z},\mathfrak{g}}$. If we restrict it to the subalgebras corresponding to the heart $\text{Rep } Q$ of the standard t -structure in $D^b(\text{Rep } Q)$, we get an isomorphism

$$(1.2) \quad \mathcal{K}_t(\mathcal{C}_Q) \simeq \mathcal{A}_t[N_-],$$

where $\mathcal{A}_t[N_-]$ is the quantum unipotent coordinate algebra associated with \mathfrak{g} , and \mathcal{C}_Q is a certain monoidal subcategory of $\mathcal{C}_{\mathbb{Z},\mathfrak{g}}$ defined by using the Auslander-Reiten (AR) quiver Γ_Q of $\text{Rep } Q$. Moreover, the isomorphism (1.2) sends the basis of $\mathcal{K}_t(\mathcal{C}_Q)$ formed by the simple (q, t) -characters to the dual canonical basis of $\mathcal{A}_t[N_-]$.

Roughly speaking, the proof in [24] consists of the following two steps:

- (i) establish the isomorphism (1.2),
- (ii) investigate the relations among the subalgebras $\mathcal{K}_t(\mathcal{D}^k \mathcal{C}_Q) \subset \mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})$ for $k \in \mathbb{Z}$.

Note that the subcategory $\mathcal{D}^k \mathcal{C}_Q \subset \mathcal{C}_{\mathbb{Z},\mathfrak{g}}$ corresponds to the shifted heart $(\text{Rep } Q)[k]$ in $D^b(\text{Rep } Q)$ and in particular we have $\mathcal{K}_t(\mathcal{D}^k \mathcal{C}_Q) \simeq \mathcal{K}_t(\mathcal{C}_Q)$ as $\mathbb{Z}[t^{\pm 1/2}]$ -algebras. As a result, we find two kinds of relations: *the quantum Serre relations* from (i) and *the quantum Boson relations* from (ii), which yields the desired presentation of $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})_{loc}$.

In this paper, we generalize the above story in a unified way including non-simply-laced cases. As a result, we find that the localized Grothendieck rings $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})_{loc}$ and $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})_{loc}$ share the same presentation, which is finally enhanced to the desired isomorphism (1.1). Our proof also consists of two steps analogous to (i) and (ii) above.

1.6. To generalize the step (i), we need to extend the definition of the subcategory \mathcal{C}_Q to the general case. Such a generalization was considered by the third named author and his collaborators in [38, 53, 51] using the *twisted AR quiver* Γ_Q associated with a Q -datum \mathcal{Q} for \mathfrak{g} , instead of the AR quiver Γ_Q associated with a Dynkin quiver Q in the last paragraph. The twisted AR quiver is a quiver encoding some combinatorial information of a certain commutation class in the Weyl group of \mathfrak{g} (not \mathfrak{g}) called a *twisted adapted class*. It was originally introduced by U. R. Suh and the third named author in [53]. The notion of a Q-datum is a combinatorial generalization of a Dynkin quiver (with a height function) introduced by the first and the third named authors in [13] to give a unified description of these twisted AR quivers. With these notions, we

²In the main body of the paper, it is denoted by $\mathcal{K}_{\mathbb{Q}(t^{1/2})}$.

can define the monoidal subcategory $\mathcal{C}_{\mathcal{Q}} \subset \mathcal{C}_{\mathbb{Z}, \mathfrak{g}}$ associated with each Q-datum \mathcal{Q} for \mathfrak{g} in a suitable way.

As a generalization of the isomorphism (1.2), we prove the following.

Theorem 1.5 (\doteq Corollary 8.7). *For each Q-datum \mathcal{Q} for \mathfrak{g} , there exists an isomorphism of $\mathbb{Z}[t^{\pm 1/2}]$ -algebras*

$$(1.3) \quad \mathcal{K}_t(\mathcal{C}_{\mathcal{Q}}) \simeq \mathcal{A}_t[N_-],$$

which sends the basis of $\mathcal{K}_t(\mathcal{C}_{\mathcal{Q}})$ formed by the simple (q, t) -characters to the dual canonical basis of $\mathcal{A}_t[N_-]$, and the basis formed by the standard (q, t) -characters to the normalized dual PBW-type basis associated with a certain reduced word corresponding to \mathcal{Q} .

It is notable that the quantum unipotent coordinate algebra $\mathcal{A}_t[N_-]$ here is associated with the simply-laced \mathfrak{g} (not \mathfrak{g}). When \mathfrak{g} is of type B, our isomorphism (1.3) coincides with the isomorphism established in the previous work [26] of the second and the fourth named authors. Hence we refer to the isomorphism (1.3) as *the HLO isomorphism* in this paper. Our proof of Theorem 1.5 generalizes the arguments in [26] with new ingredients, in particular the combinatorial theory of Q-data. The essence of the proof is a comparison between the determinantal identities satisfied by the normalized quantum unipotent minors in $\mathcal{A}_t[N_-]$ (which can be understood as a part of its quantum cluster algebra structure) and *the quantum T-systems* in $\mathcal{K}_t(\mathcal{C}_{\mathcal{Q}})$, where the latter is a newly obtained result in this paper except for types ABDE as explained in the next paragraph.

1.7. Recall that the T -systems are certain functional relations appearing in solvable lattice models (see [40] for instance). They were proved to be satisfied by the q -characters of *Kirillov-Reshetikhin modules* (KR modules for short) by Nakajima [46] for simply-laced \mathfrak{g} and by the second named author [20] for general \mathfrak{g} . For types ADE and for type B, their natural t -analogs, which we call the quantum T -systems, were established in [24] and [26] respectively, as the relations satisfied in the quantum Grothendieck ring $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$. In this paper, we prove the quantum T -systems for all \mathfrak{g} in a unified way. Strictly speaking, we establish them as a system of relations satisfied by another kind of elements $F_t(m)$ of $\mathcal{K}_t(\mathcal{C}_{\mathfrak{g}})$ which contain m corresponding to KR modules as their unique dominant monomials (see Section 6.3). Although it is not clear in our formulation (as well as that of [26] for type B) that the quantum T -systems are satisfied by the simple (q, t) -characters corresponding to KR modules, we conjecture that it is true (indeed Conjectures 1.1 and 1.2 imply $L_t(m) = F_t(m)$ for any m corresponding to KR modules).

1.8. The HLO isomorphism (1.3) itself has important corollaries. Since it translates the known positivities of the dual canonical basis of $\mathcal{A}_t[N_-]$ to the desired positivities in $\mathcal{K}_t(\mathcal{C}_{\mathcal{Q}})$, we find that (P1) and (P2) in Conjecture 1.2 hold when we restrict ourselves to the category $\mathcal{C}_{\mathcal{Q}}$ for some Q-datum \mathcal{Q} . Moreover, we can check that the specialization at $t = 1$ of the HLO isomorphism coincides with the isomorphism $K(\mathcal{C}_{\mathcal{Q}}) \simeq \mathcal{A}_{t=1}[N_-]$ induced from *the generalized quantum affine Schur-Weyl duality functors* studied in [27, 30, 38, 51, 11, 49]. It is known that the latter isomorphism sends the basis of $K(\mathcal{C}_{\mathcal{Q}})$

formed by the classes of simple modules to the specialization of the dual canonical basis of $\mathcal{A}_{t=1}[N_-]$. Thus it implies the following.

Theorem 1.6 (\doteq Corollary 9.8). *The analog of Kazhdan-Lusztig conjecture, i.e., Conjecture 1.1 holds for the category $\mathcal{C}_{\mathcal{Q}}$ for any \mathcal{Q} .*

1.9. Next we consider the analog of step (ii) for general \mathfrak{g} . To prove that the desired quantum Bosc relations are satisfied in $\mathcal{K}_t(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})$, we have to check several commutation relations between the (q, t) -characters of fundamental modules. For this purpose, we need to know the singularities of *the normalized R -matrices* between the fundamental modules. Fortunately, the denominators of these R -matrices are already known due to many works [1, 8, 12, 27, 50, 51]. Moreover, their relation to the structure constants of the quantum torus \mathcal{Y}_t was observed by the first and the third named authors in [13]. Combining these ingredients with the positivities of $\mathcal{K}_t(\mathcal{C}_{\mathcal{Q}})$ (discussed in the last paragraph), we can obtain the conclusion.

1.10. Finally, we specialize the situation to the case when $(\mathfrak{g}, \mathfrak{g})$ is of type (B_n, A_{2n-1}) . In this case, we can check that the specialization at $t = 1$ of our isomorphism (1.1) with a specific choice of \mathcal{Q} -data coincides with another isomorphism $K(\mathcal{C}_{\mathbb{Z},\mathfrak{g}}) \simeq K(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})$ obtained by M. Kashiwara, M. Kim and the third named author in [34] through a categorical relation arising from another example of the generalized quantum affine Schur-Weyl duality functors. The computation we need in the proof has already been done in the previous work [26, Section 12] by the second and the fourth named authors. Since the latter isomorphism $K(\mathcal{C}_{\mathbb{Z},\mathfrak{g}}) \simeq K(\mathcal{C}_{\mathbb{Z},\mathfrak{g}})$ respects the classes of simple modules, we immediately obtain the proof of Conjecture 1.1 for this case.

1.11. As mentioned above, in opposition to the case of simply-laced quantum affine algebras, as far as the authors know, no geometric construction of (co)standard modules involving non-simply-laced analog of quiver varieties is known or fully developed. Note however that, in addition to the recent results of Nakajima-Weekes mentioned above, geometric q -characters formulas of certain modules, including KR modules, have been obtained in [25] for non-simply-laced types. Also an approach to certain analogs of non simply-laced quiver varieties have been developed in [15]. This raises the question of a geometric interpretation of the positivity results we establish in the present paper, in terms of quiver varieties associated to non-simply-laced quantum affine algebras.

1.12. The authors plan to come back in the near future to the following related questions. The approach in [26] was based on the study of quantum cluster algebra structures. We will study the compatibility between the isomorphisms of quantum Grothendieck rings that we establish and the various involved quantum cluster algebra structures (see [54, 3]). We expect this will also provide new interesting information on related q -characters and their t -analogs, in the spirit of [25, 3]. We will also investigate the relation between on one hand the specialization of our isomorphisms to classical Grothendieck ring isomorphisms, and on the other hand the various isomorphisms obtained via the framework of the generalized quantum affine Schur-Weyl duality [28]. Eventually, we expect our approach and results to hold for integrable representations of general quantum affinizations of Kac-Moody algebras and for representations in the category \mathcal{O} of a Borel subalgebra of a quantum affine algebra [22].

Organization. This paper is organized as follows. In Section 2, we recall the definition of quantum loop algebras and some important properties of their finite-dimensional representations. In Section 3, we review the definition and some properties of the quantum Grothendieck ring of the category $\mathcal{C}_{\mathbb{Z}, \mathfrak{g}}$. Section 4 is a recollection on the combinatorial theory related to Q-data and their twisted AR quivers. In Section 5, we see several applications of the theory of Q-data to the representation theory of the quantum loop algebras. In particular, the definition of the monoidal subcategory $\mathcal{C}_{\mathcal{Q}}$ is recalled. In Section 6, after reviewing the classical T -systems in term of Q-data, we establish the quantum T -systems for all \mathfrak{g} . Section 7 is a brief review of the quantum unipotent coordinate algebra $\mathcal{A}_t[N_-]$ and its quantum cluster algebra structure. In Section 8, we prove the HLO isomorphism (1.3) for each Q-datum and obtain its corollaries. In Section 9, we deduce several commutation relations in the quantum Grothendieck ring which play an important role in the proof of our main theorem. In Section 10, we prove the desired presentation of the localized Grothendieck ring and then construct a collection of isomorphisms among the quantum Grothendieck rings. In particular, we obtain an isomorphism (1.1) and hence the desired positivities (P1) and (P2) for all \mathfrak{g} . In the last Section 11, we focus on the case when \mathfrak{g} is of type B to prove Conjecture 1.1 and also the remaining positivity (P3).

Acknowledgments. The authors are grateful to Hiraku Nakajima for helpful discussions, comments and correspondence. The authors would like also to thank Bernard Leclerc and Alex Weekes for their comments. The authors are also thankful to the anonymous referee whose suggestions improve our present paper. A part of this work was done when the first named author enjoyed short-term postdoctoral positions in Kyoto University and in Kavli IPMU during the spring-summer semester in 2020. He thanks both institutions and colleagues for their supports in the complicated situation.

R. F. was supported in part by JSPS Grant-in-Aid for Scientific Research (A) JP17H01086, (B) JP19H01782 and by JSPS Overseas Research Fellowships. D. H. was supported by the European Research Council under the European Union's Framework Programme H2020 with ERC Grant Agreement number 647353 Qaffine. S.-j. Oh was supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2019R1A2C4069647). H. O. is supported by Grant-in-Aid for Young Scientists No.19K14515.

Convention. Throughout this paper, we use the following general convention.

- For a statement P, we set $\delta(\mathbf{P})$ to be 1 or 0 according that P is true or not. As a special case, we use the notation $\delta_{i,j} := \delta(i = j)$ (Kronecker's delta).
- For a totally ordered set $J = \{\cdots < j_{-1} < j_0 < j_1 < j_2 < \cdots\}$, write

$$\prod_{j \in J}^{\rightarrow} A_j := \cdots A_{j_{-1}} A_{j_0} A_{j_1} A_{j_2} \cdots, \quad \bigotimes_{j \in J}^{\rightarrow} A_j := \cdots A_{j_{-1}} \otimes A_{j_0} \otimes A_{j_1} \otimes A_{j_2} \cdots.$$

- For an abelian category \mathcal{C} , we denote its Grothendieck group by $K(\mathcal{C})$. The class of an object $X \in \mathcal{C}$ is denoted by $[X] \in K(\mathcal{C})$. Write the subset of $K(\mathcal{C})$ formed by the classes of simple objects as $\text{Irr } \mathcal{C}$. If moreover the category \mathcal{C}

is monoidal with a bi-exact tensor product \otimes , $K(\mathcal{C})$ is endowed with a ring structure by $[X] \cdot [Y] = [X \otimes Y]$, which we call the Grothendieck ring of \mathcal{C} .

- Let \mathbb{k} be a field. For a polynomial $f(z) \in \mathbb{k}[z]$, we denote its degree by $\deg(f)$. For a rational function $f(z) \in \mathbb{k}(z)$, we denote by $[f(z)]^\pm$ its formal Laurent expansion at $z^{\pm 1} = 0$, which are elements in $\mathbb{k}((z))$ and $\mathbb{k}((z^{-1}))$, respectively.

2. QUANTUM LOOP ALGEBRAS AND THEIR FINITE-DIMENSIONAL REPRESENTATIONS

In this section, we recall the definition of quantum loop algebras and important properties of their finite-dimensional representations.

2.1. Notation. Let \mathfrak{g} be a complex finite-dimensional simple Lie algebra of rank n . As is well-known, it is classified by the Dynkin diagrams of types A_n, B_n, \dots, G_2 . Denote by r the lacing number of \mathfrak{g} , that is

$$r := \begin{cases} 1 & \text{if } \mathfrak{g} \text{ is of type } A_n, D_n, \text{ or } E_{6,7,8}, \\ 2 & \text{if } \mathfrak{g} \text{ is of type } B_n, C_n, \text{ or } F_4, \\ 3 & \text{if } \mathfrak{g} \text{ is of type } G_2. \end{cases}$$

We say that \mathfrak{g} is simply-laced (resp. non-simply-laced) if $r = 1$ (resp. $r > 1$). Let $I = \{1, 2, \dots, n\}$ be the set of Dynkin indices and $C = (c_{ij})_{i,j \in I}$ the Cartan matrix of \mathfrak{g} . We write $i \sim j$ for $i, j \in I$ if $c_{ij} < 0$. Let $D = \text{diag}(d_i \mid i \in I)$ denote the unique diagonal matrix such that $d_i \in \{1, r\}$ for each $i \in I$ and $DC = (d_i c_{ij})_{i,j \in I}$ is symmetric.

2.2. Quantum loop algebras. In what follows, for an invertible element $x \neq \pm 1$ in an integral domain and an integer k , we write the standard quantum number

$$(2.1) \quad [k]_x := \frac{x^k - x^{-k}}{x - x^{-1}}.$$

For integers $k \geq l \geq 0$, we set

$$[k]_x! := \prod_{u=1}^k [u]_x, \quad \begin{bmatrix} k \\ l \end{bmatrix}_x := \frac{[k]_x!}{[k-l]_x! [l]_x!}.$$

Let q be an indeterminate (quantum parameter) and \mathbb{k} the algebraic closure of the rational function field $\mathbb{Q}(q)$. For each $i \in I$, we define $q_i := q^{d_i} \in \mathbb{k}$.

Definition 2.1. The quantum loop algebra $U_q(L\mathfrak{g})$ is the \mathbb{k} -algebra given by the set of generators

$$\{k_i^{\pm 1} \mid i \in I\} \cup \{x_{i,k}^\pm \mid i \in I, k \in \mathbb{Z}\} \cup \{h_{i,l} \mid i \in I, l \in \mathbb{Z} \setminus \{0\}\}$$

satisfying the following relations:

- $k_i k_i^{-1} = k_i^{-1} k_i = 1, k_i k_j = k_j k_i$ for $i, j \in I$,
- $k_i x_{j,k}^\pm k_i^{-1} = q_i^{\pm c_{ij}} x_{j,k}^\pm$ for $i, j \in I$ and $k \in \mathbb{Z}$,
- $[k_i, h_{j,l}] = [h_{i,l}, h_{j,m}] = 0$ for $i, j \in I$ and $l, m \in \mathbb{Z} \setminus \{0\}$,
- $[x_i^+(z), x_j^-(w)] = \frac{\delta_{ij}}{q_i - q_i^{-1}} (\delta(z/w) \phi_i^+(w) - \delta(w/z) \phi_i^-(z))$ for $i, j \in I$,
- $(q_i^{\pm c_{ij}} z - w) x_i^\pm(z) x_j^\pm(w) = (z - q_i^{\pm c_{ij}} w) x_j^\pm(w) x_i^\pm(z)$ for $i, j \in I$,

- $(q_i^{\pm c_{ij}} z - w) \phi_i^\varepsilon(z) x_j^\pm(w) = (z - q_i^{\pm c_{ij}} w) x_j^\pm(w) \phi_i^\varepsilon(z)$ for $i, j \in I$ and $\varepsilon \in \{+, -\}$,
- $\sum_{g \in \mathfrak{S}_b} \sum_{k=0}^b (-1)^k \begin{bmatrix} b \\ k \end{bmatrix}_{q_i} x_i^\pm(z_{g(1)}) \cdots x_i^\pm(z_{g(k)}) x_j^\pm(w) x_i^\pm(z_{g(k+1)}) \cdots x_i^\pm(z_{g(b)}) = 0$, where $b := 1 - c_{ij}$, for $i, j \in I$ with $i \sim j$,

where $\delta(z), x_i^\pm(z), \phi_i^\pm(z)$ are the formal power series defined by

$$\delta(z) := \sum_{l=-\infty}^{\infty} z^l, \quad x_i^\pm(z) := \sum_{l=-\infty}^{\infty} x_{i,k}^\pm z^k, \quad \phi_i^\pm(z) := k_i^{\pm 1} \exp\left(\pm(q_i - q_i^{-1}) \sum_{l=1}^{\infty} h_{i,\pm l} z^{\pm l}\right).$$

The quantum loop algebra $U_q(L\mathfrak{g})$ is a sub-quotient of the corresponding (untwisted) quantum affine algebra and hence has a Hopf algebra structure, which is defined via another realization of $U_q(L\mathfrak{g})$ (called the Drinfeld-Jimbo presentation).

2.3. Finite-dimensional representations. A finite-dimensional $U_q(L\mathfrak{g})$ -module is said to be of type **1** if the element k_i acts as a diagonal \mathbb{k} -linear operator whose eigenvalues belong to the set $\{q^k \mid k \in \mathbb{Z}\}$ for each $i \in I$. It is well-known that the study of finite-dimensional representations of $U_q(L\mathfrak{g})$ reduces essentially to the study of the category \mathcal{C} of type **1** finite-dimensional $U_q(L\mathfrak{g})$ -modules. This is a \mathbb{k} -linear rigid monoidal tensor category. As a monoidal category, the category \mathcal{C} is not braided. Indeed, $V \otimes W$ is not isomorphic to $W \otimes V$ for general $V, W \in \mathcal{C}$. We say that V and W *commute* if $V \otimes W \cong W \otimes V$ as $U_q(L\mathfrak{g})$ -modules.

Let $V \in \mathcal{C}$. Since the elements $\{k_i^{\pm 1}, h_{i,l} \mid i \in I, l \in \mathbb{Z} \setminus \{0\}\}$ mutually commute, we have a decomposition

$$V = \bigoplus_{\gamma \in (\mathbb{k}[[z]] \times \mathbb{k}[[z^{-1}]])^I} V_\gamma$$

where, for each $\gamma = (\gamma_i^+(z), \gamma_i^-(z))_{i \in I}$, we define V_γ as the subspace of V on which each coefficient of the series $\phi_i^\pm(z) - \gamma_i^\pm(z)$ acts nilpotently for any $i \in I$. If $V_\gamma \neq 0$, it is called an ℓ -weight space and γ is called the corresponding ℓ -weight.

For a simple module L in \mathcal{C} , there uniquely exists an ℓ -weight space L_{γ_0} such that $x_{i,k}^+ L_{\gamma_0} = 0$ for all $i \in I$ and $k \in \mathbb{Z}$. The isomorphism class of a simple $U_q(L\mathfrak{g})$ -module in \mathcal{C} is determined by such γ_0 (called the ℓ -highest weight of L). Moreover we have the following.

Theorem 2.2 ([6, Theorem 3.3], [5, Theorem 12.2.6]). *If $\gamma = (\gamma_i^+(z), \gamma_i^-(z))_{i \in I}$ is an ℓ -highest weight of a simple $U_q(L\mathfrak{g})$ -module in \mathcal{C} , for each $i \in I$ there exists a unique polynomial $P_i(z) \in \mathbb{k}[z]$ with $P_i(0) = 1$ for $i \in I$ such that*

$$(2.2) \quad \gamma_i^\pm(z) = q_i^{\deg(P_i)} \left[\frac{P_i(zq_i^{-1})}{P_i(zq_i)} \right]^\pm.$$

Conversely, for any $(P_i(z))_{i \in I} \in (1 + z\mathbb{k}[[z]])^I$, we have a simple $U_q(L\mathfrak{g})$ -module in \mathcal{C} whose ℓ -highest weight γ is given by (2.2).

This I -tuple of polynomials $(P_i(z))_{i \in I}$ corresponding to a simple module L is called the *Drinfeld polynomials of L* . A non-zero vector in the ℓ -highest weight space L_γ is called an ℓ -highest weight vector of L . Since $\dim_{\mathbb{k}} L_\gamma = 1$, it is unique up to a scalar.

2.4. q -characters. Frenkel and Reshetikhin [10, Proposition 1] proved that any ℓ -weight γ of a $U_q(L\mathfrak{g})$ -module in \mathcal{C} is of the form

$$\gamma_i^\pm(z) = q_i^{\deg(Q_i) - \deg(R_i)} \left[\frac{Q_i(zq_i^{-1})R_i(zq_i)}{Q_i(zq_i)R_i(zq_i^{-1})} \right]^\pm$$

for some $(Q_i(z))_{i \in I}, (R_i(z))_{i \in I} \in (1 + z\mathbb{k}[z])^I$. For this γ , define a monomial m_γ in the Laurent polynomial ring $\mathbb{Z}[Y_{i,a}^{\pm 1} \mid i \in I, a \in \mathbb{k}^\times]$ by

$$m_\gamma := \prod_{i \in I, a \in \mathbb{k}^\times} Y_{i,a}^{q_{i,a} - r_{i,a}},$$

where $Q_i(z) = \prod_{a \in \mathbb{k}^\times} (1 - az)^{q_{i,a}}, R_i(z) = \prod_{a \in \mathbb{k}^\times} (1 - az)^{r_{i,a}}$.

Theorem 2.3 ([10, Theorem 3]). *The assignment*

$$[V] \mapsto \sum_{\gamma} \dim_{\mathbb{k}}(V_{\gamma}) m_{\gamma}$$

defines an injective algebra homomorphism $\chi_q: K(\mathcal{C}) \rightarrow \mathbb{Z}[Y_{i,a}^{\pm 1} \mid i \in I, a \in \mathbb{k}^\times]$.

This implies that the Grothendieck ring $K(\mathcal{C})$ is commutative, although the category \mathcal{C} is not braided. The map χ_q is called *the q -character homomorphism*. A monomial which is a product of only positive powers of $Y_{i,a}$'s is said to be *dominant*. Write the simple $U_q(L\mathfrak{g})$ -module in \mathcal{C} with ℓ -highest weight γ as $L(m_\gamma)$. Then, by Theorem 2.2, m_γ is dominant, and the set of dominant monomials are in bijection with $\text{Irr } \mathcal{C}$ via $m \mapsto [L(m)]$.

Following [10], for $i \in I, a \in \mathbb{k}^\times$, we set

$$A_{i,a} := Y_{i,aq_i^{-1}} Y_{i,aq_i} \left(\prod_{j: c_{ji} = -1} Y_{j,a}^{-1} \right) \left(\prod_{j: c_{ji} = -2} Y_{j,aq_i^{-1}}^{-1} Y_{j,aq_i}^{-1} \right) \left(\prod_{j: c_{ji} = -3} Y_{j,aq_i^{-2}}^{-1} Y_{j,a}^{-1} Y_{j,aq_i^2}^{-1} \right).$$

As these monomials are algebraically independent, one can define the Nakajima partial ordering \leq on the set of monomials in $\mathbb{Z}[Y_{i,a}^{\pm 1} \mid i \in I, a \in \mathbb{k}^\times]$ as follows:

$m \leq m'$ if and only if $m^{-1}m'$ is a product of elements of $\{A_{i,a} \mid i \in I, a \in \mathbb{k}^\times\}$.

Theorem 2.4 ([10, 9]). *Let m be a dominant monomial. Then all the monomials occurring in $\chi_q(L(m)) - m$ are strictly less than m with respect to the ordering \leq .*

2.5. Twists and duality. For each $a \in \mathbb{k}^\times$, we consider the \mathbb{k} -algebra automorphism τ_a of the quantum loop algebra $U_q(L\mathfrak{g})$ given by

$$\tau_a(k_i) = k_i, \quad \tau_a(x_{i,k}^\pm) = a^k x_{i,k}^\pm, \quad \tau_a(h_{i,l}) = a^l h_{i,l},$$

for $i \in I, k \in \mathbb{Z}, l \in \mathbb{Z} \setminus \{0\}$. This automorphism τ_a is a q -analog of the loop rotation $z \mapsto az$ for the loop algebra $L\mathfrak{g} = \mathfrak{g} \otimes_{\mathbb{C}} \mathbb{C}[z^\pm]$. For a finite-dimensional representation V in \mathcal{C} , the twisted representation $\tau_a^* V$ by τ_a again belongs to \mathcal{C} . Thus the assignment $V \mapsto \tau_a^* V$ defines a monoidal auto-equivalence T_a of \mathcal{C} . The functors $\{T_a\}_{a \in \mathbb{k}^\times}$ define an action of the group \mathbb{k}^\times on the monoidal category \mathcal{C} as they satisfy $T_a \circ T_b \simeq T_{ab}$ and $T_1 = \text{id}_{\mathcal{C}}$. If we denote by $y \mapsto y_a$ the \mathbb{Z} -algebra automorphism of $\mathbb{Z}[Y_{i,b}^{\pm 1} \mid i \in I, b \in \mathbb{k}^\times]$ given by $Y_{i,b} \mapsto Y_{i,ab}$, we have $T_a(L(m)) \cong L(m_a)$ for any simple module $L(m) \in \mathcal{C}$.

Recall that the monoidal category \mathcal{C} is rigid, i.e. every object $V \in \mathcal{C}$ has its left dual $\mathcal{D}(V)$ and right dual $\mathcal{D}^{-1}(V)$ in \mathcal{C} .³ Since $\mathcal{D} \circ \mathcal{D}^{-1} \simeq \mathcal{D}^{-1} \circ \mathcal{D} \simeq \text{id}_{\mathcal{C}}$, we can naturally extend $\mathcal{D}^{\pm 1}$ to a collection of functors $\{\mathcal{D}^k\}_{k \in \mathbb{Z}}$ so that we have $\mathcal{D}^k \circ \mathcal{D}^l \simeq \mathcal{D}^{k+l}$ for $k, l \in \mathbb{Z}$. Then we have $\mathcal{D}^2 \simeq T_{q^{2rh^\vee}}$ as endo-functors on \mathcal{C} , where h^\vee is the dual Coxeter number of \mathfrak{g} . See Table 1 in Section 4 below for its explicit value.

Now we consider the \mathbb{Z} -algebra automorphism \mathfrak{D} of $\mathbb{Z}[Y_{i,a}^{\pm 1} \mid i \in I, a \in \mathbb{k}^\times]$ given by

$$(2.3) \quad \mathfrak{D}(Y_{i,a}) := Y_{i^*, aq^{rh^\vee}} \quad \text{for each } i \in I, a \in \mathbb{k}^\times,$$

where $i \mapsto i^*$ denotes the involution of the set I induced from the longest element of the Weyl group of \mathfrak{g} . See Section 4.1 for its precise definition when \mathfrak{g} is simply-laced, while it is trivial when \mathfrak{g} is non-simply-laced.

Theorem 2.5 (cf. [7, Proposition 5.1(b)], [1, Appendix A], [9, Corollary 6.10]). *For any simple object $L(m) \in \mathcal{C}$ and $k \in \mathbb{Z}$, there exists an isomorphism*

$$\mathcal{D}^k(L(m)) \cong L(\mathfrak{D}^k(m))$$

of $U_q(L\mathfrak{g})$ -modules. In particular, we have

$$\chi_q(\mathcal{D}(L(m))) = \mathfrak{D}(\chi_q(L(m))) = \chi_q(L(\mathfrak{D}(m))).$$

Finally, we recall the involution ω of the quantum loop algebra $U_q(L\mathfrak{g})$ as in [4]: it is defined by

$$\omega(x_{i,k}^\pm) = -x_{i,-k}^\mp, \quad \omega(h_{i,l}) = -h_{i,-l}, \quad \omega(k_i) = k_i^{-1}$$

for $i \in I, k \in \mathbb{Z}, l \in \mathbb{Z} \setminus \{0\}$. Then, for a representation $V \in \mathcal{C}$, the q -character of the twisted representation ω^*V by ω is obtained from $\chi_q(V)$ by replacing each variable $Y_{i,a}$ by $\omega(Y_{i,a}) := Y_{i,a^{-1}}^{-1}$. In particular, the assignment $\chi_q(V) \mapsto \chi_q(\omega^*V) = \omega(\chi_q(V))$ defines a ring automorphism of $\chi_q(K(\mathcal{C}))$. Moreover, if V is a simple module $L(m)$, then we have

$$(2.4) \quad \omega^*L(m) \cong L(\omega^*m),$$

where ω^*m is the dominant monomial obtained from m by replacing each variables $Y_{i,a}$ with $Y_{i^*, a^{-1}q^{-rh^\vee}}$ (see [21]).

2.6. The category $\mathcal{C}_{\mathbb{Z}}$. In this subsection, we introduce a certain monoidal subcategory $\mathcal{C}_{\mathbb{Z}}$ of \mathcal{C} , which is of our main interest in this paper. For $i, j \in I$, write

$$d_{ij} := \min(d_i, d_j).$$

A function $\epsilon: I \rightarrow \{0, 1\}$ is called a *parity function* for \mathfrak{g} if it satisfies the condition

$$(2.5) \quad \epsilon_i \equiv \epsilon_j + d_{ij} \pmod{2} \quad \text{for any } i, j \in I \text{ with } i \sim j.$$

In what follows, we choose and fix a parity function ϵ for \mathfrak{g} . Then we define

$$(2.6) \quad \widehat{I} := \{(i, p) \in I \times \mathbb{Z} \mid p - \epsilon_i \in 2\mathbb{Z}\}.$$

Let \mathcal{Y} be the subalgebra of $\mathbb{Z}[Y_{i,a}^{\pm 1} \mid (i, a) \in I \times \mathbb{k}^\times]$ consisting of Laurent polynomials in the variables Y_{i,q^p} for $(i, p) \in \widehat{I}$. Note that we have $A_{i,q^{p+d_i}} \in \mathcal{Y}$ if and only if $(i, p) \in \widehat{I}$.

³By convention, we choose the coproduct of $U_q(L\mathfrak{g})$ opposite to that of [1, 33], that is why left and right duals are interchanged in comparison to these papers.

We denote by \mathcal{M} the set of all the dominant monomials in \mathcal{Y} . Following [23, Section 3.7], we consider the following category.

Definition 2.6. We define the category $\mathcal{C}_{\mathbb{Z}}$ as the Serre subcategory of \mathcal{C} which satisfies $\text{Irr } \mathcal{C}_{\mathbb{Z}} = \{[L(m)] \mid m \in \mathcal{M}\}$.

Proposition 2.7. *The category $\mathcal{C}_{\mathbb{Z}}$ is closed under taking tensor products and applying the duality functors $\mathcal{D}^{\pm 1}$. Moreover, the q -character homomorphism restricts to the injective algebra homomorphism $\chi_q: K(\mathcal{C}_{\mathbb{Z}}) \rightarrow \mathcal{Y}$.*

The subcategory $\mathcal{C}_{\mathbb{Z}}$ can be regarded as a skeleton of the entire rigid monoidal category \mathcal{C} . Indeed, the description of simple modules in \mathcal{C} and hence that of the Grothendieck $K(\mathcal{C})$ are essentially reduced to the description of the simple modules in $\mathcal{C}_{\mathbb{Z}}$ and that of the Grothendieck ring $K(\mathcal{C}_{\mathbb{Z}})$ thanks to the following fact.

Proposition 2.8. *Any simple representation V in \mathcal{C} can be factorized into a commutative tensor product of the form*

$$V \cong T_{a_1}(V_1) \otimes T_{a_2}(V_2) \otimes \cdots \otimes T_{a_d}(V_d)$$

with $V_k \in \mathcal{C}_{\mathbb{Z}}$ and $a_k \in \mathbb{k}^{\times}$ for $1 \leq k \leq d$ such that $a_k/a_l \notin q^{2\mathbb{Z}}$ for all $k \neq l$.

In this paper, we always work in the subcategory $\mathcal{C}_{\mathbb{Z}} \subset \mathcal{C}$. From now on, we put $Y_{i,p} := Y_{i,q^p}$ and $A_{i,p} := A_{i,q^p}$ for $(i,p) \in I \times \mathbb{Z}$ for the sake of simplicity. For example, the algebra automorphism $\mathfrak{D}: \mathcal{Y} \rightarrow \mathcal{Y}$ defined in (2.3) is given by $\mathfrak{D}(Y_{i,p}) = Y_{i^*,p+rh^{\vee}}$ in this notation. For a monomial m in $\mathcal{Y} = \mathbb{Z}[Y_{i,p}^{\pm 1} \mid (i,p) \in \widehat{I}]$, we write

$$(2.7) \quad m = \prod_{(i,p) \in \widehat{I}} Y_{i,p}^{u_{i,p}(m)}$$

with $u_{i,p}(m) \in \mathbb{Z}$.

The Grothendieck ring $K(\mathcal{C})$ is a polynomial ring in the classes of the *fundamental modules* $L(Y_{i,a})$, see [10]. This implies the following.

Proposition 2.9. *The \mathbb{Z} -algebra $K(\mathcal{C}_{\mathbb{Z}})$ is isomorphic to the polynomial ring with \widehat{I} -many variables $\mathbb{Z}[x_{i,p} \mid (i,p) \in \widehat{I}]$ via $[L(Y_{i,p})] \mapsto x_{i,p}$.*

The $U_q(L\mathfrak{g})$ -modules of the form $L(Y_{i,p})$ are exactly the fundamental modules which are in $\mathcal{C}_{\mathbb{Z}}$.

3. QUANTUM GROTHENDIECK RINGS

In this section, we recall the definition and the first properties of the quantum Grothendieck ring of the monoidal category $\mathcal{C}_{\mathbb{Z}}$. We also discuss some additional symmetry properties of quantum Grothendieck rings with respect to the dualities \mathcal{D} and ω . We keep the notations introduced in the previous section.

3.1. Inverse of quantum Cartan matrix. Let z be an indeterminate. The quantum Cartan matrix of \mathfrak{g} is the $\mathbb{Z}[z^{\pm 1}]$ -valued $I \times I$ -matrix $C(z) = (C_{ij}(z))_{i,j \in I}$ whose (i, j) -entry is given by

$$C_{ij}(z) := \begin{cases} z^{d_i} + z^{-d_i} & \text{if } i = j, \\ [c_{ij}]_z & \text{if } i \neq j, \end{cases}$$

where we used the notation (2.1). As the Cartan matrix $C = C(1)$ is invertible, so is $C(z)$. Denote its inverse by $\tilde{C}(z) = (\tilde{C}_{ij}(z))_{i,j \in I}$. Note that the entries of $\tilde{C}(z)$ are rational functions in z . Then we define a collection of integers $\{\tilde{c}_{ij}(u) \mid i, j \in I, u \in \mathbb{Z}\}$ as the coefficients of the Laurent expansions

$$\sum_{u \in \mathbb{Z}} \tilde{c}_{ij}(u) z^u := [\tilde{C}_{ij}(z)]^+.$$

The $2h^\vee$ -periodicity of these integers for simply-laced types was established in [24]. Here we list several important properties obtained in [13] for general types.

Lemma 3.1 ([13, Section 4]). *The integers $\{\tilde{c}_{ij}(u) \mid i, j \in I, u \in \mathbb{Z}\}$ satisfy the following properties:*

- (1) $\tilde{c}_{ij}(u) = 0$ if $u < d_i$, and $\tilde{c}_{ij}(d_i) = \delta_{i,j}$,
- (2) $\tilde{c}_{ij}(u + d_i) - \tilde{c}_{ij}(u - d_i) = \tilde{c}_{ji}(u + d_j) - \tilde{c}_{ji}(u - d_j)$ for all $u \in \mathbb{Z}$,
- (3) for any $u \in \mathbb{Z}$,

$$\tilde{c}_{ij}(u) = \begin{cases} \tilde{c}_{ji}(u) & \text{if } d_i = d_j, \\ \tilde{c}_{ji}(u + 1) + \tilde{c}_{ji}(u - 1) & \text{if } (d_i, d_j) = (1, 2), \\ \tilde{c}_{ji}(u + 2) + \tilde{c}_{ji}(u) + \tilde{c}_{ji}(u - 2) & \text{if } (d_i, d_j) = (1, 3), \end{cases}$$

- (4) $\tilde{c}_{ij}(u + rh^\vee) = -\tilde{c}_{ij^*}(u)$ for $u \geq 0$,
- (5) $\tilde{c}_{ij}(rh^\vee - u) = \tilde{c}_{ij^*}(u)$ for all $0 \leq u \leq rh^\vee$,
- (6) $\tilde{c}_{ij}(u) \geq 0$ for all $0 \leq u \leq rh^\vee$.

3.2. The quantum torus \mathcal{Y}_t . Let t be an indeterminate. We consider the Laurent polynomial ring $\mathbb{Z}[t^{\pm 1/2}]$, introducing a square root $t^{1/2}$ of t .

Definition 3.2 ([17]). We define the quantum torus \mathcal{Y}_t associated with \mathfrak{g} as the $\mathbb{Z}[t^{\pm 1/2}]$ -algebra presented by the set of generators $\{\tilde{Y}_{i,p}^{\pm 1} \mid (i, p) \in \hat{I}\}$ and the following relations:

- $\tilde{Y}_{i,p} \tilde{Y}_{i,p}^{-1} = \tilde{Y}_{i,p}^{-1} \tilde{Y}_{i,p} = 1$ for each $(i, p) \in \hat{I}$,
- $\tilde{Y}_{i,p} \tilde{Y}_{j,s} = t^{\mathcal{N}(i,p;j,s)} \tilde{Y}_{j,s} \tilde{Y}_{i,p}$ for each $(i, p), (j, s) \in \hat{I}$,

where $\mathcal{N}: (I \times \mathbb{Z})^2 \rightarrow \mathbb{Z}$ is defined by

$$(3.1) \quad \mathcal{N}(i, p; j, s) := \tilde{c}_{ij}(p - s - d_i) - \tilde{c}_{ij}(p - s + d_i) - \tilde{c}_{ij}(s - p - d_i) + \tilde{c}_{ij}(s - p + d_i).$$

Remark 3.3. (i) This is a remark on the convention. Our parameter t here is same as t in [24], which coincides with t^{-1} in [26].

(ii) The definition of the quantum torus in [17] is based on the analysis of vertex operators occurring in the theory of Frenkel-Reshetikhin deformed \mathcal{W} -algebras which are related to transfer-matrices of corresponding quantum integrable systems. The

construction of t -deformed Grothendieck rings for simply-laced quantum loop algebras in [55] and [47] is based on a slightly different quantum torus obtained from convolution products on Nakajima quiver varieties.

Remark 3.4. By Lemma 3.1, we have

$$(3.2) \quad \mathcal{N}(i, p; j, s) = \mathcal{N}(j, p; i, s) = -\mathcal{N}(i, s; j, p) = -\mathcal{N}(j, s; i, p)$$

for any $(i, p), (j, s) \in I \times \mathbb{Z}$, and

$$(3.3) \quad \mathcal{N}(i, p; j, s) = \tilde{c}_{ij}(p - s - d_i) - \tilde{c}_{ij}(p - s + d_i) \quad \text{if } p - s \geq \delta_{i,j}.$$

Moreover, we have

$$(3.4) \quad \tilde{Y}_{i,p} \tilde{Y}_{j,p} = \tilde{Y}_{j,p} \tilde{Y}_{i,p}$$

for $(i, p), (j, p) \in \hat{I}$ as $\mathcal{N}(i, p; j, p) = 0$.

There exists a \mathbb{Z} -algebra homomorphism $\text{ev}_{t=1}: \mathcal{Y}_t \rightarrow \mathcal{Y}$ given by

$$t^{1/2} \mapsto 1, \quad \tilde{Y}_{i,p} \mapsto Y_{i,p}.$$

This map is called *the specialization at $t = 1$* .

An element $\tilde{m} \in \mathcal{Y}_t$ is called a *monomial* if it is a product of the generators $\tilde{Y}_{i,p}$ for $(i, p) \in \hat{I}$ and $t^{\pm 1/2}$. For a monomial $\tilde{m} \in \mathcal{Y}_t$, we set $u_{i,p}(\tilde{m}) := u_{i,p}(\text{ev}_{t=1}(\tilde{m}))$ (recall the notation (2.7)). A monomial \tilde{m} in \mathcal{Y}_t is said to be *dominant* if $\text{ev}_{t=1}(\tilde{m})$ is dominant, i.e., $u_{i,p}(\tilde{m}) \geq 0$ for all $(i, p) \in \hat{I}$. Moreover, for monomials \tilde{m}, \tilde{m}' in \mathcal{Y}_t , set

$$\tilde{m} \leq \tilde{m}' \text{ if and only if } \text{ev}_{t=1}(\tilde{m}) \leq \text{ev}_{t=1}(\tilde{m}').$$

Following [17, Section 6.3], we define the \mathbb{Z} -algebra anti-involution $\overline{(\cdot)}$ on \mathcal{Y}_t by

$$t^{1/2} \mapsto t^{-1/2}, \quad \tilde{Y}_{i,p} \mapsto t \tilde{Y}_{i,p}.$$

It is easy to show that, for any monomial \tilde{m} in \mathcal{Y}_t , there uniquely exists $a \in \mathbb{Z}$ such that $t^{a/2} \tilde{m}$ is $\overline{(\cdot)}$ -invariant, and this element is denoted by $\underline{\tilde{m}}$. Note that $t^{k/2} \tilde{m} = \underline{\tilde{m}}$ for any $k \in \mathbb{Z}$. Hence $\underline{\tilde{m}}$ depends only on $\text{ev}_{t=1}(\tilde{m})$. Therefore, for every monomial m in \mathcal{Y} , the element \underline{m} is well-defined as an element of \mathcal{Y}_t . The elements of this form are called *commutative monomials*. For example, $\underline{Y_{i,p}} = t^{1/2} \tilde{Y}_{i,p}$. Note that, for every monomial m in \mathcal{Y} , we have $\underline{(m^{-1})} = (\underline{m})^{-1} (= \underline{m}^{-1})$. The commutative monomials forms a free basis of the $\mathbb{Z}[t^{\pm 1/2}]$ -module \mathcal{Y}_t . For any two monomials m, m' in \mathcal{Y} , we have

$$\underline{m \cdot m'} = t^{-\mathcal{N}(m, m')/2} \underline{m} \cdot \underline{m'} = t^{\mathcal{N}(m, m')/2} \underline{m'} \cdot \underline{m}$$

where $\mathcal{N}(m, m') \in \mathbb{Z}$ is defined as

$$(3.5) \quad \mathcal{N}(m, m') := \sum_{(i,p), (j,s) \in \hat{I}} u_{i,p}(m) u_{j,s}(m') \mathcal{N}(i, p; j, s).$$

Remark 3.5. In [17], the quantum torus \mathcal{Y}_t is a $\mathbb{Z}[t^{\pm 1}]$ -algebra. However, to guarantee the existence of commutative monomials, we need the square root of $t^{\pm 1}$. This is the reason why we add $t^{\pm 1/2}$.

For $(i, p) \in \hat{I}$, we set

$$\tilde{A}_{i,p+d_i} := \underline{A_{i,p+d_i}} \in \mathcal{Y}_t.$$

3.3. Quantum Grothendieck ring of $\mathcal{C}_{\mathbb{Z}}$. For each $i \in I$, denote by $\mathcal{K}_{i,t}$ the $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra of the quantum torus \mathcal{Y}_t generated by

$$\{\tilde{Y}_{i,p}(1 + t^{-1}\tilde{A}_{i,p+d_i}^{-1}) \mid p - \epsilon_i \in 2\mathbb{Z}\} \cup \{\tilde{Y}_{j,s}^{\pm 1} \mid (j,s) \in \hat{I}, j \neq i\}.$$

Following [47, 55, 17], the quantum Grothendieck ring of $\mathcal{C}_{\mathbb{Z}}$ is defined as

$$\mathcal{K}_t \equiv \mathcal{K}_t(\mathcal{C}_{\mathbb{Z}}) := \bigcap_{i \in I} \mathcal{K}_{i,t}.$$

By construction, the quantum Grothendieck is stable by the $(\overline{\cdot})$ -involution.

Theorem 3.6 ([17, Theorem 5.11]). *For every dominant monomial \tilde{m} in \mathcal{Y}_t , there uniquely exists an element $F_t(\tilde{m})$ of \mathcal{K}_t such that \tilde{m} is the unique dominant monomial occurring in $F_t(\tilde{m})$. The monomials \tilde{m}' occurring in $F_t(\tilde{m}) - \tilde{m}$ satisfy $\tilde{m}' < \tilde{m}$. In particular, the set $\{F_t(\underline{m}) \mid m \in \mathcal{M}\}$ forms a $\mathbb{Z}[t^{\pm 1/2}]$ -basis of \mathcal{K}_t .*

Note that $F_t(\underline{m})$ is $(\overline{\cdot})$ -invariant for any $m \in \mathcal{M}$. It is constructed by an algorithm which is a t -deformation of the Frenkel-Mukhin algorithm [9].

For a dominant monomial \tilde{m} in \mathcal{Y}_t , we set

$$E_t(\tilde{m}) := \tilde{m} \left(\prod_{p \in \mathbb{Z}} \left(\prod_{i \in I: (i,p) \in \hat{I}} \tilde{Y}_{i,p}^{u_{i,p}(\tilde{m})} \right) \right)^{-1} \prod_{p \in \mathbb{Z}} \left(\prod_{i \in I: (i,p) \in \hat{I}} F_t(\tilde{Y}_{i,p})^{u_{i,p}(\tilde{m})} \right),$$

here $\prod_i \tilde{Y}_{i,p}^{u_{i,p}(\tilde{m})}$ is well-defined by (3.4), and $\prod_i F_t(\tilde{Y}_{i,p})^{u_{i,p}(\tilde{m})}$ is well-defined by [17, Lemma 5.12].

Note that $E_t(\tilde{m})$ contains \tilde{m} as its maximal monomial. In particular, by Theorem 3.6,

$$(3.6) \quad E_t(\underline{m}) = F_t(\underline{m}) + \sum_{m' < m} C_{m,m'}(t) F_t(\underline{m}')$$

with $C_{m,m'}(t) \in \mathbb{Z}[t^{\pm 1/2}]$ (in fact, $\mathbb{Z}[t^{\pm 1}]$). Note that the set $\{E_t(\underline{m}) \mid m \in \mathcal{M}\}$ also forms a $\mathbb{Z}[t^{\pm 1/2}]$ -basis of \mathcal{K}_t since $\#\{m' \in \mathcal{M} \mid m' < m\} < \infty$ for any $m \in \mathcal{M}$ (cf. [17, Section 3.4]). For a dominant monomial \tilde{m} in \mathcal{Y}_t with $\text{ev}_{t=1}(\tilde{m}) = m$, we have

$$\text{ev}_{t=1}(E_t(\tilde{m})) = \chi_q(M(m)), \quad \text{ev}_{t=1}(\mathcal{K}_t) = \chi_q(K(\mathcal{C}_{\mathbb{Z}}))$$

[17, Theorem 6.2], here we set

$$M(m) := \bigotimes_{p \in \mathbb{Z}} \left(\bigotimes_{i \in I: (i,p) \in \hat{I}} L(Y_{i,p})^{\otimes u_{i,p}(m)} \right),$$

called a *standard module*. Note that, for any fixed $p \in \mathbb{Z}$, the isomorphism class of the tensor product $\bigotimes_{i \in I: (i,p) \in \hat{I}} L(Y_{i,p})^{\otimes u_{i,p}(m)}$ does not depend on the ordering of the factors, and it is in fact simple [9, Proposition 6.15]. Moreover, the module $M(m)$ has a simple head isomorphic to $L(m)$.⁴ The element $E_t(\underline{m})$ is called *the (q, t) -character of the standard module $M(m)$* .

⁴By convention, we choose the coproduct of $U_q(L\mathfrak{g})$ opposite to that of [1, 33] so that the module $M(m)$ is cyclic from the tensor product of ℓ -highest weight vectors.

We consider another kind of elements $L_t(\underline{m})$ in \mathcal{K}_t which is conjecturally a t -quantum version of the q -character of simple modules.

Theorem 3.7 ([47, Theorem 8.1], [17, Theorem 6.9]). *For a dominant monomial $m \in \mathcal{M}$, there exists a unique element $L_t(\underline{m})$ in \mathcal{K}_t such that*

- (S1) $\overline{L_t(\underline{m})} = L_t(\underline{m})$, and
- (S2) $L_t(\underline{m}) = E_t(\underline{m}) + \sum_{m' \in \mathcal{M}} Q_{m,m'}(t) E_t(\underline{m}')$ with $Q_{m,m'}(t) \in t\mathbb{Z}[t]$.

Moreover, we have $Q_{m,m'}(t) = 0$ unless $m' < m$. In particular, the set $\{L_t(m) \mid m \in \mathcal{M}\}$ forms a $\mathbb{Z}[t^{\pm 1/2}]$ -basis of \mathcal{K}_t .

The element $L_t(\underline{m})$ is called the (q, t) -character of the simple module $L(m)$. For example, if $m = Y_{i,p}$, we have

$$E_t(\underline{Y_{i,p}}) = F_t(\underline{Y_{i,p}}) = L_t(\underline{Y_{i,p}}).$$

But these elements differ for a general m .

In what follows, for a dominant monomial $m \in \mathcal{M}$, we will write

$$F_t(m) := F_t(\underline{m}), \quad E_t(m) := E_t(\underline{m}), \quad L_t(m) := L_t(\underline{m})$$

for simplicity.

Conjecture 3.8 ([17, Conjecture 7.3]). *For all $m \in \mathcal{M}$, we have*

$$\text{ev}_{t=1}(L_t(m)) = \chi_q(L(m)).$$

In fact, a fundamental theorem of Nakajima [47, Theorem 8.1] states that this holds true when \mathfrak{g} is simply-laced. The proof used the geometry of quiver varieties (whose theory is fully developed only for simply-laced types at the moment). This was the main motivation for this conjecture.

Thanks to the unitriangular property (S2), we can write

$$(3.7) \quad E_t(m) = L_t(m) + \sum_{m' \in \mathcal{M}: m' < m} P_{m,m'}(t) L_t(m')$$

with some $P_{m,m'}(t) \in t\mathbb{Z}[t]$ for each $m \in \mathcal{M}$. The polynomials $P_{m,m'}(t)$ are analogs of Kazhdan-Lusztig polynomials for finite-dimensional representations of quantum loop algebras. It is also expected in [17] that these polynomials have positive coefficients. By the results of Nakajima [47], it was already known for simply-laced types.

As for the fundamental modules, we have a nice result for general \mathfrak{g} as follows. Here we say that a monomial $\tilde{m} \in \mathcal{Y}_t$ is *anti-dominant* if \tilde{m}^{-1} is dominant.

Theorem 3.9 ([9, 19]). *For each $(i, p) \in \hat{I}$, the followings hold.*

- (1) *Every monomial in $L_t(Y_{i,p})$ has a positive coefficient.*
- (2) *We have $\text{ev}_{t=1}(L_t(Y_{i,p})) = \chi_q(L(Y_{i,p}))$.*
- (3) *$\tilde{Y}_{j,s}$ appearing in $L_t(Y_{i,p})$ satisfies $p \leq s \leq p + rh^\vee$. Moreover, when $s = p$ (resp. $s = p + rh^\vee$), the monomials containing $\tilde{Y}_{j,s}$ is $\underline{Y_{i,p}}$ (resp. $\underline{Y_{i^*, p+rh^\vee}^{-1}}$), which is the unique dominant (resp. anti-dominant) monomial in $L_t(Y_{i,p})$.*

Remark 3.10. The unitriangular property of the (q, t) -characters of standard modules with respect to $\overline{(\cdot)}$ is included in Theorem 3.7. That is, we have

$$E_t(m) - \overline{E_t(m)} \in \sum_{m' < m} \mathbb{Z}[t^{\pm 1}]E_t(m').$$

3.4. The automorphisms \mathfrak{D}_t and ω_t . Recall the automorphism \mathfrak{D} of \mathcal{Y} defined in Section 2.5 (see (2.3)). As its t -analog, we consider the $\mathbb{Z}[t^{\pm 1/2}]$ -algebra automorphism \mathfrak{D}_t of \mathcal{Y}_t given by

$$(3.8) \quad \mathfrak{D}_t(\tilde{Y}_{i,p}) := \tilde{Y}_{i^*, p+rh^\vee} \quad \text{for } (i, p) \in \hat{I}.$$

Clearly, we have $\text{ev}_{t=1} \circ \mathfrak{D}_t = \mathfrak{D} \circ \text{ev}_{t=1}$. Note also that $\mathfrak{D}_t \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \mathfrak{D}_t$ holds, and hence we have $\mathfrak{D}_t(\underline{m}) = \underline{\mathfrak{D}(m)}$ for any $m \in \mathcal{M}$. In particular, the set of the commutative monomials is stable under \mathfrak{D}_t . As a t -analog of Theorem 2.5, we have the following.

Lemma 3.11. *The automorphism \mathfrak{D}_t of \mathcal{Y}_t restricts to an automorphism on \mathcal{K}_t , for which we will use the same symbol. Moreover, we have*

$$(3.9) \quad \mathfrak{D}_t(L_t(m)) = L_t(\mathfrak{D}(m))$$

for any $m \in \mathcal{M}$.

Proof. Obviously we have $\mathfrak{D}_t(\tilde{A}_{i,p+d_i}) = \tilde{A}_{i^*, p+d_i+rh^\vee}$ for each $(i, p) \in \hat{I}$. Thus the automorphism \mathfrak{D}_t respects the subring $\mathcal{K}_t \subset \mathcal{Y}_t$. Since $\mathfrak{D}_t(F_t(m))$ is an element in \mathcal{K}_t which contains the unique dominant monomial $\mathfrak{D}_t(\underline{m}) = \underline{\mathfrak{D}(m)}$, we have $\mathfrak{D}_t(F_t(m)) = \underline{F_t(\mathfrak{D}(m))}$, and hence $\mathfrak{D}_t(E_t(m)) = E_t(\mathfrak{D}(m))$ for all $m \in \mathcal{M}$. Taking the fact $\mathfrak{D}_t \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \mathfrak{D}_t$ into account, we obtain $\mathfrak{D}_t(L_t(m)) = L_t(\mathfrak{D}(m))$ for each $m \in \mathcal{M}$ from the characterization in Theorem 3.7. \square

Next we define the \mathbb{Z} -algebra involution ω_t on the quantum torus \mathcal{Y}_t given by

$$\omega_t(t^{\pm 1/2}) := t^{\mp 1/2}, \quad \omega_t(\tilde{Y}_{i,p}) := \tilde{Y}_{i,-p}^{-1}$$

for $(i, p) \in \hat{I}$. This is a t -analog of the involution ω considered in Section 2.5. Note that it is well-defined because $\mathcal{N}(i, -p; j, -s) = -\mathcal{N}(i, p; j, s)$ holds by the definition (3.1). Now we prove the following assertion as a t -analog of (2.4).

Lemma 3.12. *The involution ω_t on \mathcal{Y}_t restricts to an involution on \mathcal{K}_t , for which we will use the same symbol. Moreover, we have*

$$(3.10) \quad \omega_t(L_t(m)) = L_t(\omega^* m)$$

for any $m \in \mathcal{M}$. Here $\omega^* m$ is the dominant monomial defined as in Section 2.5.

Proof. Note that we have $\omega_t \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \omega_t$ and hence ω_t stabilizes the set of commutative monomials. For each $i \in I$, the subalgebra $\mathcal{K}_{t,i} \subset \mathcal{Y}_t$ is stable under ω_t because we have

$$\omega_t(\tilde{Y}_{i,p}(1 + t^{-1}\tilde{A}_{i,p+d_i}^{-1})) = \tilde{Y}_{i,-p}^{-1}(1 + t\tilde{A}_{i,-p-d_i}) = M \cdot \tilde{Y}_{i,-p-2d_i}(1 + t^{-1}\tilde{A}_{i,-p-d_i}^{-1}),$$

where $M = t\tilde{A}_{i,-p-d_i}\tilde{Y}_{i,-p}^{-1}\tilde{Y}_{i,-p-2d_i}^{-1}$ is a product of $\tilde{Y}_{j,s}$ with $j \sim i$ and a power of t . Therefore, the quantum Grothendieck ring \mathcal{K}_t is also stable under ω_t .

To prove the equality (3.10), let us consider the *anti*-algebra automorphism $\bar{\omega}_t := \omega_t \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \omega_t$ of \mathcal{Y}_t . Note that it is $\mathbb{Z}[t^{\pm 1/2}]$ -linear and satisfies $\bar{\omega}_t \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \bar{\omega}_t$. Therefore, from Theorem 3.6 and Theorem 3.9 (3), it follows that

$$\omega_t(F_t(Y_{i,p})) = \bar{\omega}_t(F_t(Y_{i,p})) = F_t(Y_{i^*, -p - rh^\vee})$$

for each $(i, p) \in \widehat{I}$. Since $\mathcal{N}(i^*, -p - rh^\vee; j^*, -s - rh^\vee) = \mathcal{N}(j, s; i, p)$ holds by Lemma 3.1, we obtain $\bar{\omega}_t(E_t(m)) = E_t(\omega^* m)$ for any $m \in \mathcal{M}$. Then, by Theorem 3.7, we obtain $\bar{\omega}_t(L_t(m)) = L_t(\omega^* m)$ and hence the equality (3.10) for any $m \in \mathcal{M}$. \square

Remark 3.13. In contrast to (3.10), the equality

$$(3.11) \quad \omega_t(F_t(m)) = F_t(\omega^* m)$$

does not hold in general. In fact, the classical analog of (3.11) already fails. Consider $F(m) = \text{ev}_{t=1}(F_t(m))$, the element in the image of the q -character homomorphism with m as a unique dominant monomial. For $\mathfrak{g} = \mathfrak{sl}_2$, we have for $p \in \mathbb{Z}$:

$$F(Y_p) = Y_p + Y_{p+2}^{-1} = \chi_q(L(Y_p))$$

and

$$F(Y_p Y_{p+2}) = Y_p Y_{p+2} + Y_p Y_{p+4}^{-1} + Y_{p+2}^{-1} Y_{p+4}^{-1} = \chi_q(L(Y_p Y_{p+2})).$$

This implies

$$F(Y_0 Y_2^2) = F(Y_0 Y_2) F(Y_2) = Y_0 Y_2^2 + 2Y_0 Y_2 Y_4^{-1} + Y_4^{-1} + Y_0 Y_4^{-2} + Y_2^{-1} Y_4^{-2}.$$

This element has a unique dominant monomial (in fact, it is equal to $\chi_q(L(Y_0 Y_2^2))$). But its image by ω has 2 dominant monomials, $Y_{-4}^2 Y_{-2}$ and Y_{-4} . So, it is not equal to

$$F(Y_{-4}^2 Y_{-2}) = F(Y_{-4} Y_{-2}) F(Y_{-4}) - F(Y_{-4})$$

which has a monomial Y_{-2}^{-1} with coefficient -1 .

Later in the proof of Proposition 6.10, we will see that (3.11) holds when m corresponds to a Kirillov-Reshetikhin module.

4. COMBINATORICS OF Q-DATA


In this section, we introduce some combinatorial gadgets arising from what we call *Q-data* following [13]. Our exposition is slightly different from the original one in [13]: we start with the simple Lie algebra \mathfrak{g} and then introduce its unfolding (Δ, σ) as follows.

4.1. Unfoldings. For each complex finite-dimensional simple Lie algebra \mathfrak{g} , we associate a unique pair (Δ, σ) , which we call *the unfolding of \mathfrak{g}* , consisting of a simply-laced Dynkin diagram Δ and a graph automorphism σ of Δ as given in the Table 1, where $\text{id}: \Delta_0 \rightarrow \Delta_0$ is the identity map and the automorphisms \vee and $\tilde{\vee}$ are given by the blue arrows in Figure 1 below.

Let Δ_0 denote the set of vertices of the graph Δ . Under the above assignment $\mathfrak{g} \mapsto (\Delta, \sigma)$, we identify the lacing number r of \mathfrak{g} with the order of the automorphism σ , and also identify the set I of Dynkin indices of \mathfrak{g} with the set $\Delta_0 / \langle \sigma \rangle$ of σ -orbits in Δ_0 as suggested by the dotted lines in Figure 1. Then the positive integer $d_i \in \{1, r\}$ coincides with the cardinality of the σ -orbit corresponding to $i \in I$.

r	\mathfrak{g}	Δ (or \mathfrak{g})	σ	h^\vee	ℓ_0
1	A_n	A_n	id	$n+1$	$n(n+1)/2$
	D_n	D_n	id	$2n-2$	$n(n-1)$
	$E_{6,7,8}$	$E_{6,7,8}$	id	12, 18, 30	36, 63, 120
2	B_n	A_{2n-1}	\vee	$2n-1$	$n(2n-1)$
	C_n	D_{n+1}	\vee	$n+1$	$n(n+1)$
	F_4	E_6	\vee	9	36
3	G_2	D_4	$\tilde{\vee}$	4	12

TABLE 1. Unfoldings and associated numerical data

FIGURE 1. Unfoldings for non-simply-laced \mathfrak{g}

In what follows, we use the symbols i, j, \dots to denote the elements of Δ_0 . We write $i \sim j$ if they are adjacent in the graph Δ . We denote the natural quotient map $\Delta_0 \rightarrow I = \Delta_0 / \langle \sigma \rangle$ by $i \mapsto \bar{i}$. By definition, for $i \in \Delta_0$ and $i \in I$, we have $i = \bar{i}$ if and only if $i \in i$. Note that we have $i \sim j$ for $i, j \in I$ (i.e. $c_{ij} < 0$) if and only if there exist $i \in i$ and $j \in j$ satisfying $i \sim j$.

4.2. Notation for simply-laced root systems. Let (Δ, σ) be the unfolding of \mathfrak{g} . Hereafter, we always denote by \mathfrak{g} the simply-laced Lie algebra associated with the Dynkin diagram Δ . In the following paragraphs, we prepare the notation around the root system of \mathfrak{g} (not \mathfrak{g}).

Let $P = \bigoplus_{i \in \Delta_0} \mathbb{Z}\varpi_i$ denote the weight lattice of \mathfrak{g} , where $\{\varpi_i\}_{i \in \Delta_0}$ are the fundamental weights. For each $i \in \Delta_0$, let $\alpha_i := 2\varpi_i - \sum_{j \sim i} \varpi_j \in P$ denote the corresponding

simple root. We set $\mathcal{Q} := \bigoplus_{\iota \in \Delta_0} \mathbb{Z}\alpha_\iota$ and $\mathcal{Q}^+ := \sum_{\iota \in \Delta_0} \mathbb{Z}_{\geq 0}\alpha_\iota$. Let $(\cdot, \cdot): \mathcal{P} \times \mathcal{P} \rightarrow \mathbb{Q}$ denote the symmetric bilinear form defined by $(\varpi_\iota, \alpha_j) = \delta_{\iota, j}$ for $\iota, j \in \Delta_0$. The Weyl group W of \mathfrak{g} is defined as the subgroup of $\text{Aut}(\mathcal{P})$ generated by the simple reflections $\{s_\iota\}_{\iota \in \Delta_0}$, which are given by $s_\iota(\lambda) = \lambda - (\lambda, \alpha_\iota)\alpha_\iota$ for $\lambda \in \mathcal{P}$. The set of roots is defined by $R := W \cdot \{\alpha_\iota\}_{\iota \in \Delta_0}$. We have the decomposition $R = R^+ \sqcup (-R^+)$, where $R^+ := R \cap \mathcal{Q}^+$ is the set of positive roots. In the sequel, we often identify σ with the automorphism $\sigma \in \text{Aut}(\mathcal{P})$ given by $\sigma \cdot \varpi_\iota = \varpi_{\sigma(\iota)}$ for $\iota \in \Delta_0$. Then the pairing (\cdot, \cdot) on \mathcal{P} is invariant under the action of the extended group $W \rtimes \langle \sigma \rangle \subset \text{Aut}(\mathcal{P})$.

For an element $w \in W$, a sequence $\mathbf{i} = (\iota_1, \dots, \iota_l)$ of elements of Δ_0 is called a reduced word for w if it satisfies $w = s_{\iota_1} \cdots s_{\iota_l}$ and l is the smallest among all the sequences with this property. The length of a reduced word of w is called the length of w , denoted by $\ell(w) \in \mathbb{Z}_{\geq 0}$. It is well-known that there exists a unique element $w_0 \in W$ with the largest length $\ell(w_0) = \ell_0 := \#R^+$. From Table 1, we can easily check the equality $nrh^\vee = 2\ell_0$ holds for each \mathfrak{g} . Here we recall that n and h^\vee denote the rank and the dual Coxeter number of \mathfrak{g} (not \mathfrak{g}) respectively.

We define the involution $\iota \mapsto \iota^*$ on Δ_0 by the relation $w_0(\alpha_\iota) = -\alpha_{\iota^*}$. Under the assumption $\sigma \neq \text{id}$, we have $*$ = id if h^\vee is even, and $*$ = σ if h^\vee is odd. In particular, the involution $*$ on Δ_0 induces an involution on the set I , which is trivial for non-simply-laced \mathfrak{g} . This latter involution on I coincides with the involution $i \mapsto i^*$ that appeared in the definition of \mathfrak{D} (see (2.3)). In other words, we have $(\bar{i})^* = \overline{i^*}$ for $i \in \Delta_0$.

4.3. Q-data and generalized Coxeter elements. Let \mathfrak{g} be a finite-dimensional complex simple Lie algebra of rank n and (Δ, σ) the unfolding of \mathfrak{g} .

Definition 4.1. A height function on (Δ, σ) is a function $\xi: \Delta_0 \rightarrow \mathbb{Z}$ satisfying the following conditions (here we put $\xi_\iota := \xi(\iota)$):

- (1) Let $\iota, j \in \Delta_0$ with $\iota \sim j$ and $d_\iota = d_j$. Then we have $|\xi_\iota - \xi_j| = d_\iota = d_j$.
- (2) Let $i, j \in I$ with $i \sim j$ and $d_i = 1 < d_j = r$. Then there is a unique $j \in j$ such that $|\xi_i - \xi_j| = 1$ and $\xi_{\sigma^k(j)} = \xi_j - 2k$ for any $1 \leq k < r$, where $i = \{i\}$.

Such a triple $\mathcal{Q} = (\Delta, \sigma, \xi)$ is called a *Q-datum* for \mathfrak{g} .

Remark 4.2. When \mathfrak{g} is simply-laced, σ is the identity and a height function ξ on (Δ, id) is simply a function $\xi: \Delta_0 \rightarrow \mathbb{Z}$ satisfying $|\xi_\iota - \xi_j| = 1$ for each adjacent pair (ι, j) , namely a height function on a Dynkin quiver. In this sense, the notion of Q-datum is a generalization of the notion of a Dynkin quiver with a height function.

Definition 4.3. Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a Q-datum for \mathfrak{g} . A vertex $\iota \in \Delta_0$ is called a *source* of \mathcal{Q} if we have $\xi_\iota > \xi_j$ for any $j \in \Delta_0$ with $j \sim \iota$. When ι is a source of \mathcal{Q} , we define a new Q-datum $s_\iota \mathcal{Q} = (\Delta, \sigma, s_\iota \xi)$ by setting

$$(s_\iota \xi)_j := \xi_j - 2d_\iota \delta_{\iota, j} \quad \text{for } j \in \Delta_0.$$

We say that a sequence $\mathbf{i} = (\iota_1, \dots, \iota_l)$ of elements of Δ_0 is *adapted to* \mathcal{Q} if ι_k is a source of the Q-datum $s_{\iota_{k-1}} \cdots s_{\iota_2} s_{\iota_1} \mathcal{Q}$ for all $k = 1, 2, \dots, l$.

Given a Q-datum (Δ, σ, ξ) , for each $i \in I$, denote by i° the unique vertex in the σ -orbit i satisfying $\xi_{i^\circ} = \max\{\xi_\iota \mid \iota \in i\}$. We consider the following condition on \mathcal{Q} :

$$(4.1) \quad \text{for each } i \in I \text{ and } 1 \leq k < d_i, \text{ we have } \xi_{\sigma^k(i^\circ)} = \xi_{i^\circ} - 2k.$$

Proposition 4.4 ([13, Section 3.6]). *There is a unique collection $\{\tau_Q\}_Q \subset W \times \langle \sigma \rangle$ labelled by Q -data for \mathfrak{g} and characterized by the following conditions:*

- (1) *If Q satisfies (4.1), we define $\tau_Q := s_{i_1^\circ} \cdots s_{i_n^\circ} \sigma$, where (i_1, \dots, i_n) is any total ordering of I satisfying $\xi_{i_1^\circ} \geq \cdots \geq \xi_{i_n^\circ}$. (In this case, the sequence $(i_1^\circ, \dots, i_n^\circ)$ is a reduced word adapted to Q .)*
- (2) *If $\iota \in \Delta_0$ is a source of Q , we have $\tau_{s_i Q} = s_i \tau_Q s_i$.*

Remark 4.5. When \mathfrak{g} is simply-laced, a Q -datum Q for \mathfrak{g} can be seen as a Dynkin quiver Q with a height function (see Remark 4.2) and the condition (4.1) is trivially satisfied in this case. Then the element τ_Q is nothing but the unique Coxeter element adapted to the quiver Q (in the usual sense).

Definition 4.6. We refer to the unique element τ_Q in Proposition 4.4 as *the generalized Coxeter element* associated with Q .

Proposition 4.7 ([13, Proposition 3.34]). *Let Q be a Q -datum for \mathfrak{g} . Then the order of τ_Q is rh^\vee . If \mathfrak{g} is not simply-laced, we have $\tau_Q^{rh^\vee/2} = -1$.*

4.4. Repetition quivers and their Q -coordinates. Fix a simple Lie algebra \mathfrak{g} and its unfolding (Δ, σ) as above. Recall the notation $d_{ij} := \min(d_i, d_j)$ for $i, j \in I$.


Definition 4.8. Let ξ be a height function on (Δ, σ) . *The repetition quiver associated with (Δ, σ) is the quiver $\widehat{\Delta}^\sigma$ whose vertex set $\widehat{\Delta}_0^\sigma$ and arrow set $\widehat{\Delta}_1^\sigma$ are given by*

$$\begin{aligned} \widehat{\Delta}_0^\sigma &:= \{(\iota, p) \in \Delta_0 \times \mathbb{Z} \mid p - \xi_\iota \in 2d_\iota \mathbb{Z}\}, \\ \widehat{\Delta}_1^\sigma &:= \{(\iota, p) \rightarrow (j, s) \mid (\iota, p), (j, s) \in \widehat{\Delta}_0^\sigma, j \sim \iota, s - p = d_{ij}\}. \end{aligned}$$


Remark 4.9. Although the repetition quiver $\widehat{\Delta}^\sigma$ depends on the choice of height function ξ , it is not essential because any two repetition quivers defined with different choices of height functions are identical to each other after shifting uniformly by an integer. Thus we suppress the dependence on ξ in the notation $\widehat{\Delta}^\sigma$.

Example 4.10. We exhibit some examples of the repetition quivers. Here the labeling of Δ_0 is as in Figure 1.


- (1) When \mathfrak{g} is of type A_5 , its unfolding is (Δ, id) with Δ of type A_5 and the repetition quiver $\widehat{\Delta}^\sigma$ is depicted as:




- (2) When \mathfrak{g} is of type D_5 , its unfolding is (Δ, id) with Δ of type D_5 and the repetition quiver $\widehat{\Delta}^\sigma$ is depicted as:




- (3) When \mathfrak{g} is of type B_3 , its unfolding is (Δ, \vee) with Δ of type A_5 and the repetition quiver $\widehat{\Delta}^\sigma$ is depicted as:



- (4) When \mathfrak{g} is of type C_4 , its unfolding is (Δ, \vee) with Δ of type D_5 and the repetition quiver $\widehat{\Delta}^\sigma$ is depicted as:



- (5) When \mathfrak{g} is of type G_2 , its unfolding is $(\Delta, \widetilde{\vee})$ with Δ of type D_4 and the repetition quiver $\widehat{\Delta}^\sigma$ is depicted as:



Recall we have fixed a parity function $\epsilon: I \rightarrow \{0, 1\}$ for \mathfrak{g} and defined the set \widehat{I} in (2.6). In what follows, we always assume that a height function ξ on (Δ, σ) always satisfies the condition

$$(4.2) \quad \xi_i \equiv \epsilon_{\bar{i}} \pmod{2} \quad \text{for any } i \in \Delta_0.$$

Note that this assumption is not a loss of generality. Then we can define the bijection

$$f: \widehat{\Delta}_0^\sigma \rightarrow \widehat{I} \quad \text{given by } (i, p) \mapsto (\bar{i}, p),$$

which we call *the folding map*.

Next we shall attach a bijection $\phi_Q: \widehat{\Delta}_0^\sigma \rightarrow \mathbb{R}^+ \times \mathbb{Z}$ to each Q -datum $Q = (\Delta, \sigma, \xi)$ for \mathfrak{g} , which we call *the Q -coordinate of $\widehat{\Delta}^\sigma$* . Using the generalized Coxeter element τ_Q associated with Q , we define a positive root $\gamma_i^Q \in \mathbb{R}^+$ for each $i \in \Delta_0$ by

$$(4.3) \quad \gamma_i^Q := (1 - \tau_Q^{d_{\bar{i}}}) \varpi_i.$$

Then the bijection $\phi_{\mathcal{Q}}: \widehat{\Delta}_0^{\sigma} \rightarrow \mathbb{R}^+ \times \mathbb{Z}$ is defined recursively by the following rules:

- (1) $\phi_{\mathcal{Q}}(\iota, \xi_{\iota}) = (\gamma_{\iota}^{\mathcal{Q}}, 0)$ for each $\iota \in \Delta_0$.
- (2) If $\phi_{\mathcal{Q}}(\iota, p) = (\alpha, k)$, we have

$$\phi_{\mathcal{Q}}(\iota, p \pm 2d_{\iota}) = \begin{cases} (\tau_{\mathcal{Q}}^{\mp d_{\iota}} \alpha, k) & \text{if } \tau_{\mathcal{Q}}^{\mp d_{\iota}} \alpha \in \mathbb{R}^+, \\ (-\tau_{\mathcal{Q}}^{\mp d_{\iota}} \alpha, k \pm 1) & \text{if } \tau_{\mathcal{Q}}^{\mp d_{\iota}} \alpha \in -\mathbb{R}^+. \end{cases}$$

Composing with the bijection $f^{-1}: \widehat{I} \rightarrow \widehat{\Delta}_0^{\sigma}$, we get another bijection

$$\bar{\phi}_{\mathcal{Q}} := \phi_{\mathcal{Q}} \circ (f^{-1}): \widehat{I} \rightarrow \mathbb{R}^+ \times \mathbb{Z},$$

which we call *the \mathcal{Q} -coordinate of \widehat{I}* .

Remark 4.11. When \mathfrak{g} is simply-laced (or equivalently, $\sigma = \text{id}$), we regard a \mathcal{Q} -datum as a Dynkin quiver Q with a height function as in Remark 4.2. Then the corresponding coordinate map $\phi_{\mathcal{Q}}$ can be interpreted as an isomorphism between the repetitive quiver $\widehat{\Delta}^{\text{id}}$ and the Auslander-Reiten quiver of the bounded derived category of representations of the quiver Q due to Happel [16]. See [24, Section 2] or [13, Section 3] for more details.

Finally, we recall a connection between the coordinate map $\bar{\phi}_{\mathcal{Q}}$ and the inverse of quantum Cartan matrices. Let us consider the map

$$\pi: \mathbb{R}^+ \times \mathbb{Z} \rightarrow \mathbb{Q} \quad \text{given by } \pi(\alpha, k) := (-1)^k \alpha.$$

Lemma 4.12 ([13, (3.18)]). *For a \mathcal{Q} -datum \mathcal{Q} and $(i, p) \in \widehat{I}$, we have*

$$\pi(\bar{\phi}_{\mathcal{Q}}(i, p)) = \tau_{\mathcal{Q}}^{(\xi_i - p)/2} \gamma_i^{\mathcal{Q}} \quad \text{for any } i \in i.$$

Theorem 4.13 (cf. [13, Theorem 4.8]). *Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a \mathcal{Q} -datum for \mathfrak{g} . For $i, j \in I$ and any $u \in \mathbb{Z}$, we have*

$$(4.4) \quad \tilde{c}_{ij}(u) - \tilde{c}_{ij}(-u) = \begin{cases} (\varpi_{\iota}, \tau_{\mathcal{Q}}^{(u + \xi_j - \xi_i - d_i)/2} \gamma_j^{\mathcal{Q}}) & \text{if } u + \epsilon_i + \epsilon_j + d_i \in 2\mathbb{Z}, \\ 0 & \text{otherwise,} \end{cases}$$

where $\iota, j \in \Delta_0$ are any vertices such that $\iota \in i, j \in j$.

Proof. Note that the RHS of (4.4) does not depend on the choice of vertices $\iota \in i, j \in j$ (cf. [13, Lemma 4.6]). When $u \geq 0$, Lemma 3.1 (1) tells that the LHS of (4.4) is just $\tilde{c}_{ij}(u)$ and then the assertion has been proved in [13, Theorem 4.8]. Letting $c(u)$ denote the LHS of (4.4), we see that $c(u)$ is $2rh^{\vee}$ -periodic as a function in $u \in \mathbb{Z}$, namely we have $c(u + 2rh^{\vee}) = c(u)$ for all $u \in \mathbb{Z}$. Indeed, it is clear unless $-2rh^{\vee} < u < 0$ from Lemma 3.1 (4). Even if $-2rh^{\vee} < u < 0$, we have

$$c(u + 2rh^{\vee}) = \tilde{c}_{ij}(u + 2rh^{\vee}) = -\tilde{c}_{ij}(-u) = c(u)$$

thanks to Lemma 3.1 (4) & (5). On the other hand, the RHS of (4.4) is also $2rh^{\vee}$ -periodic as $\tau_{\mathcal{Q}}^{rh^{\vee}} = 1$ by Proposition 4.7. Therefore the desired equality (4.4) holds for all $u \in \mathbb{Z}$. \square

4.5. **Twisted Auslander-Reiten quivers.** Keep the setting from the last subsection.

Definition 4.14. Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a \mathcal{Q} -datum for \mathfrak{g} . The twisted Auslander-Reiten (AR) quiver of \mathcal{Q} is the full subquiver $\Gamma_{\mathcal{Q}} \subset \widehat{\Delta}^{\sigma}$ whose vertex set $(\Gamma_{\mathcal{Q}})_0$ is given by

$$(\Gamma_{\mathcal{Q}})_0 := \phi_{\mathcal{Q}}^{-1}(\mathbb{R}^+ \times \{0\}).$$

Proposition 4.15 ([13, Section 3.7]). Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a \mathcal{Q} -datum.

- (1) We have $(\Gamma_{\mathcal{Q}})_0 = \{(\iota, p) \in \widehat{\Delta}_0^{\sigma} \mid \xi_{\iota^*} - rh^{\vee} < p \leq \xi_{\iota}\}$.
- (2) For $(\iota, p) \in \widehat{\Delta}_0^{\sigma}$ with $\phi_{\mathcal{Q}}(\iota, p) = (\alpha, k)$, we have $\phi_{\mathcal{Q}}(\iota^*, p \pm rh^{\vee}) = (\alpha, k \mp 1)$.

The twisted Auslander-Reiten quiver $\Gamma_{\mathcal{Q}}$ has a remarkable combinatorial meaning in the theory of commutation classes in the Weyl group W , which will be explained later in subsection 7.2.

Now we put

$$(4.5) \quad \widehat{I}_{\mathcal{Q}} := f((\Gamma_{\mathcal{Q}})_0) = \{(\bar{i}, p) \in \widehat{I} \mid (\iota, p) \in (\Gamma_{\mathcal{Q}})_0\}$$

The following is an immediate consequence of Proposition 4.15.

Corollary 4.16. Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a \mathcal{Q} -datum for \mathfrak{g} .

- (1) If the condition (4.1) is satisfied, we have

$$\widehat{I}_{\mathcal{Q}} = \{(i, p) \in \widehat{I} \mid \xi_{(i^*)^{\circ}} - rh^{\vee} < p \leq \xi_{i^{\circ}}\}.$$

- (2) For $(i, p) \in \widehat{I}$ with $\bar{\phi}_{\mathcal{Q}}(i, p) = (\alpha, k)$, we have $\bar{\phi}_{\mathcal{Q}}(i^*, p \pm rh^{\vee}) = (\alpha, k \mp 1)$.

The following observation will be useful in the sequel.

Lemma 4.17. Given an integer $b \in \mathbb{Z}$, there is a \mathcal{Q} -datum \mathcal{Q} for \mathfrak{g} such that

$$\widehat{I}_{\mathcal{Q}} = \{(i, p) \in \widehat{I} \mid b - rh^{\vee} < p \leq b\}.$$

Proof. In view of Corollary 4.16, it suffices to show that there is a \mathcal{Q} -datum $\mathcal{Q} = (\Delta, \sigma, \xi)$ for \mathfrak{g} satisfying the condition (4.1) and that $\xi_{i^{\circ}} \in \{b, b-1\}$ for all $i \in I$. We can choose ξ explicitly case-by-case as follows. Here ϵ denotes a fixed parity function for \mathfrak{g} and we will use the labeling in Figure 1 for non-simply-laced \mathfrak{g} .

- For \mathfrak{g} simply-laced: for each $i \in \Delta_0 = I$, we set

$$\xi_i := \begin{cases} b - \epsilon_i & \text{if } b \in 2\mathbb{Z}, \\ b - 1 + \epsilon_i & \text{if } b \in 2\mathbb{Z} + 1. \end{cases}$$

This ξ corresponds to a sink-source orientation of Δ .

- For \mathfrak{g} of type B_n : if $b \in 2\mathbb{Z}$, we set

$$\xi_i := \begin{cases} b - \epsilon_{\bar{i}} & \text{if } \iota - n \in (2\mathbb{Z}_{\leq 0} - 1) \sqcup (2\mathbb{Z}_{\geq 0}), \\ b - 2 - \epsilon_{\bar{i}} & \text{if } \iota - n \in (2\mathbb{Z}_{< 0}) \sqcup (2\mathbb{Z}_{\geq 0} + 1). \end{cases}$$

If $b \in 2\mathbb{Z} + 1$, we set

$$\xi_i := \begin{cases} b - 1 + \epsilon_{\bar{i}} & \text{if } \iota - n \in (2\mathbb{Z}_{\leq 0} - 1) \sqcup (2\mathbb{Z}_{\geq 0}), \\ b - 3 + \epsilon_{\bar{i}} & \text{if } \iota - n \in (2\mathbb{Z}_{< 0}) \sqcup (2\mathbb{Z}_{\geq 0} + 1). \end{cases}$$

- For \mathfrak{g} of type C_n : if $b \in 2\mathbb{Z}$, we set

$$\xi_i := \begin{cases} b - \epsilon_{\bar{i}} & \text{if } 1 \leq i \leq n, \\ b - 2 - \epsilon_{\bar{i}} & \text{if } i = n + 1. \end{cases}$$

If $b \in 2\mathbb{Z} + 1$, we set

$$\xi_i := \begin{cases} b - 1 + \epsilon_{\bar{i}} & \text{if } 1 \leq i \leq n, \\ b - 3 + \epsilon_{\bar{i}} & \text{if } i = n + 1. \end{cases}$$

- For \mathfrak{g} of type F_4 : we set

$$(\xi_1, \dots, \xi_6) := \begin{cases} (b, b, b - 2, b - 1, b, b - 2) & \text{if } b \equiv \epsilon_1 \pmod{2}, \\ (b - 1, b - 1, b - 3, b, b - 1, b - 3) & \text{if } b \not\equiv \epsilon_1 \pmod{2}. \end{cases}$$


- For \mathfrak{g} of type G_2 : we set

$$(\xi_1, \dots, \xi_4) := \begin{cases} (b, b - 1, b - 2, b - 4) & \text{if } b \equiv \epsilon_1 \pmod{2}, \\ (b - 1, b, b - 3, b - 5) & \text{if } b \not\equiv \epsilon_1 \pmod{2}. \end{cases} \quad \square$$

Example 4.18. We exhibit explicit examples of the twisted Auslander-Reiten quivers in types A_5 and B_3 associated with some Q-data satisfying the condition in Lemma 4.17. Here the labeling of $\Delta_0 = \{1, 2, 3, 4, 5\}$ is as in Figure 1. Via the map $\bar{\phi}_{\mathcal{Q}}$, the vertices of $\Gamma_{\mathcal{Q}}$ are labeled by positive roots of type A_5 , for which we use the abbreviation $[k, l] := \alpha_k + \alpha_{k+1} + \dots + \alpha_l$ and $[k] := \alpha_k$.


- (1) Let $\mathcal{Q} = (\Delta, \text{id}, \xi)$ be the Q-datum of type A_5 given by $(\xi_1, \dots, \xi_5) = (0, -1, 0, -1, 0)$.

Then the quiver $\Gamma_{\mathcal{Q}}$ is depicted as:



- (2) Let $\mathcal{Q} = (\Delta, \vee, \xi)$ be the Q-datum of type B_3 given by $(\xi_1, \dots, \xi_5) = (-2, 0, -1, -2, 0)$.

Then the quiver $\Gamma_{\mathcal{Q}}$ is depicted as:



5. MONOIDAL SUBCATEGORIES ASSOCIATED WITH Q-DATA

In this section, we see some applications of the theory of Q-data to the representation theory of quantum loop algebras. Although most of the topics have already appeared in [13], we shall give some detailed explanations for completeness. In Section 5.1, we briefly review the notion of Q-weights. In Section 5.2, we define a subcategory $\mathcal{C}_X \subset \mathcal{C}_{\mathbb{Z}}$

supported (in a sense) on an arbitrary subset $X \subset \widehat{I}$ and give a sufficient condition for it to become a monoidal subcategory. We apply this result to the case $X = \widehat{I}_Q$ to define the subcategory \mathcal{C}_Q in Section 5.3. It generalizes the categories \mathcal{C}_Q introduced in [24] for simply-laced \mathfrak{g} associated with a Dynkin quiver Q . In Section 5.4, we introduce the truncation of q -characters and that of (q, t) -characters with respect to a general height function ξ .

5.1. **Q-weights.** First, we recall the notion of Q -weights from [13, Section 5].

Definition 5.1. Let Q be a Q -datum for \mathfrak{g} . For a monomial m in \mathcal{Y} , we define its Q -weight $\text{wt}_Q(m)$ as the element of Q given by

$$\text{wt}_Q(m) := \sum_{(i,p) \in \widehat{I}} u_{i,p}(m) \pi(\bar{\phi}_Q(i,p)).$$

For a monomial $\tilde{m} \in \mathcal{Y}_t$, we set $\text{wt}_Q(\tilde{m}) := \text{wt}_Q(\text{ev}_{t=1}(\tilde{m}))$.

By the Q -weight, the quantum torus \mathcal{Y}_t is equipped with a Q -grading.

Proposition 5.2 (cf. [13, Section 5]). *For any $(i,p) \in \widehat{I}$, we have $\text{wt}_Q(A_{i,p+d_i}) = 0$. In particular, all the elements $F_t(m), E_t(m)$ and $L_t(m)$ for $m \in \mathcal{M}$ are homogeneous with respect to the Q -weight so that*

$$\text{wt}_Q(F_t(m)) = \text{wt}_Q(E_t(m)) = \text{wt}_Q(L_t(m)) = \text{wt}_Q(m).$$

Hence the quantum Grothendieck ring \mathcal{K}_t inherits the Q -grading.

The next fact is a consequence of Corollary 4.16 (2).

Lemma 5.3. *Let Q be a Q -datum for \mathfrak{g} . For any $k \in \mathbb{Z}$ and a homogeneous element $y \in \mathcal{Y}$ (resp. \mathcal{Y}_t) with respect to wt_Q , we have*

$$\text{wt}_Q(\mathfrak{D}^k(y)) = (-1)^k \text{wt}_Q(y), \quad (\text{resp. } \text{wt}_Q(\mathfrak{D}_t^k(y)) = (-1)^k \text{wt}_Q(y)).$$

The following assertion relates the Q -weights with the integers $\mathcal{N}(i,p; j, s)$ (see (3.1)) in the defining relation of the quantum torus \mathcal{Y}_t . It is an immediate consequence of Lemma 4.12 and Theorem 4.13.

Proposition 5.4 ([13, Proposition 5.21]). *Let Q be a Q -datum for \mathfrak{g} . For any two distinct elements $(i,p), (j,s) \in \widehat{I}$ with $p \leq s$, we have*

$$\mathcal{N}(i,p; j,s) = (\text{wt}_Q(Y_{i,p}), \text{wt}_Q(Y_{j,s})).$$

5.2. **Subcategories supported on twisted-convex sets.** Let $X \subset \widehat{I}$ be a subset. Consider the \mathbb{Z} -subalgebra $\mathcal{Y}_X \subset \mathcal{Y}$ (resp. $\mathbb{Z}[t^{\pm 2/1}]$ -subalgebra $\mathcal{Y}_{t,X} \subset \mathcal{Y}_t$) generated by the variables $Y_{i,p}^{\pm 1}$ (resp. $\tilde{Y}_{i,p}^{\pm 1}$) with $(i,p) \in X$, and denote by \mathcal{M}_X the set of dominant monomials in \mathcal{Y}_X . Then we define the category \mathcal{C}_X as the Serre subcategory of $\mathcal{C}_{\mathbb{Z}}$ satisfying $\text{Irr } \mathcal{C}_X = \{[L(m)] \mid m \in \mathcal{M}_X\}$ and the $\mathbb{Z}[t^{\pm 1/2}]$ -module $\mathcal{K}_{t,X}$ as the $\mathbb{Z}[t^{\pm 1/2}]$ -submodule of \mathcal{K}_t given by

$$\mathcal{K}_{t,X} := \bigoplus_{m \in \mathcal{M}_X} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

In what follows, we shall give a sufficient condition for the subcategory \mathcal{C}_X to be monoidal (i.e. closed under taking tensor products) and at the same time for the $\mathbb{Z}[t^{\pm 1/2}]$ -submodule $\mathcal{K}_{t,X}$ to be a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra. Our discussion here is a straightforward generalization of [26, Lemmas 3.26 and 7.9].

Definition 5.5. Let (Δ, σ) be the unfolding of \mathfrak{g} .

- (1) A subset $\tilde{X} \subset \hat{\Delta}_0^\sigma$ is said to be *convex* if it satisfies the following condition: for any oriented path $(x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_l)$ in the repetition quiver $\hat{\Delta}^\sigma$, we have $\{x_1, x_2, \dots, x_l\} \subset \tilde{X}$ if and only if $\{x_1, x_l\} \subset \tilde{X}$.
- (2) We say that a subset $X \subset \hat{I}$ is *twisted-convex* if $f^{-1}(X) \subset \hat{\Delta}_0^\sigma$ is convex.

Lemma 5.6. *If $X \subset \hat{I}$ is twisted-convex, the set \mathcal{M}_X is an ideal of the partially ordered set (\mathcal{M}, \leq) , i.e., it is closed under taking smaller elements in \mathcal{M} with respect to the Nakajima ordering \leq .*

Proof. Suppose $m \in \mathcal{M}_X$ and $mM \in \mathcal{M}$ with M being a monomial in the variables $A_{i,p}^{-1}$'s. Let $A_{i,p}^{-1}$ be a factor of M and let $(i, p \pm d_i) := f^{-1}(i, p \pm d_i)$. Since the product mM is dominant, we have either (a) $Y_{i,p+d_i}$ is a factor of m , or (b) there is a factor $A_{i',p'}^{-1}$ of M which contains $Y_{i,p+d_i}$ as a factor. In the case (a), we have $(i, p+d_i) \in f^{-1}(X)$ by the assumption $m \in \mathcal{M}_X$. In the case (b), we find that there is an oriented path in the repetition quiver $\hat{\Delta}^\sigma$ from $(i, p+d_i)$ to $(i', p'+d_{i'}) := f^{-1}(i', p'+d_{i'})$. Then, we repeat the same argument replacing $A_{i,p}^{-1}$ with $A_{i',p'}^{-1}$, to find either $(i', p'+d_{i'}) \in f^{-1}(X)$, or there is yet another factor $A_{i'',p''}^{-1}$ of M with an oriented path in $\hat{\Delta}^\sigma$ from $(i', p'+d_{i'})$ to $f^{-1}(i'', p''+d_{i''})$. Repeating it finitely many times, we finally find an oriented path (possibly of length zero) in $\hat{\Delta}^\sigma$ from $(i, p+d_i)$ to a vertex in $f^{-1}(X)$. In the similar way, we find an oriented path (possibly of length zero) from a vertex in $f^{-1}(X)$ to $(i, p-d_i)$. As a result, we obtain an oriented path in $\hat{\Delta}^\sigma$ whose end points belong to $f^{-1}(X)$ and which factors through both $(i, p+d_i)$ and $(i, p-d_i)$. Then, by the convexity of $f^{-1}(X)$, we have $A_{i,p}^{-1} \in \mathcal{Y}_X$. Therefore, $M \in \mathcal{Y}_X$ and hence $mM \in \mathcal{Y}_X \cap \mathcal{M} = \mathcal{M}_X$ hold. \square

Proposition 5.7. *Assume that $X \subset \hat{I}$ is twisted-convex. Then*

- (1) *the category \mathcal{C}_X is a monoidal subcategory of $\mathcal{C}_{\mathbb{Z}}$, and*
- (2) *the $\mathbb{Z}[t^{\pm 1/2}]$ -module $\mathcal{K}_{t,X}$ is a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra of \mathcal{K}_t (hence we call it the quantum Grothendieck ring of \mathcal{C}_X).*

Moreover, under the same assumption, we have

$$\mathcal{K}_{t,X} = \bigoplus_{m \in \mathcal{M}_X} \mathbb{Z}[t^{\pm 1/2}]E_t(m) = \bigoplus_{m \in \mathcal{M}_X} \mathbb{Z}[t^{\pm 1/2}]F_t(m)$$

and hence $\text{ev}_{t=1}(\mathcal{K}_{t,X}) = \chi_q(K(\mathcal{C}_X))$.

Proof. Let $m_1, m_2 \in \mathcal{M}_X$. Suppose that a simple module $L(m)$ with $m \in \mathcal{M}$ appears as a composition factor of the tensor product module $L(m_1) \otimes L(m_2)$. Then the dominant monomial m occurs in $\chi_q(L(m_1))\chi_q(L(m_2))$. It implies that $m \leq m_1 m_2$ by Theorem 2.4. From the twisted convexity of X , we have $m \in \mathcal{M}_X$ by Lemma 5.6. This proves that $L(m_1) \otimes L(m_2) \in \mathcal{C}_X$ and hence the assertion (1). The other assertions can

be proved in a similar way using Lemma 5.6 together with Theorem 3.6, the property (3.6) and (S2) in Theorem 3.7. \square

5.3. The category \mathcal{C}_Q . Let $Q = (\Delta, \sigma, \xi)$ be a Q-datum for \mathfrak{g} . Recall the finite subset $\widehat{I}_Q = f((\Gamma_Q)_0) \subset \widehat{I}$ defined in Section 4.5. Since the set $(\Gamma_Q)_0$ is convex by Proposition 4.15 (1), the set \widehat{I}_Q is twisted-convex in the sense of Definition 5.5. Let us apply the construction in the previous subsection to the case $X = \widehat{I}_Q$. We use the notation $(\mathcal{Y}_Q, \mathcal{Y}_{t,Q}, \mathcal{M}_Q, \mathcal{C}_Q, \mathcal{K}_{t,Q})$ for simplicity to denote $(\mathcal{Y}_X, \mathcal{Y}_{t,X}, \mathcal{M}_X, \mathcal{C}_X, \mathcal{K}_{t,X})$ in this case. Thanks to Proposition 5.7, $\mathcal{C}_Q \subset \mathcal{C}_{\mathbb{Z}}$ is a monoidal subcategory and $\mathcal{K}_{t,Q} \subset \mathcal{K}_t$ is a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra. For clarity, we shall state the definition of the category \mathcal{C}_Q in a direct way.

Definition 5.8. We define the category \mathcal{C}_Q as the Serre subcategory of the category $\mathcal{C}_{\mathbb{Z}}$ satisfying $\text{Irr } \mathcal{C}_Q = \{[L(m)] \mid m \in \mathcal{M}_Q\}$, where \mathcal{M}_Q is the set of dominant monomials in the variables $Y_{i,p}$ with $(i,p) \in \widehat{I}_Q$.

Example 5.9. If we consider the Q-datum Q of type A_5 in Example 4.18 (1), the corresponding category \mathcal{C}_Q is monoidally generated by the fundamental modules associated with the variables $\{Y_{i,-2k} \mid i \in \{1, 3, 5\}, 0 \leq k \leq 2\} \cup \{Y_{i,-2k-1} \mid i \in \{2, 4\}, 0 \leq k \leq 2\}$. Similarly, if we consider the Q-datum Q of type B_3 in Example 4.18 (2), the corresponding category \mathcal{C}_Q is monoidally generated by the fundamental modules associated with the variables $\{Y_{i,-2k} \mid i \in \{1, 2\}, 0 \leq k \leq 4\} \cup \{Y_{3,-2k-1} \mid 0 \leq k \leq 4\}$.

For a positive root $\alpha \in \mathbb{R}^+$, we set

$$L^Q(\alpha) := L(Y_{i,p}), \quad L_t^Q(\alpha) := L_t(Y_{i,p}) \quad \text{where } (i,p) = \bar{\phi}_Q^{-1}(\alpha, 0).$$

Note that we have $\text{wt}_Q(L_t^Q(\alpha)) = \alpha$ by Proposition 5.2 and $\text{ev}_{t=1}(L_t^Q(\alpha)) = \chi_q(L^Q(\alpha))$ by Theorem 3.9 (2). The set $\{L^Q(\alpha) \mid \alpha \in \mathbb{R}^+\}$ gives a complete and irredundant collection of fundamental modules in \mathcal{C}_Q up to isomorphisms.

Lemma 5.10. *The followings hold.*

- (1) *The \mathbb{Z} -algebra $K(\mathcal{C}_Q)$ is isomorphic to the polynomial ring $\mathbb{Z}[x_\alpha \mid \alpha \in \mathbb{R}^+]$ in l_0 -many variables via $[L^Q(\alpha)] \mapsto x_\alpha$.*
- (2) *The set $\{L_t^Q(\alpha) \mid \alpha \in \mathbb{R}^+\}$ generates the $\mathbb{Z}[t^{\pm 1/2}]$ -algebra $\mathcal{K}_{t,Q}$.*
- (3) *For each $\alpha \in \mathbb{R}^+$ and $k \in \mathbb{Z}$, we have*

$$\mathcal{D}^k(L^Q(\alpha)) \cong L(Y_{i,p}), \quad \text{where } (i,p) = \bar{\phi}_Q^{-1}(\alpha, k).$$

Thus, the set $\{\mathcal{D}^k(L^Q(\alpha)) \mid (\alpha, k) \in \mathbb{R}^+ \times \mathbb{Z}\}$ gives a complete and irredundant collection of fundamental modules in the category $\mathcal{C}_{\mathbb{Z}}$ up to isomorphisms.

Proof. (1) and (2) are immediate from Proposition 2.9 and Proposition 5.7 respectively. (3) follows from Theorem 2.5 and Corollary 4.16 (2). \square

5.4. Truncation. Let $Q = (\Delta, \sigma, \xi)$ be a Q-datum for \mathfrak{g} . In this subsection, we apply the construction given in Section 5.2 above to the case $X = \widehat{I}_{\leq \xi} \subset \widehat{I}$, where

$$\widehat{I}_{\leq \xi} := f\left(\{(i,p) \in \widehat{\Delta}_0^\sigma \mid p \leq \xi_i\}\right).$$

We use the notation $(\mathcal{Y}_{\leq \xi}, \mathcal{Y}_{t, \leq \xi}, \mathcal{M}_{\leq \xi}, \mathcal{C}_{\leq \xi}, \mathcal{K}_{t, \leq \xi})$ to denote $(\mathcal{Y}_X, \mathcal{Y}_{t, X}, \mathcal{M}_X, \mathcal{C}_X, \mathcal{K}_{t, X})$ in this case. Since the set $\widehat{I}_{\leq \xi}$ is obviously twisted-convex, $\mathcal{C}_{\leq \xi} \subset \mathcal{C}_{\mathbb{Z}}$ is a monoidal subcategory and $\mathcal{K}_{t, \leq \xi} \subset \mathcal{K}_t$ is a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra by Proposition 5.7.

For a (non-commutative) Laurent polynomial $y \in \mathcal{Y}_t$, we denote by $y_{\leq \xi}$ the element of $\mathcal{Y}_{t, \leq \xi}$ obtained from y by discarding all the monomials containing the factors $\widetilde{Y}_{i,p}^{\pm 1}$ with $(i, p) \in \widehat{I} \setminus \widehat{I}_{\leq \xi}$. This assignment $y \mapsto y_{\leq \xi}$ defines a $\mathbb{Z}[t^{\pm 1/2}]$ -linear map $(\cdot)_{\leq \xi}: \mathcal{Y}_t \rightarrow \mathcal{Y}_{t, \leq \xi}$, which is not an algebra homomorphism. By an abuse of notation, we denote the similar \mathbb{Z} -linear homomorphism defined for \mathcal{Y} by the same symbol $(\cdot)_{\leq \xi}: \mathcal{Y} \rightarrow \mathcal{Y}_{\leq \xi}$. Then we have $\text{ev}_{t=1} \circ (\cdot)_{\leq \xi} = (\cdot)_{\leq \xi} \circ \text{ev}_{t=1}$.

Proposition 5.11. *The above linear maps $(\cdot)_{\leq \xi}$ restrict to the injective homomorphisms of algebras*

$$\mathcal{K}_{t, \leq \xi} \hookrightarrow \mathcal{Y}_{t, \leq \xi} \quad \text{and} \quad \chi_q(K(\mathcal{C}_{\leq \xi})) \hookrightarrow \mathcal{Y}_{\leq \xi}.$$

Proof. A proof can be similar to the discussions in [23, Section 6.2] and [25, Proposition 3.10], which treated special cases. In principle, the assertions follow from Theorem 3.6 and [9, Lemma 6.17] respectively, with the help of Lemma 5.6. \square

We refer to the elements $\chi_q(V)_{\leq \xi}$ with $V \in \mathcal{C}_{\leq \xi}$ as *the truncated q -characters*, and refer to the elements $L_t(m)_{\leq \xi}$ and $E_t(m)_{\leq \xi}$ with $m \in \mathcal{M}_{\leq \xi}$ as *the truncated (q, t) -characters*.

Corollary 5.12. *The above linear maps $(\cdot)_{\leq \xi}$ restrict to the injective homomorphisms of algebras*

$$\mathcal{K}_{t, \mathcal{Q}} \hookrightarrow \mathcal{Y}_{t, \mathcal{Q}} \quad \text{and} \quad \chi_q(K(\mathcal{C}_{\mathcal{Q}})) \hookrightarrow \mathcal{Y}_{\mathcal{Q}}.$$

Proof. We have to prove $(\mathcal{K}_{t, \mathcal{Q}})_{\leq \xi} \subset \mathcal{Y}_{t, \mathcal{Q}}$ and $\chi_q(K(\mathcal{C}_{\mathcal{Q}}))_{\leq \xi} \subset \mathcal{Y}_{\mathcal{Q}}$. In view of Lemma 5.10 (1) (resp. (2)), it is enough to show that the q -characters (resp. the (q, t) -characters) of fundamental modules do not involve the variables $Y_{i,p}$ (resp. $\widetilde{Y}_{i,p}$) with $(i, p) \in \widehat{I}_{\leq \xi'}$, where ξ' is another height function on (Δ, σ) given by $\xi'_i := \xi_{i^*} - rh^\vee$ for $i \in \Delta_0$. For q -characters, we can check this property from the twisted-convexity of the set $\widehat{I} \setminus \widehat{I}_{\leq \xi'}$ and the fact that the q -characters of fundamental modules are computed by Frenkel-Mukhin's algorithm, see [9, Theorem 5.9]. For (q, t) -characters, it follows from the case of q -characters via Theorem 3.9. \square

Remark 5.13. By the injectivity of $(\cdot)_{\leq \xi}$, its compatibility with $\overline{(\cdot)}$ and Theorem 3.7, the truncation of a (q, t) -character $L_t(m)_{\leq \xi}$ for each $m \in \mathcal{M}_{\mathcal{Q}}$ can be characterized in the image of truncation $(\mathcal{K}_{t, \mathcal{Q}})_{\leq \xi} \subset \mathcal{Y}_t$ by the following properties:

- (tr-S1) $\overline{L_t(m)_{\leq \xi}} = L_t(m)_{\leq \xi}$, and
- (tr-S2) $L_t(m)_{\leq \xi} - E_t(m)_{\leq \xi} \in \sum_{m' \in \mathcal{M}_{\mathcal{Q}}} t\mathbb{Z}[t]E_t(m')_{\leq \xi}$.

When the height function ξ satisfies the condition in Lemma 4.17 for a given integer $b \in \mathbb{Z}$, we obtain

$$\widehat{I}_{\leq \xi} = \widehat{I}_{\leq b} := \widehat{I} \cap (I \times \mathbb{Z}_{\leq b}).$$

Thus we will denote the corresponding truncation by $(\cdot)_{\leq b}$ instead of $(\cdot)_{\leq \xi}$ for this case. This notation is compatible with the notation in [23] where truncated q -characters for certain categories \mathcal{C}_ℓ of simply-laced quantum loop algebras are introduced.

6. QUANTUM T -SYSTEMS

In this section, we review the classical T -system identities satisfied by the q -characters of Kirillov-Reshetikhin modules and we establish a quantum analog of the T -system identities for all simple Lie algebra \mathfrak{g} in a unified way. As far as the authors know, these relations are new for non simply-laced types (except in type B). We also prove some commutation relations among the Kirillov-Reshetikhin modules involved in the T -systems and some t -commutation relations among related (q, t) -characters, which will be useful for us in the following.

6.1. Kirillov-Reshetikhin modules. For each $(i, p) \in \widehat{I}$ and $k \in \mathbb{Z}_{\geq 0}$, set

$$W_{k,p}^{(i)} := L(m_{k,p}^{(i)}), \quad m_{k,p}^{(i)} := \prod_{s=1}^k Y_{i,p+2(s-1)d_i}.$$

These simple modules are called *Kirillov-Reshetikhin modules* (or *KR modules* for short). Note that $W_{0,p}^{(i)}$ is the trivial $U_q(L\mathfrak{g})$ -module and $W_{1,p}^{(i)} = L(Y_{i,p})$ is a fundamental module.

Theorem 6.1 (cf. [46, Theorem 3.2], [20, Theorem 4.1, Lemma 4.4]). *For $(i, p) \in \widehat{I}$ and $k \in \mathbb{Z}_{\geq 0}$, the q -character of the corresponding KR module $W_{k,p}^{(i)}$ is of the form*

$$(6.1) \quad \chi_q(W_{k,p}^{(i)}) = m_{k,p}^{(i)} (1 + A_{i,p+(2k-1)d_i}^{-1} \chi)$$

where χ is a polynomial in the variables $A_{j,s+d_j}^{-1}$, $(j, s) \in \widehat{I}$.

In this paper, it will be useful to introduce yet another notation for KR modules. For two integers $a, b \in \mathbb{Z}$, we set

$$\begin{aligned} [a, b] &:= \{x \in \mathbb{Z} \mid a \leq x \leq b\}, & (a, b] &:= \{x \in \mathbb{Z} \mid a < x \leq b\}, \\ [a, b) &:= \{x \in \mathbb{Z} \mid a \leq x < b\}, & (a, b) &:= \{x \in \mathbb{Z} \mid a < x < b\}. \end{aligned}$$

We refer to subsets of these forms as *intervals*. Let (Δ, σ) be the unfolding of \mathfrak{g} and $\widehat{\Delta}^\sigma$ its repetitive quiver (see Section 4). For an interval $[a, b] \subset \mathbb{Z}$ and a vertex $\iota \in \Delta_0$, we define a dominant monomial $m^{(\iota)}[a, b] \in \mathcal{M}$ by

$$m^{(\iota)}[a, b] := \prod_{(\iota,p) \in \widehat{\Delta}_0^\sigma, p \in [a,b]} Y_{\iota,p}.$$

Note that the corresponding simple module $W^{(\iota)}[a, b] := L(m^{(\iota)}[a, b])$ is a KR module. Similarly, we define the dominant monomials $m^{(\iota)}(a, b]$, $m^{(\iota)}[a, b)$, $m^{(\iota)}(a, b)$ and denote the corresponding KR modules by $W^{(\iota)}(a, b]$, $W^{(\iota)}[a, b)$, $W^{(\iota)}(a, b)$ respectively.

In this notation, Theorem 6.1 implies the following.

Lemma 6.2. *For any $\iota \in \Delta_0$ and integers $a \leq b$, we have*

$$\begin{aligned} \chi_q \left(W^{(\iota)}[a, b] \right)_{\leq b} &= m^{(\iota)}[a, b], & \chi_q \left(W^{(\iota)}(a, b] \right)_{\leq b} &= m^{(\iota)}(a, b], \\ \chi_q \left(W^{(\iota)}[a, b) \right)_{\leq b-1} &= m^{(\iota)}[a, b), & \chi_q \left(W^{(\iota)}(a, b) \right)_{\leq b-1} &= m^{(\iota)}(a, b), \end{aligned}$$

where $(\cdot)_{\leq b}$ denotes the truncation defined in Section 5.4.

6.2. Classical T -systems. The q -characters of KR modules satisfy a system of identities called *the T -systems*.

Theorem 6.3 ([46, Theorem 1.1], [20, Theorem 3.4]). *For each $(\iota, p), (\iota, s) \in \widehat{\Delta}_0^\sigma$ with $p < s$, we have the following equality in $K(\mathcal{C}_\mathbb{Z})$:*

$$(6.2) \quad \left[W^{(\iota)}[p, s] \right] \left[W^{(\iota)}(p, s) \right] = \left[W^{(\iota)}[p, s] \right] \left[W^{(\iota)}(p, s) \right] + \prod_{j \in \Delta_0; j \sim \iota} \left[W^{(j)}(p, s) \right],$$

where the two terms on right hand side are simple tensor products of Kirillov-Reshetikhin modules.

The second term in the right hand side, there is a change of notations in comparison to [46, 20] as the indices for the product are not picked from I but from Δ_0 . This allows to have a uniform formulation of the T -systems, including non simply-laced types.

Let us consider the following two sets of dominant monomials:

$$(6.3) \quad \mathcal{M}^+(\iota; p, s) := \left\{ m^{(\iota)}[p, s], m^{(\iota)}(p, s), m^{(\iota)}(p, s) \right\} \sqcup \left\{ m^{(j)}(p, s) \mid j \sim \iota \right\},$$

$$(6.4) \quad \mathcal{M}^-(\iota; p, s) := \left(\mathcal{M}^+(\iota; p, s) \setminus \left\{ m^{(\iota)}(p, s) \right\} \right) \sqcup \left\{ m^{(\iota)}[p, s] \right\}.$$

Note that the KR modules corresponding to the dominant monomials in $\mathcal{M}^-(\iota; p, s) \cup \mathcal{M}^+(\iota; p, s)$ are exactly the ones involved in the T -system identity (6.2).

Proposition 6.4. *Let $(\iota, p), (\iota, s) \in \widehat{\Delta}_0^\sigma$ with $p < s$ as in Theorem 6.3. For $\varepsilon \in \{+, -\}$ and any $m, m' \in \mathcal{M}^\varepsilon(\iota; p, s)$, the tensor product $L(m) \otimes L(m')$ is simple. In particular, the set $\{L(m) \mid m \in \mathcal{M}^\varepsilon(\iota; p, s)\}$ forms a mutually commuting family of KR modules.*

Proof. We only consider the case $\varepsilon = +$, as a proof of the other case $\varepsilon = -$ is similar. Our basic tool is the truncation defined in Section 5.4. We divide the situation into the following four cases:

$$(6.5) \quad \{m, m'\} \subset \left\{ m^{(\iota)}[p, s], m^{(\iota)}(p, s) \right\} \sqcup \left\{ m^{(j)}(p, s) \mid j \sim \iota \right\},$$

$$(6.6) \quad \{m, m'\} \subset \left\{ m^{(\iota)}(p, s) \right\} \sqcup \left\{ m^{(j)}(p, s) \mid j \sim \iota \right\},$$

$$(6.7) \quad \{m, m'\} = \left\{ m^{(\iota)}(p, s), m^{(\iota)}(p, s) \right\},$$

$$(6.8) \quad \{m, m'\} = \left\{ m^{(\iota)}[p, s], m^{(\iota)}(p, s) \right\}.$$

For the case (6.5), noting $m^{(j)}(p, s) = m^{(j)}(p, s)$ for $j \sim \iota$, we have $\chi_q(L(m))_{\leq s} = m$ and $\chi_q(L(m'))_{\leq s} = m'$ by Lemma 6.2. Therefore we obtain

$$\chi_q(L(m) \otimes L(m'))_{\leq s} = \chi_q(L(m))_{\leq s} \cdot \chi_q(L(m'))_{\leq s} = mm'.$$

By Proposition 5.11, this shows that $L(m) \otimes L(m') \cong L(mm')$ as desired. For the case (6.6), we apply the same argument with the truncation $(\cdot)_{\leq s}$ replaced by $(\cdot)_{\leq s-1}$.

To deal with the case (6.7), we recall the involution ω defined in the last paragraph of Section 2.5. In our case, it yields

$$\omega^* W^{(\iota)}(p, s) \cong W^{(\iota^*)}(-s - rh^\vee, -p - rh^\vee), \quad \omega^* W^{(\iota)}(p, s) \cong W^{(\iota^*)}[-s - rh^\vee, -p - rh^\vee].$$

Thus we can apply the same argument as in the case (6.6) to obtain the conclusion.

The last case (6.8) was treated in [20, Theorem 6.1 (2)]. \square

6.3. Quantum T -systems. For a general dominant monomial $m \in \mathcal{M}$, the element $F_t(m)$ is given by an algorithm [17, Section 5.7.1]. As a result, we obtain the following t -analog of Theorem 6.1.

Theorem 6.5. *For $(i, p) \in \widehat{I}$ and $k \in \mathbb{Z}_{\geq 0}$, the element $F_t(m_{k,p}^{(i)})$ is of the form*

$$(6.9) \quad F_t(m_{k,p}^{(i)}) = \underline{m_{k,p}^{(i)}} (1 + \widetilde{A}_{i,p+(2k-1)d_i}^{-1} \chi)$$

where χ is a (non-commutative) polynomial in the $\widetilde{A}_{j,s+d_j}^{-1}$, $(j, s) \in \widehat{I}$.

Note that $F_t(m_{k,p}^{(i)})$ is $(\overline{\cdot})$ -invariant as a unique dominant monomial occurs, and this monomial is $(\overline{\cdot})$ -invariant. Although it is known that at $t = 1$, the q -character of $L(m_{k,p}^{(i)})$ has a unique dominant monomial, it is not clear a priori that $L_t(m_{k,p}^{(i)}) = F_t(m_{k,p}^{(i)})$. But we conjecture that they are equal.

Conjecture 6.6. *For any $(i, p) \in \widehat{I}$ and $k \in \mathbb{Z}_{\geq 0}$, we have*

$$L_t(m_{k,p}^{(i)}) = F_t(m_{k,p}^{(i)}).$$

It is known for ADE types by the results of Nakajima. In general it is clear for $k = 1$ by definition of $L_t(m_{1,p}^{(i)})$. Also one can check it for $k = 2$ by a direct computation. We will establish this result in several other situation below.

In this subsection, we establish *the quantum T -systems* in \mathcal{K}_t satisfied by the $F_t(m_{k,p}^{(i)})$. For an interval $[a, b] \subset \mathbb{Z}$ and $\iota \in \Delta_0$, we set $F_t^{(\iota)}[a, b] := F_t(m^{(\iota)}[a, b])$. We define $F_t^{(\iota)}(a, b)$, $F_t^{(\iota)}(a, b)$ and $F_t^{(\iota)}[a, b]$ in the similar way. With this notation, the next assertion follows immediately from Theorem 6.5.

Lemma 6.7. *For any $\iota \in \Delta_0$ and integers $a \leq b$, we have*

$$\begin{aligned} F_t^{(\iota)}[a, b]_{\leq b} &= \underline{m^{(\iota)}[a, b]}, & F_t^{(\iota)}(a, b)_{\leq b} &= \underline{m^{(\iota)}(a, b)}, \\ F_t^{(\iota)}[a, b]_{\leq b-1} &= \underline{m^{(\iota)}[a, b]}, & F_t^{(\iota)}(a, b)_{\leq b-1} &= \underline{m^{(\iota)}(a, b)}. \end{aligned}$$

Theorem 6.8 (Quantum T -systems). *For $(\iota, p), (\iota, s) \in \widehat{\Delta}_0^\sigma$ with $p < s$, there holds the following equality in \mathcal{K}_t :*

$$(6.10) \quad F_t^{(\iota)}[p, s] F_t^{(\iota)}(p, s) = t^x F_t^{(\iota)}[p, s] F_t^{(\iota)}(p, s) + t^y F_t(M(\iota; p, s))$$

for some $x, y \in \frac{1}{2}\mathbb{Z}$, where the dominant monomial $M(\iota; p, s) \in \mathcal{M}$ is given by

$$M(\iota; p, s) := \prod_{j \sim \iota} m^{(j)}(p, s).$$

Note that the quantum T -system is already known for ADE types [24, Proposition 5.6] and for type B [26, Theorem 9.6].

Proof. Let $i := \bar{i}$ and $k := (s - p)/d_i \in \mathbb{Z}_{\geq 1}$. Note that

$$\begin{aligned} F_t^{(i)}[p, s] &= F_t(m_{k,p}^{(i)}), & F_t^{(i)}(p, s) &= F_t(m_{k,p+2d_i}^{(i)}), \\ F_t^{(i)}[p, s] &= F_t(m_{k-1,p+2d_i}^{(i)}), & F_t^{(i)}(p, s) &= F_t(m_{k+1,p}^{(i)}). \end{aligned}$$

Using the algorithm [17, Section 5.7.1], we get a more precise result than formula (6.9), that is the analog of the statement of [20, Lemma 5.5]. Then the same proof as in [20] gives the complete list of dominant monomials occurring in the terms of the quantum T -system: $F_t(m_{k,p}^{(i)})F_t(m_{k,p+2d_i}^{(i)})$ has exactly $k + 1$ dominant monomials M, M_1, \dots, M_k where M, M_1, \dots, M_{k-1} are the dominant monomials occurring in $F_t(m_{k-1,p+2d_i}^{(i)})F_t(m_{k+1,p}^{(i)})$ and $M_k = M(i; p, s)$. But an element in \mathcal{K}_t is characterized by the multiplicities of its dominant monomials [20, Theorem 5.11], so it suffices to check that the multiplicities on the monomials match. Here these multiplicities are just certain powers of t , so we can conclude as in [26, Section 9]. \square

Remark 6.9. The exponents $x, y \in \frac{1}{2}\mathbb{Z}$ in (6.10) can be computed from the t -commutation relations in the quantum torus \mathcal{Y}_t as in [24, Proposition 5.6]. In fact, we have

$$x + 1 = y = \frac{\tilde{c}_{ii}(2(s - p) + d_i) + \tilde{c}_{ii}(2(s - p) - d_i)}{2}.$$

However we do not use this fact in this paper.

The following is the t -analog of Proposition 6.4

Proposition 6.10. *Let $(i, p), (i, s) \in \widehat{\Delta}_0^\sigma$ with $p < s$ as in Theorem 6.3. For $\varepsilon \in \{+, -\}$ and any $m, m' \in \mathcal{M}^\varepsilon(i; p, s)$, we have*

$$F_t(m)F_t(m') = t^{\mathcal{N}(m, m')} F_t(m')F_t(m)$$

in the quantum Grothendieck ring \mathcal{K}_t , where $\mathcal{N}(m, m') \in \mathbb{Z}$ is as in (3.5). In other words, the elements in $\{F_t(m) \mid m \in \mathcal{M}^\varepsilon(i; p, s)\}$ mutually commute up to powers of t .

Proof. We prove the assertions similarly as in the proof of Proposition 6.4 by using the truncation. Again we consider only the case $\varepsilon = +$ and divide the situation into four cases (6.5)–(6.8). For the first two cases (6.5) and (6.6), we simply apply Lemma 6.7 to obtain the conclusion.

To deal with the third case (6.7), we recall the \mathbb{Z} -algebra involution ω_t of \mathcal{K}_t from Section 3.4. Temporarily, let us assume the following equality

$$(6.11) \quad \omega_t \left(F_t^{(i)}[p, s] \right) = F_t^{(i^*)}[-s - rh^\vee, -p - rh^\vee]$$

holds for all $(i, p), (i, s) \in \widehat{\Delta}_0^\sigma$ with $p \leq s$. Then, as special cases, we obtain

$$\omega_t \left(F_t^{(i)}(p, s) \right) = F_t^{(i^*)}(-s - rh^\vee, -p - rh^\vee), \quad \omega_t \left(F_t^{(i)}[p, s] \right) = F_t^{(i^*)}[-s - rh^\vee, -p - rh^\vee].$$

Combining with the same argument as in the case (6.6), we get the conclusion. We postpone the proof of (6.11) to the last paragraph.

For the remaining case (6.8), we apply an \mathfrak{sl}_2 -reduction argument as in [26, Remark 9.10]: the multiplicities of commutative dominant monomials in the polynomials $F_t^{(i)}[p, s]F_t^{(i)}(p, s)$ and $F_t^{(i)}(p, s)F_t^{(i)}[p, s]$ are the same as those for the corresponding

polynomial for the \mathfrak{sl}_2 -case up to overall power of $t^{1/2}$. This follows from the fact that, in the quantum torus, the t -commutation relations between $A_{i,s}^{-1}$, $A_{i,s'}^{-1}$ and between $Y_{i,s}, A_{i,s'}^{-1}$ (with the same index $i \in I$) are the same as in the \mathfrak{sl}_2 -case (q being replaced by q_i). Consequently, as the corresponding polynomial in the \mathfrak{sl}_2 -case commute up to a power of t , we obtain the result.

Finally, we prove the above equality (6.11) by induction on $s - p \geq 0$. For the case $s = p$, it follows immediately from the fact $F_t^{(i)}[p, p] = F_t(Y_{\bar{i}, p}) = L_t(Y_{\bar{i}, p})$ and Lemma 3.12. For the induction step, we apply the *anti*-automorphism $\bar{\omega}_t := \omega_t \circ (\cdot)$ to the quantum T -system identity (6.10). Using the induction hypothesis and the t -commutativity between $F_t^{(i)}[p, s]$ and $F_t^{(i)}(p, s)$ proved in the last paragraph, we have

$$F_t^{(i^*)}[s', p'] F_t^{(i^*)}(s', p') = t^{x'} \bar{\omega}_t \left(F_t^{(i)}[p, s] \right) F_t^{(i^*)}(s', p') + t^y F_t(M(i^*; s', p'))$$

for some $x' \in \frac{1}{2}\mathbb{Z}$, where $s' := -s - rh^\vee$, $p' := -p - rh^\vee$. Comparing it with the quantum T -system identity (6.10) for (i^*, s') and (i^*, p') , we find $\bar{\omega}_t \left(F_t^{(i)}[p, s] \right) = t^{x''} F_t^{(i^*)}[s', p']$ for some $x'' \in \frac{1}{2}\mathbb{Z}$. The bar-invariance forces $x'' = 0$, which completes the proof. \square

Remark 6.11. The above argument implies that an ordered product of $\{F_t^{(j)}(p, s)\}_{j \sim i}$ with respect to any total ordering of the set $\{j \in \Delta_0 \mid j \sim i\}$ is equal to $F_t(M(i; p, s))$ up to a power of t . Therefore, the quantum T -system (6.10) can be expressed as

$$F_t^{(i)}[p, s] F_t^{(i)}(p, s) = t^a F_t^{(i)}[p, s] F_t^{(i)}(p, s) + t^b \prod_{j \sim i}^{\rightarrow} F_t^{(j)}(p, s)$$

for some $a, b \in \frac{1}{2}\mathbb{Z}$.

7. QUANTIZED COORDINATE ALGEBRAS

Let \mathfrak{g} be a complex simple Lie algebra of type ADE and Δ denote its Dynkin diagram. In this section, we collect some preliminaries on the quantum coordinate algebra of the maximal unipotent subgroup N_- associated with \mathfrak{g} and its quantum cluster algebra structure. We will freely use the notation in Section 4.2.

7.1. Quantized coordinate algebra $\mathcal{A}_v[N_-]$. Let v be an indeterminate with a formal square root $v^{1/2}$. Let $\mathcal{U}_v(\mathfrak{n}_-)$ be the $\mathbb{Q}(v^{1/2})$ -algebra presented by the set of generators $\{f_i \mid i \in \Delta_0\}$ satisfying *the quantum Serre relations*:

$$(7.1) \quad \begin{cases} f_i f_j - f_j f_i = 0 & \text{if } i \not\sim j, \\ f_i^2 f_j - (v + v^{-1}) f_i f_j f_i + f_j f_i^2 = 0 & \text{if } i \sim j. \end{cases}$$

This is regarded as a quantized enveloping algebra of the nilpotent Lie subalgebra \mathfrak{n}_- of \mathfrak{g} corresponding to the negative part. For each $i \in \Delta_0$, we write $\text{wt}(f_i) = -\alpha_i$, which endows the algebra $\mathcal{U}_v(\mathfrak{n}_-)$ with a natural $(-\mathbb{Q}^+)$ -grading.

The quantized coordinate algebra $\mathcal{A}_v[N_-]$ is a $\mathbb{Z}[v^{\pm 1/2}]$ -subalgebra of $\mathcal{U}_v(\mathfrak{n}_-)$ which is dual to the canonical integral form of $\mathcal{U}_v(\mathfrak{n}_-)$ generated by the divided powers

$\{f_v^m/[m]_v! \mid v \in \Delta_0, m \in \mathbb{Z}_{\geq 0}\}$ with respect to a suitable pairing. See Appendix A.1 for its precise definition. Here we recall that it satisfies the following properties:

$$\mathbb{Q}(v^{\pm 1/2}) \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathcal{A}_v[N_-] \simeq \mathcal{U}_v(\mathfrak{n}_-), \quad \mathbb{C} \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathcal{A}_v[N_-] \simeq \mathbb{C}[N_-],$$

where \mathbb{C} is regarded as a $\mathbb{Z}[v^{\pm 1/2}]$ -algebra via $v^{1/2} \mapsto 1$, and $\mathbb{C}[N_-]$ is the coordinate algebra of the complex unipotent algebraic group $N_- := \exp(\mathfrak{n}_-)$. Note that $\mathbb{C}[N_-]$ is commutative. We have an obvious algebra homomorphism

$$\text{ev}_{v=1}: \mathcal{A}_v[N_-] \rightarrow \mathbb{C} \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathcal{A}_v[N_-] \simeq \mathbb{C}[N_-]$$

called *the specialization at $v = 1$* .

For each $\lambda \in \mathbb{P}^+ := \sum_{v \in \Delta_0} \mathbb{Z}_{\geq 0} \varpi_v$ and Weyl group elements $w, w' \in W$, we can define a specific homogeneous element $\tilde{D}_{w\lambda, w'\lambda}$ of $\mathcal{A}_v[N_-]$ called *a normalized quantum unipotent minor*, as in [39, Section 6], [14, Section 5]. Its precise definition is recalled in Appendix A.2. From the construction, we have $\text{wt}(\tilde{D}_{w\lambda, w'\lambda}) = w\lambda - w'\lambda$ when $\tilde{D}_{w\lambda, w'\lambda} \neq 0$, and

$$\tilde{D}_{w\lambda, w'\lambda} = \begin{cases} 1 & \text{if } w\lambda = w'\lambda, \\ 0 & \text{if } w\lambda - w'\lambda \notin -\mathbb{Q}^+. \end{cases}$$

7.2. Commutation classes. Before describing the quantum cluster algebra structure of the quantized coordinate algebra $\mathcal{A}_v[N_-]$, we have to recall the generalities on the commutation classes in the Weyl group W .

Two sequences \mathbf{i} and \mathbf{i}' of elements of Δ_0 are said to be *commutation equivalent* if \mathbf{i}' is obtained from \mathbf{i} by applying a sequence of operations which transform some adjacent components (i, j) such that $i \not\sim j$ into (j, i) . This defines an equivalence relation.

For $w \in W$, let $\mathcal{I}(w) \subset \Delta_0^{\ell(w)}$ denote the set of all reduced words for w . We refer to the equivalent class containing \mathbf{i} as *the commutation class of \mathbf{i}* and denote it by $[\mathbf{i}]$. Note that the set $\mathcal{I}(w)$ is divided into a disjoint union of commutation classes. In this paper, we only consider the case $w = w_0$.

For a reduced word $\mathbf{i} = (i_1, \dots, i_{\ell_0}) \in \mathcal{I}(w_0)$, we have $\mathbb{R}^+ = \{\beta_k^{\mathbf{i}} \mid 1 \leq k \leq \ell_0\}$, where $\beta_k^{\mathbf{i}} := s_{i_1} \cdots s_{i_{k-1}}(\alpha_{i_k})$. Thus the reduced word \mathbf{i} defines a total order $<_{\mathbf{i}}$ on \mathbb{R}^+ , namely we write $\beta_k^{\mathbf{i}} <_{\mathbf{i}} \beta_l^{\mathbf{i}}$ if $k < l$. Note that the ordering $<_{\mathbf{i}}$ is *convex* in the sense that if $\alpha, \beta, \alpha + \beta \in \mathbb{R}^+$, we have either $\alpha <_{\mathbf{i}} \alpha + \beta <_{\mathbf{i}} \beta$ or $\beta <_{\mathbf{i}} \alpha + \beta <_{\mathbf{i}} \alpha$. For a commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$, we define the convex partial ordering $\preceq_{[\mathbf{i}]}$ on \mathbb{R}^+ so that

$$\alpha \preceq_{[\mathbf{i}]} \beta \text{ if and only if } \alpha <_{\mathbf{i}'} \beta \text{ for all } \mathbf{i}' \in [\mathbf{i}].$$

For a commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$ and a positive root $\alpha \in \mathbb{R}^+$, we define *the $[\mathbf{i}]$ -residue of α* , denoted by $\text{res}^{[\mathbf{i}]}(\alpha)$, to be $\nu_k \in \Delta_0$ if we have $\alpha = \beta_k^{\mathbf{i}}$ with $\mathbf{i} = (i_1, \dots, i_{\ell_0})$. Note that this is well-defined, i.e. if $\alpha = \beta_l^{\mathbf{i}'}$ for another $\mathbf{i}' = (i'_1, \dots, i'_{\ell_0}) \in [\mathbf{i}]$, then we have $i'_l = \nu_k$.

In [52], *the combinatorial Auslander-Reiten quiver $\Upsilon_{\mathbf{i}}$* was introduced for each reduced word $\mathbf{i} = (i_1, \dots, i_{\ell_0}) \in \mathcal{I}(w_0)$. By definition, it is a quiver whose vertex set is \mathbb{R}^+ and we have an arrow in $\Upsilon_{\mathbf{i}}$ from $\beta_k^{\mathbf{i}}$ to $\beta_j^{\mathbf{i}}$ if and only if $1 \leq j < k \leq \ell_0$, $i_j \sim i_k$ and there is no index j' between j and k such that $i_{j'} \in \{i_j, i_k\}$. The quiver $\Upsilon_{\mathbf{i}}$ satisfies

the following nice properties. We say that a total ordering $R^+ = \{\beta_1, \beta_2, \dots, \beta_{\ell_0}\}$ is a *compatible reading* of $\Upsilon_{\mathbf{i}}$ if we have $k < l$ whenever there is an arrow $\beta_l \rightarrow \beta_k$ in $\Upsilon_{\mathbf{i}}$.

Theorem 7.1 ([52]). *For a commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$, we have the followings:*

- (1) *If $\mathbf{i}' \in [\mathbf{i}]$, then $\Upsilon_{\mathbf{i}} = \Upsilon_{\mathbf{i}'}$. Hence $\Upsilon_{[\mathbf{i}]}$ is well-defined.*
- (2) *For $\alpha, \beta \in R^+$, we have $\alpha \preceq_{[\mathbf{i}]} \beta$ if and only if there exists an oriented path from β to α in $\Upsilon_{[\mathbf{i}]}$. In other words, the quiver $\Upsilon_{[\mathbf{i}]}$ is identical to the Hasse quiver of the partial ordering $\preceq_{[\mathbf{i}]}$.*
- (3) *A sequence $\mathbf{i}' = (\iota'_1, \dots, \iota'_{\ell_0}) \in \Delta_0^{\ell_0}$ belongs to the commutation class $[\mathbf{i}]$ if and only if there is a compatible reading $R^+ = \{\beta_1, \dots, \beta_{\ell_0}\}$ of $\Upsilon_{[\mathbf{i}]}$ such that $\iota'_k = \text{res}^{[\mathbf{i}]}(\beta_k)$ for all $1 \leq k \leq \ell_0$.*

For a reduced word $\mathbf{i} = (\iota_1, \iota_2, \dots, \iota_{\ell_0}) \in \mathcal{I}(w_0)$, we have $\mathbf{i}' = (\iota_2, \dots, \iota_{\ell_0}, \iota_1^*) \in \mathcal{I}(w_0)$ and $[\mathbf{i}] \neq [\mathbf{i}']$. This operation is referred to as a *combinatorial reflection functor* and we write $\mathbf{i}' = r_{\iota_1} \mathbf{i}$. We have the induced operation on commutation classes (i.e., $r_{\iota_1}[\mathbf{i}] := [r_{\iota_1} \mathbf{i}]$ is well-defined). The relations $[\mathbf{i}] \stackrel{r}{\sim} [r_{\iota} \mathbf{i}]$ for various $\iota \in \Delta_0$ generate an equivalence relation, called *the reflection equivalent relation* $\stackrel{r}{\sim}$, on the set of commutation classes in $\mathcal{I}(w_0)$. For a given reduced word $\mathbf{i} \in \mathcal{I}(w_0)$, the family of commutation classes $[[\mathbf{i}]] := \{[\mathbf{i}'] \mid [\mathbf{i}'] \stackrel{r}{\sim} [\mathbf{i}]\}$ is called *an r -cluster point*.

Theorem 7.2 ([53, 13]). *For each Q -datum \mathcal{Q} for \mathfrak{g} , there exists a unique commutation class $[\mathcal{Q}]$ in the set $\mathcal{I}(w_0)$ of reduced words for w_0 such that*

- (1) *$[\mathcal{Q}]$ consists of all the reduced words adapted to \mathcal{Q} in the sense of Definition 4.3,*
- (2) *there is an isomorphism $\Upsilon_{[\mathcal{Q}]} \rightarrow \Gamma_{\mathcal{Q}}$ of quivers whose underlying bijection $R^+ \rightarrow (\Gamma_{\mathcal{Q}})_0$ between the vertex sets coincides with the bijection $\phi_{\mathcal{Q}}^{-1}(\cdot, 0)$,*
- (3) *for each $\alpha \in R^+$, we have $\text{res}^{[\mathcal{Q}]}(\alpha) = \iota$ if $\phi_{\mathcal{Q}}^{-1}(\alpha, 0) = (\iota, p)$.*

Moreover, the set $\{[\mathcal{Q}] \mid \mathcal{Q} \text{ is a } Q\text{-datum for } \mathfrak{g}\}$ forms a single r -cluster point.

7.3. Quantum cluster algebra structure. In this subsection, we briefly describe the quantum cluster algebra structure on the quantum coordinate algebra $\mathcal{A}_v[N_-]$ based on [14]. Here we do not recall the general definition of a quantum cluster algebra, which is reviewed in Appendix B.

We fix a reduced word $\mathbf{i} = (\iota_1, \dots, \iota_{\ell_0}) \in \mathcal{I}(w_0)$ and use the following notation:

$$\begin{aligned} w_{\leq k} &:= s_{\iota_1} s_{\iota_2} \cdots s_{\iota_k}, & w_{\leq 0} &:= 1, \\ k^+ &:= \min(\{k+1 \leq j \leq \ell_0 \mid \iota_j = \iota_k\} \cup \{\ell_0 + 1\}), \\ k^- &:= \max(\{0\} \cup \{1 \leq j \leq k-1 \mid \iota_j = \iota_k\}), \\ k^-(\iota) &:= \max(\{0\} \cup \{1 \leq j \leq k-1 \mid \iota_j = \iota\}) \end{aligned}$$

for $k = 1, 2, \dots, \ell_0$ and $\iota \in \Delta_0$ with $\iota \sim \iota_k$.

Proposition 7.3 ([14, Proposition 10.1, Lemma 11.3]). *For a fixed reduced word $\mathbf{i} = (\iota_1, \dots, \iota_{\ell_0}) \in \mathcal{I}(w_0)$, we set*

$$J := \{1, 2, \dots, \ell_0\}, \quad J_f := \{j \in J \mid j^+ = \ell_0 + 1\}, \quad J_e := J \setminus J_f.$$

Define the \mathbb{Z} -valued $J \times J_e$ -matrix $\tilde{B}^i = (b_{st})_{s \in J, t \in J_e}$ as

$$b_{st} = \begin{cases} 1 & \text{if either (i) } t = s^+, \text{ or (ii) } t < s < t^+ < s^+ \text{ and } \iota_s \sim \iota_t, \\ -1 & \text{if either (i) } s = t^+, \text{ or (ii) } s < t < s^+ < t^+ \text{ and } \iota_s \sim \iota_t, \\ 0 & \text{otherwise.} \end{cases}$$

Define the \mathbb{Z} -valued $J \times J$ -skew-symmetric matrix $\Lambda^i = (\Lambda_{st})_{s, t \in J}$ as

$$\Lambda_{st} = (\varpi_{\iota_s} - w_{\leq s} \varpi_{\iota_s}, \varpi_{\iota_t} + w_{\leq t} \varpi_{\iota_t}) \quad \text{for } s < t.$$

Then (Λ^i, \tilde{B}^i) forms a compatible pair, which satisfies

$$\sum_{k \in J} b_{ks} \Lambda_{kt} = 2\delta_{s,t}$$

for all $s \in J_e$ and $t \in J$.

Recall the notation $\beta_k^i = w_{\leq k-1} \alpha_{\iota_k}$ for $1 \leq k \leq \ell_0$ from Section 7.2. For each $\alpha \in \mathbb{R}^+$, there exists a unique k such that $\alpha = \beta_k^i$. Then we define

$$\alpha^+ := \begin{cases} \beta_{k^+}^i & \text{if } k^+ \leq \ell_0, \\ 0 & \text{if } k^+ = \ell_0 + 1, \end{cases} \quad \alpha^- := \begin{cases} \beta_{k^-}^i & \text{if } k^- > 0, \\ 0 & \text{if } k^- = 0, \end{cases}$$

$$\alpha^-(\iota) := \begin{cases} \beta_{k^-(\iota)}^i & \text{if } k^-(\iota) > 0, \\ 0 & \text{if } k^-(\iota) = 0, \end{cases} \quad \lambda_\alpha := w_{\leq k} \varpi_{\iota_k},$$

where $\iota \in \Delta_0$ is an element such that $\iota \sim \iota_k$. Since $s_\iota \varpi_j = \varpi_j - \delta_{\iota, j} \alpha_\iota$, we have

$$(7.2) \quad \lambda_{\alpha^-} = \lambda_\alpha + \alpha$$

for each $\alpha \in \mathbb{R}^+$. Here we understand $\lambda_{\alpha^-} = \varpi_{\iota_k}$ if $\alpha^- = 0$ (i.e. if $k^- = 0$).

Recall that $\iota_k = \text{res}^{[i]}(\alpha)$ depends only on the commutation class $[i]$. It is easy to see that the assignments $\alpha \mapsto \alpha^+, \alpha^-, \alpha^-(\iota), \lambda_\alpha$ also depend only on the commutation class $[i]$. Thus Proposition 7.3 can be rephrased as follows.

Proposition 7.4. *For a fixed commutation class $[i]$ in $\mathcal{I}(w_0)$, we set*

$$J := \mathbb{R}^+, \quad J_f := \{\alpha \in \mathbb{R}^+ \mid \alpha^+ = 0\}, \quad J_e := J \setminus J_f.$$

Define the \mathbb{Z} -valued $J \times J_e$ -matrix $\tilde{B}^{[i]} = (b_{\alpha\beta})_{\alpha \in J, \beta \in J_e}$ as

$$b_{\alpha\beta} = \begin{cases} 1 & \text{if either (i) } \beta = \alpha^+, \text{ or (ii) } \beta \prec_{[i]} \alpha \prec_{[i]} \beta^+ \prec_{[i]} \alpha^+ \text{ and } \iota \sim j, \\ -1 & \text{if either (i) } \alpha = \beta^+, \text{ or (ii) } \alpha \prec_{[i]} \beta \prec_{[i]} \alpha^+ \prec_{[i]} \beta^+ \text{ and } \iota \sim j, \\ 0 & \text{otherwise.} \end{cases}$$

Here $\iota = \text{res}^{[i]}(\alpha), j = \text{res}^{[i]}(\beta)$. Define the \mathbb{Z} -valued $J \times J$ -skew-symmetric matrix $\Lambda^{[i]} = (\Lambda_{\alpha\beta})_{\alpha, \beta \in J}$ as

$$\Lambda_{\alpha\beta} = (\varpi_\iota - \lambda_\alpha, \varpi_j + \lambda_\beta) \quad \text{for } \beta \not\prec_{[i]} \alpha,$$

where $\iota = \text{res}^{[i]}(\alpha), j = \text{res}^{[i]}(\beta)$. Then $(\Lambda^{[i]}, \tilde{B}^{[i]})$ forms a compatible pair, which satisfies

$$\sum_{\gamma \in J} b_{\gamma\alpha} \Lambda_{\gamma\beta} = 2\delta_{\alpha,\beta}$$

for all $\alpha \in \tilde{J}_e$ and $\beta \in J$.

Let us fix a commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$. Associated with the \mathbb{Z} -valued $\mathbb{R}^+ \times \mathbb{R}^+$ -skew-symmetric matrix $\Lambda^{[\mathbf{i}]}$ in Proposition 7.4, we define the quantum torus $\mathcal{T}_{v, [\mathbf{i}]}$ (denoted by $\mathcal{T}(\Lambda^{[\mathbf{i}]})$ in Appendix B.1) as the $\mathbb{Z}[v^{\pm 1/2}]$ -algebra presented by the set of generators $\{X_\alpha^{\pm 1} \mid \alpha \in \mathbb{R}^+\}$ and the following relations:

- $X_\alpha X_\alpha^{-1} = X_\alpha^{-1} X_\alpha = 1$ for $\alpha \in \mathbb{R}^+$,
- $X_\alpha X_\beta = v^{\Lambda_{\alpha\beta}} X_\beta X_\alpha$ for $\alpha, \beta \in \mathbb{R}^+$.

Let $\mathcal{A}(\Lambda^{[\mathbf{i}]}, \tilde{B}^{[\mathbf{i}]})$ denote the quantum cluster algebra associated with the compatible pair $(\Lambda^{[\mathbf{i}]}, \tilde{B}^{[\mathbf{i}]})$ in Proposition 7.4. This is the $\mathbb{Z}[v^{\pm 1/2}]$ -subalgebra of the quantum torus $\mathcal{T}_{v, [\mathbf{i}]}$ generated by the union of the elements called *the quantum cluster variables*, which are obtained from all possible sequences of *mutations* (see Appendix B.2).

For $\alpha, \beta \in \mathbb{R}^+$ with $\text{res}^{[\mathbf{i}]}(\alpha) = \text{res}^{[\mathbf{i}]}(\beta) = \iota$, we write

$$\tilde{D}^{[\mathbf{i}]}(\alpha, \beta) := \tilde{D}_{\lambda_\alpha, \lambda_\beta}, \quad \tilde{D}^{[\mathbf{i}]}(\alpha, 0) := \tilde{D}_{\lambda_\alpha, \varpi_\iota}.$$

Note that $\tilde{D}^{[\mathbf{i}]}(\alpha, \beta) = 0$ unless $\beta \preceq_{[\mathbf{i}]} \alpha$. We set $\tilde{D}^{[\mathbf{i}]}(0, 0) := 1$ by convention.

Theorem 7.5 ([14, Theorem 12.3], [31, Corollary 11.2.8]). *For a commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$, there exists a $\mathbb{Z}[v^{\pm 1/2}]$ -algebra isomorphism*

$$\text{CL}: \mathcal{A}(\Lambda^{[\mathbf{i}]}, \tilde{B}^{[\mathbf{i}]}) \xrightarrow{\sim} \mathcal{A}_v[N_-]$$

given by

$$X_\alpha \mapsto \tilde{D}^{[\mathbf{i}]}(\alpha, 0)$$

for all $\alpha \in \mathbb{R}^+$. Moreover, the normalized quantum unipotent minors $\tilde{D}^{[\mathbf{i}]}(\alpha, \beta)$ and $\tilde{D}^{[\mathbf{i}]}(\alpha, 0)$ correspond to some quantum cluster variables.

In what follows, we regard $\mathcal{A}_v[N_-]$ as a subalgebra of $\mathcal{T}_{v, [\mathbf{i}]}$ through CL^{-1} .

The following system of equalities are regarded as a quantum analog of determinantal identities, which are also obtained as specific exchange relations of quantum clusters when we consider the quantum cluster algebra structure given in Theorem 7.5.

Proposition 7.6 ([14, Proposition 5.5]). *Let $[\mathbf{i}]$ be a commutation class in $\mathcal{I}(w_0)$. For any $\alpha, \beta \in \mathbb{R}^+$ with $\text{res}^{[\mathbf{i}]}(\alpha) = \text{res}^{[\mathbf{i}]}(\beta) = \iota$ and $\beta \prec_{[\mathbf{i}]} \alpha$, there exist $a, b \in \frac{1}{2}\mathbb{Z}$ such that*

$$(7.3) \quad \tilde{D}^{[\mathbf{i}]}(\alpha, \beta) \tilde{D}^{[\mathbf{i}]}(\alpha^-, \beta^-) = v^a \tilde{D}^{[\mathbf{i}]}(\alpha, \beta^-) \tilde{D}^{[\mathbf{i}]}(\alpha^-, \beta) + v^b \prod_{j \sim \iota}^{\rightarrow} \tilde{D}^{[\mathbf{i}]}(\alpha^-(j), \beta^-(j))$$

in $\mathcal{A}_v[N_-]$. Here the factors in the second term of the RHS mutually commute up to powers of $v^{1/2}$ and thus we can take any total ordering on the set $\{j \in \Delta_0 \mid j \sim \iota\}$.

7.4. Dual canonical basis. We fix a commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$ as before. For $\alpha \in \mathbb{R}^+$, we set

$$\tilde{F}(\alpha, [\mathbf{i}]) := \tilde{D}^{[\mathbf{i}]}(\alpha, \alpha^-).$$

Note that $\deg \tilde{F}(\alpha, [\mathbf{i}]) = -\alpha$. This is an analog of the dual function of a root vector corresponding to $-\alpha$. For $\mathbf{c} = (c_\alpha)_{\alpha \in \mathbf{R}^+} \in (\mathbb{Z}_{\geq 0})^{\mathbf{R}^+}$, we set

$$\nu(\mathbf{c}) := -\frac{1}{2} \sum_{\alpha, \beta \in \mathbf{R}^+} c_\alpha c_\beta (\alpha, \beta) + \sum_{\alpha \in \mathbf{R}^+} c_\alpha^2 \in \mathbb{Z}$$

and define

$$\tilde{F}(\mathbf{c}, [\mathbf{i}]) := v^{\nu(\mathbf{c})/2} \prod_{\alpha \in \mathbf{R}^+} \tilde{F}(\alpha, [\mathbf{i}])^{c_\alpha}.$$

Here we take any total ordering on $\mathbf{R}^+ = \{\beta_1, \dots, \beta_{\ell_0}\}$ such that we have $k > l$ whenever $\beta_k \prec_{[\mathbf{i}]} \beta_l$ (i.e. the opposite of $\prec_{\mathbf{i}'}$ for some $\mathbf{i}' \in [\mathbf{i}]$) for the ordered product. Since $\tilde{F}(\alpha, [\mathbf{i}])$ and $\tilde{F}(\beta, [\mathbf{i}])$ mutually commute if α and β are not comparable with respect to $\prec_{[\mathbf{i}]}$, the element $\tilde{F}(\mathbf{c}, [\mathbf{i}])$ is well-defined.

Proposition 7.7 (see, for instance, [43, Chapter 40, 41]). *The set $\{\tilde{F}(\mathbf{c}, [\mathbf{i}]) \mid \mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbf{R}^+}\}$ forms a $\mathbb{Z}[v^{\pm 1/2}]$ -basis of $\mathcal{A}_v[N_-]$.*

We refer to the basis $\{\tilde{F}(\mathbf{c}, [\mathbf{i}]) \mid \mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbf{R}^+}\}$ as a *normalized dual Poincaré-Birkhoff-Witt type basis* (henceforth, a *normalized dual PBW-type basis*). In particular, $\mathcal{A}_v[N_-]$ is generated by the set $\{\tilde{F}(\alpha, [\mathbf{i}]) \mid \alpha \in \mathbf{R}^+\} = \{\tilde{D}^{[\mathbf{i}]}(\alpha, \alpha^-) \mid \alpha \in \mathbf{R}^+\}$.

There exists a natural \mathbb{Z} -algebra anti-involution $\overline{(\cdot)}$ on the quantum torus $\mathcal{T}_{v, [\mathbf{i}]}$ given by

$$v^{\pm 1/2} \mapsto v^{\mp 1/2}, \quad X_\alpha^{\pm 1} \mapsto X_\alpha^{\pm 1}$$

for $\alpha \in \mathbf{R}^+$. This involution $\overline{(\cdot)}$ preserves the quantum cluster algebra $\mathcal{A}(\Lambda^{[\mathbf{i}]}, \tilde{B}^{[\mathbf{i}]}) \simeq \mathcal{A}_v[N_-]$ and fixes each quantum cluster variable.

Lusztig [41, 42, 43] and Kashiwara [32] have constructed a specific $\mathbb{Z}[v^{\pm 1/2}]$ -basis \mathbf{B}^* of $\mathcal{A}_v[N_-]$, which is called *the dual canonical basis*. Moreover, we can consider the normalized dual canonical basis $\tilde{\mathbf{B}}^*$. We do not recall its precise definition here. Instead, we refer the following characterization of $\tilde{\mathbf{B}}^*$ by using the normalized dual PBW-type basis.

Theorem 7.8 ([39, Theorem 4.29]). *For each commutation class $[\mathbf{i}]$ in $\mathcal{I}(w_0)$, we have*

$$\tilde{\mathbf{B}}^* = \{\tilde{G}(\mathbf{c}, [\mathbf{i}]) \mid \mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbf{R}^+}\},$$

where, for each $\mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbf{R}^+}$, $\tilde{G}(\mathbf{c}, [\mathbf{i}])$ is the unique element of $\mathcal{A}_v[N_-]$ characterized by the following conditions:

- (NDCB1) $\overline{\tilde{G}(\mathbf{c}, [\mathbf{i}])} = \tilde{G}(\mathbf{c}, [\mathbf{i}])$, and
- (NDCB2) $\tilde{G}(\mathbf{c}', [\mathbf{i}]) - \tilde{F}(\mathbf{c}, [\mathbf{i}]) \in \sum_{\mathbf{c}'' \prec_{[\mathbf{i}]} \mathbf{c}'} v\mathbb{Z}[v]\tilde{F}(\mathbf{c}'', [\mathbf{i}])$.

Here the condition $\mathbf{c}' = (c'_\alpha)_{\alpha \in \mathbf{R}^+} \prec_{[\mathbf{i}]} \mathbf{c} = (c_\alpha)_{\alpha \in \mathbf{R}^+}$ means that there exists $\beta \in \mathbf{R}^+$ such that $c'_\beta < c_\beta$ and $c'_\alpha = c_\alpha$ for all $\beta \not\prec_{[\mathbf{i}]} \alpha$.

Remark 7.9. The unitriangular property of the normalized dual PBW-type basis with respect to (\cdot) is also included in Theorem 7.8. That is, we have

$$\tilde{F}(\mathbf{c}, [\mathbf{i}]) - \overline{\tilde{F}(\mathbf{c}, [\mathbf{i}])} \in \sum_{\mathbf{c}' <_{[\mathbf{i}]} \mathbf{c}} \mathbb{Z}[v^{\pm 1}] \tilde{F}(\mathbf{c}', [\mathbf{i}]).$$

8. THE HLO ISOMORPHISMS

Throughout this section, we fix a Q-datum $\mathcal{Q} = (\Delta, \sigma, \xi)$ for \mathfrak{g} .

The aim of this section is to establish an isomorphism $\Phi_{\mathcal{Q}}$ for each Q-datum \mathcal{Q} between the quantum Grothendieck ring $\mathcal{K}_{t, \mathcal{Q}}$ of the category $\mathcal{C}_{\mathcal{Q}}$ and the quantized coordinate algebra $\mathcal{A}_v[N_-]$ associated with the simply-laced Lie algebra \mathfrak{g} unfolding \mathfrak{g} . This is a generalization of the results due to Hernandez-Leclerc [24] and Hernandez-Oya [26], and hence we call $\Phi_{\mathcal{Q}}$ the HLO isomorphism.

In Section 8.1, we prove an isomorphism $\Phi_{\mathcal{Q}}^T$ between the quantum tori $\mathcal{Y}_{t, \mathcal{Q}}$ and $\mathcal{T}_{v, \mathcal{Q}}$ from properties of the inverse of the quantum Cartan matrix. Then in Section 8.2, we verify that $\Phi_{\mathcal{Q}}^T$ induces the desired isomorphism $\Phi_{\mathcal{Q}}$ by comparing of the quantum T -system identities in $\mathcal{K}_{t, \mathcal{Q}}$ (Theorem 6.8) and the quantum determinantal identities in Proposition 7.6. In Section 8.3, as corollaries, we obtain the affirmative answer to the analog of Kazhdan-Lusztig conjecture in [17] for simple modules in the subcategory $\mathcal{C}_{\mathcal{Q}}$ (see Conjecture 3.8) and various positivities for its quantum Grothendieck ring $\mathcal{K}_{t, \mathcal{Q}}$. As far as the authors know, these are the first results in this direction beyond types ABDE. Moreover, our approach is uniform for all non simply-laced types.

Henceforth, we often write \mathcal{Q} instead of $[\mathcal{Q}]$ to simplify the notation. For example, we apply it to write $\mathcal{T}_{v, \mathcal{Q}}, \tilde{D}^{\mathcal{Q}}(\alpha, \beta)$ and $\tilde{F}(\alpha, \mathcal{Q})$ instead of $\mathcal{T}_{v, [\mathcal{Q}]}, \tilde{D}^{[\mathcal{Q}]}(\alpha, \beta)$ and $\tilde{F}(\alpha, [\mathcal{Q}])$ respectively.

8.1. An isomorphism between the quantum tori. Recall the notation for KR modules from Section 6.1. Note that, for each $(\iota, p) \in (\Gamma_{\mathcal{Q}})_0$, the dominant monomial

$$m^{(\iota)}[p, \xi_{\iota}] = \prod_{k=0}^{(\xi_{\iota} - p)/2d_{\bar{\iota}}} Y_{\bar{\iota}, p+2kd_{\bar{\iota}}}$$

belongs to $\mathcal{M}_{\mathcal{Q}}$.

The goal of this subsection is to prove the following theorem, which is a generalization of [26, Theorem 6.3].

Theorem 8.1. *For each Q-datum \mathcal{Q} for \mathfrak{g} , there is the algebra isomorphism*

$$\Phi_{\mathcal{Q}}^T: \mathcal{T}_{v, \mathcal{Q}} \rightarrow \mathcal{Y}_{t, \mathcal{Q}}$$

given by

$$v^{\pm 1/2} \mapsto t^{\pm 1/2}, \quad X_{\alpha} \mapsto \underline{m^{(\iota)}[p, \xi_{\iota}]}$$

for all $\alpha \in \mathbb{R}^+$, where $(\iota, p) = \phi_{\mathcal{Q}}^{-1}(\alpha, 0)$. Moreover, we have $\Phi_{\mathcal{Q}}^T \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \Phi_{\mathcal{Q}}^T$.

The last assertion $\Phi_{\mathcal{Q}}^T \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \Phi_{\mathcal{Q}}^T$ is obvious from the definition of $\Phi_{\mathcal{Q}}^T$. To prove the existence of $\Phi_{\mathcal{Q}}^T$, we notice that the quantum torus $\mathcal{Y}_{t, \mathcal{Q}}$ defined in Section 5.3 has

another presentation given by the set of generators $\{\underline{m}^{(\iota)}[p, \xi_\iota]^{\pm 1} \mid (\iota, p) \in (\Gamma_{\mathcal{Q}})_0\}$ and the following two relations:

- $\underline{m}^{(\iota)}[p, \xi_\iota] \cdot \underline{m}^{(\iota)}[p, \xi_\iota]^{-1} = \underline{m}^{(\iota)}[p, \xi_\iota]^{-1} \cdot \underline{m}^{(\iota)}[p, \xi_\iota] = 1$ for $(\iota, p) \in (\Gamma_{\mathcal{Q}})_0$,
- $\underline{m}^{(\iota)}[p, \xi_\iota] \cdot \underline{m}^{(j)}[s, \xi_j] = t^{\kappa(\iota, p; j, s)} \underline{m}^{(j)}[s, \xi_j] \cdot \underline{m}^{(\iota)}[p, \xi_\iota]$ for $(\iota, p), (j, s) \in (\Gamma_{\mathcal{Q}})_0$,

where

$$\kappa(\iota, p; j, s) := \mathcal{N} \left(\underline{m}^{(\iota)}[p, \xi_\iota], \underline{m}^{(j)}[s, \xi_j] \right) = \sum_{k=0}^{(\xi_\iota - p)/2d_\iota} \sum_{l=0}^{(\xi_j - s)/2d_j} \mathcal{N}(\bar{\iota}, p + 2kd_\iota; \bar{j}, s + 2ld_j).$$

Therefore, it is enough to show that, for any $(\iota, p), (j, s) \in (\Gamma_{\mathcal{Q}})_0$, we have the equality

$$(8.1) \quad \kappa(\iota, p; j, s) = \Lambda_{\alpha\beta},$$

where $\alpha, \beta \in \mathbb{R}^+$ are given by $(\alpha, 0) = \phi_{\mathcal{Q}}(\iota, p)$ and $(\beta, 0) = \phi_{\mathcal{Q}}(j, s)$. A proof of (8.1) will be given in the end of this subsection after some preparation.

Recall the notation α^- and λ_α for each $\alpha \in \mathbb{R}^+$ from Section 7.3, which is associated with the adapted class $[\mathcal{Q}]$ in the present situation.

Lemma 8.2. *Let $\alpha \in \mathbb{R}^+$ and set $\iota := \text{res}^{\mathcal{Q}}(\alpha)$.*

- (1) *We have $\alpha = \tau_{\mathcal{Q}}^{d_\iota} \alpha^-$ provided that $\alpha^- \neq 0$. In other words, we have*

$$\phi_{\mathcal{Q}}^{-1}(\alpha^-, 0) = (\iota, p + 2d_\iota) \quad \text{if } \phi_{\mathcal{Q}}^{-1}(\alpha, 0) = (\iota, p) \text{ with } p < \xi_\iota.$$

- (2) *If $\alpha^- = 0$, we have $\alpha = \gamma_\iota^{\mathcal{Q}} = \varpi_\iota - \tau_{\mathcal{Q}}^{d_\iota} \varpi_\iota$.*

- (3) *We have $\lambda_\alpha = \tau_{\mathcal{Q}}^{(\xi_\iota - p)/2 + d_\iota} \varpi_\iota$, where p is such that $(\alpha, 0) = \phi_{\mathcal{Q}}(\iota, p)$.*

Proof. The assertions (1) and (2) follow from Theorem 7.2. The assertion (3) is proved inductively by the relation (7.2) along with (1) and (2). \square

Given $i, j \in I$, we set

$$(8.2) \quad \Xi_{ij} := \max\{\xi'_j - \xi'_i \in \mathbb{Z} \mid \xi' \text{ is a height function on } (\Delta, \sigma), \iota \in i, j \in j\}.$$

We can easily check that $0 \leq \Xi_{ij} \leq rh^\vee$ holds for any $i, j \in I$.

Lemma 8.3. *Let $i, j \in I$. For any $l \in \mathbb{Z}_{\geq 0}$, we have*

$$(8.3) \quad \tilde{c}_{ij}(\Xi_{ij} - d_i - 2ld_{ij}) = 0$$

In particular, for any Q -datum $\mathcal{Q} = (\Delta, \sigma, \xi)$ for \mathfrak{g} , $\iota \in i, j \in j$ and $l \in \mathbb{Z}_{\geq 0}$, we have

$$(8.4) \quad \tilde{c}_{ij}(\xi_j - \xi_i - d_i - 2ld_{ij}) = 0.$$

Proof. For any height function ξ on (Δ, σ) , $\iota \in i$ and $j \in j$, there is $l \in \mathbb{Z}_{\geq 0}$ such that $\xi_j - \xi_i = \Xi_{ij} - ld_{ij}$. Thus (8.4) follows from (8.3). Since $\tilde{c}_{ij}(u) = 0$ for $u \leq 0$, it is enough to prove (8.3) for any integer l with $0 \leq 2ld_{ij} \leq \Xi_{ij} - d_i$. First we consider the case $d_i \geq d_j$, or equivalently $d_{ij} = d_j$. We choose a height function ξ together with $\iota \in i$ and $j \in j$ so that we have $\Xi_{ij} = \xi_j - \xi_i$ and $\xi_j \geq \xi_{j^*}$. Applying the formula (4.4) in Theorem 4.13 with $u = -(\Xi_{ij} - d_i - 2ld_j)$, we have

$$\tilde{c}_{ij}(u) - \tilde{c}_{ij}(-u) = (\varpi_\iota, \tau_{\mathcal{Q}}^{ld_j} \gamma_j^{\mathcal{Q}}).$$

By the assumption $0 \leq -u \leq \Xi_{ij} - d_i < rh^\vee$, we have $\tilde{c}_{ij}(u) = 0$ and hence the LHS is equal to $-\tilde{c}_{ij}(-u)$, which is non-positive by Lemma 3.1 (6). On the other hand, the RHS is non-negative because $\tau_{\mathcal{Q}}^{ld_j} \gamma_j^{\mathcal{Q}}$ is a positive root thanks to the fact $ld_j \leq \Xi_{ij} \leq \xi_j - \xi_{j^*} + rh^\vee$ and Proposition 4.15. Thus we get $\tilde{c}_{ij}(-u) = 0$, which proves (8.3) for the case $d_i \geq d_j$. For the other case $d_i < d_j$, we necessarily have $(d_i, d_j) = (1, r)$ and hence $d_{ij} = 1$. Moreover, we observe that $\Xi_{ij} = \Xi_{ji} - 2(r-1)$, or equivalently $\Xi_{ij} - d_i = \Xi_{ji} - d_j - r + 1$ holds by Definition 4.1. Then we get

$$\tilde{c}_{ij}(\Xi_{ij} - d_i - 2ld_{ij}) = \sum_{k=0}^{r-1} \tilde{c}_{ji}(\Xi_{ji} - d_j - 2(l+k)) = 0$$

as desired, where the first equality follows from Lemma 3.1 (3). \square

Proposition 8.4. *Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a \mathcal{Q} -datum for \mathfrak{g} . Let $(i, p), (i, p'), (j, s), (j, s') \in \widehat{\Delta}_0^\sigma$ with $p \leq p'$ and $s \leq s'$. Assuming $p - s \leq \Xi_{\bar{i}\bar{j}}$ and $s' - p' \leq \Xi_{\bar{i}\bar{j}}$, we have*

$$\mathcal{N} \left(m^{(i)}[p, p'], m^{(j)}[s, s'] \right) = \left(\tau_{\mathcal{Q}}^{(\xi_i - p)/2 + d_i} \varpi_i + \tau_{\mathcal{Q}}^{(\xi_i - p')/2} \varpi_i, \tau_{\mathcal{Q}}^{(\xi_j - s)/2 + d_j} \varpi_j - \tau_{\mathcal{Q}}^{(\xi_j - s')/2} \varpi_j \right).$$

Proof. Set $i := \bar{i}, j := \bar{j} \in I$. By the definition of $\mathcal{N}(i, p; j, s)$ in (3.1), we have

$$\begin{aligned} \mathcal{N} \left(m^{(i)}[p, p'], m^{(j)}[s, s'] \right) &= \sum_{k=0}^{(p'-p)/2d_i} \sum_{l=0}^{(s'-s)/2d_j} \mathcal{N}(i, p + 2kd_i; j, s + 2ld_j) \\ &= \sum_{k=0}^{(p'-p)/2d_i} \sum_{l=0}^{(s'-s)/2d_j} \{ \tilde{c}_{ij}((p + 2kd_i) - (s + 2ld_j) - d_i) - \tilde{c}_{ij}((p + 2kd_i) - (s + 2ld_j) + d_i) \\ &\quad - \tilde{c}_{ij}((s + 2ld_j) - (p + 2kd_i) - d_i) + \tilde{c}_{ij}((s + 2ld_j) - (p + 2kd_i) + d_i) \} \\ &= \sum_{l=0}^{(s'-s)/2d_j} \{ \tilde{c}_{ij}(p - (s + 2ld_j) - d_i) - \tilde{c}_{ij}(p' - (s' - 2ld_j) + d_i) \\ &\quad - \tilde{c}_{ij}((s' - 2ld_j) - p' - d_i) + \tilde{c}_{ij}((s + 2ld_j) - p + d_i) \} \\ &= \sum_{l=0}^{(s'-s)/2d_j} \{ -\tilde{c}_{ij}(2ld_j - s' + p' + d_i) + \tilde{c}_{ij}(2ld_j + s - p + d_i) \}, \end{aligned}$$

where the last equality follows from Lemma 8.3. On the other hand, we compute

$$\begin{aligned} &\left(\tau_{\mathcal{Q}}^{(\xi_i - p)/2 + d_i} \varpi_i + \tau_{\mathcal{Q}}^{(\xi_i - p')/2} \varpi_i, \tau_{\mathcal{Q}}^{(\xi_j - s)/2 + d_j} \varpi_j - \tau_{\mathcal{Q}}^{(\xi_j - s')/2} \varpi_j \right) \\ &= - \left(\tau_{\mathcal{Q}}^{(\xi_i - p)/2 + d_i} \varpi_i + \tau_{\mathcal{Q}}^{(\xi_i - p')/2} \varpi_i, \tau_{\mathcal{Q}}^{(\xi_j - s')/2} (1 - \tau_{\mathcal{Q}}^{(s'-s)/2 + d_j}) \varpi_j \right) \\ &= - \sum_{l=0}^{(s'-s)/2d_j} \left(\tau_{\mathcal{Q}}^{(\xi_i - p)/2 + d_i} \varpi_i + \tau_{\mathcal{Q}}^{(\xi_i - p')/2} \varpi_i, \tau_{\mathcal{Q}}^{(2ld_j + \xi_j - s')/2} \gamma_j^{\mathcal{Q}} \right) \\ &= - \sum_{l=0}^{(s'-s)/2d_j} \left(\varpi_i, \tau_{\mathcal{Q}}^{((2ld_j - s' + p - d_i) + \xi_j - \xi_i - d_i)/2} \gamma_j^{\mathcal{Q}} + \tau_{\mathcal{Q}}^{((2ld_j - s' + p' + d_i) + \xi_j - \xi_i - d_i)/2} \gamma_j^{\mathcal{Q}} \right) \end{aligned}$$

$$\begin{aligned}
&= - \sum_{l=0}^{(s'-s)/2d_j} \{ \tilde{c}_{ij}(2ld_j - s' + p - d_i) - \tilde{c}_{ij}(-2ld_j + s' - p + d_i) \\
&\quad + \tilde{c}_{ij}(2ld_j - s' + p' + d_i) - \tilde{c}_{ij}(-2ld_j + s' - p' - d_i) \} \\
&= - \sum_{l=0}^{(s'-s)/2d_j} \{ \tilde{c}_{ij}(2ld_j - s' + p' + d_i) - \tilde{c}_{ij}(-2ld_j + s' - p + d_i) \} \\
&= - \sum_{l=0}^{(s'-s)/2d_j} \{ \tilde{c}_{ij}(2ld_j - s' + p' + d_i) - \tilde{c}_{ij}(2ld_j + s - p + d_i) \}
\end{aligned}$$

where the second equality follows from (4.3), the fourth equality follows by Theorem 4.13, and we again used Lemma 8.3 for the fifth equality. From the above two computations, we obtain the conclusion. \square

Remark 8.5.

- (1) In [36], a \mathbb{Z} -valued invariant $\Lambda(V, W)$ was defined for any *rationaly renormalizable* pair (V, W) of non-zero $U_q(\mathbf{L}\mathfrak{g})$ -modules. It plays a quite important role in the theory of monoidal categorification of cluster algebras. If (V, W) is a *commuting* pair of simple modules, say $(L(m), L(m'))$, we have $\Lambda(L(m), L(m')) = \mathcal{N}(m, m')$ (see [13, Section 6]). From this point of view, Proposition 8.4 gives a computation of the invariants $\Lambda(V, W)$ for a larger class of commuting pairs of KR modules.
- (2) When $m^{(i)}[p, p']$, $m^{(j)}[s, s']$ in Proposition 8.4 are contained in some $\mathcal{M}_{\mathcal{Q}}$, the \mathbb{Z} -integer value $\mathcal{N}(m^{(i)}[p, p'], m^{(j)}[s, s'])$ coincides also with the \mathbb{Z} -integer value in [31, Proposition 10.2.3 (ii)] by [37, Theorem 4.12] (see also [35, Theorem 4.12]), since there are certain two simple commuting modules over symmetric quiver Hecke algebra of type \mathfrak{g} corresponding to $L(m^{(i)}[p, p'])$ and $L(m^{(j)}[s, s'])$ via the generalized quantum affine Schur-Weyl duality functor $F_{\mathcal{Q}}$ (see Subsection 9.2 below).

End of the proof of Theorem 8.1. To prove the equality (8.1), we apply Proposition 8.4 to the case $(i, p) = \phi_{\mathcal{Q}}^{-1}(\alpha, 0)$, $(j, s) = \phi_{\mathcal{Q}}^{-1}(\beta, 0)$, $p' = \xi_i$ and $s' = \xi_j$. Since both sides of (8.1) are skew-symmetric, we may assume that $p \leq s$. Then we have $\alpha \not\leq_{[i]} \beta$ and the conditions $p - s \leq \Xi_{\bar{i}\bar{j}}$ and $s' - p' \leq \Xi_{\bar{i}\bar{j}}$ trivially hold. Applying Proposition 8.4 and Lemma 8.2 (3), we obtain

$$\mathcal{N} \left(m^{(i)}[p, \xi_i], m^{(j)}[s, \xi_j] \right) = (\lambda_{\alpha} + \varpi_i, \lambda_{\beta} - \varpi_j) = -\Lambda_{\beta\alpha},$$

which proves (8.1). \square

8.2. The HLO isomorphisms. In this subsection, we prove the following result which generalizes [24, Theorem 6.1] (for simply-laced \mathfrak{g}) and [26, Theorem 10.1] (for \mathfrak{g} of type B) to any simple Lie algebra \mathfrak{g} . Our proof is similar to that of [26]. Let $\mathcal{K}_{t, \mathcal{Q}}^T \subset \mathcal{Y}_{t, \mathcal{Q}}$ denote the image of the truncation map $(\cdot)_{\leq \xi}: \mathcal{K}_{t, \mathcal{Q}} \hookrightarrow \mathcal{Y}_{t, \mathcal{Q}}$ (see Corollary 5.12).

Theorem 8.6. *For any Q -datum \mathcal{Q} for \mathfrak{g} , the isomorphism $\Phi_{\mathcal{Q}}^T: \mathcal{T}_{v,\mathcal{Q}} \rightarrow \mathcal{Y}_{t,\mathcal{Q}}$ in Theorem 8.1 restricts to an isomorphism*

$$\mathcal{A}_v[N_-] \rightarrow \mathcal{K}_{t,\mathcal{Q}}^T,$$

which sends the normalized dual canonical basis $\tilde{\mathbf{B}}^*$ in $\mathcal{A}_v[N_-]$ to the basis formed by the truncated (q,t) -characters of simple modules in $\mathcal{K}_{t,\mathcal{Q}}^T$. More precisely, for each $\mathbf{c} = (c_\alpha)_{\alpha \in \mathbb{R}^+} \in (\mathbb{Z}_{\geq 0})^{\mathbb{R}^+}$, we have $\Phi_{\mathcal{Q}}^T(\tilde{G}(\mathbf{c}, \mathcal{Q})) = L_t(m(\mathbf{c}))_{\leq \xi}$, where $m(\mathbf{c}) \in \mathcal{M}_{\mathcal{Q}}$ is given by

$$(8.5) \quad u_{i,p}(m(\mathbf{c})) = c_\alpha \quad \text{when } \bar{\phi}_{\mathcal{Q}}(i,p) = (\alpha, 0)$$

for each $(i,p) \in \widehat{I}_{\mathcal{Q}}$.

Proof. We prepare some notation. Recall $(\Gamma_{\mathcal{Q}})_0 = \{(i,p) \in \widehat{\Delta}_0^\sigma \mid \xi_{i^*} - rh^\vee < p \leq \xi_i\}$ from Proposition 4.15. Let

$$S_{\mathcal{Q}} := \{(i;p,s) \in \Delta_0 \times \mathbb{Z}^2 \mid (i,p), (i,s) \in \widehat{\Delta}_0^\sigma, \xi_{i^*} - rh^\vee < p \leq s \leq \xi_i + 2d_i\}$$

and choose a total ordering \prec on $S_{\mathcal{Q}}$ satisfying

$$(i;p,s) \prec (i';p',s') \text{ if either } s > s', \text{ or } s = s' \text{ and } p > p'.$$

For each $(i;p,s) \in S_{\mathcal{Q}}$, we set

$$\tilde{D}(i;p,s) := \begin{cases} \tilde{D}^{\mathcal{Q}}(\alpha, \beta) & \text{if } s \leq \xi_i, \\ \tilde{D}^{\mathcal{Q}}(\alpha, 0) & \text{if } s = \xi_i + 2d_i, \\ 1 & \text{if } p = s, \end{cases}$$

where $\alpha, \beta \in \mathbb{R}^+$ correspond to $(i,p), (i,s) \in (\Gamma_{\mathcal{Q}})_0$ respectively via the bijection $\phi_{\mathcal{Q}}^{-1}(\cdot, 0)$. Note that the equality (7.3) can be rewritten as

$$(8.6) \quad \tilde{D}(i;p,s)\tilde{D}(i;p^+,s^+) = v^a \tilde{D}(i;p,s^+)\tilde{D}(i;p^+,s) + v^b \prod_{j \sim i}^{\rightarrow} \tilde{D}(j;p^+(j),s^+(j))$$

where $p^+ := p + 2d_i$ and $p^+(j) := \min\{p' \in \mathbb{Z} \mid (j,p') \in \widehat{\Delta}_0^\sigma, p < p'\}$ (see Theorem 7.1 and Lemma 8.2). Proposition 7.6 tells that the equality (8.6) holds for each $(i;p,s) \in S_{\mathcal{Q}}$ with $p < s \leq \xi_i$ if we choose suitable $a, b \in \frac{1}{2}\mathbb{Z}$ (with an ordering of $\{j \in \Delta_0 \mid j \sim i\}$ fixed).

To prove $\Phi_{\mathcal{Q}}^T(\mathcal{A}_v[N_-]) = \mathcal{K}_{t,\mathcal{Q}}^T$, we only have to show

$$(8.7) \quad \Phi_{\mathcal{Q}}^T(\tilde{D}(i;p,s)) = F_t^{(i)}[p,s]_{\leq \xi}$$

for each $(i;p,s) \in S_{\mathcal{Q}}$ in view of Proposition 7.7. We proceed by induction with respect to the ordering \prec . In the case when $s = \xi_i + 2d_i$, we have

$$\Phi_{\mathcal{Q}}^T(\tilde{D}(i;p,s)) = \Phi_{\mathcal{Q}}^T(\tilde{D}^{\mathcal{Q}}(\alpha, 0)) = \Phi_{\mathcal{Q}}^T(X_\alpha) = \underline{m^{(i)}[p, \xi_i]} = F_t^{(i)}[p, \xi_i]_{\leq \xi} = F_t^{(i)}[p, s]_{\leq \xi}$$

thanks to Theorem 6.5, where $(\alpha, 0) = \phi_{\mathcal{Q}}(i,p)$. In the case $p = s$, it is trivial. In what follows, we focus on the other case $p < s \leq \xi_i$.

We denote by $\mathbb{F}(\mathcal{T}_{v,\mathcal{Q}})$ (resp. $\mathbb{F}(\mathcal{Y}_{t,\mathcal{Q}})$) be the skew-field of fractions of $\mathcal{T}_{v,\mathcal{Q}}$ (resp. $\mathcal{Y}_{t,\mathcal{Q}}$). We can extend the \mathbb{Z} -algebra anti-involution $\overline{(\cdot)}$ on $\mathcal{T}_{v,\mathcal{Q}}$ (resp. $\mathcal{Y}_{t,\mathcal{Q}}$) to the \mathbb{Q} -algebra

anti-involution on $\mathbb{F}(\mathcal{T}_{v,\mathcal{Q}})$ (resp. $\mathbb{F}(\mathcal{Y}_{t,\mathcal{Q}})$), denoted again by $\overline{(\cdot)}$, and extend the \mathbb{Z} -algebra isomorphism $\Phi_{\mathcal{Q}}^T$ to the \mathbb{Q} -algebra isomorphism $\Phi_{\mathcal{Q}}^T: \mathbb{F}(\mathcal{T}_{v,\mathcal{Q}}) \rightarrow \mathbb{F}(\mathcal{Y}_{t,\mathcal{Q}})$. They still satisfy $\Phi_{\mathcal{Q}}^T \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \Phi_{\mathcal{Q}}^T$.

It is shown in [14, Section 13] that the equality (8.6) (more generally (7.3)) corresponds to a specific exchange relation of quantum clusters in the quantum cluster algebra structure given in Theorem 7.5. In particular, the (non-trivial) normalized quantum unipotent minors corresponding to the following elements of $S_{\mathcal{Q}}$

$$(8.8) \quad (i; p^+, s^+), \quad (i; p, s^+), \quad (i; p^+, s), \quad (j; p^+(j), s^+(j))$$

simultaneously belong to a certain quantum cluster. Therefore the element $\tilde{D}(i; p, s)$ is characterized as the $\overline{(\cdot)}$ -invariant element of the form

$$\left(v^a \tilde{D}(i; p, s^+) \tilde{D}(i; p^+, s) + v^b \prod_{j \sim i}^{\rightarrow} \tilde{D}(j; p^+(j), s^+(j)) \right) \tilde{D}(i; p^+, s^+)^{-1}$$

for some $a, b \in \frac{1}{2}\mathbb{Z}$ in $\mathbb{F}(\mathcal{T}_{v,\mathcal{Q}})$. Since all the elements listed in (8.8) are smaller than $(i; p, s)$ with respect to \prec , we can apply the induction hypothesis to find that the element $\Phi_{\mathcal{Q}}^T(\tilde{D}(i; p, s))$ is characterized as the $\overline{(\cdot)}$ -invariant element of the form

$$\left(t^a F_t^{(i)}[p, s^+]_{\leq \xi} \cdot F_t^{(i)}[p^+, s]_{\leq \xi} + t^b \prod_{j \sim i}^{\rightarrow} F_t^{(j)}[p^+(j), s^+(j)]_{\leq \xi} \right) F_t^{(i)}[p^+, s^+]_{\leq \xi}^{-1}$$

for some $a, b \in \frac{1}{2}\mathbb{Z}$ in $\mathbb{F}(\mathcal{Y}_{t,\mathcal{Q}})$. On the other hand, by Theorem 6.8 (see also Remark 6.11), $F_t^{(i)}[s, p]_{\leq \xi}$ is the $\overline{(\cdot)}$ -invariant element of the form

$$\left(t^{a'} F_t^{(i)}[p, s]_{\leq \xi} \cdot F_t^{(i)}(p, s)_{\leq \xi} + t^{b'} \prod_{j \sim i}^{\rightarrow} F_t^{(j)}(p, s)_{\leq \xi} \right) F_t^{(i)}(p, s)_{\leq \xi}^{-1}$$

for some $a', b' \in \frac{1}{2}\mathbb{Z}$ in $\mathbb{F}(\mathcal{Y}_{t,\mathcal{Q}})$. Therefore, taking the following equalities into the account

$$\begin{aligned} F_t^{(i)}[p^+, s^+] &= F_t^{(i)}(p, s), & F_t^{(i)}[p, s^+] &= F_t^{(i)}[p, s], \\ F_t^{(i)}[p^+, s] &= F_t^{(i)}(p, s), & F_t^{(j)}[p^+(j), s^+(j)] &= F_t^{(j)}(p, s), \end{aligned}$$

where $j \sim i$, we conclude $\Phi_{\mathcal{Q}}^T(\tilde{D}(i; p, s)) = F_t^{(i)}[p, s]_{\leq \xi}$ as desired.

Next, we show the bijection between the distinguished bases. It follows from (NDCB1) in Theorem 7.8 and the identity $\Phi_{\mathcal{Q}}^T \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \Phi_{\mathcal{Q}}^T$ that the element $\Phi_{\mathcal{Q}}^T(\tilde{G}(\mathbf{c}, \mathcal{Q}))$ is $\overline{(\cdot)}$ -invariant for any $\mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbb{R}^+}$. Hence, by Theorem 3.7 (see also Remark 5.13) and Theorem 7.8, it remains to show that

$$(8.9) \quad \Phi_{\mathcal{Q}}^T \left(\tilde{F}_t(\mathbf{c}, \mathcal{Q}) \right) = E_t(m(\mathbf{c}))_{\leq \xi}$$

for every $\mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbb{R}^+}$. As a special case of (8.7), we have already proved

$$\Phi_{\mathcal{Q}}^T \left(\tilde{F}_t(\alpha, \mathcal{Q}) \right) = \Phi_{\mathcal{Q}}^T \left(\tilde{D}^{\mathcal{Q}}(\alpha, \alpha^-) \right) = F_t(Y_{i,p})_{\leq \xi}$$

for each $\alpha \in \mathbb{R}^+$, where $(i, p) = \bar{\phi}_{\mathcal{Q}}^{-1}(\alpha, 0)$. Moreover, the ordering of the product in the definition of $\tilde{F}(\mathbf{c}, \mathcal{Q})$ matches the one in the definition of $E_t(m(\mathbf{c}))$ in view of Theorem 7.1. Therefore, we find

$$\Phi_{\mathcal{Q}}^T \left(\tilde{F}(\mathbf{c}, \mathcal{Q}) \right) = t^{a(\mathbf{c})} E_t(m(\mathbf{c}))_{\leq \xi}$$

for some $a(\mathbf{c}) \in \frac{1}{2}\mathbb{Z}$. It suffices to show that $a(\mathbf{c}) = 0$ for every $\mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbb{R}^+}$. By Remark 3.10 and Remark 7.9, we have

$$\begin{aligned} \overline{E_t(m)} &\in E_t(m) + \sum_{m' < m} \mathbb{Z}[t^{\pm 1}] E_t(m'), \\ \overline{\tilde{F}(\mathbf{c}, \mathcal{Q})} &\in \tilde{F}(\mathbf{c}, \mathcal{Q}) + \sum_{\mathbf{c}' <_{\mathcal{Q}} \mathbf{c}} \mathbb{Z}[v^{\pm 1}] \tilde{F}(\mathbf{c}', \mathcal{Q}). \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \overline{\Phi_{\mathcal{Q}}^T \left(\tilde{F}(\mathbf{c}, \mathcal{Q}) \right)} &= \Phi_{\mathcal{Q}}^T \left(\overline{\tilde{F}(\mathbf{c}, \mathcal{Q})} \right) \in t^{a(\mathbf{c})} E_t(m(\mathbf{c}))_{\leq \xi} + \sum_{m' \neq m(\mathbf{c})} \mathbb{Z}[t^{\pm 1/2}] E_t(m')_{\leq \xi}, \\ \overline{\Phi_{\mathcal{Q}}^T \left(\tilde{F}(\mathbf{c}, \mathcal{Q}) \right)} &= \overline{t^{a(\mathbf{c})} E_t(m(\mathbf{c}))_{\leq \xi}} \in t^{-a(\mathbf{c})} E_t(m(\mathbf{c}))_{\leq \xi} + \sum_{m' \neq m(\mathbf{c})} \mathbb{Z}[t^{\pm 1/2}] E_t(m')_{\leq \xi}. \end{aligned}$$

Hence, $a(\mathbf{c}) = -a(\mathbf{c})$, which implies $a(\mathbf{c}) = 0$. \square

Corollary 8.7 (The HLO isomorphism). *For each \mathcal{Q} -datum \mathcal{Q} for \mathfrak{g} , there exists a \mathbb{Z} -algebra isomorphism*

$$\Phi_{\mathcal{Q}}: \mathcal{A}_v[N_-] \rightarrow \mathcal{K}_{t, \mathcal{Q}}$$

such that $(\cdot)_{\leq \xi} \circ \Phi_{\mathcal{Q}} = \Phi_{\mathcal{Q}}^T$. This isomorphism induces a bijection between the bases $\tilde{\mathbf{B}}^*$ and $\{L_t(m) \mid m \in \mathcal{M}_{\mathcal{Q}}\}$.

8.3. Corollaries of the HLO isomorphism. Fix a \mathcal{Q} -datum $\mathcal{Q} = (\Delta, \sigma, \xi)$ for \mathfrak{g} . We have the following corollaries of the HLO isomorphism $\Phi_{\mathcal{Q}}$. Since the proofs can be completely same as those in [26, Section 11.1], we omit them.

Corollary 8.8 (Positivity of the analog of Kazhdan-Lusztig polynomials). *Recall the notation in Theorem 8.6 and Corollary 8.7. We have the following:*

- (1) $\Phi_{\mathcal{Q}} \left(\tilde{F}(\mathbf{c}, \mathcal{Q}) \right) = E_t(m(\mathbf{c}))$ for all $\mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbb{R}^+}$.
- (2) For $\mathbf{c} \in (\mathbb{Z}_{\geq 0})^{\mathbb{R}^+}$ and $m \in \mathcal{M}_{\mathcal{Q}}$, write

$$\tilde{F}(\mathbf{c}, \mathcal{Q}) = \sum_{\mathbf{c}'} p_{\mathbf{c}, \mathbf{c}'}(v) \tilde{G}(\mathbf{c}', \mathcal{Q}), \quad E_t(m) = \sum_{m'} P_{m, m'}(t) L_t(m')$$

with $p_{\mathbf{c}, \mathbf{c}'}(v) \in \mathbb{Z}[v]$ and $P_{m, m'}(t) \in \mathbb{Z}[t]$. Then we have

$$p_{\mathbf{c}, \mathbf{c}'}(t) = P_{m(\mathbf{c}), m(\mathbf{c}')} (t).$$

Moreover, $P_{m(\mathbf{c}), m(\mathbf{c}')} (t) \in \mathbb{Z}_{\geq 0}[t]$.

Corollary 8.9 (Positivity of structure constants). *For $m_1, m_2 \in \mathcal{M}_Q$, write*

$$L_t(m_1)L_t(m_2) = \sum_{m \in \mathcal{M}_Q} c_{m_1, m_2}^m(t) L_t(m).$$

Then we have $c_{m_1, m_2}^m(t) \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$.

Corollary 8.10 (Positivity of the coefficients of the truncated (q, t) -characters). *For $m \in \mathcal{M}_Q$, write*

$$L_t(m)_{\leq \xi} = \sum_{m' : \text{monomial in } \mathcal{Y}_{\leq \xi}} a[m; m'] \underline{m}'$$

with $a[m; m'] \in \mathbb{Z}[t^{\pm 1/2}]$. Then we have $a[m; m'] \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$.

The next corollary gives the affirmative answer to Conjecture 6.6 in \mathcal{C}_Q .

Corollary 8.11 (The (q, t) -characters of Kirillov-Reshetikhin modules). *For a Kirillov-Reshetikhin module $W_{k,p}^{(i)} = L(m_{k,p}^{(i)})$ belonging to the subcategory \mathcal{C}_Q for some Q -datum Q for \mathfrak{g} , we have*

$$(8.10) \quad L_t(m_{k,p}^{(i)}) = F_t(m_{k,p}^{(i)}).$$

In particular, (8.10) holds true whenever $2kd_i \leq rh^\vee$.

Proof. Recall Lemma 4.17 for the last assertion. □

9. R -MATRICES AND COMMUTATION RELATIONS

In this section, we first review some properties of the normalized R -matrices $R_{V,W}(z)$ for $V, W \in \text{Irr } \mathcal{C}$ and the \mathbb{Z} -valued invariants $\mathfrak{d}(V, W)$ measuring the singularities of $R_{V,W}(z)$ introduced in [36]. Then we check that the specialization of the HLO isomorphism Φ_Q for an arbitrary Q coincides with the isomorphism between the Grothendieck rings induced from the generalized quantum affine Schur-Weyl duality functor F_Q studied in [27, 30, 38, 51, 11, 49]. From these results, we deduce several commutation relations in the quantum Grothendieck ring \mathcal{K}_t , which will be important in the sequel.

9.1. R -matrices and related invariants. Let z be an indeterminate. For any $U_q(\mathbf{Lg})$ -module $V \in \mathcal{C}$, we can consider its *affinization* V_z . This is the $\mathbb{k}[z^{\pm 1}]$ -module $V_z := V \otimes_{\mathbb{k}} \mathbb{k}[z^{\pm 1}]$ endowed with a structure of left $U_q(\mathbf{Lg})$ -module by

$$k_i(v \otimes \varphi) = (k_i v) \otimes \varphi, \quad x_{i,k}^{\pm}(v \otimes \varphi) = (x_{i,k}^{\pm} v) \otimes z^k \varphi, \quad h_{i,l}(v \otimes \varphi) = (h_{i,l} v) \otimes z^l \varphi,$$

for $v \in V$ and $\varphi \in \mathbb{k}[z^{\pm 1}]$. For $a \in \mathbb{k}^{\times}$, we have

$$T_a(V) \cong V_z / (z - a)V_z$$

as $U_q(\mathbf{Lg})$ -modules (recall the definition of T_a from Section 2.5).

Let $V, W \in \mathcal{C}$ be simple modules. We fix their ℓ -highest weight vectors $v \in V, w \in W$. It is known that there exists a unique isomorphism of $U_q(\mathbf{Lg}) \otimes_{\mathbb{k}} \mathbb{k}(z)$ -modules

$$R_{V,W}(z): (V_z \otimes W) \otimes_{\mathbb{k}[z^{\pm 1}]} \mathbb{k}(z) \rightarrow (W \otimes V_z) \otimes_{\mathbb{k}[z^{\pm 1}]} \mathbb{k}(z)$$

satisfying $R_{V,W}(z)(v_z \otimes w) = w \otimes v_z$, where $v_z := v \otimes 1 \in V_z$ (see [1, Corollary 2.5]). We call this isomorphism $R_{V,W}(z)$ *the normalized R -matrix* between V and W . Let

$d_{V,W}(z) \in \mathbb{k}[z]$ denote the monic polynomial of the smallest degree such that the image of $d_{V,W}(z)R_{V,W}(z)$ is contained in $W \otimes V_z \subset (W \otimes V_z) \otimes_{\mathbb{k}[z^{\pm 1}]} \mathbb{k}(z)$. This polynomial $d_{V,W}(z)$ is called *the denominator* of $R_{V,W}(z)$. Note that $d_{V,W}(z)$ does not depend on the choice of ℓ -highest weight vectors v and w . Moreover, we have

$$(9.1) \quad d_{T_a(V), T_b(W)}(z) = d_{V,W}((a/b)z)$$

for $a, b \in \mathbb{k}^\times$.

For the case of fundamental modules, we simply write

$$d_{i,j}(z) := d_{L(Y_{i,0}), L(Y_{j,0})}(z)$$

for any $i, j \in I$. It follows that $d_{i,j}(z) = d_{j,i}(z)$ by [1, (A.7)] (see also [13, Lemma 6.4]). The denominators $d_{i,j}(z)$ were computed in [1, 8, 12, 27, 50, 51].⁵

Proposition 9.1 ([13, Section 6.2]). *Let $i, j \in I$ with $d_i \geq d_j$. Unless $d_i = d_j = 1 < r$, we have*

$$d_{i,j}(z) = d_{j,i}(z) = \prod_{u=0}^{rh^\vee} (z - q^{u+d_i})^{\tilde{c}_{ij}(u)}.$$

Even if $d_i = d_j = 1 < r$, the denominator $d_{i,j}(z) (= d_{j,i}(z))$ divides the RHS.

Definition 9.2 (cf. [36, Section 3]). For each pair (V, W) of simple modules in \mathcal{C} , we define the $\mathbb{Z}_{\geq 0}$ -valued invariant $\mathfrak{d}(V, W)$ by

$$\mathfrak{d}(V, W) := \text{zero}_{z=1}(d_{V,W}(z)) + \text{zero}_{z=1}(d_{W,V}(z)),$$

where $\text{zero}_{z=a}(f(z))$ denotes the order of zeros of $f(z)$ at $z = a$ for $f(z) \in \mathbb{k}[z]$ and $a \in \mathbb{k}$. By definition, we have $\mathfrak{d}(V, W) = \mathfrak{d}(W, V)$.

Remark 9.3. In [36], an invariant $\mathfrak{d}(V, W)$ was defined for any *rationaly renormalizable* pair (V, W) of non-zero modules of \mathcal{C} . When both V and W are simple, it coincides with the one in Definition 9.2 thanks to [36, Proposition 3.16].

A simple $U_q(L\mathfrak{g})$ -module $V \in \mathcal{C}$ is said to be *real* if its tensor square $V \otimes V$ is also simple. For instance, every KR module is known to be real (see e.g. the proof of Proposition 6.4 above).

Theorem 9.4 ([29, Section 3.2]). *Let V, W be two simple modules in \mathcal{C} such that at least one of them is real. Then the following three conditions are mutually equivalent:*

- $\mathfrak{d}(V, W) = 0$.
- V and W commute.
- The tensor product $V \otimes W$ is simple.

In particular, we have $\mathfrak{d}(V, V) = 0$ whenever V is a real simple module.

Lemma 9.5. *For $(i, p), (j, s) \in \widehat{I}$, we have*

$$\mathfrak{d}(L(Y_{i,p}), L(Y_{j,s})) \leq \begin{cases} \tilde{c}_{ij}(|p-s| - d_i) & \text{if } |p-s| \leq rh^\vee, \\ 0 & \text{otherwise.} \end{cases}$$

The equality holds unless $d_i = d_j = 1 < r$.

⁵As we specified in the footnote in Section 3.3, our coproduct of $U_q(L\mathfrak{g})$ is opposite to the one in these references.

Proof. When $d_i \geq d_j$, this is a direct consequence of Proposition 9.1 along with (9.1). For the other case $d_i < d_j$, we use Lemma 3.1 (3) & (6). \square

Proposition 9.6. *For any $m_1, m_2 \in \mathcal{M}$, we have*

$$(\text{wt}_{\mathcal{Q}}(m_1), \text{wt}_{\mathcal{Q}}(m_2)) = \sum_{k \in \mathbb{Z}} (-1)^k \mathfrak{d} \left(L(m_1), \mathcal{D}^{k+1} L(m_2) \right).$$

Proof. This is a combination of [36, Proposition 3.22] and [13, Theorem 6.16]. \square

9.2. Generalized quantum affine Schur-Weyl dualities. Let $\mathcal{J} = \{V^x\}_{x \in J}$ be a collection of real simple modules in \mathcal{C} indexed by an arbitrary set J . We consider the graph $\Gamma_{\mathcal{J}}$ whose vertex set is J and such that the number of edges between x and y is equal to $\mathfrak{d}(V^x, V^y)$ for any $x, y \in J$. Note that the graph $\Gamma_{\mathcal{J}}$ has no edge loops by Theorem 9.4.

Given such a collection \mathcal{J} , Kang-Kashiwara-Kim [28] constructed a monoidal functor

$$F_{\mathcal{J}}: \mathcal{A}(\Gamma_{\mathcal{J}}) \rightarrow \mathcal{C}$$

called the *generalized quantum affine Schur-Weyl duality functor*, with some good properties. Here $\mathcal{A}(\Gamma_{\mathcal{J}})$ denotes the category of finite-dimensional modules over the (completed) symmetric quiver Hecke algebras (defined over \mathbb{k}) associated with the graph $\Gamma_{\mathcal{J}}$. This is a \mathbb{k} -linear abelian monoidal category and it gives a categorification of the coordinate algebra of the maximal (pro)unipotent group for the symmetric Kac-Moody algebra associated with $\Gamma_{\mathcal{J}}$. Moreover, the \mathbb{Z} -basis formed by the simple objects in $\mathcal{A}(\Gamma_{\mathcal{J}})$ coincides with the specialization of the dual canonical basis.

Now we fix a \mathcal{Q} -datum \mathcal{Q} for \mathfrak{g} . As an important example of the above construction, we consider the following collection of simple $U_q(L\mathfrak{g})$ -modules indexed by $J = \Delta_0$:

$$(9.2) \quad \mathcal{J}_{\mathcal{Q}} := \{L^{\mathcal{Q}}(\alpha_i)\}_{i \in \Delta_0},$$

where $L^{\mathcal{Q}}(\alpha)$ for $\alpha \in \mathbb{R}^+$ is the fundamental module defined in Section 5.3.

Theorem 9.7 ([27, 30, 38, 51, 11, 49]). *Let \mathcal{Q} be a \mathcal{Q} -datum for \mathfrak{g} . Then we have*

$$(9.3) \quad \mathfrak{d}(L^{\mathcal{Q}}(\alpha_i), L^{\mathcal{Q}}(\alpha_j)) = \delta(i \sim j)$$

for any $i, j \in \Delta_0$, and hence $\Gamma_{\mathcal{J}_{\mathcal{Q}}} = \Delta$. The corresponding functor $F_{\mathcal{J}_{\mathcal{Q}}}$ gives an equivalence of monoidal categories $F_{\mathcal{Q}}: \mathcal{A}(\Delta) \rightarrow \mathcal{C}_{\mathcal{Q}}$. The induced \mathbb{Z} -algebra isomorphism

$$[F_{\mathcal{Q}}]: \text{ev}_{v=1}(\mathcal{A}_v[N_-]) \simeq K(\mathcal{A}(\Delta)) \rightarrow K(\mathcal{C}_{\mathcal{Q}}),$$

is given by

$$\text{ev}_{v=1} \left(\tilde{F}(\alpha, \mathcal{Q}) \right) \mapsto [L^{\mathcal{Q}}(\alpha)]$$

for $\alpha \in \mathbb{R}^+$, which maps the dual canonical basis $\text{ev}_{v=1}(\tilde{\mathbf{B}}^*)$ specialized at $v = 1$ to the basis $\text{Irr } \mathcal{C}_{\mathcal{Q}}$ formed by the simple isomorphism classes in $\mathcal{C}_{\mathcal{Q}}$.

Recall that $\text{ev}_{t=1}(L_t(Y_{i,p})) = \chi_q(L(Y_{i,p}))$ holds for any $(i, p) \in \hat{I}$ by Theorem 3.9 (2). Therefore, as a corollary of Theorem 8.6 and Theorem 9.7, we obtain the following.

Corollary 9.8 (cf. [26, Corollary 11.8]). *The followings hold.*

- (1) *Let $\Phi_{\mathcal{Q}}|_{v=t=1}$ denote the \mathbb{Z} -algebra isomorphism $\text{ev}_{v=1}(\mathcal{A}_v[N_-]) \rightarrow \text{ev}_{t=1}(\mathcal{K}_{t,\mathcal{Q}}) \simeq K(\mathcal{C}_{\mathcal{Q}})$ induced from $\Phi_{\mathcal{Q}}$. Then we have $\Phi_{\mathcal{Q}}|_{v=t=1} = [F_{\mathcal{Q}}]$.*

(2) $\chi_q(L(m)) = \text{ev}_{t=1}(L_t(m))$ for all $m \in \mathcal{M}_{\mathcal{Q}}$.

Corollary 9.8 (2) gives the affirmative answer to Conjecture 3.8 for $\mathcal{C}_{\mathcal{Q}}$.

9.3. Some commutation relations. In this subsection, we deduce several useful commutation relations in \mathcal{K}_t from the results we have obtained in the previous subsections.

Lemma 9.9. *Let \mathcal{Q} be a Q -datum for \mathfrak{g} and $m_1, m_2 \in \mathcal{M}_{\mathcal{Q}}$. Assume that at least one of the simple modules $L(m_1)$ and $L(m_2)$ is real. If $\mathfrak{d}(L(m_1), L(m_2)) = 0$, we have the t -commutation relation*

$$L_t(m_1)L_t(m_2) = t^{\mathcal{N}(m_1, m_2)}L_t(m_2)L_t(m_1)$$

in the quantum Grothendieck ring $\mathcal{K}_t(\mathcal{C}_{\mathcal{Q}})$.

Proof. By Corollary 8.9 we have

$$L_t(m_1)L_t(m_2) = \sum_{m \in \mathcal{M}_{\mathcal{Q}}} c_{m_1, m_2}^m(t)L_t(m).$$

in \mathcal{K}_t with $c_{m_1, m_2}^m(t) \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$. Applying $\text{ev}_{t=1}$ to the both sides yields

$$\chi_q(L(m_1))\chi_q(L(m_2)) = \sum_{m \in \mathcal{M}_{\mathcal{Q}}} c_{m_1, m_2}^m(1)\chi_q(L(m))$$

by Corollary 9.8 (2). On the other hand, we have $L(m_1) \otimes L(m_2) \cong L(m_1 m_2)$ from our assumption and Theorem 9.4. Therefore we obtain $c_{m_1, m_2}^m(1) = \delta(m = m_1 m_2)$, which implies $c_{m_1, m_2}^m(t) = \delta(m = m_1 m_2) \cdot t^a$ for some $a \in \frac{1}{2}\mathbb{Z}$. Similarly, we can prove that $c_{m_2, m_1}^m(t) = \delta(m = m_1 m_2) \cdot t^b$ for some $b \in \frac{1}{2}\mathbb{Z}$. Thus the (q, t) -characters $L_t(m_1)$ and $L_t(m_2)$ commute up to a power of $t^{\pm 1/2}$, i.e. $L_t(m_1)L_t(m_2) = t^c L_t(m_2)L_t(m_1)$ for some $c \in \frac{1}{2}\mathbb{Z}$. Comparing the leading terms, we find $c = \mathcal{N}(m_1, m_2)$. \square

Lemma 9.10. *Let $(i, p), (j, s) \in \widehat{I}$. We have the following relations in \mathcal{K}_t :*

- (1) $L_t(Y_{i,p})L_t(Y_{j,s}) = t^{\mathcal{N}(i,p;j,s)}L_t(Y_{j,s})L_t(Y_{i,p})$ if $\mathfrak{d}(L(Y_{i,p}), L(Y_{j,s})) = 0$,
- (2) $L_t(Y_{i,p})L_t(Y_{i^*, p \pm rh^\vee}) = t^{\mp 1}L_t(Y_{i,p}Y_{i^*, p \pm rh^\vee}) + 1$.

Proof. We may assume $p \leq s$. First we consider the case $p + rh^\vee \leq s$. In this case, Theorem 3.9 (3) implies that the element $L_t(Y_{i,p})L_t(Y_{j,s}) - \delta((i^*, p + rh^\vee) = (j, s))$ contains no dominant monomials other than $t^{\mathcal{N}(i,p;j,s)/2}Y_{i,p}Y_{j,s}$. Hence it is equal to $t^{\mathcal{N}(i,p;j,s)/2}L_t(Y_{i,p}Y_{j,s})$ by Theorem 3.7. This shows (1) for $p + rh^\vee \leq s$ and (2) as well (note that $\mathcal{N}(i, p; i^* \pm rh^\vee) = \mp 2$ by Lemma 3.1). For the other case $p \leq s < p + rh^\vee$, we choose a suitable Q -datum \mathcal{Q} for \mathfrak{g} so that both $Y_{i,p}$ and $Y_{j,s}$ belong to $\mathcal{M}_{\mathcal{Q}}$ (recall Lemma 4.17). Under the assumption $\mathfrak{d}(L(Y_{i,p}), L(Y_{j,s})) = 0$, we obtain the relation (1) by applying Lemma 9.9. \square

Lemma 9.11. *Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a Q -datum for \mathfrak{g} . Let $(i, p), (j, s) \in \widehat{\Delta}_0^\sigma$ be two elements satisfying the condition $\xi_i \geq p \geq s > \xi_j$. Then we have the following equalities:*

- (1) $\mathfrak{d}(L(Y_{i,p}), L(Y_{j,s})) = 0$,
- (2) $(\text{wt}_{\mathcal{Q}}(Y_{i,p}), \text{wt}_{\mathcal{Q}}(Y_{j,s})) = -\mathcal{N}(i, p; j, s) = 0$,
- (3) $F_t(Y_{i,p})F_t(Y_{j,s}) = F_t(Y_{j,s})F_t(Y_{i,p})$,

where $i := \bar{i}$, $j := \bar{j}$.

Proof. Choose $a \in \mathbb{Z}_{\geq 0}$ and $b \in \mathbb{Z}_{> 0}$ so that $p = \xi_i - 2ad_i$ and $s = \xi_j + 2bd_j$. By Lemma 9.5, we have

$$\mathfrak{d}(L(Y_{i,p}), L(Y_{j,s})) = \mathfrak{d}(L(Y_{j,s}), L(Y_{i,p})) \leq \tilde{c}_{ji}(p - s - d_j) = \tilde{c}_{ji}(\xi_i - \xi_j - d_j - 2(ad_i + bd_j)).$$

The RHS is zero thanks to Lemma 8.3. This proves (1).

The first equality in (2) is due to Proposition 5.4. For the second equality, we have

$$\begin{aligned} \mathcal{N}(i, p; j, s) &= \tilde{c}_{ji}(p - s - d_j) - \tilde{c}_{ji}(p - s + d_j) \\ &= \tilde{c}_{ji}(\xi_i - \xi_j - 2ad_i - (2b + 1)d_j) - \tilde{c}_{ji}(\xi_i - \xi_j - 2ad_i - (2b - 1)d_j) \end{aligned}$$

by (3.3). The RHS is zero again by Lemma 8.3.

The last equality (3) is a consequence of (1), (2) and Lemma 9.10 (1). \square

Lemma 9.12 (see also [37]). *Let \mathcal{Q} be a Q -datum for \mathfrak{g} .*

(1) *For any simple roots $\alpha, \beta \in \mathbb{R}^+$ with $\alpha \neq \beta$, we have*

$$\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}L^{\mathcal{Q}}(\beta)) = 0.$$

(2) *For any positive roots $\alpha, \beta \in \mathbb{R}^+$ and $k \in \mathbb{Z}_{\geq 2}$, we have*

$$\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}^k L^{\mathcal{Q}}(\beta)) = 0.$$

Proof. Let $\alpha, \beta \in \mathbb{R}^+$ be distinct simple roots and put $(i, p) := \phi_{\mathcal{Q}}^{-1}(\alpha, 0)$ and $(j, s) := \phi_{\mathcal{Q}}^{-1}(\beta, 1)$. Then we have $L^{\mathcal{Q}}(\alpha) = L(Y_{i,p})$ and $\mathcal{D}L^{\mathcal{Q}}(\beta) = L(Y_{j,s})$ by Lemma 5.10 (3). Note that we have $\xi_i \geq p$ and $s > \xi_j$ by construction. If $p \geq s$, Lemma 9.11 (1) proves the assertion (1). If $p \leq s - rh^{\vee}$, we have $|p - s| > rh^{\vee}$ and hence Lemma 9.5 proves the assertion. Thus we only have to consider the case $s - rh^{\vee} < p < s$. Recall that we have $\mathcal{D}^k L(Y_{j,s}) \cong L(Y_{(j)^{k^*}, s+krh^{\vee}})$ for each $k \in \mathbb{Z}$ from Theorem 2.5. Under the assumption $s - rh^{\vee} < p < s$, we have $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}^{k+1} L^{\mathcal{Q}}(\beta)) = 0$ unless $k \in \{-1, 0\}$ by Lemma 9.5. Combining this fact with Proposition 9.6, we find

$$(\alpha, \beta) = \sum_{k \in \mathbb{Z}} (-1)^k \mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}^{k+1} L^{\mathcal{Q}}(\beta)) = -\mathfrak{d}(L^{\mathcal{Q}}(\alpha), L^{\mathcal{Q}}(\beta)) + \mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}L^{\mathcal{Q}}(\beta)).$$

On the other hand, the equality (9.3) tells that $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), L^{\mathcal{Q}}(\beta)) = -(\alpha, \beta)$ for two distinct simple roots α, β . Therefore we have $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}L^{\mathcal{Q}}(\beta)) = 0$, which proves (1).

Next we shall prove (2). Let $\alpha, \beta \in \mathbb{R}^+$ be two positive roots and $k \in \mathbb{Z}_{\geq 2}$. Putting $(i, p) := \bar{\phi}_{\mathcal{Q}}^{-1}(\alpha, 0)$, $(j, s) := \bar{\phi}_{\mathcal{Q}}^{-1}(\beta, k)$, we have $L^{\mathcal{Q}}(\alpha) = L(Y_{i,p})$ and $\mathcal{D}^k L^{\mathcal{Q}}(\beta) = L(Y_{j,s})$ by Lemma 5.10 (3). Note that we always have $p < s$. If $s - p > rh^{\vee}$, we have $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}^k L^{\mathcal{Q}}(\beta)) = 0$ by Lemma 9.5 as desired. In particular, it covers the case when $k > 2$. Thus we only have to consider the case $k = 2$ with $s - rh^{\vee} \leq p < s$. In this case, we have

- $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}^{l+1} L^{\mathcal{Q}}(\beta)) = 0$ unless $l \in \{0, 1\}$ by Lemma 9.5,
- $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}L^{\mathcal{Q}}(\beta)) = \mathfrak{d}(L(Y_{i,p}), L(Y_{j^*, s-rh^{\vee}})) = 0$ by Lemma 9.11 (1),
- $(\alpha, \beta) = -\mathcal{N}(i, p; j^*, s - rh^{\vee}) = 0$ by Lemma 5.3 and Lemma 9.11 (2).

Combining these facts with Proposition 9.6, we have $\mathfrak{d}(L^{\mathcal{Q}}(\alpha), \mathcal{D}^2 L^{\mathcal{Q}}(\beta)) = 0$. This completes the proof of (2). \square

10. ISOMORPHISMS AMONG QUANTUM GROTHENDIECK RINGS

In this section, we prove the main theorem of this paper. In Section 10.1, we observe that, for each Q-datum \mathcal{Q} , every standard (q, t) -character $E_t(m)$ can be factorized into a product of smaller standard (q, t) -characters $E_t(m_k)$ belonging to $\mathcal{K}_t(\mathcal{D}^k \mathcal{C}_{\mathcal{Q}})$ for $k \in \mathbb{Z}$. In Section 10.2, we give a presentation of the localized quantum Grothendieck ring $\mathcal{K}_t \otimes_{\mathbb{Z}[t^{\pm 1/2}]} \mathbb{Q}(t^{1/2})$. This is a generalization of the presentation obtained in [24, Section 7] for simply-laced cases. Based on this presentation, we construct in Section 10.3 a collection of isomorphisms Ψ among the quantum Grothendieck rings labeled by Q-data and prove that they respect the bases formed by the simple (q, t) -characters. For a suitable choice of Q-data, the isomorphism Ψ relates the quantum Grothendieck ring of simply-laced \mathfrak{g} with that of non-simply-laced \mathfrak{g} , which propagates the known positivity for \mathfrak{g} to the desired positivity for \mathfrak{g} . Therefore we establish the positivities (P1) and (P2) for non-simply-laced cases. This gives an affirmative answer to a long-standing expectation in [18].

10.1. A factorization of standard (q, t) -character. Let $\mathcal{Q} = (\Delta, \sigma, \xi)$ be a Q-datum for \mathfrak{g} . By Corollary 4.16 (2), the commutative monoid \mathcal{M} is naturally isomorphic to the coproduct of the submonoids $\mathfrak{D}^k \mathcal{M}_{\mathcal{Q}}$ with k running over \mathbb{Z} . In other words, each dominant monomial $m \in \mathcal{M}$ has a unique factorization

$$(10.1) \quad m = \prod_{k \in \mathbb{Z}} \mathfrak{D}^k(m_k)$$

such that $m_k \in \mathcal{M}_{\mathcal{Q}}$ for all $k \in \mathbb{Z}$. Here we have $m_k = 1$ for all but finitely many k . Thanks to Lemma 9.11 (1), $L(Y_{i,p})$ and $L(Y_{j,s})$ commute when $Y_{i,p} \in \mathfrak{D}^k \mathcal{M}_{\mathcal{Q}}$ and $Y_{j,s} \in \mathfrak{D}^l \mathcal{M}_{\mathcal{Q}}$ with $k < l$ and $p > s$. Therefore, we have

$$(10.2) \quad M(m) \cong \bigotimes_{k \in \mathbb{Z}}^{\rightarrow} \mathcal{D}^k(M(m_k))$$

as $U_q(L\mathfrak{g})$ -modules.

Here we have an analogous factorization result for its (q, t) -character $E_t(m)$.

Proposition 10.1. *For each dominant weight $m \in \mathcal{M}$, there holds*

$$(10.3) \quad E_t(m) = t^{\nu(m, \mathcal{Q})} \prod_{k \in \mathbb{Z}}^{\rightarrow} \mathfrak{D}_t^k(E_t(m_k))$$

in the quantum Grothendieck ring \mathcal{K}_t , where

$$\nu(m, \mathcal{Q}) := -\frac{1}{2} \sum_{k < l} (-1)^{k+l} (\text{wt}_{\mathcal{Q}}(m_k), \text{wt}_{\mathcal{Q}}(m_l)).$$

Proof. Thanks to Lemma 9.11 (3), we know that $F_t(Y_{i,p})$ and $F_t(Y_{j,s})$ commute when $Y_{i,p} \in \mathfrak{D}^k \mathcal{M}_{\mathcal{Q}}$ and $Y_{j,s} \in \mathfrak{D}^l \mathcal{M}_{\mathcal{Q}}$ with $k < l$ and $p > s$. It implies that the desired equality (10.3) holds up to some power of $t^{\pm 1/2}$. On the other hand, we have

$$(10.4) \quad \tilde{Y}_{i,p} \tilde{Y}_{j,s} = t^{(\text{wt}_{\mathcal{Q}}(Y_{i,p}), \text{wt}_{\mathcal{Q}}(Y_{j,s}))} \tilde{Y}_{j,s} \tilde{Y}_{i,p}$$

whenever $Y_{i,p} \in \mathfrak{D}^k \mathcal{M}_{\mathcal{Q}}$ and $Y_{j,s} \in \mathfrak{D}^l \mathcal{M}_{\mathcal{Q}}$ with $k < l$. This is a consequence of Proposition 5.4 and Lemma 9.11 (2). It implies

$$\underline{m} = t^{\nu(m, \mathcal{Q})} \prod_{k \in \mathbb{Z}}^{\rightarrow} \mathfrak{D}_t^k(\underline{m}_k),$$

which shows that the equality (10.3) exactly holds. \square

Remark 10.2. If we choose a Q-datum \mathcal{Q} as in Lemma 4.17, the above factorization results (10.2) and (10.3) are obtained immediately from the definitions and Proposition 5.4.

10.2. A presentation of quantum Grothendieck ring. In this subsection, we establish a presentation of the *localized* quantum Grothendieck ring

$$\mathcal{K}_{\mathbb{Q}(t^{1/2})} := \mathcal{K}_t \otimes_{\mathbb{Z}[t^{\pm 1/2}]} \mathbb{Q}(t^{1/2}).$$

We naturally regard \mathcal{K}_t as a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra of $\mathcal{K}_{\mathbb{Q}(t^{1/2})}$. The anti-involution $\overline{(\cdot)}$ and the automorphism \mathfrak{D} on \mathcal{K}_t are naturally extended to $\mathcal{K}_{\mathbb{Q}(t^{1/2})}$.

Hereafter, we fix a Q-datum \mathcal{Q} for \mathfrak{g} . We put

$$x_{i,k}^{\mathcal{Q}} := L_t(Y_{i,p}) \quad \text{with } (i, p) = \phi_{\mathcal{Q}}^{-1}(\alpha_i, k)$$

for each $i \in \Delta_0$ and $k \in \mathbb{Z}$. Note that we have

$$x_{i,k}^{\mathcal{Q}} = \mathfrak{D}_t^k(L_t^{\mathcal{Q}}(\alpha_i)) = \mathfrak{D}_t^k(x_{i,0}^{\mathcal{Q}})$$

by Lemma 3.11 and Lemma 5.10 (3).

The HLO isomorphism $\Phi_{\mathcal{Q}}: \mathcal{A}_v[N_-] \rightarrow \mathcal{K}_{t, \mathcal{Q}}$ in Theorem 8.6 induces the isomorphism

$$\mathcal{U}_v(\mathfrak{n}_-) \cong \mathcal{A}_v[N_-] \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathbb{Q}(v^{1/2}) \xrightarrow{\sim} \mathcal{K}_{t, \mathcal{Q}} \otimes_{\mathbb{Z}[t^{\pm 1/2}]} \mathbb{Q}(t^{1/2})$$

which sends the dual Chevalley generator $\tilde{F}(\alpha_i, \mathcal{Q}) = (1 - v^2)f_i$ to the element $x_{i,0}^{\mathcal{Q}}$ for each $i \in \Delta_0$. Thus, we obtain *the quantum Serre relations* satisfied for each $i, j \in \Delta_0$ and $k \in \mathbb{Z}$ (recall (7.1)):

$$(10.5) \quad \begin{cases} x_{i,k}^{\mathcal{Q}} x_{j,k}^{\mathcal{Q}} - x_{j,k}^{\mathcal{Q}} x_{i,k}^{\mathcal{Q}} = 0 & \text{if } i \not\sim j, \\ (x_{i,k}^{\mathcal{Q}})^2 x_{j,k}^{\mathcal{Q}} - (t + t^{-1}) x_{i,k}^{\mathcal{Q}} x_{j,k}^{\mathcal{Q}} x_{i,k}^{\mathcal{Q}} + x_{j,k}^{\mathcal{Q}} (x_{i,k}^{\mathcal{Q}})^2 = 0 & \text{if } i \sim j. \end{cases}$$

In addition, we can prove *the quantum Boson relations* hold

$$(10.6) \quad x_{i,k}^{\mathcal{Q}} x_{j,k+1}^{\mathcal{Q}} = t^{-(\alpha_i, \alpha_j)} x_{j,k+1}^{\mathcal{Q}} x_{i,k}^{\mathcal{Q}} + (1 - t^{-2}) \delta_{i,j},$$

$$(10.7) \quad x_{i,k}^{\mathcal{Q}} x_{j,l}^{\mathcal{Q}} = t^{(-1)^{k+l}(\alpha_i, \alpha_j)} x_{j,l}^{\mathcal{Q}} x_{i,k}^{\mathcal{Q}}$$

for every $i, j \in \Delta_0$ and $k, l \in \mathbb{Z}$ with $l > k + 1$. They are deduced from Lemma 9.10, Lemma 9.12 and the relation (10.4).

Definition 10.3. Let Δ be the Dynkin diagram of the simple Lie algebra \mathfrak{g} of type ADE as before. We consider the $\mathbb{Q}(t^{1/2})$ -algebra $\tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta)$ presented by the set of generators

$$\{y_{i,k} \mid i \in \Delta_0, k \in \mathbb{Z}\}$$

satisfying the following relations

(R1) for $i, j \in \Delta_0$ and $k \in \mathbb{Z}$,

$$\begin{cases} y_{i,k}y_{j,k} - y_{j,k}y_{i,k} = 0 & \text{if } i \not\sim j, \\ y_{i,k}^2y_{j,k} - (t + t^{-1})y_{i,k}y_{j,k}y_{i,k} + y_{j,k}y_{i,k}^2 = 0 & \text{if } i \sim j, \end{cases}$$

(R2) for $i, j \in \Delta_0$ and $k \in \mathbb{Z}$,

$$y_{i,k}y_{j,k+1} = t^{-(\alpha_i, \alpha_j)}y_{j,k+1}y_{i,k} + \delta_{i,j}(1 - t^{-2}),$$

(R3) for $i, j \in \Delta_0$ and $k, l \in \mathbb{Z}$ with $l > k + 1$,

$$y_{i,k}y_{j,l} = t^{(-1)^{k+l}(\alpha_i, \alpha_j)}y_{j,l}y_{i,k}.$$

Theorem 10.4. *For each Q -datum $\mathcal{Q} = (\Delta, \sigma, \xi)$ for \mathfrak{g} , there exists an isomorphism*

$$\Psi_{\mathcal{Q}}: \tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta) \rightarrow \mathcal{K}_{\mathbb{Q}(t^{1/2})}$$

of $\mathbb{Q}(t^{1/2})$ -algebras given by $y_{i,k} \mapsto x_{i,k}^{\mathcal{Q}}$ for $i \in \Delta_0$ and $k \in \mathbb{Z}$.

Proof. We have already checked the existence of the $\mathbb{Q}(t^{1/2})$ -homomorphism $\Psi_{\mathcal{Q}}$ given by the required assignment. Indeed, the relation (10.5) (resp. (10.6), (10.7)) in $\mathcal{K}_{\mathbb{Q}(t^{1/2})}$ corresponds to the defining relation (R1) (resp. (R2), (R3)) in $\tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta)$. It remains to show that $\Psi_{\mathcal{Q}}$ is bijective.

As an abuse of notation, we denote by \mathfrak{D}_t the automorphism of the $\mathbb{Q}(t^{1/2})$ -algebra $\tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta)$ given by $\mathfrak{D}_t(y_{i,k}) := y_{i,k+1}$ for $i \in \Delta_0, k \in \mathbb{Z}$. From the definition, we have $\mathfrak{D}_t \circ \Psi_{\mathcal{Q}} = \Psi_{\mathcal{Q}} \circ \mathfrak{D}_t$.

Note that the $\mathbb{Q}(t^{1/2})$ -subalgebra $\tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}^{(0)}$ of $\tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta)$ generated by $\{y_{i,0}\}_{i \in \Delta_0}$ is naturally isomorphic to $\mathcal{A}_v[N_-] \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathbb{Q}(v^{1/2})$, and the homomorphism $\Psi_{\mathcal{Q}}$ restricts to (the localized version of) the HLO isomorphism $\Phi_{\mathcal{Q}}$. Therefore, for each $m \in \mathcal{M}_{\mathcal{Q}}$, there is a unique element $P_t(m) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}^{(0)}$ such that $\Psi_{\mathcal{Q}}(P_t(m)) = E_t(m)$.

Now let $m \in \mathcal{M}$ be an arbitrary dominant monomial. Using the notation in (10.1), we define an element $P_t(m) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta)$ by

$$P_t(m) := t^{\nu(m, \mathcal{Q})} \prod_{k \in \mathbb{Z}}^{\rightarrow} \mathfrak{D}_t^k(P_t(m_k)).$$

Under the homomorphism $\Psi_{\mathcal{Q}}$, the set $\mathcal{P} := \{P_t(m) \mid m \in \mathcal{M}\}$ bijectively corresponds to the basis $\{E_t(m)\}_{m \in \mathcal{M}}$ of $\mathcal{K}_{\mathbb{Q}(t^{1/2})}$ by Proposition 10.1. In particular, the set \mathcal{P} is linearly independent over $\mathbb{Q}(t^{1/2})$.

On the other hand, using the relations (R2) and (R3), we can rewrite every monomial M in the elements of $\{y_{i,k}\}_{i \in \Delta_0, k \in \mathbb{Z}}$ as a linear combination of the elements of the form $M_{k_1}M_{k_2} \cdots M_{k_s}$ with $M_{k_u} \in \mathfrak{D}_t^{k_u} \tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}^{(0)}$ and $k_1 < k_2 < \cdots < k_s$. Therefore, \mathcal{P} forms a spanning set, and hence a basis of the $\mathbb{Q}(t^{1/2})$ -vector space $\tilde{\mathcal{A}}_{\mathbb{Q}(t^{1/2})}(\Delta)$. Since $\Psi_{\mathcal{Q}}$ induces a bijection between the bases \mathcal{P} and $\{E_t(m)\}_{m \in \mathcal{M}}$, it is an isomorphism. \square

10.3. Isomorphisms among quantum Grothendieck rings. For $s = 1, 2$, let $\mathcal{Q}^{(s)} = (\Delta^{(s)}, \sigma^{(s)}, \xi^{(s)})$ be a \mathbb{Q} -datum for a complex simple Lie algebra $\mathfrak{g}^{(s)}$. We assume that their underlying Dynkin diagrams are the same, i.e., $\Delta^{(1)} = \Delta^{(2)}$. In what follows, we use the notation $X^{(s)}$ for a mathematical object X to suggest that it is associated with $\mathfrak{g}^{(s)}$. For example, we apply it to write $\mathcal{C}_{\mathbb{Z}}^{(s)}$, $\mathcal{K}_t^{(s)}$, $\mathcal{M}^{(s)}$ etc.

From the presentation of the quantum Grothendieck ring $\mathcal{K}_{\mathbb{Q}(t^{1/2})}$ established in Theorem 10.4, we obtain the following.

Corollary 10.5. *There exists a unique $\mathbb{Q}(t^{1/2})$ -algebra isomorphism*

$$\Psi \equiv \Psi(\mathcal{Q}^{(2)}, \mathcal{Q}^{(1)}): \mathcal{K}_{\mathbb{Q}(t^{1/2})}^{(1)} \rightarrow \mathcal{K}_{\mathbb{Q}(t^{1/2})}^{(2)}$$

satisfying the following properties:

- (1) *The restriction of Ψ to $\mathcal{K}_{t, \mathcal{Q}^{(1)}} \subset \mathcal{K}_{\mathbb{Q}(t^{1/2})}^{(1)}$ coincides with the isomorphism*

$$\Phi_{\mathcal{Q}^{(2)}} \circ (\Phi_{\mathcal{Q}^{(1)}})^{-1}: \mathcal{K}_{t, \mathcal{Q}^{(1)}} \rightarrow \mathcal{K}_{t, \mathcal{Q}^{(2)}}.$$

- (2) *We have $\Psi \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \Psi$ and $\Psi \circ \mathfrak{D}_t = \mathfrak{D}_t \circ \Psi$.*

In particular, it satisfies

$$(10.8) \quad \Psi \left(x_{i,k}^{\mathcal{Q}^{(1)}} \right) = x_{i,k}^{\mathcal{Q}^{(2)}} \quad \text{for any } i \in \Delta_0 \text{ and } k \in \mathbb{Z}.$$

Proof. We define $\Psi := \Psi_{\mathcal{Q}^{(2)}} \circ (\Psi_{\mathcal{Q}^{(1)}})^{-1}$, which satisfies the required properties. \square

In what follows, we prove that the above isomorphism Ψ induces a bijection between the bases formed by the simple (q, t) -characters. First, we define the bijection $\psi_0: \mathcal{M}_{\mathcal{Q}^{(1)}} \rightarrow \mathcal{M}_{\mathcal{Q}^{(2)}}$ between the dominant monomials by the property

$$(10.9) \quad \Psi \left(L_t^{(1)}(m) \right) = L_t^{(2)}(\psi_0(m)) \quad \text{for all } m \in \mathcal{M}_{\mathcal{Q}^{(1)}}.$$

We extend this bijection ψ_0 to the bijection $\psi: \mathcal{M}^{(1)} \rightarrow \mathcal{M}^{(2)}$ by setting

$$\psi(m) := \prod_{k \in \mathbb{Z}} \mathfrak{D}^k \psi_0(m_k)$$

for each $m \in \mathcal{M}^{(1)}$, where we factorized m as in (10.1) with respect to the \mathbb{Q} -datum $\mathcal{Q}^{(1)}$. One can easily check from the property (10.8) that $\text{wt}_{\mathcal{Q}^{(1)}} = \text{wt}_{\mathcal{Q}^{(2)}} \circ \psi$ holds.

Theorem 10.6. *The isomorphism $\Psi \equiv \Psi(\mathcal{Q}^{(2)}, \mathcal{Q}^{(1)})$ in Corollary 10.5 induces a bijection between the simple (q, t) -characters. In particular, it gives the isomorphism*

$$\Psi: \mathcal{K}_t^{(1)} \rightarrow \mathcal{K}_t^{(2)}$$

between the quantum Grothendieck ring of $\mathcal{C}_{\mathbb{Z}}^{(1)}$ and that of $\mathcal{C}_{\mathbb{Z}}^{(2)}$.

Proof. It follows from the property $\Psi \circ \overline{(\cdot)} = \overline{(\cdot)} \circ \Psi$ that the element $\Psi(L_t^{(1)}(m))$ is $\overline{(\cdot)}$ -invariant for each $m \in \mathcal{M}^{(1)}$. Hence, in view of Theorem 3.7, it suffices to show

$$(10.10) \quad \Psi \left(E_t^{(1)}(m) \right) - E_t^{(2)}(\psi(m)) \in \sum_{m' \in \mathcal{M}^{(2)}} t\mathbb{Z}[t] E_t^{(2)}(m')$$

for each $m \in \mathcal{M}^{(1)}$. By Proposition 10.1, we have

$$(10.11) \quad \Psi \left(E_t^{(1)}(m) \right) = t^{\nu(\psi(m), \mathcal{Q}^{(2)})} \prod_{k \in \mathbb{Z}}^{\rightarrow} \mathfrak{D}_t^k \left(\Psi \left(E_t^{(1)}(m_k) \right) \right).$$

On the other hand, for any $m' \in \mathcal{M}_{\mathcal{Q}^{(1)}}$, the property (S2) in Theorem 3.7 implies that

$$\begin{aligned} E_t^{(1)}(m') &= L_t^{(1)}(m') + \sum_{m'' \in \mathcal{M}_{\mathcal{Q}^{(1)}}} a_{m', m''}(t) L_t^{(1)}(m''), \\ L_t^{(2)}(\psi_0(m')) &= E_t^{(2)}(\psi_0(m')) + \sum_{m'' \in \mathcal{M}_{\mathcal{Q}^{(2)}}} b_{\psi_0(m'), m''}(t) E_t^{(2)}(m'') \end{aligned}$$

hold with some $a_{m', m''}(t), b_{\psi_0(m'), m''}(t) \in t\mathbb{Z}[t]$. Then, by the property (10.9), we have

$$(10.12) \quad \Psi \left(E_t^{(1)}(m_k) \right) = E_t^{(2)}(\psi_0(m_k)) + \sum_{m' \in \mathcal{M}_{\mathcal{Q}^{(2)}}} c_{m_k, m'}(t) E_t^{(2)}(m')$$

for each $k \in \mathbb{Z}$, where

$$c_{m_k, m'}(t) := a_{m_k, \psi_0^{-1}(m')}(t) + b_{\psi_0(m_k), m'}(t) + \sum_{m'' \in \mathcal{M}_{\mathcal{Q}^{(1)}}} a_{m_k, m''}(t) b_{\psi_0(m''), m'}(t) \in t\mathbb{Z}[t].$$

Note that $\text{wt}_{\mathcal{Q}^{(2)}}(m') = \text{wt}_{\mathcal{Q}^{(2)}}(\psi_0(m_k))$ holds whenever $E_t^{(2)}(m')$ contributes non-trivially to the sum in (10.12) by Proposition 5.2. Thus, the desired property (10.10) follows from (10.11) and (10.12) by using Proposition 10.1 again. \square

As a consequence, we obtain the main result of this paper.

Corollary 10.7. *For any finite-dimensional complex simple Lie algebra \mathfrak{g} , the quantum Grothendieck ring \mathcal{K}_t has non-negative structure constants with respect to the simple (q, t) -characters $\{L_t(m)\}_{m \in \mathcal{M}}$. More precisely, if we write*

$$L_t(m_1)L_t(m_2) = \sum_{m \in \mathcal{M}} c_{m_1, m_2}^m(t) L_t(m)$$

for $m_1, m_2 \in \mathcal{M}$, we have

$$c_{m_1, m_2}^m(t) \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}].$$

In particular, the analogs of Kazhdan-Lusztig polynomials are positive, i.e.,

$$P_{m, m'}(t) \in t\mathbb{Z}_{\geq 0}[t]$$

for any $m, m' \in \mathcal{M}$ with $m' < m$ (cf. (3.7)).

Proof. When \mathfrak{g} is simply-laced, the desired positivity has been established in [47] and [55] based on the geometry of quiver varieties (see also [24, Section 5.9]). Now let \mathfrak{g} be of non-simply-laced type and (Δ, σ) denote its unfolding ($\sigma \neq \text{id}$). We choose two arbitrary Q-data of the forms $\mathcal{Q}^{(1)} = (\Delta, \text{id}, \xi^{(1)})$ and $\mathcal{Q}^{(2)} = (\Delta, \sigma, \xi^{(2)})$ so that we have $\mathfrak{g}^{(1)} = \mathfrak{g}$ (simply-laced) and $\mathfrak{g}^{(2)} = \mathfrak{g}$. Since the corresponding isomorphism $\Psi = \Psi(\mathcal{Q}^{(2)}, \mathcal{Q}^{(1)}): \mathcal{K}_t^{(1)} \rightarrow \mathcal{K}_t^{(2)}$ respects the simple (q, t) -characters (Theorem 10.6), it transforms the known positivity in $\mathcal{K}_t^{(1)}$ to the desired positivity in $\mathcal{K}_t^{(2)}$. \square

11. FURTHER RESULTS IN TYPE B

In this section, we focus on the case when \mathfrak{g} is of type B_n and hence \mathfrak{g} is of type A_{2n-1} . We show that the specialization at $t = 1$ of the isomorphism Ψ for a specific choice of Q-data coincides with the Grothendieck ring isomorphism arising from a categorical relation studied in [34]. As a consequence, we verify the analog of Kazhdan-Lusztig conjecture (Conjecture 3.8) and the desired positivity of the coefficients of simple (q, t) -characters for any simple $U_q(L\mathfrak{g})$ -modules in $\mathcal{C}_{\mathbb{Z}}$ of type B_n . Hence, we have solved the long-standing problem of determining from their highest ℓ -weight the character of simple finite-dimensional modules of type B quantum loop algebras, with a Kazhdan-Lusztig algorithm which is uniform relatively to simply-laced types (see the Introduction for more comments and references).

11.1. A comparison between type A and B. Let Δ be the Dynkin diagram of type A_{2n-1} for a fixed $n \in \mathbb{Z}_{\geq 2}$. We will use the labeling $\Delta_0 = \{1, 2, \dots, 2n-1\}$ as in Figure 1. In this subsection, as a special case of the previous subsection, we consider the following two Q-data $\mathcal{Q}^{(1)} = (\Delta, \text{id}, \xi^{(1)})$ for $\mathfrak{g}^{(1)}$ of type A_{2n-1} and $\mathcal{Q}^{(2)} = (\Delta, \vee, \xi^{(2)})$ for $\mathfrak{g}^{(2)}$ of type B_n , where

$$\xi_i^{(1)} := -\iota,$$


$$\xi_i^{(2)} := \begin{cases} -2\iota & \text{if } 1 \leq \iota < n, \\ 1 - 2n & \text{if } \iota = n, \\ 2 - 2\iota & \text{if } n < \iota \leq 2n - 1 \end{cases}$$

for $\iota \in \Delta_0 = \{1, 2, \dots, 2n-1\}$. Note that we have

$$\widehat{I}_{\mathcal{Q}^{(1)}} = (\Gamma_{\mathcal{Q}^{(1)}})_0 = \{(\iota, -\iota - 2k) \in \Delta_0 \times \mathbb{Z} \mid 0 \leq k \leq 2n - 1 - \iota\}$$

by Proposition 4.15.

Example 11.1. When $n = 3$, the twisted Auslander-Reiten quivers $\Gamma_{\mathcal{Q}^{(1)}}$ and $\Gamma_{\mathcal{Q}^{(2)}}$ are depicted respectively as follows:



By Theorem 10.6, we have the corresponding isomorphism

$$\Psi \equiv \Psi(\mathcal{Q}^{(2)}, \mathcal{Q}^{(1)}): \mathcal{K}_t^{(1)} \rightarrow \mathcal{K}_t^{(2)}$$

between the quantum Grothendieck ring of type A_{2n-1} and that of type B_n . By the construction, the restriction of Ψ to the subcategory $\mathcal{C}_{\mathcal{Q}^{(1)}}$ is $\Phi_{\mathcal{Q}^{(2)}} \circ (\Phi_{\mathcal{Q}^{(1)}})^{-1}$, which has been computed in [26]. Thus we have the following.

Theorem 11.2 ([26, Theorem 12.9]). *For any dominant monomial $m \in \mathcal{M}^{(1)}$, we have*

$$\Psi \left(L_t^{(1)}(m) \right) = L_t^{(2)}(\psi(m)),$$

where $\psi: \mathcal{M}^{(1)} \rightarrow \mathcal{M}^{(2)}$ is a bijection satisfying $\mathfrak{D} \circ \psi = \psi \circ \mathfrak{D}$ and

$$\psi(Y_{i,-i-2k}) = \begin{cases} Y_{\bar{i},-2i-4k} & \text{if } i \leq n-1 \text{ and } k \leq n-i-1, \\ Y_{n,-6n+4i+1} & \text{if } i \leq n \text{ and } k = n-i, \\ Y_{n,2n+1-4i-4k} Y_{n,-2n+3-4k} & \text{if } 2 \leq i \leq n \text{ and } n-i+1 \leq k \leq n-1, \\ Y_{n,-2n+3-4i} & \text{if } i \leq n-1 \text{ and } k = n, \\ Y_{\bar{i},4-2i-4k} & \text{if } i \leq n-2 \text{ and } n+1 \leq k, \\ Y_{\bar{i},2-2i-4k} & \text{if } i \geq n+1 \end{cases}$$

for each $(i, -i-2k) \in \widehat{I}_{\mathcal{Q}^{(1)}}$.

On the other hand, based on the generalized Schur-Weyl duality, Kashiwara-Kim-Oh [34] have obtained the following result.

Theorem 11.3 ([34, Corollary 3.2.2]). *There is an isomorphism $\tilde{\psi}: K(\mathcal{C}_{\mathbb{Z}}^{(1)}) \rightarrow K(\mathcal{C}_{\mathbb{Z}}^{(2)})$ such that we have*

$$\tilde{\psi}\left(\left[L^{(1)}(m)\right]\right) = \left[L^{(2)}(\psi(m))\right]$$

for every $m \in \mathcal{M}^{(1)}$, where ψ is the bijection in Theorem 11.2.

Proof. In [34, Section 3.2], an isomorphism $\phi_1: K(\mathcal{C}_{\mathbb{Z}}^{(1)}) \rightarrow K(\mathcal{C}_{\mathbb{Z}}^{(2)})$ is defined. It induces an bijection $\text{Irr } \mathcal{C}_{\mathbb{Z}}^{(1)} \rightarrow \text{Irr } \mathcal{C}_{\mathbb{Z}}^{(2)}$ between isomorphism classes of simple modules. From the explicit computation of ϕ_1 given in [34, Corollary 3.2.2], we see that it satisfies $[\mathfrak{D}] \circ \phi_1 = \phi_1 \circ [\mathfrak{D}]$ and its restriction to the subcategory $\mathcal{C}_{\mathcal{Q}^{(1)}}$ coincides with the homomorphism $[L(Y_{i,p})] \mapsto [L(\psi(Y_{i,p}))]$ after a suitable parameter shift (see [26, Remark 12.10]). Therefore the desired isomorphism $\tilde{\psi}$ is obtained as a parameter shift of ϕ_1 . \square

Recall that an algebra homomorphism $K(\mathcal{C}_{\mathbb{Z}}^{(1)}) \rightarrow K(\mathcal{C}_{\mathbb{Z}}^{(2)})$ is determined by the images of the classes $[L^{(1)}(Y_{i,p})]$ of fundamental modules (Proposition 2.9) and we know $\text{ev}_{t=1}(L_t^{(s)}(Y_{i,p})) = \chi_q(L(Y_{i,p}))$ for all $(i,p) \in \widehat{I}^{(s)}$ and $s = 1, 2$. Therefore, as a corollary of Theorems 11.2 and 11.3, we obtain the following.

Corollary 11.4. *The followings hold.*

- (1) Let $\Psi|_{v=t=1}$ denote the \mathbb{Z} -algebra isomorphism $K(\mathcal{C}_{\mathbb{Z}}^{(1)}) \rightarrow K(\mathcal{C}_{\mathbb{Z}}^{(2)})$ induced from $\Psi \equiv \Psi(\mathcal{Q}^{(2)}, \mathcal{Q}^{(1)})$ above. Then we have $\Psi|_{v=t=1} = \tilde{\psi}$.
- (2) We have $\chi_q(L^{(2)}(m)) = \text{ev}_{t=1}(L_t^{(2)}(m))$ for all $m \in \mathcal{M}^{(2)}$.

Corollary 11.4 (2) gives the affirmative answer to Conjecture 3.8 for every simple module $L(m)$ in $\mathcal{C}_{\mathbb{Z}}$ when \mathfrak{g} is of type B_n .

11.2. Positivity of (q, t) -characters in type B. In this subsection, we assume that \mathfrak{g} is the simple Lie algebra of type B_n . Now we obtain the following result by the same argument as Lemma 9.9 using Corollaries 10.7 and 11.4 (2).

Lemma 11.5. *Let \mathfrak{g} be of type B_n and $m_1, m_2 \in \mathcal{M}$ any dominant monomials. Assume that at least one of the simple modules $L(m_1)$ and $L(m_2)$ is real. If we have $\mathfrak{d}(L(m_1), L(m_2)) = 0$, there holds the t -commutation relation*

$$L_t(m_1 m_2) = t^{-\mathcal{N}(m_1, m_2)/2} L_t(m_1) L_t(m_2) = t^{\mathcal{N}(m_1, m_2)/2} L_t(m_2) L_t(m_1)$$

in the quantum Grothendieck ring \mathcal{K}_t .

Using this lemma, we obtain the affirmative answer to Conjecture 6.6 in type B in the following way. For an interval $[a, b]$ we set $L_t^{(i)}[a, b] := L_t(m^{(i)}[a, b])$, and similarly, we define $L_t^{(i)}(a, b]$, $L_t^{(i)}[a, b)$ and $L_t^{(i)}(a, b)$.

Theorem 11.6. *Let \mathfrak{g} be of type B_n . For any KR module $W^{(i)}[p, s]$ in $\mathcal{C}_{\mathbb{Z}}$ with $(i, p), (i, s) \in \widehat{\Delta}_0^{\sigma}$ and $p < s$, its (q, t) -character $L_t^{(i)}[p, s]$ coincides with $F_t^{(i)}[p, s]$.*

Proof. Set $i := \bar{i}$. We prove the assertion by induction on $s - p \in 2d_i \mathbb{Z}_{\geq 0}$. When $s = p$, the KR module $W^{(i)}[p, p]$ is the fundamental module $L(Y_{i,p})$ and hence $L_t(Y_{i,p}) = F_t(Y_{i,p})$ by Theorem 3.9.

Now we assume that $s - p > 0$ and the equality $L_t^{(j)}[p', s'] = F_t^{(j)}[p', s']$ holds for any $j \in \Delta_0$ and $p', s' \in \mathbb{Z}$ with $s' - p' < s - p$. If we write

$$(11.1) \quad L_t^{(i)}[p, s] L_t^{(i)}(p, s) = \sum_{m \in \mathcal{M}} c_m(t) L_t(m),$$

we have $c_m(t) \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$ by Corollary 10.7. Using Corollary 11.4 (2) and the classical T -system identity (6.2), we specialize (11.1) at $t = 1$ to find

$$(11.2) \quad c_m(1) = \begin{cases} 1 & \text{if } m = m^{(i)}[p, s] m^{(i)}(p, s) \text{ or } M(i; p, s), \\ 0 & \text{otherwise,} \end{cases}$$

where $M(i; p, s) = \prod_{j \sim i} m^{(j)}(p, s)$ as in Theorem 6.8. Note that $c_m(1) = 0$ implies that $c_m(t) = 0$, and $c_m(1) = 1$ implies that $c_m(t) = t^x$ for some $x \in \frac{1}{2}\mathbb{Z}$. Recall the set $\mathcal{M}^+(i; p, s)$ of dominant monomials defined in (6.3). Since the members of the family $\{L(m) \mid m \in \mathcal{M}^+(i; p, s)\}$ mutually commute (see Proposition 6.4), the members of the family $\{L_t(m) \mid m \in \mathcal{M}^+(i; p, s)\}$ mutually t -commute and we have

$$L_t \left(m^{(i)}[p, s] m^{(i)}(p, s) \right) = t^x L_t^{(i)}[p, s] L_t^{(i)}(p, s), \quad L_t(M(i; p, s)) = t^y \prod_{j \sim i}^{\rightarrow} L_t^{(j)}(p, s)$$

for some $x, y \in \frac{1}{2}\mathbb{Z}$ by Lemma 11.5. Here we fix an arbitrary total ordering of the set $\{j \in \Delta_0 \mid j \sim i\}$ to define the ordered product. Combining with (11.2), we obtain

$$(11.3) \quad L_t^{(i)}[p, s] L_t^{(i)}(p, s) = t^a L_t^{(i)}[p, s] L_t^{(i)}(p, s) + t^b \prod_{j \sim i}^{\rightarrow} L_t^{(j)}(p, s),$$

where the exponents $a, b \in \frac{1}{2}\mathbb{Z}$ are determined uniquely from the fact that $L_t^{(i)}[p, s]$ is the $\overline{(\cdot)}$ -invariant element of the form

$$\left(t^a L_t^{(i)}[p, s] L_t^{(i)}(p, s) + t^b \prod_{j \sim i}^{\rightarrow} L_t^{(j)}(p, s) \right) L_t^{(i)}(p, s)^{-1}$$

in the fraction field of the quantum torus \mathcal{Y}_t . In other words, we have obtained a quantum T -system identity satisfied by the (q, t) -characters of KR modules.

On the other hands, we also know the quantum T -system identity (6.10) satisfied by the corresponding $F_t(m)$'s (Theorem 6.8). The exponents $a, b \in \frac{1}{2}\mathbb{Z}$ appearing there are exactly the same as in (11.3) because the family $\{F_t(m) \mid m \in \mathcal{M}^+(i; s, p)\}$ satisfies the same t -commutation relations as the family $\{L_t(m) \mid m \in \mathcal{M}^+(i; p, s)\}$ does (see Proposition 6.10).

Now, applying the induction hypothesis to (11.3), we get

$$F_t^{(i)}[p, s]F_t^{(i)}(p, s) = t^a L_t^{(i)}[p, s]F_t^{(i)}(p, s) + t^b \prod_{j \sim i}^{\rightarrow} F_t^{(j)}(p, s)$$

with the same $a, b \in \frac{1}{2}\mathbb{Z}$ as above. Comparing this equality with (6.10), we obtain the desired conclusion $L_t^{(i)}[s, p] = F_t^{(i)}[s, p]$. \square

Theorem 11.7 (Positivity of coefficients of simple (q, t) -characters in type B). *Let \mathfrak{g} be of type B_n . For each dominant monomial $m \in \mathcal{M}$, write*

$$L_t(m) = \sum_{m' : \text{monomial in } \mathcal{Y}} a[m; m'] \underline{m}'$$

with $a[m; m'] \in \mathbb{Z}[t^{\pm 1/2}]$. Then we have

$$a[m; m'] \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}].$$

Proof. Given $m \in \mathcal{M}$, we choose integers $c < b$ so that the (q, t) -character $L_t(m)$ is a (non-commutative) Laurent polynomial in the variables $Y_{i,p}$ with $i \in I$ and $c \leq p \leq b$. Set $\widehat{\Delta}_0^\sigma[c, b] := \widehat{\Delta}_0^\sigma \cap (\Delta_0 \times [c, b])$. By Lemma 6.7 and Theorem 11.6, we have

$$L_t^{(i)}[p, b]_{\leq b} = F_t^{(i)}[p, b]_{\leq b} = \underline{m}^{(i)}[p, b]$$

for each $(i, p) \in \widehat{\Delta}_0^\sigma[c, b]$. Thus, choosing an arbitrary total ordering of the finite set $\widehat{\Delta}_0^\sigma[c, b]$, we have

$$L_t(m)_{\leq b} = L_t(m) = \sum_{m'} t^{x(m')} a[m; m'] \prod_{(i,p) \in \widehat{\Delta}_0^\sigma[c,b]}^{\rightarrow} \left(L_t^{(i)}[p, b]_{\leq b} \right)^{v_{i,p}(m')}$$

for some $x(m') \in \frac{1}{2}\mathbb{Z}$, where $v_{i,p}(m') := u_{\bar{i},p}(m') - u_{\bar{i},p+2d_{\bar{i}}}(m') \in \mathbb{Z}$. Let $l \in \mathbb{Z}_{>0}$ large enough so that $v_{i,p}(m') + l \geq 0$ holds for all $(i, p) \in \widehat{\Delta}_0^\sigma[c, b]$ and for all monomials m' with $a[m; m'] \neq 0$, and put $\mathcal{L} := \prod_{(i,p) \in \widehat{\Delta}_0^\sigma[c,b]}^{\rightarrow} (L_t^{(i)}[p, b]_{\leq b})^l$. Taking the injectivity of $(\cdot)_{\leq b}$ into account, we obtain the following equality in \mathcal{K}_t :

$$L_t(m) \cdot \mathcal{L} = \sum_{m'} t^{x'(m')} a[m; m'] L_t \left(\prod_{(i,p) \in \widehat{\Delta}_0^\sigma[c,b]} m^{(i)}[p, b]^{v_{i,p}(m') + l} \right)$$

for some $x'(m') \in \frac{1}{2}\mathbb{Z}$. By the positivity of structure constants of \mathcal{K}_t (Corollary 10.7), we obtain $t^{x'(m')} a[m; m'] \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$ and hence $a[m; m'] \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$. \square

For general \mathfrak{g} , once one proves Conjecture 6.6, the positivity of any simple (q, t) -characters (the analog of Theorem 11.7) follows by the same argument as above. We plan to pursue this direction in a future work.

11.3. Thin representations. A finite-dimensional $U_q(L\mathfrak{g})$ -module V is said to be *thin* if all ℓ -weight spaces are of dimension 1. As a corollary of Corollary 11.4 (2) and Theorem 11.7, we can conclude that the (q, t) -characters of simple thin modules are the canonical lifts of their q -characters:

Corollary 11.8. *Let \mathfrak{g} be of type B_n , and $m \in \mathcal{M}$ be a dominant monomial such that $L(m)$ is a thin $U_q(L\mathfrak{g})$ -module. Let $P(m)$ be a subset set of \mathcal{Y} consisting of Laurent monomials satisfying*

$$\chi_q(L(m)) = \sum_{m' \in P(m)} m'.$$

Then we have

$$L_t(m) = \sum_{m' \in P(m)} \underline{m'}.$$

Proof. By Theorem 11.7, we have

$$L_t(m) = \sum_{m' : \text{monomial in } \mathcal{Y}} a[m; m'](t) \underline{m'}$$

for some $a[m; m'](t) \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$. Moreover, by Corollary 11.4 (2), we have

$$\sum_{m' : \text{monomial in } \mathcal{Y}} a[m; m'](1) m' = \sum_{m' \in P(m)} m'$$

Therefore,

$$a[m; m'](t) = \begin{cases} t^{c(m; m')}, & \text{for some } c(m; m') \in \frac{1}{2}\mathbb{Z} & \text{if } m' \in P(m), \\ 0 & & \text{otherwise.} \end{cases}$$

Moreover, since $L_t(m)$ is $(\overline{\cdot})$ -invariant, we have $c(m; m') = 0$ for all $m' \in P(m)$, which completes the proof. \square

The minimal affinizations were defined by Chari [4] as generalizations of type A evaluation representations and of Kirillov-Reshetikhin modules. By [21, Theorem 3.10], all minimal affinizations are thin when \mathfrak{g} is of type B_n . Therefore Corollary 11.8 is a vast generalization of [26, Corollary 11.5].

APPENDIX A. QUANTUM UNIPOTENT MINORS

In this section, we briefly review the definitions of the quantum coordinate algebra $\mathcal{A}_v[N_-]$ and the normalized quantum unipotent minors associated with a simply-laced Lie algebra \mathfrak{g} . We will use the notation in Section 4.2.

A.1. Quantum coordinate algebra $\mathcal{A}_v[N_-]$. Let v be an indeterminate with a formal square root $v^{1/2}$. The quantized enveloping algebra $\mathcal{U}_v(\mathfrak{g})$ of \mathfrak{g} (over $\mathbb{Q}(v^{1/2})$) is the unital associative $\mathbb{Q}(v^{1/2})$ -algebra presented by the set of generators

$$\{e_\iota, f_\iota, K_\iota^{\pm 1} \mid \iota \in \Delta_0\}$$

and the following relations:

- $K_\iota K_\iota^{-1} = K_\iota^{-1} K_\iota = 1$ and $K_\iota K_j = K_j K_\iota$ for $\iota, j \in \Delta_0$,
- $K_\iota e_j = v^{(\alpha_\iota, \alpha_j)} e_j K_\iota$, $K_\iota f_j = v^{-(\alpha_\iota, \alpha_j)} f_j K_\iota$ for $\iota, j \in \Delta_0$,
- $(v - v^{-1})[e_\iota, f_j] = \delta_{\iota, j}(K_\iota - K_\iota^{-1})$ for $\iota, j \in \Delta_0$,
- $e_\iota^2 e_j - (v + v^{-1})e_\iota e_j e_\iota - e_j e_\iota^2 = f_\iota^2 f_j - (v + v^{-1})f_\iota f_j f_\iota - f_j f_\iota^2 = 0$ if $\iota \sim j$,
- $[e_\iota, e_j] = [f_\iota, f_j] = 0$ if $\iota \not\sim j$.

For each generator, we set

$$\text{wt}(e_\iota) = \alpha_\iota, \quad \text{wt}(f_\iota) = -\alpha_\iota, \quad \text{wt}(K_\iota^{\pm 1}) = 0,$$

which endows $\mathcal{U}_v(\mathfrak{g})$ with a structure of \mathbb{Q} -graded $\mathbb{Q}(v^{1/2})$ -algebra.

We denote by $\mathcal{U}_v(\mathfrak{n}_-)$ the $\mathbb{Q}(v^{1/2})$ -subalgebra of $\mathcal{U}_v(\mathfrak{g})$ generated by $\{f_\iota\}_{\iota \in \Delta_0}$. This is regarded as a quantized enveloping algebra of the nilpotent Lie subalgebra $\mathfrak{n}_- \subset \mathfrak{g}$ corresponding to the negative roots. As a $\mathbb{Q}(v^{1/2})$ -algebra, $\mathcal{U}_v(\mathfrak{n}_-)$ is presented by the set of generators $\{f_\iota\}_{\iota \in \Delta_0}$ and the quantum Serre relations (7.1).

For $\iota \in \Delta_0$, we define the $\mathbb{Q}(v^{1/2})$ -linear maps e'_ι and ${}_\iota e'$: $\mathcal{U}_v(\mathfrak{n}_-) \rightarrow \mathcal{U}_v(\mathfrak{n}_-)$ by

$$\begin{aligned} e'_\iota(xy) &= e'_\iota(x)y + v^{(\alpha_\iota, \text{wt}(x))} x e'_\iota(y), & e'_\iota(f_j) &= \delta_{\iota, j}, \\ {}_\iota e'(xy) &= v^{(\alpha_\iota, \text{wt}(y))} {}_\iota e'(x)y + x {}_\iota e'(y), & {}_\iota e'(f_j) &= \delta_{\iota, j} \end{aligned}$$

for homogeneous elements $x, y \in \mathcal{U}_v(\mathfrak{n}_-)$. There exists a unique symmetric $\mathbb{Q}(v^{1/2})$ -bilinear form $(\cdot, \cdot)_L$ on $\mathcal{U}_v(\mathfrak{n}_-)$ such that

$$(1, 1)_L = 1, \quad (f_\iota x, y)_L = \frac{1}{1 - v^2} (x, e'_\iota(y))_L, \quad (x f_\iota, y)_L = \frac{1}{1 - v^2} (x, {}_\iota e'(y))_L.$$

Then $(\cdot, \cdot)_L$ is non-degenerate.

Let $\mathcal{U}_v(\mathfrak{n}_-)_\mathbb{Z}$ be the $\mathbb{Z}[v^{\pm 1/2}]$ -subalgebra of $\mathcal{U}_v(\mathfrak{n}_-)$ generated by the divided powers $\{f_\iota^{(m)} := f_\iota^m / [m]_v! \mid \iota \in \Delta_0, m \in \mathbb{Z}_{\geq 0}\}$. Then the quantized coordinate algebra $\mathcal{A}_v[N_-]$ is defined as

$$\mathcal{A}_v[N_-] := \{x \in \mathcal{U}_v(\mathfrak{n}_-) \mid (x, \mathcal{U}_v(\mathfrak{n}_-)_\mathbb{Z})_L \subset \mathbb{Z}[v^{\pm 1/2}]\}.$$

A.2. Quantum unipotent minors. Let V be a $\mathcal{U}_v(\mathfrak{g})$ -module. For $\mu \in \mathbb{P}$, we set

$$V_\mu := \{u \in V \mid K_\iota u = v^{(\alpha_\iota, \mu)} u \text{ for all } \iota \in \Delta_0\}.$$

This is called the *weight space of V of weight μ* . A $\mathcal{U}_v(\mathfrak{g})$ -module with weight space decomposition $V = \bigoplus_{\mu \in \mathbb{P}} V_\mu$ is said to be *integrable* if e_ι and f_ι act locally nilpotently on V for all $\iota \in \Delta_0$.

For each $\lambda \in \mathbb{P}^+ := \sum_{\iota \in \Delta_0} \mathbb{Z}_{\geq 0} \varpi_\iota$, we denote by $V(\lambda)$ the (finite-dimensional) irreducible highest weight $\mathcal{U}_v(\mathfrak{g})$ -module generated by a highest weight vector $u_\lambda \in$

$V(\lambda)_\lambda$. The module $V(\lambda)$ is integrable. There exists a unique $\mathbb{Q}(v^{1/2})$ -bilinear form $(\cdot, \cdot)_\lambda^\varphi: V(\lambda) \times V(\lambda) \rightarrow \mathbb{Q}(v^{1/2})$ such that

$$(u_\lambda, u_\lambda)_\lambda^\varphi = 1, \quad (xu_1, u_2)_\lambda^\varphi = (u_1, \varphi(x)u_2)_\lambda^\varphi$$

for $u_1, u_2 \in V(\lambda)$ and $x \in \mathcal{U}_v(\mathfrak{g})$, where φ is the $\mathbb{Q}(v^{1/2})$ -algebra anti-involution on $\mathcal{U}_v(\mathfrak{g})$ defined by

$$\varphi(e_i) = f_i, \quad \varphi(f_i) = e_i, \quad \varphi(K_i) = K_i$$

for $i \in \Delta_0$. The form $(\cdot, \cdot)_\lambda^\varphi$ is non-degenerate and symmetric.

For $w \in \mathbb{W}$, we define the element $u_{w\lambda} \in V(\lambda)_{w\lambda}$ by

$$u_{w\lambda} = f_{i_1}^{(m_1)} \cdots f_{i_{l-1}}^{(m_{l-1})} f_{i_l}^{(m_l)} \cdot u_\lambda,$$

for $(i_1, \dots, i_l) \in \mathcal{I}(w)$, where $m_k := (\alpha_{i_k}, s_{i_{k+1}} \cdots s_{i_{l-1}} s_{i_l} \lambda) \in \mathbb{Z}_{\geq 0}$. It is known that this element does not depend on the choice of $(i_1, \dots, i_l) \in \mathcal{I}(w)$ and $w \in \mathbb{W}$ (depends only on $w\lambda$). See, for example, [43, Proposition 39.3.7]. Then $(u_{w\lambda}, u_{w\lambda})_\lambda^\varphi = 1$.

For $\lambda \in \mathbb{P}^+$ and $w, w' \in \mathbb{W}$, we define an element $D_{w\lambda, w'\lambda} \in \mathcal{U}_v(\mathfrak{n}_-)$ by the following property:

$$(D_{w\lambda, w'\lambda}, x)_L = (u_{w\lambda}, xu_{w'\lambda})_\lambda^\varphi$$

for $x \in \mathcal{U}_v(\mathfrak{n}_-)$. By the nondegeneracy of the bilinear form $(\cdot, \cdot)_L$, this element is uniquely determined. An element of this form is called a *quantum unipotent minor*. Moreover, we set

$$\tilde{D}_{w\lambda, w'\lambda} := v^{-(w\lambda - w'\lambda, w\lambda - w'\lambda)/4 + (w\lambda - w'\lambda, \rho)/2} D_{w\lambda, w'\lambda},$$

where $\rho := \sum_{i \in \Delta_0} \varpi_i \in \mathbb{P}^+$. This element is called a *normalized quantum unipotent minor*.

APPENDIX B. QUANTUM CLUSTER ALGEBRAS

In this subsection, we briefly review the definition of a quantum cluster algebra following [2].

B.1. Quantum seed. Let v be an indeterminate with its formal square root $v^{1/2}$. Let J be a finite set. For a \mathbb{Z} -valued $J \times J$ -skew-symmetric matrix $\Lambda = (\Lambda_{ij})_{i, j \in J}$, we define the *quantum torus* $\mathcal{T}(\Lambda)$ as the $\mathbb{Z}[v^{\pm 1/2}]$ -algebra presented by the set of generators $\{X_j^{\pm 1} \mid j \in J\}$ and the relations:

- $X_j X_j^{-1} = X_j^{-1} X_j = 1$ for $j \in J$,
- $X_i X_j = v^{\Lambda_{ij}} X_j X_i$ for $i, j \in J$.

For $\mathbf{a} = (a_j)_{j \in J} \in \mathbb{Z}^J$, we write

$$X^{\mathbf{a}} := v^{-\frac{1}{2} \sum_{i < j} a_i a_j \Lambda_{ij}} \prod_{j \in J} X_j^{a_j},$$

where we fixed an arbitrary total ordering $<$ of the set J . Note that the resulting element $X^{\mathbf{a}} \in \mathcal{T}(\Lambda)$ is independent from the choice of the total ordering $<$. Since $\mathcal{T}(\Lambda)$ is an Ore domain, it is embedded in its skew field of fractions $\mathbb{F}(\mathcal{T}(\Lambda))$.

Let $J_f \subset J$ be a subset and set $J_e := J \setminus J_f$. Let $\tilde{B} = (b_{ij})_{i \in J, j \in J_e}$ be a \mathbb{Z} -valued $J \times J_e$ -matrix whose *principal part* $(b_{ij})_{i, j \in J_e}$ is skew-symmetric. Such a matrix \tilde{B} is called an *exchange matrix*.

We say that a pair (Λ, \tilde{B}) is *compatible* if we have

$$\sum_{k \in J} b_{ki} \Lambda_{kj} = d \delta_{i,j} \quad (i \in J_e, j \in J)$$

for some positive integer d .

If (Λ, \tilde{B}) is a compatible pair, the datum $\Sigma = ((X_j)_{j \in J}, \Lambda, \tilde{B})$ is called a *quantum seed*. Then the set $\{X_j\}_{j \in J} \subset \mathbb{F}(\mathcal{T}(\Lambda))$ is called *the quantum cluster of Σ* and each element X_j is called a *quantum cluster variable*.

B.2. Mutations. Given a quantum seed $\Sigma = ((X_j)_{j \in J}, \Lambda, \tilde{B})$ and an element $k \in J_e$, we can associate a new quantum seed $\mu_k(\Sigma)$ as follows.

Define the $J \times J$ -matrix $E = (e_{ij})_{i, j \in J}$ and the $J_e \times J_e$ -matrix $F = (f_{ij})_{i, j \in J_e}$ by

$$e_{ij} := \begin{cases} \delta_{i,j} & \text{if } j \neq k, \\ -1 & \text{if } i = j = k, \\ \max(0, -b_{ik}) & \text{if } i \neq j = k, \end{cases} \quad f_{ij} := \begin{cases} \delta_{i,j} & \text{if } i \neq k, \\ -1 & \text{if } i = j = k, \\ \max(0, -b_{ik}) & \text{if } i = k \neq j. \end{cases}$$

In addition, define $\mathbf{a}' = (a'_j)_{j \in J}$ and $\mathbf{a}'' = (a''_j)_{j \in J}$ by

$$a'_j := \begin{cases} -1 & \text{if } i = k, \\ \max(0, b_{ik}) & \text{if } i \neq k, \end{cases} \quad a''_j := \begin{cases} -1 & \text{if } i = k, \\ \max(0, -b_{ik}) & \text{if } i \neq k, \end{cases}$$

Then the new datum $\mu_k(\Sigma) = ((X'_j)_{j \in J}, \Lambda', \tilde{B}')$ is given by

$$\Lambda' := E^T \Lambda E, \quad \tilde{B}' := E \tilde{B} F, \quad X'_j := \begin{cases} X^{\mathbf{a}'} + X^{\mathbf{a}''} & \text{if } j = k, \\ X_j & \text{if } j \neq k. \end{cases}$$

One can show that the datum $\mu_k(\Sigma)$ actually defines a quantum seed, which is called *the mutation of Σ in direction k* . This operation is involutive, i.e., we have $\mu_k(\mu_k(\Sigma)) = \Sigma$.

Definition B.1. Let (Λ, \tilde{B}) be a compatible pair and $\Sigma = ((X_j)_{j \in J}, \Lambda, \tilde{B})$ the associated quantum seed. *The quantum cluster algebra $\mathcal{A}_v(\Lambda, \tilde{B})$* is the $\mathbb{Z}[v^{\pm 1/2}]$ -subalgebra of the skew field $\mathbb{F}(\mathcal{T}(\Lambda))$ generated by all the quantum cluster variables in the quantum seeds obtained from Σ by any sequence of mutations.

Theorem B.2 (The quantum Laurent phenomenon [2, Corollary 5.2]). *The quantum cluster algebra $\mathcal{A}_v(\Lambda, \tilde{B})$ is contained in the quantum torus $\mathcal{T}(\Lambda)$.*

REFERENCES

- [1] Tatsuya Akasaka and Masaki Kashiwara. Finite-dimensional representations of quantum affine algebras. *Publ. Res. Inst. Math. Sci.*, 33(5):839–867, 1997.
- [2] Arkady Berenstein and Andrei Zelevinsky. Quantum cluster algebras. *Advances in Mathematics*, 195(2):405–455, 2005.
- [3] Léa Bittmann. A quantum cluster algebra approach to representations of simply laced quantum affine algebras. *Math. Z.*, 298(3-4):1449–1485, 2021.

- [4] Vyjayanthi Chari. Minimal affinizations of representations of quantum groups: the rank 2 case. *Publ. Res. Inst. Math. Sci.*, 31(5):873–911, 1995.
- [5] Vyjayanthi Chari and Andrew Pressley. *A guide to quantum groups*. Cambridge University Press, Cambridge, 1994.
- [6] Vyjayanthi Chari and Andrew Pressley. Quantum affine algebras and their representations. In *Representations of groups (Banff, AB, 1994)*, volume 16 of *CMS Conf. Proc.*, pages 59–78. Amer. Math. Soc., Providence, RI, 1995.
- [7] Vyjayanthi Chari and Andrew Pressley. Minimal affinizations of representations of quantum groups: the simply laced case. *J. Algebra*, 184(1):1–30, 1996.
- [8] Etsurō Date and Masato Okado. Calculation of excitation spectra of the spin model related with the vector representation of the quantized affine algebra of type $A_n^{(1)}$. *Internat. J. Modern Phys. A*, 9(3):399–417, 1994.
- [9] Edward Frenkel and Evgeny Mukhin. Combinatorics of q -characters of finite-dimensional representations of quantum affine algebras. *Comm. Math. Phys.*, 216(1):23–57, 2001.
- [10] Edward Frenkel and Nicolai Reshetikhin. Deformations of W -algebras associated to simple Lie algebras. *Comm. Math. Phys.*, 197(1):1–32, 1998.
- [11] Ryo Fujita. Geometric realization of Dynkin quiver type quantum affine Schur-Weyl duality. *Int. Math. Res. Not. IMRN*, (22):8353–8386, 2020.
- [12] Ryo Fujita. Graded quiver varieties and singularities of normalized R -matrices for fundamental modules. *Selecta Math. (N.S.)*, 28(1):Paper No. 2, 45, 2022.
- [13] Ryo Fujita and Se-jin Oh. Q -data and representation theory of untwisted quantum affine algebras. *Comm. Math. Phys.*, 384(2):1351–1407, 2021.
- [14] Christof Geiß, Bernard Leclerc, and Jan Schröer. Cluster structures on quantum coordinate rings. *Selecta Mathematica*, 19(2):337–397, 2013.
- [15] Christof Geiss, Bernard Leclerc, and Jan Schröer. Quivers with relations for symmetrizable Cartan matrices I: Foundations. *Invent. Math.*, 209(1):61–158, 2017.
- [16] Dieter Happel. On the derived category of a finite-dimensional algebra. *Comment. Math. Helv.*, 62(3):339–389, 1987.
- [17] David Hernandez. Algebraic approach to q, t -characters. *Adv. Math.*, 187(1):1–52, 2004.
- [18] David Hernandez. Représentations des algèbres affinisées quantiques : q, t -caractères et produit de fusion. *PhD thesis Université Paris 6*, 2004.
- [19] David Hernandez. Monomials of q and q, t -characters for non simply-laced quantum affinizations. *Math. Z.*, 250(2):443–473, 2005.
- [20] David Hernandez. The Kirillov-Reshetikhin conjecture and solutions of T -systems. *J. Reine Angew. Math.*, 596:63–87, 2006.
- [21] David Hernandez. On minimal affinizations of representations of quantum groups. *Comm. Math. Phys.*, 276(1):221–259, 2007.
- [22] David Hernandez and Michio Jimbo. Asymptotic representations and Drinfeld rational fractions. *Compos. Math.*, 148(5):1593–1623, 2012.
- [23] David Hernandez and Bernard Leclerc. Cluster algebras and quantum affine algebras. *Duke Math. J.*, 154(2):265–341, 2010.
- [24] David Hernandez and Bernard Leclerc. Quantum Grothendieck rings and derived Hall algebras. *J. Reine Angew. Math.*, 701:77–126, 2015.
- [25] David Hernandez and Bernard Leclerc. A cluster algebra approach to q -characters of Kirillov-Reshetikhin modules. *J. Eur. Math. Soc. (JEMS)*, 18(5):1113–1159, 2016.
- [26] David Hernandez and Hironori Oya. Quantum Grothendieck ring isomorphisms, cluster algebras and Kazhdan-Lusztig algorithm. *Adv. Math.*, 347:192–272, 2019.
- [27] Seok-Jin Kang, Masaki Kashiwara, and Myungho Kim. Symmetric quiver Hecke algebras and R -matrices of quantum affine algebras, II. *Duke Math. J.*, 164(8):1549–1602, 2015.
- [28] Seok-Jin Kang, Masaki Kashiwara, and Myungho Kim. Symmetric quiver Hecke algebras and R -matrices of quantum affine algebras. *Invent. Math.*, 211(2):591–685, 2018.
- [29] Seok-Jin Kang, Masaki Kashiwara, Myungho Kim, and Se-jin Oh. Simplicity of heads and socles of tensor products. *Compos. Math.*, 151(2):377–396, 2015.

- [30] Seok-Jin Kang, Masaki Kashiwara, Myungho Kim, and Se-jin Oh. Symmetric quiver Hecke algebras and R -matrices of quantum affine algebras IV. *Selecta Math. (N.S.)*, 22(4):1987–2015, 2016.
- [31] Seok-Jin Kang, Masaki Kashiwara, Myungho Kim, and Se-jin Oh. Monoidal categorification of cluster algebras. *Journal of the American Mathematical Society*, 31(2):349–426, 2018.
- [32] Masaki Kashiwara. On crystal bases of the q -analogue of universal enveloping algebras. *Duke Math. J.*, 63(2):465–516, 1991.
- [33] Masaki Kashiwara. On level-zero representations of quantized affine algebras. *Duke Math. J.*, 112(1):117–175, 2002.
- [34] Masaki Kashiwara, Myungho Kim, and Se-jin Oh. Monoidal categories of modules over quantum affine algebras of type A and B. *Proceedings of the London Mathematical Society*, 118(1):43–77, 2019.
- [35] Masaki Kashiwara, Myungho Kim, Se-jin Oh, and Euiyong Park. Monoidal categories associated with strata of flag manifolds. *Advances in Mathematics*, 328:959–1009, 2018.
- [36] Masaki Kashiwara, Myungho Kim, Se-jin Oh, and Euiyong Park. Monoidal categorification and quantum affine algebras. *Compos. Math.*, 156(5):1039–1077, 2020.
- [37] Masaki Kashiwara, Myungho Kim, Se-jin Oh, and Euiyong Park. PBW theory for quantum affine algebras. to appear in J. Eur. Math. Soc. (preprint [arXiv:2011.14253](https://arxiv.org/abs/2011.14253)), 2020.
- [38] Masaki Kashiwara and Se-jin Oh. Categorical relations between Langlands dual quantum affine algebras: doubly laced types. *J. Algebraic Combin.*, 49(4):401–435, 2019.
- [39] Yoshiyuki Kimura. Quantum unipotent subgroup and dual canonical basis. *Kyoto J. Math.*, 52(2):277–331, 2012.
- [40] Atsuo Kuniba, Tomoki Nakanishi, and Junji Suzuki. T -systems and Y -systems in integrable systems. *J. Phys. A*, 44(10):103001, 146, 2011.
- [41] George Lusztig. Canonical bases arising from quantized enveloping algebras. *Journal of the American Mathematical Society*, 3(2):447–498, 1990.
- [42] George Lusztig. Quivers, perverse sheaves, and quantized enveloping algebras. *J. Amer. Math. Soc.*, 4(2):365–421, 1991.
- [43] George Lusztig. *Introduction to quantum groups*. Modern Birkhäuser Classics. Birkhäuser/Springer, New York, 2010. Reprint of the 1994 edition.
- [44] Hiraku Nakajima. Modules of quantized Coulomb branches. in preparation.
- [45] Hiraku Nakajima. Quiver varieties and finite-dimensional representations of quantum affine algebras. *J. Amer. Math. Soc.*, 14(1):145–238, 2001.
- [46] Hiraku Nakajima. t -analogs of q -characters of Kirillov-Reshetikhin modules of quantum affine algebras. *Represent. Theory*, 7:259–274 (electronic), 2003.
- [47] Hiraku Nakajima. Quiver varieties and t -analogs of q -characters of quantum affine algebras. *Ann. of Math. (2)*, 160(3):1057–1097, 2004.
- [48] Hiraku Nakajima and Alex Weekes. Coulomb branches of quiver gauge theories with symmetrizers. to appear in J. Eur. Math. Soc. (preprint [arXiv:1907.06552](https://arxiv.org/abs/1907.06552)), 2019.
- [49] Katsuyuki Naoi. Equivalence between module categories over quiver Hecke algebras and Hernandez-Leclerc’s categories in general types. *Adv. Math.*, 389:Paper No. 107916, 47, 2021.
- [50] Se-jin Oh. The denominators of normalized R -matrices of types $A_{2n-1}^{(2)}$, $A_{2n}^{(2)}$, $B_n^{(1)}$ and $D_{n+1}^{(2)}$. *Publ. Res. Inst. Math. Sci.*, 51(4):709–744, 2015.
- [51] Se-jin Oh and Travis Scrimshaw. Categorical relations between Langlands dual quantum affine algebras: Exceptional cases. *Comm. Math. Phys.*, 368(1):295–367, 2019.
- [52] Se-jin Oh and Uhi Rinn Suh. Combinatorial Auslander-Reiten quivers and reduced expressions. *J. Korean Math. Soc.*, 56(2):353–385, 2019.
- [53] Se-jin Oh and Uhi Rinn Suh. Twisted and folded Auslander-Reiten quivers and applications to the representation theory of quantum affine algebras. *J. Algebra*, 535(1):53–132, 2019.
- [54] Fan Qin. Triangular bases in quantum cluster algebras and monoidal categorification conjectures. *Duke Math. J.*, 166(12):2337–2442, 2017.
- [55] Michela Varagnolo and Eric Vasserot. Perverse sheaves and quantum Grothendieck rings. In *Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000)*, volume 210 of *Progr. Math.*, pages 345–365. Birkhäuser Boston, Boston, MA, 2003.

(R. Fujita) RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, OIWAKE-KITASHIRAKAWA, SAKYO, KYOTO, 606-8502, JAPAN & INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, UNIVERSITÉ DE PARIS, F-75013, PARIS, FRANCE

Email address: `rfujita@kurims.kyoto-u.ac.jp`

(D. Hernandez) UNIVERSITÉ DE PARIS AND SORBONNE UNIVERSITÉ, CNRS, IMJ-PRG, IUF, F-75006, PARIS, FRANCE

Email address: `david.hernandez@u-paris.fr`

(S.-j. Oh) EWHA WOMANS UNIVERSITY SEOUL, 52 EWHAYEODAE-GIL, DAEHYEON-DONG, SEODAEMUN-GU, SEOUL, SOUTH KOREA

Email address: `sejin092@gmail.com`

(H. Oya) DEPARTMENT OF MATHEMATICAL SCIENCES, SHIBAURA INSTITUTE OF TECHNOLOGY, 307 FUKASAKU, MINUMA-KU, SAITAMA-SHI, SAITAMA, 337-8570, JAPAN

Email address: `hoya@shibaura-it.ac.jp`