

# REPRESENTATIONS OF QUANTUM AFFINIZATIONS AND FUSION PRODUCT

DAVID HERNANDEZ

ABSTRACT. In this paper we study general quantum affinizations  $\mathcal{U}_q(\hat{\mathfrak{g}})$  of symmetrizable quantum Kac-Moody algebras and we develop their representation theory. We prove a triangular decomposition and we give a classification of (type 1) highest weight simple integrable representations analog to Drinfel'd-Chari-Pressley one. A generalization of the  $q$ -characters morphism, introduced by Frenkel-Reshetikhin for quantum affine algebras, appears to be a powerful tool for this investigation. For a large class of quantum affinizations (including quantum affine algebras and quantum toroidal algebras), the combinatorics of  $q$ -characters give a ring structure  $*$  on the Grothendieck group  $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  of the integrable representations that we classified. We propose a new construction of tensor products in a larger category by using the Drinfel'd new coproduct (it can not directly be used for  $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  because it involves infinite sums). In particular we prove that  $*$  is a fusion product (a product of representations is a representation).

## CONTENTS

1. Introduction	1
2. Background	3
3. Quantum affinization $\mathcal{U}_q(\hat{\mathfrak{g}})$ and triangular decomposition	5
4. Integrable representations and category $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$	13
5. $q$ -characters	16
6. Drinfel'd new coproduct and fusion product	23
References	31

## 1. INTRODUCTION


In this paper  $q \in \mathbb{C}^*$  is not a root of unity.

V.G. Drinfel'd [Dr1] and M. Jimbo [Jim] associated, independently, to any symmetrizable Kac-Moody algebra  $\mathfrak{g}$  and  $q \in \mathbb{C}^*$  a Hopf algebra  $\mathcal{U}_q(\mathfrak{g})$  called quantum Kac-Moody algebra. The structure of the Grothendieck ring of integrable representations is well understood : it is analogous to the classical case  $q = 1$ .

The quantum algebras of finite type  $\mathcal{U}_q(\mathfrak{g})$  ( $\mathfrak{g}$  of finite type) have been intensively studied (see for example [CP4, L, R] and references therein). The quantum affine algebras  $\mathcal{U}_q(\hat{\mathfrak{g}})$  ( $\hat{\mathfrak{g}}$  affine algebra) are also of particular interest : they have two realizations, the usual Drinfel'd-Jimbo realization and a new realization (see [Dr2, Be]) as a quantum affinization of a quantum algebra of finite type  $\mathcal{U}_q(\mathfrak{g})$ . The finite dimensional representations of quantum affine algebras are the subject of intense research (see among others [AK, CP1, CP3, CP4, EM, FR, FM, N1, N2, VV2] and references therein). In particular they were classified by Chari-Pressley [CP3, CP4], and Frenkel-Reshetikhin [FR] introduced the  $q$ -characters morphism which is a powerful tool for the study of these representations (see also [Kn, FM]).

The quantum affinization process (that Drinfel'd [Dr2] described for constructing the second realization of a quantum affine algebra) can be extended to all symmetrizable quantum Kac-Moody algebras  $\mathcal{U}_q(\mathfrak{g})$  (see

[Jin, N1]). One obtains a new class of algebras called quantum affinizations : the quantum affinization of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  is denoted by  $\mathcal{U}_q(\hat{\mathfrak{g}})$ . The quantum affine algebras are the simplest examples and are very special because they are also quantum Kac-Moody algebras. When  $C$  is affine, the quantum affinization  $\mathcal{U}_q(\hat{\mathfrak{g}})$  is called a quantum toroidal algebra. It is known not to be a quantum Kac-Moody algebra but it is also of particular interest (see for example [GKV, M1, M2, N1, N3, Sa, Sc, STU, TU, VV1] and references therein). This setting is summed up in this picture :



(QKMA : Quantum Kac-Moody Algebras, QFA : Quantum Algebras of Finite type, QAA : Quantum Affine Algebras, QTA : Quantum Toroidal Algebras, QA : Quantum Affinizations; the line between the two QAA symbolizes the Drinfel'd-Beck correspondence.)

In [N1] Nakajima gave a classification of (type 1) simple integrable highest weight modules of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  when  $\mathfrak{g}$  is symmetric. The case  $C$  of type  $A_n^{(1)}$  (toroidal  $\hat{sl}_n$ -case) was also studied by Miki in [M1] (a coproduct is also used with an approach specific to the  $A_n^{(1)}$ -case; but it is technically different from the general construction proposed in this paper). In [H3] we proposed a combinatorial construction of  $q$ -characters (and also of their  $t$ -deformations) for generalized Cartan matrix  $C$  such that  $i \neq j \Rightarrow C_{i,j}C_{j,i} \leq 3$  (it includes finite and affine types except  $A_1^{(1)}, A_2^{(2)}$ ); we conjectured that they were linked with a general representation theory. But in general little is known about the representation theory outside the case of quantum affine algebras.

In this paper we study general quantum affinizations and we develop their representation theory. First we prove a triangular decomposition of  $\mathcal{U}_q(\hat{\mathfrak{g}})$ . We classify the (type 1) simple highest weight integrable representations, we define and study a generalization of the morphism of  $q$ -characters  $\chi_q$  which appears to be a natural tool for this investigation (the approach is different from [H3] because  $q$ -characters are obtained from the representation theory and not from purely combinatorial constructions). If the quantized Cartan matrix  $C(z)$  is invertible (it includes all quantum affine algebras and quantum toroidal algebras), a symmetry property of those  $q$ -characters with respect to the action of screening operators is proved (analog of the invariance for the action of the Weyl group in classical finite cases; the result is proved in [FM] for quantum affine algebras); in particular those  $q$ -characters are the combinatorial

objects considered in [H3]. Moreover we get that the image of  $\chi_q$  is a ring and we can define a formal ring structure on the Grothendieck group. Although quantum affine algebras are Hopf algebras, in general no coproduct has been defined for quantum affinizations (this point was also raised by Nakajima in [N3]). Drinfel'd gave formulas for a new coproduct which can be written for all quantum affinizations. They can not directly be used to define a tensor product of representations because they involve infinite sums. We propose a new construction of tensor products in a larger category with a generalization of the new Drinfel'd coproduct. We define a specialization process which allows us to interpret the ring structure that we defined on the Grothendieck group : we prove that it is a fusion product, that is to say that a product of representations is a representation (see [F] for generalities on fusion rings and physical motivations).

In more details, this paper is organized as follows :

in section 2 we recall backgrounds on quantum Kac-Moody algebras. In section 3 we recall the definition of quantum affinizations and we prove a triangular decomposition (theorem 3.2). Some computations are needed to prove the compatibility with affine quantum Serre relations (section 3.3); note that we get a new proof of a combinatorial identity discovered by Jing (consequence of lemma 3.10). The triangular decomposition is used in section 4.2.1 to define the Verma modules of  $\mathcal{U}_q(\hat{\mathfrak{g}})$ .

In section 4 we recall the classification of (type 1) simple integrable highest weight representations of quantum Kac-Moody algebras, and we prove such a classification for quantum affinizations (theorem 4.9; the proof is analogous to the proof given by Chari-Pressley for quantum affine algebras). The point is to give an adapted definition of a weight which we call a  $l$ -weight : we need a more precise definition than in the case of quantum affine algebras (a  $l$ -weight must be characterized by the action of  $\mathcal{U}_q(\hat{\mathfrak{h}}) \subset \mathcal{U}_q(\hat{\mathfrak{g}})$  on a  $l$ -weight space). We also give the definition of the category  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ .

In section 5 we construct  $q$ -characters of integrable modules in the category  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . New technical points are to be considered (in comparison to quantum affine algebra cases) : we have to add terms of the form  $k_\lambda$  ( $\lambda$  coweight of  $\mathcal{U}_q(\hat{\mathfrak{g}})$ ) for the well-definedness in the general case. The original definition of  $q$ -characters ([FR]) was based on an explicit formula for the universal  $\mathcal{R}$ -matrix. In general no universal  $\mathcal{R}$ -matrix has been defined for a quantum affinization. However  $q$ -characters can be obtained with a piece of the formula of a “ $\mathcal{R}$ -matrix” in the same spirit as the original approach (theorem 5.7). In section 5.5 we prove that the image of  $\chi_q$  is the intersection of the kernels of screening operators (theorem 5.15) in the same spirit as Frenkel-Mukhin [FM] did for quantum affine algebras; new technical points are involved because of the  $k_\lambda$  (we suppose that the quantized Cartan matrix  $C(z)$  is invertible). In particular it unifies this approach with [H3] and enables us to prove some results announced in [H3]. We prove that the image of  $\chi_q$  is a ring. As  $\chi_q$  is injective, we get an induced ring structure  $*$  on the Grothendieck group.

In section 6 we prove that  $*$  is a fusion product (theorem 6.2), that is to say that there is a product of modules. We use the new Drinfel'd coproduct (proposition 6.3); as it involves infinite sums, we have to work in a larger category where the tensor product is well-defined (theorem 6.7). To conclude the proof of theorem 6.2 we define specializations of certain forms which allow us to go from the larger category to  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  (section 6.5.2). We also give some concrete examples of explicit computations in section 6.6.

**Acknowledgments** : the author would like to thank Marc Rosso for his continued support and Olivier Schiffmann for his accurate remarks.

## 2. BACKGROUND

**2.1. Cartan matrix.** In this section we give some general backgrounds about Cartan matrices (for more details see [Ka]). A generalized Cartan matrix is  $C = (C_{i,j})_{1 \leq i,j \leq n}$  such that  $C_{i,j} \in \mathbb{Z}$ ,  $C_{i,i} = 2$ ,  $i \neq j \Rightarrow C_{i,j} \leq 0$ ,  $C_{i,j} = 0 \Leftrightarrow C_{j,i} = 0$ . We denote  $I = \{1, \dots, n\}$  and  $l = \text{rank}(C)$ .

In the following we suppose that  $C$  is symmetrizable, that is to say there is a matrix  $D = \text{diag}(r_1, \dots, r_n)$  ( $r_i \in \mathbb{N}^*$ ) such that  $B = DC$  is symmetric. In particular if  $C$  is symmetric then it is symmetrizable with  $D = I_n$ . For example:

$C$  is said to be of finite type if all its principal minors are in  $\mathbb{N}^*$  (see [Bo] for a classification).

$C$  is said to be of affine type if all its proper principal minor are in  $\mathbb{N}^*$  and  $\det(C) = 0$  (see [Ka] for a classification).

Let  $z$  be an indeterminate. We put  $z_i = z^{r_i}$  and for  $l \in \mathbb{Z}$ , we set  $[l]_z = \frac{z^l - z^{-l}}{z - z^{-1}} \in \mathbb{Z}[z^{\pm}]$ . Let  $C(z)$  be the quantized Cartan matrix defined by ( $i \neq j \in I$ ):

$$C_{i,i}(z) = z_i + z_i^{-1}, \quad C_{i,j}(z) = [C_{i,j}]_z$$

In sections 5.5 and 6 we suppose that  $C(z)$  is invertible. We have seen in lemma 6.9 of [H3] that the condition ( $C_{i,j} < -1 \Rightarrow -C_{j,i} \leq r_i$ ) implies that  $\det(C(z)) \neq 0$ . In particular finite and affine Cartan matrices (where we impose  $r_1 = r_2 = 2$  for  $A_1^{(1)}$ ) satisfy this condition and so the quantum affine algebras and quantum toroidal algebra are included in our study. We denote by  $\tilde{C}(z)$  the inverse matrix of  $C(z)$  and  $D(z)$  the diagonal matrix such that for  $i, j \in I$ ,  $D_{i,j}(z) = \delta_{i,j}[r_i]_z$ .

We consider a realization  $(\mathfrak{h}, \Pi, \Pi^\vee)$  of  $C$  (see [Ka]):  $\mathfrak{h}$  is a  $2n - l$  dimensional  $\mathbb{Q}$ -vector space,  $\Pi = \{\alpha_1, \dots, \alpha_n\} \subset \mathfrak{h}^*$  (set of the simple roots) and  $\Pi^\vee = \{\alpha_1^\vee, \dots, \alpha_n^\vee\} \subset \mathfrak{h}$  (set of simple coroots) and for  $1 \leq i, j \leq n$ :

$$\alpha_j(\alpha_i^\vee) = C_{i,j}$$

Denote by  $\Lambda_1, \dots, \Lambda_n \in \mathfrak{h}^*$  (resp. the  $\Lambda_1^\vee, \dots, \Lambda_n^\vee \in \mathfrak{h}$ ) the fundamental weights (resp. coweights): we have  $\alpha_i(\Lambda_j^\vee) = \Lambda_i(\alpha_j^\vee) = \delta_{i,j}$ .

Consider a symmetric bilinear form  $(,): \mathfrak{h}^* \times \mathfrak{h}^* \rightarrow \mathbb{Q}$  such that for  $i \in I$ ,  $h \in \mathfrak{h}^*$ :  $(\alpha_i, h) = h(r_i \alpha_i^\vee)$ . It is non degenerate and gives an isomorphism  $\nu: \mathfrak{h}^* \rightarrow \mathfrak{h}$ . In particular for  $i \in I$  we have  $\nu(\alpha_i) = r_i \alpha_i^\vee$  and for  $\lambda, \mu \in \mathfrak{h}^*$ ,  $\lambda(\nu(\mu)) = \mu(\nu(\lambda))$ .

Denote  $P = \{\lambda \in \mathfrak{h}^* / \forall i \in I, \lambda(\alpha_i^\vee) \in \mathbb{Z}\}$  the set of weights and  $P^+ = \{\lambda \in P / \forall i \in I, \lambda(\alpha_i^\vee) \geq 0\}$  the set of dominant weights. For example we have  $\alpha_1, \dots, \alpha_n \in P$  and  $\Lambda_1, \dots, \Lambda_n \in P^+$ . Denote  $Q = \bigoplus_{i \in I} \mathbb{Z} \alpha_i \subset P$  the root lattice and  $Q^+ = \sum_{i \in I} \mathbb{N} \alpha_i \subset Q$ . For  $\lambda, \mu \in \mathfrak{h}^*$ , write  $\lambda \geq \mu$  if  $\lambda - \mu \in Q^+$ .

If  $C$  is finite we have  $n = l = \dim(\mathfrak{h})$  and for  $\lambda \in \mathfrak{h}^*$ ,  $\lambda = \sum_{i \in I} \alpha_i^\vee(\lambda) \Lambda_i$ . In particular  $\alpha_i = \sum_{j \in I} C_{j,i} \Lambda_j$ . In general the simple roots can not be expressed in terms of the fundamental weights.

## 2.2. Quantum Kac-Moody algebra.

**Definition 2.1.** *The quantum Kac-Moody algebra  $\mathcal{U}_q(\mathfrak{g})$  is the  $\mathbb{C}$ -algebra with generators  $k_h$  ( $h \in \mathfrak{h}$ ),  $x_i^\pm$  ( $i \in I$ ) and relations:*

$$(1) \quad k_h k_{h'} = k_{h+h'}, \quad k_0 = 1$$

$$(2) \quad k_h x_j^\pm k_{-h} = q^{\pm \alpha_j(h)} x_j^\pm$$

$$(3) \quad [x_i^+, x_j^-] = \delta_{i,j} \frac{k_{r_i \alpha_i^\vee} - k_{-r_i \alpha_i^\vee}}{q_i - q_i^{-1}}$$

$$(4) \quad \sum_{r=0 \dots 1-C_{i,j}} (-1)^r \begin{bmatrix} 1 - C_{i,j} \\ r \end{bmatrix}_{q_i} (x_i^\pm)^{1-C_{i,j}-r} x_j^\pm (x_i^\pm)^r = 0 \quad (\text{for } i \neq j)$$

This algebra was introduced independently by Jimbo [Jim] and Drinfel'd [Dr1] and is also called a quantum group. It is remarkable that one can define a Hopf algebra structure on  $\mathcal{U}_q(\mathfrak{g})$  by setting :

$$\Delta(k_h) = k_h \otimes k_h$$

$$\Delta(x_i^+) = x_i^+ \otimes 1 + k_i^+ \otimes x_i^+, \quad \Delta(x_i^-) = x_i^- \otimes k_i^- + 1 \otimes x_i^-$$

$$S(k_h) = k_{-h}, \quad S(x_i^+) = -x_i^+ k_i^{-1}, \quad S(x_i^-) = -k_i^+ x_i^-$$

$$\epsilon(k_h) = 1, \quad \epsilon(x_i^+) = \epsilon(x_i^-) = 0$$

where we use the notation  $k_i^\pm = k_{\pm r_i \alpha_i^\vee}$ .

For  $i \in I$  let  $U_i$  be the subalgebra of  $\mathcal{U}_q(\mathfrak{g})$  generated by the  $x_i^\pm, k_{p\alpha_i^\vee}$  ( $p \in \mathbb{Q}$ ). Then  $U_i$  is isomorphic to  $\mathcal{U}_{q_i}(sl_2)$ , and so a  $\mathcal{U}_q(\mathfrak{g})$ -module has also a structure of  $\mathcal{U}_{q_i}(sl_2)$ -module.

**Definition 2.2.** A triangular decomposition of an algebra  $A$  is the data of three subalgebras  $(A^-, H, A^+)$  of  $A$  such that the multiplication  $x^- \otimes h \otimes x^+ \mapsto x^- h x^+$  defines an isomorphism of  $\mathbb{C}$ -vector space  $A^- \otimes H \otimes A^+ \simeq A$ .

Let  $\mathcal{U}_q(\mathfrak{g})^+$  (resp.  $\mathcal{U}_q(\mathfrak{g})^-, \mathcal{U}_q(\mathfrak{h})$ ) be the subalgebra of  $\mathcal{U}_q(\mathfrak{g})$  generated by the  $x_i^+$  (resp. the  $x_i^-$ , resp. the  $k_h$ ). We have (see [L]) :

**Theorem 2.3.**  $(\mathcal{U}_q(\mathfrak{g})^-, \mathcal{U}_q(\mathfrak{h}), \mathcal{U}_q(\mathfrak{g})^+)$  is a triangular decomposition of  $\mathcal{U}_q(\mathfrak{g})$ . Moreover  $\mathcal{U}_q(\mathfrak{h})$  (resp.  $\mathcal{U}_q(\mathfrak{g})^+, \mathcal{U}_q(\mathfrak{g})^-$ ) is isomorphic to the algebra with generators  $k_h$  (resp  $x_i^+, x_i^-$ ) and relations (1) (resp. relations (4) with  $+$ , relations (4) with  $-$ ).

### 3. QUANTUM AFFINIZATION $\mathcal{U}_q(\hat{\mathfrak{g}})$ AND TRIANGULAR DECOMPOSITION

In this section we define general quantum affinizations (without central charge), we give the relations between the currents (section 3.2) and we prove a triangular decomposition (theorem 3.2).

#### 3.1. Definition.

**Definition 3.1.** The quantum affinization of  $\mathcal{U}_q(\mathfrak{g})$  is the  $\mathbb{C}$ -algebra  $\mathcal{U}_q(\hat{\mathfrak{g}})$  with generators  $x_{i,r}^\pm$  ( $i \in I, r \in \mathbb{Z}$ ),  $k_h$  ( $h \in \mathfrak{h}$ ),  $h_{i,m}$  ( $i \in I, m \in \mathbb{Z} - \{0\}$ ) and the following relations ( $i, j \in I, r, r' \in \mathbb{Z}, m \in \mathbb{Z} - \{0\}$ ):

$$(5) \quad k_h k_{h'} = k_{h+h'}, \quad k_0 = 1, \quad [k_h, h_{j,m}] = 0, \quad [h_{i,m}, h_{j,m'}] = 0$$

$$(6) \quad k_h x_{j,r}^\pm k_{-h} = q^{\pm \alpha_j(h)} x_{j,r}^\pm$$

$$(7) \quad [h_{i,m}, x_{j,r}^\pm] = \pm \frac{1}{m} [m B_{i,j}]_q x_{j,m+r}^\pm$$

$$(8) \quad [x_{i,r}^+, x_{j,r'}^-] = \delta_{ij} \frac{\phi_{i,r+r'}^+ - \phi_{i,r+r'}^-}{q_i - q_i^{-1}}$$

$$(9) \quad x_{i,r+1}^\pm x_{j,r'}^\pm - q^{\pm B_{ij}} x_{j,r'}^\pm x_{i,r+1}^\pm = q^{\pm B_{ij}} x_{i,r}^\pm x_{j,r'+1}^\pm - x_{j,r'+1}^\pm x_{i,r}^\pm$$

$$(10) \quad \sum_{\pi \in \Sigma_s} \sum_{k=0..s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} x_{i,r_{\pi(1)}}^\pm \dots x_{i,r_{\pi(k)}}^\pm x_{j,r'}^\pm x_{i,r_{\pi(k+1)}}^\pm \dots x_{i,r_{\pi(s)}}^\pm = 0$$

where the last relation holds for all  $i \neq j$ ,  $s = 1 - C_{ij}$ , all sequences of integers  $r_1, \dots, r_s$ .  $\Sigma_s$  is the symmetric group on  $s$  letters. For  $i \in I$  and  $m \in \mathbb{Z}$ ,  $\phi_{i,m}^\pm \in \mathcal{U}_q(\hat{\mathfrak{g}})$  is determined by the formal power series in  $\mathcal{U}_q(\hat{\mathfrak{g}})[[z]]$  (resp. in  $\mathcal{U}_q(\hat{\mathfrak{g}})[[z^{-1}]]$ ):

$$\sum_{m \geq 0} \phi_{i,\pm m}^\pm z^{\pm m} = k_{\pm r_i \alpha_i^\vee} \exp(\pm (q - q^{-1}) \sum_{m' \geq 1} h_{i,\pm m'} z^{\pm m'})$$

and  $\phi_{i,m}^+ = 0$  for  $m < 0$ ,  $\phi_{i,m}^- = 0$  for  $m > 0$ .

The relations (10) are called affine quantum Serre relations. The notation  $k_i^\pm = k_{\pm r_i \alpha_i^\vee}$  is also used. We have  $k_i k_i^{-1} = k_i^{-1} k_i = 1$ ,  $k_i x_{j,m}^\pm k_i^{-1} = q^{\pm B_{ij}} x_{j,m}^\pm$ .

There is an algebra morphism  $\mathcal{U}_q(\mathfrak{g}) \rightarrow \mathcal{U}_q(\hat{\mathfrak{g}})$  defined by ( $h \in \mathfrak{h}, i \in I$ )  $k_h \mapsto k_h$ ,  $x_i^\pm \mapsto x_{i,0}^\pm$ . In particular a  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -module has also a structure of a  $\mathcal{U}_q(\mathfrak{g})$ -module.

**3.2. Relations between the currents.** For  $i \in I$ , consider the series (also called currents):

$$x_i^\pm(w) = \sum_{r \in \mathbb{Z}} x_{i,r}^\pm w^r, \phi_i^+(z) = \sum_{m \geq 0} \phi_{i,m}^+ z^m, \phi_i^-(z) = \sum_{m \geq 0} \phi_{i,-m}^- z^{-m}$$

The defining relations of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  can be written with currents ( $h, h' \in \mathfrak{h}$ ,  $i, j \in I$ ):

$$(11) \quad k_h k_{h'} = k_{h+h'}, \quad k_0 = 1, \quad k_h \phi_i^\pm(z) = \phi_i^\pm(z) k_h$$

$$(12) \quad k_h x_j^\pm(z) = q^{\pm \alpha_j(h)} x_j^\pm(z) k_h$$

$$(13) \quad \phi_i^+(z) x_j^\pm(w) = \frac{q^{\pm B_{i,j}} w - z}{w - q^{\pm B_{i,j}} z} x_j^\pm(w) \phi_i^+(z)$$

$$(14) \quad \phi_i^-(z) x_j^\pm(w) = \frac{q^{\pm B_{i,j}} w - z}{w - q^{\pm B_{i,j}} z} x_j^\pm(w) \phi_i^-(z)$$

$$(15) \quad [x_i^+(z), x_j^-(w)] = \frac{\delta_{i,j}}{q_i - q_i^{-1}} \left[ \delta\left(\frac{w}{z}\right) \phi_i^+(w) - \delta\left(\frac{z}{w}\right) \phi_i^-(z) \right]$$

$$(16) \quad (w - q^{\pm B_{i,j}} z) x_i^\pm(z) x_j^\pm(w) = (q^{\pm B_{i,j}} w - z) x_j^\pm(w) x_i^\pm(z)$$

$$(17) \quad \sum_{\pi \in \Sigma_s} \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} x_i^\pm(w_{\pi(1)}) \dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)}) \dots x_i^\pm(w_{\pi(s)}) = 0$$

where  $\delta(z) = \sum_{r \in \mathbb{Z}} z^r$ . The equation (13) (resp. equation (14)) is expanded for  $|z| < |w|$  (resp.  $|w| < |z|$ ).

Remark: in the relations (16), the terms can not be divided by  $w - q^{\pm B_{i,j}} z$ : it would involve infinite sums and make no sense.

The following equivalences are clear : (relations (5)  $\Leftrightarrow$  relations (11)) ; (relations (6)  $\Leftrightarrow$  relations (12)) ; (relations (9)  $\Leftrightarrow$  relations (16)) ; (relations (8)  $\Leftrightarrow$  relations (15)) ; (relations (10)  $\Leftrightarrow$  relations (17)).

We suppose that the relations (6) are verified and we prove the equivalence (relations (7) with  $m \geq 1 \Leftrightarrow$  relations (13)) ((relations (7) with  $m \leq -1 \Leftrightarrow$  relations (14)) is proved in an similar way): consider  $h_i^+(z) = \sum_{m \geq 1} m h_{i,m} z^{m-1}$ . The relation (7) with  $m \geq 1$  are equivalent to (expanded for  $|z| < |w|$ ):

$$[h_i^+(z), x_j^\pm(w)] = \pm [B_{i,j}]_q \frac{w^{-1} x_j^\pm(w)}{\left(1 - \frac{z}{w} q^{B_{i,j}}\right) \left(1 - \frac{z}{w} q^{-B_{i,j}}\right)}$$

It is equivalent to the data of a  $\alpha_\pm(z, w) \in (\mathbb{C}[w, w^{-1}][[z]])$  such that  $\phi_i^+(z) x_j^\pm(w) = \alpha_\pm(z, w) x_j^\pm(w) \phi_i^+(z)$ .

So it suffices to prove that this term is the  $\frac{q^{\pm B_{i,j}} w - z}{w - q^{\pm B_{i,j}} z}$  of relation (13). Let us compute this term : we

have  $\frac{\partial \phi_i^+(z)}{\partial z} = (q - q^{-1}) h_i^+(z) \phi_i^+(z)$  and so the relations (7) imply :

$$\begin{aligned} (q - q^{-1}) \phi_i^+(z) [h_i^+(z), x_j^\pm(w)] &= \frac{\partial \alpha_\pm(z, w)}{\partial z} x_j^\pm(w) \phi_i^+(z) \\ (\pm [B_{i,j}]_q \frac{w^{-1}}{\left(1 - \frac{z}{w} q^{B_{i,j}}\right) \left(1 - \frac{z}{w} q^{-B_{i,j}}\right)} \alpha_\pm(z, w) - \frac{1}{q - q^{-1}} \frac{\partial \alpha_\pm(z, w)}{\partial z}) x_j^\pm(w) \phi_i^+(z) &= 0 \\ \frac{\partial \alpha_\pm(z, w)}{\partial z} &= \pm (q^{B_{i,j}} - q^{-B_{i,j}}) \frac{w^{-1}}{\left(1 - \frac{z}{w} q^{B_{i,j}}\right) \left(1 - \frac{z}{w} q^{-B_{i,j}}\right)} \alpha_\pm(z, w) \end{aligned}$$

As  $\frac{q^{\pm B_{i,j}} w - z}{w - q^{\pm B_{i,j}} z}$  is a solution, we have  $\alpha_\pm(z, w) = \lambda(w) \frac{q^{\pm B_{i,j}} w - z}{w - q^{\pm B_{i,j}} z}$ . But at  $z = 0$  we know  $\alpha_\pm(0, w) = q^{\pm B_{i,j}}$  (relations (6)) and so  $\lambda(w) = 1$ .

**3.3. Triangular decomposition.**

3.3.1. *Statement.* Let  $\mathcal{U}_q(\hat{\mathfrak{g}})^+$  (resp.  $\mathcal{U}_q(\hat{\mathfrak{g}})^-, \mathcal{U}_q(\hat{\mathfrak{h}})$ ) be the subalgebra of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  generated by the  $x_{i,r}^+$  (resp. the  $x_{i,r}^-$ , resp. the  $k_h, h_{i,r}$ ).

**Theorem 3.2.**  $(\mathcal{U}_q(\hat{\mathfrak{g}})^-, \mathcal{U}_q(\hat{\mathfrak{h}}), \mathcal{U}_q(\hat{\mathfrak{g}})^+)$  is a triangular decomposition of  $\mathcal{U}_q(\hat{\mathfrak{g}})$ . Moreover  $\mathcal{U}_q(\hat{\mathfrak{h}})$  (resp.  $\mathcal{U}_q(\hat{\mathfrak{g}})^+, \mathcal{U}_q(\hat{\mathfrak{g}})^-$ ) is isomorphic to the algebra with generators  $k_h, h_{i,m}$  (resp  $x_{i,r}^+, x_{i,r}^-$ ) and relations (5) (resp. relations (9), (10) with +, relations (9), (10) with -).

For a quantum affine algebra ( $C$  finite) it is proved in [Be].

In this section 3.3 we prove this theorem in general. We will use the algebras  $\mathcal{U}_q^l(\hat{\mathfrak{g}}), \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  defined by :

**Definition 3.3.**  $\mathcal{U}_q^l(\hat{\mathfrak{g}})$  is the  $\mathbb{C}$ -algebra with generators  $x_{i,r}^\pm, h_{i,m}, k_h$  ( $i \in I, r \in \mathbb{Z}, m \in \mathbb{Z} - \{0\}, h \in \mathfrak{h}$ ) and relations (5), (6), (7), (8) (or relations (11), (12), (13), (14), (15)).

$\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  is the quotient of  $\mathcal{U}_q^l(\hat{\mathfrak{g}})$  by relations (9) (or relations (16)).

Note that  $\mathcal{U}_q(\hat{\mathfrak{g}})$  is a quotient of  $\mathcal{U}_q^l(\hat{\mathfrak{g}})$  and that  $(\mathcal{U}_q^{l,-}(\hat{\mathfrak{g}}), \mathcal{U}_q(\hat{\mathfrak{h}}), \mathcal{U}_q^{l,+}(\hat{\mathfrak{g}}))$  is a triangular decomposition of  $\mathcal{U}_q^l(\hat{\mathfrak{g}})$  where  $\mathcal{U}_q^{l,\pm}(\hat{\mathfrak{g}})$  is generated by the  $x_{i,r}^\pm$  without relations. In the  $sl_2$ -case we have  $\tilde{\mathcal{U}}_q(\hat{sl}_2) = \mathcal{U}_q(\hat{sl}_2)$ .

Let us sketch the proof of theorem 3.2. We use a method analog to the proof for classic cases or quantum Kac-Moody algebras (see for example the chapter 4 of [Ja]) : we have to check a compatibility condition between the relations and the product as explained in section 3.3.2. After some preliminary technical lemmas about polynomials in section 3.3.3, the heart of the proof is given in section 3.3.4 : properties of  $\mathcal{U}_q^l(\hat{\mathfrak{g}})$  (lemma 3.9) lead to a triangular decomposition of  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ . Properties of  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  proved in lemmas 3.10, 3.11 imply theorem 3.2. Note that the intermediate algebra  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  is also studied because it will be used in the last section of this paper.

Remark : lemma 3.10 gives a new proof of a combinatorial identity discovered by Jing.

The theorem 3.2 is used in section 4.2.1 to define the Verma modules of  $\mathcal{U}_q(\hat{\mathfrak{g}})$ . Let us give another consequence of theorem 3.2 : for  $i \in I$ , let  $\hat{U}_i$  be the subalgebra of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  generated by the  $x_{i,r}^\pm, k_{p\alpha_i^\vee}, h_{i,m}$  ( $r \in \mathbb{Z}, m \in \mathbb{Z} - \{0\}, p \in \mathbb{Q}$ ). We have a morphism  $\mathcal{U}_{q_i}(\hat{sl}_2) \rightarrow \hat{U}_i$  (in particular any  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -module also has a structure of  $\mathcal{U}_{q_i}(\hat{sl}_2)$ -module). Moreover theorem 3.2 implies:

**Corollary 3.4.**  $\hat{U}_i$  is isomorphic to  $\mathcal{U}_{q_i}(\hat{sl}_2)$ .

3.3.2. *General proof of triangular decompositions.* Let  $A$  be an algebra with a triangular decomposition  $(A^-, H, A^+)$ . Let  $B^+$  (resp.  $B^-$ ) be a two-sided ideal of  $A^+$  (resp.  $A^-$ ). Let  $C = A/(A.(B^+ + B^-).A)$  and denote by  $C^\pm$  the image of  $B^\pm$  in  $C$ .

**Lemma 3.5.** If  $B^+.A \subset A.B^+$  and  $A.B^- \subset B^- .A$  then  $(C^-, H, C^+)$  is a triangular decomposition of  $C$  and the algebra  $C^\pm$  is isomorphic to  $A^\pm/B^\pm$ .

*Proof:* We use the proof of section 4.21 in [Ja] : indeed the product gives an isomorphism of  $\mathbb{C}$ -vector space  $A.(B^+ + B^-).A \simeq B^+ \otimes H \otimes A^- + A^+ \otimes H \otimes B^-$ .  $\square$

3.3.3. *Technical lemmas.* Let  $i \neq j$  and  $s = 1 - C_{i,j}$ . Define  $P_\pm(w_1, \dots, w_s, z) \in \mathbb{C}[w_1, \dots, w_s, z]$  by the formula :

$$P_\pm(w_1, \dots, w_s, z) = \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} (w_1 - q^{\pm B_{i,j}} z) \dots (w_k - q^{\pm B_{i,j}} z) (w_{k+1} q^{\pm B_{i,j}} - z) \dots (w_s q^{\pm B_{i,j}} - z)$$

**Lemma 3.6.** There are polynomials  $(f_{\pm,r})_{r=1, \dots, s-1}$  of  $s-1$  variables such that:

$$P_\pm(w_1, \dots, w_s, z) = \sum_{1 \leq r \leq s-1} (w_{r+1} - q_i^{\pm 2} w_r) f_{\pm,r}(w_1, \dots, w_{r-1}, w_{r+2}, \dots, w_s, z)$$

*Proof:* It suffices to prove it for  $P_+$  (because  $P_-$  is obtained from  $P_+$  by  $q \mapsto q^{-1}$ ). First we prove that  $P_+(q_i^{-2(s-1)}w, q_i^{-2(s-2)}w, \dots, q_i^{-2}w, w, z) = 0$ . Indeed it is equal to:

$$\begin{aligned} w^s \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} q_i^{k(1-s)} (q_i^{-2(s-1)} q_i^{s-1} - \frac{z}{w}) \dots (q_i^{-2(s-k)} q_i^{s-1} - \frac{z}{w}) (q_i^{-2(s-k-1)} q_i^{1-s} - \frac{z}{w}) \dots (q_i^{1-s} - \frac{z}{w}) \\ = z^s (q_i^{1-s} - \frac{z}{w}) q^{-3s+3} M_{q_i}(\frac{z}{w} q^{3s-3}) \end{aligned}$$

where:

$$M_q(u) = \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_q q^{k(1-s)} (q^{2k} - u) (q^{2(k+1)} - u) \dots (q^{2(k+s-2)} - u)$$

Let  $\alpha_0(q), \dots, \alpha_{s-1}(q) \in \mathbb{Z}[q]$  such that  $(a-u)(a-uq^2) \dots (a-uq^{2(s-2)}) = u^{s-1} \alpha_{s-1}(q) + u^{s-2} a \alpha_{s-2}(q) + \dots + a^{s-1} \alpha_0(q)$ . So:

$$M_q(u) = \sum_{p=0 \dots s-1} \alpha_{s-p}(q) u^{s-p} \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_q q^{k(1-s+2p)}$$

And so  $M_q(u) = 0$  because of the  $q$ -binomial identity for  $p' = 1-s, 3-s, \dots, s-1$  (see [L]):

$$\sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_q q^{kp'} = 0$$

As a consequence  $P_+$  is in the kernel of the projection

$$\phi : \mathbb{C}[w_1, \dots, w_s, z] \rightarrow \mathbb{C}[w_1, \dots, w_s, z] / ((w_2 - q_i^2 w_1), \dots, (w_s - q_i^2 w_{s-1}))$$

that is to say  $P_+(w_1, \dots, w_s, z) = \sum_{1 \leq r \leq s-1} (w_{r+1} - q^{B_{i,j}} w_r) f_r(w_1, \dots, w_s, z)$  where the  $f_r \in \mathbb{C}[w_1, \dots, w_s, z]$ .

Let us prove that we can choose the  $(f_r)_{1 \leq r \leq s-1}$  so that for all  $1 \leq s \leq r-1$ ,  $f_r$  does not depend of  $w_r, w_{r+1}$ . Let  $\mathcal{A} \subset \text{Ker}(\phi)$  be the subspace of polynomials which are degree at most of 1 in each variable  $w_1, \dots, w_s$ . In particular  $P \in \mathcal{A}$ . We can decompose in a unique way  $P = \alpha + w_2 \beta + w_1 \gamma$  where  $\alpha, \gamma \in \mathbb{C}[w_3, \dots, w_s, z]$ ,  $\beta \in \mathbb{C}[w_1, w_3, \dots, w_s, z]$ . Consider  $\lambda^{(1)} = -q_i^{-2} \gamma (w_2 - q_i^2 w_1) \in \mathcal{A}$  and  $P^{(1)} = P - \lambda^{(1)} \in \mathcal{A}$ . We have in particular  $P^{(1)} = \mu_3^{(1)} + w_2 \mu_2^{(1)} + w_2 w_1 \mu_1^{(1)}$  where  $\mu_1^{(1)}, \mu_2^{(1)}, \mu_3^{(1)} \in \mathbb{C}[w_3, \dots, w_s, z]$ . In the same way we define by induction on  $r$  ( $1 \leq r \leq s-1$ ) the  $\lambda^{(r)} \in \mathcal{A}$  such that  $P^{(r)} = P^{(r-1)} - \lambda^{(r)} \in \mathcal{A}$  is of the form:

$$P^{(r)} = \mu_{r+2}^{(r)} + w_{r+1} \mu_{r+1}^{(r)} + w_{r+1} w_r \mu_r^{(r)} + \dots + w_{r+1} w_r \dots w_1 \mu_1^{(r)}$$

where for  $1 \leq r' \leq r+2$ ,  $\mu_{r'}^{(r)} \in \mathbb{C}[w_{r+2}, \dots, w_s, z]$ . Indeed in the part of  $P^{(r)}$  without  $w_{r+2}$  we can change the terms  $w_{r+1} \lambda(w_{r+3}, \dots, w_s, z)$  to  $q_i^{-2} w_{r+2} \lambda(w_{r+3}, \dots, w_s, z)$  by adding  $q_i^{-2} (w_{r+2} - q_i^2 w_{r+1}) \lambda \in \mathcal{A}$ , we can change the terms  $w_{r+1} w_r \lambda'(w_{r+3}, \dots, w_s, z)$  to  $q_i^{-4} w_{r+2} w_{r+1} \lambda'(w_{r+3}, \dots, w_s, z)$  by adding  $q_i^{-4} (w_{r+2} - q_i^2 w_{r+1}) \lambda + q_i^{-2} (w_{r+2} - q_i^2 w_{r+1}) \lambda \in \mathcal{A}$ , and so on. In particular for  $r = s-1$ :

$$P^{(s-1)} = \mu_{s+1}^{(s-1)} + \mu_s^{(s-1)} w_s + \mu_{s-1}^{(s-1)} w_s w_{s-1} + \dots + \mu_1^{(s-1)} w_s w_{s-1} \dots w_1$$

where  $\mu_{s+1}^{(s-1)}, \dots, \mu_1^{(s-1)} \in \mathbb{C}[z]$ . But :

$$0 = \phi(P^{(s-1)}) = \mu_{s+1}^{(s-1)} + \mu_s^{(s-1)} w_s + \mu_{s-1}^{(s-1)} q_i^{-2} w_s^2 + \dots + \mu_1^{(s-1)} q_i^{-2-4-\dots-2(s-1)} w_s^s$$

So for all  $1 \leq r' \leq s+1$ ,  $\mu_{r'}^{(s-1)} = 0$ , and so  $P^{(s-1)} = 0$ . In particular  $P = \lambda^{(1)} + \lambda^{(2)} + \dots + \lambda^{(s-1)}$ .  $\square$

For  $1 \leq k \leq s$  consider  $P_{\pm}^{(k)}(w_1, w_2, \dots, w_s, z) \in \mathbb{C}[w_1, \dots, w_s, z]$  defined by:

$$\begin{aligned} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} \sum_{k'=1 \dots k} (z q_i^{\pm(1-s)} - w_1) (w_2 - q_i^{\pm 2} w_1) \dots (w_{k'} - q_i^{\pm 2} w_1) (w_{k'+1} q_i^{\pm 2} - w_1) \dots (w_s q_i^{\pm 2} - w_1) \\ + (-1)^{k-1} \begin{bmatrix} s \\ k-1 \end{bmatrix}_{q_i} \sum_{k'=k \dots s} (z - w_1 q_i^{\pm(1-s)}) (w_2 - q_i^{\pm 2} w_1) \dots (w_{k'} - q_i^{\pm 2} w_1) (w_{k'+1} q_i^{\pm 2} - w_1) \dots (w_s q_i^{\pm 2} - w_1) \end{aligned}$$



**Lemma 3.7.** *i) For  $2 \leq k \leq s-1$  there are polynomials  $(f_{\pm,r}^{(k)})_{r=1,\dots,s-1}$  of  $s-1$  variables, of degree at most 1 in each variable, such that  $P_{\pm}^{(k)}(w_1, \dots, w_s, z)$  is equal to :*

$$\begin{aligned} & (z - q_i^{\pm(1-s)} w_k) f_{\pm,k-1}^{(k)}(w_1, \dots, w_{k-1}, w_{k+1}, \dots, w_s, z) + (w_{k+1} - q_i^{\pm(1-s)} z) f_{\pm,s-1}^{(k)}(w_1, \dots, w_k, w_{k+2}, \dots, w_s, z) \\ & + \sum_{1 \leq r \leq s-2, r \neq k-1} (w_{r+2} - q_i^{\pm 2} w_{r+1}) f_{\pm,r}^{(k)}(w_1, \dots, w_{r-1}, w_{r+2}, \dots, w_s, z) \end{aligned}$$

*ii) There are polynomials  $(f_{\pm,r}^{(1)})_{r=1,\dots,s-1}$  of  $s-1$  variables, of degree at most 1 in each variable, such that  $P_{\pm}^{(1)}(w_1, \dots, w_s, z)$  is equal to :*

$$(w_2 - q_i^{\pm(1-s)} z) f_{\pm,s}^{(1)}(w_3, \dots, w_s, z) + \sum_{1 \leq r \leq s-2} (w_{r+2} - q_i^{\pm 2} w_{r+1}) f_{\pm,r}^{(1)}(w_1, \dots, w_{r-1}, w_{r+2}, \dots, w_s, z)$$

*iii) There are polynomials  $(f_{\pm,r}^{(s)})_{r=1,\dots,s-1}$  of  $s-1$  variables, of degree at most 1 in each variable, such that  $P_{\pm}^{(s)}(w_1, \dots, w_s, z)$  is equal to :*

$$(z - q_i^{\pm(1-s)} w_s) f_{\pm,s-1}^{(s)}(w_1, \dots, w_{s-1}, z) + \sum_{1 \leq r \leq s-2} (w_{r+2} - q_i^{\pm 2} w_{r+1}) f_{\pm,r}^{(s)}(w_1, \dots, w_{r-1}, w_{r+2}, \dots, w_s, z)$$

*Proof:* It suffices to prove it for  $P_+^{(k)}$  (because  $P_-^{(k)}$  is obtained from  $P_+^{(k)}$  by  $q \mapsto q^{-1}$ ).

For i) : we see as in lemma 3.6 that it suffices to check that  $P_+^{(k)}(w_1, \dots, w_s, z) = 0$  if  $w_3 = q_i^2 w_2, \dots, w_k = q_i^2 w_{k-1}, w_{k+2} = q_i^2 w_{k+1}, \dots, w_s = q_i^2 w_{s-1}, z = q_i^{1-s} w_k$  and  $w_{k+1} = q_i^{1-s} z$ . It means  $w_3 = q_i^2 w_2, \dots, w_k = q_i^{2(k-2)} w_2, w_{k+1} = q_i^{2k-2-2s} w_2, \dots, w_s = q_i^{-4} w_2, z = q_i^{2k-3-s} w_2$ . So if we set  $u = w_1/w_2$  we find for  $P_+^{(k)} w_2^{-s}$ :

$$\begin{aligned} & (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} \sum_{q_i^{k'=1\dots k}} q_i^{2(k'-1)} (q_i^{2k'-2-2s} - u) (q_i^{2k'-2s} - u) \dots (q_i^{2k'-6} - u) (q_i^{2k'} - u) \dots (q_i^{2k'-2} - u) (q_i^{-2} - u) \\ & + (-1)^{k-1} \begin{bmatrix} s \\ k-1 \end{bmatrix}_{q_i} \sum_{q_i^{k'=k\dots s}} q_i^{2k'-s-1} (q_i^{2k-2s-4} - u) \dots (q_i^{2k'-2s-6} - u) (q_i^{2k'-2s} - u) \dots (q_i^{2k-4} - u) (q_i^{-2} - u) \end{aligned}$$

It is a multiple of:

$$\begin{aligned} & \frac{[s-k+1]_{q_i}}{q_i^{2k-2s-4}-u} \left[ \sum_{k'=1\dots k} \frac{q_i^{2k'-1}}{(q_i^{2k'-2}-u)(q_i^{2k'-4}-u)} \right] - \frac{[k]_{q_i}}{q_i^{2k-2}-u} q_i^s \left[ \sum_{k'=k\dots s} \frac{q_i^{2k'-1-s}}{(q_i^{2k'-2s-2}-u)(q_i^{2k'-2s-4}-u)} \right] \\ & = \frac{q_i^2 [s-k+1]_{q_i}}{(1-q_i^2)(q_i^{2k-2s-4}-u)} \left[ \sum_{k'=1\dots k} \frac{1}{q_i^{2k'-2}-u} - \frac{1}{q_i^{2k'-4}-u} \right] - \frac{q_i^2 [k]_{q_i}}{(1-q_i^2)q_i^{2k-2}-u} q_i^s \left[ \sum_{k'=k\dots s} \frac{1}{q_i^{2k'-2s-2}-u} - \frac{1}{q_i^{2k'-2s-4}-u} \right] \\ & = \frac{q_i^2 [s-k+1]_{q_i}}{(1-q_i^2)(q_i^{2k-2s-4}-u)} \left[ \frac{1}{q_i^{2k-2}-u} - \frac{1}{q_i^{-2}-u} \right] - \frac{q_i^2 [k]_{q_i}}{(1-q_i^2)(q_i^{2k-2}-u)} q_i^s \left[ \frac{1}{q_i^{-2}-u} - \frac{1}{q_i^{2k-2s-4}-u} \right] = 0 \end{aligned}$$

For ii) : as for i) we check that  $P_+^{(1)}(w_1, \dots, w_s, z) = 0$  if  $w_3 = q_i^2 w_2, \dots, w_s = q_i^2 w_{s-1}, z = q_i^{s-1} w_2$ . It means  $w_{k'} = q_i^{2(k'-2)} w_2$  for  $2 \leq k' \leq s$ . So if we set  $u = w_1/w_2$  we find for  $P_+^{(1)} w_2^{-s}$ :

$$- [s]_{q_i} (1-u) (q_i^2 - u) \dots (q_i^{2s-2} - u) + q_i^{1-s} \sum_{k'=1\dots s} q_i^{2k'-2} (q_i^{2s-2} - u) (q_i^{-2} - u) \dots (q_i^{2k'-6} - u) (q_i^{2k'} - u) \dots (q_i^{2s-2} - u)$$

It is a multiple of:  $-\frac{q_i^2 [s]_{q_i}}{q_i^{2s}-1} \left( \frac{1}{q_i^{-2}-u} - \frac{1}{q_i^{2s-2}-u} \right) + \frac{q_i^{1-s}}{1-q_i^{-2}} \left( \frac{1}{q_i^{-2}-u} - \frac{1}{q_i^{2s-2}-u} \right) = 0$ .

For iii) : as for i) we check that  $P_+^{(s)}(w_1, \dots, w_s, z) = 0$  if  $w_3 = q_i^2 w_2, \dots, w_s = q_i^2 w_{s-1}, z = q_i^{1-s} w_s$ . It means  $w_{k'} = q_i^{2(k'-2)} w_2$  for  $2 \leq k' \leq s$  and  $z = q_i^{s-3} w_2$ . The computation is analogous to i).  $\square$

**Lemma 3.8.** *For all choices of polynomials  $(f_{\pm,r}^{(k')})_{1 \leq k' \leq s, 1 \leq r \leq s-1}$  in lemma 3.7 and each  $2 \leq k \leq s$  there are polynomials  $(g_{\pm,r}^{(k)})_{r=1,\dots,s-2}$  of  $s-1$  variables such that:*

$$f_{\pm,k-1}^{(k)} - f_{\pm,s-1}^{(k-1)} = \sum_{1 \leq r \leq s-2} (w_{r+2} - q_i^{\pm 2} w_{r+1}) g_{\pm,r}^{(k)}(w_1, \dots, w_{r-1}, w_{r+2}, \dots, w_s, z)$$

*Proof:* We see as in lemma 3.6 that it suffices to check that  $f_{+,k-1}^{(k)} + f_{+,s-1}^{(k-1)} = 0$  if  $w_3 = q_i^2 w_2, \dots$ ,  $w_s = q_i^2 w_{s-1}$ . So we suppose that  $w_{k'} = q_i^{2(k'-2)}$  for all  $2 \leq k' \leq s$ . Let  $Q = w_1^{s-1}(w_2 q_i^{-2} - 1)(w_2 - 1) \dots (w_2 q_i^{2s-2}) / (q_i^2 - 1)$ . It suffices to prove that for  $2 \leq k \leq s$ , we have:

$$(18) \quad (q_i^2 - q_i^{2-2s}) f_{+,k-1}^{(k)} (Q(-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} [k]_{q_i} (q_i - q_i^{-1}))^{-1} = \frac{q_i^{k+1+s} + q_i^{-s-k+3} - q_i^{-s+k+1} - q_i^{3-k+s}}{(v q_i^{-2} - 1)(v q_i^{2k-4} - 1)(v q_i^{2s-2} - 1)}$$

$$(19) \quad (q_i^2 - q_i^{2-2s}) f_{+,s-1}^{(k-1)} (Q(-1)^{k-1} \begin{bmatrix} s \\ k-1 \end{bmatrix}_{q_i} [k-1]_{q_i} (q_i - q_i^{-1}))^{-1} = \frac{q_i^{k+1} + q_i^{-k+3} - q_i^{-2s+k+1} - q_i^{3-k+2s}}{(v q_i^{-2} - 1)(v q_i^{2k-4} - 1)(v q_i^{2s-2} - 1)}$$

because we have the relation :

$$\begin{bmatrix} s \\ k \end{bmatrix}_{q_i} [k]_{q_i} (q_i^{k+1+s} + q_i^{-s-k+3} - q_i^{-s+k+1} - q_i^{3-k+s}) = - \begin{bmatrix} s \\ k-1 \end{bmatrix}_{q_i} [k-1]_{q_i} (q_i^{k+1} + q_i^{-k+3} - q_i^{-2s+k+1} - q_i^{3-k+2s})$$

First suppose that  $3 \leq k \leq s-1$ . We have  $P_+^{(k)} = (z - q_i^{1-s} w_k) f_{+,k-1}^{(k)} + (q_i^2 w_k - q_i^{1-s} z) f_{+,s-1}^{(k)}$ . So for  $\alpha_k, \beta_k$  such that  $P_+^{(k)} = z \alpha_k + w_k \beta_k$ , we have  $f_{+,k-1}^{(k)} = \frac{q_i^2 \alpha_k + q_i^{1-s} \beta_k}{q_i^2 - q_i^{2-2s}}$  and  $f_{+,s-1}^{(k)} = \frac{q_i^{1-s} \alpha_k + \beta_k}{q_i^2 - q_i^{2-2s}}$ . But we have  $P_+^{(k)} = z(q_i^{1-s} \lambda_k + \mu_k) - w_1(\lambda_k + q_i^{1-s} \mu_k)$  where (we put  $v = w_2/w_1$ ):

$$\begin{aligned} \lambda_k &= (-1)^k w_1^{s-1} \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} \sum_{k'=1 \dots k} (v - q_i^2)(v q_i^2 - q_i^2) \dots (v q_i^{2(k'-2)} - q_i^2)(v q_i^{2k'} - 1) \dots (v q_i^{2(s-2)+2} - 1) \\ &= Q(-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} \left[ \frac{1}{v q_i^{-2} - 1} - \frac{q_i^{2k}}{v q_i^{2k-2} - 1} \right] \end{aligned}$$

$$\begin{aligned} \mu_k &= (-1)^{k-1} w_1^{s-1} \begin{bmatrix} s \\ k-1 \end{bmatrix}_{q_i} \sum_{k'=k \dots s} (v - q_i^2)(v q_i^2 - q_i^2) \dots (v q_i^{2(k'-2)} - q_i^2)(v q_i^{2k'} - 1) \dots (v q_i^{2(s-2)+2} - 1) \\ &= Q(-1)^{k-1} \begin{bmatrix} s \\ k-1 \end{bmatrix}_{q_i} \left[ \frac{q_i^{2k-2}}{v q_i^{2k-4} - 1} - \frac{q_i^{2s}}{v q_i^{2s-2} - 1} \right] \end{aligned}$$

As  $\alpha_k = q_i^{1-s} \lambda_k + \mu_k$  and  $\beta_k = -(\lambda_k + q_i^{1-s}) / (q_i^{k-2} w_2)$ , we have:

$$\begin{aligned} \alpha_k &= Q \frac{(-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} [k]_{q_i} (q_i - q_i^{-1}) ((q_i^{k+1-s} - q_i^{s+k-1}) + v(q_i^{s+k-3} + q_i^{s+3k-3} - q_i^{3k-3-s} - q_i^{s+k-1}))}{(v q_i^{-2} - 1)(v q_i^{2k-2} - 1)(v q_i^{2k-4} - 1)(v q_i^{2s-2} - 1)} \\ \beta_k &= Q \frac{(-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} [k]_{q_i} (q_i - q_i^{-1}) ((q_i^k + q_i^{2s-k+2} - q_i^{-k+2} - q_i^{k+2}) + v(-q_i^{k+2s-2} + q_i^k))}{(v q_i^{-2} - 1)(v q_i^{2k-2} - 1)(v q_i^{2k-4} - 1)(v q_i^{2s-2} - 1)} \end{aligned}$$

In particular  $(q_i^2 - q_i^{2-2s}) f_{+,k-1}^{(k)} (Q(-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} [k]_{q_i} (q_i - q_i^{-1}))^{-1}$  is:

$$\frac{(q_i^{k+1-s} + q_i^{s-k+3} - q_i^{s+k+1} - q_i^{3-k-s}) + v(q_i^{s+3k-1} + q_i^{k+1-s} - q_i^{3k-1-s} - q_i^{s+k+1})}{(v q_i^{-2} - 1)(v q_i^{2k-2} - 1)(v q_i^{2k-4} - 1)(v q_i^{2s-2} - 1)}$$

and we get formula 18 for  $k$ . Moreover  $(q_i^2 - q_i^{2-2s}) f_{+,s-1}^{(k-1)} (Q(-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} [k]_{q_i} (q_i - q_i^{-1}))^{-1}$  is:

$$\frac{(q_i^{k+2-2s} + q_i^{2s-k+2} - q_i^{-k+2} - q_i^{k+2}) + v(q_i^{k-2} + q_i^{3k-2} - q_i^{3k-2s-2} - q_i^{k+2s-2})}{(v q_i^{-2} - 1)(v q_i^{2k-2} - 1)(v q_i^{2k-4} - 1)(v q_i^{2s-2} - 1)}$$

and we get formula 19 for  $k+1$ .

So it remains to prove formula 19 with  $k=2$  and formula 18 with  $k=s$ .

$$\begin{aligned} P_+^{(1)} &= (w_2 - q_i^{(1-s)} z) f_{+,s-1}^{(1)} = -[s]_{q_i} (z q_i^{1-s} - w_1) (q_i^2 w_2 - w_1) \dots (q_i^{2s-2} w_2 - w_1) \\ &+ (z - w_1 q_i^{1-s}) \sum_{k'=1 \dots s} q_i^{2k'-2} (q_i^{-2} w_2 - w_1) \dots (q_i^{2k'-6} w_2 - w_1) (q_i^{2k'} w_2 - w_1) \dots (q_i^{2s-2} w_2 - w_1) \end{aligned}$$

$$\Rightarrow f_{+,s-1}^{(1)} = -q_i^{1-s} Q \left[ \frac{-[s]_{q_i} q_i^{1-s} (q_i^2 - 1)}{(vq_i^{-2} - 1)(v - 1)} + \sum_{k'=1 \dots s} \frac{q_i^{2k'-2}}{(q_i^{2k'-4} v - 1)(q_i^{2k'-2} v - 1)} \right]$$

And so we have for  $f_{+,s-1}^{(1)}(q_i^2 - q_i^{2-2s})(-Q[s]_{q_i}(q_i - q_i^{-1}))^{-1}$ :

$$\frac{q_i + q_i^3 - q_i^{2s+1} - q_i - 2s + 3}{(vq_i^{-2} - 1)(v - 1)(vq_i^{2s-2} v - 1)}$$

that it to say the formula 19 with  $k = 2$ .

$$\begin{aligned} P_+^{(s)} &= (z - q_i^{(1-s)} q_i^{2(s-2)} w_2) f_{+,s-1}^{(s)} = (-1)^{s-1} [s]_{q_i} (z - w_1 q_i^{1-s}) q_i^{2(s-1)} (q_i^{-2} w_2 - w_1) \dots (q_i^{2s-6} w_2 - w_1) \\ &+ (-1)^s (z q_i^{1-s} - w_1) \sum_{k'=1 \dots s} q_i^{2k'-2} (q_i^{-2} w_2 - w_1) \dots (q_i^{2k'-6} w_2 - w_1) (q_i^{2k'} w_2 - w_1) \dots (q_i^{2s-2} w_2 - w_1) \\ \Rightarrow f_{+,s-1}^{(s)} &= Q \left[ \frac{(-1)^{s-1} [s]_{q_i} q_i^{2(s-1)} (q_i^2 - 1)}{(vq_i^{2s-4} - 1)(vq_i^{2s-2} - 1)} + (-1)^s q_i^{1-s} \sum_{k'=1 \dots s} \frac{q_i^{2k'-2}}{(q_i^{2k'-4} v - 1)(q_i^{2k'-2} v - 1)} \right] \end{aligned}$$

And so we have for  $f_{+,s-1}^{(s)}(q_i^2 - q_i^{2-2s})((-1)^s Q[s]_{q_i}(q_i - q_i^{-1}))^{-1}$ :

$$\frac{q_i^{2s+1} + q_i^{3-2s} - q_i - q_i^3}{(vq_i^{-2} - 1)(vq_i^{2s-4} - 1)(vq_i^{2s-2} v - 1)}$$

that it to say the formula 18 with  $k = s$ . □

**3.3.4. Proof of theorem 3.2.** The algebras  $\mathcal{U}_q^l(\hat{\mathfrak{g}}), \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}), \mathcal{U}_q^{l,\pm}(\hat{\mathfrak{g}})$  are defined in section 3.3. Let  $\tilde{\mathcal{U}}_q^{\pm}(\hat{\mathfrak{g}}) \subset \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  be the subalgebra generated by the  $x_{i,r}^{\pm}$ . Let  $\tau_{\pm}$  be the two-sided ideal of  $\mathcal{U}_q^{l,\pm}(\hat{\mathfrak{g}})$  generated by the left terms of relations (9) (with the  $x_{i,r}^{\pm}$ ).

**Lemma 3.9.** *We have  $\tau_+ \mathcal{U}_q^l(\hat{\mathfrak{g}}) \subset \mathcal{U}_q^l(\hat{\mathfrak{g}}) \tau_+$  and  $\mathcal{U}_q^l(\hat{\mathfrak{g}}) \tau_- \subset \tau_- \mathcal{U}_q^l(\hat{\mathfrak{g}})$ . In particular  $(\tilde{\mathcal{U}}_q^-(\hat{\mathfrak{g}}), \mathcal{U}_q(\hat{\mathfrak{h}}), \tilde{\mathcal{U}}_q^+(\hat{\mathfrak{g}}))$  is a triangular decomposition of  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ .*

*Proof:* First  $\tau_+ \mathcal{U}_q(\hat{\mathfrak{h}}) \subset \mathcal{U}_q(\hat{\mathfrak{h}}) \tau_+, \mathcal{U}_q(\hat{\mathfrak{h}}) \tau_- \subset \tau_- \mathcal{U}_q(\hat{\mathfrak{h}})$  are direct consequences of relations (12), (13), (14). We have also (we use relations (15) and (13), (14)):

$$\begin{aligned} &[(w - q^{\pm B_{i,j}} z) x_i^{\pm}(z) x_j^{\pm}(w) - (q^{\pm B_{i,j}} w - z) x_j^{\pm}(w) x_i^{\pm}(z), x_k^{\mp}(u)] \\ &= (w - q^{\pm B_{i,j}} z) x_i^{\pm}(z) [x_j^{\pm}(w), x_k^{\mp}(u)] - (q^{\pm B_{i,j}} w - z) [x_j^{\pm}(w), x_k^{\mp}(u)] x_i^{\pm}(z) \\ &- (q^{\pm B_{i,j}} w - z) x_j^{\pm}(w) [x_i^{\pm}(z), x_k^{\mp}(u)] + (w - q^{\pm B_{i,j}} z) [x_i^{\pm}(z), x_k^{\mp}(u)] x_j^{\pm}(w) = 0 \end{aligned}$$

and so  $\tau_+ \mathcal{U}_q^{l,-}(\hat{\mathfrak{g}}) \subset \mathcal{U}_q^l(\hat{\mathfrak{g}}) \tau_+, \mathcal{U}_q^{l,+}(\hat{\mathfrak{g}}) \tau_- \subset \tau_- \mathcal{U}_q^l(\hat{\mathfrak{g}})$ .

The last point follows from  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}) = \mathcal{U}_q^l(\hat{\mathfrak{g}}) / (\mathcal{U}_q^l(\hat{\mathfrak{g}}) \cdot (\tau_+ + \tau_-) \cdot \mathcal{U}_q^l(\hat{\mathfrak{g}}))$ , the triangular decomposition of  $\mathcal{U}_q^l(\hat{\mathfrak{g}})$  and lemma 3.5. □

**Lemma 3.10.** *Let  $i \neq j, s = 1 - C_{i,j}$   $\mu = 1$  or  $\mu = -1$ . We have in  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  :*

$$(20) \quad \sum_{\pi \in \Sigma_s} \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} x_i^{\pm}(w_{\pi(1)}) \dots x_i^{\pm}(w_{\pi(k)}) \phi_j^{\mu}(z) x_i^{\pm}(w_{\pi(k+1)}) \dots x_i^{\pm}(w_{\pi(s)}) = 0$$

$$(21) \quad \sum_{\pi \in \Sigma_s} \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} \xi_i(w_{\pi(1)}) \dots \xi_i(w_{\pi(k)}) x_j^{\pm}(z) \xi_i(w_{\pi(k+1)}) \dots \xi_i(w_{\pi(s)}) = 0$$

where  $\xi_i(w_p) = x_i^{\pm}(w_p)$  if  $p \neq 1$  and  $\xi_i(w_1) = \phi_i^{\mu}(w_1)$ .

Remark : in particular if we multiply the equation (20) by  $(\prod_{r=1\dots s} (w_r - q_i^{s-1}z))(\prod_{1\leq r'<r\leq s} (w_r - q_i^2 w_{r'}))$  and we project it on  $x_i^\pm(w_1)\dots x_i^\pm(w_s)\phi_j^\pm(z)$  (we can use the relations (16) thanks to the multiplied polynomial), we get the combinatorial identity discovered by Jing in [Jin], which was also proved in a combinatorial way in [DJ] : for  $\pi \in \Sigma_s$  denote by  $\epsilon(\pi) \in \{1, -1\}$  the signature of  $\pi$  (we have replaced  $z \mapsto z^{-1}$ ,  $w_{k'} \mapsto w_{k'}^{-1}$  to get the formula in the same form as in [Jin]):

$$0 = \sum_{\pi \in \Sigma_s} \epsilon(\pi) \sum_{k=0\dots s} \begin{bmatrix} s \\ k \end{bmatrix}_q (z - q^{s-1}w_{\pi(1)})\dots(z - q^{s-1}w_{\pi(k)}) \\ (w_{\pi(k+1)} - q^{s-1}z)\dots(w_{\pi(s)} - q^{s-1}z) \prod_{1\leq r<r'\leq s} (w_{\pi(r)} - q^2 w_{\pi(r')})$$

*Proof:* First we prove the equation (20) with  $\mu = 1$  ( $\mu = -1$  is analog). The left term is (relations (13)):

$$\frac{\phi_j^\pm(z)}{(w_1 q^{\pm B_{i,j}} - z)\dots(w_s q^{\pm B_{i,j}} - z)} \sum_{\pi \in \Sigma_s} P_\pm(w_{\pi(1)}, \dots, w_{\pi(s)}, z) x_i^\pm(w_{\pi(1)})\dots x_i^\pm(w_{\pi(s)})$$

that is to say (see lemma 3.6):

$$\sum_{\pi \in \Sigma_s} \sum_{1\leq r\leq s-1} (w_{\pi(r+1)} - q_i^{\pm 2} w_{\pi(r)}) f_{r,\pm}(w_{\pi(1)}, \dots, w_{\pi(r-1)}, w_{\pi(r+2)}, \dots, w_{\pi(s)}, z) x_i^\pm(w_{\pi(1)})\dots x_i^\pm(w_{\pi(s)})$$

For each  $r$ , we put together the  $\pi, \pi' \in \Sigma_s$  such that  $\pi(r) = \pi'(r+1)$ ,  $\pi(r+1) = \pi'(r)$ , and  $\pi(r'') = \pi'(r'')$  for all  $r'' \neq r, r+1$ . So we get a sum of terms:

$$f_{r,\pm}(w_{\pi(1)}, \dots, w_{\pi(r-1)}, w_{\pi(r+2)}, \dots, w_{\pi(s)}, z) x_i^\pm(w_{\pi(1)})\dots x_i^\pm(w_{\pi(r-1)}) A_{\{\pi(r), \pi(r+1)\}}^\pm x_i^\pm(w_{\pi(r+2)})\dots x_i^\pm(w_{\pi(s)})$$

$$\text{where } A_{\{k,k'\}}^\pm = (w_k - q_i^{\pm 2} w_{k'}) x_i^\pm(w_{k'}) x_i^\pm(w_k) + (w_{k'} - q_i^{\pm 2} w_k) x_i^\pm(w_k) x_i^\pm(w_{k'})$$

But  $A_{\{k,k'\}}^\pm = 0$  in  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ .

Let us prove the equation (21) with  $\mu = 1$  ( $\mu = -1$  is analog). The left term is :

$$\frac{\phi_i^+(w_1)}{(w_2 q_i^{\pm 2} - w_1)\dots(w_s q_i^{\pm 2} - w_1)(z q_i^{\pm(1-s)} - w_1)} \\ \sum_{\pi \in \Sigma_{s-1}, k=1\dots s} P_\pm^{(k)}(w_1, w_{\pi(2)}, \dots, w_{\pi(s)}, z) x_i^\pm(w_{\pi(2)})\dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)})\dots x_i^\pm(w_{\pi(s)})$$

where  $\Sigma_{s-1}$  acts on  $\{2, \dots, s\}$ . With the help of lemma 3.7 and in analogy to the previous case, for each  $1 \leq k \leq s$  each  $r \neq k$ , we put together the  $\pi, \pi' \in \Sigma_s$  such that  $\pi(r) = \pi'(r+1)$ ,  $\pi(r+1) = \pi'(r)$ , and  $\pi(r'') = \pi'(r'')$  for all  $r'' \neq r, r+1$ . So the terms with polynomials  $f_{\pm, k'}^{(k)}$  with  $k' \neq s, k-1$  are erased.

$$\text{We get : } \frac{\phi_i^+(w_1)}{(w_2 q_i^{\pm 2} - w_1)\dots(w_s q_i^{\pm 2} - w_1)(z q_i^{\pm(1-s)} - w_1)} \\ \sum_{\pi \in \Sigma_{s-1}, k=1\dots s} ((z - q_i^{\pm(1-s)} w_{\pi(k)}) f_{\pm, k-1}^{(k)} + (w_{\pi(k+1)} - q_i^{\pm(1-s)} z) f_{\pm, s-1}^{(k)}) \\ x_i^\pm(w_{\pi(2)})\dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)})\dots x_i^\pm(w_{\pi(s)})$$

But this last sum is equal to :

$$\sum_{\pi \in \Sigma_{s-1}, k=2\dots s} (z - q_i^{\pm(1-s)} w_{\pi(k)}) (f_{\pm, k-1}^{(k)} - f_{\pm, s-1}^{(k-1)}) x_i^\pm(w_{\pi(2)})\dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)})\dots x_i^\pm(w_{\pi(s)})$$

where we can replace  $(z - q_i^{\pm(1-s)} w_{\pi(k)}) x_i^\pm(w_{\pi(k)}) x_j^\pm(z)$  by  $(-w_{\pi(k)} + q_i^{\pm(1-s)} z) x_j^\pm(z) x_i^\pm(w_{\pi(k)})$  (relations (16) in  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ ). As in the previous cases it follows from lemma 3.8 that this term is equal to 0.  $\square$

Let  $\tilde{\tau}_\pm$  be the two-sided ideal of  $\tilde{\mathcal{U}}_q^\pm(\hat{\mathfrak{g}})$  generated by the left terms of relations (10) with the  $x_{i,r}^\pm$ .

**Lemma 3.11.** *We have  $\tilde{\tau}_+ \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}) \subset \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}) \tilde{\tau}_+$  and  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}) \tilde{\tau}_- \subset \tilde{\tau}_- \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ .*

In particular as  $\mathcal{U}_q(\hat{\mathfrak{g}}) = \tilde{\mathcal{U}}_q^l(\hat{\mathfrak{g}})/(\tilde{\mathcal{U}}_q^l(\hat{\mathfrak{g}}).(\tilde{\tau}_+ + \tilde{\tau}_-).\tilde{\mathcal{U}}_q^l(\hat{\mathfrak{g}}))$  the result of theorem 3.2 follows from lemma 3.5 and the triangular decomposition of  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  proved in lemma 3.9.

*Proof:* First  $\tilde{\tau}_+\mathcal{U}_q(\hat{\mathfrak{h}}) \subset \mathcal{U}_q(\hat{\mathfrak{h}})\tau_+$ ,  $\mathcal{U}_q(\hat{\mathfrak{h}})\tilde{\tau}_- \subset \tilde{\tau}_-\mathcal{U}_q(\hat{\mathfrak{h}})$  are direct consequences of relations (12), (13), (14). Let us show that :

$$(22) \quad \left[ \sum_{\pi \in \Sigma_s} \sum_{k=0..s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} x_i^\pm(w_{\pi(1)}) \dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)}) \dots x_i^\pm(w_{\pi(s)}), x_l^\mp(u) \right] = 0$$

where  $i, j, l \in I$ ,  $i \neq j$ . If  $l \neq j$  and  $l \neq i$  the equation (22) follows from relations (15)). If  $l = j$ , the equation (22) follows from the identity (20) of lemma 3.10 because the left term is :

$$\sum_{\pi \in \Sigma_s} \sum_{k=0..s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} x_i^\pm(w_{\pi(1)}) \dots x_i^\pm(w_{\pi(k)}) (\delta(\frac{z}{u}) \phi_j^\pm(z) - \delta(\frac{z}{u}) \phi_j^\mp(z)) x_i^\pm(w_{\pi(k+1)}) \dots x_i^\pm(w_{\pi(s)})$$

If  $l = i$ , the equation (22) follows from the identity (21) of lemma 3.10 because the left term is :

$$\begin{aligned} & \sum_{\pi \in \Sigma_s} \sum_{k=0..s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} \left( \sum_{k'=1..k} x_i^\pm(w_{\pi(1)}) \dots x_i^\pm(w_{\pi(k'-1)}) \delta\left(\frac{w_{k'}}{u}\right) (\phi_i^\pm(w_{k'}) - \phi_i^\mp(w_{k'})) \right. \\ & \quad \left. x_i^\pm(w_{\pi(k'+1)}) \dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)}) \dots x_i^\pm(w_{\pi(s)}) \right) \\ & + \sum_{k'=k+1..s} x_i^\pm(w_{\pi(1)}) \dots x_i^\pm(w_{\pi(k)}) x_j^\pm(z) x_i^\pm(w_{\pi(k+1)}) \dots x_i^\pm(w_{\pi(k'-1)}) \\ & \quad \delta\left(\frac{w_{k'}}{u}\right) (\phi_i^\pm(w_{k'}) - \phi_i^\mp(w_{k'})) x_i^\pm(w_{\pi(k'+1)}) \dots x_i^\pm(w_{\pi(s)}) \end{aligned}$$

So we have proved the equation (22) and in particular  $\tilde{\tau}_+\tilde{\mathcal{U}}_q^-(\hat{\mathfrak{g}}) \subset \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})\tilde{\tau}_+$ ,  $\tilde{\mathcal{U}}_q^+(\hat{\mathfrak{g}})\tilde{\tau}_- \subset \tilde{\tau}_-\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ .  $\square$

#### 4. INTEGRABLE REPRESENTATIONS AND CATEGORY $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$

In this section we study highest weight representations of  $\mathcal{U}_q(\hat{\mathfrak{g}})$ . In particular theorem 4.9 is a generalization of a result of Chari-Pressley about integrable representations.

**4.1. Reminder: integrable representations of quantum Kac-Moody algebras.** In this section we review some known properties of integrable representations of  $\mathcal{U}_q(\mathfrak{g})$ .

For  $V$  a  $\mathcal{U}_q(\mathfrak{h})$ -module and  $\omega \in \mathfrak{h}^*$  we denote by  $V_\omega$  the weight space of weight  $\omega$ :

$$V_\omega = \{v \in V / \forall h \in \mathfrak{h}, k_h.v = q^{\omega(h)}v\}$$

In particular for  $v \in V_\omega$  we have  $k_i.v = q_i^{\omega(\alpha_i^\vee)}v$  and for  $i \in I$  we have  $x_i^\pm.V_\omega \subset V_{\omega \pm \alpha_i}$ .

We say that  $V$  is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable if  $V = \bigoplus_{\omega \in \mathfrak{h}^*} V_\omega$  (in particular  $V$  is of type 1).

**Definition 4.1.** A  $\mathcal{U}_q(\mathfrak{g})$ -module  $V$  is said to be integrable if  $V$  is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable,  $\forall \omega \in \mathfrak{h}^*$ ,  $V_\omega$  is finite dimensional, and for  $\mu \in \mathfrak{h}^*$ ,  $i \in I$  there is  $R \geq 0$  such that  $r \geq R \Rightarrow V_{\mu \pm r\alpha_i} = \{0\}$ .

In particular for all  $v \in V$  there is  $m_v \geq 0$  such that for all  $i \in I$ ,  $m \geq m_v$ ,  $(x_i^+)^m.v = (x_i^-)^m.v = 0$ , and  $U_i.v$  is finite dimensional.

**Definition 4.2.** A  $\mathcal{U}_q(\mathfrak{g})$ -module  $V$  is said to be of highest weight  $\omega \in \mathfrak{h}^*$  if there is  $v \in V_\omega$  such that  $V$  is generated by  $v$  and  $\forall i \in I$ ,  $x_i^+.v = 0$ .

In particular  $V = \mathcal{U}_q(\mathfrak{g})^-.v$  (theorem 2.3),  $V$  is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable, and  $V = \bigoplus_{\lambda \leq \omega} V_\lambda$ . We have (see [L]):

**Theorem 4.3.** For any  $\omega \in \mathfrak{h}^*$  there is a unique up to isomorphism simple highest weight module  $L(\omega)$  of highest weight  $\omega$ . The highest weight module  $L(\omega)$  is integrable if and only  $\omega \in P^+$ .

**4.2. Integrable representations of quantum affinizations.** In this section we generalize results of Chari-Pressley [CP3, CP4] to all quantum affinizations.

4.2.1. *l-highest weight modules.* We give the following notion of  $l$ -weight :

**Definition 4.4.** A couple  $(\lambda, \Psi)$  such that  $\lambda \in \mathfrak{h}^*$ ,  $\Psi = (\Psi_{i,\pm m}^\pm)_{i \in I, m \geq 0}$ ,  $\Psi_{i,\pm m}^\pm \in \mathbb{C}$ ,  $\Psi_{i,0}^\pm = q_i^{\pm \lambda(\alpha_i^\vee)}$  is called a  $l$ -weight.

The condition  $\Psi_{i,0}^\pm = q_i^{\pm \lambda(\alpha_i^\vee)}$  is a compatibility condition which comes from  $\phi_{i,0}^\pm = k_i^\pm$ .

We denote by  $P_l$  the set of  $l$ -weights. Note that in the finite case  $\lambda$  is uniquely determined by  $\Psi$  because  $\lambda = \sum_{i \in I} \lambda(\alpha_i^\vee) \Lambda_i$ . Analogs of those  $l$ -weights were also used in [M1] for toroidal  $\hat{sl}_n$ -cases.

**Definition 4.5.** A  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -module  $V$  is said to be of  $l$ -highest weight  $(\lambda, \Psi) \in P_l$  if there is  $v \in V$  such that  $(i \in I, r \in \mathbb{Z}, m \geq 0, h \in \mathfrak{h})$ :

$$x_{i,r}^+ \cdot v = 0, \quad V = \mathcal{U}_q(\hat{\mathfrak{g}}) \cdot v, \quad \phi_{i,\pm m}^\pm \cdot v = \Psi_{i,\pm m}^\pm v, \quad kh \cdot v = q^{\lambda(h)} \cdot v$$

In particular  $\mathcal{U}_q(\hat{\mathfrak{g}})^- \cdot v = V$  (theorem 3.2),  $V$  is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable and  $V = \bigoplus_{\lambda \leq \omega} V_\lambda$ . Note that the  $l$ -weight  $(\lambda, \Psi) \in P_l$  is uniquely determined by  $V$ . It is called the  $l$ -highest weight of  $V$ .

The notion of  $l$ -highest weight is different from the notion of highest weight for quantum affine algebras. The term ‘‘pseudo highest weight’’ is also used in the literature.

Example : for any  $(\lambda, \Psi) \in P_l$ , define the Verma module  $M(\lambda, \Psi)$  as the quotient of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  by the left ideal generated by the  $x_{i,r}^+$  ( $i \in I, r \in \mathbb{Z}$ ),  $kh - q^{\lambda(h)}$  ( $h \in \mathfrak{h}$ ),  $\phi_{i,\pm m}^\pm - \Psi_{i,\pm m}^\pm$  ( $i \in I, m \geq 0$ ). It follows from theorem 3.2 that  $M(\lambda, \Psi)$  is a free  $\mathcal{U}_q^-(\hat{\mathfrak{g}})$ -module of rank 1. In particular it is non trivial and it is a  $l$ -highest weight module of highest weight  $(\lambda, \Psi)$ . Moreover it has a unique proper submodule (mimic the classical argument in [Ka]), and :

**Proposition 4.6.** For any  $(\lambda, \Psi) \in P_l$  there is a unique up to isomorphism simple  $l$ -highest weight module  $L(\lambda, \Psi)$  of  $l$ -highest weight  $(\lambda, \Psi)$ .

4.2.2. *Integrable  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -modules.*

**Definition 4.7.** A  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -module  $V$  is said to be integrable if  $V$  is integrable as a  $\mathcal{U}_q(\mathfrak{g})$ -module.

Note that in the case of a quantum affine algebra, the two notions of integrability do not coincide. Throughout the paper only the notion of integrability of definition 4.7 is used.

For  $i \in I, r \in \mathbb{Z}$  and  $\omega \in \mathfrak{h}^*$  we have  $x_{i,r}^\pm \cdot V_\omega \subset V_{\omega \pm \alpha_i}$ . So if  $V$  is integrable, for all  $v \in V$ ,  $\hat{U}_i \cdot v$  is finite dimensional and there is  $m_0 \geq 1$  such that for all  $i \in I, r \in \mathbb{Z}, m \geq m_0 \Rightarrow (x_{i,r}^+)^m \cdot v = (x_{i,r}^-)^m \cdot v = 0$ .

**Definition 4.8.** The set  $P_l^+$  of dominant  $l$ -weights is the set of  $(\lambda, \Psi) \in P_l$  such that there exist (Drinfel'd)-polynomials  $P_i(z) \in \mathbb{C}[z]$  ( $i \in I$ ) of constant term 1 such that in  $\mathbb{C}[[z]]$  (resp. in  $\mathbb{C}[[z^{-1}]]$ ):

$$\sum_{m \geq 0} \Psi_{i,\pm m}^\pm z^{\pm m} = q_i^{\deg(P_i)} \frac{P_i(zq_i^{-1})}{P_i(zq_i)}$$

In particular for all  $i \in I$ ,  $\lambda(\alpha_i^\vee) = \deg(P_i) \geq 0$  and so  $\lambda \in P^+$  is a dominant weight.

**Theorem 4.9.** For  $(\lambda, \Psi) \in P_l$ ,  $L(\lambda, \Psi)$  is integrable if and only  $(\lambda, \Psi) \in P_l^+$ .

If  $\mathfrak{g}$  is finite (case of a quantum affine algebra) it is a result of Chari-Pressley in [CP3] (if part) and in [CP4] (only if part). Moreover in this case the integrable  $L(\lambda, \Psi)$  are finite dimensional. If  $\mathfrak{g}$  is symmetric the result is geometrically proved by Nakajima in [N1]. If  $C$  is of type  $A_n^{(1)}$  (toroidal  $\hat{sl}_n$ -case) the result is algebraically proved by Miki in [M1].

For the general case we propose a proof similar to the proof given by Chari-Pressley in the finite case. For  $\lambda \in \mathfrak{h}^*$  denote  $D(\lambda) = \{\omega \in \mathfrak{h}^*/\omega \leq \lambda\}$ .

*Proof:* The proof uses the result for  $\mathcal{U}_q(\hat{\mathfrak{sl}}_2)$  which is proved in [CP1, CP3].

First suppose that  $L = L(\lambda, \Psi)$  is integrable and for  $i \in I$  let  $L_i$  be the  $\hat{U}_i$ -submodule of  $L$  generated by the highest weight vector  $v$ . It is a  $l$ -highest weight  $\mathcal{U}_{q_i}(\hat{\mathfrak{sl}}_2)$ -module of highest weight  $(\lambda(\alpha_i^\vee), \Psi_i^\pm)$ . As  $L$  is integrable,  $L_i$  is finite dimensional. So the result for  $\mathcal{U}_{q_i}(\hat{\mathfrak{sl}}_2)$  gives  $P_i(z) \in \mathbb{C}[z]$  such that:

$$\sum_{m \geq 0} \Psi_{i, \pm m}^\pm z^{\pm m} = q_i^{\deg(P_i)} \frac{P_i(zq_i^{-1})}{P_i(zq_i)}, \quad \lambda(\alpha_i^\vee) = \deg(P_i) \geq 0$$

Now we prove that  $L = L(\lambda, \Psi) = \mathcal{U}_q(\hat{\mathfrak{g}}).v$  is integrable where  $(\lambda, \Psi) \in P_l^+$ . It suffices to prove that:

(1) For all  $\mu \leq \lambda$ , if  $L_\mu \neq \{0\}$  then there exists  $M > 0$  such that  $m > M \Rightarrow L_{\mu - m\alpha_i} = L_{\mu + m\alpha_i} = 0$  for all  $i \in I$ .

(2) For all  $\mu \leq \lambda$ ,  $\dim(L_\mu) < \infty$ .

The proof goes roughly as in section 5 of [CP3], with the following modifications :

For (1) : the existence of  $M$  for  $L_{\mu + m\alpha_i} = 0$  is clear because the weights of  $L$  are in  $D(\lambda)$ . Put  $r^\vee = \max\{-C_{i,j}/i \neq j\}$ . In particular if  $C$  is finite, we have  $r^\vee \leq 3$ . First we prove that for  $m > 0$ , the space  $L_{\mu - m\alpha_i}$  is spanned by vectors of the form  $X_1^- x_{i_1, k_1}^- \dots X_h^- x_{i_h, k_h}^- X_{h+1}^- v$  where  $\lambda - \mu = \alpha_{i_1} + \dots + \alpha_{i_h}$ ,  $k_1, \dots, k_h \in \mathbb{Z}$ ,  $X_p^-$  is of the form  $X_p^- = x_{i_1, l_{1,p}}^- \dots x_{i_h, l_{h,p}}^-$  where  $m_1 + \dots + m_{h+1} = m$  and  $m_1, \dots, m_h \leq r^\vee$  (which is the crucial condition). It is proved by induction on  $h$  (see [CP3] section 5, (e)) with the help of the relations (10). Note that in [CP3]  $r^\vee = 3$ . Now it suffices to prove that  $\hat{U}_i.v$  is finite dimensional : indeed if  $m > r^\vee h + \dim(\hat{U}_i.v)$  we have  $m_{h+1} > \dim(\hat{U}_i.v)$  and  $X_{h+1}^- v = 0$ . It is shown exactly as in lemma 2.3 of [CP2] that  $\hat{U}_i.v$  is irreducible as  $\hat{U}_i$ -module, and so is finite dimensional.

For (2) : let us write  $\lambda - \mu = \alpha_{i_1} + \dots + \alpha_{i_h}$ . The result is proved by induction on  $h$ . We have seen that  $\hat{U}_i.v$  is finite dimensional. The induction is shown exactly as in [CP3] (section 5. (b)) by considering the  $L_{\lambda - \mu + \alpha_{i_j}}$  and with the help of relation (9).  $\square$

**4.3. Category  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ .** In the following by subcategory we mean full subcategory.

**Definition 4.10.** A  $\mathcal{U}_q(\mathfrak{h})$ -module  $V$  is said to be in the category  $\mathcal{O}(\mathcal{U}_q(\mathfrak{h}))$  if:

- i)  $V$  is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable
- ii) for all  $\omega \in \mathfrak{h}^*$ ,  $\dim(V_\omega) < \infty$
- iii) there is a finite number of element  $\lambda_1, \dots, \lambda_s \in \mathfrak{h}^*$  such that the weights of  $V$  are in  $\bigcup_{j=1 \dots s} D(\lambda_j)$

A  $\mathcal{U}_q(\mathfrak{g})$ -module (resp. a  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -module) is said to be in the category  $\mathcal{O}(\mathcal{U}_q(\mathfrak{g}))$  (resp.  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ) if it is in the category  $\mathcal{O}(\mathcal{U}_q(\mathfrak{h}))$  as a  $\mathcal{U}_q(\mathfrak{h})$ -module.

In particular we have a restriction functor  $\text{res} : \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}})) \rightarrow \mathcal{O}(\mathcal{U}_q(\mathfrak{g}))$ .

For example a highest weight  $\mathcal{U}_q(\mathfrak{g})$ -module is in the category  $\mathcal{O}(\mathcal{U}_q(\mathfrak{g}))$  and the product  $\otimes$  is well-defined on  $\mathcal{O}(\mathcal{U}_q(\mathfrak{g}))$ . An integrable  $l$ -highest weight module is in the category  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . But in general a  $l$ -highest weight module is not in the category  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ , indeed ( $\mathbb{C}_r[z]$  is the space of polynomials of degree lower than  $r$ ):

**Lemma 4.11.** Consider a  $l$ -weight  $(\omega, \Psi) \in P_l$  and  $i \in I$ . If  $\dim(L(\omega, \Psi)_{\omega - \alpha_i}) = r \in \mathbb{N}$  then there is  $P(z) \in \mathbb{C}_r[z]$  such that  $P(z)\Psi_i(z) = 0$  where  $\Psi_i(z) = \sum_{r \geq 0} (\Psi_{i,r}^+ z^r - \Psi_{i,-r}^- z^{-r})$ .

In particular the existence of a  $P(z) \in \mathbb{C}[z]$  such that  $P(z)\Psi_i(z) = 0$  for all  $i \in I$  is a necessary condition for  $L(\omega, \Psi) \in \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ .

*Proof:* Let  $v_0, v_1, \dots, v_r \in L(\omega, \Psi)$  such that:

$$L(\omega, \Psi)_\omega = \mathbb{C}v_0, \quad L(\omega, \Psi)_{\omega - \alpha_i} = \mathbb{C}v_1 \oplus \dots \oplus \mathbb{C}v_r$$

For  $m \in \mathbb{Z}$  let  $\Psi_{i,m} = \Psi_{i,m}^+ - \Psi_{i,m}^-$ . As  $x_{i,m}^+ \cdot v_0 = 0$ , we have:

$$x_{i,m}^+ x_{i,m'}^- \cdot v_0 = \frac{1}{q_i - q_i^{-1}} \Psi_{i,m+m'} v_0$$

As  $x_{i,m}^- \cdot v_0 \in L(\omega, \Psi)_{\omega - \alpha_i}$  and  $x_{i,m}^+ \cdot v_j \in L(\omega, \Psi)_\omega$ , there are  $\lambda_m^j, \mu_m^j \in \mathbb{C}$  ( $m \in \mathbb{Z}, 1 \leq j \leq r$ ) such that:

$$x_{i,m}^- \cdot v_0 = \lambda_m^1 v_1 + \dots + \lambda_m^r v_r, \quad x_{i,m}^+ \cdot v_j = \mu_m^j v_0$$

In particular we have:  $\Psi_{i,m+m'} = (q_i - q_i^{-1}) \sum_{j=1 \dots r} \lambda_{m'}^j \mu_m^j$ . We set  $\lambda^j(z) = \sum_{m' \in \mathbb{Z}} \lambda_{m'}^j z^{m'}$ ,  $\Psi_i(z) = \sum_{r \geq 0} \Psi_{i,r}^+ z^r - \Psi_{i,-r}^- z^{-r}$  and we have :

$$z^{-m} \Psi_i(z) = (q_i - q_i^{-1}) \sum_{j=1 \dots r} \mu_m^j \lambda^j(z)$$

So the  $\{\Psi_i(z), z\Psi_i(z), \dots, z^r \Psi_i(z)\}$  are not linearly independent.  $\square$

## 5. $q$ -CHARACTERS

For a quantum Kac-Moody algebra, one can define a character morphism as in the classical case. For quantum affine algebras a more precise morphism, called morphism of  $q$ -characters, was introduced by Frenkel-Reshetikhin [FR] (in particular to distinguish finite dimensional representations). In this section we generalize the construction of  $q$ -characters to quantum affinizations. The technical point is to add terms  $k_\lambda$  ( $\lambda \in \mathfrak{h}^*$ ) to make it well-defined in the general case. We prove a symmetry property of  $q$ -characters that generalizes a result of Frenkel-Mukhin : the image of  $\chi_q$  is the intersection of the kernels of screening operators (theorem 5.15).

**5.1. Reminder: classical character.** Let  $\mathcal{U}_q(\mathfrak{g})$  be a quantum Kac-Moody algebra. Let  $\mathcal{E} \subset (\mathfrak{h}^*)^{\mathbb{Z}}$  be the subset of  $c : \mathfrak{h}^* \rightarrow \mathbb{Z}$  such that  $c(\lambda) = 0$  for  $\lambda$  outside the union of a finite number of sets of the form  $D(\mu)$ . For  $\lambda \in \mathfrak{h}^*$  denote  $e(\lambda) \in \mathcal{E}$  such that  $e(\lambda)(\mu) = \delta_{\lambda, \mu}$ .  $\mathcal{E}$  has a natural structure of commutative  $\mathbb{Z}$ -algebra such that  $e(\lambda)e(\mu) = e(\lambda + \mu)$  (see [Ka]).

The classical character is the map  $\text{ch} : \mathcal{O}(\mathcal{U}_q(\mathfrak{g})) \rightarrow \mathcal{E}$  such that for  $V \in \mathcal{O}(\mathcal{U}_q(\mathfrak{g}))$ :

$$\text{ch}(V) = \sum_{\omega \in \mathfrak{h}^*} \dim(V_\omega) e(\omega)$$

$\text{ch}$  is a ring morphism and  $\text{ch}(L(\omega_1)) = \text{ch}(L(\omega_2)) \Rightarrow \omega_1 = \omega_2$ .

**5.2. Formal character.** Let  $\mathcal{U}_q(\hat{\mathfrak{g}})$  be a quantum affinization. In general the map  $\text{ch} \circ \text{res}$  does not distinguish the simple integrable representations in  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . That is why Frenkel-Reshetikhin [FR] introduced the theory of  $q$ -characters for quantum affine algebras. We generalize the construction for quantum affinizations.

Let  $V$  be in  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . For  $\omega \in \mathfrak{h}^*$ , the subspace  $V_\omega \subset V$  is stable by the operators  $\phi_{i, \pm m}^\pm$  ( $i \in I, m \geq 0$ ). Moreover they commute and  $[\phi_{i,m}^\pm, k_h] = 0$ , so we have a pseudo-weight space decomposition:

$$V_\omega = \bigoplus_{\gamma / (\omega, \gamma) \in P_i} V_{\omega, \gamma}$$

where  $V_{\omega, \gamma}$  is a simultaneous generalized eigenspace:

$$V_{\omega, \gamma} = \{x \in V_\omega / \exists p \in \mathbb{N}, \forall i \in \{1, \dots, n\}, \forall m \geq 0, (\phi_{i, \pm m}^\pm - \gamma_{i, \pm m}^\pm)^p \cdot x = 0\}$$

As  $V_\omega$  is finite dimensional the  $V_{\omega, \gamma}$  are finite dimensional.



Let  $\mathcal{E}_l \subset P_l^{\mathbb{Z}}$  be the ring of maps  $c : P_l \rightarrow \mathbb{Z}$  such that  $c(\lambda, \Psi) = 0$  for  $\lambda$  outside the union of a finite number of sets of the form  $D(\mu)$ .

**Definition 5.1.** *The formal character of a module  $V$  in the category  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  is  $ch_q(V) \in \mathcal{E}_l$  defined by:*

$$ch_q(V) = \sum_{(\mu, \Gamma) \in P_l} \dim(V_{\mu, \Gamma}) e(\mu, \Gamma)$$

We have the following commutative diagram:

$$\begin{array}{ccc} \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}})) & \xrightarrow{ch_q} & \mathcal{E}_l \\ \downarrow \text{res} & & \downarrow \beta \\ \mathcal{O}(\mathcal{U}_q(\mathfrak{g})) & \xrightarrow{ch} & \mathcal{E} \end{array}$$

where  $\beta : \mathcal{E}_l \rightarrow \mathcal{E}$  is constructed from the first projection  $\pi_1 : P_l \rightarrow P$ .

**5.3. Morphism of  $q$ -characters.** The combinatorics of formal characters can be studied with a morphism of  $q$ -characters  $\chi_q$  which is defined on a category  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  :

**5.3.1. The category  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ .** Denote by  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\mathfrak{g}))$  (resp.  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ) the category of integrable representations in the category  $\mathcal{O}(\mathcal{U}_q(\mathfrak{g}))$  (resp.  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ). For example a simple integrable  $l$ -highest weight  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -modules is in  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . Moreover:

**Proposition 5.2.** *For  $V$  a module in  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  there are  $P_{(\lambda, \Psi)} \geq 0$  ( $(\lambda, \Psi) \in P_l^+$ ) such that:*

$$ch_q(V) = \sum_{(\lambda, \Psi) \in P_l^+} P_{(\lambda, \Psi)} ch_q(L(\lambda, \Psi))$$

*Proof:* We have two preliminary points:

1) a submodule, a quotient of an integrable module is integrable.

2) for  $V \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  a module and  $\mu$  a maximal weight of  $V$ , then there is  $v \in V_{\mu}$  such that  $\mathcal{U}_q(\hat{\mathfrak{g}}).v$  is a  $l$ -highest weight module : indeed for  $(\mu, \gamma) \in P_l$  such that  $V_{\mu, \gamma} \neq \{0\}$  there is  $v \in V_{\mu, \gamma} - \{0\}$  such that  $\forall i \in I, r \geq 0, \phi_{i, \pm r}^{\pm}.v = \gamma_{i, \pm r}^{\pm}.v$  (because for all  $i \in I, r \geq 0, \text{Ker}(\phi_{i, \pm r}^{\pm} - \gamma_{i, \pm r}^{\pm}) \cap V_{\mu, \gamma} \neq \{0\}$ ).

The end of the proof is essentially made in [Ka] (proposition 9.7) : first we prove that for  $\lambda \in \mathfrak{h}^*$  there exists a filtration by a sequence of submodules in  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ :  $V = V_t \supset V_{t-1} \supset \dots \supset V_1 \supset V_0 = 0$  and  $J \subset \{1, \dots, t\}$  such that:

- (i) if  $j \in J$ , then  $V_j/V_{j-1} \simeq L(\lambda_j, \Psi_j)$  for some  $(\lambda_j, \Psi_j) \in P_l^+$  such that  $\lambda_j \geq \lambda$
- (ii) if  $j \notin J$ , then  $(V_j/V_{j-1})_{\mu} = 0$  for every  $\mu \geq \lambda$

(see the lemma 9.6 of [Ka]). Next for  $(\mu, \Psi) \in P_l^+$ , fix  $\lambda$  such that  $\mu \geq \lambda$  and introduce  $P_{(\mu, \Psi)}$  the number of times  $(\mu, \Psi)$  appears among the  $(\lambda_j, \Psi_j)$  (it is independent of the choice of the filtration and of  $\mu$ ). We conclude as in proposition 9.7 of [Ka].  $\square$

**Definition 5.3.**  $QP_l^+$  is the set of  $(\mu, \gamma) \in P_l$  satisfying the following condition :

i) there exist polynomials  $Q_i(z), R_i(z) \in \mathbb{C}[z]$  ( $i \in I$ ) of constant term 1 such that in  $\mathbb{C}[[z]]$  (resp. in  $\mathbb{C}[[z^{-1}]]$ ):

$$\sum_{m \geq 0} \gamma_{i, \pm m}^{\pm} z^{\pm m} = q_i^{\deg(Q_i) - \deg(R_i)} \frac{Q_i(zq_i^{-1})R_i(zq_i)}{Q_i(zq_i)R_i(zq_i^{-1})}$$

ii) there exist  $\omega \in P^+, \alpha \in Q^+$  satisfying  $\mu = \omega - \alpha$ .

In particular  $P_l^+ \subset QP_l^+$ .

**Proposition 5.4.** *Let  $V$  be a module in  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  and  $(\mu, \gamma) \in P_l$ . If  $\dim(V_{\mu, \gamma}) > 0$  then  $(\mu, \gamma) \in QP_l^+$ .*

*Proof:* The existence of the polynomials is shown as in [FR] (proposition 1): it reduces to the  $sl_2$ -case because for  $v \in V$ ,  $\hat{U}_i.v$  is finite dimensional. The existence of  $\omega \in P$  and  $\alpha \in Q^+$  is a consequence of proposition 5.2 and theorem 4.9.  $\square$

5.3.2. *Construction of  $q$ -characters.* Consider formal variables  $Y_{i,a}^\pm$  ( $i \in I, a \in \mathbb{C}^*$ ) and  $k_\omega$  ( $\omega \in \mathfrak{h}$ ). Let  $\tilde{A}$  be the commutative group of monomials of the form  $m = \prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{u_{i,a}(m)} k_{\omega(m)}$  ( $k_0 = 1$ ) where only a finite number of  $u_{i,a}(m) \in \mathbb{Z}$  are non zero,  $\omega(m) \in \mathfrak{h}$  (the coweight of  $m$ ), and such that for  $i \in I$ :

$$\alpha_i(\omega(m)) = r_i u_i(m) = r_i \sum_{a \in \mathbb{C}^*} u_{i,a}(m)$$

The product is given by  $u_{i,a}(m_1 m_2) = u_{i,a}(m_1) + u_{i,a}(m_2)$  and  $\omega(m_1 m_2) = \omega(m_1) + \omega(m_2)$ .

For example for  $i \in I, a \in \mathbb{C}^*$ , we have  $k_{\nu(\Lambda_i)} Y_{i,a} \in \tilde{A}$  because for  $j \in I$ ,  $\alpha_j(\nu(\Lambda_i)) = \Lambda_i(\nu(\alpha_j)) = r_j \Lambda_i(\alpha_j^\vee) = r_j \delta_{i,j}$ . For  $(\mu, \Gamma) \in QP_l^+$  we define  $Y_{\mu, \Gamma} \in \tilde{A}$  by:

$$Y_{\mu, \Gamma} = k_{\nu(\mu)} \prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{\beta_{i,a} - \gamma_{i,a}}$$

where  $\beta_{i,a}, \gamma_{i,a} \in \mathbb{Z}$  are defined by  $Q_i(u) = \prod_{a \in \mathbb{C}^*} (1 - ua)^{\beta_{i,a}}$ ,  $R_i(u) = \prod_{a \in \mathbb{C}^*} (1 - ua)^{\gamma_{i,a}}$ . We have  $Y_{\mu, \Gamma} \in \tilde{A}$  because for  $i \in I$ :

$$\alpha_i(\nu(\mu)) = \mu(\nu(\alpha_i)) = r_i \mu(\alpha_i^\vee) = r_i (\deg(Q_i) - \deg(R_i)) = r_i u_i(Y_{\mu, \Gamma})$$

For  $\chi \in \tilde{A}^{\mathbb{Z}}$  we say  $\chi \in \mathcal{Y}$  if there is a finite number of element  $\lambda_1, \dots, \lambda_s \in \mathfrak{h}^*$  such that the coweights of monomials of  $\chi$  are in  $\bigcup_{j=1 \dots s} \nu(D(\lambda_j))$ . In particular  $\mathcal{Y}$  has a structure of  $\mathfrak{h}$ -graded  $\mathbb{Z}$ -algebra.

**Definition 5.5.** *The  $q$ -character of a module  $V \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  is:*

$$\chi_q(V) = \sum_{(\mu, \Gamma) \in QP_l^+} d(\mu, \Gamma) Y_{\mu, \Gamma} \in \mathcal{Y}$$

where  $d(\mu, \Gamma) \in \mathbb{Z}$  is defined by  $ch_q(V) = \sum_{(\mu, \Gamma) \in QP_l^+} d(\mu, \Gamma) e(\mu, \Gamma)$ .

We have a commutative diagram :

$$\begin{array}{ccc} \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}})) & \xrightarrow{\chi_q} & \mathcal{Y} \\ \downarrow \text{res} & & \downarrow \beta \\ \mathcal{O}_{\text{int}}(\mathcal{U}_q(\mathfrak{g})) & \xrightarrow{\text{ch}} & \mathcal{E} \end{array}$$

where for  $m \in \tilde{A}$ ,  $\beta(m) = e(\omega(m))$ .

If  $C$  is of finite type then the weight of a monomial  $m \in \mathcal{Y}$  is  $\omega(m) = \sum_{i \in I} u_i(m) \nu(\Lambda_i)$ . So we can forget the  $k_h$ , and we get the  $q$ -characters defined in [FR]. In this case the integrable simple modules are finite dimensional.

Note that in the same way one can define the  $q$ -character of a finite dimensional  $\mathcal{U}_q(\hat{\mathfrak{h}})$ -module.

5.3.3. *Morphism of  $q$ -characters.* Denote by  $\text{Rep}(\mathcal{U}_q(\mathfrak{g}))$  (resp.  $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ) the Grothendieck group generated by the modules  $V$  in  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\mathfrak{g}))$  (resp.  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ) which have a composition series (a sequence of modules  $V \supset V_1 \supset V_2 \supset \dots$  such that  $V_i/V_{i+1}$  is irreducible).

The tensor product defines a ring structure on  $\text{Rep}(\mathcal{U}_q(\mathfrak{g}))$  and  $\text{ch}$  gives a ring morphism  $\chi : \text{Rep}(\mathcal{U}_q(\mathfrak{g})) \rightarrow \mathcal{E}$ .

The  $q$ -characters are compatible with exact sequences and so we get a group morphism  $\chi_q : \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) \rightarrow \mathcal{Y}$  which is called morphism of  $q$ -characters.

**Proposition 5.6.** *The morphism  $\chi_q$  is injective and the following diagram is commutative:*

$$\begin{array}{ccc} \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) & \xrightarrow{\chi_q} & \mathcal{Y} \\ \downarrow \text{res} & & \downarrow \beta \\ \text{Rep}(\mathcal{U}_q(\mathfrak{g})) & \xrightarrow{\chi} & \mathcal{E} \end{array}$$

The commutativity of the diagram follows from the definition. To see that  $\chi_q$  is injective, let us give some definitions:

A monomial  $m \in \tilde{A}$  is said to be dominant if  $u_{i,a}(m) \geq 0$  for all  $i \in I, a \in \mathbb{C}^*$ . If a  $l$ -weight  $(\omega, \Psi)$  belongs to  $P_l^+$  then  $Y_{(\omega, \Psi)} \in \tilde{A}$  is dominant. Moreover the map  $(\omega, \Psi) \mapsto Y_{(\omega, \Psi)}$  defines a bijection between  $P_l^+$  and dominant monomials. For  $m \in \tilde{A}$  a dominant monomial we denote by  $L(m) \in \mathcal{Y}$  the  $q$ -character of  $L(\omega, \Psi)$  where  $(\omega, \Psi)$  is the corresponding dominant  $l$ -weight. In particular  $L(m) = m +$  monomials of lower weight (in the sense of the ordering on  $P$ ), and so the  $L(m)$  are linearly independent.

A module with composition series is determined in the Grothendieck group by the multiplicity of the simple modules, and we have seen that the  $\chi_q(L(\lambda, \Psi))$  ( $(\lambda, \Psi) \in P_l^+$ ) are linearly independent in  $\mathcal{Y}$ . So  $\chi_q$  is injective.

**5.4.  $q$ -characters and universal  $\mathcal{R}$ -matrix.** The original definition of  $q$ -characters ([FR]) was based on an explicit formula for the universal  $\mathcal{R}$ -matrix established in [KT, LSS, Da]. In general no universal  $\mathcal{R}$ -matrix has been defined for a quantum affinization. However  $q$ -characters can be obtained with a piece of the formula of a “ $\mathcal{R}$ -matrix” in the same spirit as the original approach:

We refer to the chapter 3 of [Gu] for general background on  $h$ -formal deformations. Consider  $\mathcal{U}_h(\hat{\mathfrak{g}})$  the  $\mathbb{C}[[h]]$ -algebra which is  $h$ -topologically generated by  $\mathfrak{h}$  and the  $x_{i,r}^\pm$  ( $i \in I, r \in \mathbb{Z}$ ),  $h_{i,m}$  ( $i \in I, m \in \mathbb{Z} - \{0\}$ ) and with the relations of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  (where we set for  $\omega \in \mathfrak{h}$ ,  $k_\omega = \exp(h\omega)$ ). The subalgebra  $\mathcal{U}_h(\hat{\mathfrak{h}}) \subset \mathcal{U}_h(\hat{\mathfrak{g}})$  is  $h$ -topologically generated by  $\mathfrak{h}$  and the  $h_{i,m}$  ( $i \in I, m \in \mathbb{Z} - \{0\}$ ).

If  $V$  is a  $\mathcal{U}_q(\hat{\mathfrak{g}})$ -module (resp.  $\mathcal{U}_q(\hat{\mathfrak{h}})$ -module) which is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable then we have an algebra morphism  $\pi_V(h) : \mathcal{U}_h(\hat{\mathfrak{g}}) \rightarrow \text{End}(V)[[h]]$  (resp.  $\pi_V(h) : \mathcal{U}_h(\hat{\mathfrak{h}}) \rightarrow \text{End}(V)[[h]]$ ) (Remark : for  $\lambda \in \mathfrak{h}^*$ ,  $\omega \in \mathfrak{h}$ ,  $v \in V_\lambda$  we set  $\omega.v = \lambda(\omega)v$ ).

Define  $\mathcal{R}^0$  and  $T$  in  $\mathcal{U}_h(\hat{\mathfrak{h}}) \hat{\otimes} \mathcal{U}_h(\hat{\mathfrak{h}}) \subset \mathcal{U}_h(\hat{\mathfrak{g}}) \hat{\otimes} \mathcal{U}_h(\hat{\mathfrak{g}})$  ( $h$ -topological completion of the tensor product) by the formula :

$$\begin{aligned} \mathcal{R}^0 &= \exp(-(q - q^{-1}) \sum_{i,j \in I, m > 0} \frac{m}{[m]_q} \tilde{B}_{i,j}(q^m) h^m h_{i,m} \otimes h_{j,-m}) \\ T &= \exp(-h \sum_{1 \leq i \leq 2n-l} \Lambda_i^\vee \otimes \nu(\alpha_i)) \end{aligned}$$

Remark : we have the usual property of  $T$  (see [FR]): for  $\lambda, \mu \in \mathfrak{h}^*$ ,  $x \in V_\lambda$ ,  $y \in V_\mu$ , we have  $T.(x \otimes y) = q^{-(\lambda, \mu)}(x \otimes y)$ . Indeed:

$$\sum_{1 \leq i \leq 2n-l} \lambda(\Lambda_i^\vee) \mu(\nu(\alpha_i)) = (\mu, \sum_{1 \leq i \leq 2n-l} \lambda(\Lambda_i^\vee) \alpha_i) = (\mu, \lambda)$$

For  $i \in I, m \in \mathbb{Z} - \{0\}$  denote  $\tilde{h}_{i,m} = \sum_{j \in I} \tilde{C}_{j,i}(q^m) h_{j,m}$ . We have an inclusion  $\tilde{A} \subset \mathcal{U}_h(\hat{\mathfrak{h}})$  because the elements  $Y_{i,a}^\pm = k_{\mp \nu(\Lambda_i)} \exp(\mp(q - q^{-1}) \sum_{m \geq 1} h^m a^{-m} \tilde{h}_{i,m}) \in \mathcal{U}_h(\hat{\mathfrak{h}})$  ( $i \in I, a \in \mathbb{C}^*$ ) are algebraically independent.

**Theorem 5.7.** *For  $V$  a finite dimensional  $\mathcal{U}_q(\hat{\mathfrak{h}})$ -module,  $((\text{Tr}_V \circ \pi_V(h)) \otimes \text{Id})(\mathcal{R}^0 T) \in \mathcal{U}_h(\hat{\mathfrak{h}})$  is equal to  $\chi_q(V)$ .*

*Proof:* For  $(\lambda, \Psi) \in P_l$  consider  $V_{(\lambda, \Psi)}$  and  $((\text{Tr}_{V_{(\lambda, \Psi)}} \circ \pi_{V_{(\lambda, \Psi)}}(h)) \otimes \text{Id})(\mathcal{R}^0 T)$ . First we see as in [FR] that the term  $\mathcal{R}_0$  gives  $\prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{u_{i,a}(\Psi)}$ . But we have:

$$\sum_{1 \leq i \leq 2n-l} \lambda(\Lambda_i^\vee) \nu(\alpha_i) = \nu \left( \sum_{1 \leq i \leq 2n-l} \lambda(\Lambda_i^\vee) \alpha_i \right) = \nu(\lambda)$$

and so  $T$  gives  $k_{-\nu(\lambda)}$ .  $\square$

In general for  $V \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  we can consider a filtration  $(V_r)_{r \geq 0}$  of finite dimensional sub  $\mathcal{U}_q(\hat{\mathfrak{h}})$ -modules of  $V$  such that  $\bigcup_{r \geq 0} V_r = V$ ; so  $\chi_q(V)$  is the ‘‘limit’’ of the  $((\text{Tr}_{V_r} \circ \pi_{V_r}(h)) \otimes \text{Id})(\mathcal{R}^0 T)$  in  $\mathcal{Y}$ .

**5.5. Combinatorics of  $q$ -characters.** In this section we prove a symmetry property of general  $q$ -characters : the image of  $\chi_q$  is the intersection of the kernels of screening operators (theorem 5.15). Our proof is analog to the proof used by Frenkel-Mukhin [FM] for quantum affine algebras; however new technical points are involved because of the  $k_\lambda$  and infinite sums. In particular it shows that those  $q$ -characters are the combinatorial objects considered in [H3] (which were constructed in the kernel of screening operators).

In sections 5.5 and 6 we suppose that  $C(z)$  is invertible (it includes the cases of quantum affine algebras and quantum toroidal algebras, see section 2). We write  $\tilde{C}(z) = \frac{\tilde{C}'(z)}{d(z)}$  where  $d(z), \tilde{C}'_{i,j}(z) \in \mathbb{Z}[z^\pm]$ . For  $r \in \mathbb{Z}$  let  $p_{i,j}(r) = [(D(z)\tilde{C}'(z))_{i,j}]_r$  where for a Laurent polynomial  $P(z) \in \mathbb{Z}[z^\pm]$  we put  $P(z) = \sum_{r \in \mathbb{Z}} [P(z)]_r z^r$ .

**5.5.1. Construction of screening operators.** Let  $\mathcal{Y}^{\text{int}} \subset \mathcal{Y}$  be the subset consisting of those  $\chi \in \mathcal{Y}$  satisfying the following property : if  $\lambda$  is the coweight of a monomial of  $\chi$  there is  $K \geq 0$  such that  $k \geq K$  implies that for all  $i \in I$ ,  $\lambda \pm kr_i \alpha_i^\vee$  is not the coweight of a monomial of  $\chi$ .

**Lemma 5.8.**  $\mathcal{Y}^{\text{int}}$  is a subalgebra of  $\mathcal{Y}$  and  $\text{Im}(\chi_q) \subset \mathcal{Y}^{\text{int}}$ .

Consider the free  $\mathcal{Y}^{\text{int}}$ -module  $\tilde{\mathcal{Y}}_i = \prod_{a \in \mathbb{C}^*} \mathcal{Y}^{\text{int}} S_{i,a}$  and the linear map  $\tilde{S}_i : \mathcal{Y}^{\text{int}} \rightarrow \tilde{\mathcal{Y}}_i$  such that, for a monomial  $m$  :

$$\tilde{S}_i(m) = m \sum_{a \in \mathbb{C}^*} u_{i,a}(m) S_{i,a}$$

In particular  $\tilde{S}_i$  is a derivation. Let us choose a representative  $a$  for each class of  $\mathbb{C}^*/q_i^{2\mathbb{Z}}$  and consider:

$$\mathcal{Y}_i = \prod_{a \in \mathbb{C}^*/q_i^{2\mathbb{Z}}} \mathcal{Y}^{\text{int}} S_{i,a}$$

For  $i \in I$  and  $a \in \mathbb{C}^*$  we set:

$$A_{i,a} = k_i Y_{i,aq_i^{-1}} Y_{i,aq_i} \prod_{j/C_{j,i} < 0, r=C_{j,i}+1, C_{j,i}+3, \dots, -C_{j,i}-1} Y_{j,aq^r}^{-1} \in \tilde{A}$$

We have  $A_{i,a} \in \tilde{A}$  because for  $j \in I$ :  $\alpha_j(r_i \alpha_i^\vee) = r_i C_{i,j} = r_j C_{j,i} = r_j u_j(A_{i,a})$ .

We would like to see  $\mathcal{Y}_i$  as a quotient of  $\tilde{\mathcal{Y}}_i$  by the relations  $S_{i,aq_i} = A_{i,a} S_{i,aq_i^{-1}}$ . But the projection is not defined for all elements of  $\tilde{\mathcal{Y}}_i$  because there are infinite sums. However if  $\chi \in \mathcal{Y}^{\text{int}}$  and  $m$  is a monomial of  $\chi$  there is a finite number of monomials in  $\chi$  of the form  $m A_{i,aq_i}^{-1} A_{i,aq_i^3}^{-1} \dots A_{i,aq_i^r}^{-1}$  or of the form  $m A_{i,aq_i^{-1}} A_{i,aq_i^{-3}}^{-1} \dots A_{i,aq_i^{-r}}^{-1}$ . So the projection on  $\mathcal{Y}_i$  is well defined on  $\tilde{S}_i(\mathcal{Y}^{\text{int}}) \subset \tilde{\mathcal{Y}}_i$ . In particular we can define by projection of  $\tilde{S}_i$  the  $i^{\text{th}}$  screening operator  $S_i : \mathcal{Y}^{\text{int}} \rightarrow \mathcal{Y}_i$ .

The original definition for the finite case is in [FR].

5.5.2. *The morphism  $\tau_i$ .* Some operators  $\tau_i$  ( $i \in I$ ) were defined for the finite case in [FM]. We generalize the construction and the properties of the operators  $\tau_i$  (lemma 5.9 and 5.10).

Let  $i \in I$ . Denote  $\mathfrak{h}_i^\perp = \{\omega \in \mathfrak{h} / \alpha_i(\omega) = 0\}$ .

Consider formal variables  $k_r^{(i)}$  ( $r \in \mathbb{Z}$ ),  $k_\omega$  ( $\omega \in \mathfrak{h}$ ),  $Y_{i,a}^\pm$  ( $a \in \mathbb{C}^*$ ),  $Z_{j,c}$  ( $j \in I - \{i\}$ ,  $c \in \mathbb{C}^*$ ). Let  $\tilde{A}^{(i)}$  be the commutative group of monomials :

$$m = k_{r(m)}^{(i)} k_{\omega(m)} \prod_{a \in \mathbb{C}^*} Y_{i,a}^{u_{i,a}(m)} \prod_{j \in I, j \neq i, c \in \mathbb{C}^*} Z_{j,c}^{z_{j,c}(m)}$$

where only a finite number of  $u_{i,a}(m)$ ,  $z_{j,c}(m)$ ,  $r(m) \in \mathbb{Z}$  are non zero,  $\omega(m) \in \mathfrak{h}_i^\perp$  and such that  $r(m) = r_i u_i(m) = r_i \sum_{a \in \mathbb{C}^*} u_{i,a}(m)$ . The product is defined as for  $\tilde{A}$ . We call  $(r(m), \omega(m)) \in \mathbb{Z} \times \mathfrak{h}_i^\perp$  the coweight of the monomial  $m$ .

Let  $\tau_i : \tilde{A} \rightarrow \tilde{A}^{(i)}$  be the group morphism defined by ( $j \in I$ ,  $a \in \mathbb{C}^*$ ,  $\lambda \in \mathfrak{h}$ ):

$$\tau_i(Y_{j,a}) = Y_{i,a}^{\delta_{i,j}} \prod_{k \neq i, r \in \mathbb{Z}} Z_{k,aq^r}^{p_{k,aq^r}^{j,k(r)}}, \quad \tau_i(k_\lambda) = k_{\alpha_i(\lambda)}^{(i)} k_{\lambda - \alpha_i(\lambda)}^{\frac{\alpha_i^\vee}{2}}$$

(note that it is a formal definition because  $Y_{j,a} k_{\nu(\Lambda_j)} \in \tilde{A}$  but  $Y_{j,a} \notin \tilde{A}$ ). It is well defined because for  $m \in \tilde{A}$ ,  $\alpha_i(\omega(m)) = r_i u_i(m)$  and  $\alpha_i(\omega(m) - \alpha_i(\omega(m)) \frac{\alpha_i^\vee}{2}) = 0$ .

**Lemma 5.9.** *The morphism  $\tau_i$  is injective and for  $a \in \mathbb{C}^*$  we have:*

$$\tau_i(A_{i,a}) = k_{2r_i}^{(i)} Y_{i,aq_i^{-1}} Y_{i,aq_i}$$

*Proof:* Let  $m \in \tilde{A}$  such that  $\tau_i(m) = 1$ . For  $a \in \mathbb{C}^*$  we have  $u_{i,a}(m) = u_{i,a}(\tau_i(m)) = 0$ . For  $k \in I$ ,  $a \in \mathbb{C}^*$  denote  $u_{k,a}(m)(z) = \sum_{r \in \mathbb{Z}} u_{k,aq^r}(m) z^r \in \mathbb{Z}[z^\pm]$ . For  $j \in I - \{i\}$ , we have :

$$0 = z_{j,aq^R}(\tau_i(m)) = \sum_{k \in I, r+r'=R} p_{k,j}(r') u_{k,aq^r}(m) = [\sum_{k \in I} \tilde{C}'_{k,j}(z) u_{k,a}(m)(z)]_R$$

As  $\tilde{C}(z)$  is invertible we get  $u_{k,a}(m) = 0$  for all  $a \in \mathbb{C}^*$ . In particular for  $j \in I$  we have  $\alpha_j(\omega(m)) = r_j u_j(m) = 0$ . But  $\omega(m) - \alpha_i(\omega(m)) \frac{\alpha_i^\vee}{2} = 0 = \omega(m)$  and so  $m = 1$ .

For the second point let  $M = \tau_i(A_{i,a})$ . First for  $b \in \mathbb{C}^*$ ,  $u_{i,b}(M) = u_{i,b}(A_{i,a}) = \delta_{a/b, q_i} + \delta_{a/b, q_i^{-1}}$ . For  $R \in \mathbb{Z}$  and  $j \neq i$  we have:

$$z_{j,aq^R}(M) = [(\tilde{C}'(z)C(z))_{i,j}]_R = [(d(z)D(z))_{i,j}]_R = 0$$

Finally we have  $r(M) = r_i \alpha_i(\alpha_i^\vee) = -2r_i$  and  $\omega(M) = r_i \alpha_i^\vee - r_i \alpha_i(\alpha_i^\vee) \frac{\alpha_i^\vee}{2} = 0$ .  $\square$

Formally we have  $\tau_i(k_i) = k_{2r_i}^{(i)}$  and for  $j \in I - \{i\}$   $\tau_i(k_j) = k_{B_{j,i}}^{(i)} k_{\alpha_j}^{(i)}$  where  $\alpha_j^{(i)} = r_j \alpha_j^\vee - \frac{B_{j,i}}{2} \alpha_i^\vee$ . This motivates the following definition: for  $(r, \omega) \in \mathbb{Z} \times \mathfrak{h}_i^\perp$  denote :

$$D(r, \omega) = \{(r', \omega') \in \mathbb{Z} \times \mathfrak{h}_i^\perp / \omega' = \omega - \sum_{j \in I, j \neq i} m_j \alpha_j^{(i)}, r' = r - \sum_{j \in I, j \neq i} B_{j,i} m_j - 2r_i k / m_j, k \geq 0\}$$

Define  $\mathcal{Y}^{\text{int},(i)} \subset (\tilde{A}^{(i)})^\mathbb{Z}$  as the set of  $\chi$  such that :

i) there is a finite number of elements  $(r_1, \omega_1), \dots, (r_s, \omega_s) \in \mathbb{Z} \times \mathfrak{h}_i^\perp$  such that the coweights of monomials of  $\chi$  are in  $\bigcup_{j=1 \dots s} D(r_j, \omega_j)$ .

ii) for  $(r, \lambda)$  the coweight of a monomial of  $\chi$  there is  $K \geq 0$  such that  $k \geq K$  implies that for all  $j \in I$ ,  $j \neq i$ ,  $(r \pm B_{j,i} k, \lambda \pm k \alpha_j^{(i)})$  and  $(r \pm 2kr_i, \lambda)$  are not the coweight of a monomial of  $\chi$ .

In particular  $\mathcal{Y}^{\text{int},(i)}$  has a structure of  $\mathbb{Z} \times \mathfrak{h}_i^\perp$ -graded  $\mathbb{Z}$ -algebra.

The morphism  $\tau_i$  can be extended to a unique morphism of  $\mathbb{Z}$ -algebra  $\tau_i : \mathcal{Y}^{\text{int}} \rightarrow \mathcal{Y}^{\text{int},(i)}$ . Denote by  $\chi_q^i$  the morphism of  $q$ -characters for the algebra  $\mathcal{U}_{q_i}(\hat{sl}_2)$ .

**Lemma 5.10.** *Consider  $V \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  and a decomposition  $\tau_i(\chi_q(V)) = \sum_k P_k Q_k$  where*

*$P_k \in \mathbb{Z}[Y_{i,a}^{\pm}, k_{\pm r_i}^{(i)}]_{a \in \mathbb{C}^*}$ ,  $Q_k$  is a monomial in  $\mathbb{Z}[Z_{j,c}^{\pm}, k_h]_{j \neq i, a \in \mathbb{C}^*, h \in \mathfrak{h}_i^{\perp}}$  and all monomials  $Q_k$  are distinct. Then there exists a  $\hat{U}_i$ -module  $\bigoplus_k V_k$  isomorphic to the restriction of  $V$  to  $\hat{U}_i$  and such that  $\chi_q^i(V_k) = P_k$ .*

*Proof:* Let  $\mathcal{U}_q(\hat{\mathfrak{h}})_i^{\perp}$  the subalgebra of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  generated by the  $k_h$  ( $h \in \mathfrak{h}_i^{\perp}$ ),  $h_{j,m}$  ( $j \neq i, m \in \mathbb{Z} - \{0\}$ ). We can apply the proof of lemma 3.4 of [FM] with  $\hat{U}_i$  and  $\mathcal{U}_q(\hat{\mathfrak{h}})_i^{\perp}$  because :

i)  $\hat{U}_i$  and  $\mathcal{U}_q(\hat{\mathfrak{h}})_i^{\perp}$  commute in  $\mathcal{U}_q(\hat{\mathfrak{g}})$

ii) The image  $\omega - \alpha_i(\omega) \frac{\alpha_i^{\vee}}{2}$  in  $\mathfrak{h}_i^{\perp}$  of  $\omega \in \mathfrak{h}$  suffices to encode the action of the  $k_h$  ( $h \in \mathfrak{h}_i^{\perp}$ ) on a vector of weight  $\nu^{-1}(\omega) = \lambda$ . Indeed for  $h \in \mathfrak{h}_i^{\perp}$ , we have:

$$\lambda(h) = (\nu^{-1}(h), \nu^{-1}(\omega)) = \nu^{-1}(h)(\omega) = \nu^{-1}(h)(\omega - \omega(\alpha_i) \frac{\alpha_i^{\vee}}{2})$$

because  $\alpha_i(h) = 0 \Rightarrow \nu^{-1}(\alpha_i^{\vee}) = 0$ . □

5.5.3.  $\tau_i$  and screening operators. In this section we prove that  $\text{Im}(\chi_q) \subset \text{Ker}(S_i)$  (proposition 5.12) with a generalization of the proof of Frenkel-Mukhin [FM].

Consider the  $\mathcal{Y}^{\text{int},(i)}$ -module  $\tilde{\mathcal{Y}}_i^{(i)} = \prod_{a \in \mathbb{C}^*} \mathcal{Y}^{\text{int},(i)} S_{i,a}$  and the linear map  $\bar{S}_i : \mathcal{Y}^{\text{int},(i)} \rightarrow \tilde{\mathcal{Y}}_i^{(i)}$  such that, for a monomial  $m$  :

$$\bar{S}_i(m) = m \sum_{a \in \mathbb{C}^*} u_{i,a}(m) S_{i,a}$$

In particular  $\bar{S}_i$  is a derivation. Consider  $\mathcal{Y}_i^{(i)} = \prod_{a \in \mathbb{C}^*/q_i^{2\mathbb{Z}}} \mathcal{Y}^{\text{int},(i)} S_{i,a}$ . The  $\bar{S}_i(\mathcal{Y}^{\text{int},(i)}) \subset \tilde{\mathcal{Y}}_i^{(i)}$  can be projected in  $\mathcal{Y}_i^{(i)}$  by the relations :

$$S_{i,aq_i} = Y_{i,aq_i} Y_{i,aq_i^{-1}} k_{2r_i}^{(i)} S_{i,aq_i^{-1}}$$

and we get a derivation that we denote also by  $\bar{S}_i : \mathcal{Y}^{\text{int},(i)} \rightarrow \mathcal{Y}_i^{(i)}$ .

We also define a map  $\tau_i : \mathcal{Y}_i \rightarrow \mathcal{Y}_i^{(i)}$  in an obvious way (with the help of lemma 5.9). We see as in lemma 5.4 of [FM] that:

**Lemma 5.11.** *We have a commutative diagram:*

$$\begin{array}{ccc} \mathcal{Y}^{\text{int}} & \xrightarrow{S_i} & \mathcal{Y}_i \\ \downarrow \tau_i & & \downarrow \tau_i \\ \mathcal{Y}^{\text{int},(i)} & \xrightarrow{\bar{S}_i} & \mathcal{Y}_i^{(i)} \end{array}$$

With the help of lemma 5.9, 5.10 and 5.11 we see as in corollary 5.5 of [FM]:

**Lemma 5.12.** *We have  $\text{Im}(\chi_q) \subset \bigcap_{i \in I} \text{Ker}(S_i)$ .*

In the following we denote  $\mathfrak{K}_i = \text{Ker}(S_i)$  and  $\mathfrak{K} = \bigcap_{i \in I} \mathfrak{K}_i$ .

**Lemma 5.13.** *An element  $\chi \in \mathcal{Y}^{\text{int}}$  is in  $\mathfrak{K}_i$  if and only if it can be written in the form  $\chi = \sum_k P_k Q_k$  where*

*$P_k \in \mathbb{Z}[k_{\nu(\Lambda_i)} Y_{i,a}(1 + A_{i,aq_i}^{-1})]_{a \in \mathbb{C}^*}$ ,  $Q_k$  is a monomial in  $\mathbb{Z}[Y_{j,a}^{\pm}, k_h]_{j \neq i, a \in \mathbb{C}^*, h \in P_i^{*,\perp}}$  and all monomials  $Q_k$  are distinct.*

*Proof:* We use the result for the  $sl_2$ -case which is proved in [FR]. First an element of this form is in  $\mathfrak{K}_i$ . Consider  $\chi \in \mathfrak{K}_i$  and write  $\tau_i(\chi) = \sum_k P'_k Q'_k$  as in lemma 5.10. From lemma 5.11 we have  $0 = \overline{S}_i(\chi) = \sum_k \overline{S}_i(P'_k) Q'_k$ . So all  $\overline{S}_i(P'_k) = 0$  and it follows from the  $sl_2$ -case that  $P'_k \in \mathbb{Z}[Y_{i,a} k_{r_i}^{(i)} + Y_{i,aq_i}^{-1} k_{-r_i}^{(i)}]_{a \in \mathbb{C}^*}$ . The lemma 5.9 lead us to the conclusion.  $\square$

5.5.4. *Description of  $\text{Im}(\chi_q)$ .* Dominant monomials are defined in section 5.3.3. We have:

**Lemma 5.14.** *An element  $\chi \in \mathfrak{K}$  has at least one dominant monomial.*

With the help of lemma 5.13 we can use the proof of lemma 5.6 of [FM] (see also the proof of theorem 4.9 in [H1] at  $t = 1$ ).

**Theorem 5.15.** *We have  $\text{Im}(\chi_q) = \mathfrak{K}$ . Moreover the elements of  $\mathfrak{K}$  are the sums:*

$$\sum_{m \text{ dominant}} \lambda_m L(m)$$

where  $\lambda_m = 0$  for  $\omega(m)$  outside the union of a finite number of sets of the form  $D(\mu)$ .

*Proof:* The inclusion  $\text{Im}(\chi_q) \subset \mathfrak{K}$  is proved in lemma 5.12. For the other one, consider  $\chi \in \mathfrak{K}$ . We can suppose that the weights of  $\chi$  are in a set  $D(\lambda)$  (because the weights of each  $L(m)$  are in a set  $D(\mu)$ ). We define by induction  $L^{(k)}(m) \in \text{Im}(\chi_q)$  ( $k \geq 0$ ) in the following way: we set  $L^{(0)} = \sum_{\omega(m)=\lambda} [\chi]_m L(m)$ . If

$L^{(k)}$  is defined, we consider the set  $\tilde{A}_{k+1}$  of monomials  $m'$  which appear in  $\chi - L^{(k)}$  such that  $\lambda - \omega(m') = m_1 r_1 \alpha_1^\vee + \dots + m_n r_n \alpha_n^\vee$  where  $m_1, \dots, m_n \geq 0$  and  $m_1 + \dots + m_n = k$ . We set:

$$L^{(k+1)} = L^{(k)} + \sum_{m' \in \tilde{A}_{k+1}} [\chi - L^{(k)}]_{m'} L(m')$$

Then we set  $L^\infty = \sum_{k \geq 0 / m \in \tilde{A}_k} [L^{(k)}]_m L(m) \in \text{Im}(\chi_q)$  and it follows from lemma 5.14 that  $L^\infty = \chi$ .  $\square$

Note that proposition 5.2 gives that for  $\chi_q(V)$  ( $V$  module in  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ) the  $\lambda_m$  are non-negative.

Remark: for  $m \in \tilde{A}$  a dominant monomial we prove in the same way that there is a unique  $F(m) \in \mathfrak{K}$  such that  $m$  has coefficient 1 in  $F(m)$  and  $m$  is the unique dominant monomial in  $F(m)$ . In the finite case an algorithm was given by Frenkel-Mukhin [FM] to compute the  $F(m)$ . In [H3] we extended the definition of the algorithm for generalized Cartan matrix and showed that it is well-defined if  $i \neq j \Rightarrow C_{i,j} C_{j,i} \leq 3$  (see also [H2] for the detailed description of this algorithm at  $t = 1$ ). Theorem 5.15 allows us to prove two results announced in [H3]: the algorithm is well defined for

$$A_1^{(1)} \text{ (with } r_1 = r_2 = 2) \text{ because } \det(C(z)) = z^4 - z^2 - z^{-2} + z^{-4} \neq 0$$

$$A_2^{(2)} \text{ (with } r_1 = 4, r_2 = 1) \text{ because } \det(C(z)) = z^5 - z - z^{-1} + z^{-5} \neq 0$$

But for  $A_1^{(1)}$  (with  $r_1 = r_2 = 1$ ) we have  $\det(C(z)) = 0$ ; we observed in [H3] that the algorithm is not well-defined in this case.

## 6. DRINFEL'D NEW COPRODUCT AND FUSION PRODUCT

Our study of combinatorics of  $q$ -characters gives a ring structure on  $\text{Im}(\chi_q)$  (corollary 6.1). As  $\chi_q$  is injective we get an induced ring structure on the Grothendieck group. In this section we prove that it is a fusion product (theorem 6.2), that is to say that the product of two modules is a module. We use the Drinfel'd new coproduct (proposition 6.3); as it involves infinite sums, we have to work in a larger category where the tensor product is well-defined (theorem 6.7). To end the proof of theorem 6.2 we define specializations of certain forms which allow us to go from the larger category to  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . Note that in our construction we do not use that  $C(z)$  is invertible.

6.1. **Fusion product.** As the  $S_i$  are derivations, theorem 5.15 gives:

**Corollary 6.1.** *Im( $\chi_q$ ) is a subring of  $\mathcal{Y}$ .*

Since  $\chi_q$  is injective on  $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ , the product of  $\mathcal{Y}$  gives an induced commutative product  $*$  on  $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . For  $(\lambda, \Psi), (\lambda', \Psi') \in P_l^+$  there are  $Q_{\lambda, \Psi, \lambda', \Psi'}(\mu, \Phi) \in \mathbb{Z}$  such that:

$$L(\lambda, \Psi) * L(\lambda', \Psi') = L(\lambda + \lambda', \Psi\Psi') + \sum_{(\mu, \Phi) \in P_l^+ / \mu < \lambda + \lambda'} Q_{\lambda, \Psi, \lambda', \Psi'}(\mu, \Phi) L(\mu, \Phi)$$

We will interpret this product as a fusion product related to the basis of simple modules : that is to say we will show that a product of modules is a module (see [F] for generalities on fusion rings and physical motivations). Let us explain it in more details : consider :

$$\text{Rep}^+(\mathcal{U}_q(\hat{\mathfrak{g}})) = \bigoplus_{(\lambda, \Psi) \in P_l^+} \mathbb{N}.L(\lambda, \Psi) \subset \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) = \bigoplus_{(\lambda, \Psi) \in P_l^+} \mathbb{Z}.L(\lambda, \Psi)$$

**Theorem 6.2.** *The subset  $\text{Rep}^+(\mathcal{U}_q(\hat{\mathfrak{g}})) \subset \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  is stable by  $*$ .*

In this section 6 we prove this theorem by interpreting  $*$  with the help of a generalization of the new Drinfel'd coproduct. Note that theorem 6.2 means that for  $(\lambda, \Psi), (\lambda', \Psi') \in P_l^+$  we have  $Q_{\lambda, \Psi, \lambda', \Psi'}(\mu, \Phi) \geq 0$ .

## 6.2. Coproduct.

6.2.1. *Reminder: case of a quantum affine algebra and Drinfel'd-Jimbo coproduct.* As said before the case of a quantum affine algebra is a very special one because there are two realizations (if we add a central charge); in particular there is a coproduct on  $\mathcal{U}_q(\hat{\mathfrak{g}})$ , a tensor product on  $\mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  and  $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  is a ring. It is the product  $*$  because it is shown in [FR] that  $\chi_q$  is a ring morphism. In particular the tensor product is commutative. So theorem 6.2 is proved in this case.

6.2.2. *General case : new Drinfel'd coproduct.* In general we have a coproduct  $\Delta_{\hat{\mathfrak{g}}} : \mathcal{U}_q(\hat{\mathfrak{h}}) \rightarrow \mathcal{U}_q(\hat{\mathfrak{h}}) \otimes \mathcal{U}_q(\hat{\mathfrak{h}})$  for the commutative algebra  $\mathcal{U}_q(\hat{\mathfrak{h}})$  defined by ( $h \in P^*, i \in I, m \neq 0$ ):

$$\Delta_{\hat{\mathfrak{g}}}(k_h) = k_h \otimes k_h, \Delta_{\hat{\mathfrak{g}}}(h_{i,m}) = 1 \otimes h_{i,m} + h_{i,m} \otimes 1$$

In particular we have ( $i \in I, m \geq 0$ ) :  $\Delta_{\hat{\mathfrak{g}}}(\phi_{i,\pm m}^{\pm}) = \sum_{0 \leq l \leq m} \phi_{i,\pm(m-l)}^{\pm} \otimes \phi_{i,\pm l}^{\pm}$ .

No coproduct has been defined for the entire  $\mathcal{U}_q(\hat{\mathfrak{g}})$ . However Drinfel'd (unpublished note, see also [DI, DF]) defined for  $\mathcal{U}_q(\hat{\mathfrak{sl}}_n)$  a map which behaves as a new coproduct adapted to the affinization realization. In this section we use those formulas for general quantum affinizations; as infinite sums are involved, we use a formal parameter  $u$  so that it makes sense.

Let  $\mathcal{C} = \mathbb{C}((u))$  be the field of Laurent series  $\sum_{r \geq R} \lambda_r u^r$  ( $R \in \mathbb{Z}, \lambda_r \in \mathbb{C}$ ). The algebra  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  is defined in section 3.3.1. Consider the  $\mathcal{C}$ -algebra  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) = \mathcal{C} \otimes \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$  (resp.  $\mathcal{U}'_q(\hat{\mathfrak{g}}) = \mathcal{C} \otimes \mathcal{U}_q(\hat{\mathfrak{g}})$ ). Let  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \hat{\otimes} \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) = (\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}) \otimes_{\mathbb{C}} \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}))((u))$  be the  $u$ -topological completion of  $\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}) \otimes_{\mathbb{C}} \tilde{\mathcal{U}}_q(\hat{\mathfrak{g}})$ . It is also a  $\mathcal{C}$ -algebra.

**Proposition 6.3.** *There is a unique morphism of  $\mathcal{C}$ -algebra  $\Delta_u : \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \rightarrow \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \hat{\otimes} \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$  such that for  $i \in I, r \in \mathbb{Z}, m \geq 0, h \in \mathfrak{h}$ :*

$$\Delta_u(x_{i,r}^+) = x_{i,r}^+ \otimes 1 + \sum_{l \geq 0} u^{r+l} (\phi_{i,-l}^- \otimes x_{i,r+l}^+)$$

$$\Delta_u(x_{i,r}^-) = u^r (1 \otimes x_{i,r}^-) + \sum_{l \geq 0} u^l (x_{i,r-l}^- \otimes \phi_{i,l}^+)$$

$$\Delta_u(\phi_{i,\pm m}^{\pm}) = \sum_{0 \leq l \leq m} u^{\pm l} (\phi_{i,\pm(m-l)}^{\pm} \otimes \phi_{i,\pm l}^{\pm}), \Delta_u(k_h) = k_h \otimes k_h$$



*Proof:* We can easily check the compatibility with relations (11), (12), (13), (14), (15), (16) because  $\Delta_u$  can also be given in terms of the currents of section 3.2: we have in  $(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \otimes_{\mathcal{C}} \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))[[z, z^{-1}]]$  :

$$\begin{aligned} \Delta_u(x_i^+(z)) &= x_i^+(z) \otimes 1 + \phi_i^-(z) \otimes x_i^+(zu), \quad \Delta_u(x_i^-(z)) = 1 \otimes x_i^-(zu) + x_i^-(z) \otimes \phi_i^+(zu) \\ \Delta_u(\phi_i^\pm(z)) &= \phi_i^\pm(z) \otimes \phi_i^\pm(zu) \end{aligned}$$

□

Remark 1 : If  $C$  is finite or simply laced then  $\Delta_u$  is compatible with the affine quantum Serre relations (relations (10)) and can be defined for  $\mathcal{U}'_q(\hat{\mathfrak{g}})$  (see [DI] for finite symmetric cases and [E, Gr] for other finite cases). We conjecture that it is also true for general  $C$ , but we do not need it for our purposes.

Remark 2 : let  $T : \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \rightarrow \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$  be the  $\mathbb{Z}$ -gradation morphism defined by  $T(x_{i,r}^\pm) = u^r x_{i,r}^\pm$ ,  $T(\phi_{i,m}^\pm) = u^m \phi_{i,m}^\pm$ ,  $T(k_n) = k_n$ . The  $u$  is put in such a way that  $\Delta_u = (\text{Id} \otimes T) \circ \Delta$  where  $\Delta$  is the usual new Drinfel'd coproduct (without  $u$ ).

Remark 3 : The map  $\Delta_u$  is not coassociative, indeed in  $(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \otimes_{\mathcal{C}} \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \otimes_{\mathcal{C}} \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))[z]$ :

$$\begin{aligned} ((\Delta_u \otimes \text{Id}) \circ \Delta_u)(\phi_i^+(z)) &= \phi_i^+(z) \otimes \phi_i^+(uz) \otimes \phi_i^+(uz) \\ ((\text{Id} \otimes \Delta_u) \circ \Delta_u)(\phi_i^+(z)) &= \phi_i^+(z) \otimes \phi_i^+(uz) \otimes \phi_i^+(u^2z) \end{aligned}$$

Remark 4 : Although it is not defined in a strict sense, the ‘‘limit’’ of  $\Delta_u$  at  $u = 1$  is coassociative. On  $\mathcal{U}_q(\hat{\mathfrak{h}})$  the limit at  $u = 1$  makes sense and is  $\Delta_{\hat{\mathfrak{h}}}$ .

**6.3. Tensor products of representations of  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ .** As the coproduct involves infinite sums, we have to introduce a category larger than  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  in order to define tensor products:

6.3.1. *The category  $\mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ .*

**Definition 6.4.** *The set of  $l, u$ -weights  $P_{l,u}$  is the set of couple  $(\lambda, \Psi(u))$  such that  $\lambda \in \mathfrak{h}^*$ ,  $\Psi(u) = (\Psi_{i,\pm m}^\pm(u))_{i \in I, m \geq 0}$ ,  $\Psi_{i,\pm m}^\pm(u) \in \mathbb{C}[u, u^{-1}]$  and  $\Psi_{i,0}^\pm(u) = q_i^{\pm \lambda(\alpha_i^\vee)}$ .*

**Definition 6.5.** *An object  $V$  of the category  $\mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  is a  $\mathcal{C}$ -vector space with a structure of  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ -module such that:*

- i)  $V$  is  $\mathcal{U}_q(\mathfrak{h})$ -diagonalizable
- ii) For all  $\lambda \in \mathfrak{h}^*$ , the sub  $\mathcal{C}$ -vector space  $V_\lambda \subset V$  is finite dimensional
- iii) there are a finite number of element  $\lambda_1, \dots, \lambda_s \in \mathfrak{h}^*$  such that the weights of  $V$  are in  $\bigcup_{j=1 \dots s} D(\lambda_j)$
- iv) for  $\lambda \in \mathfrak{h}^*$ ,  $V_\lambda = \bigoplus_{(\lambda, \Psi(u)) \in P_{l,u}} V_{(\lambda, \Psi(u))}$  where:

$$V_{\lambda, \Psi(u)} = \{x \in V_\lambda / \exists p \in \mathbb{N}, \forall i \in \{1, \dots, n\}, \forall r \geq 0, (\phi_{i,\pm r}^\pm - \Psi_{i,\pm r}^\pm(u))^p \cdot x = 0\}$$

The property iv) is added because  $\mathcal{C}$  is not algebraically closed.

The scalar extension and the projection  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \rightarrow \mathcal{U}_q(\hat{\mathfrak{g}})$  gives an injection  $i : \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}})) \rightarrow \mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  such that for  $V \in \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ ,  $i(V) = V \otimes \mathcal{C}$ .

Let  $\mathcal{E}_{l,u} \subset P_{l,u}^{\mathbb{Z}}$  be defined as  $\mathcal{E}_l$ . The formal character of a module  $V$  in the category  $\mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  is:

$$\text{ch}_{q,u}(V) = \sum_{(\mu, \Gamma(u)) \in P_{l,u}} \dim_{\mathcal{C}}(V_{\mu, \Gamma(u)}) e(\mu, \Gamma(u)) \in \mathcal{E}_{l,u}$$

We have a map  $i_{\mathcal{E}} : \mathcal{E}_l \rightarrow \mathcal{E}_{l,u}$  such that  $i_{\mathcal{E}}((\lambda, \Psi)) = (\lambda, (\Psi_{i,\pm m}^{\pm m}))$  and a commutative diagram:

$$\begin{array}{ccc} \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}})) & \xrightarrow{\text{ch}_q} & \mathcal{E}_l \\ \downarrow i & & \downarrow i_{\mathcal{E}} \\ \mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})) & \xrightarrow{\text{ch}_{q,u}} & \mathcal{E}_{l,u} \end{array}$$

In an analogous way one defines the category  $\mathcal{O}(\mathcal{U}'_q(\hat{\mathfrak{g}}))$  and a formal character  $\text{ch}_{q,u}$  on  $\mathcal{O}(\mathcal{U}'_q(\hat{\mathfrak{g}}))$ .

6.3.2. *Tensor products.* We consider subcategories of  $\mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ . Let  $R \in \mathbb{Z}$ ,  $R \geq 0$ :

**Definition 6.6.**  $\mathcal{O}^R(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  is the category of modules  $V \in \mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  such that for all  $\lambda \in \mathfrak{h}^*$ , there is a  $\mathbb{C}$ -basis  $(v_\alpha^\lambda)_\alpha$  of  $V_\lambda$  satisfying :

i) for all  $m \in \mathbb{Z}$ ,  $\alpha, \beta$ , the coefficient of  $x_{i,m}^+ \cdot v_\alpha^\lambda$  on  $v_\beta^{\lambda+\alpha_i}$  (resp. of  $x_{i,m}^- \cdot v_\alpha^\lambda$  on  $v_\beta^{\lambda-\alpha_i}$ ) is in  $\mathbb{C}[[u]]$  if  $m \geq 0$ , in  $u^{Rm}\mathbb{C}[[u]]$  if  $m \leq 0$ .

ii) for all  $m \geq 0$ ,  $\alpha, \beta$  the coefficient of  $\phi_{i,-m}^- \cdot v_\alpha^\lambda$  on  $v_\beta^\lambda$  is in  $u^{-mR}\mathbb{C}[[u]]$

iii) for all  $m \geq 0$ ,  $\alpha, \beta$  the coefficient of  $\phi_{i,m}^+ \cdot v_\alpha^\lambda$  on  $v_\beta^\lambda$  is in  $\mathbb{C}[[u]]$

Example : For  $V \in \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ , we have  $i(V) \in \mathcal{O}^0(\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}))$ .

**Theorem 6.7.** Let  $V_1 \in \mathcal{O}(\tilde{\mathcal{U}}_q(\hat{\mathfrak{g}}))$  and  $V_2 \in \mathcal{O}^R(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ . Then  $\Delta_u$  defines a structure of  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ -module on  $i(V_1) \otimes_{\mathbb{C}} V_2$  which is in  $\mathcal{O}^{R+1}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ . Moreover the  $l, u$ -weights of  $i(V_1) \otimes_{\mathbb{C}} V_2$  are of the form  $(\lambda_1 + \lambda_2, \gamma_1(z)\gamma_2(uz))$  where  $(\lambda_1, \gamma_1)$  is a  $l, u$ -weight of  $V_1$  and  $(\lambda_2, \gamma_2)$  is a  $l, u$ -weight of  $V_2$ .

Remark :  $\gamma(u)(z) = \gamma_1(z)\gamma_2(uz)$  means that for  $i \in I, m \geq 0$  :

$$\gamma_{i,\pm m}^\pm(u) = \sum_{0 \leq l \leq m} (\gamma_1)_{i,\pm l}(u)(\gamma_2)_{i,\pm(m-l)}(u)u^{\pm(m-l)}$$

*Proof:* As the definition of  $\Delta_u$  involves infinite sums, we have to prove that the action formally defined by  $\Delta_u$  makes sense on  $V'_1 \otimes_{\mathbb{C}} V_2$  where we denote  $V'_1 = i(V_1)$ . Indeed the weight spaces of  $V'_1$  and  $V_2$  are finite dimensional and for  $\lambda, \mu \in \mathfrak{h}^*$  we can use a  $\mathbb{C}$ -base  $(v_\alpha^{1,\lambda})_\alpha$  of  $(V_1)_\lambda$  as a  $\mathbb{C}$ -base of  $(V'_1)_\lambda$  and the  $\mathbb{C}$ -basis  $(v_{\alpha'}^{2,\mu})$  of  $(V_2)_\mu$  given by the definition of  $\mathcal{O}^R(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ . So consider  $\lambda, \mu \in \mathfrak{h}^*$ ,  $i \in I$  and let us investigate the coefficients ( $r \in \mathbb{Z}, m \geq 0$ ):

we have  $x_{i,r}^+ \cdot ((V'_1)_\lambda \otimes (V_2)_\mu) \subset (V'_1)_{\lambda+\alpha_i} \otimes (V_2)_\mu \oplus (V'_1)_\lambda \otimes (V_2)_{\mu+\alpha_i}$ .

on  $(V'_1)_\lambda \otimes (V_2)_{\mu+\alpha_i}$  : the coefficient of  $x_{i,m}^+ \cdot (v_\alpha^{1,\lambda} \otimes v_{\alpha'}^{2,\mu})$  on  $v_\beta^{1,\lambda} \otimes v_{\beta'}^{2,\mu+\alpha_i}$  is in  $\sum_{l \geq 0} u^{r+l}\mathbb{C}[[u]] \subset \mathbb{C}[[u]]$

if  $r \geq 0$ , in  $\sum_{l \geq 0} u^{r+l}u^{R(r+l)}\mathbb{C}[[u]] \subset u^{(R+1)r}\mathbb{C}[[u]]$  if  $r \leq 0$ .

on  $(V'_1)_{\lambda+\alpha_i} \otimes (V_2)_\mu$  : the coefficient of  $x_{i,r}^+ \cdot (v_\alpha^{1,\lambda} \otimes v_{\alpha'}^{2,\lambda})$  on  $v_\beta^{1,\lambda+\alpha_i} \otimes v_{\beta'}^{2,\mu}$  is in  $\mathbb{C}$ .

we have  $x_{i,r}^- \cdot ((V'_1)_\lambda \otimes (V_2)_\mu) \subset (V'_1)_{\lambda-\alpha_i} \otimes (V_2)_\mu \oplus (V'_1)_\lambda \otimes (V_2)_{\mu-\alpha_i}$ .

on  $(V'_1)_\lambda \otimes (V_2)_{\mu-\alpha_i}$  : the coefficient of  $x_{i,r}^- \cdot (v_\alpha^{1,\lambda} \otimes v_{\alpha'}^{2,\mu})$  on  $v_\beta^{1,\lambda} \otimes v_{\beta'}^{2,\mu-\alpha_i}$  is in  $u^r\mathbb{C}[[u]] \subset \mathbb{C}[[u]]$  if  $r \geq 0$ , in  $u^r u^{Rr}\mathbb{C}[[u]]$  if  $r \leq 0$ .

on  $(V'_1)_{\lambda-\alpha_i} \otimes (V_2)_\mu$  : the coefficient of  $x_{i,r}^- \cdot (v_\alpha^{1,\lambda} \otimes v_{\alpha'}^{2,\mu})$  on  $v_\beta^{1,\lambda-\alpha_i} \otimes v_{\beta'}^{2,\mu}$  is in  $\sum_{l \geq 0} u^l\mathbb{C}[[u]] \subset \mathbb{C}[[u]]$ .

we have  $\phi_{i,\pm m}^\pm \cdot ((V'_1)_\lambda \otimes (V_2)_\mu) \subset ((V'_1)_\lambda \otimes (V_2)_\mu)$ .

the coefficient of  $\phi_{i,m}^+ \cdot (v_\alpha^{1,\lambda} \otimes v_{\alpha'}^{2,\mu})$  on  $v_\beta^{1,\lambda} \otimes v_{\beta'}^{2,\mu}$  is in  $\sum_{0 \leq l \leq m} u^l\mathbb{C}[[u]] \subset \mathbb{C}[[u]]$ .

the coefficient of  $\phi_{i,-m}^- \cdot (v_\alpha^{1,\lambda} \otimes v_{\alpha'}^{2,\mu})$  on  $v_\beta^{1,\lambda} \otimes v_{\beta'}^{2,\mu}$  is in  $\sum_{0 \leq l \leq m} u^{-l}u^{-lR}\mathbb{C}[[u]] \subset u^{-m(R+1)}\mathbb{C}[[u]]$ .

So we have a structure of  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ -module on  $V'_1 \otimes_{\mathbb{C}} V_2$ . Let us prove that it is in  $\mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ . We verify the properties of definition 6.5: i) ii) iii) are clear because the restriction of  $\Delta_u$  to  $\mathcal{U}_q(\hat{\mathfrak{h}})$  is the restriction of  $\Delta_{\hat{\mathfrak{h}}}$ . For iv) we note that for  $(\lambda_1, \gamma_1), (\lambda_2, \gamma_2) \in P_{l,u}$ , the  $(V'_1)_{\lambda_1, \gamma_1} \otimes (V_2)_{\lambda_2, \gamma_2}$  is in the pseudo weight space of  $l, u$ -weight  $(\lambda_1 + \lambda_2, \gamma_1(z)\gamma_2(zu))$  because  $\Delta_u(\phi_i^\pm(z)) = \phi_i^\pm(z) \otimes \phi_i^\pm(zu)$  (it also proves the last point of the proposition).

Finally we see in the above computations that the coefficients verify the property of  $\mathcal{O}^{R+1}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ , so  $V_1' \otimes_C V_2$  is in  $\mathcal{O}^{R+1}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ .  $\square$

**Definition 6.8.** For  $R \geq 0$ , we denote  $\otimes_R : \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}})) \times \mathcal{O}^R(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})) \rightarrow \mathcal{O}^{R+1}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  the bilinear map constructed in theorem 6.7.

See section 6.6 for explicit examples. For  $R \geq 2$  and  $V_1, V_2, \dots, V_R \in \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ , one can define the iterated tensor product  $V_1 \otimes_{R-2} (V_2 \otimes_{R-3} (\dots \otimes_0 V_R) \dots)$  which is in  $\mathcal{O}^{R-1}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ .

#### 6.4. Simple modules of $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ .

6.4.1. *l, u-highest weight modules.* For  $(\lambda, \Psi(u)) \in P_{l,u}$ , let  $\tilde{M}(\lambda, \Psi(u))$  be the Verma  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$  module of highest weight  $(\lambda, \Psi(u))$  (it is non trivial thanks to the triangular decomposition of  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$  in lemma 3.10). So we have an analog of proposition 4.6 : for  $(\lambda, \Psi(u)) \in P_{l,u}$ , there is a unique up to isomorphism simple  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ -module  $\tilde{L}(\lambda, \Psi(u))$  of *l, u*-highest weight  $(\lambda, \Psi(u))$  that is to say that there is  $v \in \tilde{L}(\lambda, \Psi(u))$  such that  $(i \in I, r \in \mathbb{Z}, m \geq 0, h \in \mathfrak{h})$ :

$$x_{i,r}^+ \cdot v = 0, \tilde{L}(\lambda, \Psi(u)) = \tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}) \cdot v, \phi_{i,\pm m}^\pm \cdot v = \Psi_{i,\pm m}^\pm(u)v, k_h \cdot v = q^{\lambda(h)} \cdot v$$

In a similar way one define the simple  $\mathcal{U}'_q(\hat{\mathfrak{g}})$ -module  $L(\lambda, \Psi(u))$  of *l, u*-highest weight  $(\lambda, \Psi(u))$  (it is non trivial thanks to theorem 3.2).

**Lemma 6.9.** For  $(\lambda, \Psi(u)) \in P_{l,u}$  we have an isomorphism of  $\mathcal{U}'_q(\hat{\mathfrak{h}})$ -modules  $\tilde{L}(\lambda, \Psi(u)) \simeq L(\lambda, \Psi(u))$ .

*Proof:* Let  $\tilde{M}'(\lambda, \Psi(u)) \subset \tilde{M}(\lambda, \Psi(u))$  be the maximal proper  $\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})$ -submodule of  $\tilde{M}(\lambda, \Psi(u))$ . It suffices to prove that  $\tilde{\tau}_{-1}$  is included in  $\tilde{M}'(\lambda, \Psi(u))$  (see section 3.3.4; it implies that  $\tilde{L}(\lambda, \Psi(u))$  is also a  $\mathcal{U}'_q(\hat{\mathfrak{g}})$ -modules). It is a consequence of lemma 3.11.  $\square$

In particular  $\tilde{L}(\lambda, \Psi(u)) \in \mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}})) \Leftrightarrow L(\lambda, \Psi(u)) \in \mathcal{O}(\mathcal{U}'_q(\hat{\mathfrak{g}}))$  and in this case  $\text{ch}_{q,u}(\tilde{L}(\lambda, \Psi(u))) = \text{ch}_{q,u}(L(\lambda, \Psi(u)))$ .

#### 6.4.2. The category $\mathcal{O}_{\text{int}}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ .

**Definition 6.10.**  $QP_{l,u}^+$  is the set of  $(\lambda, \Psi(u)) \in P_{l,u}$  satisfying the following conditions :

i) for  $i \in I$  there exist polynomials  $Q_{i,u}(z) = (1 - za_{i,1}u^{b_{i,1}}) \dots (1 - za_{i,N_i}u^{b_{i,N_i}})$ ,  $R_{i,u}(z) = (1 - zc_{i,1}u^{d_{i,1}}) \dots (1 - zc_{i,N'_i}u^{d_{i,N'_i}})$  ( $a_{i,j}, c_{i,j} \in \mathbb{C}^*$ ,  $b_{i,j}, d_{i,j} \geq 0$ ) such that in  $\mathbb{C}[u, u^{-1}][[z]]$  (resp. in  $\mathbb{C}[u, u^{-1}][[z^{-1}]]$ ):

$$\sum_{r \geq 0} \Psi_{i,\pm r}^\pm(u) z^{\pm r} = q_i^{\deg(Q_{i,u}) - \deg(R_{i,u})} \frac{Q_{i,u}(zq_i^{-1})R_{i,u}(zq_i)}{Q_{i,u}(zq_i)R_{i,u}(zq_i^{-1})}$$

ii) there exist  $\omega \in P^+$ ,  $\alpha \in Q^+$  satisfying  $\lambda = \omega - \alpha$ .

$P_{l,u}^+$  is the set of  $(\lambda, \Psi(u)) \in QP_{l,u}^+$  such that one can choose  $R_{i,u} = 1$  (in this case we denote  $P_{i,u} = Q_{i,u}$ ).

**Lemma 6.11.** If  $(\lambda, \Psi(u)) \in P_{l,u}^+$  then  $\tilde{L}(\lambda, \Psi(u)) \in \mathcal{O}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$ . Moreover for  $(\mu, \gamma(u)) \in P_{l,u}$  we have  $\dim(\tilde{L}(\lambda, \Psi(u))_{\mu, \gamma(u)}) \neq 0 \Rightarrow (\mu, \gamma(u)) \in QP_{l,u}^+$ .

Remark : it follows from lemma 6.9 that we have the same results for  $L(\lambda, \Psi(u)) \in \mathcal{O}(\mathcal{U}'_q(\hat{\mathfrak{g}}))$ .

*Proof:* Let  $(\lambda, \Psi(u)) \in P_{l,u}^+$  and decompose  $P_{i,u}(z)$  in the form :

$$P_{i,u}(z) = P_i^{(0)}(z)P_i^{(1)}(uz) \dots P_i^{(R)}(u^R z)$$

where  $R \geq 0$ ,  $P_i^{(k)}(z) \in \mathbb{C}[z]$ ,  $P_i^{(k)}(0) = 1$  for  $0 \leq k \leq R$  ( $R$  can be taken large enough so that we have this form for all  $i \in I$ ). For  $0 \leq k \leq R$ , set  $\Psi_i^{(k)}(z) = q_i^{\deg(P_i^{(k)})} \frac{P_i^{(k)}(zq_i^{-1})}{P_i(zq_i)}$ . For  $1 \leq k \leq R$  define

$\lambda_k = \sum_{i \in I} \deg(P_i^{(k)}) \Lambda_i \in \mathfrak{h}^*$ . Set  $\lambda_0 = \lambda - \sum_{k=1 \dots R} \lambda_k$ . Then for  $0 \leq k \leq R$  the  $(\lambda_k, \Psi^{(k)}) \in P_l^+$  and we can consider  $L(\lambda_k, \Psi^{(k)}) \in \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . Let  $V \in \mathcal{O}^R(\tilde{\mathcal{U}}_q'(\hat{\mathfrak{g}}))$  be defined by :

$$V = i(L(\lambda_0, \Psi^{(0)})) \otimes_{R-1} (i(L(\lambda_1, \Psi^{(1)})) \otimes_{R-2} \dots (i(L(\lambda_{R-1}, \Psi^{(R-1)})) \otimes_0 i(L(\lambda_R, \Psi^{(R)}))) \dots)$$

Consider the  $\tilde{\mathcal{U}}_q'(\hat{\mathfrak{g}})$  submodule  $L$  of  $V$  generated by the tensor product of the highest weight vectors. It is a highest weight module of highest weight  $(\lambda, \Psi(u))$ . So  $\tilde{L}(\lambda, \Psi(u))$  is a quotient of  $L$  and so is in  $\mathcal{O}(\tilde{\mathcal{U}}_q'(\hat{\mathfrak{g}}))$ .

For the second point it follows from proposition 5.4 that the  $l, u$ -weight of  $i(L(\lambda_k, \Psi^{(k)}))$  are in  $QP_{l,u}^+$ . So with the help of the last point of theorem 6.7 we see that the  $l, u$ -weights of  $V$  are in  $QP_{l,u}^+$  and we have the property for  $\tilde{L}(\lambda, \Psi(u))$ .  $\square$

**Definition 6.12.** Let  $\mathcal{O}_{int}(\tilde{\mathcal{U}}_q'(\hat{\mathfrak{g}}))$  be the subcategory of modules  $V \in \mathcal{O}(\tilde{\mathcal{U}}_q'(\hat{\mathfrak{g}}))$  whose  $l, u$ -weights are in  $QP_{l,u}^+$ .

**Lemma 6.13.** For a module  $V \in \mathcal{O}_{int}(\tilde{\mathcal{U}}_q'(\hat{\mathfrak{g}}))$  there are  $P_{(\lambda, \Psi(u))} \geq 0$   $((\lambda, \Psi(u)) \in P_{l,u}^+)$  such that:

$$ch_{q,u}(V) = \sum_{(\lambda, \Psi(u)) \in P_{l,u}^+} P_{(\lambda, \Psi(u))} ch_{q,u}(\tilde{L}(\lambda, \Psi(u))) = \sum_{(\lambda, \Psi(u)) \in P_{l,u}^+} P_{(\lambda, \Psi(u))} ch_{q,u}(L(\lambda, \Psi(u)))$$

*Proof:* Analogous to the proof of proposition 5.2 (the second identity follows from lemma 6.9).  $\square$

## 6.5. $\mathbb{C}[u^\pm]$ -forms and specialization.

6.5.1.  $\mathbb{C}[u^\pm]$ -forms. Let  $\mathcal{U}_q^u(\hat{\mathfrak{g}}) = \mathcal{U}_q(\hat{\mathfrak{g}}) \otimes_{\mathbb{C}} \mathbb{C}[u^\pm] \subset \mathcal{U}_q'(\hat{\mathfrak{g}})$ .

**Definition 6.14.** A  $\mathbb{C}[u^\pm]$ -form of a  $\mathcal{U}_q'(\hat{\mathfrak{g}})$ -module  $V$  is a sub- $\mathcal{U}_q^u(\hat{\mathfrak{g}})$ -module  $L$  of  $V$  such that the map  $\mathcal{C} \otimes_{\mathbb{C}[u^\pm]} L \rightarrow V$  is an isomorphism of  $\mathcal{U}_q'(\hat{\mathfrak{g}})$ -module.

Note that it means that  $L$  generates  $V$  as  $\mathcal{C}$ -vector space and that some vectors which are  $\mathbb{C}[u^\pm]$ -linearly independent in  $L$  are  $\mathcal{C}$ -linearly independent in  $V$ .

Let us look at some examples:

**Proposition 6.15.** For  $(\lambda, \Psi(u)) \in P_{l,u}$  and  $v$  a highest weight vector of the Verma module  $M(\lambda, \Psi(u))$  (resp. the simple module  $L(\lambda, \Psi(u))$ ), the  $\mathcal{U}_q^u(\hat{\mathfrak{g}})$ -module  $\mathcal{U}_q^u(\hat{\mathfrak{g}}).v$  is a  $\mathbb{C}[u^\pm]$ -form of  $M(\lambda, \Psi(u))$  (resp. of  $L(\lambda, \Psi(u))$ ) which is isomorphic to the Verma (resp. the simple)  $\mathcal{U}_q^u(\hat{\mathfrak{g}})$ -module of  $l, u$ -highest weight  $(\lambda, \Psi(u))$ .

*Proof:* As  $(\lambda, \Psi(u))$  is fixed, we omit it.  $M$  is the quotient of  $\mathcal{C} \otimes_{\mathbb{C}} \mathcal{U}_q(\hat{\mathfrak{g}})$  by the relations generated by  $x_{i,r}^\pm = \phi_{i,\pm m}^\pm - \Psi_{i,\pm m}^\pm(u) = k_h - q^{\lambda(h)} = 0$ . So the relations between monomials are in  $\mathbb{C}[u^\pm]$  and  $\mathcal{U}_q^u(\hat{\mathfrak{g}}).1 \subset M$  is a  $\mathbb{C}[u^\pm]$ -form of  $M$ . Moreover those relations are the same as in the construction of the Verma  $\mathcal{U}_q^u(\hat{\mathfrak{g}})$ -module  $M^u$  as a quotient of  $\mathbb{C}[u^\pm] \otimes_{\mathbb{C}} \mathcal{U}_q(\hat{\mathfrak{g}})$ ; and so  $\mathcal{U}_q^u(\hat{\mathfrak{g}}).1 \simeq M^u$ .

Let us look at  $L$ . Denote by  $L^u$  the simple  $\mathcal{U}_q^u(\hat{\mathfrak{g}})$ -module of highest weight  $(\lambda, \Psi(u))$ . We have  $L = M/M'$  (resp.  $L^u = M^u/M'^u$ ) where  $M'$  (resp.  $M'^u$ ) in the maximal proper submodule of  $M$  (resp.  $M^u$ ).

The  $\mathcal{C}$ -subspace  $M''$  of  $M$  generated by  $M'^u$  is isomorphic to  $\mathcal{C} \otimes_{\mathbb{C}[u^\pm]} M'^u$  (because  $M^u$  is a  $\mathbb{C}[u^\pm]$ -form of  $M$ ). As  $M''$  has no vector of weight  $\lambda$ , it is a proper submodule of  $M$  and  $M'' \subset M'$ . Suppose that  $M' \neq M''$  and consider  $M'/M'' \subset M/M''$ .  $M^u/M'^u$  is a  $\mathbb{C}[u^\pm]$ -form of  $M/M''$ . Let  $v$  be a non zero highest weight vector of  $M'/M''$  and let us write:  $v = \sum_{\alpha} f_{\alpha}(u)v_{\alpha}$  where  $v_{\alpha} \in M^u/M'^u$  and  $f_{\alpha}(u) \in \mathcal{C}$

(as there is a finite number of  $f_{\alpha}(u)$ , we can suppose that they are  $\mathbb{C}[u^\pm]$ -linearly independent). For all  $i \in I, r \in \mathbb{Z}$ , we have  $x_{i,r}^+ . v = 0$  and so for all  $\alpha$ ,  $x_{i,r}^+ . v_{\alpha} = 0$ . Fix  $w_{\alpha} \in M^u$  whose image in  $M^u/M'^u$  is  $v_{\alpha}$ . As for all  $i \in I, r \in \mathbb{Z}$ ,  $x_{i,r}^+ . w_{\alpha} \in M'^u$ ,  $\mathcal{U}_q^u(\hat{\mathfrak{g}}).w_{\alpha}$  is a proper submodule of  $M^u$  and  $w_{\alpha} \in M'^u$ . So  $v = 0$ , contradiction. So  $M' = M''$ . In particular  $M' \simeq M'^u \otimes_{\mathbb{C}[u^\pm]} \mathcal{C}$ ,  $M' \cap M^u = M'^u$ .

For  $v$  a highest weight vector of  $L$ , the  $\mathcal{U}_q^u(\hat{\mathfrak{g}}).v \simeq \mathcal{U}_q^u(\hat{\mathfrak{g}}).1 = M^u/(M^u \cap M') = M^u/M'^u = L^u$  is a  $\mathbb{C}[u^\pm]$ -form of  $L$ .  $\square$

6.5.2. *Specializations.* Consider  $p : \mathcal{E}_{l,u} \rightarrow \mathcal{E}_l$  the surjection such that  $p((\lambda, \Psi(u))) = (\lambda, \Psi(1))$ .

**Lemma 6.16.** *Let  $V$  be in  $\mathcal{O}(\mathcal{U}'_q(\hat{\mathfrak{g}}))$ . If  $L$  is a  $\mathbb{C}[u^\pm]$ -form of  $V$  then the specialization  $L' = L/(1-u)L$  of  $L$  is in  $\mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$  and  $\text{ch}_q(L') = p(\text{ch}_{q,u}(V))$ .*

*Proof:* Indeed for  $(\mu, \gamma(u)) \in QP_{l,u}$  consider  $L_{\mu, \gamma(u)} = L \cap V_{\mu, \gamma(u)}$ . As  $p : L \otimes_{\mathbb{C}[u]} \mathbb{C} \rightarrow V$  is an isomorphism, we have  $V_{\mu, \gamma(u)} \simeq p^{-1}(V_{\mu, \gamma(u)}) = L_{\mu, \gamma(u)} \otimes_{\mathbb{C}[u]} \mathbb{C}$ . In particular  $L_{\mu, \gamma(u)}$  is a free  $\mathbb{C}[u^\pm]$  of rank  $\dim_{\mathbb{C}}(V_{\mu, \gamma(u)})$ . So  $\dim_{\mathbb{C}}(L'_\mu) = \dim_{\mathbb{C}}(V_\mu)$ , and  $L' \in \mathcal{O}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . We can conclude because:

$$L'_{\lambda, \gamma} = \bigoplus_{(\lambda, \gamma(u)) \in p^{-1}((\lambda, \gamma))} (L_{\lambda, \gamma(u)} / (u-1)L_{\lambda, \gamma(u)})$$

$\square$

6.5.3. *Proof of theorem 6.2.* For  $(\lambda, \Psi(u)) \in P_{l,u}^+$ , it follows from proposition 6.15 and lemma 6.16 that  $p(\text{ch}_{q,u}(L(\lambda, \Psi(u))))$  is of the form  $\text{ch}_q(L)$  where  $L \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ , that is to say  $p(\text{ch}_{q,u}(L(\lambda, \Psi(u)))) \in \text{ch}_q(\text{Rep}^+(\mathcal{U}_q(\hat{\mathfrak{g}})))$ . So (lemma 6.13) for  $V \in \mathcal{O}_{\text{int}}(\tilde{\mathcal{U}}'_q(\hat{\mathfrak{g}}))$  we have  $p(\text{ch}_{q,u}(V)) \in \text{ch}_q(\text{Rep}^+(\mathcal{U}_q(\hat{\mathfrak{g}})))$ .

Consider  $V_1, V_2 \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . We have seen that  $p(\text{ch}_{q,u}(i(V_1) \otimes_0 i(V_2))) \in \text{ch}_q(\text{Rep}^+(\mathcal{U}_q(\hat{\mathfrak{g}})))$ . But :

$$p(\text{ch}_{q,u}(i(V_1) \otimes_0 i(V_2))) = \text{ch}_q(V_1)\text{ch}_q(V_2)$$

because the specialization of  $\Delta_u$  on  $\mathcal{U}_q(\hat{\mathfrak{h}})$  at  $u = 1$  is  $\Delta_{\hat{\mathfrak{h}}}$ . This ends the proof of theorem 6.2.  $\square$

6.6. **Example.** We study in detail an example in the case  $\mathfrak{g} = \mathfrak{sl}_2$  where everything is computable thanks to Jimbo's evaluation morphism (see [CP3, CP4]). In this case we have  $\mathcal{U}_q(\hat{\mathfrak{sl}}_2) = \tilde{\mathcal{U}}_q(\hat{\mathfrak{sl}}_2)$ .

For  $a \in \mathbb{C}^*$  consider  $V = L(1-za) \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{sl}}_2))$ .  $V$  is two dimensional  $V = \mathbb{C}v_0 \oplus \mathbb{C}v_1$  and for  $r \in \mathbb{Z}$ ,  $m \geq 1$  the action of  $\mathcal{U}_q(\hat{\mathfrak{sl}}_2)$  is given in the following table:

	$v_0$	$v_1$
$x_r^+$	0	$a^r v_0$
$x_r^-$	$a^r v_1$	0
$\phi_{\pm m}^\pm$	$\pm(q - q^{-1})a^{\pm m}v_0$	$\mp(q - q^{-1})a^{\pm m}v_1$
$k^\pm$	$q^\pm v_0$	$q^\mp v_1$
$\phi^\pm(z)$	$q \frac{1-q^{-2}az}{1-az} v_0$	$q^{-1} \frac{1-q^2az}{1-az} v_1$

Remark : in the table  $\phi^\pm(z) \in \mathcal{U}_q(\hat{\mathfrak{g}})[[z^\pm]]$  acts on  $V[[z^\pm]]$ .

For  $a, b \in \mathbb{C}^*$ , let  $V = L(1-za), W = L(1-zb) \in \mathcal{O}_{\text{int}}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ . Consider basis  $V = \mathbb{C}v_0 \oplus \mathbb{C}v_1, W = \mathbb{C}w_0 \oplus \mathbb{C}w_1$  as in the previous table. The tensor product  $\otimes_0$  defines an action of  $\mathcal{U}'_q(\hat{\mathfrak{sl}}_2)$  on  $X = i(V) \otimes_{\mathbb{C}} i(W)$  (see theorem 6.7).  $X$  is a 4 dimensional  $\mathcal{C}$ -vector space of base  $\{v_0 \otimes w_0, v_1 \otimes w_0, v_0 \otimes w_1, v_1 \otimes w_1\}$ . The action of  $\mathcal{U}'_q(\hat{\mathfrak{sl}}_2)$  is given by ( $r \in \mathbb{Z}$ ):

	$v_0 \otimes w_0$	$v_1 \otimes w_0$
$x_r^+$	0	$a^r(v_0 \otimes w_0)$
$x_r^-$	$u^r b^r(v_0 \otimes w_1) + a^r q \frac{1-q^{-2}a^{-1}ub}{1-a^{-1}ub}(v_1 \otimes w_0)$	$u^r b^r(v_1 \otimes w_1)$
$\phi^\pm(z)$	$q^2 \frac{(1-q^{-2}az)(1-q^{-2}buz)}{(1-az)(1-buz)}(v_0 \otimes w_0)$	$\frac{(1-q^2az)(1-q^{-2}buz)}{(1-az)(1-buz)}(v_1 \otimes w_0)$
	$v_0 \otimes w_1$	$v_1 \otimes w_1$
$x_r^+$	$b^r u^r \frac{1-q^2a^{-1}bu}{1-uba^{-1}}(v_0 \otimes w_0)$	$a^r(v_0 \otimes w_1) + b^r u^r q \frac{1-q^{-2}a^{-1}bu}{1-uba^{-1}}(v_1 \otimes w_0)$
$x_r^-$	$a^r q^{-1} \frac{1-q^2ua^{-1}b}{1-a^{-1}ub}(v_1 \otimes w_1)$	0
$\phi^\pm(z)$	$\frac{(1-q^{-2}az)(1-q^2buz)}{(1-az)(1-buz)}(v_0 \otimes w_1)$	$q^{-2} \frac{(1-q^2az)(1-q^2buz)}{(1-az)(1-buz)}(v_1 \otimes w_1)$

Remark : in the table  $\phi^\pm(z) \in \mathcal{U}_q(\hat{\mathfrak{g}})[[z^\pm]]$  acts on  $X[[z^\pm]]$ .

Consider the  $l$ -weights  $\gamma_a, \gamma'_a, \gamma_b, \gamma'_b \in P_l$  (the  $\lambda \in \mathfrak{h}^*$  can be omitted because  $sl_2$  is finite) :

$$\gamma_a^\pm(z) = q \frac{1 - q^{-2}az}{1 - az}, \gamma'_a^\pm(z) = q^{-1} \frac{1 - q^2az}{1 - az}, \gamma_b^\pm(z) = q \frac{1 - q^{-2}bz}{1 - bz}, \gamma'_b^\pm(z) = q^{-1} \frac{1 - q^2bz}{1 - bz}$$

Consider also  $\gamma_a(z)\gamma_b(uz), \gamma'_a(z)\gamma_b(uz), \gamma_a(z)\gamma'_b(uz), \gamma'_a(z)\gamma'_b(uz) \in P_{l,u}$ . We see that :

$$\text{ch}_{q,u}(X) = e(\gamma_a(z)\gamma_b(uz)) + e(\gamma'_a(z)\gamma_b(uz)) + e(\gamma_a(z)\gamma'_b(uz)) + e(\gamma'_a(z)\gamma'_b(uz))$$

Those  $l, u$ -weights are distinct, the  $l, u$ -weights spaces are 1 dimensional:

$$X = (X)_{\gamma_a(z)\gamma_b(uz)} \oplus (X)_{\gamma'_a(z)\gamma_b(uz)} \oplus (X)_{\gamma_a(z)\gamma'_b(uz)} \oplus (X)_{\gamma'_a(z)\gamma'_b(uz)}$$

We see that  $X$  is of highest weight  $\gamma_a(z)\gamma_b(uz) \in P_{l,u}$ . Let us prove that it is simple : indeed  $X$  has no proper submodule : if for all  $r \in \mathbb{Z}$ ,  $x_r^+ \cdot (\alpha(v_1 \otimes w_0) + \beta(v_0 \otimes w_1)) = 0$ , then for all  $r \in \mathbb{Z}$ ,  $\alpha a^r + \beta b^r u^r \frac{1 - q^2 a^{-1} bu}{1 - uba^{-1}} = 0$ . In particular  $\alpha + \beta \frac{1 - q^2 a^{-1} bu}{1 - uba^{-1}} = 0$  and  $a^r - b^r u^r = 0$  for all  $r \in \mathbb{Z}$ , impossible. So  $X \simeq L(\gamma_a(z)\gamma_b(uz))$  as a  $\mathcal{U}'_q(\hat{sl}_2)$ -module. It follows from proposition 6.15 that  $\tilde{X} = \mathcal{U}_q^u(\hat{\mathfrak{g}}) \cdot (v_0 \otimes w_0) \subset X$  is a  $\mathbb{C}[u^\pm]$ -form of  $X$ .

Let us look explicitly at this  $\mathbb{C}[u^\pm]$ -form : consider  $e_1, e_2, e_3, e_4 \in \tilde{X}$  defined by:

$$e_1 = v_0 \otimes w_0, e_2 = x_0^- \cdot e_1, e_3 = -a^{-1} x_1^- \cdot e_1 + e_2, e_4 = q x_0^- \cdot e_2$$

We have the following formulas:

$$e_1 = v_0 \otimes w_0, e_2 = (v_0 \otimes w_1) + q \frac{1 - q^{-2} a^{-1} bu}{1 - a^{-1} ub} (v_1 \otimes w_0), e_3 = (1 - uba^{-1})(v_0 \otimes w_1), e_4 = (v_1 \otimes w_1)$$

Moreover the action of  $\mathcal{U}_q^u(\hat{\mathfrak{g}})$  is given by ( $r \in \mathbb{Z}$ ):

	$e_1$	$e_2$
$x_r^+$	0	$(q a^r \frac{1 - (a^{-1} ub)^{r+1}}{1 - a^{-1} ub} - q^{-1} b u a^{r-1} \frac{1 - (a^{-1} ub)^{r-1}}{1 - a^{-1} ub}) e_1$
$x_r^-$	$a^r (e_2 - \frac{1 - (a^{-1} ub)^r}{1 - a^{-1} ub} e_3)$	$(q^{-1} a^r \frac{1 - (a^{-1} ub)^{r+1}}{1 - a^{-1} ub} - q b u a^{r-1} \frac{1 - (a^{-1} ub)^{r-1}}{1 - a^{-1} ub}) e_4$
$\phi^\pm(z)$	$q^2 \frac{(1 - q^{-2} az)(1 - q^{-2} buz)}{(1 - az)(1 - buz)} e_1$	$\frac{(1 - q^2 az)(1 - q^2 buz)}{(1 - az)(1 - buz)} e_2 + \frac{az(q^2 - q^{-2})}{(1 - az)(1 - buz)} e_3$
	$e_3$	$e_4$
$x_r^+$	$b^r u^r q^{-1} (1 - q^2 a^{-1} bu) e_1$	$b^r u^r e_2 + a^r \frac{1 - (uba^{-1})^r}{1 - uba^{-1}} e_3$
$x_r^-$	$a^r q^{-1} (1 - q^2 u a^{-1} b) e_4$	0
$\phi^\pm(z)$	$\frac{(1 - q^{-2} az)(1 - q^2 buz)}{(1 - az)(1 - buz)} e_3$	$q^{-2} \frac{(1 - q^2 az)(1 - q^2 buz)}{(1 - az)(1 - buz)} e_4$

In particular we see that  $\mathbb{C}[u^\pm]e_1 \oplus \mathbb{C}[u^\pm]e_2 \oplus \mathbb{C}[u^\pm]e_3 \oplus \mathbb{C}[u^\pm]e_4$  is stable by the action of  $\mathcal{U}_q^u(\hat{\mathfrak{g}})$ , so is equal to  $\tilde{X}$ . So we have verified that  $X \simeq \tilde{X} \otimes_{\mathbb{C}[u^\pm]} \mathcal{C}$ .

Let us describe the specialization of  $\tilde{X}$  at  $u = 1$  : let  $\tilde{X}' = \mathbb{C}e_1 \oplus \mathbb{C}e_2 \oplus \mathbb{C}e_3 \oplus \mathbb{C}e_4$ . The action of  $\mathcal{U}_q(\hat{\mathfrak{g}})$  on  $\tilde{X}'$  is given by (for  $z \in \mathbb{C}, r \in \mathbb{Z}$ , we denote  $[z]'_r = \frac{1 - z^r}{1 - z} \in \mathbb{Z}[z^\pm]$  ( $z \neq 1$ ) and  $[1]'_r = r$ ):

	$e_1$	$e_2$
$x_r^+$	0	$(q a^r [a^{-1} b]'_r - q^{-1} b a^{r-1} [a^{-1} b]'_r) e_1$
$x_r^-$	$a^r (e_2 - [a^{-1} b]'_r e_3)$	$(q^{-1} a^r [a^{-1} b]'_r - q b a^{r-1} [a^{-1} b]'_r) e_4$
$\phi^\pm(z)$	$q^2 \frac{(1 - q^{-2} az)(1 - q^{-2} bz)}{(1 - az)(1 - bz)} e_1$	$\frac{(1 - q^2 az)(1 - q^2 bz)}{(1 - az)(1 - bz)} e_2 + \frac{az(q^2 - q^{-2})}{(1 - az)(1 - bz)} e_3$
	$e_3$	$e_4$
$x_r^+$	$b^r q^{-1} (1 - q^2 a^{-1} b) e_1$	$b^r e_2 + a^r [a^{-1} b]'_r e_3$
$x_r^-$	$a^r q^{-1} (1 - q^2 a^{-1} b) e_4$	0
$\phi^\pm(z)$	$\frac{(1 - q^{-2} az)(1 - q^2 bz)}{(1 - az)(1 - bz)} e_3$	$q^{-2} \frac{(1 - q^2 az)(1 - q^2 bz)}{(1 - az)(1 - bz)} e_4$

We see that  $\tilde{X}' = \mathcal{U}_q(\hat{\mathfrak{g}}).e_1$ . Moreover if  $ab^{-1} \notin \{q^2, q^{-2}\}$  :  $\tilde{X}'$  has no proper submodule because the formula  $x_m^+(ae_2 + \beta e_3) = 0$  means that for all  $r \in \mathbb{Z}$ :

$$\alpha(qa^r[a^{-1}b]_{r+1} - q^{-1}ba^{r-1}[a^{-1}b]_r) + \beta b^r q^{-1}(1 - q^2 a^{-1}b) = 0$$

which is possible only if  $ab^{-1} \in \{q^2, q^{-2}\}$  or  $\alpha = \beta = 0$ . So:

if  $ab^{-1} \notin \{q^2, q^{-2}\}$ ,  $\tilde{X}' \simeq L(\gamma_a \gamma_b)$  is simple and :

$$\text{ch}_q(V)\text{ch}_q(W) = \text{ch}_q(\tilde{X}') = \text{ch}_q(L(\gamma_a \gamma_b))$$

if  $ab^{-1} = q^2$  (resp.  $ab^{-1} = q^{-2}$ ),  $\mathbb{C}e_3 \subset \tilde{X}'$  (resp.  $\mathbb{C}((q^2 - 1)e_2 + e_3) \subset \tilde{X}'$ ) is a submodule of  $\tilde{X}'$  isomorphic to  $L(1)$  and :

$$\text{ch}_q(V)\text{ch}_q(W) = \text{ch}_q(\tilde{X}') = \text{ch}_q(L(\gamma_a \gamma_b)) + \text{ch}_q(L(1))$$

## REFERENCES

- [AK] **T. Akasaka and M. Kashiwara**, *Finite-dimensional representations of quantum affine algebras*  
Publ. Res. Inst. Math. Sci. 33, no. 5, 839–867 (1997)
- [Be] **J. Beck**, *Braid group action and quantum affine algebras*  
Comm. Math. Phys. 165, no. 3, 555–568 (1994)
- [Bo] **N. Bourbaki**, *Groupes et algèbres de Lie*  
Chapitres IV-VI, Hermann (1968)
- [CP1] **V. Chari and A. Pressley**, *Quantum Affine Algebras*  
Comm. Math. Phys. 142, 261–283 (1991)
- [CP2] **V. Chari and A. Pressley**, *Minimal affinizations of representations of quantum groups: the simply-laced case*  
Lett. Math. Phys. 35, no. 2, 99–114 (1995)
- [CP3] **V. Chari and A. Pressley**, *Quantum affine algebras and their representations*  
in Representations of groups (Banff, AB, 1994), 59–78, CMS Conf. Proc, 16, Amer. Math. Soc., Providence, RI (1995)
- [CP4] **V. Chari and A. Pressley**, *A Guide to Quantum Groups*  
Cambridge University Press, Cambridge (1994)
- [Da] **I. Damiani**, *La R-matrice pour les algèbres quantiques de type affine non tordu*.  
Ann. Sci. École Norm. Sup. (4) 31, no. 4, 493–523 (1998)
- [Dr1] **V. G. Drinfel'd**, *Quantum groups*  
Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 798–820, Amer. Math. Soc., Providence, RI, (1987)
- [Dr2] **V. G. Drinfel'd**, *A new realization of Yangians and of quantum affine algebras*  
Soviet Math. Dokl. 36, no. 2, 212–216 (1988)
- [DF] **J. Ding and I. B. Frenkel**, *Isomorphism of two realizations of quantum affine algebra  $U_q(\mathfrak{gl}(n))$*   
Comm. Math. Phys. 156, no. 2, 277–300 (1993)
- [DI] **J. Ding and K. Iohara**, *Generalization of Drinfeld quantum affine algebras*  
Lett. Math. Phys. 41, no. 2, 181–193 (1997)
- [DJ] **J. Ding and N. Jing**, *On a combinatorial identity*  
Internat. Math. Res. Notices, no. 6, 325–332 (2000)
- [E] **B. Enriquez**, *On correlation functions of Drinfeld currents and shuffle algebras*  
Transform. Groups 5, no. 2, 111–120 (2000)
- [EM] **P. I. Etingof and A. A. Moura**, *Elliptic Central Characters and Blocks of Finite Dimensional Representations of Quantum Affine Algebras*  
Represent. Theory 7, 346–373 (2003)
- [F] **J. Fuchs**, *Fusion rules in conformal field theory*  
Fortschr. Phys. 42, no. 1, 1–48 (1994)
- [FM] **E. Frenkel and E. Mukhin**, *Combinatorics of  $q$ -Characters of Finite-Dimensional Representations of Quantum Affine Algebras*  
Comm. Math. Phys., vol 216, no. 1, pp 23–57 (2001)
- [FR] **E. Frenkel and N. Reshetikhin**, *The  $q$ -Characters of Representations of Quantum Affine Algebras and Deformations of  $W$ -Algebras*  
Recent Developments in Quantum Affine Algebras and related topics, Cont. Math., vol. 248, pp 163–205 (1999)

- [Gr] **P. Grosse**, *On quantum shuffle and quantum affine algebras*  
Preprint: math.QA/0107176
- [Gu] **A. Guichardet**, *Groupes quantiques. Introduction au point de vue formel.*  
Savoirs Actuels. InterEditions, Paris; CNRS Éditions, Paris (1995)
- [GKV] **V. Ginzburg, M. Kapranov and E. Vasserot** *Langlands reciprocity for algebraic surfaces*  
Math. Res. Lett. 2, no. 2, 147–160 (1995)
- [H1] **D. Hernandez**, *t-analogues des opérateurs d'écrantage associés aux q-caractères*  
Int. Math. Res. Not., vol. 2003, no. 8, pp 451-475 (2003)
- [H2] **D. Hernandez**, *Algebraic Approach to q,t-Characters*  
Adv. Math. 187, no. 1, 1–52 (2004)
- [H3] **D. Hernandez**, *The t-analogs of q-characters at roots of unity for quantum affine algebras and beyond*  
J. Algebra 279, no. 2, 514–557 (2004)
- [Ja] **J. C. Jantzen**, *Lectures on quantum groups*  
Graduate Studies in Mathematics, 6. American Mathematical Society, Providence, RI, (1996)
- [Jim] **M. Jimbo**, *A q-difference analogue of  $U(\mathfrak{g})$  and the Yang-Baxter equation*  
Lett. Math. Phys. 10, no. 1, 63–69 (1985)
- [Jin] **N. Jing**, *Quantum Kac-Moody algebras and vertex representations*  
Lett. Math. Phys. 44, no. 4, 261–271 (1998)
- [Ka] **V. Kac**, *Infinite dimensional Lie algebras*  
3rd Edition, Cambridge University Press (1990)
- [Kn] **H. Knight**, *Spectra of tensor products of finite-dimensional representations of Yangians*  
J. Algebra 174, no. 1, 187–196 (1995)
- [KT] **S. M. Khoroshkin and V. N. Tolstoy**, *On Drinfeld's realization of quantum affine algebras*  
in Infinite-dimensional geometry in physics (Karpacz, 1992). J. Geom. Phys. 11 , no. 1-4, 445–452 (1993)
- [L] **G. Lusztig**, *Introduction to quantum groups*  
Progress in Mathematics, 110. Birkhäuser Boston, Inc., Boston, MA (1993)
- [LSS] **S. Levendorskii, Y. Soibelman and V. Stukopin**, *The quantum Weyl group and the universal quantum R-matrix for affine Lie algebra  $A_1^{(1)}$*   
Lett. Math. Phys. 27, no. 4, 253–264 (1993)
- [M1] **K. Miki**, *Representations of quantum toroidal algebra  $U_q(\mathfrak{sl}_{n+1, \text{tor}})(n \geq 2)$*   
J. Math. Phys. 41, no. 10, 7079–7098 (2000)
- [M2] **K. Miki**, *Quantum toroidal algebra  $U_q(\mathfrak{sl}_{2, \text{tor}})$  and R matrices*  
J. Math. Phys. 42, no. 5, 2293–2308 (2001)
- [N1] **H. Nakajima**, *Quiver varieties and finite-dimensional representations of quantum affine algebras*  
J. Amer. Math. Soc. 14, no. 1, 145–238 (2001)
- [N2] **H. Nakajima**, *Quiver Varieties and t-Analogs of q-Characters of Quantum Affine Algebras*  
To appear in Ann. Math. (Preprint arXiv:math.QA/0105173)
- [N3] **H. Nakajima**, *Geometric construction of representations of affine algebras*  
in “Proceedings of the International Congress of Mathematicians”, Volume I, 423–438 (2003)
- [R] **M. Rosso**, *Représentations des groupes quantiques*  
Séminaire Bourbaki exp. no. 744, Astérisque 201-203, 443-83, SMF (1992)
- [Sa] **Y. Saito**, *Quantum toroidal algebras and their vertex representations*  
Publ. Res. Inst. Math. Sci. 34, no. 2, 155–177 (1998)
- [Sc] **O. Schiffmann**, *Noncommutative projective curves and quantum loop algebras*  
Duke. Math. J. 121, no. 1, 113–168 (2004)
- [STU] **Y. Saito, K. Takemura and D. Uglov**, *Toroidal actions on level 1 modules of  $U_q(\widehat{\mathfrak{sl}}_n)$*   
Transform. Groups 3, no. 1, 75–102 (1998)
- [TU] **K. Takemura and D. Uglov**, *Representations of the quantum toroidal algebra on highest weight modules of the quantum affine algebra of type  $gl_N$*   
Publ. Res. Inst. Math. Sci. 35, no. 3, 407–450 (1999)
- [VV1] **M. Varagnolo and E. Vasserot**, *Double-loop algebras and the Fock space*  
Invent. Math. 133, no. 1, pp. 133–159 (1998)
- [VV2] **M. Varagnolo and E. Vasserot**, *Standard modules of quantum affine algebras*  
Duke Math. J. 111, no. 3, 509–533 (2002)



DAVID HERNANDEZ: ÉCOLE NORMALE SUPÉRIEURE - DMA, 45, RUE D'ULM F-75230 PARIS, CEDEX 05 FRANCE

*E-mail address:* David.Hernandez@ens.fr, URL: <http://www.dma.ens.fr/~dhernand>