DEFORMATIONS OF THE TAYLOR FORMULA

EMMANUEL FERRAND

ABSTRACT. Given a sequence $x = \{x_n,\ n \in \mathbb{N}\}$ with integer values, or more generally with values in a ring of polynomials with integer coefficients, one can form the *generalized binomial coefficients* associated to x, $\binom{n}{m}_x = \prod_{l=1}^{m} \frac{x_{n-l+1}}{x_l}$. In this note we introduce several sequences which enjoy the following remarkable feature : The fractions $\binom{n}{m}_x$ are in fact polynomials with integer coefficients.

1. INTRODUCTION

By a *deformation of the integers* we mean a sequence $x = \{x_n,\ n \in \mathbb{N}\}$ of polynomials in one or more variables and with integral coefficients, having the property that there exists some value q_0 of the variables such that $\forall n \in \mathbb{N}, x_n(q_0) = n$. The *quantum integers* $x_n = \sum_{l=0}^{n-1} q^l$ are a typical example of a deformation of the integers. Another example is given by the version of the Chebyshev polynomials defined by $x_n(cos(\theta)) = \frac{sin(n\theta)}{sin(\theta)}$.

In this note we consider some deformations of the factorial function and of the binomial coefficients which are induced by such deformations of the integers. This situation can be interpreted as a deformation of the Taylor formula, as explained below. Given a polynomial P of degree n with complex coefficients, the Taylor expansion at some point X gives

$$P(X+1) = P(X) + 1 \cdot \frac{dP}{dX}(X) + \frac{1^2}{2!} \cdot \frac{d^2P}{dX^2}(X) + ... + \frac{1^n}{n!} \cdot \frac{d^nP}{dX^n}(X).$$

In other words, if one denotes by $\tau : \mathbb{C}[X] \to \mathbb{C}[X]$ the "translation by one" operator, defined by $\tau(P)(X) = P(X+1)$, then $\tau = exp(\frac{d}{dX})$. A matrix version of this fact can be stated as follows. Denote by P and D the semi-infinite matrices whose coefficients are, respectively, $P_{i,j} = \binom{i}{j}$ and $D_{i,j} = i$ if $i = j+1$ and 0 otherwise, $(i,j) \in \mathbb{N}^2$. Then $P = exp(D)$.

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & \dots \\ 1 & 3 & 3 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \qquad D = \begin{pmatrix} 0 & 0 & 0 & 0 & \dots \\ 1 & 0 & 0 & 0 & \dots \\ 0 & 2 & 0 & 0 & \dots \\ 0 & 0 & 3 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

This suggests the following way to deform the Taylor formula. Replace the sequence $\mathbb{N}$ of the integers which appears as the non-zero coefficients of D by the terms of a sequence $x = \{x_n,\ n \in \mathbb{N}\}$ with values in some polynomial ring. Denote by D_x the corresponding

Date: July 2005.

1991 *Mathematics Subject Classification.* 05A10, 05A30, 11B39, 11B65.

Key words and phrases. Generalized binomial coefficients.

matrix. Given some integer n, define $n!_x$ to be the polynomial $n!_x = \prod_{l=1}^{n} x_l$. Define exp_x to be the formal series $exp_x(t) = \sum_{k=0}^{\infty} \frac{t^k}{k!_x}$. Observe that the matrix $exp_x(D_x)$ is well defined since, coefficients-wise, the summation is finite. Its coefficients $exp_x(D_x)_{i,j}$ will be denoted by the symbols $\binom{i}{j}_x$ and will be called *the generalized binomial coefficients associated with the sequence* x. Note that

$$\binom{i}{j}_x = \prod_{l=1}^{j} \frac{x_{i-l+1}}{x_l}$$

if $i \geq j$, and 0 otherwise.

This definition has appeared already in several contexts, see for example [KW] for an introduction to the relevant literature. Note that the fractions $\binom{i}{j}_x$ have no a priori reason to be polynomials with integer coefficients. In fact, such a phenomenon appears only for very specific sequences x.

In this note we are interested in deformations of the integers x which enjoy this property. The first part of the paper (section 2) is a variation on the classical theme of *quantum integers* and *q-binomials*. It deals with sequences that satisfy a second order linear recurrence relation. In the second part, (section 3), we deform the *integers* and the $q-binomials$ in a less standard way, using a sequence that satisfies a first order *non-linear* recurrence relation. In the third part (section 4), we introduce a sequence related to the Fermat numbers (which is not a deformation of the integers), and we show that the corresponding generalized binomial coefficients are polynomials with integer coefficients.

Let us mention that Knuth and Wilf [KW] showed that if a sequence x with integral values is a gcd-morphism (that is $x_{gcd(n,m)} = gcd(x_n, x_m)$), then the associated binomial coefficients are integers.

2. q-binomials

The properties of the so-called "quantum integers"

$$[n]_q = \sum_{l=0}^{n-1} q^l = \frac{1-q^n}{1-q}$$

and the associated "q-binomials" have been investigated long before the introduction of quantum mechanics (see [CK]). We rephrase below an approach developed by Carmichael [Car] (and probably already implicit in earlier works). It deals with a slightly more general, 2 variables version of the quantum integers.

Consider the sequence x with values in $\mathbb{Z}[a,b]$ defined by the following linear recurrence relation of order 2 :

$$x_0 = 0,\ x_1 = 1,\ x_{n+1} = a \cdot x_n + b \cdot x_{n-1}.$$

This sequence specializes to the quantum integers when $a = q+1$ and $b = -q$ (and to the usual integers for $a = 2$ and $b = -1$).

Remark. x_n is given by the following explicit formula :

$$x_n = \sum_{l=1}^{n} \binom{l-1}{n-l} a^{2l-n-1} b^{n-l}$$

as one can check by induction. □

Proposition (rephrased from [Car]).

- $x : \mathbb{N} \to \mathbb{Z}[a,b]$ is a gcd-morphism :

$$\gcd(x_n, x_m) = x_{\gcd(n,m)}.$$

- The associated binomial coefficients $\binom{n}{m}_x$ are polynomials in a and b with integral coefficients.

□

The first lines of the corresponding deformation of Pascal's triangle are as follows :

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & a & 1 & 0 & 0 & 0 \\ 1 & a^2+b & a^2+b & 1 & 0 & 0 \\ 1 & a^3+2ba & (a^2+2b)(a^2+b) & a^3+2ba & 1 & 0 \\ 1 & a^4+3ba^2+b^2 & (a^4+3ba^2+b^2)(a^2+2b) & (a^4+3ba^2+b^2)(a^2+2b) & a^4+3ba^2+b^2 & 1 \end{pmatrix}$$

Many classical sequences of integers or polynomials arise as solutions of second order recurrence relations with the appropriate initial conditions. The corresponding deformations of the Pascal triangle have often been considered separately in the literature. They receive a unified treatment through Carmichael's approach.

Example 1. For $a = b = 1$, the sequence x specializes to the *Fibonacci sequence*, and the triangle looks as follows :

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 2 & 1 & 0 & 0 & \dots \\ 1 & 3 & 6 & 3 & 1 & 0 & \dots \\ 1 & 5 & 15 & 15 & 5 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Example 2. For $a = 3$ and $b = -2$, the sequence x specializes to the *Mersenne numbers* $x_n = 2^n - 1$. The triangle then looks like

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 3 & 1 & 0 & 0 & 0 & \dots \\ 1 & 7 & 7 & 1 & 0 & 0 & \dots \\ 1 & 15 & 35 & 15 & 1 & 0 & \dots \\ 1 & 31 & 155 & 155 & 31 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Example 3. For $a = 2s$ and $b = -1$, the sequence $x_n = U_{n-1}(s)$, where U_n is the $n-$th *Chebyshev polynomial of the second kind*. This implies that, for any $(n, m) \in \mathbb{Z}^2$, the polynomial $\prod_{l=0}^{m} U_{n-l}$ is always divisible by $\prod_{l=0}^{m} U_l$ in $\mathbb{Z}[s]$.

3. Iterations of a polynomial

Fix some parameter $d \in \mathbb{N}$. Consider the polynomial

$$p(X, a_0, \ldots, a_d) = \sum_{k=0}^{d} a_k X^k$$

and the sequence x with values in $\mathbb{Z}[a_0, \ldots, a_d]$ defined by the following recurrence relation :

$$x_0 \ = 0, \ x_n = p(x_{n-1}, a_0, \ldots, a_d).$$

Note that this sequence is a deformation of the integers which encompasses the quantum integers (i.e., the case $d = 1, a_0 = 1, a_1 = q$, for which $x_n = [n]_q$).

Proposition.

- $x \ : \mathbb{N} \to \mathbb{Z}[a_0, \ldots, a_d]$ is a gcd-morphism : $x_{\gcd(n,m)} = \gcd(x_n, x_m)$.
- The associated binomial coefficients $\binom{n}{m}_x$ are polynomials of the variables $a_0, \ldots, a_d$, with integral coefficients.

Proof. Denote by ϕ_a the function $x \to p(x, a_0, \ldots, a_n)$ and by $\phi_a^{\circ n}$ its $n-$th iterate, so that $x_n = \phi_a^{\circ n}(0)$. For any $k \leq n$, $x_n = \phi_a^{\circ k}(x_{n-k})$. Writing $\phi_a^{\circ k}(x) = \phi_a^{\circ k}(0) + x \cdot Q(x)$ gives $x_n = \phi_a^{\circ k}(0) + x_{n-k} \cdot Q(x_{n-k})$. In other words, for any $k \leq n$, there exists a polynomial $R_{n,k}$ in $\mathbb{Z}[a_0, \ldots, a_n]$ such that

$$x_n = x_k + x_{n-k} \cdot R_{n,k}.$$

This implies that, for any $(k, l) \in \mathbb{Z}^2$, x_{kl} is divisible by x_k and by x_l. Furthermore this imply the following recurrence relation, from which the polynomiality of $\binom{n}{m}_x$ follows by induction :

$$\binom{n}{k}_x = x_n \cdot \frac{x_{n-1} \cdot \cdots \cdot x_{n-k+1}}{1 \cdot x_2 \cdot \cdots \cdot x_k} = \binom{n-1}{k-1}_x + R_{n,k} \cdot \binom{n-1}{k}_x.$$

Denote by d the gcd of n and k. We already know that x_d is a divisor of $\gcd(x_n, x_k)$. Write $d = \alpha \cdot n + \beta \cdot k$, with $\alpha \geq 0$ and $\beta \leq 0$, so that $x_{\alpha n} = x_d + x_{\beta k} \cdot R_{\alpha n, \beta k}$. Any common divisor of x_n and x_k is also a common divisor of $x_{\alpha n}$ and $x_{\beta k}$, and hence a divisor of x_d. This proves that $x_d = \gcd(x_n, x_k)$. $\square$

Even in the case $d = 2$, $\binom{n}{m}_x$ is a rather complicated polynomial. For example $\binom{5}{3}_x$ is of degree 11 in a_1 and of degree 21 in a_0 and a_2. If one specializes to the case $a_0 = a_1 = 1$ and $a_2 = q - 1$, the corresponding one-parameter deformation of Pascal's triangle (which is recovered at $q = 1$) looks like

$$\begin{pmatrix} 1 & 0 & 0 & \dots \\ 1 & 1 & 0 & \dots \\ 1 & 1+q & 1 & \dots \\ 1 & 1+q^2+q^3 & 1+q^2+q^3 & \dots \\ 1 & (1+q)(q^6-q^4+2q^3-q^2+1) & (1+q^2+q^3)(q^6-q^4+2q^3-q^2+1) & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Remark. Consider now a polynomial $p(X, a_1, \dots, a_d) = \sum_{k=1}^{d} a_k X^k$ whose constant term vanishes, and the sequence x with values in $\mathbb{Z}[a_0, \dots, a_d]$ defined by the following recurrence relation :

$$x_0 = a_0,\ x_n = p(x_{n-1}, a_1, \dots, a_d).$$

The corresponding $\binom{n}{m}_x$ are also polynomials in the variables $a_0, \dots, a_1$ with integral coefficients, for all $(n, m) \in \mathbb{Z}^2$. This is due to the fact that, if $n \geq m$, x_n is a multiple of x_m, which implies that $\binom{n}{m}_x$ is a multiple of $\binom{n-1}{m-1}_x$.

On the other hand, this sequence x is *not* a deformation of the integers, since $\forall n \geq m$, x_m divides x_n.

4. Fermat polynomials

The sequence of polynomials considered in this section is not a deformation of the integers, but is related to the Fermat numbers. It is defined explicitly by the formula

$$x_n = \sum_{l=0}^{n-1} (\binom{n-1}{l} \mod 2) \cdot X^l.$$

If $n > 0$, x_n is the unique element of $\mathbb{Z}[X]$ with coefficients in $\{0, 1\}$ which is congruent to $(1+X)^{n-1}$ modulo 2. The first few terms are $x_0 = 0$, $x_1 = 1$, $x_2 = 1+X$, $x_3 = 1+X^2$, $x_4 = 1 + X + X^2 + X^3$.

By a theorem of Lucas, (see, for example [GKP], ex. 61 p. 248), the parity of $\binom{n}{m}$ is determined by the binary decomposition of n and m as follows : Write $n = \sum_{l \in \mathbb{N}} \epsilon_l 2^l$ and $m = \sum_{l \in \mathbb{N}} \eta_l 2^l$, with $\epsilon_l, \eta_l \in \{0, 1\}, \forall l \in \mathbb{N}$. Then

$$\binom{n}{m} = \prod_{l \in \mathbb{N}} \binom{\epsilon_l}{\eta_l} \mod 2.$$

Since $\binom{\epsilon_l}{\eta_l} = 1 + \eta_l \cdot (\epsilon_l - 1)$, this can be rephrased in a compact way as follows. With an integer p, associate the set K_p of the exponents that appear in the binary decomposition of p, so that $n = \sum_{l \in K_n} 2^l$ and $m = \sum_{l \in K_m} 2^l$. Then $\binom{n}{m}$ is odd if and only if $K_m \subset K_n$.

For example, if $n - 1 = 2^k$ is a power of 2, $\binom{n-1}{l}$ is even for any $1 \leq l \leq 2^k - 1$. Hence $x_{2^k+1} = 1 + X^{2^k}$, and x_{2^k+1} specializes to the $k-$th *Fermat number* $1 + 2^{2^k}$ at $X = 2$. If $n = 2^k$ is a power of 2, $\binom{n-1}{l}$ is odd for any $0 \leq l \leq 2^k - 1$. Hence $x_{2^k} = \sum_0^{2^k-1} X^l = \frac{X^{2^k}-1}{X-1}$. In particular, for all $k \in \mathbb{N}, x_{2^k+1} = 2 + (X-1)x_{2^k}$.

Proposition.

- $x_{n+1} = \prod_{l \in K_n}(1 + X^{2^l})$, and x_m divides x_n in $\mathbb{Z}[X]$ if and only if $\binom{n-1}{m-1}$ is odd.
- The associated binomial coefficients $\binom{n}{m}_x$ are polynomials in X, with integral coefficients.

Proof. Observe that, for any $(l, m) \in \mathbb{N}^2$,

$$(1+X)^{2^l+m} = (1+X)^{2^l}(1+X)^m \equiv (1+X^{2^l})(1+X)^m \mod 2.$$

This imply that

$$(1+X)^{(\sum_{l \in K_n} 2^l)} \equiv \prod_{l \in K_n} (1+X^{2^l}) \mod 2.$$

On the other hand $\prod_{l \in K_n}(1+X^{2^l})$ is an element of $\mathbb{Z}[X]$ whose coefficients are in $\{0,1\}$. But x_{n+1} is by definition the unique element of $\mathbb{Z}[X]$ whose coefficients are in $\{0,1\}$ and which is congruent to $(1+X)^n$ modulo 2. Hence $x_{n+1} = \prod_{l \in K_n}(1+X^{2^l})$. From this factorization it follows that x_m divides x_n in $\mathbb{Z}[X]$ if and only if $K_{m-1} \subset K_{n-1}$. By Lucas's theorem, this last condition is equivalent to the oddity of $\binom{n-1}{m-1}$.

To prove that $\binom{n}{m}_x$ is a polynomial, we will study the exponent $\alpha_l(n,m)$ of each factor $(1+X^{2^l})$ in the decomposition $\binom{n}{m}_x = \prod_{l \in \mathbb{N}}(1+X^{2^l})^{\alpha_l(n,m)}$. Denote by $\epsilon_l : \mathbb{N} \to \{0,1\}$ the function such that $\epsilon_l(p) = 1$ iff $l \in K_p$, so that $p = \sum_{l \in \mathbb{N}} \epsilon_l(p) 2^l$. It follows that $\alpha_l(n,m) = \sum_{p=1}^{m} \epsilon_l(n-p+1-1) - \epsilon_l(p-1)$.

The function ϵ_l is periodic, of period 2^{l+1}. Hence, when estimating $\alpha_l(n,m)$, one can assume that m is smaller than 2^{l+1}. Observe that $\epsilon_l(p) = 0$ for $p \in \{0, \ldots, 2^l - 1\}$, and that $\epsilon_l(p) = 1$ for $p \in \{2^l, \ldots, 2^{l+1} - 1\}$. The sum $\sum_{p \in \{r, \ldots, r+m-1\}} \epsilon_l(p)$ over a "window" of width m is bounded from below by $max(0, m - 2^l)$. This minimal value is attained at $r = 0$. This proves that $\sum_{p=1}^{m} \epsilon_l(n-p) \geq \sum_{p=1}^{m} \epsilon_l(p-1)$, and hence that $\alpha_l(n,m) \geq 0$. □

Example. The first few lines of the corresponding triangle are as follows :

$$\begin{pmatrix}
1 & 0 & 0 & 0 & 0 & 0 & \ldots \\
1 & 1 & 0 & 0 & 0 & 0 & \ldots \\
1 & 1+X & 1 & 0 & 0 & 0 & \ldots \\
1 & 1+X^2 & 1+X^2 & 1 & 0 & 0 & \ldots \\
1 & 1+X+X^2+X^3 & (1+X^2)^2 & 1+X+X^2+X^3 & 1 & 0 & \ldots \\
1 & 1+X^4 & (1+X^2)(1+X^4) & (1+X^2)(1+X^4) & 1+X^4 & 1 & \ldots \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots
\end{pmatrix}.$$

We have seen that the specialization of x at $X = 2$ gives a sequence that interpolates in a natural way between the Fermat numbers. The specialization $1, 2, 2, 4, 2, \ldots$ at $X = 1$ is also meaningful: $x_n(1) = 2^{|K_{n-1}|}$, where $|K_{n-1}|$ denotes the number of non-vanshing terms in the binary expansion of $n-1$.

References

[Car] R.D. Carmichael, On the numerical factors of the arithmetics forms $\alpha^n \pm \beta^n$, *Ann Math* **15** (1913-1914), 30–70.

[CK] P. Cheung, V. Kac, *Quantum Calculus* Springer-Verlag New York (2002).

[GKP] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics*, second edition, Addison-Wesley, (1994).

[KW] D.E. Knuth, H.S. Wilf, The power of a prime that divides a generalized binomial coefficient *J. Reine Angew. Math* **396** (1989), 212–219.

Institut Fourier, BP 74, 38402 St Martin d'Hères Cedex, France.

E-mail address: `emmanuel.ferrand@ujf-grenoble.fr`