

A note on the nonlinear derived Cauchy problem

Frédéric Paugam

July 7, 2022

Abstract

We define and study a generalization of the analytic Cauchy problem, that specializes to the Cauchy-Kowaleskaya-Kashiwara problem in the linear case. The main leitmotive of this text is to adapt Kashiwara's formulation of this problem both to the relatively \mathcal{D} -algebraic case and to the derived analytic situation. Along the way, we define the characteristic variety of a derived nonlinear partial differential system.

Contents

1	Introduction	1
2	The relatively algebraic derived Cauchy problem	3
2.1	The classical Cauchy problem in terms of \mathcal{D} -algebras	3
2.2	Characteristic variety of a \mathcal{D} -algebra	4
2.3	The Cauchy problem in the smooth case	5
2.4	About the relatively algebraic non-characteristic derived Cauchy problem	5
3	The derived analytic Cauchy problem	6
3.1	Recollection about derived analytic geometry	6
3.2	The de Rham stack and geometry of derived nonlinear PDEs	8
3.3	The classical Cauchy problem and geometry of PDEs	9
3.4	Characteristic variety of a derived nonlinear PDE	9
3.5	About the non-characteristic derived Cauchy problem	10

1 Introduction

The classical real analytic Cauchy problem is the initial value problem (see e.g. Nirenberg's article [Nir72]):

$$\partial_t^m u = F(t, x, u, \partial_x^\alpha \partial_t^j u), \quad \partial_t^k u|_{t=0} = \varphi_k, \quad k = 0, \dots, m - 1.$$

Here x is in an open subset $\Omega \subset \mathbb{R}^n$, $t \in \mathbb{R}$ and u may be vector valued; f is a function that depends on t , x and u , and all of its derivatives of order smaller than m of the form $\partial_x^\alpha \partial_t^j u$, $|\alpha| + j \leq m$, $j < m$. If f and the φ_k 's are analytic then the problem has a unique analytic solution in a neighborhood of any initial point $(x_0, 0)$.

We will say that we are in the relatively algebraic situation of the Cauchy problem if the analytic function F is algebraic in the variable function u and its partial derivatives.

Kashiwara proposed in his memoir [Kas70] (see also [Sch13] for an exposition) a generalization of this problem adapted to the algebraic study of linear systems of partial differential equations using \mathcal{D} -modules. Let $f : X \rightarrow Y$ be a morphism of complex manifolds and \mathcal{M} and \mathcal{N} be two \mathcal{D} -modules on Y . Then one says that the Cauchy-Kowalevskaya-Kashiwara theorem is true for the triple $(f, \mathcal{M}, \mathcal{N})$ if the natural morphism

$$\nu : f^{-1} \mathbb{R} \mathcal{H}om_{\mathcal{D}}(\mathcal{M}, \mathcal{N}) \rightarrow \mathbb{R} \mathcal{H}om_{\mathcal{D}}(f_{\mathcal{D}}^* \mathcal{M}, f_{\mathcal{D}}^* \mathcal{N})$$

is an isomorphism. Kashiwara showed that this theorem is true if f is smooth and also if f is non-characteristic for a coherent \mathcal{M} and $\mathcal{N} = \mathcal{O}$.

Let us reformulate this result of Kashiwara in a nonlinear context. If $\mathcal{P}^\bullet \xrightarrow{\sim} \mathcal{M}$ is an acyclic resolution (given by a simplicial module), then we have that the natural maps between sheaves of spaces

$$\mathbb{R} \mathcal{M}ap_{s-\mathcal{D}-alg}(\mathbb{R} \text{Sym}(\mathcal{P}^\bullet), \mathcal{O}) \rightarrow \mathbb{R} \mathcal{M}ap_{\mathcal{D}}(\mathcal{M}, \mathcal{O})$$

and

$$\mathbb{R} \mathcal{M}ap_{s-\mathcal{D}-alg}(f_{\mathcal{D}}^* \mathbb{R} \text{Sym}(\mathcal{P}^\bullet), \mathcal{O}) \rightarrow \mathbb{R} \mathcal{M}ap_{\mathcal{D}}(f_{\mathcal{D}}^* \mathcal{M}, \mathcal{O})$$

are isomorphisms, so that having Kashiwara's isomorphism is equivalent, if we denote $\mathcal{A} := \mathbb{R} \text{Sym}(\mathcal{P}^\bullet) \cong \mathbb{R} \text{Sym}(\mathcal{M})$ and $\mathcal{B} = \mathcal{O}$, to having that the natural map

$$\nu : f^{-1} \mathbb{R} \mathcal{M}ap_{\mathcal{D}-alg}(\mathcal{A}, \mathcal{B}) \rightarrow \mathbb{R} \mathcal{M}ap_{\mathcal{D}-alg}(f_{\mathcal{D}}^* \mathcal{A}, f_{\mathcal{D}}^* \mathcal{B})$$

is an isomorphism of sheaves of spaces. This condition for arbitrary simplicial \mathcal{D} -algebras \mathcal{A} and \mathcal{B} will be a formulation of the nonlinear derived Cauchy problem.

Definition 1. Let $f : X \rightarrow Y$ be a morphism of complex manifolds and \mathcal{A} and \mathcal{B} be two simplicial \mathcal{D}_Y -algebras. One says that the Cauchy-Kowalevskaya-Kashiwara theorem is true for the triple $(f, \mathcal{A}, \mathcal{B})$ if the natural morphism

$$\nu : f^{-1} \mathbb{R} \mathcal{M}ap_{\mathcal{D}-alg}(\mathcal{A}, \mathcal{B}) \rightarrow \mathbb{R} \mathcal{M}ap_{\mathcal{D}-alg}(f_{\mathcal{D}}^* \mathcal{A}, f_{\mathcal{D}}^* \mathcal{B})$$

is an isomorphism of sheaves of spaces on Y .

One may ask, as a conjecture, if this result is true if $\mathcal{B} = \mathcal{O}$, and \mathcal{A} is non-characteristic for f in the sense of Subsection 2.2. We will give a proof of this conjecture in the relatively algebraic case of a smooth morphism $f : X \rightarrow Y$ (submersion).

The above derived formulation of the relatively algebraic nonlinear Cauchy problem will be precisely formulated and studied in Section 2.

The theory of \mathcal{D} -algebras is well adapted to the study of relatively algebraic nonlinear partial differential systems, because the associated jet algebras are relatively algebraic. We will give another formulation of the derived Cauchy problem adapted to general analytic nonlinear partial differential systems in Section 3, using a more geometric approach, based on the use of Simpson's de Rham spaces and of Porta's derived analytic geometry from [Por14] (see also [Por17] for a combination of those two tools).

This will lead to the following formulation of the derived Cauchy problem.

Definition 2. Let $f : X \rightarrow Y$ be a morphism of complex manifolds and Z and Z' be two derived analytic stacks over the de Rham space Y_{dR} . One says that the Cauchy-Kowalevskaya-Kashiwara theorem is true for the triple (f, Z, Z') if the natural morphism

$$\nu : f^{-1}\mathbb{R}\mathcal{M}ap_{Y_{dR}}(Z, Z') \rightarrow \mathbb{R}\mathcal{M}ap_{X_{dR}}(f_{dR}^*Z, f_{dR}^*Z')$$

is an isomorphism of sheaves of spaces on Y .

One may ask, as a conjecture, if this is true if $Z = Y_{dR}$ and Z' is non-characteristic for f in the sense of Subsection 3.4. We prove this conjecture in the particular case of a smooth submersion.

Remark: in all this note, we use derived functors without further notice.

Acknowledgments: The author thanks Pierre Schapira and Bertrand Toën for useful discussions.

2 The relatively algebraic derived Cauchy problem

2.1 The classical Cauchy problem in terms of \mathcal{D} -algebras

It is possible to relate the derived Cauchy problem to the relatively algebraic classical Cauchy problem

$$\partial_t^m u = F(t, x, u, \partial_x^\alpha \partial_t^i u), \quad \partial_t^k u|_{t=0} = \varphi_k, \quad k = 0, \dots, m - 1$$

in the complex analytic setting, where $u : Y \rightarrow E$ is a section of a (say, trivial) vector bundle of rank r . The \mathcal{D} -algebra in play is the quotient $\mathcal{A} = \text{Jet}(\mathcal{O}_E^{alg}) / (u_t^m - F)$ where $\text{Jet}(\mathcal{O}_E^{alg})$ is generated as an algebra over \mathcal{O}_Y by the coordinates of u and of its formal partial derivatives, the \mathcal{D}_Y -algebra structure being given by the usual action of partial derivatives on jet coordinates (by décalage). The sheaf of sets

$$\mathcal{H}om_{\mathcal{D}\text{-alg}}(\mathcal{A}, \mathcal{O}_Y) \subset \mathcal{H}om_{\mathcal{D}\text{-alg}}(\text{Jet}(\mathcal{O}_E^{alg}), \mathcal{O}_Y) \cong \text{Hom}_{\mathcal{O}_Y\text{-alg}}(\mathcal{O}_E^{alg}, \mathcal{O}_Y) \cong \mathcal{O}_Y^r$$

is the sheaf of solutions of the given nonlinear partial differential equation. The initial value condition corresponds to the situation where we study the pullback along the closed embedding

$$f : X = \{t = 0\} \hookrightarrow Y$$

for which

$$f_{\mathcal{D}}^*(\partial_x^\alpha u) = \partial_x^\alpha (f^* u)$$

and

$$f_{\mathcal{D}}^*(\partial_t^k u) = \partial_t^k u|_X.$$

The jet algebra $\text{Jet}(\mathcal{O}_E^{\text{alg}})$ is locally generated over $\mathcal{O}_E^{\text{alg}}$ by the coordinates $u_{t,x}^{k,\alpha}$ with standard action of $\mathcal{D}_Y = \{\sum a_{k,\alpha} \partial_t^k \partial_x^\alpha\}$ by décalage. Its pullback $f_{\mathcal{D}}^* \text{Jet}(\mathcal{O}_E^{\text{alg}})$ is generated over $f^* \mathcal{O}_E^{\text{alg}}$ by the same coordinates $u_{t,x}^{k,\alpha}$ with action of $\mathcal{D}_X = \{\sum a_\alpha \partial_x^\alpha\}$. Then the quotient \mathcal{D}_Y -algebra $\mathcal{A} = \text{Jet}(\mathcal{O}_E^{\text{alg}})/(u_t^m - F)$ is generated over $\mathcal{O}_E^{\text{alg}}$ by the coordinates $u_{t,x}^{k,\alpha}$ for $k \leq m-1$.

The natural map

$$\nu : f^{-1} \mathcal{H}om_{\mathcal{D}\text{-alg}}(\mathcal{A}, \mathcal{O}_Y) \rightarrow \mathcal{H}om_{\mathcal{D}\text{-alg}}(f_{\mathcal{D}}^* \mathcal{A}, \mathcal{O}_Y)$$

thus corresponds to taking the initial value of the solutions of the given partial differential equation, and it is an isomorphism by the classical Cauchy-Kowalevskaya theorem.

2.2 Characteristic variety of a \mathcal{D} -algebra

We propose here an adaptation of Bächtold's construction (see [Bae09]) of the characteristic variety of a (non-derived) diffiety to the derived \mathcal{D} -algebraic context.

Let \mathcal{A} be a simplicial \mathcal{D} -algebra on Y and $\mathbb{L}_{\mathcal{A}}$ be its cotangent complex. Suppose that $\mathbb{L}_{\mathcal{A}}$ is perfect (as an \mathcal{A} -module in the category of \mathcal{D} -modules, i.e., as an $\mathcal{A}[\mathcal{D}]$ -module). Let $\mathbb{T}_{\mathcal{A}}^\ell$ be the left $\mathcal{A}[\mathcal{D}]$ -module associated to its $\mathcal{A}[\mathcal{D}]$ -dual. We will call $\mathbb{T}_{\mathcal{A}}^\ell$ its tangent complex. We may locally write this complex as a complex of free $\mathcal{A}[\mathcal{D}]$ -modules of finite rank. Let $\pi : T^*Y \rightarrow Y$ be the natural projection and \mathcal{E}_Y be the sheaf of microdifferential operators on T^*Y . Define the support of a complex to be the union of the supports of its cohomology spaces. The support of the microlocalization

$$\mu(\mathbb{T}_{\mathcal{A}}^\ell) := \pi^* \mathbb{T}_{\mathcal{A}}^\ell \otimes_{\pi^* \mathcal{A}[\pi^* \mathcal{D}_Y]} (\pi^* \mathcal{A})[\mathcal{E}_Y]$$

of the tangent complex $\mathbb{T}_{\mathcal{A}}^\ell$ in

$$\text{Spec}_{T^*Y}(\pi^* \mathcal{A}) = \text{Spec}_Y(\mathcal{A}) \times_Y T^*Y$$

is called the characteristic variety of \mathcal{A} and denoted $\text{Char}(\mathcal{A})$. It may be defined as the derived space of zeroes of the $\mathcal{A} \otimes_{\mathcal{O}_Y} \mathcal{O}_{T^*Y}$ -ideal $\mathcal{I}_{\text{Char}(\mathcal{A})}$ that is the derived annihilator of the microlocalization module $\mu(\mathbb{T}_{\mathcal{A}}^\ell)$.

If $\varphi : \mathcal{A} \rightarrow \mathcal{O}_Y$ is a classical solution of \mathcal{A} (morphism of \mathcal{D} -algebras), we may pullback $\mathbb{T}_{\mathcal{A}}^\ell$ along $\varphi^* : Y \rightarrow \text{Spec}_Y(\mathcal{A})$ to get a \mathcal{D} -module on Y that is the linearization $\mathbb{T}_{\mathcal{A},\varphi}^\ell$ of the nonlinear system \mathcal{A} along its solution φ . Similarly, we may pull-back $\mu(\mathbb{T}_{\mathcal{A}}^\ell)$ to T^*Y along $\varphi^* : Y \rightarrow \text{Spec}_Y(\mathcal{A})$ to get the microlocalization $\mu(\mathbb{T}_{\mathcal{A},\varphi}^\ell)$ of the linearization, whose support gives the characteristic variety $\text{Char}_\varphi(\mathcal{A})$ of this linearization $\mathbb{T}_{\mathcal{A},\varphi}^\ell$.

We want to say that $f : X \rightarrow Y$ is non-characteristic for \mathcal{A} if, for every solution φ , $\text{Char}_\varphi(\mathcal{A})$ is non-characteristic for f in the sense of Kashiwara. This may be formulated directly in terms of the characteristic variety

$$\text{Char}(\mathcal{A}) \subset \text{Spec}_Y(\mathcal{A}) \times_Y T^*Y.$$

Indeed, we have a cartesian diagram

$$\begin{array}{ccccc} \text{Spec}_Y(\mathcal{A}) \times_Y T^*X & \xleftarrow{f_d} & \text{Spec}_Y(\mathcal{A}) \times_Y X \times_Y T^*Y & \xrightarrow{f_\pi} & \text{Spec}_Y(\mathcal{A}) \times_Y T^*Y \\ \downarrow & & \downarrow \pi & & \downarrow \\ X & \xleftarrow{id_X} & X & \xrightarrow{f} & Y \end{array}$$

and we may say f is non-characteristic for \mathcal{A} if

$$f_\pi^{-1}(\text{Char}(\mathcal{A})) \cap \text{Spec}_Y(\mathcal{A}) \times_Y T_X^*Y \subset \text{Spec}_Y(\mathcal{A}) \times_Y X \times_Y T_Y^*Y.$$

By pullback along a solution φ , we get the classical non-characteristic condition for the linearization of the given partial differential system $\mathbb{T}_{\mathcal{A},\varphi}^\ell$.

2.3 The Cauchy problem in the smooth case

Let $f : X \rightarrow Y$ be a smooth morphism (submersion). Since the Cauchy problem is a local question, we may assume that f is the natural projection $f : X = Y \times Z \rightarrow Y$ from a product space. The pullback functor $f_{\mathcal{D}}^*$ is exact.

Recall that coherent \mathcal{D} -modules have a perfect resolution. Their derived analogs are given by compact objects in the category of \mathcal{D} -modules, that are given by complexes with bounded coherent cohomology. A possible nonlinear analog of coherent \mathcal{D} -modules is given by \mathcal{D} -algebras that are homotopy finitely presented, i.e., compact objects in the homotopy category of dg- \mathcal{D} -algebras. These are given¹ by dg- \mathcal{D} -algebras that are retracts of cellular objects (see [TV07], Definition 2.1 and Proposition 2.2) whose building blocks are of the form $\text{Sym}_{\mathcal{O}}(\mathcal{E})$ for \mathcal{E} a compact \mathcal{D} -module.

In the case of such a homotopy finitely presented algebra \mathcal{A} , showing that the natural map

$$\nu : f^{-1}\mathbb{R}\text{Map}_{\mathcal{D}\text{-alg}}(\mathcal{A}, \mathcal{O}_X) \rightarrow \mathbb{R}\text{Map}_{\mathcal{D}\text{-alg}}(\mathcal{A}, \mathcal{O}_Y)$$

is an isomorphism reduces to showing that it is an isomorphism when \mathcal{A} is a free \mathcal{D} -algebra on a given coherent \mathcal{D} -module. The result then reduces to the linear case of a coherent \mathcal{D} -module, that reduces to the free case (by using a free resolution locally), that is a particular case of the relative de Rham theorem (see [Sch13], proof of Theorem 2.4.1).

¹This result was explained to the Author by Bertrand Toën.

2.4 About the relatively algebraic non-characteristic derived Cauchy problem

The argument for the non-characteristic Cauchy problem should be the same as the one used in the linear case (see [Sch13], Theorem 2.4.1): one writes the morphism as a composition of a closed (graph) immersion and a smooth (projection) morphism. The case of a closed immersion remains open because its treatment is more complicated than in the linear case.

The main point here is to be able to make the “homotopy dévissage” of the given non-linear system in linear ones (a homotopy finitely presented \mathcal{D} -algebra admits such a dévissage in terms of symmetric algebras of linear systems) compatible to the “homotopy dévissage” of the corresponding characteristic varieties.

We thus formulate it as a conjecture.

Conjecture 1 (Relatively algebraic non-characteristic derived Cauchy problem). *Let $f : Y \rightarrow X$ be a morphism of complex manifolds and suppose that \mathcal{A} is a derived \mathcal{D}_Y -algebra that is homotopy finitely presented and non-characteristic for f . Then the natural map*

$$\nu : f^{-1}\mathbb{R}\mathrm{Map}_{\mathcal{D}\text{-alg}}(\mathcal{A}, \mathcal{O}_X) \rightarrow \mathbb{R}\mathrm{Map}_{\mathcal{D}\text{-alg}}(f_{\mathcal{D}}^*\mathcal{A}, \mathcal{O}_Y)$$

is an isomorphism.

In the simplest case of the symmetric algebra $\mathcal{A} = \mathrm{Sym}(\mathcal{M})$ of a compact complex of \mathcal{D} -modules, the tangent complex $\mathbb{T}_{\mathcal{A}}^{\ell}$ is naturally identified with $\mathcal{A} \otimes_{\mathcal{O}} \mathcal{M}$, and the non-characteristic condition is equivalent to the classical one, and one can thus conclude using Kashiwara’s result.

2.5 The cellular dévissage of the derived Cauchy problem

In the more general case of a homotopy finitely presented \mathcal{D} -algebra \mathcal{A} , that is a retract of a cellular algebra $\mathcal{A}_0 = \mathcal{O} \rightarrow \mathcal{A}_1 \rightarrow \cdots \rightarrow \mathcal{A}_n = \mathcal{A}$, where there is, for each i , a homotopy cocartesian square

$$\begin{array}{ccc} \mathcal{A}_i & \longrightarrow & \mathcal{A}_{i+1} \\ \uparrow & & \uparrow \\ \mathrm{Sym}(\mathcal{M}_i) & \longrightarrow & \mathrm{Sym}(\mathcal{M}_{i+1}) \end{array}$$

with \mathcal{M}_i a compact complex of \mathcal{D} -modules, given by a complex with bounded coherent cohomology, and $\mathrm{Sym}(\mathcal{M}_i) \rightarrow \mathrm{Sym}(\mathcal{M}_{i+1})$ a cofibration.

These diagrams give cartesian diagrams

$$\begin{array}{ccc} f^{-1}\mathbb{R}\mathrm{Map}_{\mathcal{D}\text{-alg}}(\mathcal{A}_i, \mathcal{O}_X) & \longleftarrow & f^{-1}\mathbb{R}\mathrm{Map}_{\mathcal{D}\text{-alg}}(\mathcal{A}_{i+1}, \mathcal{O}_X) \\ \downarrow & & \downarrow \\ f^{-1}\mathbb{R}\mathrm{Map}_{\mathcal{D}\text{-alg}}(\mathrm{Sym}(\mathcal{M}_i), \mathcal{O}_X) & \longleftarrow & f^{-1}\mathbb{R}\mathrm{Map}_{\mathcal{D}\text{-alg}}(\mathrm{Sym}(\mathcal{M}_{i+1}), \mathcal{O}_X) \end{array}$$

and

$$\begin{array}{ccc}
\mathbb{R}\mathcal{M}ap_{\mathcal{D}\text{-alg}}(f_{\mathcal{D}}^*\mathcal{A}_i, \mathcal{O}_Y) & \longleftarrow & \mathbb{R}\mathcal{M}ap_{\mathcal{D}\text{-alg}}(f_{\mathcal{D}}^*\mathcal{A}_{i+1}, \mathcal{O}_Y) \\
\downarrow & & \downarrow \\
\mathbb{R}\mathcal{M}ap_{\mathcal{D}\text{-alg}}(f_{\mathcal{D}}^*\text{Sym}(\mathcal{M}_i), \mathcal{O}_Y) & \longleftarrow & \mathbb{R}\mathcal{M}ap_{\mathcal{D}\text{-alg}}(f_{\mathcal{D}}^*\text{Sym}(\mathcal{M}_{i+1}), \mathcal{O}_Y)
\end{array}$$

and a natural morphism ν between them.

If ν is an isomorphism at the level of the symmetric algebras, one shows by induction on i that it must be an isomorphism at the level of the cells \mathcal{A}_i . So it remains to show that one may apply Kashiwara's result, i.e., that the given \mathcal{D} -modules \mathcal{M}_i may be chosen so that the morphism f is non-characteristic for them.

3 The derived analytic Cauchy problem

It is possible to give another derived generalization of the nonlinear Cauchy problem to the non relatively algebraic situation where the function F of the Cauchy problem

$$\partial_t^m u = F(t, x, u, \partial_x^\alpha \partial_t^i u), \quad \partial_t^k u|_{t=0} = \varphi_k, \quad k = 0, \dots, m-1$$

is only analytic, using Porta's approach [Por14] to derived analytic geometry. This may be done using the geometric formulation of nonlinear partial differential systems, through the use of Simpson's de Rham spaces.

3.1 Recollection about derived analytic geometry

For the convenience of the reader, we recall from [Por17], Section 2, some basic definitions of derived analytic geometry (see also [Por14]).

Definition 3. Let \mathcal{T}_{an} be the category whose objects are Stein open subsets of \mathbb{C}^n and whose morphisms are holomorphic functions, and let τ_{an} be the analytic topology on it. A morphism in \mathcal{T}_{an} is called admissible if it is an open immersion.

Definition 4. Let \mathcal{X} be an ∞ -topos. An analytic ring in \mathcal{T} is a functor

$$\mathcal{O} : \mathcal{T}_{an} \rightarrow \mathcal{X}$$

such that \mathcal{O} commutes with products and pullbacks along admissible morphisms, and \mathcal{O} takes τ_{an} -covers to effective epimorphisms in \mathcal{X} . A morphism $f : \mathcal{O} \rightarrow \mathcal{O}'$ of analytic rings is said to be local if for every admissible morphism $\varphi : U \rightarrow V$ in \mathcal{T}_{an} , the square

$$\begin{array}{ccc}
\mathcal{O}(U) & \longrightarrow & \mathcal{O}(V) \\
\downarrow & & \downarrow \\
\mathcal{O}'(U) & \longrightarrow & \mathcal{O}'(V)
\end{array}$$

is a pullback square. A pair $(\mathcal{X}, \mathcal{O})$ is called an analytically structured ∞ -topos.

The main example of analytically structured ∞ -topos is given by the following.

Example 1. Let X be a \mathbb{C} -analytic space and X^{top} be the underlying topological space of X . Let $\mathcal{X} = \text{Sh}(X^{top})$ be the ∞ -topos of sheaves on X^{top} . One defines an analytic ring

$$\mathcal{O} : \mathcal{T}_{an} \rightarrow \mathcal{X}$$

by sending $U \in \mathcal{T}_{an}$ to the sheaf $\mathcal{O}(U)$ on X^{top} defined by

$$\mathcal{O}(U)(V) = \text{Hom}_{\text{AN}}(V, U).$$

Remark that this structure gives back the usual structure sheaf of X simply by evaluation of \mathcal{O} at the analytic affine line $U = \mathbb{A}^1$.

Definition 5. A derived \mathbb{C} -analytic space is an analytically structure ∞ -topos $(\mathcal{X}, \mathcal{O}_X)$ such that

1. locally on \mathcal{X} , $(\mathcal{X}, \pi_0(\mathcal{O}_X))$ is equivalent to a structured topos arising in Example 1;
2. the sheaves $\pi_i(\mathcal{O}_X(\mathbb{A}^1))$ are coherent as sheaves of $\pi_0(\mathcal{O}_X(\mathbb{A}^1))$ -modules.

We denote $\text{DAN}_{\mathbb{C}}$ the category of derived analytic spaces with local morphisms between them. A derived \mathbb{C} -analytic space is called Stein if $(\mathcal{X}, \pi_0(\mathcal{O}_X))$ is a Stein analytic space. We denote by $\text{DSTN}_{\mathbb{C}}$ the ∞ -category of derived Stein spaces. The category of derived analytic stacks is defined as

$$\text{DANST}_{\mathbb{C}} := \text{Sh}_{\mathcal{S}}(\text{DSTN}_{\mathbb{C}}, \tau_{an}).$$

There is a natural embedding

$$\text{DAN}_{\mathbb{C}} \rightarrow \text{DANST}_{\mathbb{C}}.$$

3.2 The de Rham stack and geometry of derived nonlinear PDEs

We recall from Porta's paper [Por17], Section 3, the definition of the de Rham stack.

Definition 6. The de Rham functor

$$(-)_{dR} : \text{DANST}_{\mathbb{C}} \rightarrow \text{DANST}_{\mathbb{C}}$$

is defined, for $U \in \text{DSTN}_{\mathbb{C}}$ by

$$X_{dR}(U) := X(t_0(U)_{red})$$

where $t_0(U)$ denote the classical truncation of a derived analytic space U and V_{red} denotes the reduced analytic space associated to a given (classical) analytic space V . There is a natural projection map

$$p : X \rightarrow X_{dR}$$

induced by the natural map

$$t_0(U)_{red} \rightarrow U.$$

Definition 7. Let $X \in \text{DANST}_{\mathbb{C}}$ be a derived analytic stack. The right adjoint p_* to the inverse image functor $p^* : \text{DANST}_{\mathbb{C}}/X_{dR} \rightarrow \text{DANST}_{\mathbb{C}}/X$ is denoted

$$\text{Jet}_X : \text{DANST}_{\mathbb{C}}/X \rightarrow \text{DANST}_{\mathbb{C}}/X_{dR}$$

and called the X -Jet functor. A derived nonlinear partial differential system on the sections of a morphism $E \rightarrow X$ of derived analytic stacks is a subspace Z of $\text{Jet}_X E$ defined over X_{dR} .

Remark 1. It is a classical result that if $X \in \text{AN}_{\mathbb{C}}$ is a smooth complex analytic space, then an \mathcal{O} -module over X_{dR} is the same as a \mathcal{D}_X -module. This result will be useful for the study of the characteristic variety of derived analytic nonlinear partial differential systems.

The non-linear analog of the derived solution space $\mathbb{R}\text{Hom}_{\mathcal{D}}(\mathcal{M}, \mathcal{O})$ of a \mathcal{D} -module \mathcal{M} is given by the derived solution space

$$\mathbb{R}\text{Map}_{X_{dR}}(X_{dR}, Z)$$

of an X_{dR} -stack Z . By replacing X_{dR} on the left by an arbitrary X_{dR} -stack, we get the derived solution space

$$\mathbb{R}\text{Map}_{X_{dR}}(Z', Z)$$

for the nonlinear partial differential system Z with values in Z' .

3.3 The classical Cauchy problem and geometry of PDEs

By translating what we did on the formulation of the relatively algebraic Cauchy problem in geometric terms, we get a natural formulation of the derived analytic Cauchy problem.

Definition 8. Let Y be a derived analytic stack. Two derived Y_{dR} -spaces Z and Z' and a morphism $f : X \rightarrow Y$ of complex analytic manifolds are said to satisfy the derived Cauchy condition if the natural morphism

$$\nu : f^{-1}\mathbb{R}\text{Map}_{Y_{dR}}(Z', Z) \longrightarrow \mathbb{R}\text{Map}_{X_{dR}}(f_{dR}^*Z', f_{dR}^*Z)$$

is an isomorphism of sheaves over Y , where $f_{dR} : X_{dR} \rightarrow Y_{dR}$ is the natural morphism between analytic de Rham spaces.

As before, one may show that if Z is the solution space of the equation

$$\partial_t^m u = F(t, x, u, \partial_x^\alpha \partial_t^i u), \quad \partial_t^k u|_{t=0} = \varphi_k, \quad k = 0, \dots, m-1$$

and $Z' = Y_{dR}$, the validity of the above result (at the level of π_0) is equivalent to the usual Cauchy-Kowalevskaya theorem, that is known to be true, even if F is an analytic function.

3.4 Characteristic variety of a derived nonlinear PDE

A definition of the characteristic variety of a nonlinear PDE was proposed by Baechtold in [Bae09] in a non-derived setting. We propose here another construction that is better adapted to the derived situation.

Let Y be a smooth analytic space and $p : Y \rightarrow Y_{dR}$ be the natural projection. Let Z be a derived space over Y_{dR} . Its relative cotangent complex gives an \mathcal{O} -module \mathbb{L}_Z over Z relative to Y_{dR} . Let $q : p^*Z \rightarrow Y$ be the natural projection and denote $\mathcal{A} := q_*\mathcal{O}_{p^*Z}$. The pullback of \mathbb{L}_Z along $p : Y \rightarrow Y_{dR}$ gives an $\mathcal{A}[\mathcal{D}]$ -module over the space Y . If we make the necessary hypothesis that this module is perfect, we may take its left dual to get another $\mathcal{A}[\mathcal{D}]$ -module \mathbb{T}_Z^ℓ called the tangent complex of the derived space. By pullback and tensorization, we may extend this to an $\mathcal{A}[\mathcal{E}_Y]$ -module over $p^*Z \times_Y T^*Y$. The support of this module will be called the characteristic variety of Z and denoted $\text{Char}(Z)$.

For $f : X \rightarrow Y$ a morphism, we have a cartesian diagram

$$\begin{array}{ccccc} p^*Z \times_Y T^*X & \xleftarrow{f_d} & p^*Z \times_Y X \times_Y T^*Y & \xrightarrow{f_\pi} & p^*Z \times_Y T^*Y \\ \downarrow & & \downarrow \pi & & \downarrow \\ X & \xleftarrow{id_X} & X & \xrightarrow{f} & Y \end{array}$$

and we may say f is non-characteristic for Z if

$$f_\pi^{-1}(\text{Char}(Z)) \cap p^*Z \times_Y T_X^*Y \subset p^*Z \times_Y X \times_Y T_Y^*Y.$$

3.5 About the non-characteristic derived Cauchy problem

We propose here a general formulation of the nonlinear derived Cauchy-Kowalevskaya-Kashiwara problem.

Conjecture 2 (Analytic non-characteristic derived Cauchy problem). *Let $f : Y \rightarrow X$ be a morphism of complex manifolds and suppose that Z is a derived Y_{dR} -stack that is homotopy finitely presented and non-characteristic for f . Then the natural map*

$$\nu : f^{-1}\mathbb{R}\text{Map}_{Y_{dR}}(Y_{dR}, Z) \longrightarrow \mathbb{R}\text{Map}_{X_{dR}}(X_{dR}, f_{dR}^*Z)$$

is an isomorphism of sheaves of spaces over Y .

It is possible that a proper definition of ‘‘homotopy finitely presented stack’’ for the above conjecture to be true involves the use of a grothendieck topology analogous to Tate’s Grothendieck topology (see [Bam14]) in the definition of derived analytic stacks, because it has better finiteness properties than the usual complex analytic topology (it is moderate in Grothendieck’s sense, and the associated topos is locally a quasi-compact topological space).

References

- [Bae09] M. Baechtold. Fold-type solution singularities and characteristic varieties of nonlinear PDEs. *Phd*, 2009.
- [Bam14] F. Bambozzi. On a generalization of affinoid varieties. *ArXiv e-prints*, January 2014.
- [Kas70] Masaki Kashiwara. Algebraic study of systems of partial differential equations. *Mém. Soc. Math. France (N.S.)*, 63:xiv+72, 1995 (1970).
- [Nir72] L. Nirenberg. An abstract form of the nonlinear Cauchy-Kowalewski theorem. *J. Differential Geometry*, 6:561–576, 1972.
- [Por14] Mauro Porta. *Géométrie analytique dérivée*. Thèse. université de Paris 7, 2014.
- [Por17] Mauro Porta. The derived riemann-hilbert correspondence, 2017.
- [Sch13] P. Schapira. An introduction to \mathcal{D} -modules. *lecture notes*, 2013.
- [TV07] Bertrand Toën and Michel Vaquié. Moduli of objects in dg-categories. *Ann. Sci. École Norm. Sup. (4)*, 40(3):387–444, 2007.