

Flat bundles, von Neumann algebras and K -theory with \mathbb{R}/\mathbb{Z} -coefficients

by

PAOLO ANTONINI, SARA AZZALI AND GEORGES SKANDALIS*

Abstract

Let M be a closed manifold and $\alpha : \pi_1(M) \rightarrow U_n$ a representation. We give a purely K -theoretic description of the associated element in the K -theory group of M with \mathbb{R}/\mathbb{Z} -coefficients ($[\alpha] \in K^1(M; \mathbb{R}/\mathbb{Z})$). To that end, it is convenient to describe the \mathbb{R}/\mathbb{Z} - K -theory as a relative K -theory of the unital inclusion of \mathbb{C} into a finite von Neumann algebra B . We use the following fact: there is, associated with α , a finite von Neumann algebra B together with a flat bundle $\mathcal{E} \rightarrow M$ with fibers B , such that $E_\alpha \otimes \mathcal{E}$ is canonically isomorphic with $\mathbb{C}^n \otimes \mathcal{E}$, where E_α denotes the flat bundle with fiber \mathbb{C}^n associated with α . We also discuss the spectral flow and rho type description of the pairing of the class $[\alpha]$ with the K -homology class of an elliptic selfadjoint (pseudo)-differential operator D of order 1.

Key Words: II_1 -factors, R/Z - K -theory, flat bundles, relative eta invariant.

Mathematics Subject Classification 2010: Primary: 46L80; Secondary: 58J28, 19K56.

Contents

1	Introduction	276
2	Preliminary constructions	279
2.1	Full modules	279
2.2	Mapping cones	280
2.3	Relative K -theory	280
2.4	The bootstrap category	282
2.5	K -theory of von Neumann algebras	282

*This research took place at the Institut de Mathématiques de Jussieu, CNRS - University Paris Diderot.

Paolo Antonini is funded by the *Projet ANR KInd*.

Sara Azzali is funded by an *INdAM-Cofund Fellowship*.

We thank these institutions for their support.

3	<i>K</i>-theory with coefficients	283
3.1	<i>K</i> -theory with rational and real coefficients	283
3.2	<i>K</i> -theory with $\mathbb{Z}/n\mathbb{Z}$ -coefficients	284
3.3	<i>K</i> -theory with \mathbb{Q}/\mathbb{Z} -coefficients	285
3.4	<i>K</i> -theory with \mathbb{R}/\mathbb{Z} -coefficients	285
3.5	Chern–Weil theory	287
3.5.1	Chern–Weil theory for K_0	287
3.5.2	Chern–Weil theory for K_1	288
3.5.3	Chern–Simons class	289
4	Atiyah’s theorem for covering spaces	290
5	$[\alpha]_{new}$ in $K^1(M; \mathbb{R}/\mathbb{Z})$ associated with a flat bundle $\alpha: \Gamma \rightarrow U(n)$	292
6	The pairing with $[D] \in K_1(M)$ and the spectral flow description	295
6.1	Various notions of connections	295
6.2	The pairing as an element of $K_1(\mathbb{Z}_{i_0})$	296
6.3	The pairing as an element of \mathbb{R}/\mathbb{Z}	297
6.4	The pairing as a rho invariant	298
	References	300

1. Introduction

Secondary invariants of geometric elliptic operators, such as the rho invariant of a unitary representation $\alpha : \Gamma \rightarrow U_n$ of the fundamental group, gain stability properties only when reduced modulo \mathbb{Z} . Indeed, Atiyah, Patodi and Singer in the seminal papers [APS2, APS3] proved that the modulo \mathbb{Z} class of the reduced rho invariant of an elliptic selfadjoint operator D can be described as the pairing of the K -homology class $[D]$ with a K -theory class $[\alpha]$ with \mathbb{R}/\mathbb{Z} -coefficients associated with α . This result is called the *index theorem for flat bundles*. The construction of $[\alpha] \in K^1(M, \mathbb{R}/\mathbb{Z})$ is using cohomology, and is based on the fact that K -theory with real coefficients is isomorphic to $H^*(M; \mathbb{R})$.

The model of $K^*(X, \mathbb{R}/\mathbb{Z})$ in [APS3] is formed from two components: the so called ‘torsion part’ $K^*(X, \mathbb{Q}/\mathbb{Z})$, defined as the inductive limit of $K^*(X, \mathbb{Z}/n!\mathbb{Z})$, and the ‘free part’ $K^*(X, \mathbb{R}) := K^*(X) \otimes \mathbb{R}$. The two parts are combined together by defining $K^*(X, \mathbb{R}/\mathbb{Z})$ as the quotient of $K^*(X, \mathbb{Q}/\mathbb{Z}) \oplus K^*(X, \mathbb{R})$ by the diagonal image of $K^*(X, \mathbb{Q}) := K^*(X) \otimes \mathbb{Q}$. The model relies therefore on the functorial properties of ordinary K -theory.

Atiyah, Patodi and Singer then suggested that a direct description can be given in terms of von Neumann algebras. This idea has been an inspiration for many authors. See in particular [DHK, DHK2, Hu, KP, Ba].

Beyond the Atiyah–Patodi–Singer (APS) one, a number of models of the \mathbb{R}/\mathbb{Z} - K -theory can be found in the literature, each one has its own features and flavor. Karoubi’s and Lott’s models are based on Chern–Weil and Chern–Simons theory [Ka, Lo]. Basu implemented the APS suggestion building a model with bundles of modules over von Neumann algebras [Ba]. Apart from the model of Karoubi, the constructions in [Lo, Ba] are based in a more or less explicit way on the notion of connection.

The purpose of this paper is to give a canonical construction of the \mathbb{R}/\mathbb{Z} - K -theory class associated with a flat bundle using operator algebraic tools, and to compute the pairing with K -homology as a Kasparov product. In particular, the models of K -theory with coefficients used here are purely operator theoretic.

Note that for a C^* -algebra A in the bootstrap category, one can define the K -theory of A with \mathbb{R} -coefficients as $K_*(A; \mathbb{R}) := K_*(A \otimes B)$, where B is any II_1 -factor: in fact, by the Künneth property, the group $K_*(A \otimes B)$ is independent of B up to a canonical isomorphism - and coincides with the APS model for A commutative.

The model for the K -theory with \mathbb{R}/\mathbb{Z} -coefficients that we use is simply the ordinary relative K -theory of the inclusion $A \hookrightarrow A \otimes B$ or, equivalently, the group $K_{*-1}(A \otimes C_{i_0})$ where C_{i_0} is the mapping cone of the unital inclusion $i_0 : \mathbb{C} \hookrightarrow B$ for any II_1 -factor B . This is immediately shown to be well defined for any C^* -algebra in the bootstrap category. Let us mention that this mapping cone has been used recently by Deeley to construct a model of K -homology with \mathbb{R}/\mathbb{Z} -coefficients [De].

To show that all the models are equivalent under canonical isomorphisms, we use also operator algebraic definitions of the K -theory with coefficients in \mathbb{Q} , $\mathbb{Z}/n\mathbb{Z}$ and \mathbb{Q}/\mathbb{Z} . Our models are direct noncommutative generalizations of the APS ones: we have the inclusion of \mathbb{C} in $n \times n$ -matrices - and inductive limits (UHF algebras), whereas the APS definition employs the K -theory of the mapping cones of maps of degree n on the circle - and projective limits.

Let M be a closed manifold with fundamental group Γ . If E_α is a flat vector bundle over M with holonomy $\alpha : \Gamma \rightarrow U(n)$, Atiyah, Patodi and Singer’s class $[\alpha]_{APS} \in K^1(M; \mathbb{R}/\mathbb{Z})$ is based on the idea that E_α defines a torsion element in the reduced K -theory of M , in fact there exists $k \in \mathbb{N}^*$ such that the sum of k copies of E_α is trivial. The class of E_α in K -theory with $\mathbb{Z}/k\mathbb{Z}$ -coefficients is then added to the transgression form corresponding to the two flat connections on the trivial bundle with fiber \mathbb{C}^{nk} thus obtained.

Our main result is the direct description of the element $[\alpha] \in K^1(M; \mathbb{R}/\mathbb{Z})$

associated with α : this is given by the pair of bundles E_α, \mathbb{C}^n with the (almost) canonical isomorphism after tensoring with a II_1 -factor B . Our construction consists of the following points:

1. there is a canonical flat bundle \mathcal{E} with fiber a II_1 -factor B associated with a morphism $u : \Gamma \rightarrow U(B)$ (where U is the unitary group of B);
2. there is a canonical isomorphism $E_\alpha \otimes \mathcal{E} \rightarrow \mathbb{C}^n \otimes \mathcal{E}$;
3. the bundle \mathcal{E} is trivial in K -theory and may actually be assumed to be trivial.

The II_1 -factor B can be taken to be $L^\infty(U_n) \rtimes \Gamma$ - i.e. the II_1 von Neumann algebra Morita equivalent to the foliated flat bundle $\tilde{M} \times_\alpha U_n$ - and the bundle \mathcal{E} encodes the bundle of frames for E_α .

The canonical isomorphism in (2) derives from the fact that a vector bundle becomes trivial when it is lifted to the bundle of frames.

The point (3) is based on the fact that a flat bundle with fiber a II_1 -factor is trivial in K -theory. This is easily seen applying Atiyah's L^2 -index theorem for covering spaces [Ati] (and its generalizations [Lü, Sc] by Lück and Schick to every trace on $C^*(\Gamma)$ - see also [Si] ⁽¹⁾), together with the property that K -homology separates points of $K^*(M; \mathbb{R})$. Conversely, Chern–Weil theory can be used to prove the same result, and this in turn gives a different proof of the L^2 -index theorem and its generalizations.

We then establish the independence from all choices involved in the construction of $[\alpha]$, and further show - using a Chern–Simons transgression argument with coefficients in von Neumann algebras - that our element is the same as the one constructed by Atiyah, Patodi and Singer.

We finally pair the K -theory class $[\alpha]$ with a K -homology class $[D] \in K_*(M)$ represented by a first order elliptic operator. The pairing is realized as a Kasparov product

$$KK^0(\mathbb{C}, C(M) \otimes C_{i_0}) \times KK^1(C(M), \mathbb{C}) \longrightarrow KK^1(\mathbb{C}, C_{i_0}) = K_1(C_{i_0}) = \mathbb{R} / \mathbb{Z} .$$

To compute the intersection product $[\alpha] \otimes [D]$, it is convenient to look at $[\alpha]$ in $KK(\mathbb{C}, Z_{i_0} \otimes C(M))$, the K -theory of the double cylinder of the inclusion $C(M) \hookrightarrow C(M) \otimes B$. This provides a path which interpolates between the operator D^α obtained as a twisted tensor product of D by the flat bundle E_α , and the operator D^n obtained by tensoring D by the trivial bundle of rank $n = \text{rk}(E_\alpha)$. The KK -product puts D^α and D^n as affiliated to the same II_∞ factor with (relatively)

¹As a side remark, we note that this result remains true for any trace on $\ell^1(\Gamma)$ by showing that Atiyah's method extends.

compact resolvent, that are bounded perturbations of each other since they have the same principal symbol. Furthermore, they have discrete spectrum and the corresponding spectral projections have integer von Neumann dimension since they are obtained by tensoring with B the Dirac operators D^α and D^n .

The pairing, as an element of \mathbb{R}/\mathbb{Z} , is given by a type II spectral flow: for every $a \in \mathbb{R}$, there is a well defined index between the two projections $\chi_{[a, \infty[}(D^\alpha)$ and $\chi_{[a, \infty[}(D^n)$ whose difference is relatively compact; this index does not depend on a up to \mathbb{Z} .

Finally, since the spectral flow of Dirac type operators is related through variational formulas to the eta invariant (see [BF, BGV, MZ2, CP1]), we get the index theorem for flat bundles $[\alpha] \otimes [D] = \xi(D^\alpha) - \xi(D^n) \in \mathbb{R}/\mathbb{Z}$.

2. Preliminary constructions

We start with some well known constructions for C^* -algebras that we shall use in the following.

2.1. Full modules

Let A be a unital C^* -algebra (or just a ring). A (right) A -module F is said to be *full* if $\text{span}\{\ell(x), x \in F, \ell \in \mathcal{L}(F, A)\} = A$ (where $\mathcal{L}(F, A)$ denotes the space of A -linear maps from F to A). If A is a non unital C^* -algebra, F is full if $\text{span}\{\ell(x), x \in F, \ell \in \mathcal{L}(F, A)\}$ is dense in A .

Lemma 2.1 *Let A be unital and F be a finitely generated projective full module over A . Let E, E' be finitely generated projective modules over A . Then $[E] = [E']$ in $K_0(A)$ if and only if $\exists k \in \mathbb{N} : E \oplus F^k \simeq E' \oplus F^k$.*

Proof: By hypothesis there exists a finitely generated projective G such that $E \oplus G \simeq E' \oplus G$. Write $1 = \sum \ell_i(x_i)$ with $\ell_1, \dots, \ell_n \in F^*$ and $x_i \in F$; thus we construct the module map $f : F^n \rightarrow A, (y_1, \dots, y_n) \mapsto \sum_i \ell_i(y_i)$ which is onto. Since G is finitely generated, there is a onto module map $g : A^m \rightarrow G$; we deduce a surjective module map $h : F^{nm} \rightarrow G$.

Then it also holds $F^{nm} \simeq G \oplus G'$ (projectivity of G), and then $E \oplus F^{nm} \simeq E' \oplus F^{nm}$.

The converse is obvious. □

Lemma 2.2 *Let A be a unital C^* -algebra, and E_1 and E_2 be finitely generated projective and full A -modules. Suppose $[E_1] = [E_2]$ in $K_0(A)$. Then there exists n such that $E_1^n \simeq E_2^n$.*

Proof: Since E_1 and E_2 are full, there exists k such that $E_1^k \oplus E_1 \simeq E_1^k \oplus E_2$ and $E_1 \oplus E_2^k \simeq E_2 \oplus E_2^k$.

Then $E_1^{a+1} \oplus E_2^b \simeq E_1^a \oplus E_2^{b+1}$, as soon as $a \geq k$ or $b \geq k$. We obtain $E_1^{2k} \simeq E_2^{2k}$. □

2.2. Mapping cones

The mapping cone of a morphism $\varphi : A \rightarrow B$ of (unital) C^* -algebras, is the algebra $C_\varphi = \{(a, \gamma) \in A \oplus C_0((0, 1], B) \mid \gamma(1) = \varphi(a)\}$.

Let now A_0, A_1, B be unital C^* -algebras.

Definition 2.3 Given two morphisms $\varphi_i : A_i \rightarrow B, i = 0, 1$, define the double cylinder algebra as

$$Z_{\varphi_0, \varphi_1} = \{(a_0, \sigma, a_1) \text{ with } a_i \in A_i, \sigma \in C([0, 1], B) \text{ s.t. } \sigma(i) = \varphi_i(a_i), i = 0, 1\}. \tag{2.1}$$

When $\varphi_0 = \varphi_1 = \varphi$ we will denote the double cylinder Z_{φ_0, φ_1} by Z_φ .

Note that for $A_0 = 0$ this construction is the cone C_{φ_1} . When $A_0 = B$ and φ_0 is the identity of B , the double cylinder is sometimes called the mapping cylinder of φ_1 .

For the double cylinder Z_φ there is a split exact sequence

$$0 \longrightarrow C_\varphi \xrightarrow{r} Z_\varphi \begin{matrix} \xleftarrow{s} \\ \xrightarrow{\pi_0} \end{matrix} A \longrightarrow 0 \tag{2.2}$$

with $r(a, \sigma) = (a, \sigma, a)$ and $\pi_1(a_0, \sigma, a_1) = a_1$, and the splitting given by $s(a) = (a, \varphi(a), a)$ so that $K_*(Z_\varphi) \simeq K_*(C_\varphi) \oplus K_*(A)$. In particular, there is a map from $K_0(Z_\varphi)$ to the summand $K_0(C_\varphi)$ of the form $x \mapsto x - (s \circ \pi_1)_* x$.

Note that if φ is injective, then

$$C_\varphi = \{\gamma \in C_0((0, 1], B) \mid \gamma(1) \in \varphi(A)\}$$

and

$$Z_\varphi = \{\gamma \in C([0, 1], B) \mid \gamma(0), \gamma(1) \in \varphi(A)\}.$$

If B is a II_1 -factor, the inclusion $i_0 : \mathbb{C} \rightarrow B$ gives $K_0(C_{i_0}) = 0$ and $K_1(Z_{i_0}) = K_1(C_{i_0}) = \mathbb{R}/\mathbb{Z}$.

2.3. Relative K -theory

Although relative K -theory coincides with the K -theory of a mapping cone, it helps in giving a more direct description of some K -theory elements. For instance, it

has been recently used by Deeley to construct a model of K -homology with \mathbb{R}/\mathbb{Z} -coefficients [De, Prop. 2.2, Ex. 5.3].

Let $\varphi : A \rightarrow B$ be a homomorphism of unital C^* -algebras. The group $K_0(\varphi)$ is given by generators and relations:

- Its generators are triples (E^+, E^-, u) where E^+, E^- are finitely generated projective A -modules, and $u : E^+ \otimes_A B \rightarrow E^- \otimes_A B$ is an isomorphism.
- Addition is given by direct sums.
- A homotopy is a triple associated with the map $C([0, 1]; A) \rightarrow C([0, 1]; B)$ induced by φ .
- Trivial elements are those triples (E^+, E^-, u) for which $u = \varphi(v)$ for some isomorphism $v : E^+ \rightarrow E^-$. The group $K_0(\varphi)$ is formed as the set of those triples modulo homotopy and addition of trivial triples.
- For non unital algebras, we just put $K_0(\varphi) = K_0(\tilde{\varphi})$, where $\tilde{\varphi} : \tilde{A} \rightarrow \tilde{B}$ is obtained by adjoining units everywhere.
- The group $K_1(\varphi)$ is also formed in this way. Its generators are pairs (u, f) where u is a unitary in $M_n(A)$ and f is a continuous path in the unitaries of $M_n(B)$ joining $\varphi(u)$ to 1_n .
- There is a natural isomorphism $K_*(\varphi) \simeq K_*(C_\varphi)$. This morphism is almost tautological for K_1 . For K_0 , it is described as follows. Assume that φ is unital and use the isomorphism $K_*(Z_\varphi) = K_*(C_\varphi) \oplus K_*(A)$ associated to the split exact sequence (2.2). The class of a triple $(E^+, E^-, U) \in K_0(\varphi)$ is then the image in the summand $K_0(C_\varphi)$ of the Z_φ -module

$$F_U = \{(x_+; f; x_-) \in E^+ \times C([0, 1]; E^+ \otimes_A B) \times E^-\};$$

$$f(0) = x_+ \otimes 1 \text{ and } Uf(1) = x_- \otimes 1\}.$$

Let $e^+, e^- \in M_n(A)$ be projections such that $E^+ = e^+ A^n$, $E^- = e^- A^n$ and $u \in M_n(B)$ such that $u^*u = \varphi(e^+)$, $uu^* = \varphi(e^-)$ and satisfies $ux = U(x)$ for $x \in e_+ A^n = E^+$. Let $f, g \in C([0, 1])$ be given by $f(t) = \cos \frac{t\pi}{2}$ and $g(t) = \sin \frac{t\pi}{2}$. Then we can write $F_U = e Z_\varphi^{2n}$ where

$$e = \left(\begin{pmatrix} e^+ & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} f^2 \varphi(e^+) & fgu^* \\ fgu & g^2 \varphi(e^-) \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & e^- \end{pmatrix} \right) \in M_{2n}(Z_\varphi).$$

2.4. *The bootstrap category*

The bootstrap category is the smallest class \mathcal{N} of separable nuclear C^* -algebras containing commutative ones and closed under KK -equivalence [Bl, 22.3.4].

Every C^* -algebra A in the bootstrap category satisfies the Künneth formula for tensor products [Bl, Theorem 23.1.3]; i.e. for every C^* -algebra B

$$0 \longrightarrow K_\bullet(A) \otimes K_\bullet(B) \xrightarrow{\alpha} K_\bullet(A \otimes B) \xrightarrow{\sigma} \text{Tor}_1^{\mathbb{Z}}(K_\bullet(A), K_\bullet(B)) \longrightarrow 0 \tag{2.3}$$

where we denote $K_\bullet = K_0 \oplus K_1$ as a graded group. Here α has degree 0, and σ has degree 1. If one of $K_\bullet(A)$ and $K_\bullet(B)$ is torsion free, then α is an isomorphism.

Remark 2.4 If A is in the bootstrap category and B is a von Neumann algebra, then $K_1(B) = 0$ and $K_0(B)$ is torsion free, see section 2.5. Therefore $K_*(A) \otimes K_0(B) \simeq K_*(A \otimes B)$.

2.5. *K-theory of von Neumann algebras*

Any von Neumann algebra A is uniquely decomposed in a product

$$A = A_{I_f} \times A_{I_\infty} \times A_{II_1} \times A_{II_\infty} \times A_{III},$$

consequently the K -groups uniquely split

$$K_*(A) = K_*(A_{I_f}) \times K_*(A_{I_\infty}) \times K_*(A_{II_1}) \times K_*(A_{II_\infty}) \times K_*(A_{III}).$$

It is easy to see that if A is properly infinite then $K_0(A) = 0$ while it is always the case that $K_1(A) = 0$. It follows that the K -theory is reduced to that of the finite piece

$$K_0(A) = K_0(A_{I_f}) \oplus K_0(A_{II_1}).$$

If A is finite, there is a unique center valued trace $\text{tr}_A^u : A \longrightarrow Z(A)$ with the property that two projections $p, q \in M_n(A)$ are equivalent if and only if $\text{tr}_A^u(p) = \text{tr}_A^u(q)$. Then it follows from the universal property of K_0 that tr_A^u defined on projections extends to a *von Neumann center valued dimension* which is an injection

$$\dim_A^u : K_0(A) \longrightarrow Z(A)_{sa} := \{a \in Z(A), a = a^*\}.$$

If A is type II_1 , this is an isomorphism.

3. K -theory with coefficients

A model for the K -theory of a C^* -algebra A with coefficients in a countable abelian group Λ is $K_*(A; \Lambda) := K_*(A \otimes B_\Lambda)$ where B_Λ is a C^* -algebra in the bootstrap category with a specified isomorphism $K_0(B_\Lambda) \simeq \Lambda$ and such that $K_1(B_\Lambda) = 0$. If Λ is uncountable, such a model can be provided using (uncountable) inductive limits.

In this section, we describe K -theory with \mathbb{R}/\mathbb{Z} -coefficients in terms of von Neumann algebras. The two models coincide for C^* -algebras in the bootstrap category. This von Neumann description is suitable for the construction of the K -theory class of a flat bundle given in section 5.

In order to relate our model with the one of Atiyah, Patodi and Singer [APS3, Sec. 5], we briefly discuss models for K -theory with coefficients in \mathbb{Q} , \mathbb{R} , $\mathbb{Z}/n\mathbb{Z}$ and \mathbb{Q}/\mathbb{Z} and compare them with the respective versions in [APS2].

3.1. K -theory with rational and real coefficients

For a (locally) compact space X , Atiyah–Patodi–Singer’s model is $K_{APS}^*(X, \mathbb{Q}) := K^*(X) \otimes \mathbb{Q}$ and $K_{APS}^*(X, \mathbb{R}) := K^*(X) \otimes \mathbb{R}$. More generally, we may set $K_*^{APS}(A, \mathbb{Q}) := K_*(A) \otimes \mathbb{Q}$ and $K_*^{APS}(A, \mathbb{R}) := K_*(A) \otimes \mathbb{R}$ for any C^* -algebra A .

Our description of K -theory with rational coefficients is the following.

Definition 3.1 Let D be the universal UHF algebra, *i.e.* the one which satisfies $K_0(D) = \mathbb{Q}$. It is obtained as an inductive limit of matrices, $D = \bigotimes_{n \in \mathbb{N}} M_n(\mathbb{C})$.

Define

$$K_*(A, \mathbb{Q}) := K_*(A \otimes D) \tag{3.1}$$

Since D is in the bootstrap category, $K_1(D) = 0$ and $K_0(D) = \mathbb{Q}$, the Künneth formula gives a natural isomorphism $K_*(A) \otimes \mathbb{Q} \rightarrow K_*(A \otimes D)$, and therefore our K -theory with rational coefficients coincides with $K_*^{APS}(A, \mathbb{Q})$.

In order to describe K -theory with real coefficients, we use the following result:

Lemma 3.2 *Let B be a II_1 -factor, and A be a C^* -algebra in the bootstrap category. Then the group $K_*(A \otimes B)$ is canonically isomorphic to $K_*(A) \otimes \mathbb{R}$, and therefore it does not depend on B up to a canonical isomorphism.*

Proof: The Künneth isomorphism

$$K_0(A) \otimes \mathbb{R} \xrightarrow{\sim} K_0(A \otimes B) \tag{3.2}$$

is given by the canonical map $[x] \otimes t \mapsto [x \otimes p_t]$. □

In particular, if B_1 is a II_1 -subfactor of B_2 , the induced map $K_*(A \otimes B_1) \longrightarrow K_*(A \otimes B_2)$ is a canonical isomorphism, independent of the inclusion $B_1 \subset B_2$. In the following, B will always denote a II_1 -factor. Then for A in the bootstrap category we can define, independently of B , a model for K -theory with coefficients in \mathbb{R} , which is naturally isomorphic to the APS one.

Definition 3.3 Let A be in the bootstrap category.

$$K_*(A, \mathbb{R}) := K_*(A \otimes B). \tag{3.3}$$

3.2. K -theory with $\mathbb{Z}/n\mathbb{Z}$ -coefficients

Let $i_n : \mathbb{C} \hookrightarrow M_n(\mathbb{C})$ be defined by $i_n(1) = \text{Id}$.

Definition 3.4

$$K_*(A, \mathbb{Z}/n\mathbb{Z}) := K_{*-1}(A \otimes C_{i_n}). \tag{3.4}$$

Let us relate it with the K -theory with $\mathbb{Z}/n\mathbb{Z}$ -coefficients as defined by Atiyah–Patodi–Singer in [APS2, Sec. 5].

Let $f_n : S^1 \rightarrow S^1$ be given by $f_n(z) = z^n$. The APS definition of the K -theory with $\mathbb{Z}/n\mathbb{Z}$ -coefficients for a (locally) compact space X is the relative K -theory of $f_n \times \text{id}_X : S^1 \times X \rightarrow S^1 \times X$, [APS2, (5.2)].

Define $f_n^* : C(S^1) \rightarrow C(S^1)$ by $f_n^*(\varphi)(z) = (\varphi \circ f_n)(z) = \varphi(z^n)$.

The K -theory of the cone algebras C_{i_n} and $C_{f_n^*}$ are computed by means of the exact sequences of the cones and give $K_1(C_{i_n}) \simeq K_0(C_{f_n^*}) \simeq \mathbb{Z}/n\mathbb{Z}$, and $K_0(C_{i_n}) \simeq K_1(C_{f_n^*}) = 0$.

Note that the inclusion $SM_n(\mathbb{C}) \rightarrow C_{i_n}$ induces the quotient map $\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$ in K_1 , and therefore the isomorphism $K_1(C_{i_n}) \simeq \mathbb{Z}/n\mathbb{Z}$ is canonical; in the same way, the isomorphism $K_0(C_{f_n^*}) \simeq \mathbb{Z}/n\mathbb{Z}$ is canonical, and there is therefore a canonical isomorphism $\psi : K_1(C_{i_n}) \rightarrow K_0(C_{f_n^*})$. By the UCT [Bl, Prop. 23.10.1], $KK^1(C_{i_n}, C_{f_n^*})$ is isomorphic to $\text{Hom}(\mathbb{Z}/n\mathbb{Z}, \mathbb{Z}/n\mathbb{Z})$. The isomorphism ψ determines therefore a generator of $KK^1(C_{i_n}, C_{f_n^*})$.

It follows that C_{i_n} and $C_{f_n^*}$ are naturally KK^1 -equivalent, or, equivalently, the algebras SC_{i_n} and $C_{f_n^*}$ are canonically KK -equivalent.

It is of course possible to give an explicit construction of the canonical element of $KK^1(C_{i_n}, C_{f_n^*})$.

As a consequence, let A be any separable C^* -algebra. Denote now with $i_{n,A} : A \hookrightarrow A \otimes M_n(\mathbb{C})$, $f_{n,A}^* : C(S^1, A) \rightarrow C(S^1, A)$ the analogous maps as above. Then $C_{i_{n,A}} = A \otimes C_{i_n}$ and $C_{f_{n,A}^*} = C_{f_n^*} \otimes A$ are canonically KK^1 -equivalent.

In particular, for a (locally) compact space X ,

$$K_{APS}^*(X, \mathbb{Z}/n\mathbb{Z}) \simeq K_{*+1}(C_0(X) \otimes C_{i_n}).$$

In other words, the definition 3.4 of K -theory of $\mathbb{Z}/n\mathbb{Z}$ coefficients coincides (for abelian C^* -algebras) with the one given in [APS2].

3.3. K -theory with \mathbb{Q}/\mathbb{Z} -coefficients

Definition 3.5 Let A be in the bootstrap category. Let D be the universal UHF algebra, and $A \hookrightarrow A \otimes D$ the obvious inclusion, define

$$K_*(A, \mathbb{Q}/\mathbb{Z}) := K_{*-1}(C_{A \hookrightarrow A \otimes D}). \tag{3.5}$$

Remark 3.6 For a (locally) compact space X , $K_{APS}^*(X, \mathbb{Q}/\mathbb{Z})$ is defined as the inductive limit of $K_{APS}^*(X, \mathbb{Z}/n!\mathbb{Z})$ [APS2, (5.3)]. More generally, for every C^* -algebra A , we may set $K_*^{APS}(A, \mathbb{Q}/\mathbb{Z}) := \varinjlim_n K_*^{APS}(A, \mathbb{Z}/n!\mathbb{Z})$. Observe that the construction in APS can be made using the projective limit. If

$$T = \varprojlim \dots S^1 \xrightarrow{\times n} S^1 \xrightarrow{\times(n+1)} S^1 \dots$$

then $K_*^{APS}(A, \mathbb{Q}/\mathbb{Z})$ is the relative K -theory of $A \otimes C(S^1) \rightarrow A \otimes C(T)$. Note also that $C(T) = C^*(\mathbb{Q})$.

We have $C_{A \hookrightarrow A \otimes D} = \varinjlim_n C_{A \hookrightarrow A \otimes M_{n!}(\mathbb{C})}$. As $K_*(A, \mathbb{Z}/n\mathbb{Z}) \simeq K_*^{APS}(A, \mathbb{Z}/n\mathbb{Z})$, using continuity of K -theory, we find $K_*(A, \mathbb{Q}/\mathbb{Z}) \simeq K_*^{APS}(A, \mathbb{Q}/\mathbb{Z})$.

3.4. K -theory with \mathbb{R}/\mathbb{Z} -coefficients

Definition 3.7 We propose the following realization of \mathbb{R}/\mathbb{Z} - K -theory of a C^* -algebra A in the bootstrap category.

$$K_*(A, \mathbb{R}/\mathbb{Z}) := K_{*+1}(\text{Cone}(A \hookrightarrow A \otimes B)),$$

where B is any II_1 -factor. Note that Definitions 3.3 and 3.7 are well posed: first of all since A is nuclear there is no ambiguity in the C^* -tensor product $A \otimes B$. Furthermore the right hand side is in both definitions independent, up to natural canonical isomorphism, of the factor B .

The latter definition is indeed a realization of \mathbb{R}/\mathbb{Z} - K -theory. The Bockstein change of coefficients is the long exact sequence associated to the mapping cone C^* -exact sequence.

Atiyah, Patodi and Singer’s description of \mathbb{R}/\mathbb{Z} - K -theory goes back to an idea of Segal [APS3, Sec. 5]. It is made up of two components. The first component is the torsion part, contained in $K^*(X, \mathbb{Q}/\mathbb{Z})$. The free part is in $K^*(X, \mathbb{R}) = K^*(X) \otimes \mathbb{R}$. More precisely

$$K_{APS}^*(X, \mathbb{R}/\mathbb{Z}) := \text{cokernel}\{(p, -j) : K^*(X, \mathbb{Q}) \longrightarrow K^*(X, \mathbb{Q}/\mathbb{Z}) \oplus K^*(X, \mathbb{R})\}$$

where p is the natural projection and j is the natural injection.

We may of course put

$$K_*^{APS}(A, \mathbb{R}/\mathbb{Z}) := \text{cokernel}\{(p, -j) : K_*(A, \mathbb{Q}) \longrightarrow K_*(A, \mathbb{Q}/\mathbb{Z}) \oplus K_*(A, \mathbb{R})\}.$$

Finally we prove:

Proposition 3.8 *For an algebra A in the bootstrap category, our realization of $K_*(A, \mathbb{R}/\mathbb{Z})$ coincides with $K_*^{APS}(A, \mathbb{R}/\mathbb{Z})$.*

Proof: Let D be the universal UHF algebra, B a II_1 -factor, and denote with $i_D : \mathbb{C} \hookrightarrow D$ and $i_B : \mathbb{C} \hookrightarrow B$ the obvious inclusions. It is enough to prove that

$$K_*(A \otimes C_{i_B}) = \text{Coker}\{(p, -j) : K_{*+1}(A \otimes D) \longrightarrow K_*(A \otimes C_{i_D}) \oplus K_{*+1}(A \otimes B)\} \tag{3.6}$$

where p is induced from the natural map $(C(M) \otimes D)(]0, 1]) \longrightarrow C_{i_D}$ and j is induced from any unital inclusion $D \hookrightarrow B$. To verify (3.6), consider the diagram

$$\begin{array}{ccc} A \otimes V_D & \xrightarrow{p} & A \otimes C_{i_D} \\ j \downarrow & & \downarrow \\ A \otimes V_B & \xrightarrow{h} & A \otimes C_{i_B} \end{array}$$

where $V_B = \{(a, s) \in C_0(]-1, 0]) \times B(]0, 1]) \text{ s.t. } i_B(a(0)) = r(0)\}$ and V_D is defined in analogous way. Note that the inclusion $A \otimes SB \rightarrow A \otimes V_B$ induces an isomorphism in K -theory as follows from the exact sequence $0 \rightarrow (A \otimes B)(]0, 1]) \rightarrow V_B \rightarrow A(]-1, 0]) \rightarrow 0$. Moreover h is surjective. Therefore, we have an associated Mayer-Vietoris exact sequence. Now $j_* : K_*(A \otimes V_D) \simeq K_{*+1}(A \otimes D) \longrightarrow K_{*+1}(A \otimes B) \simeq K_*(A \otimes V_D)$ is injective.

Finally the Mayer-Vietoris sequence gives

$$0 \longrightarrow K_*(A \otimes D) \longrightarrow K_{*+1}(A \otimes C_{i_D}) \oplus K_*(A \otimes B) \longrightarrow K_{*+1}(A \otimes C_{i_B}) \longrightarrow 0.$$

□

Remark 3.9 There are in the literature several other realizations of \mathbb{R}/\mathbb{Z} - K -theory:

Karoubi’s description is a cohomological interpretation based on Chern–Weil and Chern–Simons theory [Ka2] - this will be explained below. This was used to develop secondary characteristic classes in [Ka3, Ka4]. The description below works only for manifolds but it can be easily generalized to compact spaces by an embedding trick. Cycles for $K^1(X, \mathbb{R}/\mathbb{Z})$ are triples $((E, \nabla^E), (F, \nabla^F), \omega)$ where E, F are Hermitian bundles with Hermitian

connections ∇^E, ∇^F and ω is a transgression odd degree differential form such that

$$d\omega = \text{Ch}(\nabla^E) - \text{Ch}(\nabla^F).$$

A suitable notion of sum and equivalence of cycles is defined leading to an abelian group which is a realization of $K^1(X, \mathbb{R}/\mathbb{Z})$.

Basu’s description is a bundle theoretic relative description [Ba]. It can be thought of as a realization of the suggestion of Atiyah, Patodi and Singer to describe $K^1(X, \mathbb{R}/\mathbb{Z})$ with bundles of von Neumann algebras. Cycles are couples of finite rank vector bundles E, F such that there exists a von Neumann bundle V (relative to a semi-finite or II_∞ von Neumann algebra) such that $E \otimes V \simeq F \otimes V$. Addition of cycles is defined by direct sum and there is a natural equivalence relation. The equivalence classes form a group which is a realization of $K^1(X, \mathbb{R}/\mathbb{Z})$.

3.5. Chern–Weil theory

Let M be a closed manifold. Then the group $K_*(C(M), \mathbb{R})$ as defined above is isomorphic to $K_{APS}^*(M, \mathbb{R}) := K^*(M) \otimes \mathbb{R}$ as defined in [APS2]. This is in turn isomorphic to $H^*(M; \mathbb{R})$ using Chern–Weil theory.

Let A be a unital C^* -algebra. Let $W \rightarrow M$ be a smooth bundle of finitely generated Hilbert A -modules over the manifold M : using the Serre–Swann theorem, this is a finitely generated projective module over $C^\infty(M; A)$, endowed with a nondegenerate $C^\infty(M) \otimes A$ valued scalar product. We refer to [MF] and [Sc, Sections 2,3] for the general theory of such bundles. A connection ∇ is extended as usual to forms

$$\nabla(\omega \otimes s) = d\omega \otimes s + (-1)^{\text{deg}\omega} \omega \nabla s, \quad \omega \in \Omega^*(M), \quad s \in \Gamma(W).$$

The curvature ∇^2 is a 2-form with values in endomorphisms. If the connection is metric, the curvature is skew-adjoint.

3.5.1 CHERN–WEIL THEORY FOR K_0

If $\Omega = \nabla^2 \in \Omega^2(M; \text{End}_A(W))$ is the curvature, the exponential

$$\begin{aligned} \exp\left(\frac{\Omega}{2i\pi}\right) &:= \sum_k \frac{\Omega \wedge \cdots \wedge \Omega}{(2i\pi)^k k!} \in \Omega^{2*}(M; \text{End}_A(W)) \\ &\simeq \Omega^{2*}(M) \otimes_{C^\infty(M)} \text{End}_A(W) \end{aligned}$$

is well defined as a finite sum.

A trace $\tau : A \rightarrow \mathbb{C}$ extends to $M_n(A)$ and therefore to $\text{End}_A(E) \simeq pM_n(A)p$ for every finitely generated projective A -module $E \simeq pA^n$ (where $p \in M_n(A)$ is a projection). If $W \rightarrow M$ is a finitely generated projective A -module bundle, we thus obtain a well defined trace, which is a $C(M)$ -linear map still denoted $\tau : \text{End}_A(W) \rightarrow C(M)$. The Chern character of W associated with τ is defined as the cohomology class of the form

$$\text{Ch}_\tau(\Omega) := (\text{id} \otimes \tau) \left(\exp \left(\frac{\Omega}{2i\pi} \right) \right) \in \Omega^{2*}(M; \mathbb{C}).$$

From the trace property it follows in fact that the Chern form is closed. Its cohomology class does not depend on the connection. If τ is self adjoint, this class is seen to be real by taking ∇ to be metric.

Hilbert A -module bundles define the K -theory of a compact space with coefficients in A . Indeed $K(M; A)$ is the K -group of the category of finitely generated projective Hilbert A -module bundles. The Serre–Swann functor sending a bundle to the module of its continuous functions establishes a canonical isomorphism

$$K^0(M; A) \simeq K_0(C(M; A)) \simeq K_0(C(M) \otimes A).$$

If A is a II_1 -factor, from the commutativity of the diagram

$$\begin{array}{ccc} K^0(M) \otimes K_0(A) & \xrightarrow{\simeq} & K^0(M; A) \simeq K_0(C(M) \otimes A) \\ \text{Ch} \otimes \tau \downarrow & \swarrow \text{Ch}_\tau & \\ H^{2*}(M; \mathbb{R}) & & \end{array} \tag{3.7}$$

one gets that Ch_τ is an isomorphism.

3.5.2 CHERN–WEIL THEORY FOR K_1

The odd K -theory with coefficients in A is defined by

$$K^1(M; A) := K_c^0(M \times \mathbb{R}; A) \simeq K_0(C(M) \otimes C_0(\mathbb{R}) \otimes A) = K_0(C(M) \otimes SA)$$

where $SA = C_0(\mathbb{R}; A)$ is the suspension of A .

We then write $K^1(M; A) \times K^0(M; A) = K^0(M \times S^1; A)$. Also $\bigoplus_k H^{2k}(M \times S^1, \mathbb{C}) \simeq \bigoplus_k H^k(M, \mathbb{C})$. The map $H^{2k}(M \times S^1, \mathbb{C}) \rightarrow H^{2k-1}(M, \mathbb{C})$ is obtained by integration along the S^1 fibers. Using these identifications, if τ is a trace on A ,

we have a commuting diagram:

$$\begin{array}{ccc}
 K^0(M \times S^1; A) & \xrightarrow{\text{Ch}_\tau} & H^{2^*}(M \times S^1, \mathbb{C}) \\
 \downarrow & & \downarrow \\
 K^1(M; A) & \xrightarrow{\text{Ch}_\tau^{od}} & H^{2^*+1}(M; \mathbb{C})
 \end{array}$$

which defines the odd Ch_τ^{od} in the bottom line. In particular, if A is a II_1 -factor, then Ch_τ is an isomorphism.

Let $W \rightarrow S^1 \times M$ be a smooth bundle of finitely generated Hilbert A -modules over $S^1 \times M$ and ∇ a connection on W . Write $S^1 = \mathbb{R}/\mathbb{Z}$ and use ∇ in order to trivialize the bundle W along $[0, 1]$. We thus obtain a constant bundle W_0 over M and a family $(\nabla_u)_{u \in [0, 1]}$ of connections on W_0 . In other words, we may write (over $M \times [0, 1]$),

$$\nabla = (\nabla_u) + du \frac{\partial}{\partial u}.$$

Then $\nabla^2 = \dot{\nabla}_u \wedge du + \nabla_u^2$, whence:

$$\text{Ch}_\tau^{od}(W, \nabla) = (2\pi i)^{-1} \int_0^1 \tau \left(\dot{\nabla}_u \exp \left(\frac{\nabla_u^2}{2\pi i} \right) \right) du \in \Omega^{2^*+1}(M; \mathbb{C}).$$

Alternatively, an element of $K^1(M; A)$ is represented by a continuous (smooth) map $\alpha : M \rightarrow U_n(A)$. The corresponding bundle over $M \times S^1$ is obtained by gluing the trivial bundle $M \times [0, 1] \times A^n$ by means of α .

One then may compute Ch_τ for this bundle and get ([G, Prop. 1.2])

$$\text{Ch}_\tau(\alpha) = \sum_{k=0}^{\infty} (-1)^k \frac{k!}{(2k+1)!(2\pi i)^{k+1}} (\text{id} \otimes \tau)(\alpha^{-1} d\alpha)^{2k+1} \in \Omega^{2^*+1}(M; \mathbb{C}). \tag{3.8}$$

3.5.3 CHERN–SIMONS CLASS

Let A be a C^* -algebra and $\tau : A \rightarrow \mathbb{C}$ a trace. For a bundle E of finitely generated projective Hilbert A -modules equipped with two connections ∇_0, ∇_1 , the odd Chern–Simons differential form is

$$\text{CS}_\tau(\nabla_0, \nabla_1) := (2\pi i)^{-1} \int_0^1 \tau \left(\dot{\nabla}_u \exp \left(\frac{\nabla_u^2}{2\pi i} \right) \right) du \in \Omega^{2^*+1}(M; \mathbb{C}) \tag{3.9}$$

where $\nabla_u := (1-u)\nabla_0 + u\nabla_1$. Recall that $d(\text{CS}_\tau(\nabla_0, \nabla_1)) = \text{Ch}_\tau(\nabla_0) - \text{Ch}_\tau(\nabla_1)$.

Definition 3.10 A flat Hilbert A -module bundle is a pair (W, ∇) where W is a Hilbert A -module bundle, and ∇ is a flat connection.

A flat bundle isomorphism between two flat Hilbert A -module bundles (W, ∇^W) and (V, ∇^V) is a bundle isomorphism $\psi : W \rightarrow V$ which preserves the connection.

We fix the following notation, used repeatedly in the following. If (W, ∇^W) and (V, ∇^V) are flat Hilbert A -module bundles and $\alpha : W \rightarrow V$ is a isomorphism which does not necessarily preserve the connection, we use (3.9) to define the closed form

$$CS_\tau(\alpha^* \nabla^V, \nabla^W) \in \Omega^{odd}(M; \mathbb{C}). \tag{3.10}$$

From the above discussion we immediately get

Proposition 3.11 *If $\alpha : V \rightarrow V$ is an automorphism of a flat Hilbert A -module bundle, then α defines a class in $K_1(C(M) \otimes A)$ and the classes $Ch_\tau(\alpha)$ and $CS_\tau(\alpha^* \nabla^V, \nabla^V)$ in $H^*(M; \mathbb{C})$ coincide. \square*

4. Atiyah’s theorem for covering spaces

A key ingredient in our construction of the element $[\alpha] \in K^1(M; \mathbb{R}/\mathbb{Z})$ is the fact that every flat bundle on M with fibers a von Neumann factor of type II_1 is trivial (in K -theory). In fact this statement is equivalent to Atiyah’s theorem on covering spaces [Ati] (more precisely, to the versions of Lück and Schick in [Lü, Sc]). See also remark 4.3 below for a further generalization.

Let M be a closed manifold with $\pi_1(M) = \Gamma$. For an elliptic operator D acting on the sections of a bundle $S \rightarrow M$, let \tilde{D} be the lift on the universal covering. Recall that the index class

$$Ind \tilde{D} \in K_*(C^*\Gamma)$$

is the KK -product $[\mathcal{V}] \otimes_{C(M)} [D]$, where $\mathcal{V} = \tilde{M} \times_\Gamma C^*\Gamma$ is the so called Mishchenko bundle. Indeed, under the identification of the sections of $\mathcal{V} \otimes S$ with the Γ -invariant sections of the lifted bundle $\pi^*(\mathcal{V} \otimes S) = (C^*\Gamma \otimes \pi^*(S)) \times \tilde{M}$, one easily sees that \tilde{D} is a connection and a $C^*\Gamma$ -Fredholm operator. Its index is the

Connes–Moscovici index class given by the idempotent $\begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} + R$ for

$$R := \begin{pmatrix} S_0^2 & S_0(I + S_0)\tilde{Q} \\ S_1\tilde{D} & -S_1^2 \end{pmatrix}$$

where \tilde{Q} is any almost local parametrix of \tilde{D} and $S_0 = I - \tilde{Q}\tilde{D}$, $S_1 = I - \tilde{D}\tilde{Q}$ are smoothing.

Let now B be a II_1 -factor. A morphism $u : \Gamma \rightarrow \mathcal{U}(B)$ is a representation of Γ and therefore extends to a $*$ -morphism $\bar{u} : C^*\Gamma \rightarrow B$. Denote by E_u the bundle $E_u := \tilde{M} \times_u B$ over M with fiber B , let \mathcal{E}_u denote the corresponding (finitely generated projective right) $(C(M) \otimes B)$ -module of sections of E_u , and $[\mathcal{E}_u]$ its class in $K_0(C(M) \otimes B)$.

The von Neumann index of D (in $K_0(B)$) is

$$\text{Ind } \tilde{D}_B = \bar{u}(\text{Ind } \tilde{D}) = [\mathcal{E}_u] \otimes_{C(M)} [D]. \tag{4.1}$$

Lemma 4.1 *In $K_0(C(M) \otimes B)$ the classes \mathcal{E}_u and the trivial module $C(M) \otimes B$ coincide.*

Proof: The Chern character $\text{ch} : K_0(C(M) \otimes B) \rightarrow H^{ev}(M, \mathbb{R})$ is an isomorphism, as recalled in Section 3.5. Then the statement follows since E_u is a flat bundle. □

Remark 4.2 • Atiyah’s L^2 -index theorem (Lück’s and Schick version [Lü, Sc]) can be deduced from Lemma 4.1 with the following reasoning (cf. [Mi]). Given any trace τ on $C^*(\Gamma)$, one may construct a II_1 -factor B with a trace-preserving embedding $C^*(\Gamma) \rightarrow B$. It immediately follows from Lemma 4.1 and formula (4.1) that $\text{Ind } \tilde{D}_B = \text{Ind } D$.

- Conversely, Lemma 4.1 follows from Atiyah’s L^2 -index theorem (Lück’s and Schick version [Lü, Sc]). Indeed, for any trace τ on B , $\tau(\text{Ind } \tilde{D}_B) = \tau(\text{Ind } D \cdot [1])$, and the result follows from the fact that the pairing map $K^0(C(M)) \otimes \mathbb{R} \rightarrow \text{Hom}(K^0(M), \mathbb{R})$, $D \mapsto \langle \cdot, D \rangle$ is surjective since $K^0(M)$ is finitely generated.
- It would be interesting to find a K -theoretic proof of Lemma 4.1 with the idea: find a bigger II_1 -factor B_1 , i.e. a II_1 -factor containing B , such that the bundle $E_u \otimes_{C(M) \otimes B} C(M) \otimes B_1$ becomes explicitly trivial.

Remark 4.3 (Atiyah’s theorem for continuous traces on $\ell^1(\Gamma)$). The symmetric index of an elliptic (pseudo)-differential operator D is well defined as an element $\text{ind}_{\ell^1(\Gamma)}(\tilde{D})$ of $K_0(\ell^1(\Gamma))$ (see [CM]).

A continuous linear form on $\ell^1(\Gamma)$ is of the form $a \mapsto \sum_{g \in \Gamma} a_g f(g)$ for some $f \in \ell^\infty(\Gamma)$. The trace property holds if and only if f is constant on conjugacy classes. Let τ be a trace associated with $f \in \ell^\infty(\Gamma)$.

Now $\tau(\text{ind}_{\ell^1(\Gamma)}(\tilde{D})) = T_\tau(R)$ where T_τ is defined as follows: let first $p : \tilde{M} \rightarrow M$ be the universal covering map. Note that Γ acts freely and properly on \tilde{M} and $M = \Gamma \backslash \tilde{M}$ i.e. the fibers of p are the orbits of Γ .

Put $G = \Gamma \backslash \tilde{M}^2$ - where Γ acts diagonally on $\tilde{M} \times \tilde{M}$. Then G is a Lie groupoid and $C_c^\infty(G)$ is the algebra of Γ -invariant smooth kernels on \tilde{M} with a bounded support condition. Note that $R \in M_n(C_c^\infty(G))$.

Let $\varphi \in C_c^\infty(G)$. Let $x \in \tilde{M}$. Let $\varphi_x : \Gamma \rightarrow \mathbb{C}$ be defined by $\varphi_x(g) = \varphi(gx, x)$. For $h \in \Gamma$ and $g \in \Gamma$, we have $\varphi(ghx, hx) = \varphi(h^{-1}ghx, x)$ and therefore $\tau(\varphi_x) = \tau(\varphi_{hx})$. We then put $T_\tau(\varphi) = \int_M \tau(\varphi_x) dx$.

Now, if \tilde{D} and \tilde{Q} are taken local enough, we have $R(x, gx) = 0$ for all $x \in \tilde{M}$ and $g \neq 1$, and therefore

$$\tau(\text{ind}_{\ell^1(\Gamma)}(\tilde{D})) = \tau(1)\varepsilon(\text{ind}_{\ell^1(\Gamma)}(\tilde{D})) = \tau(1)\text{ind}(D)$$

where $\varepsilon : \Gamma \rightarrow \mathbb{C}$ is the trivial representation of Γ .

5. $[\alpha]_{new}$ in $K^1(M; \mathbb{R}/\mathbb{Z})$ associated with a flat bundle $\alpha : \Gamma \rightarrow U(n)$

The triviality property of Lemma 4.1 is crucial in what will be our definition of the element in $K^1(M; \mathbb{R}/\mathbb{Z})$ associated to a flat vector bundle.

Let $M = \Gamma \backslash \tilde{M}$ be a closed manifold with $\pi_1(M) = \Gamma$ and universal cover \tilde{M} . A flat bundle over M is just given by a homomorphism from Γ to the associated structure group.

Let B be a (finite) von Neumann algebra. A flat $C(M) \otimes B$ module with fiber B^n is given by a homomorphism u from Γ to the group $U_n(B)$ of unitaries of $M_n(B)$. The associated bundle is $\tilde{M} \times_\Gamma B^n$; the corresponding finitely generated projective module over $C(M) \otimes B$ is the set of continuous maps $f : \tilde{M} \rightarrow B^n$ such that $f(gx) = u(g)f(x)$ for all $g \in \Gamma$ and $x \in \tilde{M}$. A flat isomorphism of flat bundles associated with $u_+ : \Gamma \rightarrow U_n(B)$ and $u_- : \Gamma \rightarrow U_n(B)$ is given by a unitary $v \in U_n(B)$ intertwining the morphisms u_+ and u_- .

We first observe the following

Proposition 5.1 *Let M be a compact manifold and B be a II_1 -factor. Let \mathcal{E} be a flat $C(M) \otimes B$ module with fiber B (given by a homomorphism $u : \pi_1(M) \rightarrow U(B)$). Then there exists $\ell \in \mathbb{N}^*$ such that $\mathcal{E}^\ell \simeq (C(M) \otimes B)^\ell$.*

Proof: It follows from Lemma 4.1 that the class of \mathcal{E} agrees with that of the trivial bundle $C(M) \otimes B$ in $K_0(C(M) \otimes B)$. The conclusion follows using Lemma 2.2. □

A flat Hermitian vector bundle $E_\alpha \rightarrow M$ is given by a homomorphism $\alpha : \Gamma \rightarrow U_n$.

In the following we denote by $E^+ = C(M, E_\alpha)$ the corresponding (flat) $C(M)$ -module of sections and $E^- = C(M)^n$ the module of sections of the trivial bundle of rank n .

Proposition 5.2 *Let E_α be a flat unitary vector bundle as above.*

- a) *There exists a finite factor B , a flat $C(M) \otimes B$ module \mathcal{E} with fiber B and an isomorphism of flat bundles $v: E^- \otimes_{C(M)} \mathcal{E} \longrightarrow E^+ \otimes_{C(M)} \mathcal{E}$.*

We may of course assume B is a II_1 factor (up to tensoring by any II_1 factor). By Proposition 5.1, up to tensoring B by M_n , we have an isomorphism $\varphi: C(M) \otimes B \rightarrow \mathcal{E}$. Let w_v be the isomorphism

$$w_v := (1_{E^-} \otimes \varphi)^{-1} \circ v^{-1} \circ (1_{E^+} \otimes \varphi): E^+ \otimes B \rightarrow E^- \otimes B. \tag{5.1}$$

- b) *The class of (E^+, E^-, w_v) in $K^1(M, \mathbb{R}/\mathbb{Z})$ is independent of all choices $B, \mathcal{E}, v, \varphi$.*

Proof: a) Given a finite factor B and a flat $C(M) \otimes B$ module \mathcal{E} with fiber B associated with a morphism $u: \Gamma \rightarrow U(B)$, the flat bundle $E^- \otimes_{C(M)} \mathcal{E}$ corresponds to the morphism $g \mapsto 1 \otimes u(g) \in U_n(B) \subset M_n \otimes B$ and the flat bundle $E^+ \otimes_{C(M)} \mathcal{E}$ corresponds to the morphism $g \mapsto \alpha(g) \otimes u(g)$.

We therefore just have to construct a finite factor B with a morphism $u: \Gamma \rightarrow U(B)$ and an element $v \in U_n(B)$, such that, for all $g \in \Gamma$, we have $v(1 \otimes u(g)) = (\alpha(g) \otimes u(g))v$.

Let $\Gamma_1 = \alpha(\Gamma)$ be the image of Γ in U_n (with the discrete topology) and $K = \overline{\Gamma}_1$ its closure in U_n . Set $B_1 = L^\infty(K) \rtimes \Gamma_1$, where Γ_1 acts on K by left translation. Denote by $u: \Gamma_1 \rightarrow L^\infty(K) \rtimes \Gamma_1$ the canonical inclusion; the inclusion map $v: K \rightarrow U_n \subset M_n$ is an element of $M_n(L^\infty(K)) \subset M_n(B_1)$; by definition $(1 \otimes u_g)v(1 \otimes u_g)^{-1}$ is the function $x \mapsto v(g^{-1}x) = \alpha(g)^{-1}v(x)$, i.e. the element $(\alpha(g)^{-1} \otimes 1)v$ in $M_n \otimes L^\infty(K) \subset M_n(B_1)$.

The desired equality $v(1 \otimes u(g)) = (\alpha(g) \otimes u(g))v$ follows.

By density of Γ_1 (discrete) in K (compact), we have that B_1 is a finite factor.

- b) The choice of \mathcal{E} is equivalent to the choice of a morphism $u: \Gamma \rightarrow U(B)$; the choice of φ is equivalent to a continuous map $\psi: \tilde{M} \rightarrow U(B)$ such that $\psi(gx) = u(g)\psi(x)$ for all $g \in G$ and $x \in \tilde{M}$. The element w_v^{-1} is then the map $x \mapsto (1_{M_n} \otimes \psi(x))^{-1}v(1_{M_n} \otimes \psi(x))$ from \tilde{M} to $U_n(B)$ which satisfies $\psi(gx) = (\alpha(g) \otimes 1)\psi(x)$ and thus defines an isomorphism of bundles $E^- \otimes B \rightarrow E^+ \otimes B$.

Assume that we are given (B_j, u_j, v_j, ψ_j) with $j = 1, 2$. Put $B = B_1 \overline{\otimes} B_2$ (von Neumann tensor product of course), and define $u = u_1 \otimes u_2: g \mapsto u_1(g) \otimes u_2(g)$ and $\psi = \psi_1 \otimes \psi_2: x \mapsto \psi_1(x) \otimes \psi_2(x)$; put also $V_1 = v_1 \otimes 1_{B_2}$ and $V_2 = \beta(v_2)$, where $\beta: M_n \otimes B_2 \rightarrow M_n \otimes B_1 \otimes B_2$ is the obvious inclusion.

We find in this way two new families (B, u, V_1, ψ) and (B, u, V_2, ψ) with the desired properties.

For $x \in \tilde{M}$, we have $w_{V_1}(x) = (1_{M_n} \otimes \psi(x))^{-1} V_1^{-1} (1_{M_n} \otimes \psi(x)) = w_{v_1}(x) \otimes 1_{B_2}$; it follows that the elements of $K_1(M, \mathbb{R}/\mathbb{Z})$ corresponding to (B_1, u_1, v_1, ψ_1) and (B, u, V_1, ψ) coincide. Also $w_{V_2}(x) = \beta(w_{v_2}(x))$, and thus the elements of $K_1(M, \mathbb{R}/\mathbb{Z})$ corresponding to (B_2, u_2, v_2, ψ_2) and (B, u, V_2, ψ) coincide.

Finally, write $V_1 = V_2 W$ where W is a unitary in $M_n(Q)$ where $Q = B \cap \{u(g); g \in \Gamma\}'$. By connectedness of the unitary group of a von Neumann algebra, it follows that the elements of $K_1(M, \mathbb{R}/\mathbb{Z})$ corresponding to (B, u, V_1, ψ) and (B, u, V_2, ψ) coincide. □

Definition 5.3 Given $\alpha: \Gamma \rightarrow U_n$, E^+ and E^- as above. For any $(\mathcal{E}, \varphi, v)$ as in Proposition 5.2. We define

$$[\alpha]_{new} := [(E^+, E^-, w_v)] \in K_1(C(M), \mathbb{R}/\mathbb{Z}).$$

Theorem 5.4 Let E_α be a flat unitary vector bundle as above. Consider any other given data of: a finite factor B , a flat B -bundle $\mathcal{F} \rightarrow M$ along with an isomorphism $\theta: E^- \otimes \mathcal{F} \rightarrow E^+ \otimes \mathcal{F}$ (not required to preserve connections), and a trivialization $\psi: \mathcal{F} \rightarrow C(M) \otimes B$, and denote with $CS(\theta^* \nabla_{E^+ \otimes \mathcal{F}}; \nabla_{E^- \otimes \mathcal{F}})$ the Chern–Simons transgression form (formula 3.9). Let $w_\theta = (1_{E^+} \otimes \psi)^{-1} \circ \theta \circ (1_{E^-} \otimes \psi): E^- \otimes B \rightarrow E^+ \otimes B$. We have

$$[(E^+, E^-, w_\theta^{-1})] + j_*(\text{Ch}_\tau^{-1}(CS(\theta^* \nabla_{E^- \otimes \mathcal{F}}; \nabla_{E^+ \otimes \mathcal{F}}))) = [\alpha]_{new} \in K^1(M, \mathbb{R}/\mathbb{Z}).$$

Here j_* denotes the canonical morphism $K^1(M, \mathbb{R}) \rightarrow K^1(M, \mathbb{R}/\mathbb{Z})$.

Proof: Up to tensoring \mathcal{F} by \mathcal{E} given by proposition 5.2, we may assume that there is a flat connexion preserving isomorphism $v: E^+ \otimes \mathcal{F} \rightarrow E^- \otimes \mathcal{F}$.

Let now B and $\mathcal{F} \rightarrow M$ be fixed, and $\theta: E^- \otimes \mathcal{F} \rightarrow E^+ \otimes \mathcal{F}$ and $v: E^+ \otimes \mathcal{F} \rightarrow E^- \otimes \mathcal{F}$ two isomorphisms, such that v is connexion preserving.

Put $\ell = v \circ \theta$; it is an automorphism of $E^- \otimes \mathcal{F}$. As $v^*(\nabla_{E^- \otimes \mathcal{F}}) = \nabla_{E^+ \otimes \mathcal{F}}$, we find (using prop. 3.11)

$$CS(\theta^* \nabla_{E^+ \otimes \mathcal{F}}; \nabla_{E^- \otimes \mathcal{F}}) = CS(\ell^* \nabla_{E^- \otimes \mathcal{F}}; \nabla_{E^- \otimes \mathcal{F}}) = \text{Ch}(\ell).$$

Moreover $[(E^+, E^-, w_v)] - [(E^+, E^-, w_\theta^{-1})] = [(E^-, E^-, w_v \circ w_\theta)]$ is the image by $j: K_1(C(M); \mathbb{R}) \rightarrow K_1(C(M); \mathbb{R}/\mathbb{Z})$ of the class of the automorphism $w_v \circ w_\theta = (1_{-} \otimes \varphi)^{-1} \circ \ell \circ (1_{-} \otimes \varphi)$ of $E^- \otimes B$. This is the same class as the automorphism ℓ of $E^- \otimes \mathcal{F}$. □

The element $[\alpha]_{APS}$ is constructed in [APS3, 5] by means of an isomorphism $\theta : E^- \otimes B \rightarrow E^+ \otimes B$, where $B = M_n(\mathbb{C})$ (for a suitable n): it is defined by $[\alpha]_{APS} = [(E^+, E^-, w_\theta^{-1})] + j_*(\text{Ch}_\tau^{-1}(\text{CS}(\theta^* \nabla_{E^- \otimes B}; \nabla_{E^+ \otimes B}))$). We immediately find:

Corollary 5.5 *For every group morphism $\alpha : \Gamma \rightarrow U_n$, the element $[\alpha]_{new}$ coincides with $[\alpha]_{APS}$. □*

6. The pairing with $[D] \in K_1(M)$ and the spectral flow description

Let $[\alpha] = [(E^+, E^-, w_v)] \in K_0(C_{i_0} \otimes C(M))$ of Section 5, where C_{i_0} is the mapping cone of the unital inclusion $i_0 : \mathbb{C} \rightarrow B$ - here B is a II_1 factor. In order to stay within unital algebras, we represent $[\alpha]$ in the K -theory of the double cylinder $Z_{i_0} \otimes C(M)$, as we saw in (2.2). It is the image in the summand $K_0(C_{i_0} \otimes C(M))$ of the $Z_{i_0} \otimes C(M)$ -module

$$F_{w_v} = \{f \in C([0, 1], E^+ \otimes B); f(0) \in E^+ \otimes 1 \text{ and } w_v f(1) \in E^- \otimes 1\}.$$

The Kasparov product gives the natural pairing between the class $[\alpha] \in K_0(C_{i_0} \otimes C(M))$ and a K -homology class $[D] \in K_1(M)$. In this section we describe and interpret it. In other words, we interpret the map $K_1(M) = KK^1(C(M), \mathbb{C}) \rightarrow \mathbb{R}/\mathbb{Z}$ induced by $[\alpha]$ via the pairing

$$KK^0(\mathbb{C}, C(M) \otimes Z_{i_0}) \times KK^1(C(M), \mathbb{C}) \longrightarrow KK^1(\mathbb{C}, Z_{i_0}) = K_1(Z_{i_0}) = \mathbb{R}/\mathbb{Z}. \tag{6.1}$$

6.1. Various notions of connections

Let us start with a class $[D] \in K_1(M)$ represented by the unbounded Kasparov module (H, D) , where D is a first order self-adjoint elliptic (classical) pseudodifferential operator acting on the sections of a Hermitian vector bundle $S \rightarrow M$, and $H = L^2(M, S)$.

A particularly important case for D in our discussion is the case of a Dirac type operator, *i.e.* an operator constructed in the following way:

- We assume that M is endowed with a Riemannian metric,
- The bundle S is a *Clifford bundle* *i.e.* it is endowed with a linear bundle map $\text{cliff} : T^*M \rightarrow \text{End}(S)$ such that for every $\xi \in T_x^*M$ then $\text{cliff}(\xi) \in \text{End}(S_x)$ is skew-adjoint and satisfies $\text{cliff}(\xi)^2 = -\|\xi\|^2 1_{S_x}$; denote by $c : S \otimes T^*M \rightarrow S$ the corresponding map.

- The bundle S is further endowed with a *metric Clifford connection* ∇_S in the sense of [BGV, Def. 3.39].
- The corresponding Dirac operator is given by $D(s) = c(\nabla_S(s))$ for any smooth section s of S .

Let us distinguish different types of connections that arise in our discussion:

Definition 6.1 Let A be a unital C^* -algebra, E a finitely generated projective module endowed with a compatible Hilbert A -module structure. Let $V \rightarrow M$ be a smooth bundle with fibers E .

- If D is a first order self-adjoint elliptic (classical) pseudodifferential operator acting on the sections of a hermitian vector bundle $S \rightarrow M$, a *pseudodifferential connection* for D is a selfadjoint pseudodifferential A -linear operator D_V (in the sense of [MF]) acting on the sections of the bundle $V \otimes S$ whose principal symbol is $\sigma_{D_V} = 1_V \otimes \sigma_D$.
- If D is Dirac type, then a *Dirac type connection* for D is an operator of the form $(1_V \otimes c) \circ \nabla_{V \otimes S}$ where $\nabla_{V \otimes S}$ is a metric Clifford connection on the Clifford A -bundle $V \otimes S$.

Of course, if D is Dirac type, then a Dirac type connection for D is a particular case of a pseudodifferential connection.

6.2. The pairing as an element of $K_1(Z_{i_0})$

The product (6.1) is computed by any choice of a D -connection G in the sense of [CS, Ku] on the Hilbert Z_{i_0} -module $V = F_{w_v} \otimes_{C(M) \otimes Z_{i_0}} (L^2(M, S) \otimes Z_{i_0})$. Note that a pseudodifferential connection for D is a D -connection in the sense of [CS, Ku] (the converse is of course not true).

We have

$$[\alpha] \otimes_{C(M)} [D] = [(V, G)] \tag{6.2}$$

where we can take any G as follows. Note first that $w_v : E^+ \otimes B \rightarrow E^- \otimes B$ extends to a morphism of Hilbert B -modules, $\tilde{w}_v = w_v \otimes 1 : L^2(M; E^+ \otimes S \otimes B) \rightarrow L^2(M; E^- \otimes S \otimes B)$ since $L^2(M; E^\pm \otimes S \otimes B)$ under the obvious isomorphism to $(E^\pm \otimes B) \otimes_{C(M) \otimes B} L^2(M; S \otimes B)$.

The connection will be of the form $G = (G_t)$, i.e. $G(f)(t) = G_t(f_t)$. Since the bundle E^- is trivial, we may set $G_1 = \tilde{w}_v^{-1}(1_{\mathbb{C}^n} \otimes D \otimes 1_B)\tilde{w}_v$. To construct G_0 , we may just take any selfadjoint pseudodifferential operator D_{E^+} on $E^+ \otimes S$ whose principal symbol is $1_{E^+} \otimes \sigma_D$; we may also assume that D is local enough

(in particular if D is differential), so that there is a canonical choice for D_{E^+} , called the *twisted tensor product* of D with the flat bundle E^+ .

Put then $G_0 = D_{E^+} \otimes 1_B$. The operators G_0 and G_1 have the same principal symbol and therefore $G_1 - G_0$ is bounded. We may then choose any strongly continuous bounded path q_t of selfadjoint operators acting on $E^+ \otimes S \otimes B$ such that $q_0 = 0$, $q_1 = G_1 - G_0$ and set $G_t = G_0 + q_t$. In particular, we may define $G_t = (1-t)G_0 + tG_1$.

6.3. The pairing as an element of \mathbb{R}/\mathbb{Z}

Let us first recall a few facts about index in the type II setting:

Let B be a factor of type II_1 . Denote by τ the normalized trace on B . Let $N = B \overline{\otimes} \mathcal{L}(H)$ the associated factor of type II_∞ endowed with the trace $\text{Tr}_N = \tau \otimes \text{Tr}$ where Tr is the canonical trace on $\mathcal{L}(H)$. Let $p = 1 \otimes p_0 \in N$, where p_0 is a minimal projection in $\mathcal{L}(H)$ and identify B with pNp .

Let $P_\pm \in N$ be two projections such that $P_+ - P_-$ is in the ideal \mathcal{K}_N of generalized compact operators in N . Then $\text{ind}(P_+, P_-)$ is the index of $q : E^+ \rightarrow E^-$ where $E^\pm = P_\pm Np$ considered a Hilbert B -module and $qx = P_-x$ for $x \in E^+$. Note that q is B -Fredholm with quasi-inverse $x \mapsto P_+x$. If $P_+ - P_-$ is in the domain of Tr_N , then $\text{ind}(P_+, P_-) = \text{Tr}_N(P_+ - P_-)$. Also note that there is a path P_t of projections in N such that $P_+ - P_t$ is norm continuous and with values in \mathcal{K}_N with $P_0 = P_-$ and the projections P_+, P_1 are comparable *i.e.* $P_+ \geq P_1$, in which case the index is the class of the finite projection $P_+ - P_1$ or $P_+ \leq P_1$, in which case the $-\text{ind}(P_+, P_-)$ is the class of $P_1 - P_+$.

Let D_\pm be selfadjoint operators affiliated with N such that $D_+ - D_- \in N$ (which means that D_+ and D_- have same domain and there exists $b \in N$, with $b = b^*$ and $D_+ = D_- + b$). Note that for λ_\pm not in the spectrum of D_\pm , we have

$$(D_- - \lambda_-)^{-1} - (D_+ - \lambda_+)^{-1} = (D_+ - \lambda_+)^{-1}(b + \lambda_- - \lambda_+)(D_- - \lambda_-)^{-1},$$

therefore the resolvent of D_+ is in \mathcal{K}_N if and only if the resolvent of D_- is. Assume this is the case. For $a \in \mathbb{R}$, let χ_a be defined by

$$\chi_a(t) = 1 \text{ if } t \geq a \text{ and } \chi_a(t) = 0 \text{ if } t < a.$$

Then for all $a \in \mathbb{R}$, we have $\chi_a(D_+) - \chi_a(D_-) \in \mathcal{K}_N$ and we put

$$\text{sf}_a(D_+, D_-) = \text{ind}(\chi_a(D_+), \chi_a(D_-)).$$

Note that for all $a, b \in \mathbb{R}$, we have (by additivity of the index)

$$\text{sf}_a(D_+, D_-) - \text{sf}_b(D_+, D_-) = \text{Tr}_N((\chi_a - \chi_b)(D_+)) - \text{Tr}_N((\chi_a - \chi_b)(D_-)).$$

Let us now come to (V, G) as defined above. We easily find:

Proposition 6.2 *The class $j(\text{sf}_a(G_0, G_1)) \in \mathbb{R}/\mathbb{Z}$ is independent of $a \in \mathbb{R}$. The class $[\alpha] \otimes_{C(M)} [D]$ in \mathbb{R}/\mathbb{Z} is $j(\text{sf}_a(G_0, G_1))$.*

Proof: Since $G_0 = D_{E^+} \otimes 1_B$ and $G_1 = \tilde{w}_v^{-1}(1_{\mathbb{C}^n} \otimes D \otimes 1_B)\tilde{w}_v$, the spectra of G_0 and G_1 are discrete and the trace of all the spectral projections is integer. In other words, $\text{Tr}_N((\chi_a - \chi_b)(G_0)) \in \mathbb{Z}$ and $\text{Tr}_N((\chi_a - \chi_b)(G_1)) \in \mathbb{Z}$, and therefore $j(\text{sf}_a(G_0, G_1)) \in \mathbb{R}/\mathbb{Z}$ is independent of $a \in \mathbb{R}$.

To see that the class $[(V, G)]$ in \mathbb{R}/\mathbb{Z} is equal to $j(\text{sf}_a(G_0, G_1))$, note that:

1. Since $\text{Sp}(G_0) \cup \text{Sp}(G_1)$ is discrete, there is a smooth function $\varphi : \mathbb{R} \rightarrow [-1, +1]$ such that $\varphi = -1$ near $-\infty$, $\varphi = +1$ near $+\infty$, and $\varphi = 2\chi_a - 1$ in $\text{Sp}(G_0) \cup \text{Sp}(G_1)$. The class of the unbounded Kasparov module (V, G) is that of $(V, \varphi(G))$.
2. If $\text{sf}_a(G_0, G_1) = 0$, one can find a norm continuous path $P = (P_t)$ of projections in $\mathcal{L}(V)$ such that $t \mapsto P_t - P_0$ is a norm continuous path with values in $\mathcal{K}((L^2(M; S \otimes E^+) \otimes B))$, and such that $P_0 = \chi_a(G_0)$ and $P_1 = \chi_a(G_1)$: the class of $(V, \varphi(G))$ coincides with the class of $(V, 2P - 1)$ which is degenerate since $(2P - 1)^2 = 1$.
3. For $\lambda \in \mathbb{R}_+^*$, let E_λ be a finitely generated projective B -module with trace λ and denote $V_\lambda = C_0(]0, 1[, E_\lambda)$. Let $F_\lambda \in \mathcal{L}(V_\lambda)$ be given by $(F_\lambda \xi)(t) = (2t - 1)\xi(t)$. Then (V_λ, F_λ) is the product of the (opposite of the) Bott element in $K_1(C_0(]0, 1[) \otimes B)$ by E_λ , and its class in $K_1(C_0(]0, 1[) \otimes B) = \mathbb{R}$ is therefore equal to $-\lambda$.
4. If $\text{sf}_a(G_0, G_1) = \lambda > 0$, then the class of $(V, \varphi(G)) \oplus (V_\lambda, F_\lambda)$ is 0 by (2), hence the class of $(V, \varphi(G))$ is $j(\lambda)$. This is also true if $\text{sf}_a(G_0, G_1) < 0$ (for instance by replacing G by $-G$).

□

6.4. The pairing as a rho invariant

We will now explain how our approach can be used to establish the formula of Atiyah–Patodi–Singer in [APS3, Sec.5].

We first recall the facts about the η invariant that will be used here. A very nice survey can be found in the first chapter and appendix D of [Bo].

1. If P is an elliptic self-adjoint pseudodifferential operator of positive order, the *eta function* $s \mapsto \eta(P, s) = \text{Tr}(P|P|^{-1-s})$ has a meromorphic continuation which is regular at 0 (cf. [APS3, Thm. 4.5], [Gi2]). Put $\xi(P) = \frac{1}{2}(\eta(P) + \dimker P)$.

2. If P is differential of Dirac type, then $\eta(P) = \frac{1}{\sqrt{\pi}} \int_0^{+\infty} \text{Tr}(P e^{-tP^2}) \frac{dt}{\sqrt{t}}$ (cf [BF, Thm 2.6]).
3. The above remain true for P with coefficients in a finite von Neumann algebra B ([CG, Ram, CP1, LP, An]).
4. For P with coefficients in B , put $\xi_\varepsilon(P) = \frac{1}{2\sqrt{\pi}} \int_\varepsilon^{+\infty} \text{Tr}(P e^{-tP^2}) \frac{dt}{\sqrt{t}} + \frac{1}{2} \dim_B \ker P$. If P_1, P_0 are bounded perturbations of each other, then for any $\varepsilon > 0$, and any smooth path P_t joining P_0 with P_1 (i.e. a path of the form $t \mapsto P_0 + Q_t$ where $t \mapsto Q_t$ is smooth) we have ([CP1, Cor. 8.10])

$$\text{sf}_0(P_0, P_1) = \xi_\varepsilon(P_0) - \xi_\varepsilon(P_1) + \sqrt{\frac{\varepsilon}{\pi}} \int_0^1 \text{Tr}_N(\dot{P}_t e^{-\varepsilon P_t^2}) dt.$$

(where N is the II_∞ factor associated with B).

5. If $V \rightarrow M$ is a von Neumann bundle and P_t is a smooth family of Dirac type operators then $\sqrt{\frac{\varepsilon}{\pi}} \int_0^1 \text{Tr}_N(\dot{P}_t e^{-\varepsilon P_t^2}) dt$ converges, when $\varepsilon \rightarrow 0$, to a local term,

$$\int_M \hat{A}(M) \text{ch} S / \hat{S} \cdot \text{CS}(\nabla_0, \nabla_1)$$

see [APS2, Sec. 4], [Bo, 1.5.1].

Let $\alpha: \Gamma \rightarrow U_n$ be a morphism, denote by $E^+ = E_\alpha$ the corresponding flat vector bundle over M and E^- the trivial bundle of dimension n . Let D be an elliptic differential operator. Recall that the rho invariant of α is $\rho(\alpha, D) = \xi(D^\alpha) - n\xi(D)$.

Proposition 6.3 [APS3, Sec.5]: *If D is a Dirac type operator, then $[\alpha] \otimes_{C(M)} [D]$ is the class modulo \mathbb{Z} of $\rho(\alpha, D)$.*

Proof: If P and P' are two Dirac type operators acting on a von Neumann Clifford bundle, we put

$$\theta(P, P') = \text{sf}_0(P, P') - \xi(P) + \xi(P') = \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{\varepsilon}{\pi}} \int_0^1 \text{Tr}(\dot{P}_t e^{-\varepsilon P_t^2}) dt$$

for any smooth path P_t joining P with P' .

Fix a II_1 factor B , a flat bundle \mathcal{E} with fibers B over M , an isomorphism $\varphi: C(M) \otimes B \rightarrow \mathcal{E}$ and an isomorphism $v: E^+ \otimes_{C(M)} \mathcal{E} \rightarrow E^- \otimes_{C(M)}$ of flat bundles.

We have isomorphisms (not preserving the flat bundle structure) of the three flat bundles $E^+ \otimes_{\mathbb{C}} B$, $E^+ \otimes_{\mathbb{C}(M)} \mathcal{E} \sim_v E^- \otimes_{\mathbb{C}(M)} \mathcal{E}$ and $E^- \otimes_{\mathbb{C}} B$ given by

$$E^{\pm} \otimes B \simeq E^{\pm} \otimes_{\mathbb{C}(M)} \left(C(M) \otimes B \right) \sim_{1_{E^{\pm}} \otimes \varphi} E^{\pm} \otimes_{\mathbb{C}(M)} \mathcal{E}.$$

In other words, on the bundle $E^+ \otimes B$, we have three flat bundle structures. We thus get three Dirac type operators acting on $E^+ \otimes S \otimes B$ as twisted tensor products of D with these flat bundle structures

$$G_0 = D^{\alpha} \otimes 1_B, \quad \tilde{G} \quad \text{and} \quad G_1 = \tilde{w}_v(1_{\mathbb{C}^n} \otimes D \otimes 1_B) \tilde{w}_v^{-1}.$$

Consider the Dirac operators $P = D \otimes 1_B$ and $P' = (1_S \otimes \varphi) \tilde{D} (1_S \otimes \varphi^{-1})$ acting on $S \otimes B$.

We have $G_1 = \tilde{w}_v(1_{\mathbb{C}^n} \otimes P) \tilde{w}_v^{-1}$ and $\tilde{G} = \tilde{w}_v(1_{\mathbb{C}^n} \otimes P') \tilde{w}_v^{-1}$, therefore $\theta(G_1, \tilde{G}) = n\theta(P, P')$.

By the locality property (5) of θ , and since G_0 is the twisted tensor product of P with the flat bundle E^+ and \tilde{G} is the twisted tensor product of P' with the flat bundle E^+ , we find $\theta(G_0, \tilde{G}) = n\theta(P, P')$. And finally $\theta(G_0, G_1) = 0$. \square

REFERENCES

- An. P. Antonini, *The Atiyah Patodi Singer signature formula for measured foliations*, Ph.D Thesis, La Sapienza, Rome <http://arxiv.org/abs/0901.0143>.
- Ati. M.F. Atiyah, Elliptic operators discrete groups and von Neumann algebras. *Astérisque* **32–33** (1976), 43–72.
- APS1. M.F. Atiyah, V.K. Patodi, I.M. Singer. Spectral asymmetry and Riemannian geometry. I. *Math. Proc. Cambridge Philos. Soc.* **77** (1975), 43–49.
- APS2. ———, Spectral asymmetry and Riemannian geometry. II. *Math. Proc. Cambridge Philos. Soc.* **78** (1975), 405–432.
- APS3. ———, Spectral asymmetry and Riemannian geometry. III. *Math. Proc. Cambridge Philos. Soc.* **79** (1976), 71–99.
- Ba. D. Basu, K-theory with \mathbb{R}/\mathbb{Z} coefficients and von Neumann algebras, *K-Theory* **36**(3-4) (2005), 327–343.
- BGV. N. Berline, E. Getzler, M. Vergne, *Heat kernels and Dirac operators*, Springer-Verlag, New York 1992.
- BF. J.M. Bismut and D.S. Freed, The analysis of elliptic families II: Dirac operators, eta invariants, and the holonomy theorem of Witten, *Commun. Math. Phys.* **107** (1986), 103–163.

- Bl. B. Blackadar, *K-theory for operator algebras*, second edition, MSRI publications, Cambridge University Press.
- Bl1. ———, *Operator algebras: theory of C^* -Algebras and non Neumann algebras*, Encyclopaedia on operator algebras and non-commutative geometry **13**, Springer 2006.
- Bo. M. Bohn, On Rho invariants of fibre bundles, <http://arxiv.org/abs/0907.3530> [math.GT].
- CP1. A. Carey, J. Phillips, Spectral flow in Fredholm modules, eta invariants and the JLO cocycle, *K-Theory* **31**(2) (2004), 135–194.
- CG. J. Cheeger, M. Gromov, On the characteristic numbers of complete manifolds of bounded curvature and finite volume, in *Differential geometry and complex analysis*, Springer, Berlin 1985, 115–154.
- CM. A. Connes, H. Moscovici, Cyclic cohomology, the Novikov conjecture and hyperbolic groups, *Topology* **29**(3) (1990), 345–388.
- CS. A. Connes, G. Skandalis, The longitudinal index theorem for foliations, *Publ. Res. Inst. Math. Sci.* **20**(6) (1984).
- De. R.J. Deeley, \mathbb{R}/\mathbb{Z} -valued index theory via geometric K -homology, <http://arxiv.org/abs/1206.5662> [math.KT].
- DHK. R.G. Douglas, S. Hurder, J. Kaminker, Eta invariants and von Neumann algebras, *Bull. Amer. Math. Soc* **21**(1) (1989).
- DHK2. R.G. Douglas, S. Hurder, J. Kaminker, Cyclic cocycles, renormalization and eta-invariants, *Inv. Math.* **103**(1) (1991), 101–180.
- G. E. Getzler, The odd Chern character in cyclic homology and spectral flow, *Topology* **32**(3) (1993), 489–507.
- Gi2. P. B. Gilkey, The residue of the global eta function at the origin, *Adv. Math.* **40** (1981), 290–307.
- Gi. P. B. Gilkey, The eta invariant and secondary characteristic classes of locally flat bundles, in *Algebraic and differential topology – global differential geometry*, Teubner–Texte Math. **70**, Leipzig (1984), 49–87.
- Gr. M. Gromov, Kähler hyperbolicity and L^2 -Hodge theory, *J. Diff. Geom.* **33** (1991), 263–292.
- Hu. S. Hurder, Eta invariants and the odd index theorem for coverings. Geometric and topological invariants of elliptic operators, *Contemp. Math.* **105**, Amer. Math. Soc., Providence, RI, (1990), 47–82.
- Ka. M. Karoubi, Homologie Cyclique et K -Théorie, Astérisque **149** (1987).
- Ka2. M. Karoubi, *K-theory. An introduction*. Grundlehren der Mathematischen Wissenschaften **226**. Springer-Verlag, Berlin–New York, 1978.

- Ka3. M. Karoubi, Théorie générale des classes caractéristiques secondaires. *K-Theory* **4** (1990), 55–87.
- Ka4. M. Karoubi, Classes caractéristiques de fibrés feuilletés, holomorphes ou algébriques. *K-Theory* **8** (1994), 153–211.
- KP. J. Kaminker, V. Perera, Type II Spectral Flow and the Eta Invariant, *Canad. Math. Bull.* **43**(1) (2000), 69–73.
- Ku. D. Kucerovsky, The KK-product of unbounded modules, *K-Theory* **11**(1) (1997), 17–34.
- LP. E. Leichtnam, P. Piazza, Cut-and-Paste on foliated bundles, *Contemp. Mathem.* **366** (2005), 351–392 “Spectral Geometry of Manifolds with Boundary”, Amer. Math. Soc., Providence, RI.
- Lo. J. Lott, \mathbb{R}/\mathbb{Z} -index theory, *Comm. Anal. Geom.* **2**(2) (1994), 279–311.
- Lü. W. Lück, The relation between the Baum–Connes Conjecture and the Trace Conjecture, *Invent. Math.* **149** (2002), 123–152
- MZ1. X. Ma, W. Zhang, η -Invariant and flat vector bundles, *Chin. Ann. Math* **27B**(1) (2006), 57–72.
- MZ2. ———, η -Invariant and flat vector bundles II, in “Inspired by S. S. Chern. A memorial volume in honor of a great mathematician.” Hackensack, NJ: World Scientific. *Nankai Tracts in Mathematics* **11** (2006), 335–350.
- Mi. A. S. Miščenko, The theory of elliptic operators over C^* -algebras, *Dokl. Akad. Nauk SSSR* **239**(6) (1978), 1289–1291.
- MF. A. S. Miščenko, A. T. Fomenko, The index of elliptic operators over C^* -algebras. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* **43**(4) (1979), 831–859, 967.
- MN. I. Moriyoshi, T. Natsume, The Godbillon-Vey cyclic cocycle and longitudinal Dirac operators, *Pacific J. of Math.* **172**(2) (1996), 483–539.
- PS. P. Piazza, T. Schick, Bordism, rho-invariants and the Baum-Connes conjecture. *J. of Noncommutative Geometry* **1** (2007), 27–111.
- Pu. I. F. Putnam, An excision theorem for the K -theory of C^* algebras, *J. Operator Theory* **38** (1997), 151–171.
- Ra. I. Raeburn, K -theory and K -homology relative to a II_∞ -factor. *Proc. Amer. Math. Soc.* **71**(2) (1978), 294–298.
- Ram. M. Ramachandran, Von Neumann index theorems for manifolds with boundary, *J. Diff. Geom.* **38** (1993), 315–349.
- Sc. T. Schick, L^2 -index theorems, KK-theory, and connections, *New York J. Math.* **11** (2005), 387–443.
- Si. I.M. Singer, Some remarks on operator theory and index theory, in “ K -Theory and Operator Algebras”, *Lect. Notes Math.* **575** (1977), 128–138.

- Va. B. Vaillant, Index theory for coverings, <http://arxiv.org/abs/0806.4043>.
- Wo. K. Wojciechowski, A note on the space of pseudodifferential projections with the same principal symbol, *J. Operator Theory* **15**(2) (1986), 207–216.

PAOLO ANTONINI paolo.anton@gmail.com

Institut de Mathématiques de Jussieu, (UMR 7586), Université Paris Diderot (Paris 7)
UFR de Mathématiques, CP **7012** - Bâtiment Sophie Germain
5 rue Thomas Mann, 75205 Paris CEDEX 13, France

SARA AZZALI azzali@math.jussieu.fr

Institut de Mathématiques de Jussieu, (UMR 7586), Université Paris Diderot (Paris 7)
UFR de Mathématiques, CP **7012** - Bâtiment Sophie Germain
5 rue Thomas Mann, 75205 Paris CEDEX 13, France

GEORGES SKANDALIS skandalis@math.univ-paris-diderot.fr

Institut de Mathématiques de Jussieu, (UMR 7586), Université Paris Diderot (Paris 7)
UFR de Mathématiques, CP **7012** - Bâtiment Sophie Germain
5 rue Thomas Mann, 75205 Paris CEDEX 13, France

Received: October 11, 2013