



École doctorale de sciences mathématiques de Paris centre

THÈSE DE DOCTORAT

Discipline : Mathématiques

présentée par

Haohao LIU

April 30, 2024

**Integral points, monodromy, generic
vanishing and Fourier-Mukai transform**

dirigée par Anna CADORET

Institut de mathématiques de
Jussieu-Paris Rive gauche. UMR
7586.
Boîte courrier 247
4 place Jussieu
75 252 Paris Cedex 05

祝我的妈妈六十生日快乐！

Acknowledgments

First, I would like to thank my supervisor Anna Cadoret. It was her constant encouragement that has made this thesis produced. She is always tolerant, patient and warm-hearted. I must thank her for carefully reading the manuscript multiple times, pointing out various mistakes and making numerous constructive suggestions that have substantially improved the exposition. Many times, she indicated me how to strengthen the results by removing unnecessary conditions.

There are many professors helped me during the journey. I am grateful to Olivier Benoist, Nicolas Bergeron, Yann Brenier, Hongjia Chen, Yongqi Liang, Xinan Ma, Xiaonan Ma, Ariane Mézard, Yi Ouyang, Mao Sheng, Benoît Stroh and Zuoqin Wang. It was Professor Liang who supported me to attend a summer school on algebra and number theory, with his own funding (even if we haven't met in person till today). I benefited a lot from enlightening communication with Owen Barrett, Oren Ben-Bassat, Jonathan Block, Sébastien Boucksom, Christopher Daw, Julien Grivaux, Jack Hall, Luc Illusie, Kai-Wen Lan, Yongqiang Liu, James S. Milne, Amnon Neeman and Andrei Yafaev.

Example 3.2.2.3 was learned from Owen Barrett. I also owe a lot to Yves André, Arthur Forey, Franc Forstnerič, Robin Hartshorne, Daniel Huybrechts, Masaki Kashiwara, Joseph Lipman, Claude Sabbah, Pierre Schapira, Christian Schnell, Toshiyuki Tanisaki, Jean-Baptiste Teyssier, John Voight and Rainer Weissauer. I appreciate the patience and detailed replies of Will Sawin and Marco Maculan to my questions on their respective work. Proposition 6.7.5.4 and its proof, both due to Claude Sabbah, are explained to me by Gabriel Ribeiro. The example in Remark 3.5.3.4 is also due to Will Sawin. I thank Emiliano Ambrosi for his hospitality during my visit to the Université de Strasbourg. Peter Scholze kindly answered my question, to whom Lemma 3.2.3.11 is due. My gratitude goes to Botong Wang for answering my question on his work. I thank Lie Fu for raising the interesting Question B.5.0.2.

I express my deep gratitude to Thomas Krämer for his hospitality during my visit to the Humboldt-Universität zu Berlin. I must thank Akio Tamagawa and Benjamin Collas, for giving me a precious opportunity to visit Kyoto University and for their generous help, during and beyond my stay. The

kindness and friendliness of all the secretaries of Kokusai, RIMS are sincerely appreciated.

Numerous friends and colleagues have helped me. Stefano Alo , Jacques Audibert, Chenyu Bai, Adrien Cortes, Odylo Costa, Arnaud Eteve, Brian Hepler, Rein Janssen Groesbeek, Yueting Jiang, Arnab Kundu, Joseph Leclere, Joaqu n Lejtregger, Shang Li, Zhenghui Li, Tzu-Jan Li, Junbang Liu, Xiaolong Liu, Yachen Liu, Qiaochu Ma, Kai Mao, Nguy n M nh Linh, Yang Pei, Keyao Peng, Pietro Mesquita Piccione, Yichen Qin, Joaquim Ribeiro, Drimik Roy, Nelson Schuback, Xinyu Shao, Miao Song, Matteo Verni, Yitong Wang, Mingchen Xia, Junsheng Zhang, Ruishen Zhao, Xiaoxiang Zhou and Yicheng Zhou have kindly answered me quite a few questions. Remark 4.7.2.4 is due to Chenyu Bai. Andreas Hohl helped me with Remark 6.7.2.10. I thank Thiago Landim and Rapha l Prunier for allowing me to give a talk in the seminar they organized.

I am grateful to Long Liu for teaching me category theory and many things. Remark A.1.4.7 is told by him. I thank Gabriel Ribeiro for noting a mistake in a previous version, kindly sharing several useful notes and helping me with Lemma 6.7.5.2. In a draft version, Theorem 5.4.1.1 was stated for the categories $D_{\text{good}}^b(*)$. Gabriel Ribeiro suggested the present extension to $D_{\text{good}}(*)$, to whom I am very grateful for his suggestions and much help. I thank Alex Youcis for maintaining a comprehensive blog on arithmetic. Hui Zhang helped me out multiple times with his admirable knowledge, especially in algebraic geometry. Lemma 5.3.1.13 is told by him. Remark 5.2.0.13 is due to Yicheng Zhou, who explained many advanced concepts to me. All remaining errors are mine.

I am grateful to my girlfriend Lu Yu for her company and support. Finally, I thank my family for their infinite love.

Résumé

Cette thèse est une compilation de plusieurs résultats vaguement liés. Ils concernent la non-densité des points entiers sur les variétés algébriques, la méthode de Lawrence-Venkatesh-Sawin et la géométrie analytique complexe.

Dans Chapitre 2, parallèlement au principe alternatif d'Ullmo et Yafaev sur les points rationnels des variétés de Shimura, nous montrons que la conjecture de Lang sur les points intégraux des variétés de Shimura est soit vraie, soit très fausse.

Le Chapitre 3 est un complément à la comparaison des monodromies dans les travaux respectifs de Lawrence-Sawin et Krämer-Maculan. Nous prouvons qu'il existe de nombreux caractères, tels que le groupe de monodromie correspondant est normal dans le groupe tannakien générique.

Le Chapitre 4 contient un théorème de l'annulation générique pour les variétés dans la classe Fujiki \mathcal{C} . En particulier, cela s'applique aux variétés algébriques complexes propres lisses ainsi qu'aux variétés kählériennes compactes.

Dans Chapitre 5, nous prouvons un analogue de la formule d'inversion de Fourier pour la transformation de Fourier-Mukai sur des tores complexes. Il corrige une inexactitude dans la littérature. En application, nous retrouvons la classification de Matsushima-Morimoto des fibrés vectoriels homogènes sur des tores complexes.

Le Chapitre 6 est une transformation de Fourier-Mukai analytique sur les D -modules, dont la version algébrique a été étudiée par Laumon et Rothstein. Nous étendons leur résultat de dualité des variétés abéliennes aux tores complexes. En application, nous réprouvons le théorème de Morimoto, selon lequel sur un tore complexe, tout fibré vectoriel admettant une connexion admet une connexion intégrable.

Mots-clés

Conjecture de Lang, groupe de monodromie, annulation générique, transformation de Fourier-Mukai, D -module.

Abstract

This dissertation is a compilation of several loosely related results. They concern the nondensity of integral points on algebraic varieties, the Lawrence-Venkatesh-Sawin's method and complex analytic geometry.

In Chapter 2, parallel to Ullmo and Yafaev's alternative principle on rational points of Shimura varieties, we show that Lang's conjecture about integral points on Shimura varieties is either true or very false.

Chapter 3 is a complement to the monodromy comparison step in Lawrence-Sawin's and Krämer-Maculan's respective work. We prove that there are many characters, such that the corresponding monodromy group is normal in the generic Tannakian group.

Chapter 4 contains a generic vanishing theorem for Fujiki class \mathcal{C} . In particular, it applies to smooth proper complex algebraic varieties as well as compact Kähler manifolds.

In Chapter 5, we prove an analog of the Fourier inversion formula for the Fourier-Mukai transform on complex tori. It corrects a misstatement in the literature. As an application, we recover Matsushima-Morimoto's classification of homogeneous vector bundles on complex tori.

Chapter 6 is a lift of the analytic Fourier-Mukai to D -modules, whose algebraic version is studied by Laumon and Rothstein. We extend their duality result from abelian varieties to complex tori. As an application, we reprove Morimoto's theorem that on a complex torus, every vector bundle admitting a connection admits a flat connection.

Keywords

Lang conjecture, monodromy group, generic vanishing, Fourier-Mukai transform, D -module.

Contents

1	Introduction	13
1.1	Rational points	13
1.1.1	Mordell conjecture	13
1.1.2	Lang conjectures	14
1.1.3	Lang conjecture for Shimura varieties	16
1.2	Lawrence-Venkatesh technique	17
1.2.1	Setting	17
1.2.2	Galois representations	18
1.2.3	p -adic Hodge theory	18
1.2.4	Complex period map	19
1.2.5	p -adic period map	19
1.2.6	Ax-Schanuel property of period map	20
1.2.7	Summary	20
1.3	Lawrence-Sawin technique	21
1.3.1	Tannakian theory of sheaf convolution	21
1.3.2	Monodromy group and generic Tannakian group	23
1.3.3	Normality of geometric generic Tannakian group	24
1.3.4	Monodromy group and geometric generic Tannakian group	24
1.3.5	Summary	24
1.3.6	Normality of monodromy group	25
1.4	Generic vanishing	26
1.5	Fourier-Mukai transform	28
1.5.1	Construction	28
1.5.2	Complex tori	29
1.5.3	Homogeneous vector bundles	30
1.6	Laumon-Rothstein transform	30
1.6.1	D -modules	30
1.6.2	Construction	31
1.6.3	Schnell's proof of Fact 1.4.0.1	32
1.6.4	Complex tori	33
1.6.5	Vector bundles with connection	34
1.7	Future directions	34

1.7.1	Six-functor formalism of analytic quasi-coherent sheaves	34
1.7.2	Analytic Krämer-Weissauer's vanishing theorem	34
1.7.3	Lawrence-Venkatesh's method	35
1.8	Overview	35
2	Integral points of Shimura varieties: an "all or nothing" principle	36
2.1	Introduction	36
2.2	Locus formation	37
2.3	Shimura varieties	41
2.4	Ullmo-Yafaev alternative principle	49
2.5	"All or nothing" principle for integral points	50
3	Normality of monodromy group in generic Tannakian group	55
3.1	Introduction	55
3.1.1	Background	55
3.1.2	Statements	56
3.2	Recollections on constructible sheaves	57
3.2.1	Basics	58
3.2.2	Universal local acyclicity	61
3.2.3	Relative perverse sheaves	64
3.3	Cotori	69
3.3.1	Definition and basic properties	69
3.3.2	Cotori are Baire	71
3.4	Krämer-Weissauer theory	75
3.4.1	Generic vanishing theorem	75
3.4.2	Tannakian groups	78
3.4.3	Sheaf convolution	79
3.5	Main results	80
3.5.1	Reductivity	80
3.5.2	Fixed part	82
3.5.3	Normality	83
4	Generic vanishing theorem for Fujiki class \mathcal{C}	90
4.1	Introduction	90
4.1.1	Known results	91
4.1.2	The main result and a sketch	93
4.2	Riemann-Hilbert correspondence	94
4.2.1	Unitary local systems	94
4.2.2	Flat unitary bundles	95
4.2.3	An equivalence	96
4.3	Hodge theory and Jacobian	97
4.3.1	Regular manifolds	97
4.3.2	Jacobian	100
4.4	Albanese torus	102

4.4.1	Basics of Albanese torus	102
4.4.2	Back to Jacobian	105
4.5	Defect of semismallness	107
4.5.1	Stratifications and constructible sheaves	107
4.5.2	Equivalent definitions	109
4.5.3	Direct image of local systems	111
4.6	Generic vanishing for constructible sheaves	114
4.6.1	Thin subsets	114
4.6.2	Generic vanishing result on regular manifolds	115
4.7	Generic vanishing result for manifolds in Fujiki class \mathcal{C}	117
4.7.1	Fujiki class \mathcal{C}	117
4.7.2	Moishezon manifolds	119
5	Fourier-Mukai transform on complex tori, revisited	123
5.1	Introduction	123
5.2	Fourier-Mukai transform	125
5.3	Good modules	130
5.3.1	Functoriality	130
5.3.2	Base change theorems	137
5.3.3	Compatibility	143
5.4	Analytic Mukai duality	144
5.4.1	Statement	144
5.4.2	Proof	145
5.5	Properties of Fourier-Mukai transform	148
5.5.1	Functoriality	149
5.5.2	Unipotent vector bundles	157
5.5.3	Homogeneous vector bundles	160
6	Sheaves with connection on complex tori	164
6.1	Introduction	164
6.1.1	Background	164
6.1.2	Extension to complex tori	167
6.2	Preliminaries	169
6.2.1	Categories of splittings	170
6.2.2	Categories of twisted connection	172
6.2.3	Functors between them	174
6.3	Rothstein transform on modules with connection	177
6.3.1	Construction	177
6.3.2	Rothstein's theorem	178
6.3.3	Matsushima's theorem	179
6.4	Laumon-Rothstein sheaf of algebras	181
6.4.1	Construction	181
6.4.2	Basic properties	183
6.5	Laumon-Rothstein transform	184

6.5.1	Construction and properties	184
6.5.2	Matsushima-Morimoto theorem	186
6.6	Good modules	187
6.6.1	Definition	187
6.6.2	Basic properties	191
6.6.3	Preservation of goodness	193
6.7	Relations with other functors	195
6.7.1	Exchange of translation and multiplication	195
6.7.2	Duality	195
6.7.3	Pullback and pushout	198
6.7.4	External tensor product	201
6.7.5	Convolution and tensor product	203
A	Sheaves of modules	209
A.1	Sheaves of modules	209
A.1.1	Generalities	209
A.1.2	Pseudo-coherent modules	210
A.1.3	Analytic coherent modules	212
A.1.4	Good modules	214
A.1.5	Sections of direct sum of sheaves	217
A.2	Gabber's example	219
B	Quasi-coherent GAGA	222
B.1	Introduction	222
B.2	Review	222
B.3	GAGA for quasi-coherent modules	224
B.4	Derived category of quasi-coherent sheaves	226
B.5	Compact objects	230
C	Quasi-coherent sheaves on complex analytic spaces	231
C.1	Introduction	231
C.2	Preliminaries	231
C.3	Proof of Theorem C.1.0.2	234
D	Complex analytic geometry	236
D.1	Dimension of the fiber product	236
D.1.1	Algebraic case	236
D.1.2	Analytic case	237
D.2	Connection on line bundles	239
D.3	Jacobi inversion theorem	242
D.3.1	First proof	242
D.3.2	Second proof	243

E	<i>D</i>-modules	245
E.1	Unbounded Bernstein's equivalence	245
E.2	When is an induced <i>D</i> -module holonomic?	251
F	Group extensions of complex Lie groups	254
F.1	Introduction	254
F.2	Generalities on complex Lie groups	256
F.3	Group extensions	259
	F.3.1 Pullback and pushout	261
	F.3.2 Rudimentary classification	263
F.4	Commutative Extensions	266
	F.4.1 Generalities	266
	F.4.2 Exact sequences of Ext	271
	F.4.3 Determination of commutative extension group	275
F.5	Commutative extensions of complex tori	277
	F.5.1 Primitive cohomology classes	277
	F.5.2 The case $B = \mathbb{C}^*$	280
	F.5.3 The case $B = \mathbb{C}$	282
	F.5.4 Universal vectorial extension	283
	F.5.5 Application to the functor $\text{Ext}(A, \bullet)$	288
F.6	Extensions of complex tori are often commutative	294
F.7	Noncommutative extensions	299
	F.7.1 Lifted extensions	299
	F.7.2 Factor systems	301
	F.7.3 Non-abelian kernels and extensions of the center	306
F.8	Maximal morphisms	311
F.9	Commutative extensions of real Lie groups	313

Chapter 1

Introduction

1.1 Rational points

Intuitively, given an algebraic variety over a number field, the complexity of its geometry affects how many rational points (over finite extensions of the base field) it can possess. In Chapter 1, by an algebraic variety, we mean a geometrically integral, finite type, separated scheme over a field. An algebraic variety of dimension 1 is called a curve.

1.1.1 Mordell conjecture

Let K be a number field. Let S be a finite subset of places of K containing all infinite ones. By Siegel's theorem [Sie29, p.252], on the projective line P_K^1 with at least three punctures, or on a genus 1 curve over K with at least one puncture, there are at most finitely many $O_{K,S}$ -integral points. For curves of higher genus, Faltings's theorem (Fact 1.1.1.1) was conjectured by Mordell [Mor22, (5), p.192].

Fact 1.1.1.1 (Faltings, [Fal83, Satz 7]). *Let Y be a smooth projective curve over K of genus ≥ 2 . Then $Y(K)$ is finite.*

A sample application of Faltings's theorem is a partial solution to Fermat's Last Theorem: for every integer $n \geq 4$, there are only finitely many pairwise coprime integer solutions to the equation $x^n + y^n = z^n$. Indeed, the projective plane curve (known as the n -th Fermat curve) in $P_{\mathbb{Q}}^2$ cut out by this equation has genus $(n-1)(n-2)/2 (\geq 2)$. It is more than a decade earlier than Andrew Wiles's complete solution in 1994 to Fermat's conjecture.

Parshin [Par68] constructed a family of curves over Y , with which he showed that Fact 1.1.1.1 is a consequence of Shafarevich's conjecture for curves. This conjecture in turn follows from Shafarevich's conjecture for abelian varieties and Torelli's theorem.

We recall the statement of Shafarevich's conjecture. A smooth proper variety (resp. abelian variety) over a discrete valuation field E is said to

have good reduction if it is isomorphic to the generic fiber of a smooth proper scheme (resp. abelian scheme) over the integer ring O_E of E . By [Mil20, Prop. 6.4], there is at most one such abelian scheme. A smooth proper variety (resp. abelian variety) over K is said to have good reduction at a finite place v of the number field K if its base change to K_v has good reduction.

Fact 1.1.1.2 (Shafarevich conjecture, [Fal83, Korollar 1, p.365 (resp. Satz 6)]). *For every integer g at least 2 (resp. 1), up to K -isomorphism there are only finitely many smooth projective curves (resp. abelian varieties) defined over K of genus (resp. dimension) g , with good reduction outside S .*

In 1983, Faltings proved Shafarevich's conjecture for abelian varieties and hence Mordell's conjecture, "opening thereby a new chapter in number theory".¹ Faltings's proof can be decomposed into two parts, Facts 1.1.1.3 and 1.1.1.4.

Fact 1.1.1.3 ([Fal83, Satz 5]). *For every integer $g > 0$, up to K -isogeny there are only finitely many abelian varieties over K of dimension g , with good reduction outside S .*

Fact 1.1.1.3 is weaker than Shafarevich's conjecture for abelian varieties. Its proof is to consider the representations of the absolute Galois group Γ_K of K on the Tate modules of K -abelian varieties. For one thing, by Tate's conjecture over number fields [Fal83, Korollar 2], the Galois representation on the Tate module determines the abelian variety up to K -isogeny. For another, by Weil's conjecture proved by Deligne [Del74, Thm. 1.6], there are only finitely many such representations up to isomorphism.

Fact 1.1.1.4. *Let A be an abelian variety over K . Then up to K -isomorphism, there are only finitely many abelian varieties over K which are K -isogenous to A .*

Faltings introduced a differential height function, now known as Faltings's height, to measure the "complexity" of abelian varieties. Height function is a tool of global nature, as it collects the information at every place of the base number field. The core of the proof of Fact 1.1.1.4 is that Faltings's height does not change much under isogeny ([Fal83, Lem. 5]).

1.1.2 Lang conjectures

"One natural generalization to higher dimensions of the notion of 'curve of geometric genus $g \geq 2$ ' is 'variety of general type'." ([CHM97, p.2]). For a smooth projective variety X over a field, let ω_X be its canonical line bundle. For an integer $d \geq 0$, let $P_d(X) = h^0(X, \omega_X^{\otimes d})$ be the d -th plurigenus of

¹quotation from [Blo84, p.41]

X . The *Kodaira dimension* $\kappa(X)$ is defined to be $-\infty$ (or -1 depending on the convention) if $P_d(X) = 0$ for every integer $d > 0$; otherwise, it is the minimum real number r such that the sequence $\{P_d(X)/d^r\}_{d>0}$ is bounded. Then Kodaira dimension is the “most basic” ([Laz04, Eg. 2.1.5]) integer birational invariant of X . If $\kappa(X) = \dim X$, then X is called of *general type*. For instance, a smooth projective curve is of general type if and only if its genus is at least 2.

A high-dimensional analog of Fact 1.1.1.1 is conjectured by Lang (see, e.g., [CHM97, Conjecture A]).

Conjecture 1.1.2.1. *Let X be a positive dimensional smooth projective variety of general type over a number field K . Then $X(K)$ is not Zariski dense in X .*

Using techniques from Diophantine approximation, Faltings proves Conjecture 1.1.2.1 for subvarieties² of abelian varieties, which gives a second proof of Fact 1.1.1.1. From [Hin98, p.95], a subvariety of an abelian variety is of general type if and only if its stabilizer is finite.

Fact 1.1.2.2 ([Fal91, Thm. 1]). *Let A be an abelian variety over a number field K . Let $X \subset A$ be a subvariety of general type. Then $X(K)$ is finite.*

Based on Faltings’s work [Fal94], Moriwaki proves another particular case of Conjecture 1.1.2.1. A smooth projective variety with ample cotangent bundle is of general type.

Fact 1.1.2.3 ([Mor95, p.114]). *Let X be a smooth projective variety over a number field K . If the cotangent bundle $\Omega_{X/K}^1$ is ample and generated by global sections, then $X(K)$ is finite.*

Conjecture 1.1.2.1 is stronger than the uniformity conjecture.

Fact 1.1.2.4 ([CHM97, Thm. 1.1]). *Assume Conjecture 1.1.2.1 over every number field. Then for every number L and every integer $g \geq 2$, there is an integer $B(L, g)$ such that every smooth curve C over L of genus g , one has $\#C(L) \leq B(L, g)$.*

Conjecture 1.1.2.1 for algebraic surfaces was independently raised by Bombieri, so also known as the Bombieri-Lang conjecture. It gives a conditional solution to the Erdős-Ulam problem.

A rational distance set in \mathbb{R}^2 is a subset such that every pairwise distance between its points is rational. Erdős and Ulam conjectured in 1945 that there is no dense rational distance set in \mathbb{R}^2 .

Fact 1.1.2.5 ([Sha18, Cor. 1.4]). *Assume Conjecture 1.1.2.1 for algebraic surfaces over all number fields. Let S be an infinite rational distance set. Then either all but at most 4 points of S are on a line, or all but at most 3 points of S are on a circle.*

²Subvarieties are assumed to be closed.

A complex manifold M is called *Brody hyperbolic* if every morphism $\mathbb{C} \rightarrow M$ of complex manifolds is constant. For example, by [DR16, p.417], a compact Riemann surface is Brody hyperbolic if and only if its genus is at least 2.

Conjecture 1.1.2.6 ([Lan86, Conjecture 5.6], see also [BD21, Conjecture, p.2]). *A complex smooth projective variety is hyperbolic if and only if every subvariety is of general type.*

Conjecture 1.1.2.6 is known as the geometric Lang conjecture. It lies between algebraic geometry and complex analytic geometry. Both directions of it are unknown till now. For subvarieties of abelian varieties, Conjecture 1.1.2.6 is confirmed by [Yam19, Cor. 1.3] (and Brody's theorem [Bro78, p.213] that Brody hyperbolicity agrees with Kobayashi hyperbolicity for compact complex manifolds).

Conjecture 1.1.2.7 would follow from Conjectures 1.1.2.1 and 1.1.2.6.

Conjecture 1.1.2.7 ([Lan74, (1.3)]). *Let V be a smooth projective variety over a number field F . If a complex analytification of V is Brody hyperbolic, then $V(F)$ is finite.*

1.1.3 Lang conjecture for Shimura varieties

Shimura varieties are higher-dimensional analogs of modular curves. As Alex Youcis puts it, the reason to study Shimura varieties is multiple: They are highly symmetrical objects with rich actions of various Lie groups; They are moduli spaces of abelian varieties (with extra structures); They are moduli spaces of motives; They are objects conjectured to realize the global Langlands correspondence, *etc.* However, to define Shimura varieties requires an exceptional amount of technical sophistication. See [Mil17b] for a reference.

Let (G, X) be a Shimura datum, and let $K \leq G(\mathbb{A}_f)$ be a sufficiently small, neat, compact open subgroup. Let S be a connected component of the complex manifold $\mathrm{Sh}_K(G, X)$. From Nadel's work [Nad89, Thm. 0.2], the Baily-Borel compactification S^* of S is Brody hyperbolic. As the canonical model of $\mathrm{Sh}_K(G, X)$ exists (see, e.g., [Mil17b, p.128]), S is naturally a smooth quasi-projective variety defined over a number field F . Then Conjecture 1.1.2.7 predicts that $S(F')$ is finite for every finite extension F'/F . Similar speculation for integral points on Shimura varieties of abelian type is confirmed by Ullmo. His proof relies on Faltings's solution to Shafarevich's conjecture for abelian varieties (Fact 1.1.1.2).

Fact 1.1.3.1 ([Ull04, Thm. 3.2 (a)]). *Suppose that the Shimura datum (G, X) is of adjoint abelian type. Let $\Gamma \leq G(\mathbb{Q})$ be a net arithmetic lattice. Then for every number field F , every finite set of places Σ of F and every model \mathcal{M} of X^+/Γ over $\mathcal{O}_{F,\Sigma}$, the set $\mathcal{M}(\mathcal{O}_{F,\Sigma})$ is finite.*

Concerning the rational points on general Shimura varieties, Lang's conjecture (Conjecture 1.1.2.1) is related to an alternative principle [UY10, Thm. 1.1]. For a projective variety Z over a number field, Ullmo and Yafaev [UY10, (1)] define its *Lang locus* Z^L to be the Zariski closure of $\cup_M \overline{Z(M)}^{>0}$, where M runs through finite extensions of the definition field of Z (inside a fixed algebraic closure), and $\overline{Z(M)}^{>0}$ is the union of positive-dimensional irreducible components of the Zariski closure $\overline{Z(M)}$.

The Lang locus measures the failure of Lang's conjecture, since $Z^L = \emptyset$ if and only if Z satisfies Conjecture 1.1.2.1. For Shimura varieties, Fact 1.1.3.2 shows that Lang's conjecture is either true or very false.

Fact 1.1.3.2 (Ullmo-Yafaev's all-or-nothing principle, [UY10, Thm. 1.1]). *Let S be a (connected) Shimura variety of sufficiently high level. Then $S \cap (S^*)^L$ is either \emptyset or S .*

As Shimura varieties are not proper in general, it is equally natural to consider integral points instead of rational points. For quasi-projective varieties over \mathbb{Q} , we define an "integral Lang locus" measuring the infiniteness of integral points by choosing an integral model. This locus is independent of the choice of the model. It is empty if and only if the variety has only finitely many integral points over each number field where the variety can be defined. We give a result for integral points parallel to Fact 1.1.3.2. It shows that the Lang conjecture on integral points ([Lan91, IX, Conjecture 5.1]) is either true or very false for Shimura varieties.

Theorem (Theorem 2.5.0.12). *The integral Lang locus of a (connected) Shimura variety S is either \emptyset or S .*

In fact, we form several axioms for an abstract locus formation, and prove that such an alternative principle results from the axioms. Both Lang locus and integral Lang locus satisfy the axioms.

1.2 Lawrence-Venkatesh technique

Lawrence-Venkatesh's new proof ([LV20]) of Faltings's theorem (Fact 1.1.1.1) sheds light on Conjecture 1.1.2.1. This technique, compared with Faltings's strategy, is of local nature. We give a highly sketchy review, and refer the reader to [LV20] for more details.

1.2.1 Setting

Let K, S be as in Section 1.1.1. Let $f : X \rightarrow Y$ be a smooth proper morphism of smooth algebraic varieties over K . By enlarging S , one may choose a smooth proper morphism $\tilde{f} : \mathcal{X} \rightarrow \mathcal{Y}$ between smooth $O_{K,S}$ -schemes whose base change to K is exactly f . Lawrence-Venkatesh's idea uses the induced

variation of local Galois representations, to prove that $\mathcal{Y}(O_{K,S})$ is not Zariski dense in Y .

Remark 1.2.1.1. If Y is as in Fact 1.1.1.1, then by properness of Y over K and [Poo17, Thm. 3.2.13 (ii)], the natural map $\mathcal{Y}(O_{K,S}) \rightarrow Y(K)$ is bijective. By $\dim Y = 1$, a subset of Y which is not Zariski dense is necessarily finite. That is why one only needs nondensity of integral points in this case. Lawrence and Venkatesh [LV20] apply the following machinery to a sophisticated variant of the relative curve constructed by Parshin (and of a construction due to Kodaira).

1.2.2 Galois representations

Choose a finite place v of K with underlying rational prime p , such that p is unramified in K and no place dividing p is in S . Let K_v be the completion of K at v . Let $O_v \subset K_v$ be the integer ring. There is a natural inclusion $\mathcal{Y}(O_{K,S}) \subset \mathcal{Y}(O_v)$. For every $y \in \mathcal{Y}(O_v)$, the fiber \mathcal{X}_y is a smooth proper scheme over O_v with generic fiber X_y :

$$\begin{array}{ccc} \mathcal{X}_y & \longrightarrow & y \\ \downarrow & & \downarrow \\ \mathcal{X} & \longrightarrow & \mathcal{Y} \\ & \searrow & \downarrow \\ & & \text{Spec } O_v \end{array} \quad \begin{array}{ccc} X_y & \longrightarrow & y \\ \downarrow & & \downarrow \\ X_{K_v} & \longrightarrow & Y_{K_v} \\ & \searrow & \downarrow \\ & & \text{Spec } K_v. \end{array}$$

Let $\text{Rep}_{\mathbb{Q}_p}(\Gamma_{K_v})$ the category of (continuous) \mathbb{Q}_p -representations of the absolute Galois group Γ_{K_v} . For every integer $d \geq 0$, there is a *local* Galois representation

$$\rho_y^d : \Gamma_{K_v} \rightarrow \text{GL}(H_{\text{ét}}^d(X_{\bar{y}}/\overline{K_v}, \mathbb{Q}_p))$$

on the d -th étale cohomology group. For a locally small category \mathcal{C} , let \mathcal{C}/\sim be the set of isomorphism classes of objects of \mathcal{C} . Hence, one gets a map $\rho : \mathcal{Y}(O_v) \rightarrow \text{Rep}_{\mathbb{Q}_p}(\Gamma_{K_v})/\sim$. Representations are more or less “linear” data.

1.2.3 p -adic Hodge theory

The functor D_{cris} in p -adic Hodge theory induces a functor from the category $\text{Rep}_{\mathbb{Q}_p}(\Gamma_{K_v})$ to the category FVec_{K_v} of filtered vector spaces over K_v . Because the K_v -algebraic variety X_y has a smooth proper model \mathcal{X}_y over O_v , the p -adic Galois representation ρ_y^d is crystalline in the sense of [BC09, p.133]. By Fontaine’s conjecture proved by Faltings [Fal88, Cor., p.69], this functor sends ρ_y^d to the d -th de Rham cohomology $H_{\text{dR}}^d(X_y/K_v)$ equipped

with its Hodge filtration ([Sta24, Tag 0FM8]). The step is informally depicted below.

$$\mathcal{Y}(O_S) \xrightarrow[\text{choosing a suitable place } v|p]{\subset} \mathcal{Y}(O_v) \xrightarrow{\rho} \text{Rep}_{\mathbb{Q}_p}(\Gamma_{K_v}) / \sim \xrightarrow{D_{\text{cris}}} \text{FVec}_{K_v} / \sim .$$

Locally, one can interpret the map

$$\mathcal{Y}(O_v) \rightarrow \text{FVec}_{K_v} / \sim \quad (1.1)$$

as a period map. There is a vector bundle $V = \mathcal{H}_{\text{dR}}^d(X/Y)$ on Y , and a decreasing Hodge filtration $F^\bullet V$ by vector subbundles, whose fiber at every $y \in Y(K_v)$ is $H_{\text{dR}}^d(X_y/K_v)$ with its Hodge filtration. There is a natural *flat* connection ∇_{GM} on V , the Gauss-Manin connection.

1.2.4 Complex period map

We begin with the complex analytic analog. Consider a variation of Hodge structure $(V, F^\bullet V, \nabla)$ on a connected complex manifold Y , where V is a vector bundle, $F^\bullet V$ is a decreasing filtration of V by vector subbundles, and ∇ is a flat connection on V . (On a complex manifold, by connection we mean a holomorphic connection in the sense of [Huy05, Def. 4.2.17].) Take a base point $y_0 \in Y$ and a small open disk $\Omega \subset Y$ around y_0 . As ∇ is flat, for $y \in \Omega$, the parallel transport induces a \mathbb{C} -linear isomorphism $V_y \rightarrow V_{y_0}$. In general, the connection ∇ does not respect the filtration. Still, the fiberwise filtration $F^\bullet V_y$ is transported to a filtration on the fiber V_{y_0} , which has the same dimensional data as the filtration $F^\bullet V_{y_0}$. Let Flag/\mathbb{C} be the projective variety parameterizing the filtrations of V_{y_0} of this common dimension data. In this way, one gets a holomorphic map (only locally defined on Y), called a period map,

$$\Phi_{\mathbb{C}} : \Omega \rightarrow \text{Flag}, \quad y \mapsto \text{transport of } F^\bullet V_y \text{ to } V_{y_0}.$$

1.2.5 p -adic period map

Let k/\mathbb{Q}_p be a finite extension of \mathbb{Q}_p . Let O_k be the integer ring of k . Let m_k be the maximal ideal of O_k . Let \mathcal{Y} be a smooth O_k -scheme with generic fiber Y . Let $\bar{\mathcal{Y}}$ be the special fiber of \mathcal{Y} :

$$\begin{array}{ccccc} Y & \longrightarrow & \mathcal{Y} & \longleftarrow & \bar{\mathcal{Y}} \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \text{Spec } O_k & \longleftarrow & \text{Spec } O_k/m_k. \end{array}$$

Fix a base point $y_0 \in \mathcal{Y}(O_k)$. Denote by Ω the fiber passing y_0 of the reduction map $\mathcal{Y}(O_k) \rightarrow \bar{\mathcal{Y}}(O_k/m_k)$ to residue field, and call it the residue


disk around y_0 . Then Ω is an open neighborhood of y_0 in the k -analytic manifold Y^{an} . Consider a triple $(V, F^\bullet V, \nabla)$, where V is a vector bundle on Y , $F^\bullet V$ be a decreasing filtration on V by vector subbundles, and ∇ a flat connection. As in Section 1.2.4, one can similarly define a flag variety Flag over k , and a p -adic period map $\Phi_p : \Omega \rightarrow \text{Flag}$ which is k -analytic.

1.2.6 Ax-Schanuel property of period map

In the notation of Section 1.2.1, take $k = K_v$. Take the triple $(V, F^\bullet V, \nabla)$ to be $(\mathcal{H}_{\text{dR}}^d(X/Y) \otimes_K K_v, \text{Hodge filtration}, \nabla_{\text{GM}})$ on Y_{K_v} . When the fibers of V on a residue disk Ω are identified by the Gauss-Manin connection ∇_{GM} , the restriction of the map (1.1) to Ω coincides with the p -adic period map Φ_p . Because $\mathcal{Y}(O_v)$ is covered by finitely many residue disks, to prove that $\mathcal{Y}(O_{K,S})$ is not Zariski dense in Y , it suffices to prove the nondensity of $\Omega \cap \mathcal{Y}(O_{K,S})$. Fact 1.2.6.1 counts essentially on Bakker-Tsimerman's Ax-Schanuel type result [BT19]. Let \mathfrak{H}_p be the Zariski closure of $\Phi_p(\Omega)$ in Flag .

Fact 1.2.6.1 ([KM23, Prop. 7.10 (4)]). *Let $Z \subset \mathfrak{H}_p$ be a subvariety with $\dim \mathfrak{H}_p \geq \dim Z + \dim Y$. Then $\Phi_p^{-1}(Z)$ is not Zariski dense in Y_{K_v} .*

The situation is summarized as follows.



Take Z to be the Zariski closure of $\Phi_p(\Omega(O_{K,S}))$ in \mathfrak{H}_p . If $\dim \mathfrak{H}_p \geq \dim Z + \dim Y$, then by Fact 1.2.6.1, the subset $\Omega \cap \mathcal{Y}(O_{K,S}) \subset Y$ is not Zariski dense.

1.2.7 Summary

To show nondensity of integral points in Lawrence-Venkatesh's method, one needs to show that the dimension of the image \mathfrak{H}_p of the p -adic period map Φ_p is "large" when compared with that of $O_{K,S}$ -points. Let $\mathfrak{H}_{\mathbb{C}}$ be the image of the complex period map $\Phi_{\mathbb{C}}$ induced by $f_{\mathbb{C}} : X_{\mathbb{C}} \rightarrow Y_{\mathbb{C}}$. For one thing, as the Gauss-Manin connection ∇_{GM} is defined on K , one gets $\dim \mathfrak{H}_p \geq \dim \mathfrak{H}_{\mathbb{C}}$. Using the corresponding variation of Hodge structures, one proves that $\mathfrak{H}_{\mathbb{C}}$ contains the orbit of the base point under the monodromy action. For another thing, using Faltings's finiteness theorem (see, e.g., [LV20, Lem. 2.3]), one gets an upper bound (involving the centralizer of the crystalline Frobenius operator arising from the comparison of de Rham cohomology and crystalline cohomology) on $\Phi_p(\Omega(O_{K,S}))$.

1.3 Lawrence-Sawin technique

The technique of Lawrence-Venkatesh is a promising approach to Conjecture 1.1.2.1, because it is successfully applied in higher dimension. For example, based on this technique, Lawrence and Sawin establish in an innovative way the Shafarevich conjecture for hypersurfaces in abelian varieties. Let K, S be as in Section 1.1.1. Let A be an abelian variety over K of dimension g with good reduction outside S . A subvariety $V \subset A$ is said to have good reduction at a place $v \notin S$ of K if the Zariski closure of V in the unique abelian scheme $\mathcal{A}/\mathcal{O}_{K_v}$ with generic fiber A_{K_v} is smooth.

Fact 1.3.0.1 (Lawrence-Sawin, [LS20, Thm. 1.1]). *Suppose $\dim A \geq 4$. Fix an ample class ϕ in the Néron-Severi group of A . Then there are only finitely many hypersurfaces $H \subset A$ over K representing ϕ , with good reduction outside S , up to translation by points in $A(K)$.*

Using a similar technique, Krämer and Maculan obtained an analog for subvarieties of dimension less than half the dimension of the ambient abelian variety.

Fact 1.3.0.2 (Krämer-Maculan, [KM23, Cor., p.3]). *Fix a polynomial $P \in \mathbb{Q}[z]$ of degree $d < (g - 1)/2$ and an ample line bundle L on A . Then up to translation by points in $A(K)$, there are only finitely many nondivisible geometric complete intersections of ample divisors $X \subset A$ over K , with good reduction outside S , that have Hilbert polynomial P with respect to L and satisfy $2\chi(X \times X, \Omega_{X \times X}^d) \leq \chi_{\text{top}}(X \times X)$.*

In both cases of Facts 1.3.0.1 and 1.3.0.2, the dimension of the base algebraic variety Y in Section 1.2 is greater than 1. So nondensity of the set of integral points is strictly weaker than finiteness. An idea suggested in [LV20, Sec. 10.2] is to iterate the Lawrence-Venkatesh argument by replacing Y with the Zariski closure of integral points. In this manner, an estimate of topological monodromy group that is *uniform* in subvarieties of the variety under consideration is needed. The main novelty of [LV20] is to compare the monodromy with a Tannakian group. (The comparison involved in the proof of Fact 1.3.0.2 leans on [JKLM23].) This idea is similar to the study of monodromy groups of variation of Hodge structures via Mumford-Tate groups in [And92].

This Tannakian group arises from sheaf convolution developed fundamentally by Krämer-Weissauer [KW15b]. We give a cursory review of the comparison.

1.3.1 Tannakian theory of sheaf convolution

Tannakian formalism is a way to reconstruct a group from its representation theory. By the Tannaka-Krein duality, a compact group can be recovered

from the abelian category of its complex representations *together with the tensor product operation*.

Definition 1.3.1.1 ([DM22, Def. 2.19]). A rigid, symmetric, monoidal abelian category (C, \otimes) of unit object $\mathbb{1}$ is a *neutral Tannakian category* over a field k if it admits an exact faithful k -linear tensor functor $\omega : C \rightarrow \text{Vec}_k$ (called a fiber functor) and if $\text{End}(\mathbb{1}) = k$.

Fact 1.3.1.2 ([DM22, Thm. 2.11], [Del90, Sec. 9.2]). *Let (C, \otimes) be a neutral Tannakian category over a field k with a fiber functor $\omega : C \rightarrow \text{Vec}_k$. Then there is a natural affine group scheme $\text{Aut}^\otimes(C, \omega)$ over k (called the Tannakian group of (C, \otimes) at ω), such that ω factors through an equivalence $C \rightarrow \text{Rep}_k(\text{Aut}^\otimes(C, \omega))$ of symmetric monoidal categories. If k is algebraically closed, then $\text{Aut}^\otimes(C, \omega)$ is independent of the choice of ω up to k -isomorphism.*

We review the work of Krämer and Weissauer. Let A be a complex abelian variety. Perverse sheaves on algebraic variety are the singular version of local systems. They form a full, abelian subcategory $\text{Perv}(A)$ of the triangulated category $D_c^b(A)$ of complexes of sheaves with bounded, constructible cohomologies. This abelian category is Noetherian and Artinian. For every smooth subvariety $X \subset A$, the complex of sheaves $\mathbb{C}_X[\dim X]$ is a perverse sheaf on A .

Let $a : A \times A \rightarrow A$ be the group law. Let $p_i : A \times A \rightarrow A$ ($i = 1, 2$) be the projection to the i -th factor. The bifunctor

$$(\cdot) * (\cdot) : D_c^b(A) \times D_c^b(A) \rightarrow D_c^b(A), \quad (-, +) \mapsto Ra_*(p_1^* - \otimes^L p_2^* +)$$

is called the convolution on A . In general, $\text{Perv}(A)$ is not stable under the convolution. Still, its quotient modulo the subcategory of “negligible objects” is stable under the convolution. Let $N(A) \subset \text{Perv}(A)$ be the full subcategory comprised of (so-called negligible) objects with Euler characteristic 0.

Fact 1.3.1.3 ([Krä22, 1.b]). *The subcategory $N(A)$ is Serre (in the sense of [Sta24, Tag 02MO (1)]) in $\text{Perv}(A)$. Let $\bar{P}(A)$ be the quotient abelian category (in the sense of [Sta24, Tag 02MS]). Then the convolution descends to a bifunctor $* : \bar{P}(A) \times \bar{P}(A) \rightarrow \bar{P}(A)$. Moreover, $(\bar{P}(A), *)$ is a neutral Tannakian category.*

Every object $F \in \text{Perv}(A)$ generates a Tannakian subcategory $\langle F \rangle$ of $\bar{P}(A)$. Let $G(F)$ be the (unique up to isomorphism) Tannakian group of $\langle F \rangle$. The computation of the Tannakian group in [LS20] follows essentially the general approach in Krämer’s work [Krä22, Krä21].

1.3.2 Monodromy group and generic Tannakian group

In Lawrence-Sawin [LS20, Sec. 11], the strategy of Lawrence-Venkatesh is applied to the universal family of hypersurfaces $f : U \rightarrow \text{Hilb}$ over the corresponding Hilbert scheme:

$$\begin{array}{ccc} U & \hookrightarrow & \text{Hilb} \times A \\ & \searrow f & \downarrow \\ & & \text{Hilb} \end{array}$$

More generally, Krämer and Maculan [KM23, Sec. 1.4] consider the following setting. We work over \mathbb{C} . (One can similarly work over a base field of characteristic 0 and use perverse sheaves with ℓ -adic coefficients.) Let Y be a smooth *integral* variety. Let $X \subset A \times_{\mathbb{C}} Y$ be a subvariety, such that the projection $f : X \rightarrow Y$ is smooth of relative dimension d :

$$\begin{array}{ccc} X & \hookrightarrow & Y \times A \\ & \searrow f & \downarrow \\ A & \xleftarrow{\pi} & Y \end{array}$$

Then the flat vector bundle $(\mathcal{H}_{\text{dR}}^d(X/Y), \nabla_{\text{GM}})$ on the complex manifold Y^{an} is induced by the local system $R^d f_* \mathbb{C}_X$. To get “big monodromy”, we “twist” this local system as follows. Let

$$\Pi(A) := \text{Hom}(\pi_1(A, 0), \mathbb{C}^*)$$

denote the algebraic torus of characters of the fundamental group. For every character $\chi \in \Pi(A)$, let L_χ be the corresponding rank 1 local system on A^{an} . Consider the local system $V_\chi := R^d f_* \pi^* L_\chi$ on Y^{an} . Then $V_1 = R^d f_* \mathbb{C}_X$. Let $\text{Mon}(\chi)$ be the Zariski closure of the image of the monodromy representation of V_χ . We need to find χ such that the monodromy group $\text{Mon}(\chi)$ is big enough to carry out Lawrence-Venkatesh-Sawin’s method.

Let $\eta \in Y$ be the generic point. Denote the perverse sheaf $\mathbb{C}_{X_\eta}[d]$ on A_η by P_η . By Krämer-Weissauer’s theorem (Fact 1.4.0.1), for a generic $\chi \in \Pi(A)$, the functor

$$\omega_\chi : \langle P_\eta \rangle(\subset \bar{P}(A_\eta)) \rightarrow \text{Vec}_{\mathbb{C}}, \quad F \mapsto H^0(A_{\bar{\eta}}, F \otimes^L L_\chi)$$

is a *fiber functor*. Using $\omega_\chi(P_\eta) = V_{\chi, \bar{\eta}}$, one can compare $\text{Mon}(\chi)$ with $G(P_\eta)$.

Lemma 1.3.2.1 ([JKLM23, p.28]). *For a generic character $\chi \in \Pi(A)$, the monodromy group $\text{Mon}(\chi)$ is a closed subgroup of $G(P_\eta)$.*

To apply Bakker-Tsimerman’s Theorem (Fact 1.2.6.1), we need a lower bound on the monodromy group. Its proof uses the normality of the geometric generic Tannakian group inside the generic Tannakian group.

1.3.3 Normality of geometric generic Tannakian group

Let k be an algebraically closed field of characteristic 0 with an algebraic closure \bar{k} . Let A be an abelian variety over k . Let ℓ be a prime number. Let Λ be an algebraic closure of \mathbb{Q}_ℓ .

Fact 1.3.3.1 ([LS20, Lem. 3.7]). *Let G_k (resp. $G_{k'}$) be the Tannakian fundamental group of the category of geometrically semisimple perverse sheaves on A with coefficients in Λ (resp. summands of the pullbacks to $A_{\bar{k}}$ of geometrically semisimple perverse sheaves on A), modulo the full subcategory of “negligible objects”. Then $G_{k'}$ is naturally a normal closed subgroup of G_k , with quotient isomorphic to the Tannakian group of the neutral Tannakian category $(\text{Rep}_\Lambda(\Gamma_k), \otimes)$.*

The assumption of geometric semisimplicity in Fact 1.3.3.1 is removed in [JKLM23]. Let K/k be a field extension. Let k' be the algebraic closure of k in K . Assume that k'/k is Galois. Let $\mathcal{C} \subset \bar{P}(A)$ be a full abelian tensor subcategory. Let $\mathcal{C}_K \subset \text{Perv}(A_K)/N(A_K)$ be the full subcategory of subquotients of the pullbacks to A_K of perverse sheaves on A . Fix a fiber functor $\omega : \mathcal{C}_K \rightarrow \text{Vec}_\Lambda$.

Fact 1.3.3.2 ([JKLM23, Thm. 4.3]). *There is a short exact sequence of proalgebraic groups*

$$1 \rightarrow \text{Aut}^\otimes(\mathcal{C}_K, \omega) \rightarrow \text{Aut}^\otimes(\mathcal{C}, \omega) \rightarrow \text{Aut}^\otimes(\mathcal{C} \cap \text{Rep}_\Lambda(\text{Gal}(k'/k)), \omega) \rightarrow 1.$$

1.3.4 Monodromy group and geometric generic Tannakian group

Fact 1.3.3.1 and an analog of Larsen’s alternative ([LS20, Lem. 5.4]) permit one to get a lower bound on the monodromy group. Under certain geometric condition on the family $f : X \rightarrow Y$, by [LS20, Thm. 5.6] and [JKLM23, Thm. 4.10], for a generic character $\chi \in \Pi(A)$, the corresponding monodromy group contains the geometric generic Tannakian group, i.e., inclusions as follows:

$$\begin{array}{ccc} \text{Mon}(\chi) & \hookrightarrow & G(P_\eta) \\ \uparrow & \nearrow \text{normal} & \\ G(P_{\bar{\eta}}) & & \end{array}$$

1.3.5 Summary

In brief, in the work of Lawrence-Sawin and that of Krämer-Maculan, the crucial uniform estimation on the monodromy group follows from a comparison to the Tannakian group (of perverse sheaves on the geometric generic fiber). In fact, both the monodromy group and the Tannakian group

on the geometric generic fiber are embedded as closed subgroups in the Tannakian group on the generic fiber. The geometric generic Tannakian group is *normal* in the generic Tannakian group. This normality is used to prove that for most characters, the corresponding monodromy group contains the geometric generic Tannakian group.

1.3.6 Normality of monodromy group

Complementing Facts 1.3.3.1 and 1.3.3.2, we prove that for many characters, the associated monodromy group is also normal in the generic Tannakian group. This result poses a restriction on what the monodromy group can be.

One uses perverse sheaves on the generic fiber of an abelian scheme in Section 1.3.2. Hansen and Scholze's work [HS23] on relative perverse sheaves provides a way to study a family of perverse sheaves. Let $f : X \rightarrow Y$ be a morphism of algebraic varieties. Assume that the prime ℓ is invertible in the base field. By [HS23, Thm. 1.1], the category $D_c^b(X, \Lambda)$ has a unique t-structure, called the relative perverse t-structure, which restricts to the perverse t-structure on every geometric fiber of f . The heart $\text{Perv}(X/Y)$ is called the category of relative perverse sheaves. For every $y \in Y$, restricting to the fiber over y induces a functor $\text{Perv}(X/Y) \rightarrow \text{Perv}(X_y)$. An object of $\text{Perv}(X/Y)$ should be thought as a family of perverse sheaves. However, in general the abelian category $\text{Perv}(X/Y)$ is not Artinian.

To get an abelian category with many of the same properties familiar in the absolute setting, Hansen and Scholze add a condition, the so-called universal local acyclicity. Roughly, an object of $D_c^b(X, \Lambda)$ is universally locally acyclic if it satisfies the base change theorem. (For precision, see Definition 3.2.2.1.) The relative perverse t-structure preserves universally locally acyclic complexes. The resulting abelian subcategory $\text{Perv}^{\text{ULA}}(X/Y) \subset \text{Perv}(X/Y)$ is Noetherian, Artinian and compatible with Verdier duality. Moreover, if Y is smooth integral with generic point η , then the functor $\text{Perv}^{\text{ULA}}(X/Y) \rightarrow \text{Perv}(X_\eta)$ exhibits a Serre subcategory. In this sense, a universally locally acyclic relative perverse sheaf is determined by the perverse sheaf on the generic fiber.

Let k be an algebraically closed field of characteristic 0. Let A/k be an abelian variety. Let Y be an integral algebraic variety over k with generic point η . Let $\text{Perv}^{\text{ULA}}(A \times Y/Y)$ be the abelian category of universally locally acyclic relative perverse sheaves ([HS23]) with coefficients in Λ .

Theorem 1.3.6.1 (Theorem 3.1.2.2). *Assume $\dim A > 0$. Then there are uncountably many characters $\chi : \pi_1^{\text{ét}}(A) \rightarrow \Lambda^*$, such that the corresponding generic Tannaka group is a reductive group containing the monodromy group corresponding to χ as a closed reductive normal subgroup.*

In spirit, Theorem 1.3.6.1 is similar to that of André's normality theorem.

Fact 1.3.6.2 ([And92, Thm. 1]). *For a polarizable good variation of mixed Hodge structure over a smooth, connected, complex algebraic variety X and every x in the complement of some meager subset of X , the corresponding connected monodromy group is a normal subgroup of the corresponding derived Mumford-Tate group.*

Fact 1.3.6.2 is proved via the theorem of the fixed part due to Griffiths-Schmidt-Steenbrink-Zucker. In our case, an analog of the fixed part theorem is Theorem 3.1.2.3.

1.4 Generic vanishing

In the construction of the Tannakian category (in Fact 1.3.1.3), the existence of a fiber functor is deduced from Krämer-Weissauer's generic vanishing theorem.

Fact 1.4.0.1 (Krämer-Weissauer, [KW15b, Thm. 1.1]). *Let P be a perverse sheaf on a complex abelian variety A . Then there is a finite union $\mathcal{S}(P)$ of translates of strict algebraic subtori of $\Pi(A)$, such that for every character $\chi \in \Pi(A) \setminus \mathcal{S}(P)$ and every integer $i \neq 0$, one has $H^i(A, P \otimes^L L_\chi) = 0$.*

The proof of Fact 1.4.0.1 relies on two ingredients. The first is Deligne's [Del02] characterization of rigid symmetric monoidal abelian categories. It shows that a construction of André-Kahn [AKO02] leads to a super Tannakian category. The other is Kashiwara's conjecture for semisimple perverse sheaves. Its solution [Dri01] in turn relies on de Jong's conjecture ([BK06, Gai07]). Fact 1.4.0.1 shows that the super Tannakian category is in fact a neutral Tannakian category. As [KW15b, Sec. 3] explains, Krämer-Weissauer's theorem is a (partial) generalization of Green-Lazarsfeld's generic vanishing theorem.

Fact 1.4.0.2 ([GL87, Thm. 2]). *Let X be a compact Kähler manifold. Let*

$$w(X) = \max\{\text{codim}_X Z(\omega) : \omega \in H^0(X, \Omega_X^1) \setminus \{0\}\},$$

where $Z(\omega)$ denotes the zero-locus of a holomorphic one form ω . Then for any integers $i, j \geq 0$ with $i + j < w(X)$ and a generic line bundle $L \in \text{Pic}^0(X)$, one has $H^i(X, \Omega_X^j \otimes L) = 0$.

Green-Lazarsfeld's theorem is an analog of the Kodaira-Nakano vanishing theorem and answers a problem of Beauville [Uen83, Problem 8, p.620] affirmatively. Fact 1.4.0.1 implies generic vanishing theorem for compact Kähler manifolds whose Albanese manifolds are abelian varieties (for example, projective manifolds). In this sense, it generalizes Fact 1.4.0.2 partially. The reason is that Fact 1.4.0.1 is stated for abelian varieties. To

recover generic vanishing for all compact Kähler manifolds, one needs a generalization of Fact 1.4.0.1 for all complex tori.

Fact 1.4.0.3 (Bhatt-Schnell-Scholze, [BSS18, Thm. 1.1]). *Let P be a perverse sheaf on a complex torus A . Then there is a strict Zariski closed subset $\mathcal{S}(P)$ of the algebraic torus $\Pi(A)$ such that for every character $\chi \in \Pi(A) \setminus \mathcal{S}(P)$ and every integer $i \neq 0$, one has $H^i(A, P \otimes^L L_\chi) = 0$.*

Existing vanishing results are mainly stated for Kähler manifolds. Deligne [Del68] shows that parallel to the Kähler setting, every complex smooth proper algebraic variety (not necessarily Kähler) admits a Hodge theory. We show that generic vanishing results hold for such varieties. Instead of giving a demonstration parallel to the Kähler situation, one can give a uniform proof in Fujiki class \mathcal{C} .

A compact complex manifold is called in Fujiki class \mathcal{C} if it is the meromorphic image of a compact Kähler manifold. Compact Kähler manifolds and smooth proper complex algebraic varieties are in this class. Fujiki class \mathcal{C} admits a Hodge theory. We give a generic vanishing theorem for Fujiki class \mathcal{C} .

Theorem 1.4.0.4 (Theorem 4.7.1.3). *Let X be a complex manifold in Fujiki class \mathcal{C} with a flat unitary vector bundle F . Then for any two integers $p, q \geq 0$ with $\dim X - p - q$ larger than the defect of semismallness of an Albanese morphism of X , for a generic line bundle $L \in \text{Pic}^0(X)$, one has*

$$H^q(X, \Omega_X^p \otimes_{\mathcal{O}_X} F \otimes_{\mathcal{O}_X} L) = 0.$$

Corollary 1 (Corollary 4.7.2.6). *Let X/\mathbb{C} be a smooth proper algebraic variety with a flat unitary vector bundle F . Then for any two integers $p, q \geq 0$ with $\dim X - p - q$ larger than the defect of semismallness of an Albanese morphism of X , the locus*

$$\{L \in \text{Pic}^0(X) : H^q(X, \Omega_X^p \otimes_{\mathcal{O}_X} F \otimes_{\mathcal{O}_X} L) \neq 0\} \quad (1.2)$$

is contained in a finite union of translates of strict abelian subvarieties of the Picard variety $\text{Pic}_{X/\mathbb{C}}^0$.

The strategy of the proof of Theorem 1.4.0.4 is considering the unitary local system corresponding to F (provided by the Riemann-Hilbert correspondence). Its derived pushout along the Albanese morphism is a complex of constructible sheaves on a complex torus. For this complex of abelian sheaves, by estimating the perverse sheaf cohomologies, one deduces a generic vanishing result from Fact 1.4.0.3. This result proves Theorem 1.4.0.4 for vector bundles.

Fact 1.4.0.3 generalizes Krämer-Weissauer's theorem (Fact 1.4.0.1) to complex tori, but with a coarser control of the jump locus $\mathcal{S}(P)$ of a perverse

sheaf P . The finer control in Krämer-Weissauer's theorem results from the classification of simple perverse sheaves of Euler characteristic zero ([KW15b, Prop. 10.1 (a)]).

A question is that if Krämer-Weissauer's theorem (Fact 1.4.0.1) and the classification have generalizations to all complex tori. A positive answer would allow one to describe the failure locus of generic vanishing theorem for all compact Kähler manifolds. The proof of [KW15b, Prop. 10.1 (a)] uses Poincaré's reducibility theorem for abelian varieties, which fails for complex tori. Still, there is an independent proof due to Schnell [Sch15, Thm. 7.6] of Fact 1.4.0.1 as well as the classification. Schnell's proof is relatively elementary and makes profound use of a lift of Fourier-Mukai transform to D -modules.

1.5 Fourier-Mukai transform

Fourier-Mukai transform on abelian varieties, initiated by Mukai [Muk81], is an analog of the classical Fourier transform. For a ringed space (X, O_X) , let $\text{Mod}(O_X)$ be the category of O_X -modules. Let $D(O_X)$ be the derived category of the abelian category $\text{Mod}(O_X)$.

1.5.1 Construction

Let k be an algebraically closed field. Let A be an abelian variety over k with dual abelian variety B . Let $p_A : A \times_k B \rightarrow A$ (resp. $p_B : A \times_k B \rightarrow B$) be the projection to A (resp. B). Denote the normalized Poincaré bundle on $A \times_k B$ by \mathcal{P} .

Definition 1.5.1.1. The pair of functors

$$\begin{aligned} R\hat{S} &= Rp_{B*}(\mathcal{P} \otimes^L p_A^* \cdot) : D(O_A) \rightarrow D(O_B), \\ RS &= Rp_{A*}(\mathcal{P} \otimes^L p_B^* \cdot) : D(O_B) \rightarrow D(O_A) \end{aligned}$$

is called the *Fourier-Mukai transform* between A and B .

The Fourier-Mukai transform has found many applications in algebraic geometry: the Künneth decomposition for Chow motives [MD91], a new proof of Torelli's theorem [BP01], the study of stable bundles on elliptic surfaces [Bri98], *etc.* Motivated by noncommutative geometry, Ben-Bassat, Block and Pantev [BBBP07] study the Fourier-Mukai transform on complex tori. Similar to the classical Fourier inversion, a duality result for the Fourier-Mukai transform is stated in [Muk81, Thm. 2.2] (resp. [BBBP07, Thm. 2.1]) in the algebraic (resp. analytic) case. However, both statements are imprecise (Lemma 5.2.0.6). In the algebraic case, the minor problem is bypassed by adding a quasi-coherence condition. Let $D_{\text{qc}}(O_A) \subset D(O_A)$ be the full subcategory of objects with quasi-coherent cohomologies.

Fact 1.5.1.2 (Mukai). *The functor $R\hat{S}$ (resp. RS) restricts to a functor $D_{\text{qc}}(O_A) \rightarrow D_{\text{qc}}(O_B)$ (resp. $D_{\text{qc}}(O_B) \rightarrow D_{\text{qc}}(O_A)$). Moreover, there are canonical isomorphisms of functors*

$$\begin{aligned} RS \circ R\hat{S} &\cong T^{-g}[-1]_A^* : D_{\text{qc}}(O_A) \rightarrow D_{\text{good}}(O_A); \\ R\hat{S} \circ RS &\cong T^{-g}[-1]_B^* : D_{\text{qc}}(O_B) \rightarrow D_{\text{good}}(O_B), \end{aligned}$$

where T denotes the degree shift. In particular, $R\hat{S} : D_{\text{qc}}(O_A) \rightarrow D_{\text{qc}}(O_B)$ is an equivalence with a quasi-inverse $T^g[-1]_A^* RS$.

1.5.2 Complex tori

In the analytic setting, there are several competing definitions of “quasi-coherent” sheaves. A choice is the so-called *good* sheaves proposed by Kashiwara [Kas03, Def. 4.22]. With good sheaves, we give a way to correct [BBBP07, Thm. 2.1] in Chapter 5. First, we show that good sheaves are analytic analogs of quasi-coherent sheaves.

Proposition (GAGA, Proposition B.3.0.2). *Let X be a smooth proper complex algebraic variety. Then analytification induces an equivalence from the category of quasi-coherent O_X -module to the category of good $O_{X^{\text{an}}}$ -modules.*

Let A be a complex torus of dimension g . Let B be its dual torus. Let $D_{\text{good}}(O_A)$ be the full subcategory of $D(O_A)$ comprised of objects with good cohomologies. Notation for B are similarly understood. Set $RS : D(O_B) \rightarrow D(O_A)$ and $R\hat{S} : D(O_A) \rightarrow D(O_B)$ for the corresponding Fourier-Mukai transform.

Theorem 1.5.2.1 (Mukai, Ben-Bassat, Block, Pantev, Theorem 5.4.1.1). *The functor $R\hat{S}$ (resp. RS) restricts to a functor $D_{\text{good}}(O_A) \rightarrow D_{\text{good}}(O_B)$ (resp. $D_{\text{good}}(O_B) \rightarrow D_{\text{good}}(O_A)$). Moreover, there are canonical isomorphisms of functors*

$$\begin{aligned} RS \circ R\hat{S} &\cong T^{-g}[-1]_A^* : D_{\text{good}}(O_A) \rightarrow D_{\text{good}}(O_A); \\ R\hat{S} \circ RS &\cong T^{-g}[-1]_B^* : D_{\text{good}}(O_B) \rightarrow D_{\text{good}}(O_B). \end{aligned}$$

In particular, $R\hat{S} : D_{\text{good}}(O_A) \rightarrow D_{\text{good}}(O_B)$ is an equivalence with a quasi-inverse $T^g[-1]_A^* RS$.

Mukai’s proof for abelian varieties uses the flat base change theorem, of which we need an analytic analogue to prove Theorem 1.5.2.1. Our analytic replacement (Theorem 5.3.2.3) concerns only *smooth* base changes, but this weak version suffices for our purpose.

1.5.3 Homogeneous vector bundles

As an application of the analytic Fourier-Mukai transform, we recover Matsushima-Morimoto's classification of homogeneous vector bundles on complex tori ([Mat59, Mor59], see also [FL14, Thm. 7.1]).

The classification of vector bundles on a complex manifold X is completely worked out by Grothendieck [Gro57a] if X is the Riemann sphere $P_{\mathbb{C}}^1$, and by Atiyah [Ati57b] if X is an elliptic curve. When X is an abelian variety of higher dimension, "there are 'too' many vector bundles on X " ([Muk78, p.239]). Still, there are classification results for some special classes of vector bundles.

A vector bundle on a complex torus is called *homogeneous* if it is invariant under all translations. For example, a line bundle on A is homogeneous if and only if its isomorphism class is in $\text{Pic}^0(A)$.

An extension of finitely many O_A is called a *unipotent vector bundle* on A . By [FL14, Lem. 5.1], for every unipotent vector bundle U on A of rank r , there is a unipotent representation³ $\rho : \pi_1(A) \rightarrow \text{GL}_r(\mathbb{C})$ inducing U . More precisely, let \mathbb{C}_ρ be the local system of rank r on A corresponding to ρ . Then $\mathbb{C}_\rho \otimes_{\mathbb{C}_A} O_A$ is isomorphic to U . The extension of two homogeneous vector bundles is still homogeneous, so every unipotent vector bundle is homogeneous.

Theorem (Matsushima-Morimoto, Theorem 5.5.3.6). *A vector bundle F on the complex torus A is homogeneous if and only if F is isomorphic to $\bigoplus_{i=1}^n L_i \otimes_{\mathbb{C}_X} \mathbb{C}_{\rho_i}$, where $n \geq 0$ is an integer, $L_i \in \text{Pic}^0(X)$ and $\rho_i : \pi_1(X, 0) \rightarrow \text{GL}_{r_i}(\mathbb{C})$ is a unipotent representation of dimension r_i .*

1.6 Laumon-Rothstein transform

Laumon and Rothstein independently lift the Fourier-Mukai transform to D -modules and establish a duality result similar to [Muk81, Thm. 2.2]. The lift is referred to as the Laumon-Rothstein transform.

1.6.1 D -modules

On a complex manifold X , an O_X -module with a flat connection is called a D_X -module. A D_X -module is a flat vector bundle if and only if it is O_X -coherent. The reason that we need D -modules is twofold. For one thing, by the Riemann-Hilbert correspondence (see, e.g., [HT07, Thm. 7.2.1]), perverse sheaves on X are equivalent to regular holonomic D_X -modules. For another, Krämer-Weissauer's convolution theory relies on [KW15b,

³An action of an abstract group G on a finite dimensional vector space V by unipotent endomorphisms is called unipotent. By [Mil17a, Cor. 14.2], there is a basis of V for which G acts through upper triangular matrices with ones on the diagonal.

Prop. 10.1 (a)]. Its proof (resp. an independent proof of Schnell [Sch15]) uses characteristic cycles (resp. the Laumon-Rothstein transform) of D -modules.

1.6.2 Construction

Let A be an abelian variety over an algebraically closed field. Let B be the abelian variety dual to A . Set $g = \dim A$.

A difficulty in the theory of D -modules is that, the modules are over noncommutative ringed spaces. The Laumon-Rothstein transform turns them to modules over a commutative ringed space: the universal vector extension. By independent work of Rosenlicht [Ros58] and Serre [Ser88, Ch. VII], there is a universal vectorial extension $\pi : B^\natural \rightarrow B$, where B^\natural is a connected commutative algebraic group of dimension $2g$. Moreover, B^\natural is the moduli space of flat line bundles on A , which allows Schnell [Sch15] to apply Simpson's nonabelian Hodge theory [Sim93] there. Let $D^b(D_A)$ be the bounded derived category of the category of left D_A -modules, and let $D_{\text{qc}}^b(D_A)$ (resp. $D_c^b(D_A)$) be the full subcategory of $D^b(D_A)$ of objects with O -quasi-coherent (resp. D -coherent) cohomologies. Let $D_{\text{qc}}^b(O_{B^\natural})$ (resp. $D_c^b(O_{B^\natural})$) be the full subcategory of $D^b(O_{B^\natural})$ of objects with quasi-coherent (resp. coherent) cohomologies.

Let \mathcal{P}^\natural be the pullback of the Poincaré bundle \mathcal{P} along the morphism $\pi \times \text{Id}_A : B^\natural \times A \rightarrow B \times A$. By Mazur-Messing's theorem (see, e.g., [Lau96, Thm. 2.1.2]), B^\natural is the moduli space of flat line bundles on A . Thus, there is a flat connection ∇^\natural relative B^\natural on \mathcal{P}^\natural . Then the pair $(\mathcal{P}^\natural, \nabla^\natural)$ is naturally a $D_{B^\natural \times A/B^\natural}$ -module. Let $\tilde{\text{pr}} : B^\natural \times A \rightarrow A$ and $\tilde{\text{pr}}^\natural : B^\natural \times A \rightarrow B^\natural$ be the projections.

Definition 1.6.2.1 ([Lau96, p.14]). The Laumon-Rothstein transform between A and B is a pair of functors

$$\begin{aligned}\tilde{\mathcal{F}} &= R\tilde{\text{pr}}_*^\natural \text{DR}_{B^\natural \times A/B^\natural}((\mathcal{P}^\natural, \nabla^\natural) \otimes_{O_{B^\natural \times A}}^L \tilde{\text{pr}}^* \cdot) : D_{\text{qc}}^b(D_A) \rightarrow D_{\text{qc}}^b(O_{B^\natural}), \\ \tilde{\mathcal{F}}^\natural &= R\tilde{\text{pr}}_*((\mathcal{P}^\natural, \nabla^\natural) \otimes_{O_{B^\natural \times A}}^L L\tilde{\text{pr}}^{\natural*} \cdot) : D_{\text{qc}}^b(O_{B^\natural}) \rightarrow D_{\text{qc}}^b(D_A).\end{aligned}$$

The Laumon-Rothstein transform turns *noncommutative* D_A -modules to modules over the *commutative* ringed space B^\natural .

Fact 1.6.2.2 (Laumon-Rothstein, [Lau96, Thm. 3.2.1; Cor. 3.1.3], [Rot96, Thm. 4.5; Thm. 6.2], [Rot97]). One has $\tilde{\mathcal{F}}^\natural \tilde{\mathcal{F}} \cong T^{-g}[-1]_A^*$ on $D_{\text{qc}}^b(D_A)$ and $\tilde{\mathcal{F}} \tilde{\mathcal{F}}^\natural \cong T^{-g}[-1]_{B^\natural}^*$ on $D_{\text{qc}}^b(O_{B^\natural})$. In particular, the functor $\tilde{\mathcal{F}} : D_{\text{qc}}^b(D_A) \rightarrow D_{\text{qc}}^b(O_{B^\natural})$ is an equivalence of categories. Moreover, it restricts to an equivalence $D_c^b(D_A) \rightarrow D_c^b(O_{B^\natural})$.

1.6.3 Schnell's proof of Fact 1.4.0.1

The Laumon-Rothstein transform is a geometric tool to study generic vanishing theorems for perverse sheaves. Perverse sheaves on equivalent to regular, holonomic D -modules. The Riemann-Hilbert correspondence induces an isomorphism $\Phi : (B^\natural)^{\text{an}} \rightarrow \Pi(A)^{\text{an}}$ of complex manifolds. For a holonomic D_A -module M , by [Sch15, Sec. 3], the support of $\tilde{\mathcal{F}}(M)$ in B^\natural is identified *via* Φ with the failure locus in $\Pi(A)$ of generic vanishing for M . Schnell “deforms” the Laumon-Rothstein transform to a transform for Higgs bundles.

On a complex manifold X , in general a connection on a vector bundle is not O_X -linear. Higgs bundles can be regarded as degenerations of vector bundles with flat “linear” connection.

Definition 1.6.3.1. A Higgs bundle is a vector bundle E on X with a holomorphic one form $\phi \in \Gamma(X, \mathcal{E}nd(E))$ taking values in the bundle of endomorphisms of E such that $\phi \wedge \phi = 0$.

Deligne’s λ -connection is a notion interpolating between flat bundles and Higgs bundles.

Definition 1.6.3.2. For $\lambda \in \mathbb{C}$, a λ -connection of a vector bundle E on X is a \mathbb{C} -linear morphism of sheaves $\nabla : E \rightarrow \Omega_X^1 \otimes_{O_X} E$ such that for every open subset $U \subset X$, every $f \in O_X(U)$ and every $s \in \Gamma(U, E)$, one has $\nabla(f \cdot s) = f \cdot \nabla s + \lambda df \otimes s$. A λ -connection is called flat if its O_X -linear curvature operator $\nabla \circ \nabla : E \rightarrow \Omega_X^2 \otimes_{O_X} E$ is equal to 0.

Then 1-connection is exactly a connection, and a vector bundle with a flat 0-connection is the same as a Higgs bundle. The moduli spaces of λ -connections on projective varieties are studied in Simpson’s work [Sim97].

For a complex abelian variety A , Schnell [Sch15, Sec. 10] analyzes the moduli space $E(A)$ of line bundles on A with λ -connections. Let $\lambda : E(A) \rightarrow A_{\mathbb{C}}^1$ be the morphism taking the parameter of generalized connections. By [Sch15, Lem. 10.7, 10.9], one has $\lambda^{-1}(1) = B^\natural$ and $\lambda^{-1}(0)$ is the moduli space (i.e., $M_{\text{Dol}}(A)$ in [Sim93, p.363]) of rank 1 Higgs bundles on A . The morphism λ is real-analytically trivial, recovering the isomorphism $M_{\text{Dol}}(A) \rightarrow B^\natural$ of real Lie groups in nonabelian Hodge theory ([Sim93, p.364]).

Schnell [Sch15, Sec. 11] introduces an “extended Fourier-Mukai transform” taking values in $D(O_{E(A)})$. Restricting to $\lambda^{-1}(1)$, it coincides with the Laumon-Rothstein transform. Restricting to $\lambda^{-1}(0)$, it is essentially the Fourier transform for Higgs bundles ([Bon06, Bon10]).

Schnell deforms the holonomic D_A -module M to an O_{T^*A} -module M' , which is a “generalized” Higgs bundle (more precisely, a holonomic Higgs module as [Sab07, Example 5.1.6 (1)] shows). By definition, $\text{Supp } M'$ in T^*A has pure dimension g . Therefore, the support of the Fourier transform

of M' is a strict subset of $\lambda^{-1}(0)$. It is the intersection of the support of the extended Fourier-Mukai transform of M in $E(A)$ with $\lambda^{-1}(0)$. Using the real analytic isomorphism $\lambda^{-1}(0) \rightarrow \lambda^{-1}(1)$, Schnell proves that the support of $\tilde{\mathcal{F}}(M)$ is also a strict subset of B^\natural . Fact 1.4.0.1 follows from this strictness. Details can be found in [Sch15, Prop. 18.2].

1.6.4 Complex tori

An analytic Laumon-Rothstein transform may help to extend Schnell's method to all complex tori. By [Fav12, Thm. 3], an abelian variety A is determined by the derived category of D_A -modules. From Fact 1.6.2.2, it is also determined by the derived category of O_{B^\natural} -modules. However, this fails in the analytic case. Let A, B be complex tori dual to each other, of dimension g . By Proposition F.5.4.5 1, the universal vectorial extension $\pi : B^\natural \rightarrow B$ still exists. Contrary to the algebraic case, the complex Lie group $B^\natural = (\mathbb{C}^*)^{2g}$. Then B^\natural can only tell the dimension of A . In particular, one can no longer recover the complex structure of A from B^\natural ! That is why we need to replace B^\natural by something else in an analytic version of Fact 1.6.2.2.

In fact, we construct an O_B -subalgebra \mathcal{A}_B of $\pi_* O_{B^\natural}$, and define a pair of functors

$$\tilde{\mathcal{F}} : D(D_A) \rightarrow D(\mathcal{A}_B), \quad \tilde{\mathcal{F}}^\natural : D(\mathcal{A}_B) \rightarrow D(D_A).$$

A coherent D_A -module is called *good* if it admits global good filtration. (In the algebraic case, every coherent D -module admits a global good filtration. The complex analytic analog is false.) Let $D_{\text{good}}^b(D_A)$ (resp. $D_{O-\text{good}}(D_A)$) be the full subcategory of $D^b(D_A)$ (resp. $D(D_A)$) of objects with good (resp. O_A -good) cohomology. Theorem 1.6.4.1 is a “lift” of Theorem 1.5.2.1.

Theorem 1.6.4.1 (Theorem 6.1.2.2). *1. The pair $(\tilde{\mathcal{F}}, \tilde{\mathcal{F}}^\natural)$ is a lift of Fourier-Mukai transform in the sense that the following squares are commutative:*

$$\begin{array}{ccc} D(D_A) & \xrightarrow{\tilde{\mathcal{F}}} & D(\mathcal{A}_B) \\ \downarrow & & \downarrow \\ D(O_A) & \xrightarrow{R\hat{S}} & D(O_B), \end{array} \quad \begin{array}{ccc} D(D_A) & \xleftarrow{\tilde{\mathcal{F}}^\natural} & D(\mathcal{A}_B) \\ \downarrow & & \downarrow \\ D(O_A) & \xleftarrow{RS} & D(O_B), \end{array}$$

where the vertical functors are forgetful.

2. One has $\tilde{\mathcal{F}}\tilde{\mathcal{F}}^\natural \cong T^{-g}[-1]_B^$ on $D_{O-\text{good}}(\mathcal{A}_B)$ and $\tilde{\mathcal{F}}^\natural\tilde{\mathcal{F}} \cong T^{-g}[-1]_A^*$ on $D_{O-\text{good}}(D_A)$. Moreover, $\tilde{\mathcal{F}}^\natural\tilde{\mathcal{F}}$ preserves $D_{\text{good}}^b(D_A)$.*

1.6.5 Vector bundles with connection

A smooth vector bundle on a smooth manifold always admits a smooth connection. In the complex analytic case, a vector bundle may not admit any connection.

Fact 1.6.5.1 (Atiyah, [Ati57a, Theorems 2, 5, 6]). *Let X be a compact Kähler manifold. Let E be a vector bundle on X admitting a connection. Then for every integer $k > 0$, the k -th Chern class $c_k(E) = 0$ in $H^{2k}(X, \mathbb{R})$.*

Fact 1.6.5.1 leads to Question 1.6.5.2, which is attributed to Atiyah in [BD24, p.1].

Question 1.6.5.2. Does every vector bundle on a compact Kähler manifold admitting a connection also admit a flat connection?

Using the analytic Laumon-Rothstein transform, we recover a result of Matsushima [Mat59, Thm. 1] and Morimoto [Mor59, Thm. 2], which answers Question 1.6.5.2 affirmatively for complex tori.

Theorem. 1. (Theorem 6.3.3.1) *Let E be a coherent module on a complex torus with a connection ∇ . Then E is a homogeneous vector bundle and the pair (E, ∇) is translation invariant.*

2. (Proposition 6.5.2.1) *A homogeneous vector bundle on a complex torus admits a flat connection.*

1.7 Future directions

Several possible topics for further research are as follows. Depending on the limit author's knowledge, they vary from a vague idea to a relatively concrete plan.

1.7.1 Six-functor formalism of analytic quasi-coherent sheaves

As the proof of Theorem 1.5.2.1 needs a bit six-functor formalism in complex analytic geometry, there are a few natural questions: Does Theorem 1.5.2.1 have an analog for analytic quasi-coherent sheaves in Scholze and Clausen's sense ([Sch19] and [Sch22])? What is the relation between the notions of analytic quasi-coherence existing in the literature: the one of Scholze and Clausen, good sheaves proposed by Kashiwara (Definition A.1.4.1) and quasi-coherent sheaves in the sense of [RR74, p.100]?

1.7.2 Analytic Krämer-Weissauer's vanishing theorem

With the analytic Laumon-Rothstein transform and Theorem 1.6.4.1 at our disposal, we can study holonomic D -modules (instead of perverse sheaves

only) following Schnell [Sch15]. This shall lead to a convolution theory on complex tori, extending that on abelian varieties. The resulting analytic Krämer-Weissauer theorem would hopefully give a finer control of the loci (1.2) for not only projective manifolds but also compact Kähler manifolds.

1.7.3 Lawrence-Venkatesh’s method

Faltings [Fal83] deduced Mordell’s conjecture (Fact 1.1.1.1) from Shafarevich’s conjecture [Fal83, Satz 6]. By Shafarevich’s conjecture, for any integers $g \geq 1$ and $n \geq 3$, the Siegel variety $A_{g,n}$ (a Shimura variety parametrizing principally polarized abelian varieties of dimension g with a level n -structure) has only finitely many integral points ([Ull04, Prop. 3.1 (a)]). Now that Lawrence-Venkatesh’s method [LV20] can recover Faltings’s theorem, a natural question is if it can also prove the finiteness of integral points of $A_{g,n}$.

The situation should be compared to that in [LS20, p.7], where the authors considered the universal hypersurface inside a *constant* abelian scheme and compared its Tannakian group with the monodromy group. Similarly, to study Shafarevich’s conjecture, we can consider the convolution of relative perverse sheaves ([HS23]) on the universal abelian variety over $A_{g,n}$. Then we may calculate the Tannakian group associated with the universal theta divisor and try to relate it to the corresponding monodromy group.

1.8 Overview

The thesis consists of several independent chapters. Chapter 2 contains an arithmetic result related to Conjecture 1.1.2.1. The geometric foundation of the work [LS20, KM23] has inspired the study in Chapters 3, 4 and 6. Chapter 3 is related to the monodromy comparison part of [LS20, KM23]. Chapters 4, 5 and 6 are of complex analytic nature, completely independent of arithmetic. Appendix A reviews generalities of sheaves of modules over ringed spaces and supplements Chapter 5. Appendix B extend the classical GAGA theorem from coherent sheaves to quasi-coherent sheaves. Appendix C shows that quasi-coherent sheaves on complex analytic spaces form an abelian category. Appendix D complements Chapter 4 by giving more details. Appendix E concerns basics of D -modules and adds a detail on the Laumon-Rothstein theorem (Fact 1.6.2.2). Appendix F details a construction used in Chapter 6 and investigates related group-theoretic problems.

Chapter 2

Integral points of Shimura varieties: an “all or nothing” principle

2.1 Introduction

A complex analytic space X is called *Brody hyperbolic*, if every morphism $\mathbb{C} \rightarrow X$ is constant. For example, by [Cos05, p.78], a genus g compact Riemann surface is Brody hyperbolic if and only if $g \geq 2$. Conjecture 2.1.0.1 predicts that hyperbolicity (geometric property) restricts the behavior of rational points (arithmetic result).

Conjecture 2.1.0.1 (Lang, [Lan74, (1.3)], [Lan86, p.160]). *Let X be an integral projective variety over a number field $F(\subset \mathbb{C})$. If the complex analytification $X(\mathbb{C})$ is Brody hyperbolic, then the set of rational points $X(F)$ is finite.*

Ullmo and Yafaev [UY10] study Conjecture 2.1.0.1 in the case of Shimura varieties. Let (G, X) be a Shimura datum (in the sense of [Mil17b, Def. 5.5]). Let $K \leq G(\mathbb{A}_f)$ be a compact open subgroup. For every connected component $S \subset \mathrm{Sh}_K(G, X)$, denote the Baily-Borel compactification of S by S^* . Fact 2.1.0.2 is derived from [Nad89, Thm. 0.2] in the paragraph following [UY10, Thm. 2.1].

Fact 2.1.0.2 (Nadel). *There is an open subgroup $K' \leq K$, such that for every induced finite étale cover $S' \rightarrow S$, the Baily-Borel compactification S'^* is Brody hyperbolic.*

For one thing, by Fact 2.1.0.2, shrinking K to a sufficiently small open subgroup, one may and will assume that the Shimura variety S is Brody hyperbolic. For another, S^* has a natural structure of projective variety over a number field $F(\subset \mathbb{C})$. Then Conjecture 2.1.0.1 predicts $S(F')$ to be

finite for every finite extension F'/F . Ullmo and Yafaev [UY10] introduce “Lang locus” (Example 2.2.0.2) for algebraic varieties over $\bar{\mathbb{Q}}$ to measure the failure of Conjecture 2.1.0.1. In particular, the Lang locus of an algebraic variety over $\bar{\mathbb{Q}}$ is empty if and only if it has only finitely many rational points over every number field where it can be defined. The Lang locus of Shimura varieties satisfies an alternative principle.

Fact 2.1.0.3 ([UY10, Thm. 1.1]). *Let S be a Shimura variety of sufficiently high level. Then its Lang locus is either empty or full S .*

As Ullmo and Yafaev put it, Fact 2.1.0.3 means that for Shimura varieties, Conjecture 2.1.0.1 is either true or very false.

As Shimura varieties are not proper in general, it is natural to consider integral points. Conjecture 2.1.0.1 predicts that S has only finitely many integral points. We derive an analogue of Fact 2.1.0.3 for integral points. We define a notion of “integral Lang locus” (Definition 2.5.0.1) for algebraic varieties over $\bar{\mathbb{Q}}$ that measures the failure of finiteness of integral points.

Theorem (Theorem 2.5.0.12). *The integral Lang locus of a Shimura variety S is either empty or full S .*

Notation and conventions

Let $\bar{\mathbb{Q}}$ be the algebraic closure of \mathbb{Q} in \mathbb{C} . Let \mathbb{A}_f be the ring of finite adèles of \mathbb{Q} . Unless otherwise specified, an algebraic variety means a finite type, separated, geometrically reduced scheme over a field. The closure of a subset of an algebraic variety is taken in the Zariski topology. A subvariety is assumed to be Zariski closed. A Zariski-closed subset of a variety is endowed with the reduced induced closed subscheme structure, hence a subvariety.

By an *étale cover* $X \rightarrow Y$, we mean a finite étale morphism between integral algebraic varieties. In particular, it is surjective. If $\text{Aut}(X/Y)$ acts transitively on each fiber, then $X \rightarrow Y$ is called a *Galois cover*, of Galois group $\text{Aut}(X/Y)$. For a topological space X , we write $X^{>0}$ for the union of irreducible components of positive Krull dimension. Then for every subspace $Y \subset X$, one has $Y^{>0} \subset X^{>0}$.

2.2 Locus formation

We shall show that an alternative principle (Corollary 2.4.0.3) for an abstract locus is a consequence of some axioms.

Suppose that for every integral algebraic variety X over $\bar{\mathbb{Q}}$, we define a subvariety $X^L \subset X$. For a reducible algebraic variety Z over $\bar{\mathbb{Q}}$, let $Z = \cup_{i=1}^n Z_i$ be the decomposition into irreducible components. Set $Z^L := \cup_{i=1}^n Z_i^L$. Suppose that the formation $(\cdot)^L$ satisfies Assumption 2.2.0.1.

Assumption 2.2.0.1. For any integral algebraic varieties X, Y over $\bar{\mathbb{Q}}$:

1. (Dimension) If $X^L \neq \emptyset$, then every irreducible component of X^L has positive dimension;
2. (Inheritance) For every closed immersion $i : X \rightarrow Y$ over $\bar{\mathbb{Q}}$, one has $i(X^L) \subset Y^L$;
3. (Cover) For every étale cover $f : X \rightarrow Y$ over $\bar{\mathbb{Q}}$, one has $f(X^L) \subset Y^L$;
4. (Iteration) One has $X^L \subset (X^L)^L$;
5. (Birational) For every finite birational morphism $f : X \rightarrow Y$ over $\bar{\mathbb{Q}}$, one has $f(X^L) \subset Y^L$.

For every integral algebraic variety X over $\bar{\mathbb{Q}}$, by Assumption 2.2.0.1 2, one has $X^L \supset (X^L)^L$. From Assumption 2.2.0.1 4, one has $X^L = (X^L)^L$.

Example 2.2.0.2. The Lang locus defined in [UY10, Sec. 2.2] satisfies Assumption 2.2.0.1. For every integral algebraic variety X over $\bar{\mathbb{Q}}$, by [Sta24, Tag 01ZM (1), Tag 01ZQ], there exist a number field F , an algebraic variety X_F over F and an isomorphism $X_F \otimes_F \bar{\mathbb{Q}} \rightarrow X$ over $\bar{\mathbb{Q}}$. For each finite subextension M/F , let $X(X_F, M)$ be the image of the natural injection¹ $X_F(M) \rightarrow X(\bar{\mathbb{Q}})$. The *Lang locus* of X relative to X_F is defined to be the Zariski closure of

$$\cup_M \overline{X(X_F, M)}^{>0}$$

in X , where M runs through all finite subextensions of F . By Lemma 2.2.0.3, the Lang locus depends only on X . From [UY10, Lemmes 2.3, 2.5], the Lang locus satisfies Assumption 2.2.0.1. It measures the failure of finiteness of rational points, since $X^L = \emptyset$ if and only if $X_F(M)$ is finite for every finite subextension M/F .

Lemma 2.2.0.3. *The Lang locus of X is independent of the choice of X_F .*

Proof. Take another model $X_{F'}$ over a number field F' . There is a $\bar{\mathbb{Q}}$ -isomorphism $X_F \otimes_F \bar{\mathbb{Q}} \rightarrow X_{F'} \otimes_{F'} \bar{\mathbb{Q}}$. Because $X_{F'}$ is separated, by [Gro65, Prop. 4.8.13], the morphism is defined over a number field F'' containing both F and F' . For every finite extension M/F , there is a number field M' containing M and F'' . Then $X(X_F, M) \subset X(X_{F'}, M')$, so $\overline{X(X_F, M)}^{>0} \subset \overline{X(X_{F'}, M')}^{>0}$ and hence the Lang locus relative to X_F is contained in that relative to $X_{F'}$. The reverse inclusion follows by symmetry. \square

¹The natural map $X_F(M) \rightarrow X_F$ is not injective in general. For instance, let $F = \mathbb{Q}$, $M = \mathbb{Q}(\sqrt{2})$ and $X_F = \mathbf{A}_{\mathbb{Q}}^1$. Then $\pm\sqrt{2} \in X_F(M)$ are mapped to the same closed point of X_F corresponding to the maximal ideal $(x^2 - 2) \subset \mathbb{Q}[x]$.

Remark 2.2.0.4. 1. The Lang locus X^L in Example 2.2.0.2 is slightly different from the “lieu de Lang” X_F^L (a Zariski closed subset of X_F) defined by [UY10, (1)]. Let $\phi : X \rightarrow X_F$ be the natural morphism of schemes. For every finite extension M/F , let $X_F[M]$ be the image of the natural map $X_F(M) \rightarrow X_F$. Then $\phi(X(X_F, M)) = X_F[M]$. Because ϕ is integral, and surjective integral morphisms preserve the dimension, one has $\phi(\overline{X(X_F, M)}) = \overline{X_F[M]}$ and $\phi(\overline{X(X_F, M)}^{>0}) = \overline{X_F[M]}^{>0}$. Hence $\phi(X^L) = X_F^L$.

2. For a finite birational morphism $f : X \rightarrow Y$ of integral algebraic varieties over $\bar{\mathbb{Q}}$, it is not clear whether the Lang locus of Y is the image of the Lang locus of X (even if this is stated in [UY10, p.697]). That is why we require only inclusion but not equality in Assumption 2.2.0.1 5.

We gather some consequences of Assumption 2.2.0.1.

Lemma 2.2.0.5. *Let X be an algebraic variety over $\bar{\mathbb{Q}}$.*

1. *If $X = \cup_{i=1}^r Z_i$, where each Z_i is a subvariety of X , then $X^L = \cup_{i=1}^r Z_i^L$.*
2. *If Z is an irreducible component of X^L , then $Z^L = Z$.*
3. *If $f : X \rightarrow Y$ is Galois cover over $\bar{\mathbb{Q}}$, then $f^{-1}(f(X^L)) = X^L$. If $Z \subset Y$ is an irreducible subvariety, and Z' is an irreducible component of $f^{-1}(Z)$, then $f(f^{-1}(Z)^L) = f(Z'^L)$.*

Proof.

1. By Assumption 2.2.0.1 2, one has $\cup_{i=1}^r Z_i^L \subset X^L$. If Y is an irreducible component of X , then there exists an index i such that $Y \subset Z_i$. From Assumption 2.2.0.1 2, one has $Y^L \subset Z_i^L$ and hence $X^L \subset \cup_{i=1}^r Z_i^L$.
2. Write $X^L = \cup_{i=1}^n Z_i$ for the decomposition into irreducible components with $Z_1 = Z$. By Assumption 2.2.0.1 4, one has

$$Z \subset X^L = (X^L)^L = \cup_{i=1}^n Z_i^L.$$

As Z is irreducible, there is an index i such that $Z \subset Z_i^L \subset Z_i$. As $Z = Z_1$ is an irreducible component of X^L , one has $i = 1$ and $Z = Z^L$.

3. For every $x \in f^{-1}(f(X^L))$, there is $x' \in X^L$ with $f(x') = f(x)$. Let Θ be the Galois group of $f : X \rightarrow Y$. There is $\theta \in \Theta$ with $\theta(x') = x$, so $x \in X^L$ by Assumption 2.2.0.1 2. Therefore, $f^{-1}(f(X^L)) = X^L$. Since Θ permutes transitively the irreducible components of $f^{-1}(Z)$, one has $f^{-1}(Z) = \Theta \cdot Z'$. By Part 1, one has $f^{-1}(Z)^L = \Theta \cdot Z'^L$ and hence $f(f^{-1}(Z)^L) = f(Z'^L)$.

□

Given an étale cover $f : X \rightarrow Y$ over $\bar{\mathbb{Q}}$, the induced morphism $X^L \rightarrow Y^L$ may not be surjective. We introduce a sublocus that lifts along all étale covers.

For an integral algebraic variety X over $\bar{\mathbb{Q}}$, define its *locus at infinite level* by

$$X^{L\infty} := \cap_{f:T \rightarrow X} f(T^L),$$

where $f : T \rightarrow X$ runs through all étale covers of X . By Assumption 2.2.0.1 3, the sublocus $X^{L\infty}$ is a subvariety of X^L . As X is topologically Noetherian, and $f(T^L) \subset X$ is closed for every such $f : T \rightarrow X$, there exists a particular cover $f_1 : X_1 \rightarrow X$ with $f_1(X_1^L) = X^{L\infty}$. For every étale cover $X_2 \rightarrow X_1$, the composition $X_2^L \rightarrow X_1^L \xrightarrow{f_1} X^{L\infty}$ is still surjective.

Remark 2.2.0.6. By Assumption 2.2.0.1 1, if $X^{L\infty} \neq \emptyset$, then its irreducible components are *positive dimensional*.

For a reducible algebraic variety Y over $\bar{\mathbb{Q}}$, let $Y = \cup_{i=1}^n Y_i$ be the decomposition into irreducible components. Define $Y^{L\infty} = \cup_{i=1}^n Y_i^{L\infty}$, which is a subvariety of Y^L .

Lemma 2.2.0.7. *Let $f : T \rightarrow S$ be an étale cover over $\bar{\mathbb{Q}}$. Then $f^{-1}(S^{L\infty}) = T^{L\infty}$. In particular, $T^{L\infty} = T$ is equivalent to $S^{L\infty} = S$, and $S^{L\infty} = S^L$ implies $T^{L\infty} = T^L$.*

Proof. • We show $T^{L\infty} \subset f^{-1}(S^{L\infty})$.

Fix $t \in T^{L\infty}$, and set $s = f(t)$. For every étale cover $g : S' \rightarrow S$, there is a commutative diagram

$$\begin{array}{ccc} & T' & \\ g' \swarrow & & \searrow f' \\ T & & S' \\ f \searrow & & \swarrow g \\ & S & \end{array}$$

where each arrow is an étale cover. There is $t' \in T'^L$ with $g'(t') = t$. Then by Assumption 2.2.0.1 3, one has $s' := f'(t') \in S'^L$ and $s = g(s') \in g(S'^L)$. Hence $s \in S^{L\infty}$.

• We show $T^{L\infty} \supset f^{-1}(S^{L\infty})$.

Take $t \in f^{-1}(S^{L\infty})$. Then $s := f(t) \in S^{L\infty}$. For every étale cover $u : Z \rightarrow T$, there is an étale cover $v : N \rightarrow Z$ such that the composition $N \xrightarrow{v} Z \xrightarrow{u} T \xrightarrow{f} S$ is a Galois cover. One has

$$\begin{aligned} (u \circ v)^{-1}(t) &\subset (f \circ u \circ v)^{-1}(s) \subset (f \circ u \circ v)^{-1}(S^{L\infty}) \\ &\stackrel{(a)}{\subset} (f \circ u \circ v)^{-1}((f \circ u \circ v)(N^L)) = N^L, \end{aligned}$$

where (a) uses Lemma 2.2.0.5 3. Thus, one has $u^{-1}(t) \subset v(N^L) \subset Z^L$ and $t \in u(Z^L)$. Hence $t \in T^{L\infty}$.

- The equality $T^{L\infty} = T$ is equivalent to $S^{L\infty} = S$.

If $T^{L\infty} = T$, then $S^{L\infty} = f(f^{-1}(S^{L\infty})) = f(T^{L\infty}) = f(T) = S$. Conversely, if $S^{L\infty} = S$, then $T^{L\infty} = f^{-1}(S^{L\infty}) = f^{-1}(S) = T$.

- The equality $S^{L\infty} = S^L$ implies $T^{L\infty} = f^{-1}(S^{L\infty}) = f^{-1}(S^L) \stackrel{(b)}{\supset} T^L$, where (b) uses Assumption 2.2.0.1 3. Hence $T^{L\infty} = T^L$.

□

2.3 Shimura varieties

We review some basic facts about Shimura varieties, the main objects of interest in this note. We use essentially results on the geometry of Hecke correspondences and special subvarieties from [UY10, UY14].

Basics

Let G be an affine algebraic group over \mathbb{Q} .

Definition 2.3.0.1 ([Pin90, Sec. 0.1, p.13]). For every prime number p , choose an embedding $\bar{\mathbb{Q}} \rightarrow \bar{\mathbb{Q}}_p$.

1. For an element $g = (g_p)_p \in \mathrm{GL}_n(\mathbb{A}_f)$, let $\Gamma_p \leq \bar{\mathbb{Q}}_p^\times$ be the subgroup generated by all eigenvalues of $g_p \in \mathrm{GL}_n(\mathbb{Q}_p)$. If the intersection of the torsion subgroups

$$\cap_p (\bar{\mathbb{Q}}^\times \cap \Gamma_p)_{\mathrm{tor}} = \{1\}$$

for p running through all primes, then g is called *neat*.

2. An element of $G(\mathbb{A}_f)$ is *neat* if its image under some faithful algebraic representation of $G \rightarrow \mathrm{GL}_n/\mathbb{Q}$ is neat.
3. A subgroup of $G(\mathbb{A}_f)$ is *neat* if all its elements are neat.

Fact 2.3.0.2 ([Pin90, p.13]).

1. Let $K \leq G(\mathbb{A}_f)$ be a compact open subgroup. Then there is an open normal subgroup $K' \leq K$ that is neat.
2. Let $K \leq G(\mathbb{A}_f)$ be a neat subgroup. Then $K \cap G(\mathbb{Q})$ is a neat subgroup of $G(\mathbb{Q})$ (in the sense of [Mil17b, p.34]).

Let (G, X) be a Shimura datum. The set $G(\mathbb{R})$ is naturally a (real) Lie group. For a Lie group L , let L^+ be its identity component. Let G^{ad} be the quotient of G by its center. Set $G(\mathbb{R})_+$ to be the preimage of $G^{\text{ad}}(\mathbb{R})^+$ under the natural morphism $G(\mathbb{R}) \rightarrow G^{\text{ad}}(\mathbb{R})$ of Lie groups. Then $G(\mathbb{Q})^+ \subset G(\mathbb{Q})_+ \subset G(\mathbb{Q})$. By [Noo06, p.168] and [Mil17b, Prop. 5.9], X is naturally a finite disjoint union of isomorphic hermitian symmetric domains. Let X^+ be a connected component of X . By [Mil17b, Prop. 5.7 (b)], the stabilizer of X^+ in $G(\mathbb{Q})$ is $G(\mathbb{Q})_+ := G(\mathbb{Q}) \cap G(\mathbb{R})_+$.

Let $K \leq G(\mathbb{A}_f)$ be a compact open subgroup. From Fact 2.3.0.2 1, by passing to an open subgroup of K , we may and *always* assume that K is neat. Then by [Pin90, Prop. 3.3 (b)], $\text{Sh}_K(G, X) := G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K$ is naturally a complex manifold. For every $g \in G(\mathbb{A}_f)$, put $\Gamma_g := gKg^{-1} \cap G(\mathbb{Q})_+$ and $S_g := \Gamma_g \backslash X^+$. By Fact 2.3.0.2 2 and [Mil17b, Prop. 4.1], Γ_g is a neat (hence torsion-free) arithmetic subgroup of $G(\mathbb{Q})$ (in the sense of [Mil17b, p.33]). From [Mil17b, Prop. 3.1], $S_g = [X^+, g]_K$ is naturally a connected complex manifold. Let C be a set of representatives for the double coset space $G(\mathbb{Q})_+ \backslash G(\mathbb{A}_f) / K$. From [Mil17b, Lemmas 5.12 and 5.13], the set C is finite, and as complex manifold² $\text{Sh}_K(G, X) = \sqcup_{g \in C} S_g$.

By [Mil17b, Thm. 3.12; Cor. 3.16], the complex manifold S_g has a canonical structure of a complex algebraic variety. The algebraic variety S_g is an irreducible, *smooth* arithmetic locally symmetric variety ([Mil17b, p.58]). It is Zariski-open in its Baily-Borel compactification S_g^* ([Mil17b, p.40]), which is a projective variety. Thus, $\text{Sh}_K(G, X)$ is also a smooth quasi-projective (reducible) complex algebraic variety.

Let $E(G, X) \subset \mathbb{Q}$ be the reflex field of the Shimura datum (G, X) (in the sense of [Mil17b, Def. 12.2]). By [Mil17b, Rk. 12.3 (a)], $E(G, X)$ is a number field. From [Mil99, Rk. 2.4 (b)] and [Mil17b, p.128], $\text{Sh}_K(G, X)$ admits a unique (up to a unique isomorphism) *canonical model* over $E(G, X)$ (in the sense of [Mil17b, Def. 12.8]). Hence a *smooth* quasi-projective variety $\text{Sh}_K(G, X)$ over $E(G, X)$. By [Del71b, Cor. 5.4], for every morphism of Shimura data $f : (G', X') \rightarrow (G, X)$ and every compact open subgroup $K \leq G(\mathbb{A}_f)$ containing $f(K')$, the induced morphism $\text{Sh}_{K'}(G', X') \rightarrow \text{Sh}_K(G, X)$ is defined over a number field. Assume that f is the identity. Then the induced morphism (denoted by $p_{K', K}$) is finite étale and defined over $E(G, X)$. For every irreducible component $S' \subset \text{Sh}_{K'}(G', X')$, its image $S \subset \text{Sh}_K(G, X)$ is an irreducible component, and the restriction $S' \rightarrow S$ is an étale cover defined over a finite extension of $E(G, X)$. From [CK16, p.1901], when K' is normal in K , this étale cover is Galois.

By [Moo98b, p.282] and [GN20, Remark (3), p.56], every connected component $S \subset \text{Sh}_K(G, X)$ and its inclusion $S \rightarrow S^*$ to the Baily-Borel compactification are defined over a finite abelian extension of $E(G, X)$.

²But compare with [EY03, Sec. 5, p.634] and [Noo06, p.169].

Such an S is called a Shimura variety³ associated with (G, X, K) .

Hecke correspondences

By [Mil17b, Thm. 13.6], for every $g \in G(\mathbb{A}_f)$, there is an isomorphism $T(g) : \text{Sh}_K(G, X) \rightarrow \text{Sh}_{g^{-1}Kg}(G, X)$ of algebraic varieties over $E(G, X)$. For every $h \in G(\mathbb{A}_f)$, the morphism $T(g)$ sends the connected component $[X^+, h]_K \subset \text{Sh}_K(G, X)$ isomorphically to $[X^+, hg]_{g^{-1}Kg} \subset \text{Sh}_{g^{-1}Kg}(G, X)$. The algebraic correspondence

$$\text{Sh}_K(G, X) \xleftarrow{p_{K \cap gKg^{-1}, K}} \text{Sh}_{K \cap gKg^{-1}}(G, X) \xrightarrow{p_{K \cap gKg^{-1}, gKg^{-1}}} \text{Sh}_{gKg^{-1}}(G, X) \xrightarrow{T(g)} \text{Sh}_K(G, X)$$

over $E(G, X)$ is denoted by $T_g^\mathbb{A}$, and called the *adelic Hecke correspondence* induced by g .

Let $S = (K \cap G(\mathbb{Q})_+) \backslash X^+$. For every $q \in G(\mathbb{Q})_+$, let $S_q = (K \cap q^{-1}Kq \cap G(\mathbb{Q})_+) \backslash X^+$. Then S_q is the connected component $[X^+, 1]$ of $\text{Sh}_{K \cap q^{-1}Kq}(G, X)$ (resp. $\text{Sh}_K(G, X)$). The map Id_{X^+} (resp. $X^+ \rightarrow X^+, x \mapsto q \cdot x$) induces an étale cover $\alpha_q : S_q \rightarrow S$ (resp. $\beta_q : S_q \rightarrow S$). There is a commutative diagram

$$\begin{array}{ccccc} S & \xleftarrow{\alpha_q} & S_q & \xrightarrow{\beta_q} & S \\ \downarrow & & \downarrow & & \downarrow \\ \text{Sh}_K(G, X) & \xleftarrow{p_{K \cap q^{-1}Kq, K}} & \text{Sh}_{K \cap q^{-1}Kq}(G, X) & \xrightarrow{T(q^{-1})p_{K \cap q^{-1}Kq, q^{-1}Kq}} & \text{Sh}_K(G, X) \end{array}$$

of complex manifolds. Therefore, the correspondence

$$S \xleftarrow{\alpha_q} S_q \xrightarrow{\beta_q} S$$

is algebraic and defined over a number field. It is called the *(rational) Hecke correspondence* induced by q , and denoted by T_q .

Let $\{q_i\}_{i=1}^n$ be elements of $G(\mathbb{Q})_+ \cap KgK$ satisfying

$$G(\mathbb{Q})_+ \cap KgK = \sqcup_{i=1}^n \Gamma q_i^{-1} \Gamma, \Gamma := K \cap G(\mathbb{Q})_+.$$

By [KY14, p.881], the correspondence on $[X^+, 1] \subset \text{Sh}_K(G, X)$ induced by $T_g^\mathbb{A}$ decomposes as $\sum_{i=1}^n T_{q_i}$. For instance, the correspondences $T_1^\mathbb{A}$ and T_1 are the identity.

³By [Moo98b, Prop. 2.9] and [Mil17b, Thm. 5.17], when K is sufficiently small, S is a connected Shimura variety (in the sense of [Mil17b, Def. 4.10]).

Special subvarieties

Definition 2.3.0.3. [Moo98a, Def. 2.5] An irreducible subvariety $Z \subset \text{Sh}_K(G, X)$ over \mathbb{C} is called special, if there exists a connected, reductive algebraic subgroup $H \leq G$ defined over \mathbb{Q} , an element $g \in G(\mathbb{A}_f)$ and a connected component D_H^+ of

$$D_H := \{x \in X \mid h_x : \text{Res}_{\mathbb{C}/\mathbb{R}}(\mathbb{G}_m) \rightarrow G_{\mathbb{R}} \text{ factors through } H_{\mathbb{R}}\},$$

such that $Z(\mathbb{C})$ is the image of $D_H^+ \times gK$ in $\text{Sh}_K(G, X)(\mathbb{C}) = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f)/K$.

By [Moo98a, 2.4], D_H is a finite union of $H(\mathbb{R})$ -conjugacy classes. Let C be the $H(\mathbb{R})$ -conjugacy class containing D_H^+ . Then (H, C) is a Shimura subdatum⁴ of (G, X) . Then from [Del71b, Cor. 5.4] and [Moo98a, Rk. 2.6], every special subvariety of $\text{Sh}_K(G, X)$ is defined over a number field.

Example 2.3.0.4. 1. A complex point $s \in \text{Sh}_K(G, X)$ is a special subvariety, if and only if there is a special point $x \in X$ (in the sense of [Mil17b, Def. 12.5]) and $g \in G(\mathbb{A}_f)$ with $s = [x, g]_K$.

2. When $H = G$, the corresponding special subvarieties of $\text{Sh}_K(G, X)$ are precisely the connected components.

For every $g \in G(\mathbb{A}_f)$ and every irreducible subvariety $Z \subset \text{Sh}_K(G, X)$ over \mathbb{C} , Z is special if and only if $T(g)(Z)$ is special in $\text{Sh}_{g^{-1}Kg}(G, X)$. By [Moo98a, Sec. 2.9], an irreducible component of the intersection of a family of special subvarieties of $\text{Sh}_K(G, X)$ over \mathbb{C} is again special. Therefore, for a complex, irreducible subvariety $Y \subset \text{Sh}_K(G, X)$, there is a smallest special subvariety $Z_Y \subset \text{Sh}_K(G, X)$ containing Y . We say that Y is *Hodge generic* in Z_Y . Let $\mathbb{S} := \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$ be the Deligne torus.

Definition 2.3.0.5. The *generic Mumford-Tate group* (denoted by $\text{MT}(X)$) of the Shimura datum (G, X) is the smallest closed subgroup H of G over \mathbb{Q} , such that every $h : \mathbb{S} \rightarrow G_{\mathbb{R}}$ in X factors through $H_{\mathbb{R}}$. If $\text{MT}(X) = G$, then the Shimura datum (G, X) is called irreducible.

The subgroup $\text{MT}(X) \leq G$ is normal, connected and reductive. By [Che09, Def. 1.3.3], $(\text{MT}(X), X)$ is a Shimura subdatum of (G, X) . Fact 2.3.0.6 characterizes special subvarieties as Hecke image of irreducible components of a Shimura subvariety. Recall that $K \leq G(\mathbb{A}_f)$ is a *neat*, compact open subgroup. For $g \in G(\mathbb{A}_f)$, the quotient $S_g = \Gamma_g \backslash X^+$ is an irreducible component of $\text{Sh}_K(G, X)$.

Fact 2.3.0.6 ([UY10, Lem. 2.7]). *Let $(H, X_H) \subset (G, X)$ be an irreducible Shimura subdatum. Let X_H^+ be a connected component of X_H contained in*

⁴in the sense of [CLZ16, p.894]

X^+ . Set $\Gamma_{H,g} = gKg^{-1} \cap H(\mathbb{Q})_+$ and $\tilde{Z}_g := \Gamma_{H,g} \backslash X_H^+$ (an irreducible component of $\mathrm{Sh}_{gKg^{-1} \cap H(\mathbb{A}_f)}(H, X_H)$). Then the image Z_g of \tilde{Z}_g under the \mathbb{C} -morphism

$$\mathrm{Sh}_{gKg^{-1} \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \mathrm{Sh}_K(G, X), \quad [x, h] \mapsto [x, hg]$$

is a special subvariety of S_g . The induced morphism $\pi : \tilde{Z}_g \rightarrow Z_g$ is finite and birational. Conversely, every special subvariety of S_g arises in this way.

Remark 2.3.0.7. If the special subvariety Z_g in Fact 2.3.0.6 is normal, then by Zariski's main theorem (see, e.g., [Liu06, Cor. 4.6]), $\pi : \tilde{Z}_g \rightarrow Z_g$ is an isomorphism.

Let $S = S_g$ be a Shimura variety associated with (G, X, K) .

Lemma 2.3.0.8. *Let $Z \subset S$ be a special subvariety over $\bar{\mathbb{Q}}$. Let $\pi : \tilde{Z} \rightarrow Z$ be a finite birational morphism given by Fact 2.3.0.6. Then there is a Galois cover $f : S' \rightarrow S$ over $\bar{\mathbb{Q}}$ with $S'^L = S'^{L_\infty}$, such that for every irreducible component $Z' \subset f^{-1}(Z)$, one has $Z'^L = Z'^{L_\infty}$ and $f : Z' \rightarrow Z$ factors through an étale cover $Z' \rightarrow \tilde{Z}$.*

Proof. The Hecke isomorphism $T(g) : \mathrm{Sh}_{gKg^{-1}}(G, X) \rightarrow \mathrm{Sh}_K(G, X)$ sends $[X^+, 1]_{gKg^{-1}}$ to S_g . It keeps the special subvarieties. Thus, one may assume $g = 1$ (by replacing K with gKg^{-1}). Let $(H, X_H) \subset (G, X)$ be an irreducible Shimura subdatum inducing Z via Fact 2.3.0.6. Then the restriction

$$\pi : \tilde{Z} := [X_H^+, 1]_{K \cap H(\mathbb{A}_f)} \rightarrow Z$$

of $\mathrm{Sh}_{K \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \mathrm{Sh}_K(G, X)$ is finite and birational.

The system $\{[X_H^+, 1]_U\}$ (U running through all open subgroups of $K \cap H(\mathbb{A}_f)$) is cofinal in all the étale covers of \tilde{Z} . So there is an open subgroup $K_{0,H} \leq K$ such that the étale cover $g_0 : \tilde{Z}_0 := [X_H^+, 1]_{K_{0,H}} \rightarrow \tilde{Z}$ satisfies $g_0(\tilde{Z}_0^L) = \tilde{Z}^{L_\infty}$. Similarly, there is an open subgroup $K_1 \leq K$ such that $K_1 \cap H(\mathbb{A}_f) \subset K_{0,H}$ and the étale cover $f_1 : S_1 := ([X^+, 1]_{K_1}) \rightarrow S$ satisfies $f_1(S_1^L) = S^{L_\infty}$. By Lemma 2.2.0.7, one has $S_1^L = S_1^{L_\infty}$.

- There is an open subgroup $K_2 \leq K_1$, such that $K_2 \cap H(\mathbb{A}_f) = K_1 \cap H(\mathbb{A}_f)$ and the natural morphism $i_1 : \mathrm{Sh}_{K_2 \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \mathrm{Sh}_{K_2}(G, X)$ is a closed immersion.⁵

Indeed, by [Del71b, Prop. 1.15], there is a compact open subgroup $K_m \leq G(\mathbb{A}_f)$ containing $K_1 \cap H(\mathbb{A}_f)$, such that the morphism

$$i_2 : \mathrm{Sh}_{K \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \mathrm{Sh}_{K_m}(G, X)$$

⁵In [DJK20, p.6], it is claimed to hold for $K_2 = K_1$. This is a typo, as the first paragraph of [Mil17b, p.59] explains.

is a closed immersion. Let $K_2 = K_1 \cap K_m$. Then $K_2 \cap H(\mathbb{A}_f) = K_1 \cap H(\mathbb{A}_f)$ and $i_2 = p_{K_2, K_m} i_1$. Since $p_{K_2, K_m} : \mathrm{Sh}_{K_2}(G, X) \rightarrow \mathrm{Sh}_{K_m}(G, X)$ is separated, by magic square, i_1 is a closed immersion.

Then the morphism $\tilde{Z}_2 := [X_H^+, 1]_{K_2 \cap H} \rightarrow S_2 := [X^+, 1]_{K_2}$ is a closed immersion. The induced morphism $\pi_2 : \tilde{Z}_2 \rightarrow (f_1 f_2)^{-1}(Z)$ is a closed immersion.

- The closed immersion π_2 identifies \tilde{Z}_2 with an irreducible component of $(f_1 f_2)^{-1}(Z)$.

Since \tilde{Z}_2 is irreducible, it is contained in an irreducible component $C \subset (f_1 f_2)^{-1}(Z)$. As π is birational, by [Sta24, Tag 0BAC], there is a nonempty open subset $\tilde{U} \subset \tilde{Z}$, such that $U := \pi(\tilde{U})$ is open in Z and $\pi|_{\tilde{U}} : \tilde{U} \rightarrow U$ is an isomorphism. Consider the commutative square

$$\begin{array}{ccc} g_2^{-1}(U) & \longrightarrow & (f_1 f_2)^{-1}(U) \\ \downarrow & & \downarrow \\ \tilde{U} & \xrightarrow{\pi|_{\tilde{U}}} & U. \end{array}$$

The morphism $g_2^{-1}(\tilde{U}) \rightarrow (f_1 f_2)^{-1}(U) \xrightarrow{f_1 f_2} U$ (resp. $f_1 f_2 : (f_1 f_2)^{-1}(U) \rightarrow U$) is a base change of the étale morphism $g_2 : \tilde{Z}_2 \rightarrow Z_2$ (resp. $f_1 f_2 : S_2 \rightarrow S$), so it is étale. By [Sta24, Tag 03PC (10)], the morphism $\pi_2|_{g_2^{-1}(\tilde{U})} : g_2^{-1}(\tilde{U}) \rightarrow (f_1 f_2)^{-1}(U)$ is étale. From [Sta24, Tag 03PC (9)], $g_2^{-1}(\tilde{U})$ is an open subset of $(f_1 f_2)^{-1}(U)$, hence a nonempty open subset of C . Since C is irreducible, $g_2^{-1}(\tilde{U})$ is dense in C . Therefore, $C \subset \tilde{Z}_2$.

There is a normal, open subgroup $K' \leq K$ with $K' \subset K_2$. Let $f_3 : S' := [X^+, 1]_{K'} \rightarrow S_2$ be the induced étale cover. Since K' is normal in K , the composition $f (= f_1 f_2 f_3) : S' \rightarrow S$ is a Galois cover. Since $S_1^L = S_1^{L_\infty}$, by Lemma 2.2.0.7, one has $S'^L = S'^{L_\infty}$.

Let \tilde{Z}_3 be an irreducible component of $f_3^{-1}(\tilde{Z}_2)$. The morphism $f_3 : f_3^{-1}(\tilde{Z}_2) \rightarrow \tilde{Z}_2$ is a base change of the étale cover $f_3 : S_3 \rightarrow S_2$, so it is finite and étale. The algebraic variety \tilde{Z}_2 is smooth, so is $f_3^{-1}(\tilde{Z}_2)$. Therefore, \tilde{Z}_3 is open in $f_3^{-1}(\tilde{Z}_2)$. The morphism $g_3 : \tilde{Z}_3 \rightarrow \tilde{Z}_2$ is finite étale, and \tilde{Z}_2 is connected, so g_3 is surjective. The situation is depicted as a commutative diagram

$$\begin{array}{ccccccc}
\tilde{Z}_3 & \hookrightarrow & f_3^{-1}(\tilde{Z}_2) & \hookrightarrow & f^{-1}(Z) & \hookrightarrow & S' \\
& \searrow g_3 & \downarrow & \square & \downarrow & \square & \downarrow f_3 \\
& & \tilde{Z}_2 & \xrightarrow{\pi_2} & (f_1 f_2)^{-1}(Z) & \hookrightarrow & S_2 \\
& & \downarrow & & \downarrow & \square & \downarrow f_2 \\
& & \tilde{Z}_0 & & & & S_1 \\
& & \downarrow g_0 & & & & \downarrow f_1 \\
& & \tilde{Z} & \xrightarrow{\pi} & Z & \hookrightarrow & S
\end{array}$$

Then \tilde{Z}_3 is an étale cover of \tilde{Z} and an irreducible component of $f^{-1}(Z)$. The Galois group of the Galois cover $f : S' \rightarrow S$ permutes the irreducible components of $f^{-1}(Z)$, so they have similar properties. \square

Lemma 2.3.0.9 is used in the proof of Theorem 2.4.0.1.

Lemma 2.3.0.9. *If $S^{L\infty} \neq \emptyset$ is a finite union of special subvarieties of S , then $S^L = S$.*

Proof. Write $S^{L\infty} = \cup_{i=1}^n Z_i$ for the decomposition into irreducible components. By assumption, for every $1 \leq i \leq n$, the subvariety $Z_i \subset S$ is special. Let $\pi_i : \tilde{Z}_i \rightarrow Z_i$ be a finite birational morphism given by Fact 2.3.0.6. Let $f_i : S_i \rightarrow S$ be a Galois cover corresponding to π_i given by Lemma 2.3.0.8. There is a Galois cover $f : S' \rightarrow S$, such that for every $1 \leq i \leq n$, there is an étale cover $g_i : S' \rightarrow S_i$ with $f_i g_i = f$. Then $S'^L = S'^{L\infty}$. Hence $S'^L = (S'^L)^L = (S'^{L\infty})^L$.

1. One has $S^{L\infty} \subset \cup_{i=1}^n \pi_i(\tilde{Z}_i^{L\infty})$.

Indeed, one has $f(S'^L) = S^{L\infty}$. For every irreducible component $C \subset S'^L$, the subset $f(C)$ of $S^{L\infty}$ is irreducible. Then there is $1 \leq i \leq n$ with $f(C) \subset Z_i$. Thus, $g_i(C)$ is an irreducible subset of $f_i^{-1}(Z_i)$. There is an irreducible component $Z' \subset f_i^{-1}(Z_i)$ containing $g_i(C)$. By Lemma 2.3.0.8, the morphism $f_i : Z' \rightarrow Z_i$ factors through an étale cover $Z' \rightarrow \tilde{Z}_i$. Therefore, Z' and $g_i^{-1}(Z')$ are smooth. One has

$$g_i^{-1}(Z') \subset f^{-1}(Z_i) \subset f^{-1}(S^{L\infty}) \stackrel{(a)}{=} S'^L,$$

where (a) uses Lemma 2.2.0.7. Then C is an irreducible component of $g_i^{-1}(Z')$, hence an étale cover of Z' . One has $f(C^L) \subset \pi_i(\tilde{Z}_i^{L\infty})$. Thus, 1 is proved.

2. One has $\tilde{Z}_1^{L\infty} = \tilde{Z}_1$.

From 1, one has $Z_1 \subset \bigcup_{i=1}^n \pi_i(\tilde{Z}_i^{L_\infty})$. Since Z_1 is irreducible, there is $1 \leq j \leq n$ with $Z_1 \subset \pi_j(\tilde{Z}_j^{L_\infty}) \subset Z_j$. As Z_1 is an irreducible component of S^{L_∞} , one has $j = 1$. Then $\dim \tilde{Z}_1^{L_\infty} \geq \dim Z_1 = \dim \tilde{Z}_1$. The irreducibility of \tilde{Z}_1 proves 2.

3. For every $q \in G(\mathbb{Q})_+$, one has $T_q Z_1 \subset S^L$.

Let $(H, X_H) \subset (G, X)$ be an irreducible Shimura subdatum inducing Z_1 via Fact 2.3.0.6. Then $\tilde{Z}_1 = [X_H^+, 1]_{K \cap H(\mathbb{A}_f)}$. For every irreducible component $Z_q \subset \alpha_q^{-1}(Z_1)$, there is an irreducible component \tilde{Z}_q of $\text{Sh}_{K \cap q^{-1}Kq \cap H(\mathbb{A}_f)}(H, X_H)$ with the following properties:

- The morphism $\text{Sh}_{K \cap q^{-1}Kq \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \text{Sh}_{K \cap H(\mathbb{A}_f)}(H, X_H)$ restricts to an étale cover $\alpha'_q : \tilde{Z}_q \rightarrow \tilde{Z}_1$.
- The image of \tilde{Z}_q under the morphism $\text{Sh}_{K \cap q^{-1}Kq \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \text{Sh}_{K \cap q^{-1}Kq}(G, X)$ is Z_q .

Conjugating by q gives another irreducible Shimura subdatum $(qHq^{-1}, q \cdot X_H) \subset (G, X)$, and a morphism of Shimura data $(H, X_H) \rightarrow (qHq^{-1}, q \cdot X_H)$. Let \tilde{Z}'_q be the image of \tilde{Z}_q under the induced morphism $\text{Sh}_{K \cap q^{-1}Kq \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \text{Sh}_{K \cap qH(\mathbb{A}_f)q^{-1}}(qHq^{-1}, q \cdot X_H)$. Then \tilde{Z}'_q is an irreducible component of

$$\text{Sh}_{K \cap qH(\mathbb{A}_f)q^{-1}}(qHq^{-1}, q \cdot X_H),$$

and the restriction $\beta'_q : \tilde{Z}_q \rightarrow \tilde{Z}'_q$ is an étale cover. By Fact 2.3.0.6, the morphism $\text{Sh}_{K \cap qH(\mathbb{A}_f)q^{-1}}(qHq^{-1}, q \cdot X_H) \rightarrow \text{Sh}_K(G, X)$ restricts to a finite birational morphism $\pi'_q : \tilde{Z}'_q \rightarrow \beta_q(Z_q)$. Consider the commutative diagram

$$\begin{array}{ccccc} & & \tilde{Z}_q & & \\ & \swarrow & \downarrow & \searrow & \\ \tilde{Z}_q & & Z_q & & \tilde{Z}'_q \\ \downarrow \pi_1 & \swarrow \alpha_q & & \searrow \beta_q & \downarrow \pi'_q \\ Z_1 & & & & \beta_q(Z_q). \end{array}$$

From 2 and Lemma 2.2.0.7, one has $\tilde{Z}'_q = \tilde{Z}_q^{L_\infty} = \tilde{Z}_q^{L'}$. Then $\beta_q(Z_q) = \pi'_q(\tilde{Z}'_q) = \pi'_q(\tilde{Z}_q^{L'}) \stackrel{(a)}{\subset} \beta_q(Z_q)^L$, where (a) uses Assumption 2.2.0.1 5. By Assumption 2.2.0.1 2, one has $\beta_q(\tilde{Z}_q) \subset S^L$. Thus, 3 is proved.

Since Z_1 is a special subvariety of S , from [KUY18, Lem. 2.5], Z_1 contains a special point z . By [LZ19, Rk. 2.7], $\{T_q z\}_{q \in G(\mathbb{Q})_+}$ is dense in the complex manifold $S(\mathbb{C})$. By 3, the Zariski closed subset $S^L \subset S$ contains $\{T_q z\}_{q \in G(\mathbb{Q})_+}$. Hence $S^L = S$. \square

2.4 Ullmo-Yafaev alternative principle

In Theorem 2.4.0.1, we show that an alternative principle results from Assumption 2.2.0.1. Let $S = S_g = (gKg^{-1} \cap G(\mathbb{Q})_+) \backslash X^+$ be a Shimura variety associated with (G, X, K) .

Theorem 2.4.0.1 (Ullmo-Yafaev alternative). *Either $S^{L\infty} = \emptyset$ or $S^{L\infty} = S$.*

Proof. By Hecke isomorphisms, one may assume $g = 1$ and $S = [X^+, 1]_K$. By Lemma 2.2.0.7, one may replace S by an étale cover induced by an open subgroup of K . One may thereby assume $S^L = S^{L\infty} \neq \emptyset$. For every irreducible component $Z \subset S^L$, by Assumption 2.2.0.1 1 (resp. Lemma 2.2.0.5 2), one has $\dim(Z) > 0$ (resp. $Z^L = Z$).

1. The subvariety $Z \subset S$ is special.

Let $S_M \subset S$ be the smallest special subvariety containing Z . From Fact 2.3.0.6, there is a Shimura subdatum $(H, X_H) \subset (G, X)$, such that the restriction $\pi : \tilde{S}_M := [X_H^+, 1]_{K \cap H(\mathbb{A}_f)} \rightarrow S_M$ of $\text{Sh}_{K \cap H(\mathbb{A}_f)}(H, X_H) \rightarrow \text{Sh}_K(G, X)$ is finite birational.

Take a Galois cover $f : S' \rightarrow S$ given by Lemma 2.3.0.8 for the special subvariety $S_M \subset S$. Since f is finite surjective, there is an irreducible component $T \subset f^{-1}(Z)$ with $f(T) = Z$.

Since $Z \subset S^L$ is an irreducible component, T is an irreducible component of

$$f^{-1}(S^L) = f^{-1}(S^{L\infty}) \stackrel{(a)}{=} S'^{L\infty} \stackrel{(b)}{=} S'^L.$$

Here (a) and (b) use Lemma 2.2.0.7. Then by Lemma 2.2.0.5 2, one has $T^L = T$. There is an irreducible component $S'_M \subset f^{-1}(S_M)$ containing T .

By Lemma 2.3.0.8, one has $S'^L_M = S'^{L\infty}_M$, and $f : S'_M \rightarrow S_M$ factors through an étale cover

$$g : S'_M \rightarrow \tilde{S}_M.$$

2. One has $g(T) \subset \tilde{S}_M^{L\infty}$.

Consider the commutative diagram

$$\begin{array}{ccccccc} T & \hookrightarrow & S'_M & \hookrightarrow & f^{-1}(S_M) & \hookrightarrow & S' \\ & & \downarrow g & & \downarrow & \square & \downarrow f \\ & & \tilde{S}_M & \xrightarrow{\pi} & S_M & \hookrightarrow & S. \end{array}$$

One has

$$T = T^L \subset S'^L_M = S'^{L\infty}_M.$$

Hence $g(T) \subset g(S'^{L\infty}_M) = \tilde{S}_M^{L\infty}$. Thus, 2 is proved.

3. The nonempty, irreducible, closed subset $g(T) \subset \tilde{S}_M^{L\infty}$ is Hodge generic in \tilde{S}_M .

Since π is finite surjective, there is an irreducible component $\tilde{Z} \subset \pi^{-1}(Z)$ with $\pi(\tilde{Z}) = Z$. For every special subvariety $V \subset \tilde{S}_M$ containing $g(T)$, by [KY14, p.879], $\pi(V) \subset S$ is a special subvariety containing $\pi g(T) = f(T) = Z$. Hence $\pi(V) = S_M$. Therefore, $\dim V \geq \dim S_M = \dim \tilde{S}_M$. Since \tilde{S}_M is irreducible, one has $V = \tilde{S}_M$. Thus, 3 is proved.

By 2, 3 and Lemma 2.4.0.2, one has $\tilde{S}_M = \tilde{S}_M^{L\infty} = \tilde{S}_M^L$. One has $S_M = \pi(\tilde{S}_M) = \pi(\tilde{S}_M^L) \stackrel{(a)}{\subset} S_M^L \subset S^L$, where (a) uses Assumption 2.2.0.1 5. Since Z is an irreducible component of S^L and S_M is irreducible, one has $Z = S_M$. Thus, 1 is proved.

By 1, the locus S^L is a finite union of special subvarieties. From Lemma 2.3.0.9, one has $S^{L\infty} = S$. \square

Lemma 2.4.0.2 (Ullmo-Yafaev). *Let $S = [X^+, 1]_K \subset \text{Sh}_K(G, X)$. If $S^{L\infty}$ contains a nonempty, irreducible closed subset that is Hodge generic in S , then $S^{L\infty} = S$.*

Proof. For every $q \in G(\mathbb{Q})^+$, by Lemma 2.2.0.7, one has $T_q S^{L\infty} = \beta_q(\alpha_q^{-1}(S^{L\infty})) = \beta_q(S_q^{L\infty}) = S^{L\infty}$. Write $S^{L\infty} = U_1 \cup U_2$, where U_1 is the union of irreducible components of $S^{L\infty}$ that are Hodge generic in S , and U_2 is the union of the remaining irreducible components. By Remark 2.2.0.6 and assumption, one has $\dim U_1 > 0$.

Let C be an irreducible component of $T_q U_2$. Then there is an irreducible subvariety $C_q \subset S_q$ with $\beta_q(C_q) = C$ and $\alpha_q(C_q) \subset U_2$. Then $\alpha_q(C_q)$ is not Hodge generic in S . Thus, there is a strict, special subvariety $V \subset S$ containing $\alpha_q(C_q)$. Then $C \subset T_q(\alpha_q(C_q)) \subset T_q V$. There is an irreducible component $W \subset T_q V$ containing C . By [LZ19, Remark 2.7], W is a special subvariety of S . Since $\dim W \leq \dim V < \dim S$, the subvariety $C \subset S$ is not Hodge generic. As every irreducible component of $T_q U_2$ is not Hodge generic in S , and $U_1 \subset T_q S^{L\infty} = T_q U_1 \cup T_q U_2$, one has $U_1 \subset T_q U_1$. By $\dim U_1 > 0$ and [UY10, Thm. 1.2], one has $U_1 = S$ and $S^{L\infty} = S$. \square

Corollary 2.4.0.3 ([UY10, Thm. 1.1]). *If a Shimura variety S over $\bar{\mathbb{Q}}$ is of sufficiently high level, then either $S^L = \emptyset$ or $S^L = S$.*

Proof. As the level is high, one has $S^L = S^{L\infty}$. The result follows from Theorem 2.4.0.1. \square

2.5 “All or nothing” principle for integral points

We define an locus concerning integral points, analogous to the Lang locus concerning rational points. We verify Assumption 2.2.0.1 for this locus.

Then an alternative principle follows.

Let X be an integral algebraic variety over $\bar{\mathbb{Q}}$. As in Example 2.2.0.2, there is a number field $F \subset \bar{\mathbb{Q}}$, an integral algebraic variety X_F over F and an isomorphism $X_F \otimes_F \bar{\mathbb{Q}} \rightarrow X$ over $\bar{\mathbb{Q}}$. For every finite set Σ of places of F including all archimedean ones, let $O_{F,\Sigma}$ be the ring of Σ -integers. When Σ is sufficiently large, there exists an integral scheme \mathcal{X} that is finite type and separated over $O_{F,\Sigma}$, whose generic fiber is X_F . (From [Har77, III, Prop. 9.7], \mathcal{X} is flat over $O_{F,\Sigma}$.) We call \mathcal{X} an integral model for X relative to (F, Σ) . By a finite extension $(M, \Omega)/(F, \Sigma)$, we mean a finite extension M/F together with a finite set Ω of places of M containing all the places above Σ .

For every $(M, \Omega)/(F, \Sigma)$, let $X(\mathcal{X}, M, \Omega)$ be the image of the injection

$$\mathcal{X}(O_{M,\Omega}) \rightarrow X(\bar{\mathbb{Q}}), \quad x \mapsto x|_{\text{Spec } \bar{\mathbb{Q}}}.$$

Definition 2.5.0.1. Let \mathcal{X}^I be the Zariski closure of

$$\bigcup_{(M,\Omega)/(F,\Sigma)} \overline{X(\mathcal{X}, M, \Omega)}^{>0}$$

inside X , where (M, Ω) runs through all finite extensions of (F, Σ) . We call \mathcal{X}^I the *integral Lang locus* of X relative to (\mathcal{X}, F, Σ) .

The integral Lang locus \mathcal{X}^I is a subvariety of the Lang locus of X .

Lemma 2.5.0.2. *Given models \mathcal{X}_i over O_{F_i, Σ_i} ($i = 1, 2$) for X , one has $\mathcal{X}_1^I = \mathcal{X}_2^I$.*

Proof. By [Gro66, Cor. 8.8.2.5], there is a common finite extension (F_3, Σ_3) of (F_i, Σ_i) ($i = 1, 2$), such that there is an O_{F_3, Σ_3} -isomorphism

$$\mathcal{X}_1 \otimes_{O_{F_1, \Sigma_1}} O_{F_3, \Sigma_3} \rightarrow \mathcal{X}_2 \otimes_{O_{F_2, \Sigma_2}} O_{F_3, \Sigma_3}$$

extending the isomorphism between the generic fibers. For every finite extension $(M_1, \Omega_1)/(F_1, \Sigma_1)$, there is a common finite extension (M_2, Ω_2) of (F_3, Σ_3) and (M_1, Ω_1) . Then

$$\mathcal{X}_1(O_{M_1, \Omega_1}) \subset \mathcal{X}_1(O_{M_2, \Omega_2}) = \mathcal{X}_2(O_{M_2, \Omega_2}),$$

so $X(\mathcal{X}_1, M_1, \Omega_1) \subset X(\mathcal{X}_2, M_2, \Omega_2)$. Therefore,

$$\overline{X(\mathcal{X}_1, M_1, \Omega_1)}^{>0} \subset \overline{X(\mathcal{X}_2, M_2, \Omega_2)}^{>0} \subset \mathcal{X}_2^I.$$

Hence $\mathcal{X}_1^I \subset \mathcal{X}_2^I$. The other inclusion follows by symmetry. \square

By Lemma 2.5.0.2, one may use the notation X^I for \mathcal{X}^I and call it *integral Lang locus* of X . We extend the definition to reducible algebraic varieties as in Section 2.2.

Remark 2.5.0.3. Assume that X is proper over $\bar{\mathbb{Q}}$. Then there is an integral model (\mathcal{X}, F, Σ) for X , such that \mathcal{X} is proper over $O_{F, \Sigma}$. By [Poo17, Thm. 3.2.13 (ii)], X^I coincides with the Lang locus of X .

Definition 2.5.0.4. [Ull04, Déf. 2.3] An integral algebraic variety X over $\bar{\mathbb{Q}}$ is called arithmetically hyperbolic if $X^I = \emptyset$.

An integral algebraic variety X over $\bar{\mathbb{Q}}$ is arithmetically hyperbolic if and only if for one (hence for every by Lemma 2.5.0.2) model (\mathcal{X}, F, Σ) , the set of integral points $\mathcal{X}(O_{M, \Omega})$ is finite for every finite extension $(M, \Omega)/(F, \Sigma)$ (so [Ull04, Lem. 2.4] follows from Lemma 2.5.0.2).

Example 2.5.0.5. Let $X = \mathbf{P}^1 \setminus \{0, 1, \infty\} = Y(2)$ be a modular curve over $\bar{\mathbb{Q}}$. Its Baily-Borel compactification is $X^* = \mathbf{P}^1$, and the Lang locus of X is full. By the Siegel-Mahler theorem (see, e.g., [HS00, Thm. D.8.1]), X is arithmetically hyperbolic.

A complex analytic space is called *Kobayashi hyperbolic*, if its Kobayashi pseudo-distance (in the sense of [Kob98, p.50]) is a metric. Every Kobayashi hyperbolic, complex analytic space is Brody hyperbolic. Conversely, Brody [Bro78, p.213] proves that every compact, Brody hyperbolic complex analytic space is Kobayashi hyperbolic. In view of Remark 2.5.0.3, Conjecture 2.5.0.6 implies Conjecture 2.1.0.1.

Conjecture 2.5.0.6 ([Lan91, IX, Conjecture 5.1], [Ull04, Conjecture 2.5]). *Let X be a quasi-projective, integral algebraic variety over $\bar{\mathbb{Q}}$. If the complex analytic space $X(\mathbb{C})$ is Kobayashi hyperbolic, then X is arithmetically hyperbolic.*

Fact 2.5.0.7 is an evidence of Conjecture 2.5.0.6. It relies on Faltings's solution [Fal83, Satz 6] to Shafarevich's conjecture.

Fact 2.5.0.7 ([Ull04, Thm. 3.2 (a)]). *Let (G, X) be an adjoint Shimura datum of abelian type (in the sense of [Ull04, p.4118]). Let $K \leq G(\mathbb{A}_f)$ be a neat compact open subgroup. Then every irreducible component of $\text{Sh}_K(G, X)_{\bar{\mathbb{Q}}}$ is arithmetically hyperbolic.*⁶

We prove that an alternative principle holds for integral points on Shimura varieties, by checking Assumption 2.2.0.1. Since an irreducible component of X^I with dimension 0 is an isolated point, Assumption 2.2.0.1 1 holds. Lemma 2.5.0.8 verifies Assumptions 2.2.0.1 2, 3 and 5.

Lemma 2.5.0.8. *Let $f : Z_1 \rightarrow Z_2$ be a morphism of integral algebraic varieties over $\bar{\mathbb{Q}}$. If f has finite geometric fibers, then $f(Z_1^I) \subset Z_2^I$.*

⁶By [Moo98b, 2.17], the model over $\bar{\mathbb{Q}}$ defined by Faltings [Fal82] (used in [Ull04, Thm. 3.2 (a)]) is the scalar extension of the canonical model along $E(G, X) \rightarrow \bar{\mathbb{Q}}$.

Proof. One may choose a number field F , a finite set Σ of places of F containing all the archimedean ones, a model \mathcal{Z}_i over $O_{F,\Sigma}$ for Z_i ($i = 1, 2$) and an $O_{F,\Sigma}$ -morphism $f' : \mathcal{Z}_1 \rightarrow \mathcal{Z}_2$ whose base change to F is f . For every finite extension $(M, \Omega)/(F, \Sigma)$, one has $f'(\mathcal{Z}_1(O_{M,\Omega})) \subset \mathcal{Z}_2(O_{M,\Omega})$, so $f(Z_1(\mathcal{Z}_1, M, \Omega)) \subset Z_2(\mathcal{Z}_2, M, \Omega)$. Hence

$$f(\overline{Z_1(\mathcal{Z}_1, M, \Omega)}) \subset \overline{Z_2(\mathcal{Z}_2, M, \Omega)}.$$

Let $C \subset \overline{Z_1(\mathcal{Z}_1, M, \Omega)}$ be an irreducible component of positive dimension. Then $f(C)$ is irreducible but not a singleton. (For otherwise, C is a finite set by assumption, which is a contradiction). Hence

$$f(C) \subset \overline{Z_2(\mathcal{Z}_2, M, \Omega)}^{>0} \subset Z_2^I.$$

Therefore, $f(\overline{Z_1(\mathcal{Z}_1, M, \Omega)}^{>0}) \subset Z_2^I$ and $f(Z_1^I) \subset Z_2^I$. \square

Corollary 2.5.0.9 ([Ull04, Prop. 2.6]). *A locally closed subvariety of an arithmetically hyperbolic variety is also arithmetically hyperbolic.*

Proof. It follows from Lemma 2.5.0.8. \square

Lemma 2.5.0.10 verifies Assumption 2.2.0.1 4 for integral Lang loci.

Lemma 2.5.0.10. *Let X be an integral algebraic variety over $\bar{\mathbb{Q}}$. Then $X^I \subset (X^I)^I$.*

Proof. Write $X^I = \cup_{i=1}^n Y_i$ as the union of irreducible components. Take a model (\mathcal{X}, F, Σ) for X . Let \mathcal{Y}_i be the scheme-theoretic image of the composition $Y_i \rightarrow X \rightarrow \mathcal{X}$, which is model of Y_i relative to (F, Σ) . For every finite extension $(M, \Omega)/(F, \Sigma)$, the Zariski closed subset $\overline{X(\mathcal{X}, M, \Omega)} \subset X$ is the disjoint union of $\overline{X(\mathcal{X}, M, \Omega)}^{>0}$ with a finite set $\{p_1, \dots, p_t\} \subset X(\bar{\mathbb{Q}})$.

Consider $x \in \mathcal{X}(O_{M,\Omega})$, i.e., a section $x : \text{Spec}(O_{M,\Omega}) \rightarrow \mathcal{X}$ to the structure morphism $\mathcal{X} \rightarrow \text{Spec}(O_{M,\Omega})$. If $x|_{\text{Spec } \bar{\mathbb{Q}}} \notin \{p_1, \dots, p_t\}$, then

$$x|_{\text{Spec } \bar{\mathbb{Q}}} \in \overline{X(\mathcal{X}, M, \Omega)}^{>0} \subset X^I.$$

Thus, there exists an index $1 \leq i \leq n$ with $x|_{\text{Spec } \bar{\mathbb{Q}}} \in Y_i$. Since \mathcal{Y}_i is Zariski closed in \mathcal{X} , the section x factors through \mathcal{Y}_i , i.e., $x \in \mathcal{Y}_i(O_{M,\Omega})$. Therefore,

$$X(\mathcal{X}, M, \Omega) \subset \cup_{i=1}^n \mathcal{Y}_i(\mathcal{Y}_i, M, \Omega) \cup \{p_1, \dots, p_t\}.$$

Then

$$\overline{X(\mathcal{X}, M, \Omega)}^{>0} \subset \cup_{i=1}^n \overline{\mathcal{Y}_i(\mathcal{Y}_i, M, \Omega)}^{>0} \subset \cup_{i=1}^n \mathcal{Y}_i^I = (X^I)^I,$$

so $X^I \subset (X^I)^I$. \square

Lemma 2.5.0.11 implies [Ull04, Prop. 2.8].

Lemma 2.5.0.11 (Chevalley-Weil). *If $f : X \rightarrow Y$ is an étale cover over $\bar{\mathbb{Q}}$, then $f(X^I) = Y^I$. In particular, $X^{I_\infty} = X^I$. Moreover, $X^I = X$ (resp. $X^I = \emptyset$) is equivalent to $Y^I = Y$ (resp. $Y^I = \emptyset$).*

Proof. By Lemma 2.5.0.8, one has $f(X^I) \subset Y^I$. There is a number field F , a finite set Σ of places of F containing all the archimedean ones, and a finite étale $O_{F,\Sigma}$ -morphism $f' : \mathcal{X} \rightarrow \mathcal{Y}$ between models whose base change to the generic fiber recovers f .

For every finite extension $(M, \Omega)/(F, \Sigma)$, by the Chevalley-Weil theorem (see, e.g., [SBW89, p.50]), there is a finite extension $(M', \Omega')/(M, \Omega)$ with $Y(\mathcal{Y}, M, \Omega) \subset f(X(\mathcal{X}, M', \Omega'))$. Since zero dimensional schemes are discrete,

$$\overline{Y(\mathcal{Y}, M, \Omega)}^{>0} \subset f(\overline{X(\mathcal{X}, M', \Omega')})^{>0} \subset f(X^I).$$

Hence $Y^I \subset f(X^I)$. □

Theorem 2.5.0.12. *The integral Lang locus of a Shimura variety S is either empty or whole S .*

Proof. By Lemmas 2.5.0.8 and 2.5.0.10, the formation of the integral Lang locus $(\cdot)^I$ satisfies Assumption 2.2.0.1. The result is a combination of Theorem 2.4.0.1 and Lemma 2.5.0.11. □

Chapter 3

Normality of monodromy group in generic Tannakian group

3.1 Introduction

3.1.1 Background

Constructing local systems (or ℓ -adic lisse sheaves) with a prescribed monodromy group is an important problem having a long history.

In positive characteristics, Katz and his collaborators exhibit local systems whose monodromy groups are the simple algebraic group G_2 ([Kat88, 11.8]), $2.J_2$ ([KRL19]), the finite symplectic groups ([KT19b]), the special unitary groups ([KT19a]), *etc.* In particular, the exceptional Lie groups appear unexpectedly in algebraic geometry.

In characteristic 0, such constructions help to understand Galois groups of number fields. Dettweiler and Reiter [DR10] prove the existence of a local system on $\mathbf{P}_{\mathbb{Q}}^1 \setminus \{0, 1, \infty\}$ whose monodromy group is G_2 . It produces a motivic Galois representations with image dense in G_2 . Their proof relies on Katz's middle convolution of perverse sheaves. Yun [Yun14] constructs local systems with some other exceptional groups as monodromy groups. As applications, he answers a long standing question of Serre, and solves new cases of the inverse Galois problem. His construction uses the geometric Langlands correspondence.

A new proof of Mordell's conjecture [IV20], and its potential generalization to higher dimensional varieties over number fields, rely on the existence of local systems with big monodromy over the variety in question. Lawrence and Sawin [LS20] use this technique to prove Shafarevich's conjecture for hypersurfaces in abelian varieties. Krämer and Maculan [KM23] apply roughly the same strategy to obtain an arithmetic finiteness result for

very irregular varieties of dimension less than half the dimension of their Albanese variety. In both cases, the construction of local systems uses perverse sheaves.

In [LS20], that construction rests on comparing the monodromy group with the Tannakian group from Krämer-Weissauer's convolution theory [KW15b]. As [JKLM23, p.4] comments, this comparison is similar to the study of monodromy groups *via* Mumford-Tate groups in [And92].

We briefly outline their argument. On an abelian variety A , a quotient of the abelian category $\text{Perv}(A)$ (of perverse sheaves) is a Tannakian category under sheaf convolution. Let X be an irreducible algebraic variety with generic point η . Let K be a universally locally acyclic, relative perverse sheaf on the constant abelian scheme $p_X : A \times X \rightarrow X$ (intuitively, a family of perverse sheaves on A parameterized by X). The Tannakian group $G(K|_{A_{\bar{\eta}}})$ of $K|_{A_{\bar{\eta}}} \in \text{Perv}(A_{\bar{\eta}})$ is normal in the Tannakian group $G(K|_{A_{\eta}})$ of $K|_{A_{\eta}} \in \text{Perv}(A_{\eta})$ ([LS20, Lem. 3.7], [JKLM23, Thm. 4.3]). This normality is used to prove that for most character sheaves L_{χ} on A , the monodromy groups $\text{Mon}(K \otimes p_A^* L_{\chi})$ contain $G(K|_{A_{\bar{\eta}}})$. Then Lawrence-Venkatesh's machinery works for these twists $K \otimes p_A^* L_{\chi}$.

3.1.2 Statements

In the main result (Theorem 3.1.2.2), we prove that the generic Tannakian group of a semisimple, relative perverse sheaf is reductive. Moreover, for many characters, the monodromy group is a normal subgroup of this reductive group. This normality puts further restriction on the monodromy group. Using Krämer's method ([Krä22, Thm. 6.2.1]), Lawrence and Sawin [LS20, Lem. 4.6] even show that the geometric generic Tannakian group is simple.

Setting 3.1.2.1. Let k be an algebraically closed field of characteristic 0. Let X be an integral algebraic variety over k with generic point η . Let A be an abelian variety over k . Denote by $p_X : A \times X \rightarrow X$ and $p_A : A \times X \rightarrow A$ the projections.

Let ℓ be a prime number. Let $\bar{\mathbb{Q}}_{\ell}$ be an algebraic closure of \mathbb{Q}_{ℓ} . Let $D_c^b(A \times X)$ be the triangulated category of bounded constructible $\bar{\mathbb{Q}}_{\ell}$ -sheaves on $A \times X$. Let $\text{Perv}^{\text{ULA}}(A \times X/X) \subset D_c^b(A \times X)$ be the full subcategory of p_X -universally locally acyclic (ULA, Definition 3.2.2.1) relative perverse sheaves (Definition 3.2.3.2). It is an abelian category. Let $\pi_1^{\text{ét}}(A)$ be the étale fundamental group of A based at the geometric origin point. For every character $\chi : \pi_1^{\text{ét}}(A) \rightarrow \bar{\mathbb{Q}}_{\ell}^{\times}$, let $\chi_{\eta} : \pi_1^{\text{ét}}(A_{\eta}) \rightarrow \bar{\mathbb{Q}}_{\ell}^{\times}$ be the pullback of χ along $(p_A|_{A_{\eta}}) : \pi_1^{\text{ét}}(A_{\eta}) \rightarrow \pi_1^{\text{ét}}(A)$. Fix $K \in \text{Perv}^{\text{ULA}}(A \times X/X)$ which is a semisimple object of $D_c^b(A \times X)$ (in the sense of Definition 3.2.1.3). Let $\text{Mon}(K, \chi_{\eta})$ be the corresponding monodromy group (Definition 3.4.3.4).

Let $G_{\omega_\chi}(K|_{A_\eta})$ be the Tannakian monodromy group (Definition 3.4.2.1) of $K|_{A_\eta}$, referred to as the *generic Tannakian group*.

Theorem 3.1.2.2 (Theorems 3.5.1.1, 3.5.3.1). *Assume $\dim A > 0$. Then there are uncountably many characters $\chi : \pi_1^{\text{ét}}(A) \rightarrow \bar{\mathbb{Q}}_\ell^\times$, such that $G_{\omega_\chi}(K|_{A_\eta})$ is a well-defined reductive group. It contains $\text{Mon}(K, \chi_\eta)$ as a closed, reductive, normal subgroup.*

The line of the proof of Theorem 3.1.2.2 is similar to that of André's normality theorem [And92, Thm. 1]. André proves that for a polarizable good variation of mixed Hodge structure, the connected monodromy group outside a meager locus is normal in the derived Mumford-Tate group. As [And92, p.10] explains, the normality is a consequence of the theorem of the fixed part due to Griffiths-Schmidt-Steenbrink-Zucker. In our case, an analog of the theorem of the fixed part is Theorem 3.1.2.3.

Let $\mathcal{C}(A)_\ell$ be the cotorus parameterizing pro- ℓ characters of $\pi_1^{\text{ét}}(A)$ (Definition 3.3.1.3). For every $\chi_{\ell'} \in \mathcal{C}(A)_{\ell'}$ and every $\chi_\ell \in \mathcal{C}(A)_\ell$, set $\chi = \chi_{\ell'} \chi_\ell$.

Theorem 3.1.2.3 (Theorem 3.5.2.1). *Assume that X is smooth. Then there is a subobject $K^0 \subset K$ in $\text{Perv}^{\text{ULA}}(A \times X/X)$ with the following property: For every character $\chi_{\ell'} : \pi_1^{\text{ét}}(A) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ of finite order prime to ℓ , there is a nonempty Zariski open subset $U \subset \mathcal{C}(A)_\ell$, such that for every $\chi_\ell \in U$, one has*

$$H^0(A_{\bar{\eta}}, K^0|_{A_{\bar{\eta}}} \otimes^L L_{\chi_\eta}) = H^0(A_{\bar{\eta}}, K|_{A_{\bar{\eta}}} \otimes^L L_{\chi_\eta})^{\Gamma_{k(\eta)}}.$$

The proof of Theorem 3.1.2.3 uses the projection $p_A : A \times X \rightarrow A$, which restricts our results to constant abelian schemes. We leave the question whether Theorem 3.1.2.2 has an analog for relative perverse sheaves on an arbitrary (non-constant) abelian scheme.

Notation and conventions

An object of an abelian category is *semisimple* if it is the direct sum of finitely many simple objects. An abelian category is *semisimple* if every object is semisimple. For a field k , its absolute Galois group is denoted by Γ_k . An algebraic variety means a scheme of finite type and separated over k . A linear algebraic group is *reductive*, if its identity component is reductive (in the sense of [Mil17a, 6.46, p.135]). For a topological group, $\bar{\mathbb{Q}}_\ell$ -characters are assumed to be continuous. For an irreducible algebraic variety X (on which ℓ is invertible) and a $\bar{\mathbb{Q}}_\ell$ -character χ of its étale fundamental group $\pi_1^{\text{ét}}(X)$, let L_χ be the corresponding rank one lisse $\bar{\mathbb{Q}}_\ell$ -sheaf on X .

3.2 Recollections on constructible sheaves

No originality is claimed in Section 3.2. Let k be a field. Let ℓ be a prime number invertible in k . Fix an algebraic closure \bar{k} of k . For every algebraic

variety X over k , denote by $D_c^b(X) := D_c^b(X, \bar{\mathbb{Q}}_\ell)$ the triangulated category of complexes of $\bar{\mathbb{Q}}_\ell$ -sheaves on X with bounded constructible cohomologies defined in [BBDG82, p.74]. Let $\mathbb{D}_X : D_c^b(X) \rightarrow D_c^b(X)^{\text{op}}$ be the Verdier duality functor. The heart of the standard t-structure on $D_c^b(X)$ is denoted by $\text{Cons}(X)$, which is the category of constructible $\bar{\mathbb{Q}}_\ell$ -sheaves on X . For $F \in \text{Cons}(X)$, set $\text{Supp } F := \{x \in X \mid F_x \neq 0\}$ to be its support. Then $\text{Supp } F$ is a quasi-constructible subset of X in the sense of [Gro66, 10.1.1]. Let $\text{Loc}(X) \subset \text{Cons}(X)$ be the full subcategory of lisse $\bar{\mathbb{Q}}_\ell$ -sheaves on X . For every integer n , let $\mathcal{H}^n : D_c^b(X) \rightarrow \text{Cons}(X)$ be the functor taking the n -th cohomology sheaf.

For every subset $S \subset X$, let \bar{S} be its Zariski closure. Let ${}^pD^{\leq 0}(X) \subset D_c^b(X)$ be the full subcategory of objects K with $\dim \bar{\text{Supp}} \mathcal{H}^n K \leq -n$ for every integer n . Let ${}^pD^{\geq 0}(X) \subset D_c^b(X)$ be the full subcategory of objects K with $\mathbb{D}_X K \in {}^pD^{\leq 0}(X)$. Then $({}^pD^{\leq 0}(X), {}^pD^{\geq 0}(X))$ defines the (absolute) perverse t-structure on $D_c^b(X)$, whose heart is denoted by $\text{Perv}(X)$. The functor \mathbb{D}_X interchanges ${}^pD^{\leq 0}(X)$ and ${}^pD^{\geq 0}(X)$. For every integer n , let ${}^p\mathcal{H}^n : D_c^b(X) \rightarrow \text{Perv}(X)$ be the functor taking the n -th perverse cohomology sheaf. For a morphism $f : X' \rightarrow X$ of schemes and $K \in D_c^b(X)$, set $K|_{X'} := f^*K$.

3.2.1 Basics

Fact 3.2.1.1 (Projection formula, [FK88, Rk. (2), p.100], [Sta24, Tag 0F10 (1)]). *Let $f : X \rightarrow Y$ be a morphism of algebraic varieties over \bar{k} . Let $L \in D_c^b(Y)$ be an object with $\mathcal{H}^n L \in \text{Loc}(X)$ for every integer n . Then there is a natural isomorphism $(Rf_* -) \otimes^L L \rightarrow Rf_*(- \otimes^L f^*L)$ of functors $D_c^b(X) \rightarrow D_c^b(Y)$.*

Let X be an algebraic variety over k .

Fact 3.2.1.2 ([FK88, Prop. 12.10]). *For every $F \in \text{Cons}(X)$, there is a nonempty Zariski open subset $U \subset X$ with $F|_U \in \text{Loc}(U)$.*

Definition 3.2.1.3 ([BC18, Def. 78]). An object $K \in D_c^b(X)$ is called semisimple if it is isomorphic to a finite direct sum of degree shifts of semisimple objects of $\text{Perv}(X)$.

If $K \in D_c^b(X)$ is semisimple, then it is isomorphic to $\bigoplus_{n \in \mathbb{Z}} {}^p\mathcal{H}^n(K)[-n]$ in $D_c^b(X)$, and each ${}^p\mathcal{H}^n(K)$ is a semisimple object of $\text{Perv}(X)$. A degree shift of a semisimple object of $D_c^b(X)$ is still semisimple.

Example 3.2.1.4. Every perverse cohomology sheaf of a semisimple object of $D_c^b(X)$ is semisimple. By contrast, its cohomology sheaves may be no longer semisimple in $D_c^b(X)$.

Consider $k = \mathbb{C}$ and $X = A^1$. Let $j : U = A^1 \setminus \{0, 1\} \rightarrow X$ be the inclusion. Then the topological fundamental group $\pi_1^{\text{top}}(U^{\text{an}}, -1)$ is the free

group generated by two loops a and b , surrounding 0 and 1 respectively. There is a unique morphism

$$\pi_1^{\text{top}}(U^{\text{an}}, -1) \rightarrow \text{SL}_2(\mathbb{Z}) \quad (3.1)$$

sending a, b to

$$A = \begin{pmatrix} -1 & -2 \\ 0 & -1 \end{pmatrix}, \quad B = \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix}$$

respectively. By Grauert-Remmert's theorem (see, e.g., [GR71, XII, Cor. 5.2]), the étale fundamental group $\pi_1^{\text{ét}}(U, -1)$ is the profinite completion of $\pi_1^{\text{top}}(U^{\text{an}}, -1)$. Since $\text{SL}_2(\mathbb{Z}_\ell)$ is a profinite group, the morphism (3.1) extends naturally to a continuous morphism

$$\pi_1^{\text{ét}}(U, -1) \rightarrow \text{SL}_2(\mathbb{Z}_\ell) \hookrightarrow \text{GL}_2(\bar{\mathbb{Q}}_\ell^2). \quad (3.2)$$

The representation (3.2) is irreducible. Otherwise, assume that $v := (x, y)^T \neq 0 \in \bar{\mathbb{Q}}_\ell^2$ generates a 1-dimensional subrepresentation. Then $Av = (-x - 2y, -y)^T$ is parallel to v . Therefore, $y = 0$. Similarly, $Bv = (-x, -2x - y)^T$ is parallel to v , then $x = 0$, a contradiction.

Let L be the rank two *simple lisse* $\bar{\mathbb{Q}}_\ell$ -sheaf on U corresponding to (3.2). Then L^{an} is the local system on U^{an} corresponding to (3.1). For every small open ball $B_0 \subset X^{\text{an}}$ centered at 0, the \mathbb{C} -vector space $H^0(B_0, j_*^{\text{an}} L^{\text{an}})$ is the kernel of the linear operator $A - 1$ on the stalk L_{-1}^{an} . Since $A - 1$ is invertible, one has $H^0(B_0, j_*^{\text{an}} L^{\text{an}}) = 0$. Therefore, the stalk $(j_*^{\text{an}} L^{\text{an}})_0 = 0$. Similarly, the stalk $(j_*^{\text{an}} L^{\text{an}})_1 = 0$. In conclusion, the natural morphism $j_!^{\text{an}} L^{\text{an}} \rightarrow j_*^{\text{an}} L^{\text{an}}$ is an isomorphism in $\text{Cons}(X^{\text{an}})$.

We prove that $H^1(U^{\text{an}}, L^{\text{an}}) = H^1(\pi_1^{\text{top}}(U^{\text{an}}, -1), L_{-1}^{\text{an}})$ is nonzero. Define a map $f : \pi_1^{\text{top}}(U^{\text{an}}, -1) \rightarrow \bar{\mathbb{Q}}_\ell^2$ inductively. Set $f(e) = 0$, $f(a) = f(b) = (1, 0)^T$, $f(a^{-1}) = -A^{-1}f(a)$, and $f(b^{-1}) = -B^{-1}f(b)$. Once f is defined for every element of $\pi_1^{\text{top}}(U^{\text{an}}, -1)$ with length $n \geq 1$, we define it on elements of length $n + 1$ as follows. For every element $g \in \pi_1^{\text{top}}(U^{\text{an}}, -1)$ of length n , set

$$\begin{aligned} f(ag) &= Af(g) + f(a), & f(bg) &= Bf(g) + f(b), \\ f(a^{-1}g) &= A^{-1}f(g) + f(a^{-1}), & f(b^{-1}g) &= B^{-1}f(g) + f(b^{-1}). \end{aligned}$$

The map f is a crossed homomorphism. It is not a boundary, because the equation $(A - 1)x = (B - 1)x = (1, 0)^T$ admits no solution in $\bar{\mathbb{Q}}_\ell^2$.

Therefore, L^{an} is in the cohomology support loci of U^{an} (in the sense of [BLSW17, p.295]). From [HT07, Example 8.1.35 (ii)], one has $j_! L[1] \in \text{Perv}(X)$. By [BLSW17, p.299], $j_!^{\text{an}} L^{\text{an}}[1]$ is not semisimple in $\text{Perv}(X^{\text{an}})$.

By [BBDG82, Thm. 4.3.1 (ii)], the intermediate extension $K := j_{!*} L[1]$ is a simple object of $\text{Perv}(X)$. We claim that $\mathcal{H}^{-1}K$ is not semisimple in $D_c^b(X)$. From [HT07, Prop. 8.2.11], K is isomorphic to $\tau^{\leq -1} Rj_* L[1]$,

where $\tau^{\leq -1} : D_c^b(X) \rightarrow D_c^b(X)$ is the truncation functor with respect to the standard t-structure. Thus, $\mathcal{H}^{-1}K$ is isomorphic to $\mathcal{H}^{-1}(Rj_*L[1]) = j_*L$ in $\text{Cons}(X)$. Then $(\mathcal{H}^{-1}K)^{\text{an}}$ is isomorphic to $j_!^{\text{an}}L^{\text{an}}$ in $\text{Cons}(X^{\text{an}})$. From [Kat90, p.375], one has $(\mathcal{H}^{-1}K)[1] \in \text{Perv}(X)$. Since $(\mathcal{H}^{-1}K)^{\text{an}}[1]$ is not semisimple in $\text{Perv}(X^{\text{an}})$, by [Kat90, Lem. 12.7.1.1], $(\mathcal{H}^{-1}K)[1]$ is not semisimple in $\text{Perv}(X)$. The claim is proved.

Lemma 3.2.1.5 is used in the proof of Theorem 3.5.1.1.

Lemma 3.2.1.5. *Let $U \subset X$ be an open subset of X . Then the functor $(-)|_U : \text{Perv}(X) \rightarrow \text{Perv}(U)$ sends every simple object of $\text{Perv}(X)$ to a simple or zero object of $\text{Perv}(U)$. In particular, the functor $(-)|_U : D_c^b(X) \rightarrow D_c^b(U)$ preserves semisimplicity.*

Proof. Let K be a simple object of $\text{Perv}(X)$. By [BBDG82, Thm. 4.3.1 (ii)], there is an irreducible, locally closed and geometrically smooth subvariety $j : V \rightarrow X$ and a simple lisse \mathbb{Q}_ℓ -sheaf on V , such that K is isomorphic to $j_!L[\dim V]$. If V is disjoint from U , then $K|_U = 0$. Otherwise, take a geometric point \bar{x} on $V \cap U$. From [GR71, V, Prop. 8.2], the morphism $\pi_1^{\text{ét}}(U \cap V, \bar{x}) \rightarrow \pi_1^{\text{ét}}(V, \bar{x})$ is surjective. Thus, the composite representation $\pi_1^{\text{ét}}(U \cap V, \bar{x}) \rightarrow \text{GL}(L_{\bar{x}})$ is also simple, i.e., the lisse \mathbb{Q}_ℓ -sheaf $L|_{U \cap V}$ is simple. Let $h : U \cap V \rightarrow U$ be the base change of $j : V \rightarrow X$ along the inclusion $U \rightarrow X$. Then $K|_U$ is isomorphic to $h_!L|_{U \cap V}[\dim(U \cap V)]$, hence simple in $\text{Perv}(U)$. \square

When $k = \mathbb{C}$, Fact 3.2.1.6 1 follows from Kashiwara's conjecture for semisimple perverse sheaves and the paragraph following [BBDG82, Thm. 6.2.5]. Kashiwara's conjecture is formulated in [Kas98, Sec. 1]; see also [Dri01, Sec. 1.2, 1]. It is reduced to de Jong's conjecture by Drinfeld [Dri01], which in turn is proved in [BK06] and [Gai07]. The case of general k follows via Fact 3.2.1.7.

Fact 3.2.1.6. *Let k be an algebraically closed field of characteristic 0. Let $f : X \rightarrow Y$ be a proper morphism of algebraic varieties over k . Let K be a semisimple object of $D_c^b(X)$.*

1. (Decomposition theorem) *Then Rf_*K is a semisimple object of $D_c^b(Y)$.*
2. (Global invariant cycle theorem, [BBDG82, Cor. 6.2.8]) *Let i be an integer. By Fact 3.2.1.2, there is a nonempty connected open subset $V \subset Y$ such that $\mathcal{H}^i Rf_*K|_V$ is a lisse sheaf. Then for every $y \in V(k)$, the canonical map*

$$H^i(X, K) \rightarrow H^i(X_y, K|_{X_y})^{\pi_1^{\text{ét}}(V, y)}$$

is surjective.

Fact 3.2.1.7. *Let E/F be an extension of algebraically closed fields. Let X be an algebraic variety over F . Then:*

1. ([JKLM23, proof of Lem. A.1]) *The functor $(-)|_{X_E} : D_c^b(X) \rightarrow D_c^b(X_E)$ is fully faithful. It induces an exact functor $\text{Perv}(X) \rightarrow \text{Perv}(X_E)$.*
2. ([BBDG82, Thm. 4.3.1 (ii)]) *An object of $\text{Perv}(X)$ is simple if and only if its image under $(-)|_{X_E} : \text{Perv}(X) \rightarrow \text{Perv}(X_E)$ is simple.*

Lemma 3.2.1.8. *Let L be a lisse $\bar{\mathbb{Q}}_\ell$ -sheaf of rank one on X . Then $- \otimes^L L : D_c^b(X) \rightarrow D_c^b(X)$ is an equivalence of categories. It is t -exact for the perverse t -structures.*

Proof. Let L^{-1} be the lisse sheaf dual to L . By associativity of the derived tensor product \otimes^L , the pair of functors $(- \otimes^L L, - \otimes^L L^{-1})$ is an equivalence.

1. Right t -exactness: The functor is t -exact for the standard t -structures. Thus, for every $K \in {}^pD^{\leq 0}(X)$ and every integer n , one has $\mathcal{H}^n(K \otimes^L L) = \mathcal{H}^n(K) \otimes^L L$. Therefore, one has $\text{Supp } \mathcal{H}^n(K \otimes^L L) = \text{Supp } \mathcal{H}^n(K)$. Thus, $K \otimes^L L \in {}^pD^{\leq 0}(X)$.
2. Left t -exactness: By Part 1, for every $K \in {}^pD^{\geq 0}(X)$, one has $L^{-1} \otimes^L \mathbb{D}_X K \in {}^pD^{\leq 0}(X)$. By [KW01, II, Cor. 7.5 f)], one has isomorphisms

$$\mathbb{D}_X(K \otimes^L L) \rightarrow R\mathcal{H}om(L, \mathbb{D}_X K) \rightarrow L^{-1} \otimes^L \mathbb{D}_X K$$

in $D_c^b(X)$. Therefore, one gets $K \otimes^L L \in {}^pD^{\geq 0}(X)$.

□

3.2.2 Universal local acyclicity

In Section 3.2.2, all schemes are assumed to be quasi-compact and quasi-separated. For a scheme X and a geometric point \bar{x} on X , denote by $O_{X, \bar{x}}^{\text{sh}}$ the strict henselization (in the sense of [Sta24, Tag 04GQ (3)]) of $O_{X, \bar{x}}$. Set $X_{(\bar{x})} := \text{Spec } O_{X, \bar{x}}^{\text{sh}}$.

Let $f : X \rightarrow S$ be a separated morphism of finite presentation between $\mathbb{Z}[1/\ell]$ -schemes.

Definition 3.2.2.1 ([Sta24, Tag 0GJM], [Bar23, Def. 1.2]). Let K be an object of $D_c^b(X)$.

- If for every geometric point \bar{x} on X and every geometric point \bar{t} on $S_{(\bar{s})}$ with $\bar{s} = f(\bar{x})$, the canonical morphism $R\Gamma(X_{(\bar{x})}, K) \rightarrow R\Gamma(X_{(\bar{x})} \times_{S_{(\bar{s})}} \bar{t}, K)$ is an isomorphism, then K is called f -locally acyclic.

- If for every morphism $S' \rightarrow S$ of schemes, in the cartesian square

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ f' \downarrow & \square & \downarrow f \\ S' & \xrightarrow{g} & S \end{array} \quad (3.3)$$

g'^*K is f' -locally acyclic, then K is called f -universally locally acyclic (f -ULA). Let $D^{\text{ULA}}(X/S) \subset D_c^b(X)$ be the full subcategory of f -ULA objects.

By [HS23, Thm. 4.4], an object $K \in D_c^b(X)$ is f -ULA if and only if K is universally locally acyclic in the sense of [HS23, Def. 3.2]. Thus, the notation $D^{\text{ULA}}(X/S)$ agrees with that in [HS23]. It is a triangulated subcategory of $D_c^b(X)$.

Fact 3.2.2.2.

1. ([Bar23, Lem. 3.4]) If $S = \text{Spec } k$, then $D^{\text{ULA}}(X/k) = D_c^b(X)$.
2. ([Bar23, Cor. 3.10 (i)]) If $f : X \rightarrow S$ is an isomorphism, then $D^{\text{ULA}}(X/S) \subset D_c^b(X)$ is the full subcategory of objects whose cohomology sheaves are lisse.
3. ([HS23, Prop. 3.4 (i)]) Let $g : S' \rightarrow S$ be a morphism of $\mathbb{Z}[1/\ell]$ -schemes. Then in the notation of (3.3), the functor $g'^* : D_c^b(X) \rightarrow D_c^b(X')$ restricts to a functor $D^{\text{ULA}}(X/S) \rightarrow D^{\text{ULA}}(X'/S')$.
4. ([Ric14, Lem. 3.15], [Bar23, Lem. 3.3 (i), (ii)]) Let $f' : Y \rightarrow S$ be a separated morphism of finite presentation between $\mathbb{Z}[1/\ell]$ -schemes. Let $h : X \rightarrow Y$ be a morphism of S -schemes. If h is smooth (resp. proper), then the functor $h^* : D_c^b(Y) \rightarrow D_c^b(X)$ (resp. $Rh_* : D_c^b(X) \rightarrow D_c^b(Y)$) restricts to a functor $D^{\text{ULA}}(Y/S) \rightarrow D^{\text{ULA}}(X/S)$ (resp. $D^{\text{ULA}}(X/S) \rightarrow D^{\text{ULA}}(Y/S)$).
5. ([HS23, p.643]) Let $g : S \rightarrow T$ be a smooth morphism of $\mathbb{Z}[1/\ell]$ -schemes. Then $D^{\text{ULA}}(X/S) \subset D^{\text{ULA}}(X/T)$.
6. ([Zhu17, Thm. A.2.5 (4)]) Let $f_i : X_i \rightarrow S$ ($i = 1, 2$) be a separated morphism of finite presentation between $\mathbb{Z}[1/\ell]$ -schemes. Let $K_i \in D^{\text{ULA}}(X_i/S)$. Then $K_1 \boxtimes_S K_2 \in D^{\text{ULA}}(X_1 \times_S X_2/S)$.

Although the derived external tensor product preserves universal local acyclicity (Fact 3.2.2.2 6), Example 3.2.2.3 shows that the derived tensor product of two ULA complexes may not be ULA.

Example 3.2.2.3. Let $f : X \rightarrow S$ be the morphism $\mathbf{A}_{\mathbb{C}}^2 \rightarrow \mathbf{A}_{\mathbb{C}}^1$, $(a, b) \mapsto a + b$. Let $i : A \rightarrow X$ (resp. $j : B \rightarrow X$) be the inclusion of a -axis (resp. b -axis) of X . Then $fi : A \rightarrow S$ (resp. $fj : B \rightarrow S$) is an isomorphism. Let $K = i_* \bar{\mathbb{Q}}_{\ell, A}$ and $K' = j_* \bar{\mathbb{Q}}_{\ell, B}$. By Fact 3.2.2.2 2 and 4, one has $K, K' \in D^{\text{ULA}}(X/S)$. As $K \otimes^L K'$ is the skyscraper supported at the origin of X with stalk $\bar{\mathbb{Q}}_{\ell}$, it is not f -ULA. (Otherwise, from Remark 3.2.2.4, the skyscraper viewed as a sheaf on A is fi -ULA, which contradicts Fact 3.2.2.2 2.)

Remark 3.2.2.4. Let $i : Z \rightarrow X$ be a closed immersion. Let $K \in D_c^b(Z)$. By definition, if $i_* K$ is f -locally acyclic, then K is fi -locally acyclic. Therefore, by the base change theorem, if $i_* K$ is f -ULA, then K is fi -ULA.

Lemma 3.2.2.5. Assume that S is irreducible with generic point η . Let $K \in D^{\text{ULA}}(X/S)$. If $K|_{X_{\bar{\eta}}} = 0$ in $D_c^b(X_{\bar{\eta}})$, then $K = 0$.

Proof. It suffices to prove that for every $s \in S$, one has $K|_{X_{\bar{s}}} = 0$ in $D_c^b(X_{\bar{s}})$. By [Gro61c, Prop. 7.1.9], there is a discrete valuation ring R and a separated morphism $g : \text{Spec}(R) = S' \rightarrow S$, sending the generic (resp. closed) point ξ (resp. r) of S' to η (resp. s). Let $i : R \rightarrow R^h$ be the henselization of R (in the sense of [Sta24, Tag 04GQ (1)]). By [Sta24, Tag 0AP3], R^h is a discrete valuation ring. From [Mil80, I, Exercise 4.9], the local morphism i is injective. Then $i^* : \text{Spec}(R^h) \rightarrow S'$ preserves the generic (resp. closed) point. Replacing R by R^h , one may assume further that R is henselian.

Consider the following cartesian squares

$$\begin{array}{ccccc} X'_{\bar{r}} & \xrightarrow{\bar{i}} & X'_{(\bar{r})} & \xleftarrow{\bar{j}} & X'_{\bar{\xi}} \\ \downarrow & & \downarrow & & \downarrow \\ \bar{r} & \longrightarrow & S'_{(\bar{r})} & \longleftarrow & \bar{\xi}, \end{array}$$

where every vertical morphism is a base change of $f : X \rightarrow S$. In the notation of (3.3), let $R\Phi : D^+(X') \rightarrow D^+(X'_{\bar{r}})$ be the vanishing cycle functor. Let $R\Psi : D^+(X') \rightarrow D^+(X'_{\bar{\xi}})$ be the nearby cycle functor. Set $K' = g'^* K$. By definition, one has $R\Psi(K') = \bar{i}^* R\bar{j}_*(K'|_{X'_{\bar{\xi}}})$. As R is henselian, from [Ill06, (1.1.3)], there is a natural exact triangle $K'|_{X'_{\bar{r}}} \rightarrow R\Psi(K') \rightarrow R\Phi(K') \xrightarrow{\pm 1}$ in $D^+(X'_{\bar{r}})$. Since $K'|_{X'_{\bar{\xi}}}$ is a pullback of $K|_{X_{\bar{\eta}}} = 0$, one has $K'|_{X'_{\bar{\xi}}} = 0$ and $R\Psi(K') = 0$. By [Ill06, Cor. 3.5], the universal local acyclicity of K implies $R\Phi(K') = 0$. Therefore, one gets $K'|_{X'_{\bar{r}}} = 0$.

Since $K'|_{X'_{\bar{r}}}$ is the pullback of $K|_{X_{\bar{s}}}$ under the field extension $k(\bar{r})/k(\bar{s})$, by Fact 3.2.1.7 1, one gets $K|_{X_{\bar{s}}} = 0$. \square

3.2.3 Relative perverse sheaves

Let $f : X \rightarrow S$ be a morphism of algebraic varieties over k . In particular, f is separated and of finite presentation. Set $K_{X/S} := Rf^! \mathbb{Q}_\ell \in D_c^b(X)$ to be the relative dualizing complex. The functor

$$\mathbb{D}_{X/S}(-) = R\mathcal{H}om_{\mathbb{Q}_\ell}(-, K_{X/S}) : D_c^b(X) \rightarrow D_c^b(X)^{\text{op}}$$

is called the relative Verdier duality. There is a canonical morphism of functors $\text{Id}_{D_c^b(X)} \rightarrow \mathbb{D}_{X/S} \circ \mathbb{D}_{X/S}$ ([KL85, (1.1.5)]).

Fact 3.2.3.1 is stated for ∞ -categories in [HS23], but holds for the underlying triangulated categories (described in [HRS23, Lem. 7.9]) by [HS23, Footnote 1].

Fact 3.2.3.1.

1. ([HS23, Thm. 1.1]) *There is a unique t-structure $({}^{p/S}D^{\leq 0}(X/S), {}^{p/S}D^{\geq 0}(X/S))$ on $D_c^b(X)$, called the relative perverse t-structure, with the following property: An object $K \in D_c^b(X)$ lies in ${}^{p/S}D^{\leq 0}(X/S)$ (resp. ${}^{p/S}D^{\geq 0}(X/S)$) if and only if for every geometric point $\bar{s} \rightarrow S$, the restriction $K|_{X_{\bar{s}}}$ lies in ${}^pD^{\leq 0}(X_{\bar{s}})$ (resp. ${}^pD^{\geq 0}(X_{\bar{s}})$). In particular, for every $s \in S$, the functor $(-)|_{X_s} : D_c^b(X) \rightarrow D_c^b(X_s)$ is t-exact, where the source (resp. target) is equipped with the relative (resp. absolute) perverse t-structure.*
2. ([HS23, Thm. 1.9]) *The relative perverse t-structure on $D_c^b(X)$ restricts to a t-structure $({}^{p/S}D^{\text{ULA}, \leq 0}(X/S), {}^{p/S}D^{\text{ULA}, \geq 0}(X/S))$ on $D^{\text{ULA}}(X/S)$.*
3. ([HS23, Prop. 3.4]) *The functor $\mathbb{D}_{X/S}$ preserves $D^{\text{ULA}}(X/S)$, and the morphism $\text{Id}_{D^{\text{ULA}}(X/S)} \rightarrow \mathbb{D}_{X/S} \circ \mathbb{D}_{X/S}$ of functors $D^{\text{ULA}}(X/S) \rightarrow D^{\text{ULA}}(X/S)$ is an isomorphism. The formation of $\mathbb{D}_{X/S} : D^{\text{ULA}}(X/S) \rightarrow D^{\text{ULA}}(X/S)^{\text{op}}$ commutes with any base change in S , so $\mathbb{D}_{X/S}$ exchanges ${}^{p/S}D^{\text{ULA}, \leq 0}(X/S)$ with ${}^{p/S}D^{\text{ULA}, \geq 0}(X/S)$.*

Definition 3.2.3.2. Let $\text{Perv}(X/S)$ (resp. $\text{Perv}^{\text{ULA}}(X/S)$) be the heart of the relative perverse t-structure on $D_c^b(X)$ (resp. $D^{\text{ULA}}(X/S)$).

By Fact 3.2.3.1 1, an object $K \in D_c^b(X)$ lies in $\text{Perv}(X/S)$ if and only if for every geometric point $\bar{s} \rightarrow S$, one has $K|_{X_{\bar{s}}} \in \text{Perv}(X_{\bar{s}})$.

Example 3.2.3.3.

1. ([HS23, p.632]) If $S = \text{Spec}(k)$, then $\text{Perv}(X/k) = \text{Perv}(X)$.
2. If f is universally injective, then $\text{Perv}(X/S) = \text{Cons}(X)$.
3. ([Bar23, Cor. 3.10 (ii)]) If f is smooth of relative dimension r , then the functor $(-)[r] : \text{Loc}(X) \rightarrow D_c^b(X)$ factors through $\text{Perv}^{\text{ULA}}(X/S)$.

Example 3.2.3.4. Let $i : Y \rightarrow X$ be a closed immersion of S -schemes, with $Y \rightarrow S$ smooth of relative dimension d and with geometrically connected fibers. If L is a lisse $\bar{\mathbb{Q}}_\ell$ -sheaf on Y , then $i_* L[d] \in \text{Perv}^{\text{ULA}}(X/S)$.

Indeed, by Fact 3.2.2.2 2, one has $L \in D^{\text{ULA}}(Y/Y)$. From the smoothness of $Y \rightarrow S$ and Fact 3.2.2.2 5, one has $L \in D^{\text{ULA}}(Y/S)$. Using the properness of $i : Y \rightarrow X$ and Fact 3.2.2.2 4, one has $i_* L[d] \in D^{\text{ULA}}(X/S)$. For every geometric point $\bar{s} \rightarrow S$, let $i_{\bar{s}} : Y_{\bar{s}} \rightarrow X_{\bar{s}}$ be the base change of i along the morphism $X_{\bar{s}} \rightarrow X$. By the proper base change theorem, $i_* L[d]|_{X_{\bar{s}}} = (i_{\bar{s}})_*(L|_{Y_{\bar{s}}})[d] \in \text{Perv}(X_{\bar{s}})$. Therefore, $i_* L[d] \in \text{Perv}^{\text{ULA}}(X/S)$.

Fact 3.2.3.5 ([HS23, Thm. 1.10 (ii)]). *Assume that S is geometrically unibranch and irreducible with generic point η . Then the functor*

$$(-)|_{X_\eta} : \text{Perv}^{\text{ULA}}(X/S) \rightarrow \text{Perv}(X_\eta)$$

is exact and fully faithful, and its essential image is stable under subquotients.

Lemma 3.2.3.6. *If S is geometrically unibranch and irreducible, then $\text{Perv}^{\text{ULA}}(X/S)$ is a Serre subcategory of $\text{Perv}(X/S)$.*

Proof. By definition, $\text{Perv}^{\text{ULA}}(X/S)$ is a strictly full subcategory of $\text{Perv}(X/S)$. By Fact 3.2.3.1 2 and [BBDG82, Thm. 1.3.6], $\text{Perv}^{\text{ULA}}(X/S) \subset \text{Perv}(X/S)$ is an abelian subcategory and closed under extensions in $D^{\text{ULA}}(X/S)$. As $D^{\text{ULA}}(X/S) \subset D_c^b(X)$ is a triangulated subcategory, $\text{Perv}^{\text{ULA}}(X/S)$ is closed under extensions in $\text{Perv}(X/S)$. Because S is geometrically unibranch, from the proof of [HS23, Thm. 6.8 (ii)], $\text{Perv}^{\text{ULA}}(X/S)$ is closed under subquotients in $\text{Perv}(X/S)$. By [Sta24, Tag 02MP], it is a Serre subcategory. \square

Lemma 3.2.3.7 is stated without proof for regular schemes S in [HS23, p.636].

Lemma 3.2.3.7. *Assume that S is smooth over k of equidimension d . Then the shifted inclusion*

$$(-)[d] : D^{\text{ULA}}(X/S) \rightarrow D_c^b(X) \quad (3.4)$$

is t -exact, where $D^{\text{ULA}}(X/S)$ (resp. $D_c^b(X)$) is equipped with the relative (resp. absolute) perverse t -structure. In particular, it induces an exact functor

$$(-)[d] : \text{Perv}^{\text{ULA}}(X/S) \rightarrow \text{Perv}(X). \quad (3.5)$$

Proof. 1. The functor $(-)[d] : D_c^b(X) \rightarrow D_c^b(X)$ is right t -exact, where the source (resp. target) is equipped with the relative (resp. absolute) perverse t -structure. For every geometric point \bar{s} on S , the functor

$(-)|_{X_{\bar{s}}} : D_c^b(X) \rightarrow D_c^b(X_{\bar{s}})$ is t-exact for the standard t-structures. Then for every integer n and every $K \in {}^{p/S}D^{\leq 0}(X/S)$, one has $\mathcal{H}^n(K[d])|_{X_{\bar{s}}} = \mathcal{H}^{n+d}(K|_{X_{\bar{s}}})$. Hence

$$X_{\bar{s}} \cap \text{Supp } \mathcal{H}^n(K[d]) = \text{Supp } \mathcal{H}^{n+d}(K|_{X_{\bar{s}}}).$$

As $K|_{X_{\bar{s}}} \in {}^pD^{\leq 0}(X_{\bar{s}})$, one has $\dim \text{Supp } \mathcal{H}^{n+d}(K|_{X_{\bar{s}}}) \leq -n - d$. By Lemma 3.2.3.10 3, one has

$$\dim \text{Supp } \mathcal{H}^n(K[d]) \leq -n.$$

From Lemma 3.2.3.10 1, the Zariski closure of $\text{Supp } \mathcal{H}^n(K[d])$ in X has dimension at most $-n$. Hence $K[d] \in {}^pD^{\leq 0}(X)$.

2. The functor (3.4) is left t-exact. One may assume that k is algebraically closed. For every $K \in {}^{p/S}D^{\text{ULA}, \geq 0}(X/S)$, by smoothness of S and the proof of [Bar23, Cor. 3.8], $\mathbb{D}_X(K[d])$ is (noncanonically) isomorphic to $(\mathbb{D}_{X/S}K)[d]$ in $D_c^b(X)$. From Fact 3.2.3.1 3, $\mathbb{D}_{X/S}K \in {}^{p/S}D^{\text{ULA}, \leq 0}(X/S)$. By Part 1, one has $(\mathbb{D}_{X/S}K)[d] \in {}^pD^{\leq 0}(X)$. Hence $K[d] \in {}^pD^{\geq 0}(X)$.

□

Remark 3.2.3.8. In Lemma 3.2.3.7, the functor $(-)[d] : D_c^b(X) \rightarrow D_c^b(X)$ may not send $\text{Perv}(X/S)$ to $\text{Perv}(X)$. Indeed, let $k = \mathbb{C}$, and let $f : X = 0 \rightarrow S = A_{\mathbb{C}}^1$ be the inclusion of the origin. By Example 3.2.3.3 1, the relative perverse t-structure on $D_c^b(X)$ coincides with the standard one (which is also the absolute perverse t-structure). Then $\text{Perv}(X/S) = \text{Perv}(X)$.

Lemma 3.2.3.9. *If S is integral with generic point η and $\dim S = d$, then the functor $(-)|_{X_{\eta}}[-d] : D_c^b(X) \rightarrow D_c^b(X_{\eta})$ is t-exact for the absolute perverse t-structures. In particular, it restricts to an exact functor*

$$(-)|_{X_{\eta}}[-d] : \text{Perv}(X) \rightarrow \text{Perv}(X_{\eta}). \quad (3.6)$$

Proof. 1. Right t-exactness: For every $K \in {}^pD^{\leq 0}(X)$ and every integer n , one has $\text{Supp } \mathcal{H}^n(K|_{X_{\eta}}[-d]) = \text{Supp } \mathcal{H}^{n-d}(K|_{X_{\eta}}) = X_{\eta} \cap \text{Supp } \mathcal{H}^{n-d}(K)$. By Lemma 3.2.3.10 4, one has

$$\dim \text{Supp } \mathcal{H}^n(K|_{X_{\eta}}[-d]) \leq \dim \text{Supp } (\mathcal{H}^{n-d}(K)) - d \leq -n.$$

From Lemma 3.2.3.10 1, one has $K|_{X_{\eta}}[-d] \in {}^pD^{\leq 0}(X_{\eta})$.

2. Left t-exactness: For every $K \in D_c^b(X)$ and every integer n , one has

$$\text{Supp } \mathcal{H}^n(\mathbb{D}_{X_{\eta}}(K|_{X_{\eta}}[-d])) = \text{Supp } \mathcal{H}^n((\mathbb{D}_X K)|_{X_{\eta}}[-d]). \quad (3.7)$$

Indeed, from [Del77, Thm. 2.13, p.242], by shrinking S to a nonempty open subset, one may assume that $K \in D^{\text{ULA}}(X/S)$. By the proof of [Bar23, Cor. 3.8], one has $\mathbb{D}_X K = (\mathbb{D}_{X/S} K)(d)[2d]$. From Fact 3.2.3.1 3, $(\mathbb{D}_X K)|_{X_\eta}[-d]$ is a Tate twist of $\mathbb{D}_{X_\eta}(K|_{X_\eta}[-d])$, which proves (3.7).

Now assume $K \in {}^p D^{\geq 0}(X)$. Then $\mathbb{D}_X K \in {}^p D^{\leq 0}(X)$. From Part 1, one has $(\mathbb{D}_X K)|_{X_\eta}[-d] \in {}^p D^{\leq 0}(X_\eta)$. By (3.7), one has $\mathbb{D}_{X_\eta}(K|_{X_\eta}[-d]) \in {}^p D^{\leq 0}(X_\eta)$, or equivalently, $K|_{X_\eta}[-d] \in {}^p D^{\geq 0}(X_\eta)$. □

By convention, the dimension of an empty space is $-\infty$.

Lemma 3.2.3.10. *Let X be a scheme of finite type over a field F . Let C be a quasi-constructible subset of X .*

1. *Then $\dim C = \dim \bar{C}$.*
2. *Let $\{B_i\}_{i=1}^n$ be finitely many locally closed subsets of X and $B = \cup_{i=1}^n B_i$. Then $\dim B = \max_{i=1}^n \dim B_i$.*

Let $f : X \rightarrow Y$ be a morphism between schemes of finite type over F .

3. *Let $n \geq 0$ be an integer such that $\dim(C \cap f^{-1}(y)) \leq n$ for every $y \in Y$. Then $\dim C \leq \dim Y + n$.*
4. *Assume that Y is integral with generic point η . Then $\dim Y + \dim(C \cap X_\eta) \leq \dim C$.*

Proof.

1. As X is a Noetherian scheme, the topological subspace C is Noetherian. Therefore, C is the union of finitely many irreducible components. Thus, one may assume further that C is nonempty and irreducible. Then the reduced induced closed subscheme \bar{C} of X is integral and of finite type over F . By [Bor91, AG. Prop. 1.3], C contains a nonempty open subset of \bar{C} . By [Har77, II, Exercise 3.20 (e)], one has $\dim C = \dim \bar{C}$.
2. For every $1 \leq i \leq n$, since $B_i \subset B$, one has $\dim B_i \leq \dim B$. Then $\max_i \dim B_i \leq \dim B$. As B_i is quasi-constructible in X , by 1, one has $\dim B_i = \dim \bar{B}_i$. As $\{\bar{B}_i\}_{i=1}^n$ is a finite closed cover of \bar{B} , one gets $\dim B \leq \dim \bar{B} = \max_i \dim \bar{B}_i = \max_i \dim B_i$.
3. By 2, one may assume that C is locally closed in X . Taking irreducible components, one may assume further that C is irreducible. Let Z be the Zariski closure of $f(C)$ in Y . Then Z is irreducible. With reduced induced subscheme structures, one views C and Z as integral schemes

of finite type over F . Moreover, $f : X \rightarrow Y$ induces a dominant morphism $g : C \rightarrow Z$ over F . By [Har77, II, Exercise 3.22 (b)], for every $y \in f(C) = g(C)$, one has

$$n \geq \dim C \cap f^{-1}(y) = \dim g^{-1}(y) \geq \dim C - \dim Z.$$

Hence $\dim C \leq \dim Z + n \leq \dim Y + n$.

4. As in the proof of 3, one may assume that C is an irreducible, locally closed subset of X and view C as an integral scheme of finite type over F . One may assume that $C \cap X_\eta$ is nonempty. As C_η is homeomorphic to $C \cap X_\eta$, the morphism $C \rightarrow Y$ induced by f is dominant. Thus, by [Har77, II, Exercise 3.22 (c)], one gets $\dim C \cap X_\eta = \dim C_\eta = \dim C - \dim Y$.

□

Lemma 3.2.3.11. *Assume that S is smooth over k , integral with generic point η and $\dim S = d$. Then:*

1. *Let $A \in \text{Perv}^{\text{ULA}}(X/S)$, and let $B[d]$ be a subquotient of $A[d]$ in $\text{Perv}(X)$. If the image $B|_{X_\eta} \in \text{Perv}(X_\eta)$ of $B[d]$ under the functor (3.6) is zero, then $B[d] = 0$ in $\text{Perv}(X)$.*
2. *The functor (3.5) identifies $\text{Perv}^{\text{ULA}}(X/S)$ as a Serre subcategory of $\text{Perv}(X)$.*

Proof.

1. By regularity of S and [HS23, Cor. 1.12], one has $B \in D^{\text{ULA}}(X/S)$. Since $B|_{X_\eta} = 0$, by Lemma 3.2.2.5, one has $B = 0$.
2. It follows from the definition that the functor (3.5) is fully faithful. Its essential image is closed under extensions in $\text{Perv}(X)$, because $\text{Perv}^{\text{ULA}}(X/S)$ is closed under extensions in the triangulated subcategory $D^{\text{ULA}}(X/S)$ of $D_c^b(X)$.

We claim that the essential image is closed under taking subobjects. Take $K \in \text{Perv}^{\text{ULA}}(X/S)$ and a subobject $L[d]$ of $K[d] \in \text{Perv}(X)$. As S is integral, Lemma 3.2.3.9 shows that $L|_{X_\eta} \subset K|_{X_\eta}$ is a subobject in $\text{Perv}(X_\eta)$. By smoothness of S and Fact 3.2.3.5, there is a subobject $L' \subset K$ in $\text{Perv}^{\text{ULA}}(X/S)$ with $L'|_{X_\eta} = L|_{X_\eta}$. Set $M = K/L' \in \text{Perv}^{\text{ULA}}(X/S)$. Let $N[d]$ be the image of $L[d]$ under the morphism $K[d] \rightarrow M[d]$ in $\text{Perv}(X)$. As the sequence

$$0 \rightarrow L'[d] \cap L[d] \rightarrow L[d] \rightarrow N[d] \rightarrow 0$$

is exact in $\text{Perv}(X)$, by Lemma 3.2.3.9, the sequence

$$0 \rightarrow L'|_{X_\eta} \cap L|_{X_\eta} \rightarrow L|_{X_\eta} \rightarrow N|_{X_\eta} \rightarrow 0$$

is exact in $\text{Perv}(X_\eta)$. Hence $N|_{X_\eta} = 0$. Since $N[d]$ is a subobject of $M[d] \in \text{Perv}(X)$, by Part 1, one has $N[d] = 0$. Then $L[d] \subset L'[d]$ is a subobject in $\text{Perv}(X)$. Since $(L'[d])/(L[d])$ is a quotient of $L'[d]$ in $\text{Perv}(X)$ and $(L'|_{X_\eta})/(L|_{X_\eta}) = 0$ in $\text{Perv}(X_\eta)$, one gets $(L'[d])/(L[d]) = 0$ in $\text{Perv}(X)$. Therefore, $L[d] = L'[d]$. The claim is proved.

Similarly, the essential image is closed under taking quotients. By [Sta24, Tag 02MP], the essential image is a Serre subcategory of $\text{Perv}(X)$.

□

3.3 Cotori

3.3.1 Definition and basic properties

For a profinite group G , let $\mathcal{C}(G)$ be the group of ℓ -adic characters, i.e., continuous morphisms $\chi : G \rightarrow \bar{\mathbb{Q}}_\ell^\times$. Then $\chi(G)$ are compact subgroup of $\bar{\mathbb{Q}}_\ell^\times$.

Lemma 3.3.1.1.

1. Let C be a compact subset of $\bar{\mathbb{Q}}_\ell$. Then there is a finite subextension E of $\bar{\mathbb{Q}}_\ell/\mathbb{Q}_l$ with $C \subset E$.
2. Let $G \leq \bar{\mathbb{Q}}_\ell^\times$ be a compact subgroup. Then there is a finite subextension E of $\bar{\mathbb{Q}}_\ell/\mathbb{Q}_l$ with $G \subset O_E^\times$.
3. In 2, let $G^{(\ell)}$ (resp. $G^{(\ell')}$) be the ℓ -Sylow subgroup (resp. maximal prime-to- ℓ quotient) of G . Then the topological group $G \xrightarrow{\sim} G^{(\ell)} \times G^{(\ell')}$, and $G^{(\ell')}$ is finite.

Proof. 1. Otherwise, there is a sequence of elements x_1, x_2, \dots in C with $[\mathbb{Q}_l(x_{n+1}) : \mathbb{Q}_l] > [\mathbb{Q}_l(x_n) : \mathbb{Q}_l]$ for every integer $n > 0$. Let $B \subset C$ be the (infinite) set of elements of this sequence. For every subset $S \subset B$, every finite subextension F/\mathbb{Q}_l , the set $S \cap F$ is finite, so closed in F . Therefore, S is closed in $\bar{\mathbb{Q}}_\ell$. In particular, the set B is closed and hence compact in C . Every subset of B is closed in B , so B is discrete. Thus, B is finite, a contradiction.

2. By 1, there is a finite subextension E of $\bar{\mathbb{Q}}_\ell/\mathbb{Q}_l$ containing G . By [Ser64, Thm. 1 2, p.122], one has $G \subset O_E^\times$.
3. By 2 and [Ser64, Cor., p.155], G is an ℓ -adic Lie group. From Lazard's theorem (see, e.g., [GSK09, p.711]), there is a pro- ℓ open subgroup $U \leq G$. By [RZ10, Cor. 2.3.6 (b)], there is an ℓ -Sylow subgroup $H \leq$

G containing U . Since G is compact, $[G : U]$ is finite. Thus, the group G/H is finite of order prime to ℓ . By [RZ10, Prop. 2.3.8], G is isomorphic to $G/H \times H$. Since G is commutative, by [RZ10, Cor. 2.3.6 (c)], G has exactly one ℓ -Sylow subgroup. \square

Let $\mathcal{C}(G)_{\ell'}$ (resp. $\mathcal{C}(G)_{\ell}$) be the subgroup of characters of finite order prime to ℓ (resp. that are pro- ℓ). Then there is a canonical isomorphism $\mathcal{C}(G)_{\ell} \xrightarrow{\sim} \mathcal{C}((G^{(\ell)})^{\text{ab}})$. By Lemma 3.3.1.1 3, one has $\mathcal{C}(G) = \mathcal{C}(G)_{\ell'} \times \mathcal{C}(G)_{\ell}$.

We review the contents of [GL96, Sec. 3.2]. Fix an integer $n \geq 0$. Let A_n be a free $\hat{\mathbb{Z}}$ -module of rank n . Let $\{\gamma_1, \dots, \gamma_n\}$ be a \mathbb{Z}_{ℓ} -basis of $A_n^{(\ell)}$. Let $\mathcal{R} = \{O_E : E/\mathbb{Q}_{\ell} \text{ is a finite subextension of } \bar{\mathbb{Q}}_{\ell}\}$, which is a directed set under inclusion. For every $R \in \mathcal{R}$, let m_R be the maximal ideal of R . Let $R[[A_n^{(\ell)}]] := \varprojlim_{i,j \geq 1} (R/m_R^i)[A_n^{(\ell)}/\ell^j]$ be the completed group ring. There is a canonical injective morphism $A_n^{(\ell)} \rightarrow R[[A_n^{(\ell)}]]^{\times}$ of groups.

Fact 3.3.1.2 ([GL96, p.509]). *The ring $R[[A_n^{(\ell)}]]$ is a Noetherian, regular, complete, local domain of Krull dimension $1 + n$. There is an isomorphism of topological rings*

$$R[[A_n^{(\ell)}]] \rightarrow R[[X_1, \dots, X_n]], \quad \gamma_i \mapsto 1 + X_i. \quad (3.8)$$

Gabber and Loeser introduce a scheme of ℓ -adic characters.

Definition 3.3.1.3. Write $R_n = \bar{\mathbb{Q}}_{\ell} \otimes_{\mathbb{Z}_{\ell}} \mathbb{Z}_{\ell}[[A_n^{(\ell)}]]$. Define the “cotorus” associated with A_n to be $\mathcal{C}_{\ell} := \text{Spec } R_n$.

By [GL96, Prop. A.2.2.3 (ii)], the scheme \mathcal{C}_{ℓ} is integral and regular. Its set of closed points coincides with $\mathcal{C}_{\ell}(\bar{\mathbb{Q}}_{\ell})$, and it is Zariski dense in \mathcal{C}_{ℓ} . When $n > 0$, the $\bar{\mathbb{Q}}_{\ell}$ -scheme \mathcal{C}_{ℓ} is **not** locally of finite type.

Lemma 3.3.1.4. *Every character $\chi : A_n^{(\ell)} \rightarrow \bar{\mathbb{Q}}_{\ell}^{\times}$ extends canonically to a surjective morphism $R_n \rightarrow \bar{\mathbb{Q}}_{\ell}$ of $\bar{\mathbb{Q}}_{\ell}$ -algebras.*

Proof. There is a finite subextension E/\mathbb{Q}_{ℓ} in $\bar{\mathbb{Q}}_{\ell}$ containing all the $\chi(\gamma_i)$. Then for every $f = \sum_{\alpha \in \mathbb{N}^n} c_{\alpha} X^{\alpha} \in \mathbb{Z}_{\ell}[[X_1, \dots, X_n]]$, by completeness of E , the series $\sum_{\alpha \in \mathbb{N}^n} c_{\alpha} \prod_{i=1}^n (\chi(\gamma_i) - 1)^{\alpha_i}$ converges in E . Denote its limit by $f(\chi(\gamma_1) - 1, \dots, \chi(\gamma_n) - 1)$. The composition $\mathbb{Z}_{\ell}[[A_n^{(\ell)}]] \rightarrow E$ of (3.8) followed by

$$\mathbb{Z}_{\ell}[[X_1, \dots, X_n]] \rightarrow E, \quad f \mapsto f(\chi(\gamma_1) - 1, \dots, \chi(\gamma_n) - 1)$$

extends χ . It induces the stated surjection. The construction is independent of the choice of the \mathbb{Z}_{ℓ} -basis of $A_n^{(\ell)}$. \square

By Lemma 3.3.1.4, for every $\chi \in \mathcal{C}(A_n)_{\ell}$, the corresponding character $A_n^{(\ell)} \rightarrow \bar{\mathbb{Q}}_{\ell}^{\times}$ induces a surjection $R_n \rightarrow \bar{\mathbb{Q}}_{\ell}$. Let $\Psi(\chi)$ be the corresponding element of $\mathcal{C}_{\ell}(\bar{\mathbb{Q}}_{\ell})$. Hence a map

$$\Psi : \mathcal{C}(A_n)_{\ell} \rightarrow \mathcal{C}_{\ell}(\bar{\mathbb{Q}}_{\ell}). \quad (3.9)$$

Fact 3.3.1.5 ([GL96, p.519]). *The map (3.9) is bijective.*

3.3.2 Cotori are Baire

The objective of Section 3.3.2 is Lemma 3.3.2.10, used in the proof of Theorem 3.5.3.1. We show that over an uncountable algebraically closed field, a reasonable scheme has uncountably many rational points outside a countable union of strict closed subsets. Fix an uncountable, algebraically closed field k .

Baire schemes

Definition 3.3.2.1. A k -scheme X is called k -Baire, if its dimension $\dim X$ is finite and $X(k) \setminus \bigcup_{i \geq 1} Z_i(k)$ is uncountable for every countable sequence $\{Z_i\}_{i \geq 1}$ of closed subschemes of X with $\dim Z_i < \dim X$ for all i . A k -algebra R is called k -Baire if $\text{Spec}(R)$ is k -Baire.

The underlying reduced induced closed subscheme $X_{\text{red}} \rightarrow X$ induces a bijection $X_{\text{red}}(k) \rightarrow X(k)$, so X is k -Baire if and only X_{red} is k -Baire. A Noetherian k -scheme of dimension 1 with uncountably many k -points is k -Baire.

Remark 3.3.2.2. Let $k = \mathbb{C}$. Let X be a complex algebraic variety with $\dim X > 0$. The analytification X^{an} of X is locally compact Hausdorff. Then by the Baire category theorem (see, e.g., [Wil70, Cor. 25.4 a)], X is \mathbb{C} -Baire.

Lemma 3.3.2.3. *Let $f : X \rightarrow Y$ be a finite surjective morphism of k -schemes. If Y is k -Baire, then so is X .*

Proof. Let $\{Z_i\}_i$ be a sequence of closed subschemes of X with $\dim Z_i < \dim X$. Then for every integer $i \geq 1$, since f is a closed morphism, $Y_i := f(Z_i)$ is closed in Y . Endow each Y_i with the reduced induced structure. Let $Z'_i := f^{-1}(Y_i) = Y_i \times_Y X$. Then there is a canonical closed immersion $Z_i \rightarrow Z'_i$. The restriction $Z_i \rightarrow Y_i$ of f is a finite surjective morphism. By [Sta24, Tag 0ECG], one has $\dim X = \dim Y$ and $\dim Y_i = \dim Z_i$. In particular, $\dim X$ is finite and $\dim Y_i < \dim Y$.

As k is algebraically closed, the induced map $X(k) \rightarrow Y(k)$ is surjective. Then the induced map

$$X(k) \setminus (\bigcup_{i \geq 1} Z'_i(k)) \rightarrow Y(k) \setminus (\bigcup_i Y_i(k))$$

is surjective. Because Y is k -Baire, the target is uncountable. Then $X(k) \setminus (\bigcup_{i \geq 1} Z_i(k))$ is also uncountable, as it contains the source. \square

Lemma 3.3.2.4. *Let X be a Noetherian k -scheme.*

1. Then X is k -Baire if and only if X has an irreducible component C with $\dim C = \dim X$, such that the underlying reduced induced closed subscheme C is k -Baire.
2. Assume that $n := \dim X - 1$ is finite. If X has uncountably many (pairwise set-theoretically distinct) irreducible, k -Baire, closed subschemes of dimension n , then X is k -Baire.

Proof. 1. Assume that there is such a component C . Consider a sequence of closed subschemes $\{Z_i\}_{i \geq 1}$ of X with $\dim Z_i < \dim X$ for all $i \geq 1$. Then for every $i \geq 1$, one has $\dim C \cap Z_i \leq \dim Z_i < \dim X = \dim C$. Since C is k -Baire, the set $C(k) \setminus \cup_i (C \cap Z_i)(k)$ is uncountable. Therefore, $X(k) \setminus \cup_i Z_i(k)$ is also uncountable.

Assume that every component of X of maximum dimension is not k -Baire. As X is Noetherian, one can write $X = \cup_{j=1}^n C_j$ as a finite union of the irreducible components. For every j with $\dim C_j = \dim X$, the scheme C_j is not k -Baire. Therefore, there is a sequence $\{Z_i^j\}_{i \geq 1}$ of closed subschemes of C_j such that $\dim Z_i^j < \dim C_j$ for all i and $C_j(k) \setminus \cup_i Z_i^j(k)$ is countable. The finite family of components C_k with $\dim C_k < \dim X$, joint with the sequences $\{Z_i^j\}_i$ for all j with $\dim C_j = \dim X$, gives a countable family $\{Z_s\}_s$ of closed subschemes of X with $\dim Z_s < \dim X$ for all s . Then $X(k) \setminus (\cup_s Z_s(k))$ is countable, so X is not k -Baire.

2. Consider a sequence of closed subschemes $\{Z_i\}_{i \geq 1}$ of X with $\dim Z_i < \dim X$ for all $i \geq 1$. Every Z_i is a Noetherian scheme, so it has only finitely many irreducible components. The set of irreducible components of the family $\{Z_i\}_i$ is countable. Thus, one may assume that every Z_i is irreducible. By assumption, X has an n -dimensional, irreducible, k -Baire closed subscheme X' which is set-theoretically distinct from any Z_i . For every $i \geq 1$, because $\dim X' = n \geq \dim Z_i$ and Z_i is irreducible, one has $X' \not\subset Z_i$ and $X' \cap Z_i \neq X'$. Since X' is irreducible, one has $\dim(X' \cap Z_i) < \dim X'$. As X' is k -Baire, the set $X'(k) \setminus \cup_{i \geq 1} (X' \times_X Z_i)(k)$ is uncountable, which is a subset of $X(k) \setminus \cup_{i \geq 1} Z_i(k)$. Therefore, X is k -Baire. □

Lemma 3.3.2.5 is well-known.

Lemma 3.3.2.5. *If X is a finite type k -scheme with $\dim X > 0$, then X is k -Baire.*

Proof. Since X is of finite type over k , its dimension m is finite and X has only finitely many irreducible components. Replacing X with an irreducible

component of dimension m , one may that assume X is irreducible. Then by [Har77, Exercise 3.20 (e), p.94], every nonempty open subset of X has dimension m . Replacing X by an affine open, one may assume that X is affine. By Noether's normalization lemma, there is a finite surjective morphism $p : X \rightarrow \mathbf{A}_k^m$ over k . By Lemma 3.3.2.3, one may assume $X = \mathbf{A}_k^m$.

By induction on $m > 0$, we prove that \mathbf{A}_k^m is k -Baire. When $m = 1$, $\dim \mathbf{A}_k^1 = 1$ and $\mathbf{A}_k^1(k)$ is uncountable, so \mathbf{A}_k^1 is k -Baire. Assume the statement for $m - 1$ with $m \geq 2$. The set of hyperplanes in \mathbf{A}_k^m is uncountable. By the inductive hypothesis, every hyperplane is k -Baire. From Lemma 3.3.2.4 2, so is \mathbf{A}_k^m . The induction is completed. \square

Baireness of cotori

We show that every positive dimensional cotorus is $\bar{\mathbb{Q}}_\ell$ -Baire.

Definition 3.3.2.6 ([BGR84, Def. 1, p.205]). Let A be a k -algebra, and let $A[X] \rightarrow B$ be an injective ring map. We say that B is k -Rückert over A if there is a nonempty family W of monic polynomials in $A[X]$ such that the following axioms are fulfilled:

1. If $f, g \in A[X]$ are monic polynomials with $fg \in W$, then $f, g \in W$.
2. For every $w \in W$, the A -algebra B/w is isomorphic to $A[X]/w$.
3. For every $b \in B \setminus \{0\}$, there is an automorphism σ of the k -algebra B and a unit $u \in B^\times$ such that $u\sigma(b) \in W$.

Remark 3.3.2.7. From Axiom 1, one gets $1 \in W$. If $W = \{1\}$, then by Axiom 3, for every $b \in B \setminus \{0\}$, one has $b \in B^\times$, i.e., B is a field. Conversely, if B is a field, then B is k -Rückert over A with $W = \{1\}$.

If $W \neq \{1\}$, then $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is surjective. Indeed, take $w(\neq 1) \in W$. By Axiom 2, there is an A -isomorphism $B/w \rightarrow A[X]/w$, hence an isomorphism $\text{Spec}(A[X]/w) \rightarrow \text{Spec}(B/w)$ of $\text{Spec}(A)$ -schemes. Because w is a monic polynomial different from 1, the ring map $A \rightarrow A[X]/w$ is injective and finite. The induced morphism $\text{Spec}(A[X]/w) \rightarrow \text{Spec}(A)$ is surjective, so $\text{Spec}(B/w) \rightarrow \text{Spec}(A)$ is surjective.

For a commutative ring R and an ideal $I \subset R$, let $V_R(I) = \text{Spec } R/I \subset \text{Spec } R$. For $r \in R$, let $V_R(r) = V_R(rR)$. Lemma 3.3.2.8 is used in the induction step of the proof of Lemma 3.3.2.10.

Lemma 3.3.2.8. *Let A be Noetherian k -algebra of dimension n . Let B be a domain, but not a field, containing $A[X]$. Assume that B is k -Rückert over A .*

1. *The ring B is Noetherian of dimension $n + 1$.*

2. Suppose that A is k -Baire. Let S be an uncountable subset of A such that for every $s \in S$, one has $\dim V_A(s) = n - 1$. Suppose that the family $\{V_A(s)\}_{s \in S}$ is pairwise disjoint. Then B is k -Baire.

Proof. For every $b \in B \setminus (B^\times \cup \{0\})$, by Axiom 3, there is an automorphism σ of the k -algebra B and a unit $u \in B^\times$ such that $w := u\sigma(b)$ is in W . Since b is not a unit, one has $w \neq 1$. By Axiom 2, the A -algebra B/w is isomorphic to $A[X]/w$. Since $w(\neq 1)$ is a monic polynomial over A , the ring map $A \rightarrow A[X]/w$ is injective finite.

1. One has

$$\dim B/b = \dim B/w = \dim A[X]/w \stackrel{(a)}{=} \dim A = n, \quad (3.10)$$

where (a) uses [Sta24, Tag 00OK]. The domain B is not a field, so $\dim B = n + 1$. By [BGR84, Prop. 2, p.206], the ring B is Noetherian.

2. The morphism $\operatorname{Spec} A[x]/w \rightarrow \operatorname{Spec} A$ is finite surjective. Then by Lemma 3.3.2.3, the algebra $A[X]/w$ is k -Baire. As σ is over k , the k -algebra B/b is isomorphic to B/w . Then B/b is k -Baire.

For every $s \in S$, one has $\dim V_A(s) < \dim A$, so $s \neq 0$. As B is not a field, from Remark 3.3.2.7, the morphism $\operatorname{Spec}(B) \rightarrow \operatorname{Spec}(A)$ is surjective. The preimage of $V_A(s)$ under the surjection $\operatorname{Spec}(B) \rightarrow \operatorname{Spec}(A)$ is $V_B(s)$, so $V_B(s)$ is nonempty. In particular, $s \notin B^\times$ and B/s is k -Baire. Moreover, the family $\{V_B(s)\}_{s \in S}$ is pairwise disjoint. By (3.10), one gets $\dim V_B(s) = n$.

By Part 1, B is Noetherian. Then for every $s \in S$, by Lemma 3.3.2.4 1, there is a k -Baire irreducible component $C_s \subset \operatorname{Spec}(B/s)$ of dimension n . The family $\{C_s\}_{s \in S}$ is pairwise disjoint. From Lemma 3.3.2.4 2, B is k -Baire.

□

Set $S_n := \bar{\mathbb{Q}}_\ell \otimes_{\mathbb{Z}_\ell} \mathbb{Z}_\ell[[X_1, \dots, X_n]]$. By [GL96, Prop. 3.2.2 (1)], the natural morphism $S_n \rightarrow \bar{\mathbb{Q}}_\ell[[X_1, \dots, X_n]]$ is injective. Then the isomorphism (3.8) identifies R_n with the $\bar{\mathbb{Q}}_\ell$ -subalgebra $S_n \subset \bar{\mathbb{Q}}_\ell[[X_1, \dots, X_n]]$.

Fact 3.3.2.9. For every integer $n \geq 0$,

1. ([GL96, Thm. A.2.1, Prop. A.2.2.1]) the ring S_n is a Noetherian, regular, Jacobson domain of Krull dimension n ;
2. ([GL96, Prop. A.2.2.2, proof of A.2.2.3 (ii)]) S_{n+1} is $\bar{\mathbb{Q}}_\ell$ -Rückert over S_n .

Lemma 3.3.2.10. For every integer $n \geq 1$, the algebra S_n is $\bar{\mathbb{Q}}_\ell$ -Baire.

Proof. Since $\bar{\mathbb{Q}}_\ell$ is a flat \mathbb{Z}_ℓ -module, the injection $\mathbb{Z}_\ell[X_1, \dots, X_n] \rightarrow \mathbb{Z}_\ell[[X_1, \dots, X_n]]$ induces an injection $\bar{\mathbb{Q}}_\ell[X_1, \dots, X_n] \rightarrow S_n$. The natural morphism

$$\mathrm{Spec}(\bar{\mathbb{Q}}_\ell[[X_1, \dots, X_n]]) \rightarrow \mathbf{A}_{\bar{\mathbb{Q}}_\ell}^n \quad (3.11)$$

of $\bar{\mathbb{Q}}_\ell$ -schemes factors through a morphism $p_n : \mathrm{Spec}(S_n) \rightarrow \mathbf{A}_{\bar{\mathbb{Q}}_\ell}^n$.

Let $\mathcal{M} = \cup_E m_E$, where E runs through all finite subextensions of $\bar{\mathbb{Q}}_\ell \subset \bar{\mathbb{Q}}_\ell$, and m_E is the maximal ideal of the ring of integers of E . Then \mathcal{M} is the maximal ideal of the integral closure $\bar{\mathbb{Z}}_\ell$ of \mathbb{Z}_ℓ inside $\bar{\mathbb{Q}}_\ell$. By [Rob00, Prop., p.128], the residue field $\bar{\mathbb{Z}}_\ell/\mathcal{M}$ is an algebraic closure of the finite field F_ℓ , so it is countable. As \mathbb{Z}_ℓ is uncountable, so is the set \mathcal{M} .

For every $(a_1, \dots, a_n) \in \mathcal{M}^n$, there is a surjective morphism of $\bar{\mathbb{Q}}_\ell$ -algebras:

$$\bar{\mathbb{Q}}_\ell[[X_1, \dots, X_n]] \rightarrow \bar{\mathbb{Q}}_\ell, \quad f \mapsto f(a_1, \dots, a_n).$$

Its kernel is a $\bar{\mathbb{Q}}_\ell$ -point of $\mathrm{Spec}(\bar{\mathbb{Q}}_\ell[[X_1, \dots, X_n]])$, whose image under (3.11) is $(a_1, \dots, a_n) \in \mathbf{A}_{\bar{\mathbb{Q}}_\ell}^n(\bar{\mathbb{Q}}_\ell)$. Hence $\mathcal{M}^n \subset p_n(\mathrm{Spec}(S_n)(\bar{\mathbb{Q}}_\ell))$. In particular, $\mathrm{Spec}(S_n)(\bar{\mathbb{Q}}_\ell)$ is uncountable.

By induction on $n > 0$, we prove that S_n is $\bar{\mathbb{Q}}_\ell$ -Baire, and $\{V_{S_n}(X_1 - a)\}_{a \in \mathcal{M}}$ is a pairwise disjoint family of $(n - 1)$ -dimensional subsets. When $n = 1$, by Fact 3.3.2.9 1, S_1 is $\bar{\mathbb{Q}}_\ell$ -Baire. Moreover, $\{V_{S_1}(X_1 - a)\}_{a \in \mathcal{M}}$ is a pairwise distinct family of closed point of $\mathrm{Spec}(S_1)$. The statement is proved for $n = 1$. Assume the statement for $n - 1$ with $n \geq 2$. By Fact 3.3.2.9, (3.10), and Lemma 3.3.2.8 2, the statement holds for n . The induction is completed. \square

3.4 Kramer-Weissauer theory

Let k be a field of characteristic 0. Let Vec_k be the category of finite dimensional k -vector spaces. Choose an algebraic closure \bar{k} of k . Let $\mathrm{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k)$ be the category of continuous, finite dimensional $\bar{\mathbb{Q}}_\ell$ -representations of Γ_k . Let A be an abelian variety over k . Recall that $\pi_1^{\acute{e}t}(A_{\bar{k}})$ is a free $\hat{\mathbb{Z}}$ -module of rank $2 \dim A$. With the notation of Section 3.3, set

- $\mathcal{C}(A) = \mathcal{C}(\pi_1^{\acute{e}t}(A_{\bar{k}}))$: the group of characters $\pi_1^{\acute{e}t}(A_{\bar{k}}) \rightarrow \bar{\mathbb{Q}}_\ell^*$;
- $\mathcal{C}(A)_{\ell'} = \mathcal{C}(\pi_1^{\acute{e}t}(A_{\bar{k}}))_{\ell'}$: the group of characters of finite order prime to ℓ ;
- $\mathcal{C}(A)_\ell$: the cotorus assigned to $\pi_1^{\acute{e}t}(A_{\bar{k}})$.

3.4.1 Generic vanishing theorem

For an object $K \in \mathrm{Perv}(A)$, set

$$\mathcal{S}(K) := \{\chi \in \mathcal{C}(A) \mid H^i(A_{\bar{k}}, K \otimes^L L_\chi) \neq 0 \text{ for some integer } i \neq 0\}.$$

Fact 3.4.1.1 ([KW15b, Thm. 1.1], [Wei16, Vanishing Theorem, p.561; Thm. 2]). *For every perverse sheaf $K \in \text{Perv}(A)$ and every character $\chi_{\ell'} \in \mathcal{C}(A)_{\ell'}$, the set*

$$\{\chi_{\ell} \in \mathcal{C}(A)_{\ell}(\bar{\mathbb{Q}}_{\ell}) \mid \chi_{\ell'} \chi_{\ell} \in \mathcal{S}(K)\}$$

is the set of $\bar{\mathbb{Q}}_{\ell}$ -points of a strict Zariski closed subset of the scheme $\mathcal{C}(A)_{\ell}$.

We review [KW15a, p.725]. For every $K \in \text{Perv}(A)$, its Euler characteristic satisfies

$$\chi(A, K) := \sum_{i \in \mathbb{Z}} (-1)^i \dim_{\bar{\mathbb{Q}}_{\ell}} H^i(A_{\bar{k}}, K) \geq 0. \quad (3.12)$$

Let $N(A) \subset \text{Perv}(A)$ be the full subcategory of objects K with $\chi(A, K) = 0$. From the additivity of the function $\chi(A, -) : \text{Ob}(\text{Perv}(A)) \rightarrow \mathbb{N}$ and (3.12), $N(A)$ is a Serre subcategory of $\text{Perv}(A)$. Let $\bar{P}(A) := \text{Perv}(A)/N(A)$ be the quotient abelian category. For every $\chi \in \mathcal{C}(A)$, set

$$\mathcal{E}^{\chi}(A_{\bar{k}}) = \{K \in \text{Perv}(A_{\bar{k}}) \mid H^i(A_{\bar{k}}, K \otimes^L L_{\chi}) = 0, \quad \forall i \in \mathbb{Z} \setminus \{0\}\}.$$

Then $\mathcal{E}^{\chi}(A_{\bar{k}})$ is closed under extensions in $\text{Perv}(A_{\bar{k}})$. Let $P^{\chi}(A) \subset \text{Perv}(A)$ be the full subcategory of objects K with $Q \in \mathcal{E}^{\chi}(A_{\bar{k}})$ for every simple subquotient Q of $K|_{A_{\bar{k}}}$ in $\text{Perv}(A_{\bar{k}})$.

By [BBDG82, Thm. 4.3.1 (i)], every object $K \in \text{Perv}(A)$ is Noetherian and Artinian. For every $\chi_{\ell'} \in \mathcal{C}(A)_{\ell'}$, by Fact 3.4.1.1 and Lemma 3.4.1.2 1, the set $\{\chi_{\ell} \in \mathcal{C}(A)_{\ell}(\bar{\mathbb{Q}}_{\ell}) \mid K \in P^{\chi_{\ell'} \chi_{\ell}}(A)\}$ is the set of $\bar{\mathbb{Q}}_{\ell}$ -points of a strict Zariski closed subset of $\mathcal{C}(A)_{\ell}$.

Lemma 3.4.1.2. *Let \mathcal{A} be an abelian category, and let $X \in \mathcal{A}$ be a Noetherian and Artinian object.*

1. *Let Y be a simple subquotient of X . Then there is a composite series of X with one graded piece isomorphic to Y . In particular, up to isomorphism X has only finitely many simple subquotients.*
2. *If every subobject of X admits a direct complement, then X is semisimple.*

Proof.

1. There is a subobject $i : X_0 \subset X$ and a quotient $q : X_0 \rightarrow Y$ in \mathcal{A} . Let $N = \ker(q)$. By [Sta24, Tag 0FCH, Tag 0FCI], both N and X/X_0 are Noetherian and Artinian. From [Sta24, Tag 0FCJ], they admit composite series. A composite series of X/X_0 is equivalent to a filtration $X_0 \subset X_1 \subset X_2 \subset \cdots \subset X_n = X$ by subobjects such that X_i/X_{i-1} is simple for every $1 \leq i \leq n$. This filtration and every composite series of N glue to a composite series of X with a step $N \subset X_0$, whose factor is isomorphic to Y . By the Jordan-Hölder lemma [Sta24, Tag 0FCK], up to isomorphism Y has finitely many choices.

2. One may assume that $X \neq 0$. Let \mathcal{P} be the family of nonzero semisimple subobjects of X . By [Sta24, Tag 0FCJ], X has a nonzero simple subobject, so \mathcal{P} is nonempty. Since X is Noetherian, the family \mathcal{P} has a maximal element $i : X_0 \rightarrow X$. By assumption, there is a subobject $F \subset X$ with $X_0 \oplus F = X$. Then $F = 0$. (Otherwise, by [Sta24, Tag 0FCJ], F has a nonzero simple subobject F_0 . Then $X_0 \oplus F_0 \in \mathcal{P}$ is strictly larger than X_0 , which is a contradiction.) Therefore, i is an isomorphism and X is semisimple.

□

Remark 3.4.1.3. In a Noetherian and Artinian abelian category, an object may have infinitely many distinct (non semisimple) subobjects up to isomorphism.

Lemma 3.4.1.4. *Let \mathcal{A} be a Noetherian and Artinian abelian category. Let \mathcal{E} be a class of objects of \mathcal{A} closed under isomorphisms. Let $\mathcal{S} \subset \mathcal{A}$ be the full subcategory of objects every nonzero simple subquotient of which is in \mathcal{E} .*

1. *Then \mathcal{S} is a Serre subcategory of \mathcal{A} .*
2. *If further \mathcal{E} is closed under extensions, then $\mathcal{S} \subset \mathcal{E}$.*

Proof.

1. (a) We prove that \mathcal{S} is closed under subquotients. Let X be an object of \mathcal{S} with a subquotient Y . Every simple subquotient of Y is that of X , hence in \mathcal{E} . Thus, $Y \in \mathcal{S}$.

Let $0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$ be a short exact sequence in \mathcal{A} with $L, N \in \mathcal{S}$. Let Q be a nonzero simple subquotient of M . We prove that $Q \in \mathcal{E}$.

- (b) First, assume that Q is a quotient of M . The natural morphism $L \rightarrow Q$ is either an epimorphism or zero, in which case Q is a simple quotient of L or N respectively. Hence $Q \in \mathcal{E}$.
- (c) Now assume that Q is general. There is a subobject $M_0 \subset M$ and an epimorphism $M_0 \rightarrow Q$. Then

$$0 \rightarrow f^{-1}(M_0) \rightarrow M_0 \rightarrow g(M_0) \rightarrow 0$$

is a short exact sequence in \mathcal{A} with $f^{-1}(M_0)$ (resp. $g(M_0)$) a subobject of L (resp. N). From Part 1a, both $f^{-1}(M_0)$ and $g(M_0)$ are in \mathcal{S} . From Part 1b, one has $Q \in \mathcal{E}$.

From Part 1c, one has $M \in \mathcal{S}$ and \mathcal{S} is closed under extensions. The result follows from [Sta24, Tag 02MP].

2. By [Sta24, Tag 0FCJ], every object $X \in \mathcal{S}$ admits a filtration in \mathcal{A}

$$0 \subset X_1 \subset X_2 \subset \cdots \subset X_n = X$$

by subobjects such that each X_i/X_{i-1} is a simple subquotient of X . Then $X_i/X_{i-1} \in \mathcal{E}$. As \mathcal{E} is closed under extensions, one has $X \in \mathcal{E}$. \square

By Lemma 3.4.1.4 1, for every $\chi \in \mathcal{C}(A)$, $P^\chi(A) \subset \text{Perv}(A)$ is a Serre subcategory. From Lemma 3.4.1.4 2, for every $K \in P^\chi(A)$ and every integer $i \neq 0$, one has

$$H^i(A_{\bar{k}}, K \otimes^L L_\chi) = 0. \quad (3.13)$$

From the proof of [LS20, Lem. 3.4 (3)], the functor

$$\omega_\chi : P^\chi(A) \rightarrow \text{Vec}_{\bar{\mathbb{Q}}_\ell}, \quad K \mapsto H^0(A_{\bar{k}}, K \otimes^L L_\chi) \quad (3.14)$$

is exact. Let $N^\chi(A)$ be the full subcategory of $P^\chi(A)$ of objects in $N(A)$. For every $K \in N^\chi(A)$, by [KW15b, Cor. 4.2], one has $\chi(A, K \otimes^L L_\chi) = 0$. From (3.13), one has $H^0(A_{\bar{k}}, K \otimes^L L_\chi) = 0$. By [Sta24, Tag 02MS], the functor ω_χ factors uniquely through an exact functor (still denoted by ω_χ)

$$P^\chi(A)/N^\chi(A) \rightarrow \text{Vec}_{\bar{\mathbb{Q}}_\ell}. \quad (3.15)$$

3.4.2 Tannakian groups

Let (\mathcal{C}, \otimes) a neutral Tannakian category (in the sense of [DM22, Def. 2.19]) over an algebraically closed field Q of characteristic 0, with a fiber functor $\omega : \mathcal{C} \rightarrow \text{Vec}_Q$. Let $\text{Aut}^\otimes(\mathcal{C}, \omega)$ be the corresponding affine group scheme over Q . By [Del90, Sec. 9.2, p.187], up to isomorphism of group schemes, $\text{Aut}^\otimes(\mathcal{C}, \omega)$ is independent of the choice of ω . (See [Wib22, Thm. 1.2] for an elementary proof.)

For an object $K \in \mathcal{C}$, let $\iota : \langle K \rangle \hookrightarrow \mathcal{C}$ be the full subcategory whose objects are the subquotients of $\{(K \oplus K^\vee)^{\otimes n}\}_{n \geq 1}$. Then $(\langle K \rangle, \otimes)$ is a neutral Tannakian subcategory of \mathcal{C} (in the sense of [Mil07, 1.7]), for which $\omega_\iota : \langle K \rangle \rightarrow \text{Vec}_Q$ is a fiber functor. The group scheme $\text{Aut}^\otimes(\langle K \rangle, \omega_\iota)$ is the image of the natural morphism $\text{Aut}^\otimes(\mathcal{C}, \omega) \rightarrow \text{GL}(\omega(K))$.

Definition 3.4.2.1. The algebraic group $\text{Aut}^\otimes(\langle K \rangle, \omega_\iota)$ is called the Tannakian monodromy group of K at ω and is denoted by $G_\omega(K)$.

By [Sim92, p.69], $G_\omega(K)$ is reductive if and only if K is semisimple in \mathcal{C} .

Example 3.4.2.2. With tensor product, $\text{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k)$ is a neutral Tannakian category over $\bar{\mathbb{Q}}_\ell$. The forgetful functor $\omega : \text{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k) \rightarrow \text{Vec}_{\bar{\mathbb{Q}}_\ell}$ is a fiber functor. The Tannakian monodromy group of an object $\rho : \Gamma_k \rightarrow \text{GL}(V)$ at ω is the Zariski closure of $\rho(\Gamma_k) \subset \text{GL}(V)$.

3.4.3 Sheaf convolution

Let $m : A \times_k A \rightarrow A$ be the group law on A . Let $p_i : A \times_k A \rightarrow A$ be the projection to i -th factor ($i = 1, 2$). The bifunctor

$$* : D_c^b(A) \times D_c^b(A) \rightarrow D_c^b(A), \quad - * + := Rm_*(p_1^* - \otimes^L p_2^* +)$$

is called the *convolution* on A .

Example 3.4.3.1. For every closed reduced subvariety $i : X \rightarrow A$, let $\delta_X := i_* \bar{\mathbb{Q}}_{\ell, X} \in D_c^b(A)$. Then for every closed point $x \in A$, one has $\delta_x * \delta_X = \delta_{x+X}$.

By [Wei11] and [JKLM23, Sec. 3.1], the pair $(D_c^b(A), *)$ is a rigid, symmetric monoidal category, with unit δ_0 . For every $K \in D_c^b(A)$, its adjoint dual is $K^\vee := [-1]_A^* \mathbb{D}_A K$.

Fact 3.4.3.2 ([KW15b, proof of Thm. 13.2], [LS20, Lem. 3.4 (4)], [JKLM23, Prop. 3.1]). *The convolution on $D_c^b(A)$ induces a bifunctor $\bar{P}(A) \times \bar{P}(A) \rightarrow \bar{P}(A)$, $(-, +) \mapsto {}^p\mathcal{H}^0(- * +)$ fitting into a commutative square*

$$\begin{array}{ccc} \mathrm{Perv}(A) \times \mathrm{Perv}(A) & \xrightarrow{*} & D_c^b(A) \\ \downarrow & & \downarrow {}^p\mathcal{H}^0 \\ \bar{P}(A) \times \bar{P}(A) & \dashrightarrow & \bar{P}(A). \end{array}$$

It makes $\bar{P}(A)$ a neutral Tannakian category over $\bar{\mathbb{Q}}_\ell$. For every $\chi \in \mathcal{C}(A)$, the subcategory $P^\chi(A)/N^\chi(A) \subset \bar{P}(A)$ is a Tannakian subcategory, on which (3.15) is a fiber functor.

Example 3.4.3.3. [KW15a, Example 7.1] Fix a closed point $x \in A$. Then $\delta_x \in \mathrm{Perv}(A)$. Then $\mathcal{S}(\delta_x)$ is empty and for every $\chi \in \mathcal{C}(A)$, one has $\delta_x \in P^\chi(A)$. If x is a torsion point of order n , then $G_{\omega_\chi}(\delta_x)$ is isomorphic to \mathbb{Z}/n . If x is not a torsion point, then $G_{\omega_\chi}(\delta_x)$ is isomorphic to $\mathbb{G}_{m/\bar{\mathbb{Q}}_\ell}$.

Let $\psi : \pi_1^{\text{ét}}(A) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ be a character, and set $\psi' = \psi|_{\pi_1^{\text{ét}}(A_{\bar{k}})}$. The functor

$$\omega_\psi : \mathrm{Perv}(A) \rightarrow \mathrm{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k), \quad K \mapsto H^0(A_{\bar{k}}, K \otimes^L L_\psi)$$

fits into a commutative square

$$\begin{array}{ccc} \mathrm{Perv}(A) & \xrightarrow{\omega_\psi} & \mathrm{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k) \\ \uparrow & & \downarrow \omega \\ P^{\psi'}(A) & \xrightarrow{(3.14)} & \mathrm{Vec}_{\bar{\mathbb{Q}}_\ell} \end{array}$$

The quotient functor $P^{\psi'}(A)/N^{\psi'}(A) \rightarrow \mathrm{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k)$ of $\omega_\psi|_{P^{\psi'}(A)}$ induces a morphism of affine groups schemes

$$\omega_\psi^* : \mathrm{Aut}^\otimes(\mathrm{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k), \omega) \rightarrow \mathrm{Aut}^*(P^{\psi'}(A)/N^{\psi'}(A), \omega_{\psi'}). \quad (3.16)$$

Definition 3.4.3.4. For every $K \in \text{Perv}(A)$, let $\text{Mon}(K, \psi)$ be the Tannakian monodromy group of $\omega_\psi(K)$ in $\text{Rep}_{\bar{\mathbb{Q}}_\ell}(\Gamma_k)$.

For every $K \in P^{\psi'}(A)$, the functor $\omega_\psi|_{\langle K \rangle} : \langle K \rangle \rightarrow \langle \omega_\psi(K) \rangle$ induces a closed immersion of linear algebraic groups $\omega_\psi^* : \text{Mon}(K, \psi) \rightarrow G_{\omega_{\psi'}}(K)$, which is the projection of (3.16) in $\text{GL}(\omega_{\psi'}(K))$.

3.5 Main results

Consider Setting 3.1.2.1. For every character $\chi \in \mathcal{C}(A)$, denote the pullback of χ along $(p_A|_{A_\eta})_* : \pi_1^{\text{ét}}(A_\eta) \rightarrow \pi_1^{\text{ét}}(A)$ by $\chi_\eta : \pi_1^{\text{ét}}(A_\eta) \rightarrow \bar{\mathbb{Q}}_\ell^\times$. Then the restriction $\chi_\eta|_{\pi_1^{\text{ét}}(A_{\bar{\eta}})}$ is identified with χ via the isomorphism $(p_A|_{A_{\bar{\eta}}})_* : \pi_1^{\text{ét}}(A_{\bar{\eta}}) \rightarrow \pi_1^{\text{ét}}(A)$. Let $K \in \text{Perv}(A \times X/X)$ be an object which is semisimple in $D_c^b(A \times X)$.

We shall prove that the monodromy group of K is normal in its Tannakian group. By the normality criterion (Lemma 3.5.0.1), it suffices to show that the monodromy is reductive, and to consider the monodromy fixed part of *all* the representations of the Tannakian group. Such representations are from perverse sheaves.

Lemma 3.5.0.1. *Let G be a linear algebraic group over an algebraically closed field C . Let H be a closed, reductive subgroup of G . If for every $V \in \text{Rep}_C(G)$, the subspace V^H is G -stable, then H is normal in G .*

Proof. By [Gro06, Cor. 2.4] and reductivity, H is observable in G (in the sense of [BBHM63, p.134]). From [And21, Prop. C.3], H is normal in G . \square

3.5.1 Reductivity

Theorem 3.5.1.1. *For every $\chi \in \mathcal{C}(A) \setminus \mathcal{S}(K|_{A_\eta})$, the monodromy group $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ is reductive.*

Proof. By Lemma 3.2.1.5, when X is replaced by a nonempty open subset, the semisimplicity of K in $D_c^b(A \times X)$ is preserved. Moreover, the $\Gamma_{k(\eta)}$ -representation $\omega_{\chi_\eta}(K|_{A_\eta})$ and hence the group $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ remain unchanged. Thus, by [Sta24, Tag 056V], one may assume that X is smooth. As K is semisimple in $D_c^b(A \times X)$, from Lemma 3.2.1.8, so is $K \otimes^L p_A^* L_\chi$. By Fact 3.2.1.6 1, the object $Rp_{X*}(K \otimes^L p_A^* L_\chi)$ is semisimple in $D_c^b(X)$.

By the proper base change theorem (see, e.g., [Sta24, Tag 095T]), for every integer n , one has

$$\mathcal{H}^n Rp_{X*}(K \otimes^L p_A^* L_\chi)_{\bar{\eta}} = H^n(A_{\bar{\eta}}, K|_{A_{\bar{\eta}}} \otimes^L L_\chi).$$

Since $\chi \notin \mathcal{S}(K|_{A_\eta})$, when $n \neq 0$, one has $H^n(A_{\bar{\eta}}, K|_{A_{\bar{\eta}}} \otimes^L L_\chi) = 0$. By Fact 3.2.1.2, there is a nonempty open subset U_0 (resp. U_n for every integer $n \neq 0$) of X such that $[\mathcal{H}^0 R p_{X*}(K \otimes^L p_A^* L_\chi)]|_{U_0}$ is a lisse $\bar{\mathbb{Q}}_\ell$ -sheaf (resp. $[\mathcal{H}^n R p_{X*}(K \otimes^L p_A^* L_\chi)]|_{U_n} = 0$). The set

$$J := \{n \in \mathbb{Z} : \mathcal{H}^n R p_{X*}(K \otimes^L p_A^* L_\chi) \neq 0\}$$

is finite and X is irreducible, so $U := U_0 \cap \bigcap_{n \in J} U_n$ is a nonempty open subset of X . Shrinking X to U , one may assume further that $\mathcal{H}^n R p_{X*}(K \otimes^L p_A^* L_\chi) = 0$ for every integer $n \neq 0$, and that $\mathcal{H}^0 R p_{X*}(K \otimes^L p_A^* L_\chi)$ is a lisse $\bar{\mathbb{Q}}_\ell$ -sheaf on X .

Thus, the semisimple object $R p_{X*}(K \otimes^L p_A^* L_\chi)[\dim X]$ of $D_c^b(X)$ lies in $\text{Perv}(X)$, so it is semisimple in $\text{Perv}(X)$. By [Ach21, Prop. 3.4.1], the object $R p_{X*}(K \otimes^L p_A^* L_\chi)$ of $\text{Loc}(X)$ is semisimple. Therefore, the corresponding representation

$$\pi_1^{\text{ét}}(X, \bar{\eta}) \rightarrow \text{GL}(H^0(A_{\bar{\eta}}, K|_{A_{\bar{\eta}}} \otimes^L L_\chi))$$

is semisimple. Because X is smooth, the natural morphism $\eta_* : \Gamma_{k(\eta)} \rightarrow \pi_1^{\text{ét}}(X, \bar{\eta})$ is surjective. Then the composition $\Gamma_{k(\eta)} \rightarrow \text{GL}(H^0(A_{\bar{\eta}}, K|_{A_{\bar{\eta}}} \otimes^L L_\chi))$, i.e., the representation $\omega_{\chi_\eta}(K|_{A_\eta})$, is semisimple. Consequently, the algebraic group $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ is reductive. \square

Example 3.5.1.2. In Theorem 3.5.1.1, the algebraic group $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ may not be semisimple. Let X be a smooth, projective, integral algebraic curve over k of genus 1. Then $\pi_1^{\text{ét}}(X, \bar{\eta}) \cong \hat{\mathbb{Z}}^2$. There exists a character $\sigma : \pi_1^{\text{ét}}(X, \bar{\eta}) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ of infinite order. Let $A = \text{Spec}(k)$. Then $\mathcal{C}(A) = \{1\}$ and $\text{Mon}(L_\sigma|_{A_\eta}, 1) = \mathbb{G}_{m/\bar{\mathbb{Q}}_\ell}$ is an algebraic torus.

Remark 3.5.1.3. In view of Example 3.2.1.4, the semisimplicity of $\mathcal{H}^0 R p_{X*}(K \otimes^L p_A^* L_\chi)$ in $D_c^b(X)$ is not clear *a priori*. That is why we exclude characters in the spectrum $\mathcal{S}(K|_{A_\eta})$ in Theorem 3.5.1.1.

Remark 3.5.1.4. Let $i : Y \rightarrow A \times X$ be a closed subvariety, such that the induced morphism $f : Y \rightarrow X$ is smooth with connected fibers of dimension d :

$$\begin{array}{ccc} Y & \xhookrightarrow{i} & A \times X \\ \downarrow f & \swarrow p_X & \downarrow p_A \\ X & & A. \end{array}$$

By Example 3.2.3.4, one has $K := i_* \bar{\mathbb{Q}}_{\ell, Y}[d] \in \text{Perv}^{\text{ULA}}(A \times X/X)$. By Fact 3.2.1.6 1, it is semisimple in $D_c^b(A \times X)$. Assume that X is smooth. Then for every $\chi \in \mathcal{C}(A) \setminus \mathcal{S}(K|_{A_\eta})$, the algebraic group $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ coincides with the Zariski closure of the image of the monodromy representation of the lisse $\bar{\mathbb{Q}}_\ell$ -sheaf $R^d f_* i^* p_A^* L_\chi$ on X , which is studied in [KM23, Sec. 1.4] (but with coefficient \mathbb{C} instead of $\bar{\mathbb{Q}}_\ell$).

3.5.2 Fixed part

Theorem 3.1.2.3 follows from Theorem 3.5.2.1 and Fact 3.4.1.1, because the union in Condition 1 of Theorem 3.5.2.1 is in fact a finite union.

Theorem 3.5.2.1. *Assume that X is smooth and $K \in \text{Perv}^{\text{ULA}}(A \times X/X)$. Then there exists a subobject $K^0 \subset K$ in $\text{Perv}^{\text{ULA}}(A \times X/X)$ such that for every $\chi \in \mathcal{C}(A)$ with*

1. $\chi \notin \cup_{j \in \mathbb{Z}} \mathcal{S}({}^p\mathcal{H}^j(Rp_{A*}K))$,
2. $K|_{A_\eta} \in P^\chi(A_\eta)$ and
3. ${}^p\mathcal{H}^0(Rp_{A*}K) \in P^\chi(A)$,

one has $\omega_{\chi_\eta}(K^0|_{A_\eta}) = \omega_{\chi_\eta}(K|_{A_\eta})^{\Gamma_{k(\eta)}}$.

Proof. By properness of $p_X : A \times X \rightarrow X$ and Fact 3.2.2.2 4, one has $Rp_{X*}K \in D^{\text{ULA}}(X/X)$. Then from Fact 3.2.2.2 2, the sheaf $\mathcal{H}^0 Rp_{X*}K$ is lisse. Since X is smooth, by [Sta24, Tag 0BQM], the canonical morphism $\Gamma_{k(\eta)} \rightarrow \pi_1^{\text{ét}}(X, \bar{\eta})$ is surjective. Thus, from Fact 3.2.1.6 2, the natural map

$$H^0(A \times X, K \otimes^L p_A^* L_\chi) \rightarrow \omega_{\chi_\eta}(K|_{A_\eta})^{\Gamma_{k(\eta)}} \quad (3.17)$$

is surjective.

By Fact 3.2.1.1, one has

$$H^0(A \times X, K \otimes^L p_A^* L_\chi) = H^0(A, (Rp_{A*}K) \otimes^L L_\chi). \quad (3.18)$$

By Condition 1, for any integers $i \neq 0$ and j , one has

$$H^i(A, {}^p\mathcal{H}^j(Rp_{A*}K) \otimes^L L_\chi) = 0.$$

By Lemma 3.2.1.8, the spectral sequence in [Max19, Rk. 8.1.14 (6)] becomes

$$E_2^{i,j} = H^i(A, {}^p\mathcal{H}^j(Rp_{A*}K) \otimes^L L_\chi) \Rightarrow H^{i+j}(A, (Rp_{A*}K) \otimes^L L_\chi).$$

It degenerates at page E_2 . Hence

$$H^0(A, (Rp_{A*}K) \otimes^L L_\chi) = H^0(A, ({}^p\mathcal{H}^0 Rp_{A*}K) \otimes^L L_\chi). \quad (3.19)$$

Set $K^1 := p_A^* {}^p\mathcal{H}^0(Rp_{A*}K) \in D_c^b(A \times X)$. By Fact 3.2.2.2 1, one has ${}^p\mathcal{H}^0(Rp_{A*}K) \in D^{\text{ULA}}(A/k)$. From Fact 3.2.2.2 3, one gets $K^1 \in D^{\text{ULA}}(A \times X/X)$. For every $x \in X(k)$, the restriction $p_A|_{A_x} : A_x \rightarrow A$ is an isomorphism of abelian varieties over k , so the functor $(p_A|_{A_x})^* : \text{Perv}(A) \rightarrow \text{Perv}(A_x)$ is an equivalence of abelian categories. It sends ${}^p\mathcal{H}^0(Rp_{A*}K)$ to $K^1|_{A_x}$, so $K^1|_{A_x} \in \text{Perv}(A_x)$ and hence $K^1 \in \text{Perv}^{\text{ULA}}(A \times X/X)$. From

$K^1|_{A_\eta} = (p_A|_{A_\eta})^* {}^p\mathcal{H}^0(Rp_{A*}K)$ and Condition 3, one has $K^1|_{A_\eta} \in P^\chi(A_\eta)$. Then

$$\omega_\chi(K^1|_{A_\eta}) = H^0(A, {}^p\mathcal{H}^0(Rp_{A*}K) \otimes^L L_\chi). \quad (3.20)$$

Every fiber of $p_A : A \times X \rightarrow A$ has dimension $\dim X$, so by [BBDG82, 4.2.4], the functor

$$Rp_{A*}[-\dim X] : D_c^b(A \times X) \rightarrow D_c^b(A)$$

is left t-exact for the absolute perverse t-structures. From smoothness of X and Lemma 3.2.3.7, one has $K[\dim X] \in \text{Perv}(A \times X)$ and so $Rp_{A*}K \in {}^pD^{\geq 0}(A)$. Taking the perverse truncation, one has ${}^p\tau^{\leq 0}(Rp_{A*}K) = {}^p\mathcal{H}^0(Rp_{A*}K)$. Via the adjunction formula (see, e.g., [KW01, p.107]), the natural morphism

$${}^p\tau^{\leq 0}(Rp_{A*}K) \rightarrow Rp_{A*}K$$

in $D_c^b(A)$ (from the definition of t-structure) induces a morphism $h : K^1 \rightarrow K$ in $D_c^b(A \times X)$. Then h is a morphism in $\text{Perv}^{\text{ULA}}(A \times X/X)$. Let K^0 be the image of h in the abelian category $\text{Perv}^{\text{ULA}}(A \times X/X)$. By Fact 3.2.3.1 1, the functor $\text{Perv}(A \times X/X) \rightarrow \text{Perv}(A_\eta)$ is exact. Then $K^0|_{A_\eta}$ is the image of $h|_{A_\eta} : K^1|_{A_\eta} \rightarrow K|_{A_\eta}$ in $\text{Perv}(A_\eta)$.

Because $P^\chi(A_\eta)$ is an abelian subcategory of $\text{Perv}(A_\eta)$, by Condition 2, the image of $h|_{A_\eta}$ in $P^\chi(A_\eta)$ is still $K^0|_{A_\eta}$. As the functor (3.14) is exact, the image of $\omega_\chi(h|_{A_\eta}) : \omega_\chi(K^1|_{A_\eta}) \rightarrow \omega_\chi(K|_{A_\eta})$ is $\omega_\chi(K^0|_{A_\eta})$. Combining (3.17), (3.18), (3.19) with (3.20), one gets $\omega_\chi(K^0|_{A_\eta}) = \omega_{\chi_\eta}(K|_{A_\eta})^{\Gamma_{k(\eta)}}$. \square

3.5.3 Normality

By [JKLM23, Thm. 4.3], for every character $\chi \in \mathcal{C}(A)$, the geometric generic Tannakian group $G_{\omega_\chi}(K|_{A_\eta})$ is a normal closed subgroup of the generic Tannakian group $G_{\omega_\chi}(K|_{A_\eta})$. Theorem 3.5.3.1 shows that for uncountably many characters, the corresponding monodromy group is also a normal closed subgroup of the generic Tannakian group.

For every $\chi_{\ell'} \in \mathcal{C}(A)_{\ell'}$ and every $\chi_\ell \in \mathcal{C}(A)_\ell$, set $\chi = \chi_{\ell'}\chi_\ell$.

Theorem 3.5.3.1. *Assume $K \in \text{Perv}^{\text{ULA}}(A \times X/X)$ and $\dim A > 0$. Then for every $\chi_{\ell'} \in \mathcal{C}(A)_{\ell'}$, there is an uncountable subset $E \subset \mathcal{C}(A)_\ell(\bar{\mathbb{Q}}_\ell)$, such that for every $\chi_\ell \in E$,*

- *one has $K|_{A_\eta} \in P^\chi(A_\eta)$,*
- *the algebraic group $G_{\omega_\chi}(K|_{A_\eta})$ is reductive,*
- *and $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ is a normal closed subgroup of $G_{\omega_\chi}(K|_{A_\eta})$.*

We sketch the proof of Theorem 3.5.3.1. For every representation V of the Tannakian group $G(K|_{A_\eta})$ and every $\chi_{\ell'} \in \mathcal{C}(A)_{\ell'}$, by Theorem 3.1.2.3, there is a strict Zariski closed subset B_V of the cotorus $\mathcal{C}(A)_\ell$, such that for every $\chi_\ell \in (\mathcal{C}(A)_\ell \setminus B_V)(\bar{\mathbb{Q}}_\ell)$, the monodromy invariant $V^{\text{Mon}(K|_{A_\eta}, \chi_\eta)}$ is a $G(K|_{A_\eta})$ -subrepresentation. Choose $E = \mathcal{C}(A)_\ell(\bar{\mathbb{Q}}_\ell) \setminus \cup_V B_V(\bar{\mathbb{Q}}_\ell)$. From Lemma 3.5.0.1, normality holds when $\chi_\ell \in E$.

Proof. Both $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ and $G_{\omega_\chi}(K|_{A_\eta})$ depend only on the generic fiber of $p_X : A \times X \rightarrow X$. Therefore, shrinking X to a nonempty open subset does not change them. Thus, one may assume that X is smooth.

Claim 3.5.3.2. The object $K|_{A_\eta} \in \text{Perv}(A_\eta)$ is semisimple.

From Claim 3.5.3.2 and Lemma 3.5.3.6 1, the object $K|_{A_\eta} \in \bar{P}(A_\eta)$ is also semisimple. Therefore, a (hence every) Tannakian group of the neutral Tannakian category $\langle K|_{A_\eta} \rangle \subset \bar{P}(A_\eta)$ is a *reductive*, algebraic group over $\bar{\mathbb{Q}}_\ell$. Then by Lemma 3.5.3.5, there is a countable sequence of objects $\{\bar{K}_i\}_{i \geq 1}$, such that every object of $\langle K|_{A_\eta} \rangle$ is isomorphic to some \bar{K}_i . To apply Theorem 3.1.2.3, we need semisimple objects of $D_c^b(A \times X)$.

Claim 3.5.3.3. For every object $N \in \langle K|_{A_\eta} \rangle$, there is $L \in \text{Perv}^{\text{ULA}}(A \times X/X)$ that is semisimple in $D_c^b(A \times X)$, such that $L|_{A_\eta}$ is isomorphic to N in $\bar{P}(A_\eta)$.

From Claim 3.5.3.3, for every integer $i \geq 1$, there is $K_i \in \text{Perv}^{\text{ULA}}(A \times X/X)$ that is semisimple in $D_c^b(A \times X)$ with $K_i|_{A_\eta}$ isomorphic to \bar{K}_i in $\bar{P}(A_\eta)$. From smoothness of X and Theorem 3.1.2.3, there is a subobject $K_i^0 \subset K_i$ in $\text{Perv}^{\text{ULA}}(A \times X/X)$ and a strict Zariski closed subset $B_i \subset \mathcal{C}(A)_\ell$, such that for every $\chi_\ell \in (\mathcal{C}(A)_\ell \setminus B_i)(\bar{\mathbb{Q}}_\ell)$, one has $K_i|_{A_\eta} \in P^\times(A_\eta)$ and

$$\omega_{\chi_\eta}(K_i|_{A_\eta})^{\Gamma_{k(n)}} = \omega_{\chi_\eta}(K_i^0|_{A_\eta}). \quad (3.21)$$

Set $E := \mathcal{C}(A)_\ell(\bar{\mathbb{Q}}_\ell) \setminus \cup_{i \geq 1} B_i(\bar{\mathbb{Q}}_\ell)$. From Lemma 3.3.2.10 and the assumption $\dim A > 0$, the set E is uncountable. For every $\chi_\ell \in E$, one has $K|_{A_\eta} \in P^\times(A_\eta)$. For every $i \geq 1$, by $\chi_\ell \notin B_i(\bar{\mathbb{Q}}_\ell)$ and (3.21), the subspace $\omega_{\chi_\eta}(K_i|_{A_\eta})^{\text{Mon}(K|_{A_\eta}, \chi_\eta)}$ is $G_{\omega_\chi}(K|_{A_\eta})$ -stable. By Theorem 3.5.1.1 and Lemma 3.5.0.1, the subgroup $\text{Mon}(K|_{A_\eta}, \chi_\eta)$ of $G_{\omega_\chi}(K|_{A_\eta})$ is *normal*. \square

Proof of Claim 3.5.3.2. For every subobject $M \subset K|_{A_\eta}$ in $\text{Perv}(A_\eta)$, by Fact 3.2.3.5 and the smoothness of X , there is a subobject $K' \subset K$ in $\text{Perv}^{\text{ULA}}(A \times X/X)$ with $K'|_{A_\eta} = M$. By Lemma 3.2.3.7, the morphism $K'[\dim X] \rightarrow K[\dim X]$ is a monomorphism in $\text{Perv}(A \times X)$. Because K is semisimple in $D_c^b(A \times X)$, its shift $K[\dim X]$ is semisimple in $\text{Perv}(A \times X)$. Thus, there is a subobject $N \subset K[\dim X]$ in $\text{Perv}(A \times X)$ with

$$K[\dim X] = (K'[\dim X]) \oplus N.$$

Then $K = K' \oplus (N[-\dim X])$ in $D_c^b(A \times X)$. For every integer j , let ${}^{p/X}\mathcal{H}^j : D_c^b(A \times X) \rightarrow \text{Perv}(A \times X/X)$ be the j -th cohomology functor associated with the relative perverse t-structure. If $j \neq 0$, then

$$0 = {}^{p/X}\mathcal{H}^j(K) = 0 \oplus {}^{p/X}\mathcal{H}^j(N[-\dim X])$$

in $\text{Perv}(A \times X/X)$. Hence ${}^{p/X}\mathcal{H}^j(N[-\dim X]) = 0$ and

$$N[-\dim X] \in \text{Perv}(A \times X/X).$$

Consequently, $K|_{A_\eta} = M \oplus (N|_{A_\eta}[-\dim X])$ in $\text{Perv}(A_\eta)$. By [BBDG82, Thm. 4.3.1 (i)], the abelian category $\text{Perv}(A_\eta)$ is Noetherian and Artinian. As every subobject of $K|_{A_\eta}$ in $\text{Perv}(A_\eta)$ admits a direct complement, the semisimplicity follows from Lemma 3.4.1.2 2. \square

Proof of Claim 3.5.3.3. From Lemma 3.5.3.5, the object $N \in \bar{P}(A_\eta)$ is semisimple. There is an integer $n \geq 0$ such that N is a subquotient of $(K|_{A_\eta} \oplus K|_{A_\eta}^\vee)^{*n}$ in $\bar{P}(A_\eta)$.

We “globalize” the fiberwise convolution functors as follows. Define a bifunctor

$$\begin{aligned} *_X : D_c^b(A \times X) \times D_c^b(A \times X) &\rightarrow D_c^b(A \times X), \\ (-, +) &\mapsto R(m \times \text{Id}_X)_*(p_{13}^* - \otimes^L p_{23}^* +), \end{aligned} \quad (3.22)$$

where p_{ij} are the projections on $A \times A \times X$. By the proper base change theorem, for every $x \in X(k)$, one has $(- *_X +)|_{A_x} \xrightarrow{\sim} (-|_{A_x}) * (+|_{A_x})$ as bifunctors $D_c^b(A \times X) \times D_c^b(A \times X) \rightarrow D_c^b(A_x)$. Therefore, one has $(- *_X +)|_{A_\eta} \xrightarrow{\sim} (-|_{A_\eta}) * (+|_{A_\eta})$ as bifunctors $D_c^b(A \times X) \times D_c^b(A \times X) \rightarrow D_c^b(A_\eta)$.

The bifunctor (3.22) restricts to a bifunctor $D^{\text{ULA}}(A \times X/X) \times D^{\text{ULA}}(A \times X/X) \rightarrow D^{\text{ULA}}(A \times X/X)$. Indeed, for any $K', K'' \in D^{\text{ULA}}(A \times X/X)$, by Fact 3.2.2.2 6, one has

$$p_{13}^* K' \otimes^L p_{23}^* K'' \in D^{\text{ULA}}(A \times A \times X/X).$$

By Fact 3.2.2.2 4, one gets $K' *_X K'' \in D^{\text{ULA}}(A \times X/X)$.

Set $K^\vee := ([-1]_A \times \text{Id}_X)^* \mathbb{D}_{A \times X/X} K$. By Fact 3.2.3.1 3, one has $K^\vee \in \text{Perv}^{\text{ULA}}(A \times X/X)$ and $(K^\vee)|_{A_\eta} = (K|_{A_\eta})^\vee$. Then

$$(K \oplus K^\vee)^{*xn} \in D^{\text{ULA}}(A \times X/X).$$

Set $M := {}^{p/X}\mathcal{H}^0((K \oplus K^\vee)^{*xn}) \in \text{Perv}^{\text{ULA}}(A \times X/X)$. Then $M|_{A_\eta} = {}^p\mathcal{H}^0([K|_{A_\eta} \oplus (K|_{A_\eta})^\vee]^{*n})$ in $\text{Perv}(A_\eta)$. By Lemma 3.5.3.6 3, there is a semisimple subquotient L' of $M|_{A_\eta}$ in $\text{Perv}(A_\eta)$, whose image in $\bar{P}(A_\eta)$ is N . By smoothness of X and Fact 3.2.3.5, there is a semisimple subquotient L of M in $\text{Perv}^{\text{ULA}}(A \times X/X)$ with $L|_{A_\eta} = L'$. By smoothness of X and Lemma 3.2.3.11 2, the object $L[\dim X]$ is semisimple in $\text{Perv}(A \times X)$. Then L is semisimple in $D_c^b(A \times X)$. \square

Remark 3.5.3.4. When $A = \operatorname{Spec}(k)$, the bifunctor (3.22) becomes $\otimes^L : D_c^b(X) \times D_c^b(X) \rightarrow D_c^b(X)$. The derived tensor product may not preserve semisimplicity in the category $D_c^b(X)$. That is why we need semisimplicity in $\operatorname{Perv}^{\operatorname{ULA}}(A \times X/X)$ in the last paragraph of the proof of Claim 3.5.3.3.

In fact, consider $k = \mathbb{C}$, $X = A^1$ and $U = X \setminus \{0\}$. Let $j : U \rightarrow X$ be the inclusion. Then $\pi_1^{\text{ét}}(U, 1) = \hat{\mathbb{Z}}$. The unique surjective morphism $\pi_1^{\text{ét}}(U, 1) \rightarrow \mathbb{Z}/2$ corresponds to a rank one lisse $\bar{\mathbb{Q}}_\ell$ -sheaf L on U . Then $L \otimes^L L \cong \bar{\mathbb{Q}}_{\ell,U}$, and L^{an} is a $\bar{\mathbb{Q}}_\ell$ -local system on $U^{\text{an}} = \mathbb{C} \setminus \{0\}$.

Let U_0 be a punctured ball in $X^{\text{an}} = \mathbb{C}$ centered at 0 containing 1. One has $H^0(U_0, L^{\text{an}}) = (L^{\text{an}})^{\pi_1^{\text{top}}(U_0, 1)} = 0$, and $H^1(U_0, L^{\text{an}})$ coincides with the group cohomology $H^1(\pi_1^{\text{top}}(U_0, 1), L_1^{\text{an}})$, where the $\pi_1^{\text{top}}(U_0, 1) = \mathbb{Z}$ -action on the stalk L_1^{an} is the monodromy. For every crossed homomorphism $f : \mathbb{Z} \rightarrow L_1^{\text{an}}$, every integer j , one has $f(1+j) = f(1) - f(j)$. Therefore, when j is even (resp. odd), $f(j)$ is 0 (resp. $f(1)$). In particular, f is a boundary and hence $H^1(\pi_1^{\text{top}}(U_0, 1), L_1^{\text{an}}) = 0$. Thus, L^{an} is *not* in the cohomology support loci of U_0 . From [BLSW17, p.299], $j_1^{\text{an}} L^{\text{an}}[1]$ is a simple object of $\operatorname{Perv}(X^{\text{an}})$. Set $M := j_! L[1]$. By [BBDG82, p.150], the natural morphism $j_{!*} L[1] \rightarrow M$ is an isomorphism in $D_c^b(X)$. In particular, M is a *simple* object of $\operatorname{Perv}(X)$.

From [KW01, II, Cor. 7.5 g)], one has

$$N := M \otimes^L M = j_!(L \otimes^L j^* j_! L)[2] = j_! \bar{\mathbb{Q}}_{\ell,U}[2].$$

By [HT07, Example 8.1.35 (ii)], one has $N[-1] \in \operatorname{Perv}(X)$. Let $i : 0 \rightarrow A^1$ be the inclusion. From the short exact sequence

$$0 \rightarrow j_! \bar{\mathbb{Q}}_{\ell,U} \rightarrow \bar{\mathbb{Q}}_{\ell,X} \rightarrow i_*(\bar{\mathbb{Q}}_{\ell,0}) \rightarrow 0$$

in $\operatorname{Cons}(X)$, one gets an exact sequence

$${}^p\mathcal{H}^0(\bar{\mathbb{Q}}_{\ell,X}) \rightarrow {}^p\mathcal{H}^0(i_*(\bar{\mathbb{Q}}_{\ell,0})) \rightarrow {}^p\mathcal{H}^1(j_! \bar{\mathbb{Q}}_{\ell,U}) \rightarrow {}^p\mathcal{H}^1(\bar{\mathbb{Q}}_{\ell,X}) \rightarrow {}^p\mathcal{H}^1(i_*(\bar{\mathbb{Q}}_{\ell,0}))$$

in $\operatorname{Perv}(X)$. Since $i_*(\bar{\mathbb{Q}}_{\ell,0}), \bar{\mathbb{Q}}_{\ell,X}[1] \in \operatorname{Perv}(X)$, it gives a short exact sequence

$$0 \rightarrow i_*(\bar{\mathbb{Q}}_{\ell,0}) \rightarrow N[-1] \rightarrow \bar{\mathbb{Q}}_{\ell,X}[1] \rightarrow 0$$

in $\operatorname{Perv}(X)$. This sequence does not split as $N[-1]$ is supported on U . Therefore, $N[-1]$ is not a semisimple object of $\operatorname{Perv}(X)$. It follows that $M \otimes^L M$ is not semisimple in $D_c^b(X)$. (This sequence also shows that the support of a perverse sheaf may be smaller than that of a subquotient.)

For a category \mathcal{C} , let \mathcal{C}/\sim be the class of isomorphism classes of objects in \mathcal{C} .

Lemma 3.5.3.5. *Let (\mathcal{C}, \otimes) be a neutral Tannakian category over k with a fiber functor $\omega : \mathcal{C} \rightarrow \operatorname{Vec}_k$. Assume that $\operatorname{Aut}^{\otimes}(\mathcal{C}, \omega)$ is a reductive, algebraic group over k . Then the underlying abelian category is semisimple, and \mathcal{C}/\sim is countable.*

Proof. Set $G = \text{Aut}^\otimes(\mathcal{C}, \omega)$. Let $\text{Rep}(G)$ be the category of k -rational representations of G . Then \mathcal{C} is equivalent to $\text{Rep}(G)$. Because k has characteristic zero, by [Mil17a, Cor. 22.43], the abelian category $\text{Rep}(G)$ is semisimple. As k is algebraically closed, by [AHR20, Thm. 2.16], there is an at most countable set X^+ and for every $\lambda \in X^+$, a unital k -algebra \mathcal{A}^λ with the following property: The set $\text{Irr}(G)$ of isomorphism classes of simple objects of $\text{Rep}(G)$ is in bijection with the set of pairs (λ, E) , where $\lambda \in X^+$ and E is an isomorphism class of simple left \mathcal{A}^λ -modules. For every $\lambda \in X^+$, from [AHR20, Lem. 2.19], the algebra \mathcal{A}^λ is semisimple. Then by [Lan02, XVII, Thm. 4.3, Cor. 4.5], the set of isomorphism classes of simple left \mathcal{A}^λ -modules is finite. Therefore, $\text{Irr}(G)$ is at most countable. Consequently, $\text{Rep}(G)/\sim$ is countable. \square

Lemma 3.5.3.6. *Let \mathcal{A} be an abelian category. Let $\mathcal{B} \subset \mathcal{A}$ be a Serre subcategory. Consider the quotient functor $F : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{B}$.*

1. *Let $X \in \mathcal{A}$. Let $i : Y \rightarrow F(X)$ be a monomorphism in \mathcal{A}/\mathcal{B} . Then there is a monomorphism $j : Z \rightarrow X$ in \mathcal{A} and an isomorphism $u : Y \rightarrow F(Z)$ in \mathcal{A}/\mathcal{B} fitting into a commutative diagram in \mathcal{A}/\mathcal{B}*

$$\begin{array}{ccc} & F(Z) & \\ u \nearrow & & \searrow F(j) \\ Y & \xrightarrow{i} & F(X). \end{array}$$

Dually, up to isomorphism every quotient in \mathcal{A}/\mathcal{B} lifts to a quotient in \mathcal{A} . In particular, if $X \in \mathcal{A}$ is a simple object, then $F(X)$ is either simple or zero in \mathcal{A}/\mathcal{B} .

2. *Let $V \in \mathcal{A}$ be a Noetherian and Artinian object. If $F(V)$ is simple in \mathcal{A}/\mathcal{B} , then there is a simple subquotient W of V in \mathcal{A} such that $F(W)$ is isomorphic to $F(V)$ in \mathcal{A}/\mathcal{B} .*
3. *Assume that \mathcal{A} is Noetherian and Artinian. Let $X \in \mathcal{A}$. If Y is a simple subquotient of $F(X)$ in \mathcal{A}/\mathcal{B} , then there is a simple subquotient W of X , with $F(W)$ isomorphic to Y in \mathcal{A}/\mathcal{B} .*

Proof.

1. By the construction in the proof of [Sta24, Tag 02MS] and the right calculus of fractions in [Sta24, Tag 04VB], there is a diagram

$$\begin{array}{ccc} & M & \\ f \swarrow & & \searrow g \\ Y & & X \end{array}$$

in \mathcal{A} , such that $F(f)$ is an isomorphism and $F(g) = i \circ F(f)$ in \mathcal{A}/\mathcal{B} . Therefore, $F(g)$ is a monomorphism. Since F is exact, one has $F(\ker(g)) = \ker(F(g)) = 0$, so $\ker(g) \in \mathcal{B}$. Let $q : M \rightarrow M/\ker(g)$ be the epimorphism in \mathcal{A} , and let $j : M/\ker(g) \rightarrow X$ be the monomorphism in \mathcal{A} induced by g . Then $F(q)$ is an isomorphism in \mathcal{A}/\mathcal{B} . Set $u : Y \rightarrow F(M/\ker(g))$ to be the morphism $F(q) \circ F(f)^{-1}$ in \mathcal{A}/\mathcal{B} . Then u is an isomorphism with the stated property.

2. Let \mathcal{P} be the family of subobjects V' of V in \mathcal{A} with $V/V' \in \mathcal{B}$. Then \mathcal{P} is nonempty since $V \in \mathcal{P}$. As V is Artinian in \mathcal{A} , there is a minimal object $U \in \mathcal{P}$. Moreover, the morphism $F(U) \rightarrow F(V)$ is an isomorphism in \mathcal{A}/\mathcal{B} . Let \mathcal{Q} be the family of subobjects of $U \in \mathcal{A}$ lying in \mathcal{B} . Then \mathcal{Q} is nonempty since $0 \in \mathcal{Q}$. As V is Noetherian in \mathcal{A} , so is U . Thus, \mathcal{Q} has a maximal object U_0 . Then $W := U/U_0$ is a subquotient of $V \in \mathcal{A}$ and the morphism $F(U) \rightarrow F(W)$ is an isomorphism in \mathcal{A}/\mathcal{B} . In particular, $W \neq 0$ in \mathcal{A} .

We claim that W is simple in \mathcal{A} . Indeed, let $U' \rightarrow W$ be a subobject in \mathcal{A} . Then there is a subobject U'' of U in \mathcal{A} containing U_0 with $U''/U_0 = U'$. As $F(U'')$ is a subobject of a simple object $F(U)$ in \mathcal{A}/\mathcal{B} , either the morphism $F(U'') \rightarrow F(U)$ is an isomorphism or $F(U'') = 0$. If $F(U'') = 0$, then $U'' \in \mathcal{B}$ and $U'' \in \mathcal{Q}$. Since U_0 is maximal in \mathcal{Q} , one has $U_0 = U''$, so $U' = 0$. If $F(U'') \rightarrow F(U)$ is an isomorphism, then $U/U'' \in \mathcal{B}$. Since the sequence

$$0 \rightarrow U/U'' \rightarrow V/U'' \rightarrow V/U \rightarrow 0$$

is exact in \mathcal{A} , and \mathcal{B} is closed under extensions, one gets $V/U'' \in \mathcal{B}$ and $U'' \in \mathcal{P}$. Since U is minimal in \mathcal{P} , one has $U'' = U$. The morphism $U' \rightarrow W$ is thus an isomorphism in \mathcal{A} . The claim is proved.

3. By 1, there is a subquotient Z of X in \mathcal{A} with $F(Z)$ isomorphic to Y . Then $F(Z)$ is simple in \mathcal{A}/\mathcal{B} . By assumption, Z is Noetherian and Artinian in \mathcal{A} . Thus from 2, there is a simple subquotient W of Z in \mathcal{A} with $F(W)$ isomorphic to $F(Z)$ and to Y in \mathcal{A}/\mathcal{B} .

□

Example 3.5.3.7. Let $s : X \rightarrow A \times X$ be a section to $p_X : A \times X \rightarrow X$. Let F be a lisse $\bar{\mathbb{Q}}_\ell$ -sheaf on X , and let $\sigma : \pi_1^{\text{ét}}(X, \bar{\eta}) \rightarrow \text{GL}(F_{\bar{\eta}})$ be the corresponding monodromy representation. By Fact 3.2.2.2 2, one has $F \in D^{\text{ULA}}(X/X)$. Then from Fact 3.2.2.2 4, one has $K := Rs_* F \in D^{\text{ULA}}(A \times X/X)$. For every $x \in X(k)$, by the proper base change theorem, $K|_{A_x} \in D_c^b(A_x)$ is the skyscraper supported at the closed point $s(x) \in A_x$ with stalk F_x . Thus, $K|_{A_x} \in \text{Perv}(A_x)$ and $K \in \text{Perv}^{\text{ULA}}(A \times X/X)$. Moreover, $K|_{A_{\bar{\eta}}} is the skyscraper supported at $s(\bar{\eta}) \in A_{\bar{\eta}}$ with stalk $F_{\bar{\eta}}$. Therefore, the generic$

and the geometric generic Tannakian groups agree and are computed in Example 3.4.3.3.

For every $\chi \in \mathcal{C}(A)$, by Fact 3.2.1.1, one has $K \otimes^L p_A^* L_\chi = Rs_*(F \otimes^L s^* p_A^* L_\chi)$. Thus, $Rp_{X*}(K \otimes^L p_A^* L_\chi) = F \otimes^L s^* p_A^* L_\chi$ is a lisse $\bar{\mathbb{Q}}_\ell$ -sheaf on X . The corresponding $\pi_1^{\text{ét}}(X, \bar{\eta})$ -representation is the tensor product of σ with the character

$$\pi_1^{\text{ét}}(X, \bar{\eta}) \xrightarrow{(p_A^{\text{cs}})^*} \pi_1^{\text{ét}}(A, p_{As}(\bar{\eta})) \xrightarrow{\sim} \pi_1^{\text{ét}}(A, 0) \xrightarrow{\chi} \bar{\mathbb{Q}}_\ell^\times.$$

The $\Gamma_{k(\eta)}$ -representation induced by pulling back along $\eta_* : \Gamma_{k(\eta)} \rightarrow \pi_1^{\text{ét}}(X, \bar{\eta})$ is $\omega_{\chi_\eta}(K|_{A_\eta})$.

Assume that F is semisimple in $\text{Loc}(X)$. Then $F[\dim X]$ is a semisimple object of $\text{Perv}(X)$. By Fact 3.2.1.6 1, the object K is semisimple in $D_c^b(A \times X)$.

Chapter 4

Generic vanishing theorem for Fujiki class \mathcal{C}

4.1 Introduction

Recall the historical origin of generic vanishing results. In the last paragraph of [Enr39], Enriques gave an upper bound on the dimension of the paracanonical system of curves on some class of algebraic surfaces. However, in [Enr49, p.354] he pointed out a mistake in the proof of his result as well as a similar theorem by Severi [Sev42]. Catanese [Cat83, p.103] posed Conjecture 4.1.0.1.

Conjecture 4.1.0.1. *For a smooth projective surface S/\mathbb{C} without irrational pencils, the dimension of the paracanonical system $\{K_S\}$ is at most the geometric genus $p_g(S)$.*

In 1987, Green and Lazarsfeld [GL87, Theorem 4.2] provided a positive answer to Conjecture 4.1.0.1. Its proof uses a result ([GL87, Prop. 4.1]) of generic vanishing type.

As is explained in [Uen83, pp.619–620], the dimension of $\{K_S\}$ in Conjecture 4.1.0.1 is related to Conjecture 4.1.0.2, which is also of generic vanishing type.

Conjecture 4.1.0.2 ([Uen83, Problem 8, p.620]). *Let X be a projective manifold and $\alpha : X \rightarrow \text{Alb}(X)$ be an Albanese morphism. If $\dim \alpha(X) > 1$, then $H^1(X, L) = 0$ for generic $L \in \text{Pic}^0(X)$.*

Green and Lazarsfeld [GL87] proved a strengthening of Conjecture 4.1.0.2. Since then, the theory of generic vanishing results has been very much investigated and numerous authors have contributed to its development, so the overview in Section 4.1.1 is by no means complete.

For a finitely generated \mathbb{Z} -module H , let H_{tor} be the submodule of H comprised of torsion elements and $H_{\text{free}} := H/H_{\text{tor}}$. We use the words

“locally free sheaf” and “vector bundle” interchangeably. Let $F \rightarrow X$ be a (holomorphic) vector bundle on a complex manifold. The dimension of a complex space always means the complex dimension. For any three integers $p, q, m \geq 0$, the corresponding jumping locus is defined as

$$S_m^{p,q}(X, F) := \{L \in \text{Pic}^0(X) : h^q(X, \Omega_X^p \otimes_{O_X} L \otimes_{O_X} F) \geq m\}.$$

For simplicity, p (resp. m , resp. F) is omitted when $p = 0$ (resp. $m = 1$, resp. $F = O_X$). Roughly speaking, generic vanishing results show that these loci are small (in some sense) and study their structure when F is flat unitary (in the sense of Definition 4.2.2.2).

4.1.1 Known results

Let X be a connected compact Kähler manifold, $\alpha : X \rightarrow \text{Alb}(X)$ be the Albanese map associated with some base point and $F \rightarrow X$ be a flat unitary vector bundle. Each locus $S_m^{p,q}(X, F)$ is an analytic subset of the complex torus $\text{Pic}^0(X)$ (see the proof of Theorem 4.7.1.3 1) and “generic” means outside a strict analytic subset. In the literature, generic vanishing results concerning $S^q(X, F)$ (resp. $S^{p,q}(X, F)$) are usually called of Kodaira type (resp. Nakano type). Such results typically involve the following invariants:

- $\dim \alpha(X)$;
- $w(X) := \max\{\text{codim}(Z(\eta), X) : 0 \neq \eta \in H^0(X, \Omega_X^1)\}$, where $Z(\eta)$ denotes the zero-locus of the 1-form η ;
- the defect of semismallness $r(\alpha)$ of α (in the sense of Definition 4.5.2.1).

Using deformation theory of cohomology groups, Green and Lazarsfeld [GL87, Remarks (1), p.401] proves Fact 4.1.1.1, which is of Kodaira type and implies Conjecture 4.1.0.2.

Fact 4.1.1.1. *For every integer $k \geq 0$, one has*

$$\text{codim}_{\text{Pic}^0(X)}(S^k(X, F)) \geq \dim \alpha(X) - k.$$

In particular, if $k < \dim \alpha(X)$, then $H^k(X, F \otimes_{O_X} L) = 0$ for a generic line bundle $L \in \text{Pic}^0(X)$.

Green and Lazarsfeld also give a Nakano-type generic vanishing theorem.

Fact 4.1.1.2 ([GL87, Remarks (1), p.404]). *For any integers $i, j \geq 0$ with $i + j < w(X)$, one has $S^{i,j}(X, F) \neq \text{Pic}^0(X)$.*

In another direction, there are known results concerning the structure of the jumping loci.

Fact 4.1.1.3 ([GL91, Thm. 0.1 (1)]). *For any two integers $k, m \geq 0$, the subset $S_m^k(X)$ is a finite union of translates of subtori of $\text{Pic}^0(X)$.*

Beauville and Catanese conjectured that for every integer $q \geq 0$, $S^q(X)$ is a finite union of *torsion* translates of subtori of $\text{Pic}^0(X)$ ([Cat91, Problem 1.25] and [Bea92, p.1]). When X is a projective manifold, this conjecture is proved by Simpson [Sim93, Sec. 5].

Fact 4.1.1.4 (Simpson). *If X is furthermore projective, then for any two integers $k, m \geq 0$, the locus $S_m^k(X)$ is a finite union of torsion translates of subtori of $\text{Pic}^0(X)$.*

Some arguments of [Sim93] are of arithmetic nature, so they do not apply to the Kähler case. Campana [Cam01, Sec. 1.5.2] provided a partial answer for not only Kähler manifolds but also for Fujiki class \mathcal{C} (Definition 4.7.1.1).

Later on, Wang [Wan16, Cor. 1.4] answered affirmatively Beauville and Catanese's conjecture in full generality.

Fact 4.1.1.5 (Wang). *For any three integers $p, q, m \geq 0$, the subset $S_m^{p,q}(X)$ of $\text{Pic}^0(X)$ is a finite union of torsion translates of subtori.*

Hacon [Hac04, Cor. 4.2] uses Fourier-Mukai transforms of coherent modules on complex abelian varieties to recover Fact 4.1.1.1 when X is a projective manifold. This algebraic viewpoint sheds new insight on this topic. Similarly, as a byproduct of the theory on convolution of perverse sheaves on abelian varieties, Krämer and Weissauer obtain a Nakano-type generic vanishing theorem. The proof of [KW15b, Thm. 3.1] gives Fact 4.1.1.6.

Fact 4.1.1.6. *If furthermore the Albanese torus $\text{Alb}(X)$ is algebraic, then for any two integers $p, q \geq 0$ with $p + q < \dim X - r(\alpha)$, the locus $S^{p,q}(X, F)$ is contained in a finite union of translates of strict subtori of $\text{Pic}^0(X)$.*

Around the same time, by different methods Popa and Schnell [PS13, Thm. 1.2] obtained precise codimension bounds.

Fact 4.1.1.7. *If furthermore X is a projective manifold, then*

$$\text{codim}_{\text{Pic}^0(X)}(S^{p,q}(X)) \geq |p + q - \dim X| - r(\alpha)$$

for any two integers $p, q \geq 0$. Moreover, for every X there exist p and q for which the inequality becomes an equality.

4.1.2 The main result and a sketch

An algebraic variety means an integral scheme of finite type and separated over a field. Even though not necessarily Kähler, a complex smooth proper algebraic variety also admits Hodge theory ([Del68, Prop. 5.3], [Del71a, Thm. 3.2.5]). It is natural to ask if generic vanishing results also hold for such varieties. The aim of this note is to show that generic vanishing result is not only true for Kähler manifolds, but also for complex manifolds in Fujiki class \mathcal{C} . This class contains compact Kähler manifolds as well as smooth proper algebraic varieties.

Here is the main result that is of Nakano type.

Theorem (Theorem 4.7.1.3). *Let X be an n -dimensional complex manifold in Fujiki class \mathcal{C} with an Albanese morphism $\alpha : X \rightarrow \text{Alb}(X)$, and let F be a flat unitary vector bundle on X . Then for any two integers $p, q \geq 0$ with $p + q < n - r(\alpha)$, the locus $S^{p,q}(X, F)$ is a strict analytic subset of the complex torus $\text{Pic}^0(X)$.*

For smooth proper algebraic varieties, the following finer result follows from Corollary 4.7.2.6 and Lemma 4.6.1.2. It is not immediate from previously known generic vanishing results.

Corollary 4.1.2.1. *Let X be an n -dimensional smooth proper, complex algebraic variety with an algebraic Albanese morphism $\alpha : X \rightarrow \text{Alb}(X)$. Let \mathcal{L} be a unitary local system on the analytification X^{an} , and let $F = \mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_{X^{\text{an}}}$ be the corresponding holomorphic vector bundle. Then, for any two integers $p, q \geq 0$ with $p + q < n - r(\alpha)$, the subset $S^{p,q}(X, F)$ is contained in a finite union of translates (torsion translates if \mathcal{L} is semisimple of geometric origin in the sense of [BBDG82, p.163]) of strict abelian subvarieties of the Picard variety $\text{Pic}_{X/\mathbb{C}}^0$.*

Here is the outline of the proof of Theorem 4.7.1.3. By the Riemann-Hilbert correspondence restricted to unitary objects, we pass from flat unitary vector bundles to unitary local systems. The corresponding cohomology groups are related by Hodge decomposition (Fact 4.7.1.2). In this way, the initial generic vanishing problem for a flat unitary vector bundle twisted by line bundles is reduced to a generic vanishing problem for a unitary local system twisted by rank 1 local systems.

By pushing forward along the Albanese map, the problem about the local system on a manifold in Fujiki class \mathcal{C} is converted to a problem about a complex of sheaves on a complex torus. The last problem is solved by Krämer and Weissauer [KW15b] for perverse sheaves (on complex abelian varieties) and by the subsequent generalization (to all complex tori) due to Bhatt, Schnell and Scholze [BSS18].

This text is organized as follows. Section 4.2 reviews the unitary Riemann-Hilbert correspondence. Section 4.3 and 4.4 construct the Jacobian

and the Albanese map for regular manifolds, relaxing the usual Kähler condition. Several definitions of defect of semismallness are proved to be equivalent in Section 4.5. The work of Krämer and Weissauer on generic vanishing for perverse sheaves is recalled in Section 4.6. Finally in Section 4.7, the previous results are applied to prove the main result, Theorem 4.7.1.3, for Fujiki class \mathcal{C} .

4.2 Riemann-Hilbert correspondence

In Section 4.2, we review how the classical Riemann-Hilbert correspondence restricts to an equivalence between unitary local systems and flat unitary vector bundles on complex manifolds. The reason to introduce this restricted equivalence is that unitary local systems on manifolds in Fujiki class \mathcal{C} admit Hodge decomposition (Fact 4.7.1.2).

4.2.1 Unitary local systems

Let X be a path-connected, locally path-connected and locally simply connected topological space with a base point $x_0 \in X$. Let $\text{Loc}(X)$ be the category of local systems (of finite dimensional \mathbb{C} -vector spaces) on X . Let $\pi_1(X, x_0)$ be the fundamental group of X at x_0 and $\text{Rep}_{\mathbb{C}}(\pi_1(X, x_0))$ be the category of its finite dimensional complex representations. By [Del70, Cor. 1.4, p.4], the functor taking the stalk at x_0 gives rise to an equivalence

$$\text{Loc}(X) \rightarrow \text{Rep}_{\mathbb{C}}(\pi_1(X, x_0)) \quad (4.1)$$

compatible with tensor products. The image under (4.1) of a local system on X is called the corresponding monodromy representation.

Every compact subgroup of $\text{GL}_r(\mathbb{C})$ can be conjugated into the unitary subgroup $U_r(\mathbb{C})$. Therefore, for every representation $\rho : \pi_1(X, x_0) \rightarrow \text{GL}(V)$, the following conditions are equivalent:

1. The closure of $\rho(\pi_1(X, x_0))$ inside $\text{GL}(V)$ is compact;
2. There is a hermitian inner product $h : V \otimes_{\mathbb{C}} \bar{V} \rightarrow \mathbb{C}$ such that $\rho(\pi_1(X, x_0))$ is contained in the corresponding unitary group $U(V, h)$.

When the conditions hold, ρ is called *unitary*. Let $\text{Rep}_{\mathbb{C}}^u(\pi_1(X, x_0)) \subset \text{Rep}_{\mathbb{C}}(\pi_1(X, x_0))$ be the full subcategory of unitary representations. Let $\text{Loc}^u(X)$ be the full subcategory of $\text{Loc}(X)$ corresponding to $\text{Rep}_{\mathbb{C}}^u(\pi_1(X, x_0))$ via the equivalence (4.1). Its objects are called unitary local systems on X . Every unitary local system is semisimple, since every unitary representation is so.

4.2.2 Flat unitary bundles

Let $E \rightarrow X$ be a holomorphic vector bundle on a complex manifold with a hermitian metric h . By [Huy05, Prop. 4.2.14], there exists a unique hermitian connection ∇_h that is compatible with the holomorphic structure (in the sense of [Huy05, Def. 4.2.12], i.e., $\nabla^{0,1} = \bar{\partial}^E$), which is called the Chern connection of (E, h) . The corresponding curvature form, called the Chern curvature, is an $\text{End}(E, h)$ -valued $(1, 1)$ -form, (see, e.g., [Huy05, Prop. 4.3.8 iii]).

For every integer $k \geq 0$ (resp. any two integers $i, j \geq 0$), let A_X^k (resp. $A_X^{i,j}$) be the sheaf of *smooth* k (resp. (i, j)) forms on X . Then there is a direct sum decomposition $A_X^k = \bigoplus_{i+j=k} A_X^{i,j}$. In general, a (smooth) *flat* connection ∇ on E that is compatible with the holomorphic structure needs not to be a holomorphic connection (in the sense of [Huy05, Def. 4.2.17]).

Lemma 4.2.2.1. *Let $E \rightarrow X$ be a holomorphic vector bundle with a flat connection $\nabla : E \rightarrow E \otimes_{A_X^0} A_X^1$. If ∇ is compatible with the holomorphic structure, then ∇ is a holomorphic connection.*

Proof. Take a local holomorphic frame $\{e_1, \dots, e_r\}$ of E , and denote the corresponding local smooth connection matrix 1-form by Ω . As $\nabla^{0,1} = \bar{\partial}^E$, one has $\Omega^{0,1} = 0$. By flatness, $d\Omega + \Omega \wedge \Omega = 0$. Taking the $(1, 1)$ part of it, one gets $\bar{\partial}\Omega = 0$, i.e., Ω is a holomorphic form. This shows that ∇ is holomorphic. \square

Let $\text{Mod}(O_X)$ be the category of O_X -modules, and let $\text{VB}(X) \subset \text{Mod}(O_X)$ be the full subcategory of finite locally free O_X -modules. Let $\text{DE}(X)$ be the category of holomorphic vector bundles with a flat *holomorphic* connection. Forgetting the connection gives a functor $\text{DE}(X) \rightarrow \text{VB}(X)$. Let $\text{DE}^u(X) \subset \text{DE}(X)$ be the full subcategory comprised of objects (F, ∇) such that there exists a hermitian metric on F whose Chern connection is ∇ .

Definition 4.2.2.2. An object in the essential image of $\text{DE}^u(X)$ under the forgetful functor $\text{DE}(X) \rightarrow \text{VB}(X)$ is called a *flat unitary* vector bundle on X .

From [Huy05, Eg. 4.2.15], the trivial line bundle O_X is flat unitary. By Lemma 4.2.2.1, a holomorphic vector bundle is flat unitary if and only if it admits a hermitian metric whose Chern connection is flat.

4.2.3 An equivalence

Let X be a connected complex manifold. By the Riemann-Hilbert correspondence ([Del70, Thm. 2.17, p.12]), the pair of functors

$$\mathrm{Loc}(X) \rightarrow \mathrm{DE}(X), \quad \mathcal{L} \mapsto (\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_X, \mathrm{Id}_{\mathcal{L}} \otimes d); \quad (4.2)$$

$$\mathrm{DE}(X) \rightarrow \mathrm{Loc}(X), \quad (E, \nabla) \mapsto \ker(\nabla) \quad (4.3)$$

forms an equivalence of categories. It is compatible with tensor products and preserves the rank.

Theorem 4.2.3.1 (Unitary Riemann-Hilbert correspondence). *The equivalence (4.2), (4.3) restricts to an equivalence between $\mathrm{Loc}^u(X)$ and $\mathrm{DE}^u(X)$.*

Proof. First, we prove that the functor (4.2) sends $\mathrm{Loc}^u(X)$ to $\mathrm{DE}^u(X)$. Consider a unitary local system \mathcal{L} on X . Since the corresponding monodromy representation is unitary, we may choose a hermitian inner product h_{x_0} on the stalk \mathcal{L}_{x_0} such that the representation factors through $U(\mathcal{L}_{x_0}, h_{x_0})$. For any $x \in X$, choose a path γ from x_0 to x and propagate h_{x_0} along this curve, i.e., using the linear isomorphism $\gamma_* : \mathcal{L}_{x_0} \rightarrow \mathcal{L}_x$ induced by γ , we translate h_{x_0} to a hermitian inner product h_x of \mathcal{L}_x . This h_x is independent of the choice of γ by assumption. Hence a positive definite hermitian form h on \mathcal{L} that is invariant under the monodromy action. Then h extends naturally to a (smooth) hermitian metric h' on the associated holomorphic vector bundle $\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_X$ on X and the corresponding flat holomorphic connection $\mathrm{Id}_{\mathcal{L}} \otimes d$ is a hermitian connection. Therefore, $\mathrm{Id}_{\mathcal{L}} \otimes d$ is the Chern connection of $(\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_X, h')$ and $(\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_X, \mathrm{Id}_{\mathcal{L}} \otimes d) \in \mathrm{DE}^u(X)$.

Conversely, we prove that the functor (4.3) sends $\mathrm{DE}^u(X)$ to $\mathrm{Loc}^u(X)$. Consider a holomorphic hermitian vector bundle (E, h) on X whose Chern connection ∇_h is flat. Around every point we can find a local ∇_h -horizontal holomorphic frame $\{e_1, \dots, e_r\}$ of E . For any $1 \leq i, j \leq r$, since the connection ∇_h is compatible with h , we have

$$d[h(e_i, e_j)] = h(\nabla_h e_i, e_j) + h(e_i, \nabla_h e_j) = 0.$$

Therefore, the local function $h(e_i, e_j)$ is locally constant and the parallel transport along every closed path on X preserves the hermitian inner products on the fibers of E . The sheaf $\ker(\nabla_h)$ of horizontal sections of E forms a local system on X , whose stalks are exactly the fibers of E . Thus, it admits a monodromy-invariant positive definite hermitian form and is consequently unitary.¹ \square

¹The definition of unitary local system in [Tim87, p.152] seems to forget this invariance.

4.3 Hodge theory and Jacobian

In Section 4.3, we review the definition of Jacobian and show that for every complex manifold admitting Hodge theory (Definition 4.3.1.1), its Jacobian has nice expected properties.

4.3.1 Regular manifolds

Let X be a complex manifold. Let $d : A_X^\bullet \rightarrow A_X^{\bullet+1}$ be the exterior derivative. Then $d = \partial + \bar{\partial}$, where $\partial : A_X^{\bullet,\bullet} \rightarrow A_X^{\bullet+1,\bullet}$ and $\bar{\partial} : A_X^{\bullet,\bullet} \rightarrow A_X^{\bullet,\bullet+1}$ are the $(1,0)$ and $(0,1)$ part of d respectively. For every $\mathcal{E} \in \text{Loc}^u(X)$, every integer $k \geq 0$, define a decreasing filtration of $A_X^k \otimes_{\mathbb{C}} \mathcal{E}$ by

$$F^p = F^p(A_X^k \otimes_{\mathbb{C}} \mathcal{E}) := \oplus_{i \geq p} A_X^{i,k-i} \otimes_{\mathbb{C}} \mathcal{E}. \quad (4.4)$$

Then $(d \otimes \text{Id}_{\mathcal{E}})(F^p) \subset F^p$. Therefore, this filtration induces a spectral sequence, called the Frölicher spectral sequence:

$$E_1^{p,q} = H^q(X, \Omega_X^p \otimes_{\mathbb{C}} \mathcal{E}) \Rightarrow H^{p+q}(X, \mathcal{E}), \quad (4.5)$$

where the differential $d_1^{p,q} : E_1^{p,q} \rightarrow E_1^{p+1,q}$ is induced by the operator $\partial : A_X^{p,q} \rightarrow A_X^{p+1,q}$ on X . It is the classical notion in [Voi02, Sec. 8.3.3] when \mathcal{E} is the constant sheaf \mathbb{C}_X .

Although the Hodge theory for the first cohomology groups H^1 suffices for most properties of the Jacobian and the Albanese, in the sequel we mainly work with manifolds admitting Hodge theory in all degrees. Such manifolds are called “regular” for convenience.

Definition 4.3.1.1 (Regular manifold, [DGMS75, 5.21 (2)]). Assume that X is compact. Let $\mathcal{E} \in \text{Loc}^u(X)$. If the following conditions are satisfied:

1. The corresponding spectral sequence (4.5) degenerates at page E_1 ;
2. For every integer $k \geq 0$, the filtration induced by $F^\bullet(A_X^\bullet \otimes_{\mathbb{C}} \mathcal{E})$ on $H^k(X, \mathcal{E})$ gives a complex Hodge structure of weight k , in particular a Hodge decomposition

$$H^k(X, \mathcal{E}) = \oplus_{p+q=k} H^q(X, \Omega_X^p \otimes_{\mathbb{C}} \mathcal{E}); \quad (4.6)$$

3. For any integers $p, q \geq 0$, the conjugation map induces a \mathbb{C} -anti-linear isomorphism

$$H^q(X, \Omega_X^p \otimes_{\mathbb{C}} \mathcal{E}) \rightarrow H^p(X, \Omega_X^q \otimes_{\mathbb{C}} \mathcal{E}^\vee),$$

where $\mathcal{E}^\vee = \text{Hom}(\mathcal{E}, \mathbb{C}_X)$ is the dual local system.

Then X is called \mathcal{E} -regular (and simply *regular* when $\mathcal{E} = \mathbb{C}_X$).

For instance, classical Hodge theory asserts that compact Kähler manifolds are regular (see e.g., [Voi02, Sec. 6.1.3]). Because of Fact 4.3.1.2, regular manifolds are also called $\partial\bar{\partial}$ -manifolds.

Fact 4.3.1.2 ($\partial\bar{\partial}$ -lemma, [DGMS75, 5.14, 5.21], [Var86, Prop. 3.4], [Huy05, Cor. 3.2.10]). *Assume that X is compact. Then X is regular if and only if for every d -closed smooth (p, q) -form η on X , the following conditions are equivalent:*

1. η is d -exact;
2. η is ∂ -exact;
3. η is $\bar{\partial}$ -exact;
4. η is $\partial\bar{\partial}$ -exact.

If the above conditions hold and η is real, then there is a real smooth $(p-1, q-1)$ -form ρ on X with $\eta = i\partial\bar{\partial}\rho$.

Remark 4.3.1.3. For Fact 4.3.1.2, it is important that the decomposition (4.6) is induced by the filtration (4.4). In fact, [COUV16, Prop. 4.3] constructs a non-regular, compact complex manifold X of dimension 3, such that the spectral sequence (4.5) for $\mathcal{E} = \mathbb{C}_X$ degenerates at page E_1 , with numerical Hodge symmetry $h^{p,q}(X) = h^{q,p}(X)$ for any two integers $p, q \geq 0$. In this case, there is a non canonical decomposition of the form (4.6).

For the rest of Section 4.3.1, **we assume that X is a regular manifold**. For every integer $k \geq 0$ (resp. any two integers $p, q \geq 0$), the space of global ∂ -closed, $\bar{\partial}$ -closed smooth k (resp. (p, q)) forms on X is denoted by $Z^k(X)$ (resp. $Z^{p,q}(X)$). For any two integers $p, q \geq 0$, the Dolbeault cohomology group $H^q(X, \Omega_X^p)$ is denoted by $H^{p,q}(X)$.

Corollary 4.3.1.4. *For any integers $p, q \geq 0$ and $k := p+q$, there is a canonical commutative diagram*

$$\begin{array}{ccc} Z^{p,q}(X) & \hookrightarrow & Z^k(X) \\ \downarrow & & \downarrow \\ H^{p,q}(X) & \xrightarrow{\iota^{p,q}} & H^k(X), \end{array}$$

where the first row is the natural inclusion and each vertical map is surjective. Moreover,

$$H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} \text{im}(\iota^{p,q}), \quad (4.7)$$

where each $\text{im}(\iota^{p,q})$ can be identified with $H^{p,q}(X)$. The complex conjugation map $Z^{p,q}(X) \rightarrow Z^{q,p}(X)$ descends to a \mathbb{C} -antilinear isomorphism $H^{p,q}(X) \rightarrow H^{q,p}(X)$ (Hodge symmetry).

Proof. For each $\bar{\partial}$ -closed (p, q) -form η on X , $\partial\eta$ is a d -closed, ∂ -exact $(p + 1, q)$ -form. By Fact 4.3.1.2, there is a $(p, q - 1)$ -form ρ on X with $\partial\eta + \partial\bar{\partial}\rho = 0$, then the (p, q) -form $\eta + \bar{\partial}\rho$ is in $Z^{p,q}(X)$. Therefore, the map taking Dolbeault cohomology class $Z^{p,q}(X) \rightarrow H^{p,q}(X)$ is surjective.

Note that $\eta + \bar{\partial}\rho$ is d -closed. Its de Rham cohomology class is independent of the choice of ρ . Indeed, if ρ' is another $(p, q - 1)$ -form with $\eta + \bar{\partial}\rho'$ also d -closed, then $\bar{\partial}(\rho - \rho')$ is d -closed and $\bar{\partial}$ -exact. By Fact 4.3.1.2, it is d -exact.

Thus the map

$$\iota^{p,q} : H^q(X, \Omega_X^p) \rightarrow H_{\text{dR}}^{p+q}(X, \mathbb{C}), \quad [\eta] \mapsto [\eta + \bar{\partial}\rho]$$

is a well-defined \mathbb{C} -linear map. By a third application of Fact 4.3.1.2, the map $\iota^{p,q}$ is injective. Thus, $H^{p,q}$ is identified with $\text{im}(\iota^{p,q})$.

We claim that the sum $\sum_{p+q=k} \text{im}(\iota^{p,q})$ is direct. In fact, if $\alpha^{p,q} \in Z^{p,q}(X)$ for each pair (p, q) with $p + q = k$ and the de Rham class of $\sum_{p+q=k} \alpha^{p,q}$ is 0 in $H_{\text{dR}}^k(X, \mathbb{C})$, then there is a $(k - 1)$ -form β on X with $d\beta = \sum_{p+q=k} \alpha^{p,q}$. Thus,

$$\alpha^{p,q} = \partial(\beta^{p-1,q}) + \bar{\partial}(\beta^{p,q-1}).$$

The ∂ -exact form $\partial(\beta^{p-1,q})$ is thereby $\bar{\partial}$ -closed, so d -closed. By Fact 4.3.1.2 again, $\partial(\beta^{p-1,q})$ is $\bar{\partial}$ -exact, hence $[\alpha^{p,q}] = 0$ in $H^{p,q}(X)$ for every (p, q) . The claim is proved.

By assumption,

$$\dim_{\mathbb{C}} H^k(X, \mathbb{C}) = \sum_{p+q=k} \dim_{\mathbb{C}} H^{p,q}(X),$$

hence the decomposition (4.7). In particular, the map taking de Rham cohomology class $Z^k(X) \rightarrow H^k(X, \mathbb{C})$ is surjective. The complex conjugate of $Z^{p,q}(X)$ is exactly $Z^{q,p}(X)$, the Hodge symmetry follows. \square

Lemma 4.3.1.5 is used in the proof of Corollary 4.3.2.2.

Lemma 4.3.1.5. *For every integer $k \geq 0$, the map $H^k(X, \mathbb{C}) \rightarrow H^k(X, \mathcal{O}_X)$ induced by the inclusion $\mathbb{C} \rightarrow \mathcal{O}_X$ coincides with the projection $H^k(X, \mathbb{C}) \rightarrow H^{0,k}(X)$ given by the Hodge decomposition (4.7).*

Proof. Consider the following commutative diagram

$$\begin{array}{ccccccc} \mathbb{C}_X & \hookrightarrow & A_X^0 & \xrightarrow{d} & A_X^1 & \xrightarrow{d} & \dots \\ \downarrow & & \downarrow p^{0,0} & & \downarrow p^{0,1} & & \\ \mathcal{O}_X & \hookrightarrow & A_X^{0,0} & \xrightarrow{\bar{\partial}} & A_X^{0,1} & \xrightarrow{\bar{\partial}} & \dots \end{array}$$

The first row is an acyclic resolution of \mathbb{C}_X by (smooth) Poincaré lemma, and the second row is the Dolbeault resolution. The first vertical map is the inclusion and each $p^{0,j} : A_X^j \rightarrow A_X^{0,j}$ is taking the $(0, j)$ -part of a j -form. It is a morphism of complexes. Taking global sections, the induced map on k -th cohomology groups is the first map in the statement.

For a class $[\alpha] \in H^k(X, \mathbb{C})$, we may assume that the representative k -form α is ∂ -closed and $\bar{\partial}$ -closed by Corollary 4.3.1.4. Then its image under the first map $H^k(X, \mathbb{C}) \rightarrow H^k(X, O_X)$ is represented by the $(0, k)$ -part of α , which is still ∂ -closed and $\bar{\partial}$ -closed. This describes exactly the projection induced by the Hodge decomposition (4.7). \square

4.3.2 Jacobian

For a connected compact complex manifold X , let $b_1(X) := \dim_{\mathbb{C}} H^1(X, \mathbb{C})$ be its first Betti number. The exponential short exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow O_X \xrightarrow{f \mapsto \exp(2\pi i f)} O_X^* \rightarrow 1$$

induces a long exact sequence

$$H^0(X, O_X) \xrightarrow{f \mapsto \exp(2\pi i f)} H^0(X, O_X^*) \rightarrow H^1(X, \mathbb{Z}) \rightarrow H^1(X, O_X) \rightarrow H^1(X, O_X^*) \xrightarrow{\delta} H^2(X, \mathbb{Z}). \quad (4.8)$$

Set $\text{Pic}(X) := H^1(X, O_X^*)$ for the Picard group, $\text{NS}(X) := \text{im}(\delta)$ for the Néron-Severi group, $\text{Pic}^0(X) = \ker(\delta)$ and $\text{Pic}^\tau(X) := \delta^{-1}(H^2(X, \mathbb{Z})_{\text{tor}})$. As X is compact connected, one has $H^0(X, O_X) = \mathbb{C}$, $H^0(X, O_X^*) = \mathbb{C}^*$ and the first map in (4.8) is surjective. Accordingly, the third map $H^1(X, \mathbb{Z}) \rightarrow H^1(X, O_X)$ is injective and

$$\text{Pic}^0(X) = \frac{H^1(X, O_X)}{H^1(X, \mathbb{Z})}. \quad (4.9)$$

If X is a complex torus, then $H^2(X, \mathbb{Z})$ is torsion free and

$$\text{Pic}^0(X) = \text{Pic}^\tau(X). \quad (4.10)$$

For general X , let $\text{Loc}^1(X)$ (resp. $\text{Loc}^{u,1}(X)$) be the set of isomorphism classes of rank-1 (resp. and unitary) local systems on X . Then $\text{Loc}^1(X)$ is a group under tensor product and $\text{Loc}^{u,1}(X)$ is a subgroup. For each $\mathcal{L} \in \text{Loc}^1(X)$, $L := \mathcal{L} \otimes_{\mathbb{C}} O_X$ is a flat line bundle on X . By [Dem12, Ch. V, § 9], $L \in \text{Pic}^\tau(X)$, whence a group morphism

$$\text{Loc}^1(X) \rightarrow \text{Pic}^\tau(X), \quad \mathcal{L} \mapsto \mathcal{L} \otimes_{\mathbb{C}} O_X. \quad (4.11)$$

Remark 4.3.2.1. Theorem 4.2.3.1 implies that a line bundle on X is flat unitary if and only if its class in $\text{Pic}(X)$ lies in the image of the restriction of (4.11):

$$\text{Loc}^{u,1}(X) \rightarrow \text{Pic}^\tau(X). \quad (4.12)$$

The image of (4.12) may not be contained in $\text{Pic}^0(X)$. For instance, let X be an Enriques surface, then $\pi_1(X, x_0) = \mathbb{Z}/2$, $\#\text{Loc}^{u,1}(X) = \mathbb{Z}/2$. By Corollary 4.4.2.2 below, the map (4.12) is an isomorphism, while $\text{Pic}^0(X)$ is trivial.

Corollary 4.3.2.2. *Assume that X is regular. Then $\text{Pic}^\tau(X)$ has a natural structure of compact complex Lie group with identity component $\text{Pic}^0(X)$ that is a complex torus of dimension $b_1(X)/2$. Moreover, $\pi_0(\text{Pic}^\tau(X)) = \text{NS}(X)_{\text{tor}}$.*

Proof. The inclusion $\mathbb{R} \subset O_X$ induces an \mathbb{R} -linear map

$$\phi : H^1(X, \mathbb{R}) \rightarrow H^1(X, O_X). \quad (4.13)$$

Because of Lemma 4.3.1.5 and the Hodge symmetry in Corollary 4.3.1.4, taking complex conjugate inside $H^1(X, \mathbb{C})$ induces an \mathbb{R} -linear map

$$\bar{\phi} : H^1(X, \mathbb{R}) \rightarrow H^0(X, \Omega_X^1). \quad (4.14)$$

If $\xi \in \ker(\phi)$, then the image of ξ under the injection $H^1(X, \mathbb{R}) \rightarrow H^1(X, \mathbb{C})$ is $\phi(\xi) + \bar{\phi}(\xi) = 0$, so $\xi = 0$. This shows that ϕ is injective. But $\dim_{\mathbb{R}} H^1(X, \mathbb{R}) = \dim_{\mathbb{R}} H^1(X, O_X) = b_1(X)$, so ϕ is a linear isomorphism.

The map $H^1(X, \mathbb{Z}) \rightarrow H^1(X, O_X)$ in (4.8) factors through ϕ . Since $H^1(X, \mathbb{Z})$ is a full lattice of $H^1(X, \mathbb{R})$, it remains a full lattice in $H^1(X, O_X)$. Therefore, the quotient $\text{Pic}^0(X)$ is a complex torus of dimension $b_1(X)/2$. The \mathbb{Z} -module $\text{Pic}^0(X)$ is divisible, so the short exact sequence

$$0 \rightarrow \text{Pic}^0(X) \rightarrow \text{Pic}^\tau(X) \rightarrow \text{NS}(X)_{\text{tor}} \rightarrow 0$$

splits. Therefore, there is a natural structure of compact complex Lie group on $\text{Pic}^\tau(X)$ satisfying the stated properties. \square

The complex torus $\text{Pic}^0(X)$ in Corollary 4.3.2.2 is called the *Jacobian* of the regular manifold X .

Example 4.3.2.3. Here are two examples showing how Corollary 4.3.2.2 fails for non-regular compact complex manifolds.

1. Let X be a Hopf surface ([Huy05, Example 3.3.2]). The Betti number $b_1(X) = 1$, $H^1(X, \mathbb{Z}) = \mathbb{Z}$ and $H^1(X, O_X) = \mathbb{C}$, so the complex manifold $\text{Pic}^0(X) = \mathbb{C}/\mathbb{Z}$ is not compact. However, by [Kod64], the Frölicher spectral sequence of \mathbb{C}_X degenerates.
2. Let Y be a Calabi-Eckmann manifold ([BS17, Sec 1.2]). Then $H^1(Y, O_Y) = \mathbb{C}$ and Y is simply connected, so $H^1(Y, \mathbb{Z}) = 0$ and $b_1(Y) = 0$, but $\text{Pic}^0(Y) = \mathbb{C}$ is not compact and $b_1(Y)/2 < \dim \text{Pic}^0(Y)$.

4.4 Albanese torus

We turn to the conception of Albanese torus and Albanese map. They help to reduce some problems about general complex manifolds to those about complex tori. They are also tools to study the Jacobian. Again, Section 4.4 conveys the fact that Hodge theory guarantees the usual properties of the Albanese torus and Albanese map.

Fix a connected regular manifold X and a base point $x_0 \in X$.

4.4.1 Basics of Albanese torus

From [Uen06, Cor. 9.5, p.101], every element of $H^0(X, \Omega_X^1)$ is d -closed, so there is a well-defined natural map

$$\iota : H_1(X, \mathbb{Z}) \rightarrow H^0(X, \Omega_X^1)^\vee, \quad [\gamma] \mapsto (\beta \mapsto \int_\gamma \beta), \quad (4.15)$$

where γ runs through closed paths on X . Set

$$\text{Alb}(X) = H^0(X, \Omega_X^1)^\vee / \text{im}(\iota). \quad (4.16)$$

Lemma 4.4.1.1. *On $\text{Alb}(X)$, there is a natural structure of $h^{1,0}(X)$ -dimensional complex torus with $H_1(\text{Alb}(X), \mathbb{Z}) = \text{im}(\iota)$.*

Proof. Using the \mathbb{R} -linear isomorphism (4.14) and de Rham isomorphism

$$H_{\text{dR}}^1(X, \mathbb{R}) \rightarrow H^1(X, \mathbb{R}),$$

the map (4.15) is identified with the natural map $H_1(X, \mathbb{Z}) \rightarrow H_{\text{dR}}^1(X, \mathbb{R})^\vee$. The latter extends to an \mathbb{R} -linear isomorphism $H_1(X, \mathbb{R}) \rightarrow H_{\text{dR}}^1(X, \mathbb{R})^\vee$ by Poincaré duality. Therefore,

$$\ker(\iota) = H_1(X, \mathbb{Z})_{\text{tor}} \quad (4.17)$$

and $\text{im}(\iota)$ is a full lattice in $H^0(X, \Omega_X^1)^\vee$ isomorphic to $H_1(X, \mathbb{Z})_{\text{free}}$. Thus, the quotient $\text{Alb}(X)$ is a complex torus with the stated properties. \square

The complex torus $\text{Alb}(X)$ in Lemma 4.4.1.1 is called the *Albanese torus* of X . For each $x \in X$, choose two paths γ_x, γ'_x connecting x_0 to x . Then the composition γ of γ_x followed by the reverse of γ'_x is a closed path on X and

$$\int_{\gamma_x} \bullet - \int_{\gamma'_x} \bullet = \int_\gamma \bullet = \iota([\gamma])$$

belongs to $\text{im}(\iota)$. Therefore, $[\int_{\gamma_x} \bullet] = [\int_{\gamma'_x} \bullet]$ in $\text{Alb}(X)$. As $[\int_{\gamma_x} \bullet]$ is independent of the choice of γ_x , we write it as $\int_{x_0}^x \bullet$. For the fixed base point $x_0 \in X$, the associated Albanese morphism is

$$\alpha_{X, x_0} : X \rightarrow \text{Alb}(X), \quad x \mapsto \int_{x_0}^x \bullet.$$

The subscripts X and x_0 are omitted when they are clear from the context.

Proposition 4.4.1.2.

1. The Albanese map $\alpha_{X,x_0} : X \rightarrow \text{Alb}(X)$ is a morphism of complex manifolds and the formation of Albanese map is functorial for the pair (X, x_0) .
2. The induced morphism $\alpha_{x_0,*} : H_1(X, \mathbb{Z}) \rightarrow H_1(\text{Alb}(X), \mathbb{Z})$ is surjective with kernel $H_1(X, \mathbb{Z})_{\text{tor}}$.
3. The morphism α_{x_0} satisfies the following universal property: every morphism of pointed complex manifolds $(X, x_0) \rightarrow (A, 0)$ with A a complex torus factors uniquely through a morphism of complex tori $\text{Alb}(X) \rightarrow A$. In particular, the complex subtorus of $\text{Alb}(X)$ generated by $\alpha_{x_0}(X)$ is $\text{Alb}(X)$.
4. The pullback morphism $\alpha_{x_0}^* : H^1(\text{Alb}(X), \mathbb{Z}) \rightarrow H^1(X, \mathbb{Z})$ is an isomorphism of weight 1 \mathbb{Z} -Hodge structures independent of the choice of x_0 .
5. The pullback $\alpha_{x_0}^* : \text{Pic}^0(\text{Alb}(X)) \rightarrow \text{Pic}^0(X)$ is an isomorphism of complex tori independent of the choice of x_0 . In particular, the complex tori $\text{Alb}(X)$ and $\text{Pic}^0(X)$ are dual to each other (in the sense of [BL04, p.34]).

Proof.

1. When X is Kähler, it is proved in [Huy05, Prop. 3.3.8]. The general case is similar.
2. By Lemma 4.4.1.1, $H_1(\text{Alb}(X), \mathbb{Z}) = \text{im}(\iota)$. Let $\gamma : [0, 1] \rightarrow X$ be a closed path on X based at x_0 . It defines a path

$$\zeta : [0, 1] \rightarrow H^0(X, \Omega_X^1)^\vee, \quad \zeta(t) = \int_{\gamma(0)}^{\gamma(t)} \bullet,$$

where the integral is along a part of γ . Then

$$\zeta \pmod{\text{im}(\iota)} = \alpha_{x_0} \circ \gamma : [0, 1] \rightarrow \text{Alb}(X).$$

Therefore, $\alpha_{x_0,*}[\gamma] = \zeta(1) - \zeta(0) = \int_\gamma \bullet = \iota([\gamma])$. Hence a commutative triangle

$$\begin{array}{ccc} & \text{im}(\iota) = H_1(\text{Alb}(X), \mathbb{Z}) & \\ \alpha_{x_0,*} \nearrow & & \nwarrow \\ H_1(X, \mathbb{Z}) & \xrightarrow{\quad \iota \quad} & H^0(X, \Omega_X^1)^\vee \end{array}$$

Therefore, $\alpha_{x_0,*}$ is surjective and $\ker(\alpha_{x_0,*}) = \ker(\iota) = H_1(X, \mathbb{Z})_{\text{tor}}$, where the last equality uses (4.17).

3. The universal property follows from Point 1. Let T be the complex subtorus of $\text{Alb}(X)$ generated by $\alpha_{x_0}(X)$. Then the pointed morphism $\alpha_{x_0} : (X, x_0) \rightarrow (T, 0)$ factors through $\alpha_{x_0} : (X, x_0) \rightarrow (\text{Alb}(X), 0)$, so $T = \text{Alb}(X)$.
4. From [BL04, Thm. 1.4.1 b)], the map $\alpha_{x_0}^* : H^{1,0}(\text{Alb}(X)) \rightarrow H^{1,0}(X)$ is a \mathbb{C} -linear isomorphism. By [BL04, Sec 1.3, p.13], $H^1(\text{Alb}(X), \mathbb{Z})$ is naturally isomorphic to $\text{Hom}(\text{im}(\iota), \mathbb{Z})$. By Poincaré duality, the latter is identified with $H^1(X, \mathbb{Z})$, so

$$\alpha_{x_0}^* : H^1(\text{Alb}(X), \mathbb{Z}) \rightarrow H^1(X, \mathbb{Z})$$

is an isomorphism of weight 1 \mathbb{Z} -Hodge structures. Up to translation, different base points give rise to the same Albanese map. More precisely, for $x \in X$, $T_{\alpha_x(x_0)} \circ \alpha_{x_0} = \alpha_x$, where

$$T_a : \text{Alb}(X) \rightarrow \text{Alb}(X), \quad u \mapsto u + a$$

is the translation by a on $\text{Alb}(X)$. The independence stated in Point 4 follows.

5. As the isomorphism (4.9) is functorial in X , there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} H^1(\text{Alb}(X), \mathbb{Z}) & \longrightarrow & H^1(\text{Alb}(X), \mathcal{O}_{\text{Alb}(X)}) & \longrightarrow & \text{Pic}^0(\text{Alb}(X)) & \longrightarrow & 0 \\ \downarrow \alpha_{x_0}^* & & \downarrow \alpha_{x_0}^* & & \downarrow \alpha_{x_0}^* & & \\ H^1(X, \mathbb{Z}) & \longrightarrow & H^1(X, \mathcal{O}_X) & \longrightarrow & \text{Pic}^0(X) & \longrightarrow & 0. \end{array}$$

By Point 4, the left two vertical maps are isomorphisms independent of x_0 . Therefore, the right vertical map is an isomorphism independent of x_0 . As $\text{Alb}(X)$ is a complex torus, by [BL04, Proposition 2.4.1], $\text{Pic}^0(\text{Alb}(X))$ is the dual torus of $\text{Alb}(X)$. As $\alpha_{x_0}^* : \text{Pic}^0(\text{Alb}(X)) \rightarrow \text{Pic}^0(X)$ is an isomorphism, $\text{Pic}^0(X)$ is dual to $\text{Alb}(X)$.

□

Remark 4.4.1.3. By [Uen06, Cor. 9.5, p.101], for every connected regular manifold X the formation of $\text{Alb}(X)$ and α_{x_0} agrees with the construction in [Bla56, §2]. Then [Bla56, p.163] gives another proof of the universal property stated in Proposition 4.4.1.2 3.

EXAMPLE 4.3.2.3 1 (CONTINUED). If X were a Hopf surface, then $H_1(X, \mathbb{Z}) = \mathbb{Z}$ and $H^0(X, \Omega_X^1) = 0$. Equation (4.16) would define a point and Proposition 4.4.1.2 2 would fail.

4.4.2 Back to Jacobian

Albanese torus helps to understand the Jacobian. Corollary 4.4.2.1 is used to show the jumping loci are analytic subsets.

Corollary 4.4.2.1 (Universal line bundle). *There exists a unique (up to isomorphism) line bundle L on $X \times \text{Pic}^0(X)$ such that its pullback module to $\{x_0\} \times \text{Pic}^0(X)$ is trivial and for every point $y \in \text{Pic}^0(X)$, the isomorphism class of the pullback line bundle $L|_{X \times \{y\}}$ in $\text{Pic}(X)$ is y .*

Proof. Consider the map

$$f = \alpha_{x_0} \times \text{Id}_{\text{Pic}^0(X)} : X \times \text{Pic}^0(X) \rightarrow \text{Alb}(X) \times \text{Pic}^0(X).$$

By Proposition 4.4.1.2 5 and [GH78, Lemma, p.328], there is a holomorphic line bundle \mathcal{P} on $\text{Alb}(X) \times \text{Pic}^0(X)$ that is trivial on $\{0\} \times \text{Pic}^0(X)$ such that for every $y \in \text{Pic}^0(X)$, the line bundle $\mathcal{P}|_{\text{Alb}(X) \times \{y\}}$ is of class y in $\text{Pic}^0(\text{Alb}(X))$. Let $L = f^*\mathcal{P}$, then $L|_{\{x_0\} \times \text{Pic}^0(X)} = f^*(\mathcal{P}|_{\{0\} \times \text{Pic}^0(X)})$ is trivial. For every $y \in \text{Pic}^0(X)$, the line bundle

$$L|_{X \times \{y\}} = f^*(\mathcal{P}|_{\text{Alb}(X) \times \{y\}}) = \alpha_{x_0}^*(\mathcal{P}|_{\text{Alb}(X) \times \{y\}})$$

is of class y in $\text{Pic}^0(X)$. The existence is proved. The uniqueness follows from [BL04, Cor. A.9]. \square

Let

$$\text{Char}(X) = \text{Hom}(H_1(X, \mathbb{Z}), \mathbb{C}^*)$$

be the group of characters of the first homology of X . By [Hat05, Cor. A.8, A.9], the abelian group $H_1(X, \mathbb{Z})$ is finitely generated. From [Mil17a, Ch. 12 b.], $\text{Char}(X)$ has a natural structure of diagonalizable algebraic group over \mathbb{C} , with identity component $\text{Char}^\circ(X)$ isomorphic to $\mathbb{G}_m^{b_1(X)}$. Moreover, $\text{Char}^u(X) := \text{Hom}(H_1(X, \mathbb{Z}), S^1)$ is a real Lie subgroup of $\text{Char}(X)$ of dimension $b_1(X)$. There is a canonical group isomorphism by taking character sheaves

$$\text{Char}^u(X) \rightarrow \text{Loc}^{u,1}(X), \quad \chi \mapsto \mathcal{L}_\chi. \quad (4.18)$$

Set $T(X) := \text{Hom}(H_1(X, \mathbb{Z})_{\text{free}}, S^1)$. Then $T(X)$ is the identity component of $\text{Char}^u(X)$. From Corollary 4.3.2.2, composing the isomorphism (4.18) and the map (4.12) gives a morphism of real Lie groups

$$T(X) \rightarrow \text{Pic}^0(X). \quad (4.19)$$

In Corollary 4.4.2.2 1, the isomorphism allows one to identify certain characters with topologically trivial line bundles. This identification is used in the proof of Theorem 4.7.1.3. When X is in Fujiki Class \mathcal{C} (resp. Kähler), Corollary 4.4.2.2 2 is also in [Ara90, Lem. 2] (resp. the proof of [Wan16, Cor. 1.4]).

Corollary 4.4.2.2.

1. The morphism (4.19) is an isomorphism of real Lie groups.
2. The map (4.12) is a group isomorphism and $\text{NS}(X)_{\text{tor}} = H^2(X, \mathbb{Z})_{\text{tor}}$. In particular, every element of $\text{Pic}^\tau(X)$ is a flat unitary line bundle.

Proof.

1. Lemma 4.4.1.1 gives an identification $H_1(X, \mathbb{Z})_{\text{free}} = \text{im}(\iota)$. By [BL04, Prop. 2.2.2], the natural group morphism

$$\text{Hom}(\text{im}(\iota), S^1) \rightarrow \text{Pic}^0(\text{Alb}(X)) \quad (4.20)$$

defined via factors of automorphy ([BL04, p.30]) is an isomorphism. The map (4.19) is the composition of (4.20) with the isomorphism $\alpha_{x_0}^* : \text{Pic}^0(\text{Alb}(X)) \rightarrow \text{Pic}^0(X)$ in Proposition 4.4.1.2 5.

To sum it up:

$$\begin{array}{ccccc} \text{Char}^u(\text{Alb}(X)) = \text{Hom}(\text{im}(\iota), S^1) & \xrightarrow{\sim} & T(X) & \hookrightarrow & \text{Char}^u(X) = \text{Loc}^{u,1}(X) \\ \downarrow (4.20) & & \downarrow (4.19) & & \downarrow (4.12) \\ \text{Pic}^0(\text{Alb}(X)) & \xrightarrow[\sim]{\alpha_{x_0}^*} & \text{Pic}^0(X) & \hookrightarrow & \text{Pic}^\tau(X). \end{array}$$

2. The commutative diagram of abelian sheaves on X

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{R} & \longrightarrow & S^1 \longrightarrow 0 \\ & & \downarrow \text{Id} & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathcal{O}_X & \xrightarrow{f \mapsto \exp(2\pi i f)} & \mathcal{O}_X^* \longrightarrow 0 \end{array}$$

has exact rows. Moreover, the \mathbb{Z} -module \mathbb{R} is injective. Therefore, there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} & & 0 & \longrightarrow & \text{Hom}(H_1(X, \mathbb{Z})_{\text{tor}}, S^1) & \xrightarrow{\sim} & \text{Ext}_{\mathbb{Z}}^1(H_1(X, \mathbb{Z})_{\text{tor}}, \mathbb{Z}) \longrightarrow 0 \\ & & & & \uparrow r & & \uparrow \sim \\ 0 & \rightarrow & \text{Hom}(H_1(X, \mathbb{Z}), \mathbb{Z}) & \rightarrow & \text{Hom}(H_1(X, \mathbb{Z}), \mathbb{R}) & \longrightarrow & \text{Char}^u(X) \longrightarrow \text{Ext}_{\mathbb{Z}}^1(H_1(X, \mathbb{Z}), \mathbb{Z}) \longrightarrow 0 \\ & & \uparrow \sim & & \uparrow \sim & & \uparrow \sim \\ 0 & \longrightarrow & H^1(X, \mathbb{Z}) & \longrightarrow & H^1(X, \mathbb{R}) & \longrightarrow & H^1(X, S^1) \longrightarrow H^2(X, \mathbb{Z}) \\ & & \downarrow \text{Id} & & \downarrow (4.13) & & \downarrow \\ 0 & \longrightarrow & H^1(X, \mathbb{Z}) & \longrightarrow & H^1(X, \mathcal{O}_X) & \longrightarrow & H^1(X, \mathcal{O}_X^*) \xrightarrow{\delta} H^2(X, \mathbb{Z}) \end{array}$$

where r is the restriction, the vertical morphisms in the middle are from [Hat05, Thm. 3.2] and $\text{im}(\xi) = H^2(X, \mathbb{Z})_{\text{tor}}$ by [Hat05, p.196]. Hence an isomorphism $\psi : \text{Hom}(H_1(X, \mathbb{Z})_{\text{tor}}, S^1) \rightarrow H^2(X, \mathbb{Z})_{\text{tor}}$ fitting into a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & T(X) & \longrightarrow & \text{Loc}^{u,1}(X) & \xrightarrow{r} & \text{Hom}(H_1(X, \mathbb{Z})_{\text{tor}}, S^1) \longrightarrow 0 \\
& & \downarrow (4.19) & & \downarrow (4.12) & & \downarrow \psi \\
0 & \longrightarrow & \text{Pic}^0(X) & \longrightarrow & \text{Pic}^\tau(X) & \xrightarrow{\delta} & H^2(X, \mathbb{Z})_{\text{tor}} \longrightarrow 0,
\end{array}$$

where the first row is exact. Thus, δ is surjective and the second row is also exact. By the five lemma, the middle vertical map (4.12) is an isomorphism.

□

4.5 Defect of semismallness

We review the defect of semismallness of a morphism, an invariant introduced by de Cataldo and Migliorini that plays a crucial role in the decomposition theorem and Lefschetz's theorem. It appears in Fact 4.1.1.6 and Theorem 4.7.1.3. Its main property that we need is Proposition 4.5.3.2. Complex analytic spaces are in the sense of [GR84, p.7].

4.5.1 Stratifications and constructible sheaves

We refer to [BF84, Sec. 2.1] for the definitions of **constructible stratifications** and **Whitney stratifications** of a complex analytic space.

Theorem 4.5.1.1 is about the semicontinuity of fiber dimension. Although it is well-known, a short proof is included due to the lack of reference. Its analogue in algebraic geometry is a celebrated theorem of Chevalley [Gro66, Cor. 13.1.5].

Theorem 4.5.1.1 (Analytic Chevalley theorem). *Let $f : X \rightarrow Y$ be a proper morphism of reduced complex analytic spaces. For every integer $n \geq 0$, let $Y_n = \{y \in Y : \dim f^{-1}(y) = n\}$ and $Y_{\geq n} = \cup_{m \geq n} Y_m$. Then $Y_{\geq n}$ is an analytic subset of Y . In particular, $\{Y_n\}_{n \in \mathbb{N}}$ is a constructible stratification of Y .*

Proof. Let $F_n := \{x \in X : \dim_x f^{-1}(f(x)) \geq n\}$. By [Fis76, Thm. 3.6, p.137], F_n is an analytic subset of X . By the definition of global dimension [GR84, p.94], one has $Y_{\geq n} = f(F_n)$. By Remmert theorem (see, e.g., [Whi72, Thm. 4A, p.150]), the subset $Y_{\geq n}$ is analytic in Y . □

Definition 4.5.1.2. ([BF84, p.125]) Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces. If two Whitney stratifications $\mathfrak{X} : X = \sqcup_{\alpha} X_{\alpha}$ and $\mathfrak{Y} : Y = \sqcup_{\lambda} Y_{\lambda}$ satisfy that:

1. For each α , there is λ with $f(X_{\alpha}) \subset Y_{\lambda}$;
2. For each pair (α, λ) with $f(X_{\alpha}) \subset Y_{\lambda}$, the restricted morphism $f : X_{\alpha} \rightarrow Y_{\lambda}$ is smooth.

Then such a pair $(\mathfrak{X}, \mathfrak{Y})$ is called a Whitney stratification of f .

Fact 4.5.1.3 ([Hir77, Thm. 1], [BF84, Lem. 2.4], [GM88, Thm, p.43]). *Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces. Suppose that \mathfrak{X} , (resp. \mathfrak{Y}) is a constructible stratification of X (resp. Y), then there exists a Whitney stratification $(\mathfrak{X}', \mathfrak{Y}')$ of f such that \mathfrak{X}' (resp. \mathfrak{Y}') refines \mathfrak{X} (resp. \mathfrak{Y}).*

Corollary 4.5.1.4 is useful but implicit in the literature.

Corollary 4.5.1.4. *Let X be a complex analytic space. For finitely many constructible stratifications of X , there exists a Whitney stratification of X refining all of them.*

Proof. It suffices to consider the case of two constructible stratifications \mathfrak{X}_1 and \mathfrak{X}_2 of X . By Fact 4.5.1.3, there is a Whitney stratification $(\mathfrak{X}, \mathfrak{X}')$ of Id_X such that \mathfrak{X} (resp. \mathfrak{X}') refines \mathfrak{X}_1 (resp. \mathfrak{X}_2). Moreover, \mathfrak{X} refines \mathfrak{X}' by Definition 4.5.1.2. Hence a Whitney stratification \mathfrak{X} refining both \mathfrak{X}_1 and \mathfrak{X}_2 . \square

For a complex analytic space X , using analytic constructible stratifications, one can define constructible sheaves. Let $D_c^b(X)$ be the triangulated category of complexes of sheaves of \mathbb{C} -vector spaces whose cohomology is bounded and constructible (see, e.g., [Dim04, p.82]).

Fact 4.5.1.5 ([KS90, Prop. 8.5.7 (b)], [Dim04, Thm. 4.1.5 (b)]). *Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces and $\mathcal{K} \in D_c^b(X)$. If f is proper on $\text{Supp}(\mathcal{K})$, then $Rf_*\mathcal{K} \in D_c^b(Y)$.*

Corollary 4.5.1.6. *Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces and $\mathcal{K} \in D_c^b(X)$. Then there exists a Whitney stratification $(\mathfrak{X}, \mathfrak{Y})$ of f such that for every integer i and every stratum S of \mathfrak{Y} , the restriction $\mathcal{H}^i(Rf_*\mathcal{K})|_S$ is a local system on S .*

Proof. By Fact 4.5.1.5, $Rf_*\mathcal{K} \in D_c^b(Y)$. In particular, there are only finitely many $j \in \mathbb{Z}$ with $\mathcal{H}^j(Rf_*\mathcal{K}) \neq 0$. For each such j , there is an admissible partition (in the sense of [Dim04, p.81]) \mathcal{P}_j on Y such that the restriction of $\mathcal{H}^j(Rf_*\mathcal{K})$ to each stratum of \mathcal{P}_j is a local system. By Corollary 4.5.1.4, there exists a Whitney stratification \mathfrak{Y}^0 of Y refining the finitely many \mathcal{P}_j . By Fact 4.5.1.3, there is a Whitney stratification $(\mathfrak{X}, \mathfrak{Y})$ of f satisfying the properties. \square

4.5.2 Equivalent definitions

The defect of semismallness measures how far a morphism of complex manifolds is from being semismall (see, e.g., [KW01, Def. 7.3, p.156]). However, in the literature there exist multiple seemingly different definitions. We review some of them and show that they are equivalent.

Definition 4.5.2.1. Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds with $\dim X = n$.

- ([EV89, Def. 1.1]) Define

$$r_1(f) = \max_Z (\dim Z - \dim f(Z) - \operatorname{codim}_X(Z)), \quad (4.21)$$

where Z runs through all irreducible analytic subsets of X .

- ([Max19, Def. 9.3.7]) For a Whitney stratification $(X = \sqcup S_\alpha, Y = \sqcup T_\lambda)$ of f , we choose a point $y_\lambda \in T_\lambda$ in each stratum, and define

$$r_2(f) = \max_\lambda \{2 \dim f^{-1}(y_\lambda) + \dim T_\lambda - n\}. \quad (4.22)$$

(By convention, the empty space has dimension $-\infty$.)

- ([dCM05, Def. 4.7.2]) For each integer $i \geq 0$, let $Y_i = \{y \in Y : \dim f^{-1}(y) = i\}$. Define

$$r_3(f) = \max_{i \geq 0} (2i + \dim Y_i - n).$$

- ([PS13, Def. 2.8]) For each integer $i \geq 0$, let $Y_{\geq i} = \{y \in Y : \dim f^{-1}(y) \geq i\}$ for each $i \geq 0$. Define

$$r_4(f) = \max_{i \geq 0} (2i + \dim Y_{\geq i} - n).$$

- ([dCM09, Sec. 3.3.2, part 2]) Define $r_5(f) = \dim X \times_Y X - n$.

- ([Wil16, Sec 3.2]) Define

$$r_6(f) = \max\{i \in \mathbb{Z} : {}^p\mathcal{H}^i(Rf_*\mathbb{C}_X[n]) \neq 0\}.$$

Proposition 4.5.2.2. *The first five numbers in Definition 4.5.2.1 are all equal.*

This common integer is called the *defect of semismallness* of f and denoted by $r(f)$. We shall show $r(f) = r_6(f)$ in Proposition 4.5.3.2 2.

Proof.

- $r_3(f) = r_4(f)$: As each Y_i is a subset of $Y_{\geq i}$, one has $r_3(f) \leq r_4(f)$. There are only finitely many integers $i \geq 0$ with $Y_{\geq i}$ nonempty, so the maximum defining $r_4(f)$ is attained at some $i_0(\geq 0)$. Then

$$2(i_0 + 1) + \dim Y_{\geq i_0+1} \leq 2i_0 + \dim Y_{\geq i_0}.$$

Since $Y_{\geq i_0} = Y_{\geq i_0+1} \cup Y_{i_0}$, one has $\dim Y_{\geq i_0} = \dim Y_{i_0}$. Then

$$r_4(f) = 2i_0 + \dim Y_{\geq i_0} - n \leq r_3(f).$$

Therefore, $r_3(f) = r_4(f)$.

- $r_2(f) = r_5(f)$: By Thom's first isotopy lemma (see, e.g., [Mat12, Prop. 11.1]), for every λ , the restriction $f|_{f^{-1}(T_\lambda)} : f^{-1}(T_\lambda) \rightarrow T_\lambda$ is a topologically locally trivial fibration. Therefore, $\dim f^{-1}(y_\lambda)$ is independent of $y_\lambda \in T_\lambda$ and

$$\dim f^{-1}(T_\lambda) \times_{T_\lambda} f^{-1}(T_\lambda) = \dim T_\lambda + 2 \dim f^{-1}(y_\lambda). \quad (4.23)$$

As $\{f^{-1}(T_\lambda) \times_{T_\lambda} f^{-1}(T_\lambda)\}_\lambda$ is a locally finite partition of $X \times_Y X$ into locally closed subsets (in the analytic Zariski topology), one has

$$\dim X \times_Y X = \max_\lambda [\dim f^{-1}(T_\lambda) \times_{T_\lambda} f^{-1}(T_\lambda)]. \quad (4.24)$$

Plugging (4.23) into (4.24) we get $r_5(f) = r_2(f)$. In particular, $r_2(f)$ is independent of the choice of the stratifications.

- $r_1(f) \leq r_2(f)$: For every irreducible analytic subset $Z \subset X$, $f(Z)$ is an irreducible analytic subset of Y . Then $\{Y \setminus f(Z), f(Z)\}$ is a constructible stratification of Y . Fact 4.5.1.3 yields a Whitney stratification $(X = \sqcup S_\alpha, Y = \sqcup T_\lambda)$ of f with $Y = \sqcup T_\lambda$ refining $\{Y \setminus f(Z), f(Z)\}$. There exists λ_0 such that T_{λ_0} is an open subset of $f(Z)$, hence $\dim T_{\lambda_0} \leq \dim f(Z)$. Then $f^{-1}(T_{\lambda_0}) \cap Z$ is a nonempty open subset of Z . Therefore,

$$\dim Z = \dim(f^{-1}(T_{\lambda_0}) \cap Z) \leq \dim f^{-1}(T_{\lambda_0}).$$

Then

$$2 \dim Z - \dim f(Z) \leq 2 \dim f^{-1}(T_{\lambda_0}) - \dim T_{\lambda_0} = 2 \dim f^{-1}(y_{\lambda_0}) + \dim T_{\lambda_0}.$$

This shows $r_1(f) \leq r_2(f)$. In particular, the maximum in (4.21) is indeed attained.

- $r_2(f) \leq r_1(f)$: Fix a Whitney stratification $Y = \sqcup_\lambda T_\lambda$ defining $r_2(f)$. For every λ with $f^{-1}(y_\lambda)$ nonempty, $\overline{T_\lambda}$ is an analytic subset of Y of dimension $\dim T_\lambda$. Then $f^{-1}(\overline{T_\lambda})$ is a nonempty analytic subset of

X . Let Z_0 be an irreducible component of $f^{-1}(\overline{T_\lambda})$ with $\dim Z_0 = \dim f^{-1}(\overline{T_\lambda})$. Then $f(Z_0) \subset \overline{T_\lambda}$ and $\dim f(Z_0) \leq \dim T_\lambda$. Therefore,

$$2 \dim f^{-1}(y_\lambda) + \dim T_\lambda = 2 \dim f^{-1}(T_\lambda) - \dim T_\lambda \leq 2 \dim Z_0 - \dim f(Z_0).$$

This shows $r_2(f) \leq r_1(f)$.

- $r_2(f) \leq r_3(f)$: By Theorem 4.5.1.1, $\{Y_i\}$ is a constructible stratification of Y . By Fact 4.5.1.3, there is a Whitney stratification $(X = \sqcup S_\alpha, Y = \sqcup T_\lambda)$ of f such that the stratification $Y = \sqcup T_\lambda$ refines $Y = \sqcup_i Y_i$. For every λ , there is i_0 with $T_\lambda \subset Y_{i_0}$. In particular, for every $y_\lambda \in T_\lambda$, one has $\dim f^{-1}(y_\lambda) = i_0$, so

$$2 \dim f^{-1}(y_\lambda) + \dim T_\lambda \leq 2i_0 + \dim Y_{i_0}.$$

This shows $r_2(f) \leq r_3(f)$.

- $r_3(f) \leq r_2(f)$: For every integer $i \geq 0$ with Y_i nonempty, $Y_i = \sqcup_\lambda (Y_i \cap T_\lambda)$ is a constructible stratification, so there is an index λ_0 with $\dim(Y_i \cap T_{\lambda_0}) = \dim Y_i$. Then $\dim Y_i \leq \dim T_{\lambda_0}$. One may take $y_{\lambda_0} \in Y_i \cap T_{\lambda_0}$. Then

$$2i + \dim Y_i \leq 2 \dim f^{-1}(y_{\lambda_0}) + \dim T_{\lambda_0},$$

which shows $r_3(f) \leq r_2(f)$.

□

From the diagonal inclusion $X \rightarrow X \times_Y X$, one gets $\dim X \leq \dim X \times_Y X$, so $r(f) = r_5(f) \geq 0$. If $r(f) = 0$, then f is said to be semismall.

Example 4.5.2.3.

1. If $f : X \rightarrow Y$ is a proper morphism of complex manifolds that is flat of relative dimension r , then $r(f) = r$.
2. Let X be projective manifold such that $-K_X$ is nef and $\alpha : X \rightarrow \text{Alb}(X)$ be the Albanese map associated with some base point. Then $r(\alpha) = \dim X - \dim \alpha(X)$ by [LTZZ10, Theorem].

4.5.3 Direct image of local systems

Defect of semismallness is an important invariant appearing in the decomposition of direct image of perverse sheaves. Proposition 4.5.3.2 is an elementary instance. We begin with a well-known estimation of cohomological dimension of a complex analytic space, used in the proof of Proposition 4.5.3.2. An analogue for topological manifolds is [KS90, Prop. 3.2.2 (iv)].

Lemma 4.5.3.1. *Let X be a paracompact (in the sense of [Mun00, Def., p.253]) complex analytic space of complex dimension n . Then $H^q(X, F) = 0$ for every abelian sheaf F on X and every integer $q > 2n$.*

Proof. By [GR84, Prop., p.94], there is an open covering $\{U_\alpha\}_\alpha$ of X such that for each α , there is a finite morphism $f_\alpha : U_\alpha \rightarrow B_\alpha$ of complex analytic spaces to an open ball $B_\alpha \subset \mathbb{C}^n$. As X is Hausdorff paracompact, by [Mun00, Lemma 41.6], there exists a locally finite open covering $\{V_\alpha\}$ on X such that $\overline{V_\alpha} \subset U_\alpha$ for each α .

From [Mun00, p.314], for every α , the topological dimension ([Mun00, Def., p.305]) $\text{covdim}(B_\alpha) = 2n$. By [KK83, Prop. 51 A.2], the topological space X is metrizable. From [Mun00, Thm. 32.2], each U_α is normal. Therefore, by [Eng95, Thm. 3.3.10, p.200], $\text{covdim}(U_\alpha) \leq 2n$. By [Eng95, Theorem 3.1.3, p.169], $\text{covdim}(\overline{V_\alpha}) \leq 2n$. Similarly, X is normal, so $\text{covdim}(X) \leq 2n$ by [Eng95, Thm. 3.1.10, p.172]. By Alexandroff theorem (see, e.g., [Bre12, p.122]), the cohomological dimension ([Eng95, p.75]) $\dim_{\mathbb{Z}} X \leq 2n$. \square

The category $D_c^b(X)$ has a natural perverse t-structure (p being the middle perversity)

$$({}^p D^{\leq 0}(X), {}^p D^{\geq 0}(X)),$$

whose heart $\text{Perv}(X)$ is a \mathbb{C} -linear abelian category ([BBDG82], see also [HT07, Thm. 8.1.27]). An object of $\text{Perv}(X)$ is called a perverse sheaf on X . For every integer i , the functor taking the i -th perverse cohomology sheaf is denoted by ${}^p \mathcal{H}^i : D_c^b(X) \rightarrow \text{Perv}(X)$. For any two integers $a \leq b$, set

$$\begin{aligned} {}^p D^{[a,b]}(X) &:= \{\mathcal{K} \in D_c^b(X) : {}^p \mathcal{H}^i(\mathcal{K}) = 0, \forall i \notin [a, b]\}; \\ D^{[a,b]}(X) &:= \{\mathcal{K} \in D_c^b(X) : \mathcal{H}^i(\mathcal{K}) = 0, \forall i \notin [a, b]\}. \end{aligned}$$

Verdier duality $\mathcal{D}_X : D_c^b(X) \rightarrow D_c^b(X)$ is a contravariant auto-equivalence that interchanges ${}^p D^{\leq 0}(X)$ and ${}^p D^{\geq 0}(X)$ (see, e.g., [HT07, p.192]).

Proposition 4.5.3.2 is an analytic analogue of [dCM03, Prop. 10.0.7]. It allows local coefficients and in our case permits to descend some problems about local systems on X to problems about complexes of sheaves on Y . The proof is different from that in [dCM03], in particular it does not use the decomposition theorem [dCM03, Thm. 10.0.6].

Proposition 4.5.3.2. *Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds, where X is of pure dimension n . Let \mathcal{L} a local system on X . Then:*

1. $Rf_*(\mathcal{L}[n]) \in {}^p D^{[-r(f), r(f)]}(Y)$. In particular, $Rf_* \mathcal{L}[n] \in \text{Perv}(Y)$ when f is moreover semismall.

2. When $\mathcal{L} = \mathbb{C}_X$, for $j = \pm r(f)$, one has ${}^p\mathcal{H}^j(Rf_*\mathbb{C}_X[n]) \neq 0$. Hence $r(f) = r_6(f)$.

Proof. From Corollary 4.5.1.6, there exists a Whitney stratifications $(X = \sqcup_{\alpha} X_{\alpha}, Y = \sqcup_{\lambda} Y_{\lambda})$ of f such that for every λ , every integer j , the restriction $\mathcal{H}^j(Rf_*\mathcal{L}[n])|_{Y_{\lambda}}$ is a local system. For each λ , choose a point $y_{\lambda} \in Y_{\lambda}$.

1. First, we show that $Rf_*\mathcal{L}[n] \in {}^pD^{\leq r(f)}(Y)$. Fix an integer i . If $\dim Y_{\lambda} > r(f) - i$, then by (4.22), one has $i + n > 2 \dim f^{-1}(y_{\lambda})$. Since the fiber $f^{-1}(y_{\lambda})$ is a compact complex analytic space, by Lemma 4.5.3.1,

$$H^{i+n}(f^{-1}(y_{\lambda}), \mathcal{L}|_{f^{-1}(y_{\lambda})}) = 0.$$

By proper base change theorem (see, e.g., [Mil13, Thm. 17.2]),

$$\mathcal{H}^i(Rf_*\mathcal{L}[n])_{y_{\lambda}} = H^{i+n}(f^{-1}(y_{\lambda}), \mathcal{L}|_{f^{-1}(y_{\lambda})}).$$

So $\mathcal{H}^i(Rf_*\mathcal{L}[n]) = 0$ on every stratum Y_{λ} with $\dim Y_{\lambda} > r(f) - i$. Therefore, $\dim \text{Supp } \mathcal{H}^i(Rf_*\mathcal{L}[n]) \leq r(f) - i$ and hence $Rf_*\mathcal{L}[n] \in {}^pD^{\leq r(f)}(Y)$.

It remains to show $Rf_*\mathcal{L}[n] \in {}^pD^{\geq -r(f)}(Y)$. By what we have proved, $Rf_*\mathcal{L}^{\vee}[n] \in {}^pD^{\leq r(f)}(Y)$. Since $\mathcal{D}_X(\mathcal{L}[n]) = \mathcal{L}^{\vee}[n]$, one has

$$Rf_*\mathcal{L}^{\vee}[n] = Rf_*\mathcal{D}_X(\mathcal{L}[n]) = \mathcal{D}_Y(Rf_*\mathcal{L}[n]).$$

The last equality uses Verdier's duality (see, e.g., [Max19, Prop. 5.3.9]). This shows $Rf_*\mathcal{L}[n] \in {}^pD^{\geq -r(f)}(Y)$.

2. By (4.22), there exists λ_0 with $r(f) = 2 \dim f^{-1}(y_{\lambda_0}) + \dim Y_{\lambda_0} - n$. In particular, $f^{-1}(y_{\lambda_0})$ is nonempty. Let $i_0 = r(f) - \dim Y_{\lambda_0}$, then $i_0 + n = 2 \dim f^{-1}(y_{\lambda_0})$. By proper base change theorem again,

$$\mathcal{H}^{i_0}(Rf_*\mathbb{C}[n])_{y_{\lambda_0}} = H^{i_0+n}(f^{-1}(y_{\lambda_0}), \mathbb{C}) \neq 0.$$

Therefore, $Y_{\lambda_0} \subset \text{Supp } \mathcal{H}^{i_0}(Rf_*\mathbb{C}[n])$ and hence

$$\dim \text{Supp } \mathcal{H}^{i_0}(Rf_*\mathbb{C}[n]) \geq \dim Y_{\lambda_0} = r(f) - i_0.$$

Then $Rf_*\mathbb{C}[n] \notin {}^pD^{\leq r(f)-1}(Y)$. Together with Point 1, this shows

$${}^p\mathcal{H}^{r(f)}(Rf_*\mathbb{C}_X[n]) \neq 0.$$

The other part follows from Verdier's duality.

□

4.6 Generic vanishing for constructible sheaves

In Section 4.6, we review the generic vanishing theorem for (complexes of) constructible sheaves on a complex torus. The case of abelian varieties is treated in [KW15b] and the general case in [BSS18]. We shall reduce the generic vanishing problem on a manifold in Fujiki class \mathcal{C} to results on its Albanese torus.

4.6.1 Thin subsets

To state Krämer-Weissauer's theorem, we recall the terminology "thin subset" introduced in [KW15b, p.532 and p.536].

Fix a complex torus A . Then $\text{Char}(A)$ has a natural structure of algebraic torus over \mathbb{C} of dimension $2 \dim A$ and $T(A) = \text{Char}^u(A)$ is a real Lie subgroup of $\text{Char}(A)$. For each complex subtorus $B \subset A$, let $K(B)$ be the kernel of the morphism of algebraic tori $\text{Char}(A) \rightarrow \text{Char}(B)$ induced by functoriality. The induced morphism $\pi_1(B, 0) \rightarrow \pi_1(A, 0)$ is injective with torsion-free cokernel of rank $2 \dim A - 2 \dim B$, so $K(B)$ is an algebraic subtorus of $\text{Char}(A)$.

Definition 4.6.1.1. A thin subset of $\text{Char}(A)$ is a finite union of translates $\chi_i \cdot K(A_i)$ for certain characters $\chi_i \in \text{Char}(A)$ and certain *nonzero* complex subtori $A_i \subset A$. If every χ_i can be chosen to be a torsion point of $\text{Char}(A)$, then such a thin subset is called *arithmetic*.

A thin subset of $\text{Char}(A)$ is strict and Zariski closed. If the complex torus A is nonzero and *simple*, then a subset of $\text{Char}(A)$ is thin if and only if it is finite.

For each complex subtorus $B \subset A$, we have a functorial commutative diagram

$$\begin{array}{ccccccc}
 \text{Char}^u(A) & \xrightarrow{(4.18)} & \text{Loc}^{u,1}(A) & \xrightarrow{(4.12)} & \text{Pic}^\tau(A) & \xleftarrow{(4.10)} & \text{Pic}^0(A) \\
 \downarrow \phi & & \downarrow & & \downarrow & & \downarrow \psi \\
 \text{Char}^u(B) & \longrightarrow & \text{Loc}^{u,1}(B) & \longrightarrow & \text{Pic}^\tau(A) & \longleftarrow & \text{Pic}^0(B)
 \end{array} \tag{4.25}$$

where all the horizontal maps are isomorphisms by Corollary 4.4.2.2 2.

A subset of $\text{Pic}^0(A)$ is called (arithmetic and) thin, if it is the intersection of $\text{Char}^u(A)$ with a (arithmetic and) thin subset of $\text{Char}(A)$ when $\text{Pic}^0(A)$ is identified with $\text{Char}^u(A)$ via the diagram (4.25).

Lemma 4.6.1.2. *Every thin subset of $\text{Pic}^0(A)$ is a finite union of translates of strict complex subtori.*

Proof. Let B be a subtorus of A . As the induced morphism $\pi_1(B, 0) \rightarrow \pi_1(A, 0)$ is injective with torsion-free cokernel of rank $2(\dim A - \dim B)$,

the restriction morphism $\phi : \text{Char}^u(A) \rightarrow \text{Char}^u(B)$ in (4.25) is surjective, and its kernel $K(B) \cap \text{Char}^u(A)$ is the group of unitary characters of $\pi_1(A, 0)/\pi_1(B, 0)$. Therefore, the kernel of the morphism $\psi : \text{Pic}^0(A) \rightarrow \text{Pic}^0(B)$ in (4.25) is a complex subtorus of dimension $\dim A - \dim B$. \square

For a connected regular manifold X , let $\alpha : X \rightarrow \text{Alb}(X)$ be its Albanese morphism corresponding to some base point. Then α induces a morphism $\alpha^* : \text{Char}(\text{Alb}(X)) \rightarrow \text{Char}(X)$ of algebraic groups. By Proposition 4.4.1.2 2, this map identifies $\text{Char}(\text{Alb}(X))$ with the identity component $\text{Char}^\circ(X)$ of $\text{Char}(X)$. Thus we can define thin subsets of $\text{Char}^\circ(X)$. By Proposition 4.4.1.2 5, $\text{Pic}^0(X)$ is naturally identified with $\text{Pic}^0(\text{Alb}(X))$, thus we can define (arithmetic and) thin subsets of $\text{Pic}^0(X)$.

4.6.2 Generic vanishing result on regular manifolds

Roughly speaking, Krämer-Weissauer's theorem controls the failure of vanishing for perverse sheaves on complex tori, measured by the following loci.

Let X be a compact complex manifold of dimension d . For any integers $k \geq 0$, i and for every $\mathcal{K} \in D_c^b(X)$, consider the cohomology support locus

$$\Sigma^i(X, \mathcal{K}) := \{\chi \in \text{Char}(X) : H^i(X, \mathcal{L}_\chi \otimes \mathcal{K}) \neq 0\}.$$

Let $\Sigma^{\neq 0}(X, \mathcal{K}) := \cup_{i \neq 0, i \in \mathbb{Z}} \Sigma^i(X, \mathcal{K})$. Similarly, let $\Sigma^{> j}(X, \mathcal{K}) := \cup_{i > j} \Sigma^i(X, \mathcal{K})$ for every integer j . By Verdier's duality, $H^{2d-i}(X, \mathcal{K}^\vee \otimes \mathcal{L}_{\chi^{-1}})$ is the \mathbb{C} -linear dual of $H^i(X, \mathcal{K} \otimes \mathcal{L}_\chi)$. Therefore,

$$\Sigma^{2d-i}(X, \mathcal{K}^\vee) = \{\chi^{-1} : \chi \in \Sigma^i(X, \mathcal{K})\}. \quad (4.26)$$

Fact 4.6.2.1. *Let X be a compact Kähler manifold, and let $\mathcal{K} \in D_c^b(X)$. Then:*

1. ([Wan16, p.547]) *For every integer i , the subset $\Sigma^i(X, \mathcal{K})$ of $\text{Char}(X)$ is Zariski closed.*
2. ([BSS18, Thm. 1.1]) *If X is a complex torus, and if $\mathcal{K} \in \text{Perv}(X)$, then $\Sigma^{\neq 0}(X, \mathcal{K})$ is a strict subset of $\text{Char}(X)$.*
3. ([KW15b, Thm. 1.1 and Lem. 11.2 (c)]) *If further X is a complex abelian variety, then $\Sigma^{\neq 0}(X, \mathcal{K})$ is contained in a thin (and arithmetic when \mathcal{K} is semisimple of geometric origin) subset of $\text{Char}(X)$.*

Corollary 4.6.2.2. *Let X be a compact Kähler manifold, and let $\mathcal{K} \in D_c^b(X)$. Then:*

1. *There are only finitely many integers i such that $\Sigma^i(X, \mathcal{K}) \neq \emptyset$. In particular, $\Sigma^{\neq 0}(X, \mathcal{K})$ and for every integer j , $\Sigma^{> j}(X, \mathcal{K})$ are Zariski closed in $\text{Char}(X)$.*

2. If X is a complex torus, and if $\mathcal{K} \in {}^pD^{\leq m}(X)$ for some integer m , then $\Sigma^{>m}(X, \mathcal{K}) \neq \text{Char}(X)$.
3. If X is a complex abelian variety, and $\mathcal{K} \in {}^pD^{\leq m}(X)$ for some integer m , then $\Sigma^{>m}(X, \mathcal{K})$ is contained in a thin (and arithmetic when \mathcal{K} is semisimple of geometric origin) subset of $\text{Char}(X)$.

Proof. The proof is sketched in [KW15b, p.533].

1. There exist two integers $c < d$ such that $\mathcal{K} \in D^{[c,d]}(X)$. Applying [KS90, Proposition 10.2.12] to the proper morphism $X \rightarrow p$, where p is a point, one gets two integers $a < b$ such that $Rf_*(D^{[c,d]}(X)) \subset D^{[a,b]}(p)$. For every character sheaf \mathcal{L} on X , the functor $* \otimes^L \mathcal{L} : D_c^b(X) \rightarrow D_c^b(X)$ is t-exact with respect to the standard t-structure. Consequently, $\mathcal{K} \otimes^L \mathcal{L} \in D^{[c,d]}(X)$ and hence $Rf_*(\mathcal{K} \otimes^L \mathcal{L}) \in D^{[a,b]}(p)$. For all integers $i \notin [a, b]$, $\Sigma^i(X, \mathcal{K}) = \emptyset$. This shows the first part of the assertion. The second part of the assertion follows from Fact 4.6.2.1 1.
2. By shifting degree, one may assume $m = 0$. For every character sheaf \mathcal{L} on X , the functor $* \otimes^L \mathcal{L} : D_c^b(X) \rightarrow D_c^b(X)$ is t-exact with respect to the perverse t-structure ([KW15b, Prop. 4.1]). Hence for every integer j , ${}^p\mathcal{H}^j(\mathcal{K} \otimes^L \mathcal{L}) = {}^p\mathcal{H}^j(\mathcal{K}) \otimes^L \mathcal{L}$. Consider the subset

$$W = \bigcup_{j \in \mathbb{Z}} \Sigma^{\neq 0}(X, {}^p\mathcal{H}^j(\mathcal{K})) \quad (4.27)$$

of $\text{Char}(X)$. It is in fact a finite union, because by [Dim04, Remark 5.1.19], ${}^p\mathcal{H}^j(K) \neq 0$ for only finitely many integers j . By Fact 4.6.2.1 2, $W \neq \text{Char}(X)$.

For every $\chi \in \text{Char}(X) \setminus W$, consider the Grothendieck spectral sequence from [dCM09, p.545]

$$E_2^{i,j} = H^i(X, {}^p\mathcal{H}^j(\mathcal{K}) \otimes^L \mathcal{L}_\chi) \Rightarrow H^{i+j}(X, \mathcal{K} \otimes^L \mathcal{L}_\chi). \quad (4.28)$$

For any integers $i \neq 0$ and j , one has $H^i(X, {}^p\mathcal{H}^j(\mathcal{K}) \otimes^L \mathcal{L}_\chi) = 0$, so the spectral sequence (4.28) degenerates (in the sense of [Sta24, Tag 011O (2)]) at page E_2 and hence

$$H^j(X, \mathcal{K} \otimes^L \mathcal{L}_\chi) = H^0(X, {}^p\mathcal{H}^j(\mathcal{K}) \otimes^L \mathcal{L}_\chi)$$

for every integer j . Now that $\mathcal{K} \in {}^pD^{\leq 0}(X)$, for every $i > 0$ one has ${}^p\mathcal{H}^i(\mathcal{K}) = 0$ and hence $H^i(X, \mathcal{K} \otimes^L \mathcal{L}_\chi) = 0$. This shows $\chi \notin \Sigma^{>0}(X, \mathcal{K})$. One concludes that $\Sigma^{>0}(X, \mathcal{K}) \subset W$.

3. As ${}^p\mathcal{H}^j(\mathcal{K}) \neq 0$ for only finitely many integers j , by Fact 4.6.2.1 3, the subset W defined by (4.27) is contained in a thin (and arithmetic when \mathcal{K} is semisimple of geometric origin) subset of $\text{Char}(X)$.

□

Theorem 4.6.2.3 is a generic vanishing result for local systems on a manifold admitting Hodge theory. When X is a projective manifold, [PS13, Theorem 1.5] gives a dimension estimate of $\Sigma^k(X, \mathbb{C}_X)$.

Theorem 4.6.2.3. *Let X be a connected regular manifold of dimension n . Let $\alpha : X \rightarrow \text{Alb}(X)$ be the Albanese map associated with some base point and \mathcal{E} be a local system on X . Let k be an integer either $< n - r(\alpha)$ or $> n + r(\alpha)$. Then:*

1. $\Sigma^k(X, \mathcal{E}) \cap \text{Char}^\circ(X)$ is a strict Zariski closed subset of $\text{Char}^\circ(X)$.
2. If furthermore $\text{Alb}(X)$ is algebraic, then $\Sigma^k(X, \mathcal{E}) \cap \text{Char}^\circ(X)$ is contained in a thin subset of $\text{Char}^\circ(X)$.

Proof. In view of (4.26), one may assume $k > d + r(\alpha)$. Set $\mathcal{K} := R\alpha_*\mathcal{E}[d + r(\alpha)]$. We first prove

$$\Sigma^k(X, \mathcal{E}) \cap \text{Char}^\circ(X) = \Sigma^{k-d-r(\alpha)}(\text{Alb}(X), \mathcal{K}) \subset \Sigma^{>0}(\text{Alb}(X), \mathcal{K}). \quad (4.29)$$

This is used in the proof of both 1 and 2.

Indeed, by Proposition 4.5.3.2, the complex of sheaves \mathcal{K} lies in ${}^pD^{\leq 0}(\text{Alb}(X))$. For every $\chi \in \text{Char}^\circ(X)$, let \mathcal{D}_χ (resp. \mathcal{L}_χ) be the corresponding character sheaf on $\text{Alb}(X)$ (resp. on X). Then $\alpha^*\mathcal{D}_\chi = \mathcal{L}_\chi$. By [KW01, Cor. 7.5 (g), p.109], $R\alpha_*(\mathcal{E} \otimes^L \mathcal{L}_\chi) = (R\alpha_*\mathcal{E}) \otimes^L \mathcal{D}_\chi$ in $D_c^b(\text{Alb}(X))$. It follows that

$$H^k(X, \mathcal{E} \otimes \mathcal{L}_\chi) = H^k(\text{Alb}(X), (R\alpha_*\mathcal{E}) \otimes^L \mathcal{D}_\chi) = H^{k-d-r(\alpha)}(\text{Alb}(X), \mathcal{K} \otimes^L \mathcal{D}_\chi),$$

whence (4.29). Now Point 1 follows from Fact 4.6.2.1 1 and Corollary 4.6.2.2 2, and Point 2 follows from Corollary 4.6.2.2 3. □

4.7 Generic vanishing result for manifolds in Fujiki class \mathcal{C}

In Section 4.7.1, we recall the definition of Fujiki class \mathcal{C} , the object of central interest in this note. Then we restrict mainly to algebraic varieties in Section 4.7.2.

4.7.1 Fujiki class \mathcal{C}

Definition 4.7.1.1 (Fujiki class \mathcal{C} , [Uen80, Def. 1]). A compact complex manifold is called in Fujiki class \mathcal{C} if it is the meromorphic image of a compact Kähler manifold.

Every compact Kähler manifold is in Fujiki class \mathcal{C} . The reason why Fujiki class \mathcal{C} is interesting is two-fold. For one thing, this class is large enough in practice. For another, in this class there is a Hodge theory with unitary local systems as coefficients.

Fact 4.7.1.2 ([Tim87, Cor. 5.3], [Ara90, Thm. 1, Thm. 2, Cor. 2]). *Let X be a complex manifold in Fujiki class \mathcal{C} . Then, for every unitary local system \mathcal{E} on X , X is \mathcal{E} -regular.*

In particular, from Fact 4.7.1.2, every manifold in Fujiki class \mathcal{C} is regular. As is explained in Section 4.3 and Section 4.4, the Jacobian and Albanese of a complex manifold in Fujiki class \mathcal{C} behave well.

Theorem 4.7.1.3. *Let X be an n -dimensional complex manifold in Fujiki class \mathcal{C} , and let $\alpha : X \rightarrow \text{Alb}(X)$ be the Albanese map associated with some base point. Let $E \rightarrow X$ be a flat unitary holomorphic vector bundle. Then for any integers $p, q \geq 0$, one has:*

1. *The locus $S^{p,q}(X, E)$ is an analytic subset of $\text{Pic}^0(X)$.*
2. *$S^{n-p, n-q}(X, E^\vee) = \{L \in \text{Pic}^0(X) \mid L^\vee \in S^{p,q}(X, E)\}$.*
3. *If $p + q < n - r(\alpha)$ or $p + q > n + r(\alpha)$, then $S^{p,q}(X, E)$ is contained in a strict (and thin when $\text{Alb}(X)$ is algebraic) subset of $\text{Pic}^0(X)$.*

Proof.

1. The projection $p_2 : X \times \text{Pic}^0(X) \rightarrow \text{Pic}^0(X)$ is a regular family in the sense of [GR84, p.207]. Let $p_1 : X \times \text{Pic}^0(X) \rightarrow X$ be the other projection. Let \mathcal{P} be the universal line bundle on $X \times \text{Pic}^0(X)$ given by Corollary 4.4.2.1. Applying the upper semi-continuity theorem ([GR84, p.210]) to the vector bundle $\mathcal{P} \otimes p_1^* \Omega_X^p$ and the regular family p_2 , one gets that $S^{p,q}(E)$ is an analytic subset of $\text{Pic}^0(X)$.
2. By Serre duality (see, e.g., [Huy05, Prop. 4.1.15]), for every $L \in \text{Pic}(X)$, there is a perfect pairing

$$H^q(X, \Omega_X^p \otimes_{O_X} L \otimes_{O_X} E) \times H^{n-q}(X, \Omega_X^{n-p} \otimes_{O_X} L^\vee \otimes_{O_X} E^\vee) \rightarrow \mathbb{C},$$

so $L \in S^{p,q}(X, E)$ if and only if $L^\vee \in S^{n-p, n-q}(X, E^\vee)$.

3. By Theorem 4.2.3.1, there is a unitary local system \mathcal{E} on X such that $\mathcal{E} \otimes_{\mathbb{C}} O_X$ is isomorphic to E . For each $\chi \in \text{Char}(X)$, let $L_\chi := \mathcal{L}_\chi \otimes_{\mathbb{C}} O_X$. Then the isomorphism (4.19) of real Lie groups is given by $\chi \mapsto L_\chi$. Moreover, the Hodge decomposition (4.6) for $\mathcal{E} \otimes_{\mathbb{C}} \mathcal{L}_\chi$ provided by Fact 4.7.1.2 is

$$H^k(X, \mathcal{E} \otimes_{\mathbb{C}} \mathcal{L}_\chi) = \bigoplus_{p+q=k} H^q(X, \Omega_X^p \otimes_{\mathbb{C}} \mathcal{E} \otimes_{\mathbb{C}} \mathcal{L}_\chi) = H^q(X, \Omega_X^p \otimes_{O_X} E \otimes_{O_X} L_\chi).$$

Therefore, under the isomorphism (4.19), one has

$$\Sigma^k(X, \mathcal{E}) \cap T(X) = \cup_{p+q=k} S^{p,q}(X, E). \quad (4.30)$$

The result follows from Theorem 4.6.2.3.

□

Remark 4.7.1.4. Theorem 4.7.1.3 3 extends Fact 4.1.1.6 from Kähler manifolds to Fujiki class \mathcal{C} . As $\dim X - r(\alpha) \leq \dim \alpha(X)$, the numerical hypothesis in Theorem 4.7.1.3 is more restrictive than that in Fact 4.1.1.1. An example from [GL87, Remark, p.401] is reconsidered in the last paragraph of [KW15b, Sec. 3], to show that the bound $p + q < \dim X - r(\alpha)$ is optimal for Fact 4.1.1.6.

4.7.2 Moishezon manifolds


Moishezon manifolds are examples of manifolds in Fujiki class \mathcal{C} .

Definition 4.7.2.1 (Moishezon manifold, [MM07, Def. 2.2.12]). A connected compact complex manifold X is called Moishezon if it has $\dim X$ algebraically independent meromorphic functions.

In fact, according to [MM07, Thm. 2.2.16], for every Moishezon manifold X , there is a proper modification $\pi : X' \rightarrow X$ with X' a projective manifold. In particular, X is the meromorphic image of a projective manifold, hence in Fujiki class \mathcal{C} . Conversely, a connected compact complex manifold that is the meromorphic image of a projective manifold is Moishezon by the proof of [Voi02, Cor. 12.12]. For more references, see [JM22, Sec. 1].

The intersection of the two subclasses, Kähler and Moishezon, is exactly the class of projective manifolds. More precisely, Moishezon's theorem (see, e.g., [Voi02, Thm. 12.13]) asserts that a Moishezon manifold is Kähler if and only if it is projective. A Moishezon manifold may not be homotopy equivalent to a Kähler manifold ([Ogu94, Thm. 1]). Kodaira-Spencer stability theorem (see, e.g., [Voi02, Thm. 9.1]) shows that small deformations of a Kähler manifold are Kähler. Similarly, small deformations of a regular manifold are regular ([AT13, Cor. 3.7]). By contrast, there is a small deformation of a Moishezon manifold that is not in Fujiki class \mathcal{C} ([Cam91, Sec. 0]). In particular, there exists a regular manifold that is not in Fujiki class \mathcal{C} .

Moishezon manifolds are abundant. For example, for every smooth proper complex algebraic variety X , its analytification X^{an} is a Moishezon manifold ([Har77, p.442]). Hironaka ([Hir60], see also [Har77, p.443]) gives examples of Moishezon manifolds that are not algebraic, and smooth proper algebraic varieties that are not projective. The situation is depicted below. Every inclusion in this graph is strict.



We need Proposition 4.7.2.2 on the algebraicity of Picard torus and Albanese torus to compare them with the Picard variety and Jacobian variety of an algebraic variety.

Proposition 4.7.2.2. *If X is a Moishezon manifold, then $\text{Alb}(X)$ and $\text{Pic}^0(X)$ are complex abelian varieties dual to each other.*

Proof. By [MM07, Thm. 2.2.16], X admits a proper modification $\pi : X' \rightarrow X$ with X' a projective manifold. By [Voi02, Prop. 7.16], the Jacobian $\text{Pic}^0(X')$ is projective. From Proposition 4.4.1.2 5, the torus $\text{Alb}(X')$ is dual to $\text{Pic}^0(X')$, so $\text{Alb}(X')$ is algebraic. By [Uen06, Prop. 9.12, p.107], the morphism $\pi_* : \text{Alb}(X') \rightarrow \text{Alb}(X)$ given by Proposition 4.4.1.2 1 is an isomorphism. \square

Remark 4.7.2.3. By [BL04, p.70], the analytic dual torus of a complex abelian variety is an abelian variety. Moreover, by [MRM74, p.86], the (algebraic) dual abelian variety (defined in [MRM74, p.78]) of a complex abelian variety coincides with its analytic dual torus, so we do not distinguish the two duals in this case.

Remark 4.7.2.4. Another proof of Proposition 4.7.2.2 is as follows. From Lemma D.3.0.1, there is an integer $n \geq 1$ such that $\text{Alb}(X)$ is the image of X^n under certain morphism. As the product of finitely many Moishezon manifolds, X^n is a Moishezon manifold. Then the complex torus $\text{Alb}(X)$ is Moishezon, so projective by Moishezon Theorem.

Let X be a smooth proper complex algebraic variety of dimension n with a base point $x_0 \in X(\mathbb{C})$. Let Sch/\mathbb{C} (resp. Set) be the category of \mathbb{C} -schemes (resp. sets). The fppf-sheaf associated to the functor

$$P_{X/\mathbb{C}} : (\text{Sch}/S)^{\text{op}} \rightarrow \text{Set}, \quad T \mapsto \text{Pic}(X \times_{\mathbb{C}} T)$$

is called the *relative Picard functor* of X ([BLR90, Def. 2, p.201]). From [BLR90, p.211, p.231 and p.233], the relative Picard functor of X is represented by a smooth group scheme $\text{Pic}_{X/\mathbb{C}}$ over \mathbb{C} . In particular, the group $\text{Pic}_{X/\mathbb{C}}(\mathbb{C}) = \text{Pic}(X)$. By [BLR90, Thm. 3, p.232], the identity component $\text{Pic}_{X/\mathbb{C}}^0$ of $\text{Pic}_{X/\mathbb{C}}$ is proper over \mathbb{C} , hence a complex abelian variety called the *Picard variety* of X .

From [Ser58, Thm. 5], there is a complex abelian variety $\text{Alb}(X)$ with a \mathbb{C} -morphism $\alpha_{X,x_0} : (X, x_0) \rightarrow (\text{Alb}(X), 0)$ of pointed varieties satisfying the following universal property (similar to that stated in Proposition 4.4.1.2 3): every \mathbb{C} -morphism of pointed varieties $(X, x_0) \rightarrow (A, 0)$ with A a complex abelian variety factors uniquely through a morphism of abelian varieties $\text{Alb}(X) \rightarrow A$. Such morphism α_{x_0} is unique up to a unique isomorphism. We call $\text{Alb}(X)$ the *algebraic Albanese variety* of X and $\alpha_{X,x_0} : (X, x_0) \rightarrow (\text{Alb}(X), 0)$ the *algebraic Albanese morphism* corresponding to x_0 .

For every \mathcal{O}_X -module F , let F^{an} be the corresponding $\mathcal{O}_{X^{\text{an}}}$ -module defined in [GR71, Exp. XII, 1.3]. Hence a functor

$$\text{Mod}(\mathcal{O}_X) \rightarrow \text{Mod}(\mathcal{O}_{X^{\text{an}}}), \quad F \mapsto F^{\text{an}}.$$

By Serre's GAGA [GR71, Exp. XII, Thm. 4.4], the natural group morphism

$$\text{Pic}(X) \rightarrow \text{Pic}(X^{\text{an}}), \quad L \mapsto L^{\text{an}}$$

is an isomorphism. Corollary 4.7.2.5 2 of GAGA type compares the algebraic Picard variety and the analytic Jacobian. Once again, it is well-known, but a proof is given for the lack of reference.

Corollary 4.7.2.5.

1. *The analytification of $\text{Alb}(X)$ (resp. $\alpha_{X,x_0} : X \rightarrow \text{Alb}(X)$) is $\text{Alb}(X^{\text{an}})$ (resp. $\alpha_{X^{\text{an}},x_0} : X^{\text{an}} \rightarrow \text{Alb}(X^{\text{an}})$).*
2. *The analytification of $\text{Pic}_{X/\mathbb{C}}^0$ is $\text{Pic}^0(X^{\text{an}})$.*

Proof.

1. Since X^{an} is a Moishezon manifold, by Proposition 4.7.2.2, its Albanese torus $\text{Alb}(X^{\text{an}})$ is projective. By Chow's theorem [BL04, Cor. A.4], the map $\alpha_{X^{\text{an}},x_0}$ is algebraic. By Proposition 4.4.1.2 3, every algebraic morphism $(X, x_0) \rightarrow (A, 0)$ to a complex abelian variety A factors uniquely through an analytic (hence algebraic by Chow's theorem again) morphism of complex tori $\text{Alb}(X^{\text{an}}) \rightarrow A^{\text{an}}$. The result follows.
2. By [Moc12, Prop. A.6], the (algebraic) dual abelian variety of $\text{Pic}_{X/\mathbb{C}}^0$ is $\text{Alb}(X)$. By Proposition 4.4.1.2 5, $\text{Pic}^0(X^{\text{an}})$ is the (analytic) dual torus of $\text{Alb}(X^{\text{an}}) = \text{Alb}(X)^{\text{an}}$, so $\text{Pic}^0(X^{\text{an}})$ is the analytification of $\text{Pic}_{X/\mathbb{C}}^0$.

□

Identifying $\text{Pic}_{X/\mathbb{C}}^0$ with $\text{Pic}^0(X^{\text{an}})$ via Corollary 4.7.2.5 2, one can define thin subsets of $\text{Pic}_{X/\mathbb{C}}^0$. Define the defect of semismallness of a proper morphism $f : M \rightarrow N$ between complex algebraic varieties by $r(f) = r(f^{\text{an}})$. With this terminology, we get the following generic vanishing result for smooth proper algebraic varieties.

Corollary 4.7.2.6. *Let \mathcal{E} be a unitary local system on X^{an} , and let $E = \mathcal{E} \otimes_{\mathbb{C}} \mathcal{O}_{X^{\text{an}}}$ be the corresponding holomorphic vector bundle. Then for any integers $p, q \geq 0$ with $p + q > n + r(\alpha)$ or $p + q < n - r(\alpha)$, the locus $S^{p,q}(X^{\text{an}}, E)$ is contained in a thin (and arithmetic when \mathcal{E} is semisimple of geometric origin in $D_c^b(X^{\text{an}})$) subset of $\text{Pic}_{X/\mathbb{C}}^0$.*

Proof. By Corollary 4.7.2.5 1, the analytification $\alpha_{X, x_0}^{\text{an}} : X^{\text{an}} \rightarrow \text{Alb}(X)^{\text{an}}$ coincides with $\alpha_{X^{\text{an}}, x_0} : X^{\text{an}} \rightarrow \text{Alb}(X^{\text{an}})$, and by definition, $r(\alpha) = r(\alpha^{\text{an}})$. From Theorem 4.7.1.3 3, the locus $S^{p,q}(X^{\text{an}}, E)$ is contained in a thin subset of $\text{Pic}^0(X)$.

What remains to show is the assertion in the parentheses. Assume that \mathcal{E} is semisimple of geometric origin. By the decomposition theorem [BBDG82, Thm. 6.2.5], $\mathcal{K} := R\alpha_*\mathcal{E}[n + r(\alpha)]$ is semisimple of geometric origin in $D_c^b(\text{Alb}(X^{\text{an}}))$. By Theorem 4.7.1.3 2, one may assume that $p + q > n + r(\alpha)$, so that

$$S^{p,q}(X^{\text{an}}, E) \subset \Sigma^{p+q}(X^{\text{an}}, \mathcal{E}) \cap T(X) \subset \Sigma^{>0}(\text{Alb}(X), \mathcal{K}),$$

where the first inclusion follows from (4.30) and the second from (4.29). From Corollary 4.6.2.2 3, $\Sigma^{>0}(\text{Alb}(X), \mathcal{K})$ is contained in an arithmetic thin subset of $\text{Pic}_{X/\mathbb{C}}^0$. □

Remark 4.7.2.7. By Chow's theorem, every analytic subset of X^{an} is algebraic. Therefore, $D_c^b(X^{\text{an}})$ coincides with $D_c^b(X(\mathbb{C}), \mathbb{C})$ defined in [BBDG82, p.66] using algebraic Whitney stratifications.

Chapter 5

Fourier-Mukai transform on complex tori, revisited

5.1 Introduction

For a ringed space (Z, \mathcal{O}_Z) , let $D(Z)$ be the derived category of the abelian category of \mathcal{O}_Z -modules. A scheme of finite type and separated over a field is called an algebraic variety. For two algebraic varieties (resp. complex analytic spaces) M, N , let $p_M : M \times N \rightarrow M$ and $p_N : M \times N \rightarrow N$ be the projections. For an object $K \in D(M \times N)$, the *integral transform* $\phi_K^{[M \rightarrow N]} : D(M) \rightarrow D(N)$ with integral kernel K is defined as

$$\phi_K^{[M \rightarrow N]}(\cdot) = R p_{N,*}(K \otimes^L p_M^* \cdot). \quad (5.1)$$

When Z is a complex analytic space, let $D_{\text{good}}(Z) \subset D(Z)$ be the full subcategory consisting of complexes whose cohomology sheaves are good (Definition A.1.4.1). Roughly speaking, an analytic sheaf of modules is good if it can be approximated by coherent submodules. For a complex torus X of dimension g , let \hat{X} be the dual complex torus. Let \mathcal{P} be the normalized¹ Poincaré line bundle on $X \times \hat{X}$. Define functors $RS : D(\hat{X}) \rightarrow D(X)$ and $R\hat{S} : D(X) \rightarrow D(\hat{X})$ by $RS = \phi_{\mathcal{P}}^{[\hat{X} \rightarrow X]}$, $R\hat{S} = \phi_{\mathcal{P}}^{[X \rightarrow \hat{X}]}$. The pair $(RS, R\hat{S})$ is called the *Fourier-Mukai transform* of X . Theorem 5.1.0.1 establishes an analog of the Fourier inversion formula for this pair.

Theorem 5.1.0.1 (Theorem 5.4.1.1). *The functor $R\hat{S}$ (resp. RS) restricts to a functor $D_{\text{good}}(X) \rightarrow D_{\text{good}}(\hat{X})$ (resp. $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(X)$). Moreover, there are natural isomorphisms of functors*

$$\begin{aligned} RS \circ R\hat{S} &\cong [-1]_X^*[-g] : D_{\text{good}}(X) \rightarrow D_{\text{good}}(X), \\ R\hat{S} \circ RS &\cong [-1]_{\hat{X}}^*[-g] : D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(\hat{X}), \end{aligned}$$

¹i.e., both pullback modules $\mathcal{P}|_{X \times 0}$ and $\mathcal{P}|_{0 \times \hat{X}}$ are trivial

where $[-g]$ denotes degree shift.

Theorem 5.1.0.1 is a complex analytic variant of [Muk81, Thm. 2.2] (Statement 5.2.0.4, which has a minor problem for lack of quasi-coherence condition). For complex tori, a parallel false assertion is made as [BBBP07, Thm. 2.1] (Statement 5.2.0.5). Theorem 5.1.0.1 shows that “good sheaves” on complex manifolds serve as substitutes for “quasi-coherent sheaves” on algebraic varieties in this case. As an application, we recover Matsushima-Morimoto’s classification of homogeneous vector bundles on complex tori.

Theorem (Theorem 5.5.3.6). *A vector bundle F on the complex torus X is translation invariant if and only if there is an integer $n \geq 0$, unipotent vector bundles² U_1, \dots, U_n on X and $P_1, \dots, P_n \in \text{Pic}^0(X)$, such that F is isomorphic to $\bigoplus_{i=1}^n (P_i \otimes U_i)$.*

Notation and conventions

For a topological space M , the category of abelian sheaves on M is denoted by $\text{Ab}(M)$. The category of ringed spaces is denoted by RingS . For a ringed space (X, O_X) , let $\text{Mod}(O_X)$ be the category of O_X -modules. The full subcategory of $\text{Mod}(O_X)$ comprised of quasi-coherent (resp. coherent) O_X -modules in the sense of Definition A.1.1.1 3 (resp. 6) is denoted by $\text{Qch}(X)$ (resp. $\text{Coh}(X)$). For a closed subset $Z \subset X$, let $\text{Coh}_Z(X) \subset \text{Coh}(X)$ be the full subcategory consisting of modules with support contained in Z .

Given a symbol $* \in \{\emptyset, +, -, b\}$, the notation $D^*(X)$ refers to the unbounded/bounded below/bounded above/bounded derived category of $\text{Mod}(O_X)$ in order. The full subcategory of $D^*(X)$ consisting of the complexes whose cohomologies are coherent (resp. quasi-coherent) is denoted by $D_c^*(X)$ (resp. $D_{\text{qc}}^*(X)$). Denote by $R\mathcal{H}om_X : D(X)^{\text{op}} \times D(X) \rightarrow D(X)$ the internal hom bifunctor constructed in [Sta24, Tag 08DH].

For a locally ringed space X and $x \in X$, let $i_x : (x, O_{X,x}) \rightarrow (X, O_X)$ be the canonical morphism of locally ringed spaces. For an $O_{X,x}$ -module M , the O_X -module $(i_x)_* M$ is denoted by M_x .

All complex analytic spaces (in the sense of [KK83, Def. 43.2]) are assumed to be paracompact. Let An be the category of complex analytic spaces. The dimension of a complex manifold always refers to the complex dimension, which is assumed to be finite.

When X is an abelian variety (resp. complex torus), its dual abelian variety (resp. complex torus) is denoted by \hat{X} . The normalized Poincaré bundle on $X \times \hat{X}$ is denoted by \mathcal{P} . For $y \in \hat{X}$ (resp. $x \in X$), let P_y (resp. P_x) denote the line bundle $\mathcal{P}|_{X \times y}$ (resp. $\mathcal{P}|_{x \times \hat{X}}$).

²Definition 5.5.2.6

5.2 Fourier-Mukai transform

Complex tori are generalizations of complex abelian varieties. Every complex torus of dimension 1 is an abelian variety. By contrast, for every integer $g \geq 2$, a very general complex torus of dimension g is not³ an abelian variety (see, e.g., [BZ23b, p.21]).

The Fourier-Mukai transform is an analog of the classical Fourier transform. It is proposed by Mukai [Muk81] on abelian varieties and complex tori. Let k be an algebraically closed field. Let X be an abelian variety over k (resp. a complex torus) of dimension g . Write RS and $R\hat{S}$ for $\phi_{\mathcal{P}}^{[\hat{X} \rightarrow X]}$ and $\phi_{\mathcal{P}}^{[X \rightarrow \hat{X}]}$ respectively. The pair $(RS, R\hat{S})$ is called the Fourier-Mukai transform of X . The functor RS (resp. $R\hat{S}$) restricts to a functor $D^b(\hat{X}) \rightarrow D^b(X)$ (resp. $D^b(X) \rightarrow D^b(\hat{X})$).

Let X be an abelian variety. The usual exchange of translation and time shifting (resp. multiplication and convolution) of Fourier transform finds analog for Fourier-Mukai transform, namely the exchange of translation and line bundle twisting (resp. tensor product and Pontrjagin product) in [Muk81, (3.1) (resp. (3.7))]. Moreover, Mukai proves a duality theorem similar to the classical Fourier inversion formula.

Fact 5.2.0.1. *[Algebraic Mukai duality] There are canonical isomorphisms of functors*

$$\begin{aligned} RS \circ R\hat{S} &\cong [-1]_X^*[-g] : D_{\text{qc}}(X) \rightarrow D_{\text{qc}}(X); \\ R\hat{S} \circ RS &\cong [-1]_{\hat{X}}^*[-g] : D_{\text{qc}}(\hat{X}) \rightarrow D_{\text{qc}}(\hat{X}). \end{aligned}$$

In particular, the functor $RS : D_{\text{qc}}(\hat{X}) \rightarrow D_{\text{qc}}(X)$ is an equivalence of categories, with a quasi-inverse $[-1]_{\hat{X}}^ \circ R\hat{S}[g]$.*

Example 5.2.0.2 ([Muk81, Eg. 2.6]). For every $y \in \hat{X}(k)$, one has $RS(k_y) = P_y$ and $R\hat{S}(P_y) = k_{-y}[-g]$.

Remark 5.2.0.3. Combining Fact 5.2.0.1, the natural equivalence $D(\text{Qch}(X)) \rightarrow D_{\text{qc}}(X)$ ([BN93, Cor. 5.5]) with the compatibility of derived direct images [TT90, Cor. B.9], one gets [Rot96, Mukai's Theorem, p.569] stated for $D^b(\text{Qch}(*))$ instead of $D_{\text{qc}}(*)$. The quasi-coherence restriction is essential for Čech resolution with respect to affine covers in [Rot96, p.571].

The proof of Fact 5.2.0.1 uses projection formula and the flat base change theorem ([Lip60, Prop. 3.9.4; Prop. 3.9.5]). Compared with Fact

³To the contrary, it is incorrectly implied in [BBR94, p.151] that every complex torus of dimension 2 admits a compatible structure of algebraic complex surface. In fact, it fails for each 2-dimensional complex torus X that is not a projective manifold. For otherwise, assume there is a complex algebraic surface V with $V^{\text{an}} \cong X$. Then V is proper by [GR71, XII, Prop. 3.2 (v)]. In consequence, the algebraic variety V is projective by [Har77, p.357]. Thus, X is a projective manifold, a contradiction.

5.2.0.1, the original statement (Statement 5.2.0.4) has no quasi-coherence restriction.

Statement 5.2.0.4 ([Muk81, Thm. 2.2]). The functor RS gives an equivalence of categories between $D(\hat{X})$ and $D(X)$, and its quasi-inverse is $[-1]_{\hat{X}}^* \circ R\hat{S}[g]$.

In [BBBP07, Thm. 2.1], an assertion similar to Statement 5.2.0.4 is made for complex tori.

Statement 5.2.0.5. Let X be a complex torus. Then the integral transform $RS : D^b(\hat{X}) \rightarrow D^b(X)$ is an equivalence of triangulated categories.

However, Lemma 5.2.0.6 shows that Statement 5.2.0.4 (resp. Statement 5.2.0.5) holds if and only if $g = 0$.

Lemma 5.2.0.6 ([th16]). *Let X be an abelian variety or a complex torus. If the functor $RS : D^b(\hat{X}) \rightarrow D^b(X)$ is an equivalence of categories, then $g = 0$.*

Proof. When X is a complex torus, let $k = \mathbb{C}$. In both cases, let $F = k_0^{\mathbb{N}}$ be the product of a countable infinite family of k_0 in $\text{Mod}(O_{\hat{X}})$. Since $k^{\mathbb{N}} = k^{\oplus I}$ as a k -module for some index set I , the direct sum sheaf $k_0^{\oplus I}$ is isomorphic to F . Therefore, by [Sta24, Tag 07D9 (2)], F is the direct sum of I copies of k_0 in $D^b(\hat{X})$. We claim that F is the product of \mathbb{N} copies of k_0 in $D^b(\hat{X})$.

By [Gro57b, p.129], the abelian category $\text{Mod}(O_{\hat{X},0})$ satisfies the AB 4*) axiom. From [Sta24, Tag 07KC (2)], the inclusion $\text{Mod}(O_{\hat{X},0}) \rightarrow D^b(\text{Mod}(O_{\hat{X},0}))$ commutes with countable products. Let $i : 0 \rightarrow \hat{X}$ be the closed immersion. Since $i_* : \text{Mod}(O_{\hat{X},0}) \rightarrow \text{Mod}(O_{\hat{X}})$ is exact, there is a commutative square

$$\begin{array}{ccc} \text{Mod}(O_{\hat{X},0}) & \xrightarrow{i_*} & \text{Mod}(O_{\hat{X}}) \\ \downarrow & & \downarrow \\ D^b(\text{Mod}(O_{\hat{X},0})) & \xrightarrow{Ri_*} & D^b(\hat{X}). \end{array}$$

Since $Ri_* : D^b(\text{Mod}(O_{\hat{X},0})) \rightarrow D^b(\hat{X})$ has a left adjoint, it commutes with products. As $F = i_*(k^{\mathbb{N}})$, the claim is proved.

As $RS : D^b(\hat{X}) \rightarrow D^b(X)$ is an equivalence, inside $D^b(X)$, the object $RS(F)$ is the direct sum of I copies of $RS(k_0)$, as well as the product of \mathbb{N} copies of $RS(k_0)$. By Example 5.2.0.2 (when X is an abelian variety) and Lemma 5.2.0.8 (when X is a complex torus), one has $RS(k_0) = O_X$. Therefore, $RS(F)$ is isomorphic to $O_X^{\oplus I}$ and to $O_X^{\mathbb{N}}$ in $\text{Mod}(O_X)$.

Assume the contrary $g > 0$. Then there is a nonempty *connected* open subset $V \subset X$, such that $O_X(V)$ is an integral domain but not a field. In particular, the ring $O_X(V)$ is not Artinian. By [Har77, II, Exercise 1.11] (when X is an abelian variety) and Corollary A.1.5.4 (when X is a complex

torus), the $O_X(V)$ -module $\Gamma(V, RS(F))$ is isomorphic to $O_X(V)^{\oplus I}$ and to $O_X(V)^{\mathbb{N}}$. However, this contradicts Fact 5.2.0.7. \square

Fact 5.2.0.7 ([Len68, Thm, p.211]). *If A is a commutative ring such that $A^{\mathbb{N}}$ is a free A -module, then A is Artinian.*

For algebraic varieties, the analog of Lemma 5.2.0.8 follows from the flat base change theorem and the projection formula.

Lemma 5.2.0.8. *Let X, Y be two complex analytic spaces, let $K \in D(X \times Y)$, and let $x \in X$. Consider the closed embedding $h_x : Y \rightarrow X \times Y$, $y \mapsto (x, y)$. Then $\phi_K^{[X \rightarrow Y]}(\mathbb{C}_x) = Lh_x^* K$.*

Proof. Let $p : X \times Y \rightarrow X$, $q : X \times Y \rightarrow Y$ be the two projections. Denote the closed embedding of complex analytic spaces $x \rightarrow X$ by j_x . The cartesian square

$$\begin{array}{ccc} Y & \xrightarrow{p_0} & x \\ \downarrow h_x & \square & \downarrow j_x \\ X \times Y & \xrightarrow{p} & X \end{array}$$

in the category \mathbf{An} induces a natural morphism $\phi : p^* \mathbb{C}_x \rightarrow Rh_{x,*} O_Y$ in $\text{Mod}(O_{X \times Y})$. Both sheaves are supported on $\{x\} \times Y$.

For two (Hausdorff) locally convex topological vector spaces E, F over \mathbb{C} , the completed projective topological tensor product $E \hat{\otimes}_{\mathbb{C}} F$ is defined in [Gro55, Ch. I, Déf. 2, p.32]. For every $y \in Y$, by [GR84, p.27], the stalk $O_{X \times Y, (x, y)} = O_{X, x} \hat{\otimes}_{\mathbb{C}} O_{Y, y}$. Then

$$(p^* \mathbb{C}_x)_{(x, y)} = \mathbb{C} \otimes_{O_{X, x}} O_{X \times Y, (x, y)} = O_{Y, y}.$$

Therefore, $\phi_{(x, y)} : (p^* \mathbb{C}_x)_{(x, y)} \rightarrow (h_{x,*} O_Y)_{(x, y)}$ is an isomorphism. Thus, ϕ is an isomorphism.

By [Sta24, Tag 0B55], the natural morphism $(Rh_{x,*} O_Y) \otimes^L K \rightarrow Rh_{x,*}(Lh_x^* K)$ is an isomorphism. Then

$$\begin{aligned} \phi_K^{[X \rightarrow Y]}(\mathbb{C}_x) &= Rq_*(p^* \mathbb{C}_x \otimes^L K) \cong Rq_*(Rh_{x,*} O_Y \otimes^L K) \\ &\cong Rq_* Rh_{x,*}(Lh_x^* K) \cong R(qh_x)_*(Lh_x^* K) = Lh_x^* K. \end{aligned}$$

\square

The minor problem with Statement 5.2.0.4 occurs in the proof of [Muk81, Prop. 1.3], when the flat base change theorem [Har66, Prop. 5.12] stated for objects of $D_{qc}(\ast)$ is applied to objects in $D^-(\ast)$. Similarly, the minor problem with Statement 5.2.0.5 originates from a lack of certain analytic quasi-coherence in the wrong Statement 5.2.0.9 (a counterpart of [Muk81, Prop. 1.3]). A modification of Statement 5.2.0.9 is Proposition 5.4.2.3.

Statement 5.2.0.9 ([BBBP07, p.427]). If M , N , and P are compact complex manifolds and $K \in D^b(M \times N)$ and $L \in D^b(N \times P)$, then one has a natural isomorphism of functors from $D^b(M)$ to $D^b(P)$:

$$\phi_L^{[N \rightarrow P]} \circ \phi_K^{[M \rightarrow N]} \cong \phi_{K * L}^{[M \rightarrow P]},$$

where

$$K * L = Rp_{M \times P*}(p_{M \times N}^* K \otimes^L p_{N \times P}^* L) \in D^b(M \times P),$$

and $p_{M \times N}$, $p_{M \times P}$, $p_{N \times P}$ are the natural projections $M \times N \times P \rightarrow M \times N$, etc.

When X is an abelian variety of positive dimension, by Fact 5.2.0.1, $RS(F)$ is the product of \mathbb{N} copies of O_X in $\text{Qch}(X)$. It is not isomorphic to $O_X^{\mathbb{N}}$ by Lemma 5.2.0.10.

Lemma 5.2.0.10. *Let X be an integral scheme with generic point η . If the O_X -module $O_X^{\mathbb{N}}$ is quasi-coherent, then the natural morphism $\eta \rightarrow X$ is an isomorphism.*

Proof. Consider an arbitrary affine open $U = \text{Spec}(A) \subset X$. Then A is an integral domain of fraction field $\kappa(\eta)$. We show that the natural inclusion $A \rightarrow \kappa(\eta)$ is an isomorphism.

For otherwise, there exists $f \in A \setminus (A^* \cup \{0\})$. Let $D_f \subset U$ be the corresponding standard open subset. Note $\Gamma(U, O_X^{\mathbb{N}}) = A^{\mathbb{N}}$ and $\Gamma(D_f, O_X^{\mathbb{N}}) = (A_f)^{\mathbb{N}}$. As $O_X^{\mathbb{N}} \in \text{Qch}(X)$, the natural A_f -module morphism $\Gamma(U, O_X^{\mathbb{N}})_f \rightarrow \Gamma(D_f, O_X^{\mathbb{N}})$ is an isomorphism. Or equivalently, the natural map $\phi : (A^{\mathbb{N}})_f \rightarrow (A_f)^{\mathbb{N}}$ is an isomorphism.

In particular, there exists $a = (a_0, a_1, \dots) \in A^{\mathbb{N}}$ and an integer $m \geq 0$ such that $\phi(a/f^m) = (1/f^i)_{i \geq 0}$. Then $a_{m+1} = f^{-1}$ in A_f . There exists an integer $n \geq 0$ such that $(a_{m+1}f - 1)f^n = 0$ in A . Since A is a domain, $a_{m+1}f - 1 = 0$ in A . This contradicts the fact that $f \notin A^*$.

Therefore, the natural morphism $\eta \rightarrow U$ is an isomorphism. The proof is completed as U is taken arbitrarily. \square

Lemma 5.2.0.11 computes the derived restriction of a relatively flat module, which is a partial converse to [Huy06, Lemma 3.31] in the analytic setting.

Lemma 5.2.0.11. *Let $f : S \rightarrow X$ be a flat morphism of complex analytic spaces, and let K be an O_S -module flat over X . For $x \in f(S)$, let $i_x : S_x \rightarrow S$ be the inclusion of the fiber over x . Then $Li_x^* K = i_x^* K$.*

Proof. To simplify the notation, we denote i_x by i . By [Sta24, Tag 0B55], the natural morphism

$$Ri_* O_{S_x} \otimes_{O_S}^L K \rightarrow Ri_*(Li^* K) \tag{5.2}$$

is an isomorphism. They are supported on S_x , since for every integer n one has

$$H^n(Ri_*(Li^*K)) = Ri_*H^n(Li^*K) = i_*H^n(Li^*K).$$

For every $s \in S_x$, the morphism $j : (s, O_{S,s}) \rightarrow S$ of ringed spaces is flat and $j^* : \text{Mod}(O_S) \rightarrow \text{Mod}(O_{S,s})$ is taking the stalk at s . Let m_x be the maximal ideal of $O_{X,x}$. As the ring map $f_s^\# : O_{X,x} \rightarrow O_{S,s}$ is flat, one has

$$O_{S_x,s} = (O_{X,x}/m_x) \otimes_{O_{X,x}} O_{S,s} = (O_{X,x}/m_x) \otimes_{O_{X,x}}^L O_{S,s}. \quad (5.3)$$

By [Sta24, Tag 079U], one has

$$\begin{aligned} Lj^*(Ri_*O_{S_x} \otimes_{O_S}^L K) &= Lj^*Ri_*O_{S_x} \otimes_{O_{S,s}}^L Lj^*K \\ &= O_{S_x,s} \otimes_{O_{S,s}}^L K_s = [(O_{X,x}/m_x) \otimes_{O_{X,x}}^L O_{S,s}] \otimes_{O_{S,s}}^L K_s \\ &= (O_{X,x}/m_x) \otimes_{O_{X,x}}^L (O_{S,s} \otimes_{O_{S,s}}^L K_s) \\ &= (O_{X,x}/m_x) \otimes_{O_{X,x}}^L K_s = (O_{X,x}/m_x) \otimes_{O_{X,x}} K_s, \end{aligned} \quad (5.4)$$

where the third (resp. fourth, resp. last) equality uses (5.3) (resp. Lemma 5.4.2.1, resp. the flatness of the $O_{X,x}$ -module K_s).

Then for every integer $n \neq 0$, every $s \in S_x$, the stalk

$$[H^n(Ri_*O_{S_x} \otimes_{O_S}^L K)]_s = H^n[Lj^*(Ri_*O_{S_x} \otimes_{O_S}^L K)] = H^n((O_{X,x}/m_x) \otimes_{O_{X,x}} K_s) = 0,$$

where the second equality uses (5.4). Hence

$$i_*H^n(Li^*K) = H^n[Ri_*(Li^*K)] \cong H^n(Ri_*O_{S_x} \otimes_{O_S}^L K) = 0,$$

where the second equality uses (5.2). Thus, for every integer $n \neq 0$, $H^n(Li^*K) = 0$ in $\text{Mod}(O_{S_x})$. \square

Remark 5.2.0.12. Lemmas 5.2.0.8 and 5.2.0.11 yield an analytic version of [Huy06, Eg. 5.4 vi]): Let X, Y be two complex analytic spaces. Let $x \in X$ and K be an $O_{X \times Y}$ -module flat over X . Then $\phi_K^{[X \rightarrow Y]}(\mathbb{C}_x) = K|_{\{x\} \times Y}$.

Remark 5.2.0.13. Here is an example showing the necessity of the flatness of f in Lemma 5.2.0.11.

Let $A = \mathbb{C}[t]$ and $B = \mathbb{C}[x, y]/xy$. Then the B -module xB (resp. yB) is isomorphic to B/y (resp. B/x). Let $S = \text{Spec}(B)$ and $X = \text{Spec}(A) = A_{\mathbb{C}}^1$. The morphism $A \rightarrow B$ of k -algebras defined by $t \mapsto x$ induces a morphism $f : S \rightarrow X$ of schemes. Let K be the coherent O_S -module corresponding to the B -module B/y . Then K is flat over X , because the ring map composition $A \rightarrow B \rightarrow B/y$ is an isomorphism. Let $i : S_0 \rightarrow S$ be the inclusion of the fiber over $0 \in X(\mathbb{C})$. Then i is a closed immersion defined by ideal $xB \subset B$, so Li^*K is induced by $K \otimes_B^L (B/x)$. By [Osb12, Exercise 9, b), p.72],

$$\text{Tor}_2^B(B/y, B/x) = (yB) \otimes_B (xB) \cong (B/x) \otimes_B (B/y) = B/(x, y) = \mathbb{C}.$$

In particular, $Li^*K \neq i^*K$. Taking analytification one gets $L(i^{\text{an}})^*K^{\text{an}} \neq (i^{\text{an}})^*K^{\text{an}}$.

Corollary 5.2.0.14 follows from Lemma 5.2.0.11, and it is an analytic counterpart of [Huy06, Example 5.4 vi)].

Corollary 5.2.0.14. *In Lemma 5.2.0.8, if $K \in \text{Mod}(O_{X \times Y})$ is flat over X , then $\phi_K^{[X \rightarrow Y]}(\mathbb{C}_x) = i^*K$.*

By Corollary 5.2.0.14 and Theorem 5.4.1.1, Example 5.2.0.2 remains true when X is a complex torus.

5.3 Good modules

As Section 5.2 explains, to obtain an analytic analogue of Fact 5.2.0.1, it is necessary to find a substitute for quasi-coherence on complex manifolds. We show that goodness introduced by Kashiwara (Definition A.1.4.1) can be used as such.

5.3.1 Functoriality

In Corollary 5.3.1.16, we prove that goodness is preserved by integral transforms. To prove this, we show that goodness is preserved by the operations involved in (5.1).

Example 5.3.1.1. [Har66, Example 1., p.68] Let $f : X \rightarrow Y$ be a morphism of ringed spaces. Then the derived pullback $Lf^* : D(Y) \rightarrow D(X)$ (constructed in [Spa88, Prop. 6.7 (a)]) is bounded above (in the sense of [Lip60, 1.11.1]), and the derived pushout $Rf_* : D(X) \rightarrow D(Y)$ is bounded below.

Proposition 5.3.1.2 (Pullback). *Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces. Then $Lf^* : D(Y) \rightarrow D(X)$ restricts to a functor*

1. $D_c^b(Y) \rightarrow D_c^b(X)$ when Y is a complex manifold or f is flat;
2. $D_{\text{good}}(Y) \rightarrow D_{\text{good}}(X)$.

Proof.

1. Because Y is smooth or f is flat, by Lemma 5.3.1.3, the morphism f has finite tor-dimension. Thus, Lf^* restricts to a functor $D^b(Y) \rightarrow D^b(X)$.

Consider $F \in D_c^b(Y)$. To prove that $Lf^*F \in D_c^b(X)$, by [Har66, I, Prop. 7.3 (i)], one may assume $F \in \text{Coh}(Y)$. This case is proved by Lemma A.1.3.3.

2. (a) Let $G \in D_{\text{good}}^-(Y)$. By Example 5.3.1.1, Lemma A.1.4.3 3 and a dual of [Har66, Prop. 7.3 (ii)], to prove $Lf^*G \in D_{\text{good}}(X)$, one may assume $G \in \text{Good}(Y)$. Let U be a relatively compact open subset of X . Then $f(\bar{U})$ is a compact subset of Y , so contained in a relatively compact open subset V of Y . Since G is good, its restriction $G|_V = \sum_{i \in I} G_i$ is the sum of a directed family of coherent \mathcal{O}_V -submodules of $G|_V$. Let $g : f^{-1}(V) \rightarrow V$ be the base change of f along the inclusion $V \rightarrow Y$. As Lf^* commutes with colimits, one has

$$(Lf^*G)|_{f^{-1}(V)} = \text{colim}_{i \in I} Lg^*G_i.$$

For every integer n , in $\text{Mod}(\mathcal{O}_{f^{-1}(V)})$ one has

$$\begin{aligned} H^n(Lf^*G)|_{f^{-1}(V)} &= H^n((Lf^*G)|_{f^{-1}(V)}) \\ &= H^n(\text{colim}_{i \in I} Lg^*G_i) = \text{colim}_{i \in I} H^n(Lg^*G_i). \end{aligned}$$

Since G_i is coherent, by Lemma A.1.3.3, the $\mathcal{O}_{f^{-1}(V)}$ -module $H^n(Lg^*G_i)$ is coherent. By Lemma A.1.4.3 3, the $\mathcal{O}_{f^{-1}(V)}$ -module $H^n(Lf^*G)|_{f^{-1}(V)}$ is good. Since \bar{U} is a compact subset of $f^{-1}(V)$, the subset U is relatively compact in $f^{-1}(V)$. Hence, $H^n(Lf^*G)|_U$ is the sum of a directed family of coherent submodules. Hence $Lf^*G \in D_{\text{good}}(X)$.

- (b) Then consider the general case $C \in D_{\text{good}}(Y)$. For every integer $m \geq 0$, the m -th canonical truncation ([Sta24, Tag 0118 (4)]) $C_m := \tau^{\leq m}C$ is in $D_{\text{good}}^-(Y)$. From the proof of [Lip60, Prop. 2.5.5], there is a bounded above complex of flat \mathcal{O}_Y -modules Q_m with a quasi-isomorphism $Q_m \rightarrow C_m$ that is functorial in C_m . Moreover, the complex $Q := \text{colim}_m Q_m$ is K-flat (in the sense of [Spa88, Def. 5.1]), and the canonical morphism $Q \rightarrow C$ is a quasi-isomorphism. Because $Lf^* : D(Y) \rightarrow D(X)$ admits a right adjoint, it commutes with colimits. Thus, the resulting morphisms

$$\text{colim}_m Lf^*Q_m \rightarrow Lf^*Q \rightarrow Lf^*C$$

are isomorphisms in $D(X)$.

Let $\text{Ch}(\text{Mod}(\mathcal{O}_X))$ be the category of chain complexes over $\text{Mod}(\mathcal{O}_X)$. The directed set \mathbb{N} can be seen naturally as a category. Define a functor $\mathbb{N} \rightarrow \text{Ch}(\text{Mod}(\mathcal{O}_X))$, $m \mapsto f^*Q_m$. Because $\text{Mod}(\mathcal{O}_X)$ is a Grothendieck abelian category, for every integer n , by [Hov99, Lem. 1.5], the natural morphism

$$\text{colim}_m H^n(f^*Q_m) \rightarrow H^n(\text{colim}_m f^*Q_m)$$

in $\text{Mod}(O_X)$ is an isomorphism. Hence an isomorphism $H^n(Lf^*C) \cong \text{colim}_m H^n(Lf^*Q_m)$ in $\text{Mod}(O_X)$. Since $Q_m \in D_{\text{good}}^-(Y)$, by Case 2a, the O_X -module $H^n(Lf^*Q_m)$ is good. By Lemma A.1.4.3 3, so is the O_X -module $H^n(Lf^*C)$.

□

The tor-dimension $\text{tor-dim } f$ of a morphism $f : X \rightarrow Y$ of ringed spaces is defined to be the lower dimension (in the sense of [Lip60, 1.11.1]) of the functor $Lf^* : D^-(Y) \rightarrow D(X)$. If f is flat, then $\text{tor-dim } f = 0$. If f has finite tor-dimension, then $Lf^* : D^-(Y) \rightarrow D(X)$ restricts to a functor $D^b(Y) \rightarrow D^b(X)$. The weak dimension $\text{wgld}(R)$ of a commutative ring R is defined to be the supremum of flat dimension of all R -modules.

Lemma 5.3.1.3. *Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces, with Y a complex manifold. Then f has finite tor-dimension.*

Proof. From [Lip60, (2.7.6.4)], one only needs to show that for every $x \in X$, the flat dimension of the $O_{Y,f(x)}$ -module $O_{X,x}$ is uniformly bounded. By definition, the flat dimension of every $O_{Y,f(x)}$ -module is bounded by the weak dimension of the ring $O_{Y,f(x)}$. Because Y is a complex manifold, the local ring $O_{Y,f(x)}$ is Noetherian regular. By Lemma 5.3.1.4, $\text{wgld } O_{Y,f(x)}$ is the Krull dimension of $O_{Y,f(x)}$, which coincides with the dimension of the complex manifold Y near $f(x)$. □

Lemma 5.3.1.4 (Serre). *Let R be a commutative, Noetherian, regular local ring. Then $\text{wgld}(R)$ coincides with the Krull dimension of R , hence finite.*

Proof. From [Osb12, Cor. 4.21], the weak dimension coincides with the global dimension of R . By Serre's theorem (see, e.g., [Osb12, p.332]), the global dimension equals the Krull dimension, which is finite. □

Proposition 5.3.1.5 (Tensor product). *Let X be a complex analytic space. Then the bifunctor (constructed in [Spa88, Thm. A. (ii)]) $\otimes^L : D(X) \times D(X) \rightarrow D(X)$ restricts to a bifunctor*

1. $D^b(X) \times D^b(X) \rightarrow D^b(X)$ (resp. $D_c^b(X) \times D_c^b(X) \rightarrow D_c^b(X)$) when X is a complex manifold;
2. $D_{\text{good}}(X) \times D_{\text{good}}(X) \rightarrow D_{\text{good}}(X)$.

Proof.

1. The weak dimension of a ringed space (M, O_M) is defined to be $\sup_{x \in M} \text{wgld}(O_{M,x})$. By [HT07, (C.2.20)], to prove the statement for $D^b(X)$, it suffices to bound the weak dimension of X . As X is smooth, for every $x \in X$, the stalk $O_{X,x}$ is a Noetherian, regular local ring. Thus, by Lemma 5.3.1.4, its weak dimension $\text{wgld}(O_{X,x})$ is equal to

the dimension of the complex manifold X near x . Therefore, the weak dimension of X is at most $\dim X$

Consider any $F, G \in D_c^b(X)$. To prove that $F \otimes^L G \in D_c^b(X)$, by [Har66, I, Prop. 7.3 (i)], one may assume $F, G \in \text{Coh}(X)$. Then the conclusion follows from [GH78, 4., p.700].

2. Take $F, G \in D_{\text{good}}(X)$. To prove that $F \otimes^L G \in D_{\text{good}}(X)$, as in the proof of Proposition 5.3.1.2 2, one may assume that $F, G \in D_{\text{good}}^-(X)$. By a dual of [Har66, I, Prop. 7.3 (ii)], one may assume that $F, G \in \text{Good}(X)$. Let U be a relatively compact open subset of X .

For every integer n , we claim that the O_U -module $H^n(F \otimes_{O_X}^L G)|_U$ is good. By assumption, the restrictions $F|_U = \sum_{i \in I} F_i$ and $G|_U = \sum_{j \in J} G_j$ can be written as sums of directed families of coherent submodules. By [Sta24, Tag 08DJ], the functor $\otimes_{O_U}^L(G|_U) : D(U) \rightarrow D(U)$ has a right adjoint, so

$$(F \otimes^L G)|_U = \text{colim}_{i \in I} [F_i \otimes^L (G|_U)]. \quad (5.5)$$

By [Sta24, Tag 05NI (2)], there exists a complex C^\bullet of flat O_U -modules and a quasi-isomorphism $C^\bullet \rightarrow G|_U$. Then for every $i \in I$, in $D(U)$

$$F_i \otimes_{O_U} C^\bullet \xrightarrow{\sim} F_i \otimes_{O_U}^L G|_U. \quad (5.6)$$

Define a functor $I \rightarrow \text{Ch}(\text{Mod}(O_X))$ by $i \mapsto F_i \otimes C^\bullet$. By [Hov99, Lem. 1.5], the natural morphism

$$\text{colim}_{i \in I} H^n(F_i \otimes C^\bullet) \rightarrow H^n(\text{colim}_{i \in I} (F_i \otimes C^\bullet))$$

in $\text{Mod}(O_U)$ is an isomorphism. Combining it with (5.5) and (5.6), one gets an isomorphism in $\text{Mod}(O_U)$

$$\text{colim}_{i \in I} H^n(F_i \otimes_{O_U}^L G|_U) \rightarrow H^n(F \otimes_{O_X}^L G)|_U.$$

Because $\text{Good}(U)$ is closed under colimits in $\text{Mod}(O_U)$ by Lemma A.1.4.3 3, one may assume that $F|_U$ is coherent. Similarly, one may assume further that $G|_U$ is coherent. Then the claim follows from Lemma A.1.3.4.

□

Remark 5.3.1.6. Proposition 5.3.1.5 2 can also be derived from Proposition 5.3.1.2 2 as in the proof of [Bjö93, Thm. 3.2.13 (3)].

As the proof of Theorem 5.3.1.7 is lengthy, we split it into a series of lemmas.

Theorem 5.3.1.7 (Pushout). *Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces. If $\dim X$ is finite, then $Rf_* : D(X) \rightarrow D(Y)$ restricts to a functor $D_{\text{good}}(X) \rightarrow D_{\text{good}}(Y)$ (resp. $D_{\text{good}}^b(X) \rightarrow D_{\text{good}}^b(Y)$).*

Proof. By Lemma 5.3.1.11, the functor Rf_* restricts to a functor $D^b(X) \rightarrow D^b(Y)$. We show that $Rf_*F \in D_{\text{good}}(Y)$ for every $F \in D_{\text{good}}(X)$. By [Har66, I, Prop. 7.3 (iii)], Lemmas 5.3.1.11 and A.1.4.3 3, one may assume that $F \in \text{Good}(X)$. For every relatively compact open subset $V \subset Y$, its closure \bar{V} is compact in Y . As f is proper, the preimage $f^{-1}(\bar{V})$ is compact. Thus, $U := f^{-1}(V)$ is a relatively compact open subset of X . Since F is good, $F|_U = \text{colim}_{i \in I} F_i$, where $\{F_i\}_{i \in I}$ is a directed family of coherent \mathcal{O}_U -submodules of $F|_U$. Let $g : U \rightarrow V$ be the base change of f . Fix an integer n . By Lemma 5.3.1.9, in $\text{Mod}(\mathcal{O}_V)$

$$(R^n f_* F)|_V = R^n g_*(F|_U) = \text{colim}_{i \in I} R^n g_* F_i.$$

As a base change of f , the morphism g is proper. Then by Fact 5.3.1.8, for every $i \in I$, the \mathcal{O}_V -module $R^n g_* F_i$ is coherent. By $\text{Coh}(V) \subset \text{Good}(V)$ and Lemma A.1.4.3 3, the \mathcal{O}_V -module $(R^n f_* F)|_V$ is good. Therefore, $Rf_* F \in D_{\text{good}}(Y)$. □

Fact 5.3.1.8 (Grauert direct image theorem, see e.g., [GR84, p.207]). *Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces. Then $Rf_* : D(X) \rightarrow D(Y)$ restricts to a functor $\text{Coh}(X) \rightarrow D_c(Y)$.*

Lemma 5.3.1.9. *Let $f : X \rightarrow Y$ be a proper map between locally compact, Hausdorff spaces. Then for every integer $n \geq 0$, the functor $R^n f_* : \text{Ab}(X) \rightarrow \text{Ab}(Y)$ commutes with filtrant colimits.*

Proof. Let $(F_i, f_{ij})_{i \in I}$ be a filtrant inductive system with colimit F in $\text{Ab}(X)$. Since the abelian category $\text{Ab}(Y)$ is Grothendieck, the filtrant colimit $G = \text{colim}_{i \in I} R^n f_* F_i$ exists and there is a canonical morphism $\phi : G \rightarrow R^n f_* F$ in $\text{Ab}(Y)$. For every $y \in Y$, the functor $\text{Ab}(Y) \rightarrow \text{Ab}$ taking the stalk at y commutes with colimits, so $G_y = \text{colim}_{i \in I} (R^n f_* F_i)_y$. By [Mil13, Thm. 17.2], for every i the stalk $(R^n f_* F_i)_y = H^n(X_y, F_i|_{X_y})$. Then by [God58, Thm. 4.12.1], the morphism $\phi_y : G_y \rightarrow (R^n f_* F)_y$ is an isomorphism. Therefore, ϕ is an isomorphism. □

The proof of Fact 5.3.1.10 is similar to that of [KS90, Prop. 3.2.2].

Fact 5.3.1.10. *Let X be a locally compact, Hausdorff topological space which is countable at infinity. Suppose that there is an integer $n \geq 0$ such that every point of X has an open neighborhood homeomorphic to a locally closed subset of \mathbb{R}^n . Then for every abelian sheaf F on X and every integer $j > n$, one has $H^j(X, F) = 0$.*

Lemma 5.3.1.11. *Let X be a complex analytic space of finite dimension n . Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces. Then for an object $E \in D(X)$ with $H^m(E) = 0$ for every integer $m > 0$, one has $H^i(Rf_*E) = 0$ for every integer $i > 2n$. In particular, the functor $Rf_* : D(X) \rightarrow D(Y)$ is bounded.*

Proof. For every open subset $V \subset Y$ and every \mathcal{O}_X -module M , from $i > 2n$ and Fact 5.3.1.10, one has $H^i(f^{-1}(V), M) = 0$. Applying Lemma 5.3.1.14 to the functor $\Gamma(f^{-1}(V), \cdot) : \text{Mod}(\mathcal{O}_X) \rightarrow \text{Ab}$, one gets

$$H^i(R\Gamma(f^{-1}(V), E)) = H^i(R\Gamma(f^{-1}(V), \tau^{\geq 1}E)) = 0.$$

By Lemma 5.3.1.13, the \mathcal{O}_Y -module $H^i(Rf_*E) = 0$. □

Remark 5.3.1.12. The finite dimension condition in Lemma 5.3.1.11 is necessary. For every integer $m \geq 1$, let T_m be a complex torus of dimension m , and let $f_m : T_m \rightarrow \text{Specan } \mathbb{C}$ be the canonical morphism. Let $f : X \rightarrow Y$ be $\sqcup_{m \geq 1} f_m : \sqcup_{m \geq 1} T_m \rightarrow \sqcup_{m \geq 1} \text{Specan } \mathbb{C}$. Then f is proper. For every integer $q \geq 1$, the sheaf $R^q f_* \mathcal{O}_X \neq 0$.

Lemma 5.3.1.13 is a derived version of [Har77, III, Prop. 8.1].

Lemma 5.3.1.13. *Let $f : X \rightarrow Y$ be a continuous map of topological spaces. Then for every integer i and every $F \in D(\text{Ab}(X))$, the sheaf $H^i(Rf_*F)$ on Y is the sheaf associated to the abelian presheaf $V \mapsto H^i R\Gamma(f^{-1}(V), F)$.*

Proof. By [Spa88, Thm. D], there is a quasi-isomorphism $F \rightarrow I$, where I is a K-injective complex of abelian sheaves on X . Then the canonical morphism $Rf_*F \rightarrow f_*I$ is an isomorphism in $D(\text{Ab}(Y))$. By [Mur06, Lem. 3], $H^i(Rf_*F)$ is the sheaf associated the presheaf

$$V \mapsto H^i(\Gamma(V, f_*I)) = H^i(\Gamma(f^{-1}(V), I)) = H^i(R\Gamma(f^{-1}(V), F)).$$

□

Lemma 5.3.1.14. *Let X be a ringed space as in Fact 5.3.1.10. Let $F : \text{Mod}(\mathcal{O}_X) \rightarrow \text{Ab}$ be an additive functor. Assume that F commutes with countable products, and there is an integer $N \geq 0$ with $R^p F(M) = 0$ for every integer $p \geq N$ and every $M \in \text{Mod}(\mathcal{O}_X)$. Then the right derived functor $RF : D(X) \rightarrow D(\text{Ab})$ exists. Moreover, for any integers $i \geq j$, the natural transformation*

$$H^i(RF \cdot) \rightarrow H^i(RF(\tau^{\geq j-N+1} \cdot)) : D(X) \rightarrow \text{Ab}$$

is an isomorphism.

Proof. The existence of N and [Wei95, Cor. 10.5.11] show that $RF : D^+(X) \rightarrow D^+(\text{Ab})$ extends to a right derived functor $RF : D(X) \rightarrow D(\text{Ab})$ of F .

For every integer m and every $E \in D(X)$, set $E_m := \tau^{\geq -m} E$. Then $\{E_m\}_{m \in \mathbb{Z}}$ forms an inverse system in $D(X)$. Let n be as in Fact 5.3.1.10. Then for every open subset $U \subset X$, any integers $p(> n)$ and q , one has $H^p(U, H^q(E)) = 0$. Then by [Sta24, Tag 0D64], the canonical morphism $E \rightarrow R \lim_m E_m$ is an isomorphism in $D(X)$. Since F commutes with countable products, from [Sta24, Tag 08U1], in $D(\text{Ab})$ one has $RF(E) \xrightarrow{\sim} R \lim_m RF(E_m)$. For every integer i , by [Sta24, Tag 08U5], there is a short exact sequence in the category Ab

$$0 \rightarrow R^1 \lim_m H^{i-1}(RF(E_m)) \rightarrow H^i(RF(E)) \rightarrow \lim_m H^i(RF(E_m)) \rightarrow 0. \quad (5.7)$$

We claim that $R^1 \lim_m H^{i-1}(RF(E_m)) = 0$.

For every integer $m \geq N - i$, by [Sta24, Tag 08J5], there is an exact triangle

$$H^{-m}(E)[m] \rightarrow E_m \rightarrow E_{m-1} \xrightarrow{+1} H^{-m}(E)[m+1] \quad (5.8)$$

in $D(X)$. By assumption, one has

$$\begin{aligned} H^i(RF(H^{-m}(E)[m])) &= R^{i+m}F((H^{-m}(E))) = 0; \\ H^i(RF(H^{-m}(E)[m+1])) &= R^{i+m+1}F((H^{-m}(E))) = 0. \end{aligned}$$

Taking the long exact sequence associated with (5.8), one concludes that the canonical morphism $H^i(RF(E_m)) \rightarrow H^i(RF(E_{m-1}))$ in Ab is an isomorphism. Since the inverse system $\{H^i(RF(E_m))\}_{m \geq 1}$ is constant starting with $m = N - i - 1$, it satisfies the Mittag-Leffler condition in the sense of [Sta24, Tag 02N0]. From [Sta24, Tag 07KW (3)], one obtains

$$R^1 \lim_m H^i(RF(E_m)) = 0,$$

which proves the claim.

When $i \geq j$, as the inverse system is constant from $m = N - j - 1$, one has $\lim_m H^i(RF(E_m)) = H^i[RF(E_{N-j-1})]$. Then the sequence (5.7) induces an isomorphism $H^i(RF(E)) \rightarrow H^i(RF(\tau^{\geq j-N+1} E))$. \square

Remark 5.3.1.15. In the statement of Lemma 5.3.1.14, because $\text{Mod}(O_X)$ is a Grothendieck abelian category, it has enough injectives. By [Ver66, p.338], the total right derived functor $RF : D^+(X) \rightarrow D^+(\text{Ab})$ exists (even if F may not be left exact).

Corollary 5.3.1.16. *Let X, Y be complex manifolds (resp. complex analytic spaces), with X compact and Y finite dimensional. If F is an object of $D_c^b(X \times Y)$ (resp. $D_{\text{good}}(X \times Y)$), then $\phi_F^{[X \rightarrow Y]}$ restricts to a functor $D_c^b(X) \rightarrow D_c^b(Y)$ (resp. $D_{\text{good}}(X) \rightarrow D_{\text{good}}(Y)$).*

Proof. Because X is compact, its dimension is finite and the projection $X \times Y \rightarrow Y$ is proper. Thus, $X \times Y$ is finite dimensional. The result is a combination of Proposition 5.3.1.2 1 (resp. 2), Proposition 5.3.1.5 1 (resp. 2), Fact 5.3.1.8 and Lemma 5.3.1.11 (resp. Theorem 5.3.1.7). \square

Remark 5.3.1.17. Although we don't need the functors $R\mathcal{H}om$, $f_!$ and $f^!$, it is interesting to know whether they preserve goodness or not.

5.3.2 Base change theorems

As a replacement for the (algebraic) flat base change theorem (used in Mukai's proof of Fact 5.2.0.1), we give an analytic smooth base change theorem. It is a consequence of Theorem 5.3.2.3 and Fact 5.3.2.2.

Consider a cartesian square in the category \mathbf{An} :

$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ \downarrow f' & \square & \downarrow f \\ S' & \xrightarrow{g} & S. \end{array} \quad (5.9)$$

Then [Sta24, Tag 08HY] gives a natural transformation of functors $D(X) \rightarrow D(S')$

$$Lg^* Rf_* \rightarrow Rf'_* Lg'^*, \quad (5.10)$$

coming from the adjunction in [Sta24, Tag 079W].

Smooth base change

Definition 5.3.2.1. A morphism $g : S' \rightarrow S$ of complex analytic spaces is called locally product, if for every $s' \in S'$, there is an open neighborhood U of $s' \in S'$ and a complex analytic space Z , such that $g(U)$ is open in S and there is a $g(U)$ -isomorphism $U \rightarrow g(U) \times Z$.

By [CD94, II, Cor. 2.7], a locally product morphism is flat.

Fact 5.3.2.2 ([Gro61b, Thm. 3.1]). *A morphism of complex analytic spaces is smooth (in the sense of in the sense of [Gro61b, Déf. 3.2]) if and only if it is a submersion (in the sense of [Fis76, p.100]). In particular, a smooth morphism is locally product.*

Theorem 5.3.2.3. *Consider the square (5.9) with both $\dim X$ and $\dim X'$ finite, $f : X \rightarrow S$ proper and $g : S' \rightarrow S$ locally product. Then (5.10) restricts to an isomorphism of functors $D_{\text{good}}(X) \rightarrow D_{\text{good}}(S')$.*

We begin the proof with several lemmas.

Definition 5.3.2.4. A morphism of complex analytic spaces $g : S' \rightarrow S$ is said to satisfy property \mathcal{Q}_S if for every *proper* morphism $f : X \rightarrow S$ of complex analytic spaces, every coherent \mathcal{O}_X -module F and every integer $i \geq 0$, the base change morphism $g^* R^i f_* F \rightarrow R^i f'_*(g'^* F)$ induced by (5.9) is an isomorphism in $\text{Mod}(\mathcal{O}_{S'})$.

Lemma 5.3.2.5 shows that the property \mathcal{Q} is local on the source and the target.

Lemma 5.3.2.5. *Let $g : S' \rightarrow S$ and be a morphism of complex analytic spaces.*

1. *Let $h : S'' \rightarrow S'$ be another morphism of complex analytic spaces. If g and h satisfy \mathcal{Q}_S and $\mathcal{Q}_{S'}$ respectively, then gh satisfies \mathcal{Q}_S .*
2. *Assume that $\{S'_i\}_{i \in I}$ (resp. $\{S_j\}_{j \in J}$) is an open covering of S' (resp. S) such that for every $i \in I$ (resp. $j \in J$), the morphism $g|_{S'_i} : S'_i \rightarrow S$ (resp. $g^{-1}(S_j) \rightarrow S_j$) satisfies \mathcal{Q}_S (resp. \mathcal{Q}_{S_j}). Then g satisfies \mathcal{Q}_S .*
3. *If g is an open embedding of complex analytic spaces, then g satisfies \mathcal{Q}_S .*

Proof. 1. The proof is similar to that of [Day23, Lem. 2.13 (2)].

2. It follows from the local nature of sheaves.

3. The proof is similar to that of [Har77, III, Cor. 8.2].

□

Lemma 5.3.2.6. *Let $f : X \rightarrow S$ be a proper morphism of complex analytic spaces, with S Stein. Then for every coherent \mathcal{O}_X -module F and every integer $n \geq 0$, one has $H^n(X, F) = H^0(S, R^n f_* F)$.*

Proof. By properness of f and Fact 5.3.1.8, the \mathcal{O}_S -module $R^n f_* F$ is coherent. As S is Stein, from Cartan's Theorem B (see, e.g., [KK83, Sec. 52, Thm. B]), for every integer $m > 0$ one has $H^m(S, R^n f_* F) = 0$. The conclusion follows from [Sta24, Tag 01F4 (2)]. □

Remark 5.3.2.7. As an application of Lemma 5.3.2.6, we give an enhancement of Lemma 5.3.1.11 for good modules. Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces with $\dim X$ finite. Then for every good \mathcal{O}_X -module G and every integer $n > \dim X$, one has $R^n f_* G = 0$.

Assume first that G is coherent. For every Stein open subset $V \subset Y$, from Cartan's Theorem A (see e.g., [GR04, Theorem A, p.XVIII]), the restriction $R^n f_* G|_V$ is generated by sections $H^0(V, R^n f_* G|_V)$. By Lemma 5.3.2.6, one has

$$H^0(V, R^n f_* G|_V) = H^n(f^{-1}(V), G|_{f^{-1}(V)}),$$

which vanishes by [Rei64, Cor., p.2333]. Thus, $R^n f_* G|_V = 0$. Hence $R^n f_* G = 0$.

Assume now that $G \in \text{Good}(X)$ is arbitrary. For every relatively compact open subset $W \subset Y$, the open subset $f^{-1}(W)$ of X is relatively compact. Then there is a directed family of coherent submodules $\{G_i\}_{i \in I}$ of $G|_{f^{-1}(W)}$ such that $G|_{f^{-1}(W)} = \text{colim}_{i \in I} G_i$. By Lemma 5.3.1.9, one gets $(R^n f_* G)|_W = \text{colim}_{i \in I} R^n(f|_{f^{-1}(W)})_* G_i = 0$. Hence $R^n f_* G = 0$.

Lemma 5.3.2.8. *Let X, Y be complex analytic spaces, with Y Stein. Let $p : X \times Y \rightarrow X$ be the projection. Then for every coherent \mathcal{O}_X -module F and every integer $i \geq 0$, the natural morphism $H^i(X, F) \hat{\otimes}_{\mathbb{C}} \mathcal{O}_Y(Y) \rightarrow H^i(X \times Y, p^* F)$ of locally convex topological vector spaces is an isomorphism.*

Proof. Choose a Stein covering \mathcal{U} of X . Let C^\bullet be the Čech complex of F relative to \mathcal{U} . Then $H^i(C^\bullet) = H^i(X, F)$. By [EP⁺96, Prop. 4.1.5], for every integer q , the q -th term C^q of the complex C^\bullet is a Fréchet space. Moreover, $\{U \times Y : U \in \mathcal{U}\}$ forms a Stein covering of $X \times Y$. By [EP⁺96, Prop. 4.2.3; Thm. 4.2.4], the Čech complex of $p^* F$ relative to this Stein covering is $C^\bullet \hat{\otimes}_{\mathbb{C}} \mathcal{O}(Y)$. Therefore, $H^i(C^\bullet \hat{\otimes}_{\mathbb{C}} \mathcal{O}(Y)) = H^i(X \times Y, p^* F)$. By [EP⁺96, Prop. 4.1.5], $\mathcal{O}(Y)$ is a unital Fréchet nuclear algebra, so from [EP⁺96, Thm. A1.6 (d)], the functor $* \hat{\otimes}_{\mathbb{C}} \mathcal{O}(Y)$ preserves exact sequences, hence commutes with taking cohomology groups of the Čech complexes. \square

We consider the special case of products.

Corollary 5.3.2.9. *Let S, Z be two complex analytic spaces. Then the projection $S \times Z \rightarrow S$ satisfies \mathcal{Q}_S .*

Proof. Fix a proper morphism $X \rightarrow S$ of complex analytic spaces and a coherent \mathcal{O}_X -module F . By Lemma 5.3.2.5, we may assume that S, Z are Stein spaces. Then the result follows from Lemma 5.3.2.6, Lemma 5.3.2.8 and [EP⁺96, Prop. 4.2.3; Thm. 4.2.4]. \square

Corollary 5.3.2.10. *Every locally product morphism $g : S' \rightarrow S$ of complex analytic spaces satisfies \mathcal{Q}_S .*

Proof. Fix $s' \in S'$, and let $s = g(s')$. Since g is locally product, there is an open neighborhood U (resp. V) of $s' \in S'$ (resp. $s \in S$), a complex analytic space Z and an isomorphism $\psi : U \rightarrow Z \times V$ of complex analytic spaces such that the diagram

$$\begin{array}{ccc} U & \xrightarrow{\psi} & Z \times V \\ \downarrow g|_U & \swarrow p_2 & \\ V & & \end{array}$$

commutes, where p_2 is the projection to the second factor. By Corollary 5.3.2.9, $g|_U : U \rightarrow V$ satisfies \mathcal{Q}_V . By Lemma 5.3.2.5, the morphism $g : S' \rightarrow S$ satisfies \mathcal{Q}_S . \square

Proof of Theorem 5.3.2.3. The morphism f' is a base change of f , hence a proper morphism. Because $\dim X, \dim X'$ are finite, by Theorem 5.3.1.7 and Proposition 5.3.1.2 2, the functors Lg^*Rf_* and $Rf'_*Lg'^*$ restrict to functors $D_{\text{good}}(X) \rightarrow D_{\text{good}}(S')$.

For every $K \in D_{\text{good}}(X)$, we prove that the base change morphism $Lg^*Rf_*K \rightarrow Rf'_*Lg'^*K$ in $D(S')$ is an isomorphism. By Lemma 5.3.1.11, the functors $Rf_* : D(X) \rightarrow D(S)$ and $Rf'_* : D(X') \rightarrow D(S')$ are bounded. From [Har66, I, Prop. 7.1 (iii)] and Lemma A.1.4.3 3, one may assume that $K \in \text{Good}(X)$. For every $s' \in S'$, there is a relatively compact open neighborhood $V \subset S$ of $g(s')$. The preimage $f^{-1}(V)$ is a relatively compact open subset of X . Consider the base change of the square (5.9) along the open embedding $V \rightarrow S$:

$$\begin{array}{ccc} f^{-1}(V) \times_V g^{-1}V & \xrightarrow{v'} & f^{-1}(V) \\ \downarrow u' & \square & \downarrow u \\ g^{-1}(V) & \xrightarrow{v} & V. \end{array}$$

Because g is locally product, so is v . One can write $K|_{f^{-1}(V)} = \text{colim}_{i \in I} G_i$, where $\{G_i\}_{i \in I}$ is a directed family of coherent submodules of $K|_{f^{-1}(V)}$. By Lemma 5.3.1.9, the natural morphism

$$(g^*R^i f_* K)|_{g^{-1}(V)} \rightarrow R^i f'_*(g'^* K)|_{g^{-1}(V)} \quad (5.11)$$

in $\text{Mod}(O_{g^{-1}(V)})$ is the colimit of the morphisms

$$v^* R^i u_* G_i \rightarrow R^i u'_* v'^* G_i.$$

By Corollary 5.3.2.10, for all $i \in I$, they are isomorphisms. Then (5.11) is an isomorphism. \square

Remark 5.3.2.11. In the proof of [BBR94, Lem. 5], an analytic flat base change result is applied without further justification. In [MS08, p.153], a flat base change theorem for cartesian squares in the category of complex manifolds is stated, referring to [Spa88] for the proof. However, the cited result [Spa88, Prop. 6.20] is for cartesian squares in the category RingS . In general, a cartesian square in the category of complex manifolds is not cartesian in RingS . For example, the complex vector space \mathbb{C}^2 is the product of two copies of \mathbb{C} in the category of complex manifolds, but is not the product even in the subcategory $\text{LRS} \subset \text{RingS}$ of locally ringed space.⁴

In fact, by [Gil11, Cor. 5], the product E of two copies of \mathbb{C} in LRS exists. By the universal property of E , there is a unique morphism $f : \mathbb{C}^2 \rightarrow E$ in

⁴By contrast, every cartesian square in the category of schemes remains cartesian in LRS ([Sta24, Tag 01JN]).

LRS induced by the two projections $p_i : \mathbb{C}^2 \rightarrow \mathbb{C}$. Let $o = f(0) \in E$. We claim that the local ring $O_{E,o}$ is not Noetherian.

The local ring $A := O_{\mathbb{C},0} = \mathbb{C}\{z\}$ is the ring of convergent power series. Let $B = A \otimes_{\mathbb{C}} A$. Let $\epsilon : B \rightarrow A$ be the surjective (diagonal) morphism defined by $\epsilon(f \otimes g) = fg$. Set $I = \ker(\epsilon)$. Let $c : A \rightarrow \mathbb{C}$ be the ring map taking the constant term. Then $c\epsilon : B \rightarrow \mathbb{C}$ is surjective, so $m = \ker(c\epsilon)$ is a maximal ideal of B containing I . Set $S = B \setminus m$. Then $O_{E,o} = S^{-1}B$. From [Tu97, p.367], I/I^2 is a free B/I -module of *infinite* rank. Thus, $S^{-1}(I/I^2) = (S^{-1}I)/(S^{-1}I^2)$ is a free $S^{-1}(B/I) = (S^{-1}B)/(S^{-1}I)$ -module of infinite rank. In particular, the ideal $S^{-1}I$ of the ring $S^{-1}B$ is not finitely generated. The claim is proved.

By [GH78, p.679], the ring $\mathbb{C}\{x, y\}$ is Noetherian. Thus, the local morphism $f_0^\# : O_{E,o} \rightarrow O_{\mathbb{C}^2,0} = \mathbb{C}\{x, y\}$ is not an isomorphism. Hence, f is not an isomorphism in LRS.

Non-smooth base change

Remark 5.3.2.12. A base change theorem for algebraic varieties may not have a direct generalization to complex analytic spaces. For example, the affine base change theorem [Sta24, Tag 02KG] fails for morphism of Stein manifolds. In the cartesian square (5.9), assume that $S = \text{Specan } \mathbb{C}$ is a point, $X = \mathbb{C}$ and S' is a positive-dimensional complex manifold. Then there is an open subset $U \subset S'$ isomorphic to an open ball in \mathbb{C}^n with $n > 0$. On the one hand, by Cartan's Theorem B, $Rf_*O_X = f_*O_X = O_{\mathbb{C}}(\mathbb{C})$. Thus, $g^*Rf_*O_X$ is a free $O_{S'}$ -module of infinite rank $\dim_{\mathbb{C}} O_{\mathbb{C}}(\mathbb{C})$. From Corollary A.1.5.4, $\Gamma(U, g^*Rf_*O_X) = O_U(U) \otimes_{\mathbb{C}} O_{\mathbb{C}}(\mathbb{C})$. On the other hand, one has $f'^{-1}(U) = U \times \mathbb{C}$ and $g'^*O_X = O_{X'}$, so

$$\begin{aligned} \Gamma(U, f'_*g'^*O_X) &= \Gamma(f'^{-1}(U), O_{X'}) \\ &\stackrel{(a)}{=} \Gamma(U \times \mathbb{C}, O_{U \times \mathbb{C}}) = O_U(U) \hat{\otimes}_{\mathbb{C}} O_{\mathbb{C}}(\mathbb{C}), \end{aligned}$$

where (a) uses [EP⁺96, p.75]. The natural morphism $\Gamma(U, g^*Rf_*O_X) \rightarrow \Gamma(U, f'_*g'^*O_X)$ is not an isomorphism, so the base change morphism $g^*f_*O_X \rightarrow f'_*g'^*O_X$ is not an isomorphism.

Lemma 5.3.2.13 is used in the proof of Proposition 5.5.1.2.

Lemma 5.3.2.13 (Base change). *Consider the cartesian square (5.9) with $\dim X, \dim S'$ finite and f flat proper. Then (5.10) induces an isomorphism $Lg^*Rf_* \rightarrow Rf'_*Lg'^*$ of functors $D_{\text{good}}(X) \rightarrow D_{\text{good}}(S')$.*

Proof. Because $\dim X$ is finite, by Theorem 5.3.1.7 and Proposition 5.3.1.2 2, the functor $Lg^*Rf_* : D(X) \rightarrow D(S')$ restricts to a functor $D_{\text{good}}(X) \rightarrow D_{\text{good}}(S')$. Consider the following commutative diagram

$$\begin{array}{ccccc}
& & g' & & \\
& \nearrow & & \searrow & \\
X' & \xrightarrow{i'} & S' \times X & \xrightarrow{p'} & X \\
\downarrow f' & & \downarrow \text{Id}_{S'} \times f & & \downarrow f \\
S' & \xrightarrow{i} & S' \times S & \xrightarrow{p} & S, \\
& \nwarrow & & \nearrow & \\
& & g & &
\end{array}$$

where the morphism $i : S' \rightarrow S' \times S$ is defined by $i(s') = (s', g(s'))$, and $p : S' \times S \rightarrow S$ is the projection. Then i is a closed embedding of complex analytic spaces.

Because p is locally product, by Theorem 5.3.2.3, the natural transformation $Lp^*Rf_* \rightarrow R(\text{Id}_{S'} \times f)_*Lp'^* : D_{\text{good}}(X) \rightarrow D_{\text{good}}(S' \times S)$ is an isomorphism. Because f is flat proper, so is $\text{Id}_{S'} \times f$. Moreover, $\dim(S' \times X) = \dim S' + \dim X$ is finite. Thus, there are isomorphism of functors $D_{\text{good}}(X) \rightarrow D_{\text{good}}(S')$

$$\begin{aligned}
Lg^*Rf_* &\cong Li^*Lp^*Rf_* \xrightarrow{\sim} Li^*R(\text{Id}_{S'} \times f)_*Lp'^* \\
&\stackrel{(a)}{\xrightarrow{\sim}} Rf'_*Li'^*Lp'^* \cong Rf'_*Lg'^*,
\end{aligned} \tag{5.12}$$

where the isomorphism (a) uses Lemma 5.3.2.14 2. By [Sta24, Tag 0E47], the isomorphism (5.12) is induced by (5.10). \square

Lemma 5.3.2.14. *In the cartesian square (5.9), assume that g is a closed embedding of complex analytic spaces. Then:*

1. *The base change morphism $f^*g_*O_{S'} \rightarrow g'_*O_{X'}$ in $\text{Mod}(O_X)$ is an isomorphism.*
2. *If f is flat proper and X has finite dimension, then (5.10) is an isomorphism.*

Proof. 1. Let I be the kernel of the canonical surjection $O_S \rightarrow g_*O_{S'}$ in $\text{Mod}(O_S)$. Since $f^* : \text{Mod}(O_S) \rightarrow \text{Mod}(O_X)$ is right exact, the sequence

$$f^*I \rightarrow O_X \rightarrow f^*g_*O_{S'} \rightarrow 0$$

is exact in $\text{Mod}(O_X)$. Because g is a closed embedding, by [Gro61a, Remarque 2.10], the square (5.9) is cartesian in the category RingS . Then from [Gro61a, 9-05], the cokernel of the morphism $f^*I \rightarrow O_X$ in $\text{Mod}(O_X)$ is $g'_*O_{X'}$. Therefore, the morphism $f^*g_*O_{S'} \rightarrow g'_*O_{X'}$ is an isomorphism.

2. As g is a closed embedding, the functor $g_* : \text{Ab}(S') \rightarrow \text{Ab}(S)$ is exact and $g^{-1}g_* = \text{Id}_{\text{Ab}(S')}$. Therefore, the functor $Rg_* = g_* : D(S') \rightarrow$

$D(S)$ is conservative in the sense of [Rie17, p.180]. Thus, it suffices to show that the natural transformation

$$Rg_*Lg^*Rf_*E \rightarrow Rg_*Rf'_*Lg'^*E \xrightarrow{\sim} Rf_*Rg'_*Lg'^*E \quad (5.13)$$

of functors $D(X) \rightarrow D(S)$ is an isomorphism. By [Sta24, Tag 0B55], the natural morphisms

$$\begin{aligned} (Rg_*O_{S'}) \otimes_{O_S}^L Rf_*E &\rightarrow Rg_*Lg^*Rf_*E, \\ (Rg'_*O_{X'}) \otimes_{O_X}^L E &\rightarrow Rg'_*Lg'^*E \end{aligned}$$

are isomorphisms. One has

$$Rg'_*O_{X'} = g'_*O_{X'} \xleftarrow{(a)} f^*g_*O_{S'} \xrightarrow{(b)} Lf^*Rg_*O_{S'},$$

where (a) uses Point 1, and (b) uses the flatness of f . Thus, the natural transformation (5.13) becomes

$$(Rg_*O_{S'}) \otimes_{O_S}^L Rf_*E \rightarrow Rf_*(Lf^*Rg_*O_{S'} \otimes_{O_X}^L E).$$

It is an isomorphism by the finiteness of $\dim X$, the properness of f and Fact 5.3.2.15. □

From Fact 5.3.1.10, one gets Fact 5.3.2.15 as a special case of [Spa88, Prop. 6.18]. A slight variant can also be derived from [KS90, Prop. 2.6.6] and Lemma 5.4.2.1.

Fact 5.3.2.15 (Projection formula). *Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces. If $\dim X$ is finite, then there is a canonical isomorphism $(Rf_! -) \otimes_{O_Y}^L (+) \rightarrow Rf_!(- \otimes_{O_X}^L Lf^* +)$ of bifunctors $D(X) \times D(Y) \rightarrow D(Y)$.*

5.3.3 Compatibility

For a complex algebraic variety X , let $\psi_X : X^{\text{an}} \rightarrow X$ be its complex analytification. With quasi-coherence condition, the algebraic and analytic integral transforms are compatible.

Corollary 5.3.3.1. *Let X, Y be two complex algebraic varieties, with X proper. Then for every $K \in D_{\text{qc}}(X \times Y)$, the natural square*

$$\begin{array}{ccc} D(X) & \xrightarrow{\phi_K^{[X \rightarrow Y]}} & D(Y) \\ \downarrow \psi_X^* & & \downarrow \psi_Y^* \\ D(X^{\text{an}}) & \xrightarrow{\phi_{K^{\text{an}}}^{[X^{\text{an}} \rightarrow Y^{\text{an}}]}} & D(Y^{\text{an}}), \end{array}$$

restricts to a commutative square

$$\begin{array}{ccc}
D_{\text{qc}}(X) & \xrightarrow{\phi_K^{[X \rightarrow Y]}} & D_{\text{qc}}(Y) \\
\downarrow \psi_X^* & & \downarrow \psi_Y^* \\
D_{\text{good}}(X^{\text{an}}) & \xrightarrow{\phi_{K^{\text{an}}}^{[X^{\text{an}} \rightarrow Y^{\text{an}}]}} & D_{\text{good}}(Y^{\text{an}}).
\end{array} \tag{5.14}$$

Proof. From [Sta24, Tag 08DW (1)], [Sta24, Tag 08DX (1)] and [Sta24, Tag 08D5 (1)], the functor $\phi_K^{[X \rightarrow Y]}$ restricts to a functor $D_{\text{qc}}(X) \rightarrow D_{\text{qc}}(Y)$. By Corollary 5.3.1.16 and compactness of X^{an} , the functor $\phi_{K^{\text{an}}}^{[X^{\text{an}} \rightarrow Y^{\text{an}}]}$ restricts to a functor $D_{\text{good}}(X^{\text{an}}) \rightarrow D_{\text{good}}(Y^{\text{an}})$. By Lemma B.2.0.2, the functor ψ_X^* (resp. ψ_Y^*) restricts to a functor $D_{\text{qc}}(X) \rightarrow D_{\text{good}}(X^{\text{an}})$ (resp. $D_{\text{qc}}(Y) \rightarrow D_{\text{good}}(Y^{\text{an}})$).

By [Sta24, Tag 0D5S] (resp. [Sta24, Tag 079U]), analytification commutes with derived pullback (resp. tensor product). As X is proper over \mathbb{C} , the projection $p_Y : X \times Y \rightarrow Y$ is proper. By Proposition B.3.0.1, analytification commutes with derived direct image. Thus, the square (5.14) is commutative. \square

Remark 5.3.3.2. Fact B.2.0.1 (Theorem B.4.0.2) proves Corollary 5.4.1.2 (resp. Theorem 5.4.1.1) for complex tori that are *algebraic*. Because if X is a complex abelian variety, then every functor in the square

$$\begin{array}{ccc}
D_c^b(X) & \xrightarrow{R\hat{S}} & D_c^b(\hat{X}) \\
\downarrow \psi_X^* & & \downarrow \psi_{\hat{X}}^* \\
D_c^b(X^{\text{an}}) & \xrightarrow{R\hat{S}} & D_c^b(\hat{X}^{\text{an}})
\end{array}
\quad
\begin{array}{ccc}
D_{\text{qc}}(X) & \xrightarrow{R\hat{S}} & D_{\text{qc}}(\hat{X}) \\
\downarrow \psi_X^* & & \downarrow \psi_{\hat{X}}^* \\
D_{\text{qc}}(X^{\text{an}}) & \xrightarrow{R\hat{S}} & D_{\text{qc}}(\hat{X}^{\text{an}})
\end{array}$$

is an equivalence. In fact, by [Huy06, Def. 5.1] and the natural equivalence $D^b(\text{Coh}(X)) \rightarrow D_c^b(X)$ in [FJJ⁺71, Exp. II, Cor. 2.2.2.1], the functor $R\hat{S} : D(X) \rightarrow D(\hat{X})$ restricts to a functor $D_c^b(X) \rightarrow D_c^b(\hat{X})$. The functor on the top of the square is an equivalence by Fact 5.2.0.1. From Fact B.2.0.1, the vertical functors are also equivalences. From Corollary 5.3.1.16, the functor $R\hat{S}$ restricts to a functor $D_c^b(X^{\text{an}}) \rightarrow D_c^b(\hat{X}^{\text{an}})$. The commutativity of the square follows from Corollary 5.3.3.1.

5.4 Analytic Mukai duality

5.4.1 Statement

Let X be a complex torus of dimension g .

Theorem 5.4.1.1 (Mukai, Ben-Bassat, Block, Pantev). *There are natural isomorphisms of functors*

$$\begin{aligned} RS \circ R\hat{S} &\xrightarrow{\sim} [-1]_X^*[-g] : D_{\text{good}}(X) \rightarrow D_{\text{good}}(X); \\ R\hat{S} \circ RS &\xrightarrow{\sim} [-1]_{\hat{X}}^*[-g] : D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(\hat{X}). \end{aligned}$$

In particular, $RS : D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(X)$ is an equivalence of categories, with a quasi-inverse $[-1]_{\hat{X}}^* R\hat{S}[g]$.

Corollary 5.4.1.2. ⁵ *The functors $RS : D_c^b(\hat{X}) \rightarrow D_c^b(X)$ and $R\hat{S} : D_c^b(X) \rightarrow D_c^b(\hat{X})$ are equivalences of triangulated categories.*

Proof. It follows from Corollary 5.3.1.16 and Theorem 5.4.1.1. \square

Remark 5.4.1.3. A Mukai duality for complex tori similar to Corollary 5.4.1.2 is stated in [Blo10, p.314], with $D^b(\text{Coh}(*))$ at the place of $D_c^b(*)$. However, Prof. Jonathan Block told the author that here we should stick to $D_c^b(*)$. In fact, in general the abelian category $\text{Coh}(X)$ does not have enough injectives, so it is unclear how to define the derived direct image involved in [Blo10, p.314]. Moreover, recently Prof. Alexey Bondal announced⁶ that for a generic complex torus X of dimension > 2 , the natural functor $D^b(\text{Coh}(X)) \rightarrow D_c^b(X)$ is *not* an equivalence.

5.4.2 Proof

We follow the strategy of [BBBP07, Thm. 2.1] to prove Theorem 5.4.1.1.

Preliminaries

Lemma 5.4.2.1 (Associativity). *Let A, B be two sheaves of rings on a topological space X . For $M \in D(\text{Mod}(A))$, $N \in D(\text{BiMod}(A, B))$,⁷ and $K \in D(\text{Mod}(B))$, there is a canonical isomorphism $M \otimes_A^L (N \otimes_B^L K) = (M \otimes_A^L N) \otimes_B^L K$ in $D(\text{BiMod}(A, B))$.*

Proof. By [Sta24, Tag 06YF], there exists a quasi-isomorphism $M' \rightarrow M$ (resp. $K' \rightarrow K$) in $D(\text{Mod}(A))$ (resp. $D(\text{Mod}(B))$), where M' (resp. K') is a K-flat complex of A (resp. B) modules. From [Sta24, Tag 06YH], one has

$$\begin{aligned} M \otimes_A^L (N \otimes_B^L K) &= M' \otimes_A (N \otimes_B^L K) \\ &= M' \otimes_A (N \otimes_B K') = (M' \otimes_A N) \otimes_B K' \\ &= (M \otimes_A^L N) \otimes_B K' = (M \otimes_A^L N) \otimes_B^L K. \end{aligned}$$

\square

⁵[PPS17, Thm. 13.1] relies on Statement 5.2.0.5.

⁶<https://www.mathnet.ru/eng/present35371>

⁷Here, $\text{BiMod}(A, B)$ denotes the category of sheaves of (A, B) -bimodules.

Lemma 5.4.2.2, an analytic analog of [Muk81, Example 1.2], exhibits the derived pullback and direct image as particular examples of integral transforms.

Lemma 5.4.2.2. *Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces. Let $i : \Gamma_f \rightarrow X \times Y$ be the inclusion of the graph of f . Set $F = i_* O_{\Gamma_f} \in \text{Mod}(O_{X \times Y})$. Then there are canonical isomorphism of functors*

$$\phi_F^{[X \rightarrow Y]} \xrightarrow{\sim} Rf_* : D(X) \rightarrow D(Y); \quad (5.15)$$

$$\phi_F^{[Y \rightarrow X]} \xrightarrow{\sim} Lf^* : D(Y) \rightarrow D(X). \quad (5.16)$$

Proof. Let $g : \Gamma_f \rightarrow X$ be the projection. Since g is an isomorphism of complex analytic spaces, one has a canonical isomorphism

$$Lg^* \xrightarrow{\sim} R(g^{-1})^* \quad (5.17)$$

of functors $D(X) \rightarrow D(\Gamma_f)$. Consider the following diagram

$$\begin{array}{ccc} \Gamma_f & \xrightarrow{i} & X \times Y \\ \downarrow g & \swarrow p_X & \searrow p_Y \\ X & \xrightarrow{f} & Y. \end{array}$$

As i is a closed embedding of complex analytic spaces, by [Sta24, Tag 0B55], the natural transformation

$$Ri_* O_{\Gamma_f} \otimes^L Lp_X^*(\cdot) \rightarrow Ri_* Li^* Lp_X^*(\cdot) \quad (5.18)$$

is an isomorphism of functors $D(X) \rightarrow D(X \times Y)$. One has

$$\begin{aligned} \phi_F^{[X \rightarrow Y]} &:= R p_{Y*}(F \otimes^L p_X^* \cdot) = R p_{Y*}(Ri_* O_{\Gamma_f} \otimes^L Lp_X^* \cdot) \\ &\stackrel{(a)}{\xrightarrow{\sim}} R p_{Y*} Ri_* Li^* Lp_X^* \stackrel{(b)}{\xrightarrow{\sim}} R p_{Y*} Ri_* Lg^* \\ &\stackrel{(c)}{\xrightarrow{\sim}} R p_{Y*} Ri_* R(g^{-1})^* \stackrel{(d)}{\xrightarrow{\sim}} Rf_*, \end{aligned}$$

where (a) (resp. (c)) uses (5.18) (resp. (5.17)), and (b), (d) are from [Spa88, Thm. A (iii)].

Thus, (5.15) is proved. The proof of (5.16) is similar. \square

Proposition 5.4.2.3 is the first ingredient of the proof of Theorem 5.4.1.1, which expresses the composition of two integral transforms as another integral transform.

Proposition 5.4.2.3. *Let M, N, P be complex analytic spaces, with M, N compact and $\dim P$ finite. Let p_{ij} be the projections of the product $M \times N \times P$. For $K \in D_{\text{good}}(M \times N)$ and $L \in D(N \times P)$, set*

$$H = Rp_{13*}(p_{12}^*K \otimes^L p_{23}^*L) (\in D(M \times P)).$$

Then there is a natural isomorphism $\phi_L^{[N \rightarrow P]} \phi_K^{[M \rightarrow N]} \xrightarrow{\sim} \phi_H^{[M \rightarrow P]}$ of functors $D_{\text{good}}(M) \rightarrow D(P)$.

Proof. Let

$$\begin{aligned} a : M \times N &\rightarrow M, & b : N \times P &\rightarrow P, \\ p : M \times N &\rightarrow N, & q : N \times P &\rightarrow N, \\ u : M \times P &\rightarrow M, & v : M \times P &\rightarrow P \end{aligned}$$

be projections.

The morphism q is locally product. Properness of p follows from the compactness of M . By Propositions 5.3.1.2 2 and 5.3.1.5 2, the functor $K \otimes^L a^* \cdot : D(M) \rightarrow D(M \times N)$ restricts to a functor $D_{\text{good}}(M) \rightarrow D_{\text{good}}(M \times N)$. Then one can apply Theorem 5.3.2.3 to the cartesian square

$$\begin{array}{ccc} M \times N \times P & \xrightarrow{p_{12}} & M \times N \\ p_{23} \downarrow & \square & \downarrow p \\ N \times P & \xrightarrow{q} & N, \end{array}$$

so the base change natural transformation induces an isomorphism

$$q^*Rp_*(K \otimes^L a^* \cdot) \rightarrow Rp_{23*}p_{12}^*(K \otimes^L a^* \cdot) \quad (5.19)$$

of functors $D_{\text{good}}(M) \rightarrow D_{\text{good}}(N \times P)$. Thus, one has isomorphisms

$$\begin{aligned} \phi_L^{[N \rightarrow P]} \phi_K^{[M \rightarrow N]} &= Rb_*[L \otimes^L q^*Rp_*(K \otimes^L a^* \cdot)] \\ &\stackrel{(a)}{\xrightarrow{\sim}} Rb_*[L \otimes^L Rp_{23*}p_{12}^*(K \otimes^L a^* \cdot)] \\ &\stackrel{(b)}{\xrightarrow{\sim}} Rb_*Rp_{23*}[p_{23}^*L \otimes^L p_{12}^*(K \otimes^L a^* \cdot)] \\ &\cong Rp_{3*}[p_{23}^*L \otimes^L p_{12}^*(K \otimes^L a^* \cdot)] \\ &\cong Rv_*Rp_{13*}(p_{12}^*K \otimes^L p_{23}^*L \otimes^L p_1^* \cdot) \\ &\stackrel{(c)}{\xleftarrow{\sim}} Rv_*[H \otimes^L u^* \cdot] = \phi_H^{[M \rightarrow P]}, \end{aligned}$$

of functors $D_{\text{good}}(M) \rightarrow D(P)$ where (a) uses (5.19), and (b) (resp. (c)) is from the compactness of M (resp. N) and Fact 5.3.2.15. \square

Fact 5.4.2.4, the other ingredient of the proof of Theorem 5.4.1.1, calculates the cohomology of the Poincaré bundle.

Fact 5.4.2.4 ([Kem91, Thm. 3.15]). *Let X be a complex torus of dimension g . Let $p_X : X \times \hat{X} \rightarrow X$, $p_{\hat{X}} : X \times \hat{X} \rightarrow \hat{X}$ be the two projections. Then for the normalized Poincaré bundle \mathcal{P} , one has $Rp_{X*}\mathcal{P} = \mathbb{C}_0[-g]$ in $D^b(X)$ and $Rp_{\hat{X}*}\mathcal{P} = \mathbb{C}_0[-g]$ in $D^b(\hat{X})$.*

Proof of Theorem 5.4.1.1

By Corollary 5.3.1.16, the functor RS (resp. $R\hat{S}$) restricts to a functor $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(X)$ (resp. $D_{\text{good}}(X) \rightarrow D_{\text{good}}(\hat{X})$). Let p_{ij} be the projections of $X \times X \times \hat{X}$. Set

$$H = Rp_{12,*}(p_{13}^*\mathcal{P} \otimes^L p_{23}^*\mathcal{P}).$$

By Propositions 5.3.1.2 1 and 5.3.1.5 1, Fact 5.3.1.8 and Lemma 5.3.1.11, one has $H \in D_c^b(X \times X)$. By Proposition 5.4.2.3, one has an isomorphism of $RS \circ R\hat{S} \xrightarrow{\sim} \phi_H^{[X \rightarrow X]}$ of functors $D_{\text{good}}(X) \rightarrow D_{\text{good}}(X)$. Let $m : X \times X \rightarrow X$ be the group law. Since the $\mathcal{O}_{X \times X \times \hat{X}}$ -module $p_{13}^*\mathcal{P}$ is flat, one has $p_{13}^*\mathcal{P} \otimes^L p_{23}^*\mathcal{P} = p_{13}^*\mathcal{P} \otimes p_{23}^*\mathcal{P}$. By [BL04, Lem. 14.1.7],⁸ the $\mathcal{O}_{X \times X \times \hat{X}}$ -module $p_{13}^*\mathcal{P} \otimes p_{23}^*\mathcal{P}$ is isomorphic to $(m \times \text{Id}_{\hat{X}})^*\mathcal{P}$. Then $H \xrightarrow{\sim} Rp_{12,*}(m \times \text{Id}_{\hat{X}})^*\mathcal{P}$.

Because the morphism m is smooth, applying Theorem 5.3.2.3 to the cartesian square

$$\begin{array}{ccc} X \times X \times \hat{X} & \xrightarrow{m \times \text{Id}_{\hat{X}}} & X \times \hat{X} \\ p_{12} \downarrow & \square & \downarrow p_X \\ X \times X & \xrightarrow{m} & X \end{array}$$

in the category An , one has an isomorphism $m^*Rp_{X,*}\mathcal{P} \rightarrow H$ in $D_c^b(X \times X)$. Let $i : \Gamma_{[-1]} \rightarrow X \times X$ be the inclusion of the graph of $[-1]_X : X \rightarrow X$. From Fact 5.4.2.4, one has $H \xrightarrow{\sim} m^*\mathbb{C}_0[-g] = i_*\mathcal{O}_{\Gamma_{[-1}}}[-g]$. By Lemma 5.4.2.2, there is an isomorphism $\phi_H^{[X \rightarrow X]} \xrightarrow{\sim} [-1]_X^*[-g]$ of functors $D(X) \rightarrow D(X)$, which shows the isomorphism $RS \circ R\hat{S} \xrightarrow{\sim} [-1]_X^*[-g]$ of functors $D_{\text{good}}(X) \rightarrow D_{\text{good}}(X)$. The proof of the second isomorphism is similar.

5.5 Properties of Fourier-Mukai transform

For later reference purposes, we check that each result starting from Theorem 2.2 to (3.12') in [Muk81] has an analytic version. We only indicate the necessary modifications in statements and proofs.

⁸It is stated for abelian varieties, but its proof works for complex tori.

For a complex torus X , let g_X be its dimension. Let $(RS_X, R\hat{S}_X)$ be the Fourier-Mukai transform of X . The subscripts are omitted when there is only one complex torus in context. Let $p_X : X \times \hat{X} \rightarrow X$, $p_{\hat{X}} : X \times \hat{X} \rightarrow \hat{X}$ be the projections. For a morphism $\phi : X \rightarrow Y$ of complex tori, let $\hat{\phi} : \hat{Y} \rightarrow \hat{X}$ be the dual morphism.

5.5.1 Functoriality

Exchange of translations and twists

For every point x of the complex torus X , let $T_x : X \rightarrow X$, $x' \mapsto x' + x$ be the translation by x .

Proposition 5.5.1.1. *For every $x \in X$ and every $\hat{x} \in \hat{X}$, there are canonical isomorphisms*

$$RS \circ T_{\hat{x}}^* \cong (\cdot \otimes_{O_X} P_{-\hat{x}}) \circ RS, \quad (5.20)$$

$$RS \circ (\cdot \otimes_{O_{\hat{X}}} P_x) \cong T_x^* \circ RS \quad (5.21)$$

of functors $D(\hat{X}) \rightarrow D(X)$.

Proof. We prove (5.20). From [BL04, Cor. A.9], one gets

$$T_{(0,-\hat{x})}^* \mathcal{P} \xrightarrow{\sim} \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_X^* P_{-\hat{x}}; \quad (5.22)$$

$$T_{(x,0)}^* \mathcal{P} \xrightarrow{\sim} \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* P_x. \quad (5.23)$$

Then there are isomorphisms

$$\begin{aligned} RS(T_{\hat{x}}^* \cdot) &= Rp_{X*}(\mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* T_{\hat{x}}^* \cdot) \\ &= Rp_{X*}(\mathcal{P} \otimes_{O_{X \times \hat{X}}} T_{(0,\hat{x})}^* p_{\hat{X}}^* \cdot) \\ &= Rp_{X*} T_{(0,\hat{x})}^* (T_{(0,-\hat{x})}^* \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\ &\xrightarrow{\sim} Rp_{X*} R(T_{(0,-\hat{x})})^* (T_{(0,-\hat{x})}^* \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\ &\cong Rp_{X*} (T_{(0,-\hat{x})}^* \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\ &\stackrel{(a)}{\xrightarrow{\sim}} Rp_{X*} (p_X^* P_{-\hat{x}} \otimes \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\ &\stackrel{(b)}{\xleftarrow{\sim}} P_{-\hat{x}} \otimes Rp_{X*} (\mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\ &= P_{-\hat{x}} \otimes RS(\cdot) \end{aligned}$$

of functors $D(\hat{X}) \rightarrow D(X)$, where (a) (resp. (b)) uses (5.22) (resp. Fact 5.3.2.15).

We prove (5.21) as follows:

$$\begin{aligned}
RS(P_x \otimes \cdot) &= Rp_{X*}(\mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^*(P_x \otimes \cdot)) \\
&= Rp_{X*}(\mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* P_x \otimes p_{\hat{X}}^* \cdot) \\
&\stackrel{(a)}{\sim} Rp_{X*}(T_{(x,0)}^* \mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\
&= Rp_{X*} T_{(x,0)}^* (\mathcal{P} \otimes_{O_{X \times \hat{X}}} T_{(-x,0)}^* p_{\hat{X}}^* \cdot) \\
&\sim Rp_{X*} R(T_{(-x,0)})_* (\mathcal{P} \otimes_{O_{X \times \hat{X}}} T_{(-x,0)}^* p_{\hat{X}}^* \cdot) \\
&\cong R(T_{-x})_* Rp_{X*}(\mathcal{P} \otimes_{O_{X \times \hat{X}}} p_{\hat{X}}^* \cdot) \\
&\cong T_x^* RS(\cdot),
\end{aligned}$$

where (a) uses (5.23). □

Exchange of the direct image and the inverse image

A result similar to Proposition 5.5.1.2 is stated as [Lau96, Prop. 1.3.1]. As Laumon omits its proof, we give one. The Fourier-Mukai transform is functorial.

Proposition 5.5.1.2. *For a morphism $\phi : Y \rightarrow X$ of complex tori, there are canonical isomorphisms of functors*

$$L\phi^* \circ RS_X \cong RS_Y \circ R\phi_* : D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(Y), \quad (5.24)$$

$$R\phi_* \circ RS_Y \cong RS_X \circ L\hat{\phi}^*(\cdot)[g_X - g_Y] : D_{\text{good}}(\hat{Y}) \rightarrow D_{\text{good}}(X). \quad (5.25)$$

Proof. The isomorphism (5.25) follows from (5.24) as follows. There are isomorphisms

$$\begin{aligned}
R\phi_* RS_Y &\stackrel{(a)}{\sim} [-1]_X^* RS_X R\hat{S}_X R\phi_* RS_Y(\cdot)[g_X] \\
&\stackrel{(b)}{\sim} [-1]_X^* RS_X L\hat{\phi}^* R\hat{S}_Y RS_Y(\cdot)[g_X] \\
&\stackrel{(c)}{\sim} [-1]_X^* RS_X L\hat{\phi}^* [-1]_Y^*(\cdot)[g_X - g_Y] \\
&= RS_X L\hat{\phi}^*(\cdot)[g_X - g_Y]
\end{aligned}$$

of functors $D_{\text{good}}(\hat{Y}) \rightarrow D_{\text{good}}(X)$, where (a) and (c) use Theorem 5.4.1.1, and (b) uses (5.24).

To prove (5.24), we show

$$(\phi \times \text{Id}_{\hat{X}})^* \mathcal{P}_X \cong (\text{Id}_Y \times \hat{\phi})^* \mathcal{P}_Y. \quad (5.26)$$

Set $L := (\phi \times \text{Id}_{\hat{X}})^* \mathcal{P}_X \otimes_{O_{Y \times \hat{X}}} (\text{Id}_Y \times \hat{\phi})^* \mathcal{P}_Y^{-1}$. By definition, on the one hand for every $\hat{x} \in \hat{X}$, one has $L|_{Y \times \hat{x}} \xrightarrow{\sim} \phi^* P_{\hat{x}} \otimes P_{\hat{\phi}(\hat{x})}^{-1} \xrightarrow{\sim} O_Y$; on the other hand,

one has $L|_{0 \times \hat{X}} \xrightarrow{\sim} \hat{\phi}^* O_{\hat{Y}} \xrightarrow{\sim} O_{\hat{X}}$. By the seesaw principle [BL04, Cor. A.9], these imply $L \xrightarrow{\sim} O_{Y \times \hat{X}}$.

By applying Theorem 5.3.2.3 to the cartesian square

$$\begin{array}{ccc} Y \times \hat{X} & \xrightarrow{p_2} & \hat{X} \\ \text{Id}_Y \times \hat{\phi} \downarrow & \square & \downarrow \hat{\phi} \\ Y \times \hat{Y} & \xrightarrow{p_{\hat{Y}}} & \hat{Y} \end{array}$$

in the category An , the base change natural transformation

$$p_{\hat{Y}}^* R\hat{\phi}_* \rightarrow R(\text{Id}_Y \times \hat{\phi})_* p_2^* \quad (5.27)$$

induces an isomorphism of functors $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(Y \times \hat{Y})$. By Propositions 5.3.1.2 2 and 5.3.1.5 2, the functor $\mathcal{P}_X \otimes p_{\hat{X}}^*(\cdot) : D(\hat{X}) \rightarrow D(X \times \hat{X})$ restricts to a functor $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(X \times \hat{X})$. Because p_X is smooth proper, by applying Lemma 5.3.2.13 to the cartesian square

$$\begin{array}{ccc} Y \times \hat{X} & \xrightarrow{\phi \times \text{Id}_{\hat{X}}} & X \times \hat{X} \\ p_1 \downarrow & \square & \downarrow p_X \\ Y & \xrightarrow{\phi} & X \end{array}$$

in the category An , the base change natural transformation induces an isomorphism

$$L\phi^* R p_{X*}(\mathcal{P}_X \otimes p_{\hat{X}}^*(\cdot)) \rightarrow R p_{1*} L(\phi \times \text{Id}_{\hat{X}})^*(\mathcal{P}_X \otimes p_{\hat{X}}^*(\cdot)) \quad (5.28)$$

of functors $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(Y)$.

There are isomorphisms

$$\begin{aligned} L\phi^* \circ R S_X &= L\phi^* R p_{X*}(\mathcal{P}_X \otimes p_{\hat{X}}^*(\cdot)) \\ &\stackrel{(a)}{\xrightarrow{\sim}} R p_{1*} L(\phi \times \text{Id}_{\hat{X}})^*(\mathcal{P}_X \otimes p_{\hat{X}}^*(\cdot)) \\ &\cong R p_{1*} [L(\phi \times \text{Id}_{\hat{X}})^* \mathcal{P}_X \otimes^L L(\phi \times \text{Id}_{\hat{X}})^* p_{\hat{X}}^*(\cdot)] \\ &\cong R p_{1*} [(\phi \times \text{Id}_{\hat{X}})^* \mathcal{P}_X \otimes p_2^*(\cdot)] \\ &\stackrel{(b)}{\xrightarrow{\sim}} R p_{1*} [(\text{Id}_Y \times \hat{\phi})^* \mathcal{P}_Y \otimes p_2^*(\cdot)] \\ &\cong R p_{Y*} R(\text{Id}_Y \times \hat{\phi})_* [L(\text{Id}_Y \times \hat{\phi})^* \mathcal{P}_Y \otimes p_2^*(\cdot)] \\ &\stackrel{(c)}{\xleftarrow{\sim}} R p_{Y*} [\mathcal{P}_Y \otimes R(\text{Id}_Y \times \hat{\phi})_* p_2^*(\cdot)] \\ &\stackrel{(d)}{\xleftarrow{\sim}} R p_{Y*} [\mathcal{P}_Y \otimes p_{\hat{Y}}^* R\hat{\phi}_*(\cdot)] \\ &= R S_Y R\hat{\phi}_* \end{aligned}$$

of functors $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(Y)$, where (a) (resp. (b), resp. (c), resp. (d)) uses (5.28) (resp. (5.26), resp. Fact 5.3.2.15, resp. (5.27)). This proves (5.24). \square

Remark 5.5.1.3. In Proposition 5.5.1.2, if ϕ is an isogeny, then

$$\begin{aligned}\phi^* \circ RS_X &\cong RS_Y \circ \hat{\phi}_* : D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(Y); \\ \phi_* \circ RS_Y &\cong RS_X \circ \hat{\phi}^* : D_{\text{good}}(\hat{Y}) \rightarrow D_{\text{good}}(X).\end{aligned}$$

In fact, ϕ is finite flat and $g_Y = g_X$. By [GR04, Thm. 4, p.47], the functor $\phi_* : \text{Mod}(Y) \rightarrow \text{Mod}(X)$ is exact, so $R\phi_* = \phi_*$ as a functor $D(Y) \rightarrow D(X)$. By the flatness, the inverse image $\phi^* : \text{Mod}(X) \rightarrow \text{Mod}(Y)$ is exact and $L\phi^* = \phi^*$ as a functor $D(X) \rightarrow D(Y)$.

In [Muk81, (3.4)], for an isogeny $\phi : Y \rightarrow X$ of abelian varieties, the derived functor $R\phi_* : D_{\text{qc}}(Y) \rightarrow D_{\text{qc}}(X)$ is also written as ϕ_* , but for a different reason [Sta24, Tag 08D7].

For the first half of [Muk81, Prop. 3.11 (4)], the result [MRM74, Sec. 23, Lem. 3] cited in its proof still holds for complex tori, with a similar (and simpler) proof.

Exchange of the Pontrjagin product and the tensor product

Let p_i be the two projections $X \times X \rightarrow X$. Define a bifunctor $*^R : D(X) \times D(X) \rightarrow D(X)$ by $- *^R + = Rm_*(p_1^* - \otimes^L p_2^* +)$. As in Corollary 5.3.1.16, the bifunctor $*^R$ restricts to a bifunctor $D_{\text{good}}(X) \times D_{\text{good}}(X) \rightarrow D_{\text{good}}(X)$ (resp. $D_c^b(X) \times D_c^b(X) \rightarrow D_c^b(X)$).

Fact 5.5.1.4 ([Muk81, (3.7)]). *For every $F \in D_{\text{good}}(\hat{X})$, there are canonical isomorphisms*

$$\begin{aligned}RS(F *^R \cdot) &\cong RS(F) \otimes^L RS(\cdot), \\ RS(F \otimes^L \cdot) &\cong RS(F) *^R RS(\cdot)[g]\end{aligned}$$

of functors $D_{\text{good}}(\hat{X}) \rightarrow D_{\text{good}}(X)$.

Commutativity with external tensor product

Let M, N be two complex analytic spaces. Let $p : M \times N \rightarrow M$ and $q : M \times N \rightarrow N$ be the projections. The bifunctor $D(M) \times D(N) \rightarrow D(M \times N)$, $(-, +) \mapsto (p^* -) \otimes^L (q^* +)$ is denoted by $(\cdot) \boxtimes^L (\cdot)$.

Proposition 5.5.1.5. *Let X, Y be two complex tori and $Z = X \times Y$. Then there is a canonical isomorphism $RS_Z(- \boxtimes^L +) = RS_X(-) \boxtimes^L RS_Y(+)$ of bifunctors $D_{\text{good}}(\hat{X}) \times D_{\text{good}}(\hat{Y}) \rightarrow D_{\text{good}}(Z)$.*

Proof. By the seesaw principle, one has $\mathcal{P}_Z \xrightarrow{\sim} \mathcal{P}_X \boxtimes^L \mathcal{P}_Y$. Then there are canonical isomorphisms

$$\begin{aligned}
RS_Z(- \boxtimes^L +) &= Rp_{Z*}[\mathcal{P}_Z \otimes^L Lp_Z^*(- \boxtimes^L +)] \\
&\xrightarrow{\sim} Rp_{Z*}[(\mathcal{P}_X \boxtimes^L \mathcal{P}_Y) \otimes^L (Lp_X^*(-) \boxtimes^L Lp_Y^*(+))] \\
&\xrightarrow{\sim} R(p_X \times p_Y)_*[(\mathcal{P}_X \otimes^L Lp_X^*(-)) \boxtimes^L (\mathcal{P}_Y \otimes^L Lp_Y^*(+))] \\
&\stackrel{(a)}{\xrightarrow{\sim}} Rp_{X*}(\mathcal{P}_X \otimes^L Lp_X^*(-)) \boxtimes^L Rp_{Y*}(\mathcal{P}_Y \otimes^L Lp_Y^*(+)) \\
&= RS_X(-) \boxtimes^L RS_Y(+)
\end{aligned}$$

of bifunctors $D_{\text{good}}(\hat{X}) \times D_{\text{good}}(\hat{Y}) \rightarrow D_{\text{good}}(Z)$, where (a) uses Lemma 5.5.1.6 2. \square

Lemma 5.5.1.6.

1. Let X, Y, T be complex analytic spaces, with X, T finite dimensional. Let $f : X \rightarrow Y$ be a proper morphism. Then there is a canonical isomorphism

$$Rf_*(-) \boxtimes^L (+) \rightarrow R(f \times \text{Id}_T)_*(- \boxtimes^L +)$$

of bifunctors $D_{\text{good}}(X) \times D(T) \rightarrow D(Y \times T)$.

2. Let $f_i : X_i \rightarrow Y_i$ ($i = 1, 2$) be proper morphism of complex analytic spaces. If X_1, X_2 and Y_1 are finite dimensional, then there is a canonical isomorphism

$$(Rf_{1*}-) \boxtimes^L (Rf_{2*}+) \rightarrow R(f_1 \times f_2)_*(- \boxtimes^L +)$$

of bifunctors $D_{\text{good}}(X_1) \times D_{\text{good}}(X_2) \rightarrow D_{\text{good}}(Y_1 \times Y_2)$.

Proof.

1. Consider the notation in the commutative diagram

$$\begin{array}{ccccc}
& & X \times T & \xrightarrow{u} & X \\
& \swarrow v & \downarrow f \times \text{Id}_T & \square & \downarrow f \\
T & \xleftarrow{q} & Y \times T & \xrightarrow{p} & Y,
\end{array}$$

where u, v, p and q are projections. Since $v = q \circ (f \times \text{Id}_T)$, there is a canonical isomorphism $v^* \xrightarrow{\sim} L(f \times \text{Id}_T)^* q^*$ of functors $D(T) \rightarrow D(X \times T)$. As $f \times \text{Id}_T$ is a base change of f , it is also proper. As $\dim(X \times T)$ is finite, by Fact 5.3.2.15, the canonical morphism

$$[R(f \times \text{Id}_T)_* u^* -] \otimes^L q^* + \rightarrow R(f \times \text{Id}_T)_*[u^* - \otimes^L v^* +] \quad (5.29)$$

of bifunctors $D(X) \times D(T) \rightarrow D(Y \times T)$ is an isomorphism.

By Theorem 5.3.2.3, one has a canonical isomorphism

$$p^* Rf_* \rightarrow R(f \times \text{Id}_T)_* u^* : D_{\text{good}}(X) \rightarrow D_{\text{good}}(Y \times T). \quad (5.30)$$

Therefore, there are canonical isomorphisms

$$\begin{aligned} (Rf_* -) \boxtimes^L + &= (p^* Rf_* -) \otimes^L q^* + \\ &\stackrel{(a)}{\sim} [R(f \times \text{Id}_T)_* u^* -] \otimes^L q^* + \\ &\stackrel{(b)}{\sim} R(f \times \text{Id}_T)_* [u^* - \otimes v^* +] \\ &= R(f \times \text{Id}_T)_* (- \boxtimes^L +), \end{aligned}$$

of bifunctors $D_{\text{good}}(X) \times D(T) \rightarrow D(Y \times T)$, where (a) (resp. (b)) uses (5.30) (resp. (5.29)).

2. Since $\dim(X_1 \times X_2)$ is finite, as in Corollary 5.3.1.16, the bifunctor $R(f_1 \times f_2)_* (- \boxtimes^L +)$ restricts to a bifunctor $D_{\text{good}}(X_1) \times D_{\text{good}}(X_2) \rightarrow D_{\text{good}}(Y_1 \times Y_2)$.

As $\dim Y_1, \dim X_2$ are finite, by Point 1, there are canonical isomorphisms of bifunctors

$$\begin{aligned} (Rf_{1*} -) \boxtimes^L + &\rightarrow R(f_1 \times \text{Id}_{X_2})_* (- \boxtimes^L +) : D_{\text{good}}(X_1) \times D(X_2) \rightarrow D(Y_1 \times X_2), \\ (Rf_{1*} -) \boxtimes^L (Rf_{2*} +) &\rightarrow R(\text{Id}_{Y_1} \times f_2)_* [(Rf_{1*} -) \boxtimes^L +] : D(X_1) \times D_{\text{good}}(X_2) \rightarrow D(Y_1 \times Y_2). \end{aligned}$$

Then there is a canonical isomorphism of bifunctors

$$\begin{aligned} (Rf_{1*} -) \boxtimes^L (Rf_{2*} +) &\rightarrow R(\text{Id}_{Y_1} \times f_2)_* [(Rf_{1*} -) \boxtimes^L +] \\ &\rightarrow R(\text{Id}_{Y_1} \times f_2)_* R(f_1 \times \text{Id}_{X_2})_* (- \boxtimes^L +) \\ &\rightarrow R(f_1 \times f_2)_* (- \boxtimes^L +) : D_{\text{good}}(X_1) \times D_{\text{good}}(X_2) \rightarrow D_{\text{good}}(Y_1 \times Y_2). \end{aligned}$$

□

Skew commutativity with duality

We summarize classical facts about the duality theory on complex manifolds.

Fact 5.5.1.7. *Let X be a complex manifold of pure dimension n , and let $\omega_X = \bigwedge^n \Omega_X$ be the canonical line bundle.*

1. ([RR70, p.81; p.90]) *The dualizing functor $D_X = R\mathcal{H}om_X(\cdot, \omega_X)[n] : D(X) \rightarrow D(X)$ restricts to a functor $D_c(X) \rightarrow D_c(X)$ and the natural transformation $\text{Id} \rightarrow D_X \circ D_X : D_c(X) \rightarrow D_c(X)$ is an isomorphism.*

If X is compact, then D_X exchanges⁹ $D_c^+(X)$ with $D_c^-(X)$, and induces an equivalence $D_c^b(X) \rightarrow D_c^b(X)$.

2. ([RRV71, p.264]) There is a canonical isomorphism $R\mathcal{H}om_X(-, +) \rightarrow D_X(- \otimes^L D_X +)$ of bifunctors $D_c(X) \times D_c^+(X) \rightarrow D(X)$.
3. ([RRV71, p.264], [Bjö93, p.122]) Let $f : X \rightarrow Y$ be a proper morphism of complex manifolds. Then there is a canonical isomorphism of functors $Rf_* D_X \rightarrow D_Y Rf_* : D_c(X) \rightarrow D(Y)$.

Proposition 5.5.1.8 ([Muk81, (3.8)]). *There are canonical isomorphisms of functors*

$$\begin{aligned} D_X \circ RS &\xrightarrow{\sim} ([-1]_X^* \circ RS \circ D_{\hat{X}})[g] : D_c^+(\hat{X}) \rightarrow D_c^-(X); \\ D_{\hat{X}} \circ R\hat{S} &\xrightarrow{\sim} ([-1]_{\hat{X}}^* \circ R\hat{S} \circ D_X)[g] : D_c^+(X) \rightarrow D_c^-(\hat{X}). \end{aligned}$$

We make some preparation for the proof of Proposition 5.5.1.8. Lemma 5.5.1.9 is an adaption of [Har66, Ch.II, Prop. 5.8] and [Sta24, Tag 0C6I].

Lemma 5.5.1.9. *Let $f : X \rightarrow Y$ be a flat morphism of complex analytic spaces. Then:*

1. *There is a canonical natural transformation of bifunctors*

$$f^* R\mathcal{H}om_Y(-, +) \rightarrow R\mathcal{H}om_X(f^* -, f^* +) : D(Y) \times D(Y) \rightarrow D(X). \quad (5.31)$$

2. *The natural transformation (5.31) restricts to an isomorphism of bifunctors $D_c^-(Y) \times D(Y) \rightarrow D(X)$.*

Proof. Set $G \in D(Y)$.

1. By [Spa88, Thm. D], there is a *functorial* quasi-isomorphism $G \rightarrow G'$, where G' is a K-injective complex over $\text{Mod}(O_Y)$. There are natural transformations of functors $D(Y) \rightarrow D(X)$

$$\begin{aligned} f^* R\mathcal{H}om_Y(\cdot, G) &\rightarrow f^* \mathcal{H}om_Y(\cdot, G') \rightarrow \mathcal{H}om_X(f^* \cdot, f^* G') \\ &\rightarrow R\mathcal{H}om_X(f^* \cdot, f^* G') \xleftarrow{\sim} R\mathcal{H}om_X(f^* \cdot, f^* G). \end{aligned}$$

2. By [Har66, I, Examples 1], the (contravariant) functors

$$f^* R\mathcal{H}om_Y(\cdot, G), R\mathcal{H}om_X(f^* \cdot, f^* G) : D(Y) \rightarrow D(X)$$

are bounded below. Consider $F \in D_c^-(Y)$. To show the natural morphism $f^* R\mathcal{H}om_Y(F, G) \rightarrow R\mathcal{H}om_X(f^* F, f^* G) : D_c^-(Y) \rightarrow D(X)$

⁹By [FS13, p.4971], in general the functor $R\mathcal{H}om_X(\cdot, \omega_X) : D(X) \rightarrow D(X)$ does not exchange $D_c^{b, \leq 0}(X)$ and $D_c^{b, \geq 0}(X)$.

is an isomorphism, by [Har66, I, Prop. 7.1 (ii)], one may assume $F \in \text{Coh}(Y)$. By [Sta24, Tag 08DL], one may shrink Y to open subsets. Thus, from Lemma A.1.3.1, one may assume that there is a quasi-isomorphism $K \rightarrow F$, where K is a complex of finite free \mathcal{O}_Y -modules. The morphism f is flat, so $f^*K \rightarrow f^*F \rightarrow 0$ is a globally free resolution of f^*F . The morphism (5.31) is identified with $f^*\mathcal{H}om_Y(K, G) \rightarrow \mathcal{H}om_X(f^*K, f^*G)$, which is an isomorphism. \square

Lemma 5.5.1.10. *Let $E \rightarrow X$ be a holomorphic vector bundle on a complex manifold, and let E^\vee be the dual vector bundle. Then there is an isomorphism of functors $E^\vee \otimes D_X \cdot \rightarrow D_X(E \otimes \cdot) : D(X) \rightarrow D(X)$.*

Proof. Since E is a vector bundle, one has isomorphisms

$$E \otimes \cdot \xrightarrow{\sim} \mathcal{H}om_X(E^\vee, \cdot) \xrightarrow{\sim} R\mathcal{H}om_X(E^\vee, \cdot)$$

of functors $D(X) \rightarrow D(X)$. Then

$$D_X(E \otimes \cdot) = R\mathcal{H}om_X(R\mathcal{H}om_X(E^\vee, \cdot), \omega_X)[\dim X].$$

As E^\vee is a perfect object of $D(X)$ (in the sense of [Sta24, Tag 08CM]), by [Sta24, Tag 0G40], one has $D_X(E \otimes \cdot) = R\mathcal{H}om_X(\cdot, \omega_X)[\dim X] \otimes^L E^\vee = E^\vee \otimes D_X \cdot$. \square

Corollary 5.5.1.11. *Let $f : X \rightarrow Y$ be a flat morphism of complex manifolds of relative dimension n . Write $\omega_f = \omega_X \otimes_{\mathcal{O}_X} f^*\omega_Y^\vee$ for the relative dualizing line bundle. Then there is a canonical isomorphism of functors $D_X f^* D_Y \rightarrow \omega_f \otimes_{\mathcal{O}_X} f^*(\cdot)[n] : D_c^-(Y) \rightarrow D_c^-(X)$.*

Proof. One has

$$\begin{aligned} D_X f^* D_Y \mathcal{O}_Y &= D_X(f^* R\mathcal{H}om_Y(\mathcal{O}_Y, \omega_Y)[\dim Y]) = D_X(f^* \omega_Y[\dim Y]) \\ &= R\mathcal{H}om_X(f^* \omega_Y, \omega_X)[\dim X - \dim Y] \stackrel{(a)}{=} \mathcal{H}om_X(f^* \omega_Y, \omega_X)[n] \\ &= f^* \omega_Y^\vee \otimes_{\mathcal{O}_X} \omega_X[n] = \omega_f[n], \end{aligned} \tag{5.32}$$

where (a) uses that $f^* \omega_Y$ is a line bundle on X .

By Fact 5.5.1.7 1 and 2, there is an isomorphism $D_Y \xrightarrow{\sim} R\mathcal{H}om_Y(\cdot, D_Y \mathcal{O}_Y)$ of functors $D_c^-(Y) \rightarrow D_c^+(Y)$. From Lemma 5.5.1.9 2, there are isomorphisms

$$f^* D_Y \xrightarrow{\sim} f^* R\mathcal{H}om_Y(\cdot, D_Y \mathcal{O}_Y) \xrightarrow{\sim} R\mathcal{H}om_X(f^* \cdot, f^* D_Y \mathcal{O}_Y)$$

of functors $D_c^-(Y) \rightarrow D_c^+(X)$. Then by Fact 5.5.1.7 1 and 2 again, there are isomorphisms

$$\begin{aligned} D_X f^* D_Y &\xrightarrow{\sim} f^*(\cdot) \otimes^L D_X f^* D_Y \mathcal{O}_Y \\ &\stackrel{(a)}{=} f^*(\cdot) \otimes_{\mathcal{O}_X}^L \omega_f[n] \stackrel{(b)}{=} f^*(\cdot) \otimes_{\mathcal{O}_X} \omega_f[n] \end{aligned}$$

of functors $D_c^-(Y) \rightarrow D_c^-(X)$, where (a) (resp. (b)) equality uses (5.32) (resp. local freeness of ω_f). \square

Lemma 5.5.1.12. *There is an isomorphism $Rp_{X*}(\mathcal{P}^{-1} \otimes^L p_{\hat{X}}^* \cdot) = [-1]_X^* RS$ of functors $D(\hat{X}) \rightarrow D(X)$.*

Proof. By [BL04, Cor. A.9], one has $\mathcal{P}^{-1} \xrightarrow{\sim} ([-1]_X \times [1]_{\hat{X}})^* \mathcal{P}$. Since $p_{\hat{X}} \circ ([-1]_X \times [1]_{\hat{X}}) = p_{\hat{X}}$, there are isomorphisms

$$\begin{aligned} Rp_{X*}(\mathcal{P}^{-1} \otimes^L p_{\hat{X}}^* \cdot) &\xrightarrow{\sim} Rp_{X*}([-1]_X \times [1]_{\hat{X}})^*(\mathcal{P} \otimes^L p_{\hat{X}}^* \cdot) \\ &\xleftarrow{\sim} [-1]_X^* Rp_{X*}(\mathcal{P} \otimes^L p_{\hat{X}}^* \cdot) = [-1]_X^* RS \end{aligned}$$

of functors $D(\hat{X}) \rightarrow D(X)$. \square

Proof of Proposition 5.5.1.8. By Fact 5.5.1.7 1 and 3, There are isomorphisms

$$D_X \circ RS = D_X Rp_{X*}(\mathcal{P} \otimes^L p_{\hat{X}}^* \cdot) \xrightarrow{\sim} Rp_{X*} D_{X \times \hat{X}}(\mathcal{P} \otimes^L p_{\hat{X}}^* \cdot)$$

of functors $D_c^+(\hat{X}) \rightarrow D_c^-(X)$. From Lemma 5.5.1.10, there is an isomorphism $D_{X \times \hat{X}}(\mathcal{P} \otimes^L p_{\hat{X}}^* \cdot) \xrightarrow{\sim} \mathcal{P}^{-1} \otimes^L D_{X \times \hat{X}} p_{\hat{X}}^* \cdot$ of functors $D(\hat{X}) \rightarrow D(X \times \hat{X})$. By Fact 5.5.1.7 1, the functor $D_{\hat{X}}$ restricts to a functor $D_c^+(\hat{X}) \rightarrow D_c^-(\hat{X})$, whence Corollary 5.5.1.11 yields an isomorphism $D_{X \times \hat{X}} p_{\hat{X}}^* = (p_{\hat{X}}^* D_{\hat{X}} \cdot)[g]$ of functors $D_c^+(\hat{X}) \rightarrow D_c^-(X \times \hat{X})$. Therefore, there are isomorphisms

$$D_X \circ RS \xrightarrow{\sim} Rp_{X*}(\mathcal{P}^{-1} \otimes^L p_{\hat{X}}^* D_{\hat{X}} \cdot)[g] \xrightarrow{(a)} [-1]_X^* RS(D_{\hat{X}} \cdot)[g]$$

of functors $D_c^+(\hat{X}) \rightarrow D_c^-(X)$, where (a) uses Lemma 5.5.1.12.

The second isomorphism follows from the first by swapping X and \hat{X} . \square

5.5.2 Unipotent vector bundles

Definition 5.5.2.1 ([Muk81, Def. 2.3]). We say that W.I.T. (weak index theorem) holds for a coherent module F on the complex torus X if there is an integer $i(F)$ such that $H^i R\hat{S}(F) = 0$ for every integer $i \neq i(F)$. In that case, the integer $i(F)$ is called the *index* of F and the coherent module $\hat{F} := H^{i(F)} R\hat{S}(F)$ on \hat{X} is called the Fourier transform of F . We say that I.T. (index theorem) holds for F if there is an integer i_0 such that for every $L \in \text{Pic}^0(X)$ and every integer $i \neq i_0$, one has $H^i(X, F \otimes_{O_X} L) = 0$.

Fact 5.5.2.2 ([Nak94, p.80]). *Let F be a coherent O_X -module, then I.T. holds for F if and only if W.I.T holds for F and \hat{F} is locally free on \hat{X} .*

Example 5.5.2.4 show that that the word ‘‘Artinian’’ in Statement 5.5.2.3 is a typo. It should be ‘‘finite length’’ ([Muk78, Thm. 4.12 (1)]).

Statement 5.5.2.3 ([Muk81, Eg. 2.9]). Let X be an abelian variety. Let $\text{Mod}^{\text{Ar}}(O_{\hat{X},0}) \subset \text{Mod}(O_{\hat{X},0})$ be the full subcategory comprised of Artinian $O_{\hat{X},0}$ -modules. Then the functor $\text{Mod}(O_X) \rightarrow \text{Mod}(O_{\hat{X},0})$ taking the stalk at 0 restricts to an equivalence $\text{Coh}_0(\hat{X}) \rightarrow \text{Mod}^{\text{Ar}}(O_{\hat{X},0})$ of categories.

Example 5.5.2.4. When $\dim X = 1$, the ring $O_{\hat{X},0}$ is a discrete valuation ring (DVR). Let $\mathbb{C}(\hat{X})$ be the fraction field of $O_{\hat{X},0}$ (or equivalently, the field of rational functions on \hat{X}). By Lemma 5.5.2.5 2, the $O_{\hat{X},0}$ -module $\mathbb{C}(\hat{X})/O_{\hat{X},0}$ is Artinian but not finitely generated, so cannot be the stalk at $0 \in \hat{X}$ of any coherent $O_{\hat{X}}$ -module.

Lemma 5.5.2.5. Let R be a DVR with a uniformizer π and fraction field K , then:

1. For every nonzero proper R -submodule $M \subsetneq K$, there is an integer n such that $M = \pi^n R$.
2. The R -module K/R is Artinian but not finitely generated.

Definition 5.5.2.6. A vector bundle U on a complex analytic space M is called unipotent if it has a filtration by vector subbundles

$$0 = U_0 \subset U_1 \subset \cdots \subset U_{n-1} \subset U_n = U$$

such that $U_i/U_{i-1} \cong O_M$ for all $1 \leq i \leq n$. Denote the full subcategory of $\text{Coh}(M)$ consisting of unipotent vector bundles by $\text{Uni}(M)$.

By [FL14, Lem. 5.1], every unipotent vector bundle on a complex torus admits a flat holomorphic connection whose underlying local system is unipotent.

Proposition 5.5.2.7. 1. W.I.T. with index g holds for every unipotent vector bundle on X .

2. The functor $H^g R\hat{S} : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{\hat{X}})$ restricts to an equivalence $\text{Uni}(X) \rightarrow \text{Coh}_0(\hat{X})$, with a quasi-inverse $H^0 R\hat{S} = R\hat{S} : \text{Coh}_0(\hat{X}) \rightarrow \text{Uni}(X)$.
3. For every unipotent vector bundle $U \rightarrow X$ and every integer $i \geq 0$, one has $H^i(X, U) = \text{Ext}_{O_{\hat{X},0}}^i(\mathbb{C}, \hat{U})$.

Proof. 1. Because $R\hat{S}$ is a triangulated functor, the full subcategory of $\text{Coh}(X)$ comprised of modules satisfying W.I.T. of a fixed index is closed under extensions. By Lemma 5.2.0.8 and Theorem 5.4.1.1, one has $R\hat{S}(O_X) = R\hat{S}RS(\mathbb{C}_0) \xrightarrow{\sim} \mathbb{C}_0[-g]$. Then W.I.T. with index g holds for O_X , so it holds for every unipotent vector bundle on X .

2. By Point 1, one has an isomorphism of functors $H^g R\hat{S} \xrightarrow{\sim} R\hat{S}[g] : \text{Uni}(X) \rightarrow \text{Mod}(O_{\hat{X}})$. The full subcategory of $\text{Mod}(O_X)$ comprised of modules F with $\text{Supp}(H^g R\hat{S}(F)) \subset \{0\}$ is closed under extensions and contains O_X , so it contains Uni_X . Since $\text{Uni}(X) \subset \text{Coh}(X)$, the functor $H^g R\hat{S} : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{\hat{X}})$ restricts to a functor $\text{Uni}(X) \rightarrow \text{Coh}_0(\hat{X})$.

For every $F \in \text{Coh}_0(\hat{X})$, the restriction $\text{Supp}(p_X^* F \otimes \mathcal{P}) \rightarrow X$ of p_X is finite. By [GR04, Thm. 4, p.47], one has $RS(F) = H^0 RS(F)$. By Lemma 5.5.2.8 3, the $O_{\hat{X}}$ -module F has a filtration with successive quotients isomorphic to \mathbb{C}_0 . Then $RS(F)$ has a filtration with successive quotients isomorphic to $RS(\mathbb{C}_0) = O_X$. By [Gro60, Ch. 0, 5.4.9], every term of this filtration is finite locally free. Therefore, $RS(F) \in \text{Uni}(X)$ and RS restricts to a functor $\text{Coh}_0(\hat{X}) \rightarrow \text{Uni}(X)$. By Theorem 5.4.1.1, the functor $H^g R\hat{S} : \text{Uni}(X) \rightarrow \text{Coh}_0(\hat{X})$ is an equivalence with a quasi-inverse RS .

3. It follows from [Muk81, Prop. 2.7] and Point 1. □

For a commutative ring R , let $\text{Mod}_f(R) \subset \text{Mod}(R)$ be the full subcategory comprised of R -modules of finite length. Lemma 5.5.2.8 1 confirms a guess in [Gro61a, 9-12] for complex field.

Lemma 5.5.2.8. *Let X be a complex analytic space. Let $x \in X$.*

1. *The functor $i_x^{-1} : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{X,x})$ taking the stalk at x restricts to a functor $\text{Coh}_x(X) \rightarrow \text{Mod}_f(O_{X,x})$. In particular, if X is a singleton, then $\dim_{\mathbb{C}} O_X$ is finite.*
2. *The functor $i_{x,*} : D(O_{X,x}) \rightarrow D(O_X)$ restricts to a functor $\text{Mod}_f(O_{X,x}) \rightarrow \text{Coh}_x(X)$.*
3. *The functor $i_x^{-1} : \text{Coh}_x(X) \rightarrow \text{Mod}_f(O_{X,x})$ is an equivalence.*

Proof. 1. For every $F \in \text{Coh}_x(X)$, to prove that F_x is a finite length $O_{X,x}$ -module, one may assume that $F_x \neq 0$. As F is a finite type O_X -module, F_x is a finite $O_{X,x}$ -module. Then $\text{Supp}_{O_{X,x}}(F_x)$ is nonempty. Let m_x be the maximal ideal of $O_{X,x}$. For every $f \in m_x$, there is an open neighborhood U of $x \in X$ such that f is the stalk of some $\bar{f} \in O_X(U)$. Then \bar{f} vanishes on $\text{Supp}(F)$. By the Rückert Nullstellensatz (see, e.g., [GR84, p.67]), there is an integer $n \geq 1$ such that $\bar{f}^n F = 0$ near x . In particular, $f \in \sqrt{\text{Ann}_{O_{X,x}}(F_x)}$. Therefore,

$$m_x \subset \sqrt{\text{Ann}_{O_{X,x}}(F_x)}.$$

By [GR84, Corollary, p.44], the ideal m_x is finitely generated, so there is an integer $N \geq 1$ with $m_x^N \subset \text{Ann}_{O_{X,x}}(F_x)$. By [Sta24, Tag 00L6], $\text{Supp}_{O_{X,x}}(F_x)$ is the unique closed point of $\text{Spec}(O_{X,x})$. By [Sta24, Tag 00L5], the $O_{X,x}$ -module F_x has finite length. The second statement follows from Lemma 5.5.2.9.

2. Up to isomorphism, the only simple $O_{X,x}$ -module is the residue field \mathbb{C} . Every $M \in \text{Mod}_f(O_{X,x})$ has a composite series with successive quotients isomorphic to \mathbb{C} . Thus, M_x has a filtration with successive quotients isomorphic to \mathbb{C}_x . Since \mathbb{C}_x is coherent, by [Sta24, Tag 01BY (4)], M_x is coherent. Therefore, $i_{x,*}$ restricts to a functor $\text{Mod}_f(O_{X,x}) \rightarrow \text{Coh}_x(X)$.
3. Let $i_x : (x, O_{X,x}) \rightarrow (X, O_X)$ be the canonical morphism of locally ringed spaces. There is a canonical isomorphism $i_x^*(i_x)_* \xrightarrow{\sim} \text{Id}_{\text{Mod}(O_{X,x})}$ of functors $\text{Mod}(O_{X,x}) \rightarrow \text{Mod}(O_{X,x})$. By adjunction, $(i_x)_* : \text{Mod}(O_{X,x}) \rightarrow \text{Mod}(O_X)$ is fully faithful. By Point 2, pushout $(i_x)_*$ restricts to a functor $\text{Mod}_f(O_{X,x}) \rightarrow \text{Coh}_x(O_X)$. For every object F of $\text{Coh}_x(O_X)$, by Point 1, F_x is an object of $\text{Mod}_f(O_{X,x})$. The adjunction morphism $F \rightarrow (i_x)_*(F_x)$ is an isomorphism. Thus, $(i_x)_* : \text{Mod}_f(O_{X,x}) \rightarrow \text{Coh}_x(O_X)$ is essentially surjective and hence an equivalence. Therefore, the functor $i_x^* : \text{Coh}_x(O_X) \rightarrow \text{Mod}_f(O_{X,x})$ (taking the stalk at x) is an equivalence. □

Lemma 5.5.2.9. *Let $F \rightarrow A$ be a ring map, with F a field and (A, m) an Artinian local ring. If $\dim_F A/m$ is finite, then $\dim_F A$ is finite.*

Proof. Because A is an Artinian local ring, by [Ati69, Prop. 8.4], there is an integer $n > 0$ with $m^n = 0$. For every integer $i \geq 0$, the A -module m^i is finitely generated, so the A/m -module m^i/m^{i+1} is finitely generated. Thus, $\dim_F m^i/m^{i+1} = \dim_F A/m \cdot \dim_{A/m} m^i/m^{i+1}$ is finite. Then $\dim_F A = \sum_{i=0}^n \dim_F m^i/m^{i+1}$ is finite. □

5.5.3 Homogeneous vector bundles

Definition 5.5.3.1. A vector bundle E on the complex torus X is *homogeneous* if for every $x \in X$, one has $T_x^*E \cong E$. Let $H(X) \subset \text{Coh}(X)$ be the full subcategory comprised of homogeneous vector bundles.

For a complex analytic space M , let $\text{Coh}_f(M) \subset \text{Coh}(M)$ be the full subcategory consisting of objects with finite support.

Proposition 5.5.3.2. 1. *For every integer i , the functor $H^i R\hat{S} : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{\hat{X}})$ restricts to a functor $H(X) \rightarrow \text{Coh}_f(\hat{X})$.*

2. *W.I.T. holds for every homogeneous vector bundle on X with index g .*

3. The functor $H^g R\hat{S} : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{\hat{X}})$ restricts to an equivalence of categories $H(X) \rightarrow \text{Coh}_f(\hat{X})$, with a quasi-inverse $H^0 RS$.

Proof. 1. Let E be a homogeneous vector bundle on X . By Corollary 5.3.1.16, the $O_{\hat{X}}$ -module $H^i R\hat{S}(E)$ is coherent. For every $x \in X$, by Proposition 5.5.1.1, one has $R\hat{S}(E) \xrightarrow{\sim} R\hat{S}(T_{-x}^* E) \xrightarrow{\sim} P_x^* \otimes R\hat{S}(E)$, so $H^i R\hat{S}(E) \xrightarrow{\sim} P_x^* \otimes H^i R\hat{S}(E)$. From Lemma 5.5.3.4, the support of $H^i R\hat{S}(E)$ is finite.

2. For every integer $i \neq g$, by Point 1, one has $H^i R\hat{S}(E) \in \text{Coh}_f(\hat{X})$ and

$$\begin{aligned} 0 &= H^{i-g}([-1]_X^* E) \\ &= H^i([-1]_X^* E[-g]) \\ &\stackrel{(a)}{\xrightarrow{\sim}} H^i RS \circ R\hat{S}(E) \\ &= H^i R p_{X*}(\mathcal{P} \otimes^L p_{\hat{X}}^* R\hat{S}(E)) \\ &\stackrel{(b)}{\xrightarrow{\sim}} H^0 R p_{X*}(\mathcal{P} \otimes^L p_{\hat{X}}^* H^i R\hat{S}(E)) \\ &= H^0 RS(H^i R\hat{S}(E)), \end{aligned}$$

where (a) (resp. (b)) uses Theorem 5.4.1.1 (resp. [GR04, Thm. 4, p.47]).

It remains to prove that for every $F \in \text{Coh}_f(\hat{X})$ with $H^0 RS(F) = 0$, one has $F = 0$. Since F is the direct sum of finitely many coherent submodules whose supports are singletons, one may assume that $\text{Supp}(F)$ is a singleton. By Proposition 5.5.1.1, one may assume that $F \in \text{Coh}_0(\hat{X})$. From Proposition 5.5.2.7 2, one has $F = 0$.

3. By Point 1, the functor $H^g R\hat{S} : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{\hat{X}})$ restricts to a functor $H(X) \rightarrow \text{Coh}_f(\hat{X})$. From Point 2, one has an isomorphism of functors $H^g R\hat{S} \cong R\hat{S}[g] : H(X) \rightarrow \text{Coh}_f(\hat{X})$.

By Propositions 5.5.1.1 and 5.5.2.7, the functor $H^0 RS : \text{Mod}(O_{\hat{X}}) \rightarrow \text{Mod}(O_X)$ restricts to a functor $H^0 RS = RS : \text{Coh}_f(\hat{X}) \rightarrow H(X)$. By Theorem 5.4.1.1, the functor $H^g R\hat{S} : H(X) \rightarrow \text{Coh}_f(\hat{X})$ is an equivalence with a quasi-inverse $H^0 RS$.

□

For a sheaf of module F on a complex analytic space, denote the torsion part of F (in the sense of [CD94, p.60]) by $T(F)$.

Lemma 5.5.3.3. *Let X be a compact Kähler manifold. Let F be a coherent O_X -module. Then for every irreducible component $C \subset \text{Supp}(F)$, there is a connected compact Kähler manifold Z and a morphism $h : Z \rightarrow X$, such that $h(Z) = C$ and $h^* F / T(h^* F)$ is a vector bundle on Z of positive rank.*

Proof. By [GR84, p.76], $\text{Supp}(F)$ is an analytic subset of X . Because X is a Kähler manifold, with the induced reduced complex structure, the subspace C is a Kähler space in the sense of [Var89, II, 1.3]. Let $i : C \rightarrow X$ be the inclusion. Set

$$D = \{x \in C : i^*F \text{ is not locally free at } x\}.$$

From [Ros68, Prop. 3.1], D is a strict analytic subset of C . By Rossi's theorem (see, e.g. [Rie71, Thm. 2]), there is a reduced irreducible complex analytic space W and a proper modification $f : W \rightarrow C$, such that $W \setminus f^{-1}(D) \rightarrow C \setminus D$ is biholomorphic and $E := N/T(N)$ is a vector bundle on W , where $N = f^*i^*F$. From [GD71, Cor. 5.2.4.1], one has $\text{Supp}(N) = W$. From [CD94, I, Thm. 9.12], one gets $\text{Supp}(T(N)) \neq W$. Therefore, the rank r of the vector bundle E is positive.

Since $f : W \rightarrow C$ is bimeromorphic, the space W is in the Fujiki class \mathcal{C} (defined in [Fuj78, p.34]). By [Fuj78, Lem. 4.6, 1)], there is a connected compact Kähler manifold Z with a surjective morphism $g : Z \rightarrow W$. Denote the composition $Z \xrightarrow{g} W \xrightarrow{f} C \xrightarrow{i} X$ by h . Then $h(Z) = C$. As E is flat over O_W , by [Sta24, Tag 05NJ], applying g^* to the natural short exact sequence

$$0 \rightarrow T(N) \rightarrow N \rightarrow E \rightarrow 0$$

in $\text{Mod}(O_W)$, one gets a short exact sequence in $\text{Mod}(O_Z)$:

$$0 \rightarrow g^*T(N) \rightarrow h^*F \rightarrow g^*E \rightarrow 0.$$

As g^*E is torsion free, $g^*T(N) \supset T(h^*F)$. One has $g^*T(N) \subset T(g^*N) = T(h^*F)$. Therefore, $T(h^*F) = g^*T(N)$ and $h^*F/T(h^*F) = g^*E$ is a vector bundle on Z of rank $r > 0$. \square

Lemma 5.5.3.4. *Let M be a coherent sheaf on the complex torus X . If $M \otimes P \cong M$ for all $P \in \text{Pic}^0(X)$, then $\text{Supp}(M)$ is finite.*

Proof. Suppose the contrary that $\text{Supp}(M)$ is infinite. With the reduced induced complex structure, the complex subspace $\text{Supp}(M)$ has positive dimension. Let C be an irreducible component of $\text{Supp}(M)$ of maximal dimension. Take a morphism $h : Z \rightarrow X$ provided by Lemma 5.5.3.3. Then the rank r of the vector bundle $E := h^*M/T(h^*M)$ is positive. As $h(Z) = C$, the morphism of complex tori $h^* : \text{Pic}^0(X) \rightarrow \text{Pic}^0(Z)$ is nonzero. In particular, there is $L \in \text{Pic}^0(X)$ such that the line bundle $(h^*L)^{\otimes r}$ is nontrivial.

On the other hand, we claim that the line bundle $(h^*L)^{\otimes r}$ is trivial. Indeed, by assumption $M \otimes L \cong M$, so $h^*M \otimes h^*L \cong h^*M$. Since $T(h^*M \otimes h^*L) = T(h^*M) \otimes h^*L$, one gets $E \otimes h^*L \cong E$. Taking the determinant of both sides, one has $\det(E) \otimes (h^*L)^{\otimes r} \cong \det(E)$. As $\det(E)$ is an invertible sheaf, the line bundle $(h^*L)^{\otimes r}$ on Z is trivial. The claim is proved, which gives a contradiction. \square

Remark 5.5.3.5. The proof of [Muk81, Lem. 3.3] (the algebraic counterpart of Lemma 5.5.3.4) relies on the following fact: Every positive dimensional projective variety contains a projective curve. By contrast, every simple non-algebraic complex torus contains no 1-dimensional analytic subset ([Pil00, Lem. 4.3]).

The classification of homogeneous vector bundles on complex tori is due to Matsushima [Mat59] and Morimoto [Mor59]. Using the Fourier-Mukai transform, Mukai [Muk81, p.159] proves an analog for abelian varieties. We can similarly recover Matsushima-Morimoto's theorem.

Theorem 5.5.3.6. *A vector bundle F on the complex torus X is homogeneous if and only if there is an integer $n \geq 0$, unipotent vector bundles U_1, \dots, U_n on X and $P_1, \dots, P_n \in \text{Pic}^0(X)$, such that F is isomorphic to $\bigoplus_{i=1}^n P_i \otimes U_i$.*

Proof. It follows from Propositions 5.5.1.1, 5.5.2.7 2 and 5.5.3.2 3. □

Chapter 6

Sheaves with connection on complex tori

6.1 Introduction

6.1.1 Background

Mukai [Muk81, Sec. 2] introduces an analog of the Fourier transform for sheaves of modules on abelian varieties, known as the *Fourier-Mukai transform*. Laumon [Lau96] and Rothstein [Rot96] study independently its lift to sheaves with connection (integrable or not). They both prove the Fourier inversion formula for the lift. Laumon [Lau96, Thm. 6.3.3] applies it to investigate generalized 1-motives. Meanwhile, as an application, Rothstein [Rot96, Thm. 3.2] recovers Matsushima's theorem ([Mat59]): every vector bundle on an abelian variety admitting a connection is translation invariant. Schnell's work [Sch15] about holonomic D -modules on abelian varieties relies upon the lift of the Fourier-Mukai transform.

Let k be an algebraically closed field. Let A, B be abelian varieties over k dual to each other. Set $g = \dim A$. Let p_A (resp. p_B) denote the projection from $A \times B$ to A (resp. B). Let \mathcal{P} be the normalized Poincaré line bundle on $A \times B$. We adopt the following sign convention for the Fourier-Mukai transform:

$$\begin{aligned} R\mathcal{S}_1 &= Rp_{A*}(\mathcal{P} \otimes^L p_B^* \cdot) : D(O_B) \rightarrow D(O_A); \\ R\mathcal{S}_2 &= Rp_{B*}(\mathcal{P}^{-1} \otimes^L p_A^* \cdot) : D(O_A) \rightarrow D(O_B), \end{aligned} \tag{6.1}$$

For a triangulated category, let T denote the degree shift automorphism. For an algebraic variety V over k , denote by $D_{\text{qc}}(O_V) \subset D(O_V)$ (resp. $D_c^b(O_V) \subset D^b(O_V)$) the full subcategory of objects whose cohomologies are quasi-coherent (resp. coherent) O_V -modules. Mukai establishes an analog of the Fourier inversion formula for this triangulated subcategory.

Fact 6.1.1.1 (Mukai, [Muk81, Thm. 2.2], [Rot96, p.569]). 1. *There are natural isomorphisms of functors $RS_1 \circ RS_2 \cong T^{-g}$ on $D_{\text{qc}}(O_A)$ and $RS_2 \circ RS_1 \cong T^{-g}$ on $D_{\text{qc}}(O_B)$. In particular, $RS_1 : D_{\text{qc}}(O_B) \rightarrow D_{\text{qc}}(O_A)$ is an equivalence of triangulated categories, with a quasi-inverse $T^g RS_2$.*

2. *The functor $RS_1 : D(O_B) \rightarrow D(O_A)$ restricts to an equivalence $D_c^b(O_B) \rightarrow D_c^b(O_A)$.*

Let $0 \rightarrow H^0(A, \Omega_A^1) \rightarrow B^\natural \xrightarrow{p} B \rightarrow 0$ be the universal vectorial extension of B (constructed in [Ros58, Prop. 11]). For an algebraic variety V , denote the forgetful functor $D(D_V) \rightarrow D(O_V)$ by for_V . Let $D_{\text{qc}}(D_A) \subset D(D_A)$ (resp. $D_c^b(D_A) \subset D^b(D_A)$) be the full subcategory of objects whose cohomologies are quasi-coherent O_A -modules (resp. coherent D_A -modules). Laumon and Rothstein lift the Fourier-transform to D -modules and establish a duality result similar to Fact 6.1.1.1.

Fact 6.1.1.2 (Laumon, Rothstein).

1. *There are functors $RS_1 : D(O_{B^\natural}) \rightarrow D(D_A)$ and $RS_2 : D(D_A) \rightarrow D(O_{B^\natural})$ fitting into commutative squares*

$$\begin{array}{ccc} D_{\text{qc}}(O_{B^\natural}) & \xrightarrow{RS_1} & D_{\text{qc}}(D_A) \\ \downarrow Rp_* & & \downarrow \text{for}_A \\ D_{\text{qc}}(O_B) & \xrightarrow{RS_1} & D_{\text{qc}}(O_A), \end{array} \quad \begin{array}{ccc} D_{\text{qc}}(O_{B^\natural}) & \xleftarrow{RS_2} & D_{\text{qc}}(D_A) \\ \downarrow Rp_* & & \downarrow \text{for}_A \\ D_{\text{qc}}(O_B) & \xleftarrow{RS_2} & D_{\text{qc}}(O_A). \end{array}$$

2. (Remark 6.1.1.4) *There are natural isomorphisms of functors $RS_1 RS_2 \cong T^{-g}$ on $D_{\text{qc}}(D_A)$ and $RS_2 RS_1 \cong T^{-g}$ on $D_{\text{qc}}(O_{B^\natural})$, hence an equivalence $RS_1 : D_{\text{qc}}(O_{B^\natural}) \rightarrow D_{\text{qc}}(D_A)$.*

3. ([Lau96, Cor. 3.1.3], [Rot96, Thm. 6.2]) *The functor $RS_1 : D(O_{B^\natural}) \rightarrow D(D_A)$ restricts to an equivalence $RS_1 : D_c^b(O_{B^\natural}) \rightarrow D_c^b(D_A)$.*

Remark 6.1.1.3. Laumon and Rothstein use apparently different definitions for the functors on D -modules. We sketch why the two definitions agree.

In the notation of [Lau96, p.14] and [Vig21, Sec. 2.1.1], one has functors $\tilde{\mathcal{F}} : D_{\text{qc}}^b(D_A) \rightarrow D_{\text{qc}}^b(O_{B^\natural})$ and $\tilde{\mathcal{F}}^\natural : D_{\text{qc}}^b(O_{B^\natural}) \rightarrow D_{\text{qc}}^b(D_A)$ defined by composition

$$\begin{array}{ccc} D_{\text{qc}}^b(D_A) & \xrightarrow{\tilde{\mathcal{F}}} & D_{\text{qc}}^b(O_{B^\natural}) \\ \tilde{\text{pr}}^!(B^\natural) \downarrow & & \uparrow \tilde{\text{pr}}^!_{+/B^\natural} \\ D_{\text{qc}}^b(D_{B^\natural \times A/B^\natural}) & \xrightarrow{L_{(\tilde{\mathcal{P}}, \tilde{\nabla}) \otimes_{O_{B^\natural \times A}}}} & D_{\text{qc}}^b(D_{B^\natural \times A/B^\natural}); \end{array} \quad \begin{array}{ccc} D_{\text{qc}}^b(O_{B^\natural}) & \xrightarrow{\tilde{\mathcal{F}}^\natural} & D_{\text{qc}}^b(D_A) \\ \tilde{\text{pr}}^{\natural, b}_{/B^\natural} \downarrow & & \uparrow \tilde{\text{pr}}^{\natural, b}_{+/B^\natural} \\ D_{\text{qc}}^b(D_{A \times B^\natural/B^\natural}) & \xrightarrow{L_{(\tilde{\mathcal{P}}, \tilde{\nabla}) \otimes_{O_{A \times B^\natural}}}} & D_{\text{qc}}^b(D_{A \times B^\natural/B^\natural}). \end{array}$$

Applying the projection formula (to $\text{Id}_A \times p : A \times B^\natural \rightarrow A \times B$) and the flat base change to the cartesian square

$$\begin{array}{ccc} A \times B^\natural & \xrightarrow{\text{Id}_A \times p} & A \times B \\ \downarrow p_{B^\natural} & & \downarrow p_B \\ B^\natural & \xrightarrow{p} & B, \end{array}$$

one gets an isomorphism for ${}_A\tilde{\mathcal{F}}^\natural \cong \mathcal{F}' R p_*$ of functors $D_{\text{qc}}^b(O_{B^\natural}) \rightarrow D_{\text{qc}}^b(O_A)$. This shows the compatibility with the Fourier-Mukai transform, as well as that $[-1]_A^* \tilde{\mathcal{F}}^\natural$ is the restriction of [Rot96, (4.16)] to $D_{\text{qc}}^b(O_{B^\natural})$. (The sign is due to different conventions.) Fact 6.1.1.2 1 is not mentioned in [Lau96], and is implicitly used in the derivation of [Rot96, (2.25)]. For Rothstein's definition, the compatibility can be proved as in Proposition 6.3.1.2.

Remark 6.1.1.4. The direct image functor $p_{A*} : \text{Mod}(O_{A \times B}) \rightarrow \text{Mod}(O_A)$ restricts to a left exact functor $p_{A*} : \text{Qch}(O_{A \times B}) \rightarrow \text{Qch}(O_A)$. Let $R(\text{qc})p_{A*} : D(\text{Qch}(O_{A \times B})) \rightarrow D(\text{Qch}(O_A))$ be the right derived functor of the restriction. Denote the functor

$$R(\text{qc})p_{A*}(\mathcal{P} \otimes_{O_{A \times B}} p_B^* \pi_* \cdot) : D(\text{Qch}(O_{B^\natural})) \rightarrow D(\text{Mod}_{\text{qc}}(D_A))$$

by $R(\text{qc})S_1$. Strictly speaking, [Rot96, Thm. 4.5] and [Rot97] demonstrate that the functor $R(\text{qc})S_1$ is an equivalence. In comparison, Laumon's result [Lau96, Thm. 3.2.1] is stated for bounded derived categories D_{qc}^b and needs the characteristic of k to be 0.

We sketch how to get Fact 6.1.1.2 2 from Rothstein's original statement. For every algebraic variety V , by [Sta24, Tag 077P (1)], the abelian category $\text{Qch}(O_V)$ has enough injectives. Furthermore, from [Con00, Lem. 2.1.3], the inclusion $\iota_V : \text{Qch}(O_V) \rightarrow \text{Mod}(O_V)$ preserves injectives. Let $\text{Mod}(O_B)_{\text{sp}}$ be as in Example 6.2.1.5 (resp. $\text{Mod}(O_{A \times B})_{-1-\text{cxi}}$ denote $\text{Mod}(O_{A \times B})_{\pi_B, -\pi_B^* 1-\text{cxi}}$). Let $\text{Qch}(O_B)_{\text{sp}}$ (resp. $\text{Qch}(O_{A \times B})_{-1-\text{cxi}}$) be the full subcategory of quasi-coherent objects.

Then the exact functor $\pi_* : \text{Qch}(O_{B^\natural}) \rightarrow \text{Qch}(O_B)_{\text{sp}}$ is the restriction of $R\pi_* : D(O_{B^\natural}) \rightarrow D(\text{Mod}(O_B)_{\text{sp}})$. Using [Lip60, Prop. 3.9.2] and [Har66, I, Prop. 7.1 (iii)], one proves that the canonical square

$$\begin{array}{ccc} D(\text{Qch}(O_{B^\natural})) & \xrightarrow{L\iota_{B^\natural}} & D_{\text{qc}}(O_{B^\natural}) \\ \downarrow \pi_* & & \downarrow R\pi_* \\ D(\text{Qch}(O_B)_{\text{sp}}) & \longrightarrow & D_{\text{qc}}(\text{Mod}(O_B)_{\text{sp}}) \end{array}$$

is commutative. Similarly, using [Kas04, Remark 3.2], one proves that the canonical square

$$\begin{array}{ccc}
D(\mathrm{Qch}(O_{A \times B})_{-1-\mathrm{c} \times \mathrm{n}}) & \longrightarrow & D_{\mathrm{qc}}(\mathrm{Mod}(O_{A \times B})_{-1-\mathrm{c} \times \mathrm{n}}) \\
\downarrow p_{A*} & & \downarrow R p_{A*} \\
D(\mathrm{Mod}_{\mathrm{qc}}(D_A)) & \xrightarrow{L\iota'_A} & D_{\mathrm{qc}}(D_A)
\end{array}$$

is commutative. Therefore, the following square is commutative

$$\begin{array}{ccc}
D(\mathrm{Qch}(O_{B^\natural})) & \xrightarrow{R(\mathrm{qc})S_1} & D(\mathrm{Mod}_{\mathrm{qc}}(D_A)) \\
\downarrow L\iota_{B^\natural} & & \downarrow L\iota'_A \\
D_{\mathrm{qc}}(O_{B^\natural}) & \xrightarrow{RS_1} & D_{\mathrm{qc}}(D_A).
\end{array} \tag{6.2}$$

By [Sta24, Tag 09T4] and Theorem E.1.0.4, the two vertical functors in (6.2) are equivalences. As $R(\mathrm{qc})S_1$ is an equivalence, so is the bottom row.

Remark 6.1.1.5. From [Sch14, p.97] and the square in [HT07, p.38], the bifunctor \otimes_O on relative D -modules is compatible with that on the underlying O -modules. However, the following triangles

$$\begin{array}{ccc}
D_{\mathrm{qc}}^b(D_{A \times B^\natural/B^\natural}) & \xrightarrow{(\tilde{\mathcal{P}}, \tilde{\nabla}) \otimes_{O_{A \times B^\natural}}^L \cdot} & D_{\mathrm{qc}}^b(D_{A \times B^\natural/B^\natural}) \\
\searrow \text{for} & & \nearrow (\tilde{\mathcal{P}}, \tilde{\nabla}) \otimes_{O_{A \times B^\natural}}^L \cdot \\
& D_{\mathrm{qc}}^b(O_{A \times B^\natural}) & \\
D_{\mathrm{qc}}^b(O_{A \times B^\natural}) & \xrightarrow{(\tilde{\mathcal{P}}, \tilde{\nabla}) \otimes_{O_{A \times B^\natural}}^L \cdot} & D_{\mathrm{qc}}^b(D_{A \times B^\natural/B^\natural}) \\
\searrow \tilde{\mathcal{P}} \otimes_{O_{A \times B^\natural}}^L \cdot & & \nearrow \text{for} \\
& D_{\mathrm{qc}}^b(O_{A \times B^\natural}) &
\end{array}$$

are not commutative in general. Thus, the first remark in [Vig21, p.58] is not true. In particular, the last but one equations in the proofs of [Vig21, Propositions 2.2.12 and 2.2.13] are wrong. Similarly, in [Lau96, p.14], the relative integrable connection on $(\tilde{\mathcal{P}}, \tilde{\nabla}) \otimes_{O_{B^\natural \times A}} \tilde{\mathrm{pr}}^{\natural*} M^\natural$ is induced by not only $\tilde{\nabla}$, but also the canonical relative connection on $\tilde{\mathrm{pr}}^{\natural*} M^\natural$.

6.1.2 Extension to complex tori

Let X, Y be complex tori dual to each other and of dimension g . Define the analytic Fourier-Mukai transform $RS_1 : D(O_X) \rightarrow D(O_Y)$ and $RS_2 : D(O_Y) \rightarrow D(O_X)$ by formulae similar to (6.1). For a complex manifold Z , let $D_{\mathrm{good}}(O_Z) \subset D(O_Z)$ be the full subcategory of objects whose cohomologies are good O_Z -modules (in the sense of [Kas03, Def. 4.22]). In [BBBP07, Thm. 2.1], a result similar to Fact 6.1.1.1 is established for complex tori.

Fact 6.1.2.1 (Mukai, Ben-Bassat, Block, Pantev).

1. (Theorem 5.4.1.1) *There are natural isomorphisms of functors*

$$\begin{aligned} RS_1 RS_2 &\cong T^{-g} : D_{\text{good}}(O_Y) \rightarrow D_{\text{good}}(O_Y), \\ RS_2 RS_1 &\cong T^{-g} : D_{\text{good}}(O_X) \rightarrow D_{\text{good}}(O_X). \end{aligned}$$

In particular, $RS_1 : D_{\text{good}}(O_X) \rightarrow D_{\text{good}}(O_Y)$ is an equivalence of categories with a quasi-inverse $T^g RS_2$.

2. ([PPS17, Thm. 13.1]) *The functor $RS_1 : D(O_X) \rightarrow D(O_Y)$ restricts to an equivalence $D_c^b(O_X) \rightarrow D_c^b(O_Y)$.*

We lift the analytic Fourier-Mukai transform to D -modules, and give an analog of Fact 6.1.1.2. Good D -modules are reviewed in Section 6.6.1. For a complex manifold Z and an O_Z -algebra \mathcal{R} , let $D_{O\text{-good}}(\mathcal{R}) \subset D(\mathcal{R})$ (resp. $D_{\text{good}}^b(\mathcal{R}) \subset D^b(\mathcal{R})$) be the full subcategory of objects whose cohomologies are good over O_Z (resp. \mathcal{R}).

Theorem 6.1.2.2.

- (Prop. 6.5.1.2) *There is a canonical commutative O_X -algebra \mathcal{A}_X , such that the functors RS_1 and RS_2 lift naturally to triangulated functors $RS_1 : D(\mathcal{A}_X) \rightarrow D(D_Y)$ and $RS_2 : D(D_Y) \rightarrow D(\mathcal{A}_X)$ respectively.*
- (Thm. 6.5.1.3) *The functors RS_i restrict to equivalences $RS_1 : D_{O\text{-good}}(\mathcal{A}_X) \rightarrow D_{O\text{-good}}(D_Y)$ and $RS_2 : D_{O\text{-good}}(D_Y) \rightarrow D_{O\text{-good}}(\mathcal{A}_X)$.*
- (Thm. 6.6.3.1) *The functors RS_i restricts to equivalences $RS_1 : D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(D_Y)$ and $RS_2 : D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_X)$.*

Remark 6.1.2.3. Arinkin [Fav12, Thm. 3] uses Fact 6.1.1.2 3 to show that an abelian variety A can be recovered from the triangulated category $D_c^b(D_A)$. By Proposition F.5.4.7, however, for a complex abelian variety A , the complex Lie group $(A^\natural)^{\text{an}}$ (associated with A^\natural) is isomorphic to $(\mathbb{C}^*)^{2g}$. So an analytic version of Fact 6.1.1.2 3 needs a modification.

The proof of Fact 6.1.1.2 due to Laumon [Lau96] and that of Rothstein [Rot97] are different. Let $\pi^\natural : A^\natural \rightarrow \text{Spec}(k)$ be the structural morphism. As an immediate step, Laumon [Lau96, Thm. 2.4.1] proves that the adjunction morphism $O_{\text{Spec}(k)} \rightarrow R\pi_*^\natural O_{A^\natural}$ is an isomorphism in $D_{\text{qc}}^b(O_{\text{Spec}(k)})$. By contrast, when $k = \mathbb{C}$, the adjunction morphism $O_{\text{Specan}(\mathbb{C})} \rightarrow R(\pi^\natural)_*^{\text{an}} O_{(A^\natural)^{\text{an}}}$ is not an isomorphism. Still, the proof of [Rot97] works for complex tori. We follow it closely, except that the underived Fourier-Mukai transforms [Rot97, (2.14), (2.15)] are ignored. Instead, we define the corresponding functors on the derived categories directly. We should notice four misprints therein.

- [Rot97, (2.11)] should be

$$\widetilde{\pi_{12}}^* \mathcal{P} = [(1_X \times \tilde{m})^* \mathcal{P}] \otimes_{\mathcal{O}_{X \times Y \times \tilde{Y}}} [\widetilde{\pi_{13}}^* \tilde{\mathcal{P}}^{-1}],$$

where $\widetilde{\pi_{ij}}$ denotes the projections on $X \times Y \times \tilde{Y}$ and \tilde{Y} is the first-order neighborhood of 0 in Y .

- [Rot97, (2.23)] should be

$$\pi_{13}^* \mathcal{P}^{-1} \otimes \pi_{23}^* \mathcal{P} = (\epsilon_X \times 1_Y)^* \mathcal{P},$$

where π_{ij} denotes the projections on $X \times X \times Y$.

- In [Rot97, (2.24)], the starting equation should be

$$\tilde{\pi}_{12}^* \mathcal{O}_\Delta \otimes \tilde{\pi}_{13}^* \tilde{\mathcal{P}}^{-1} \otimes \tilde{\pi}_{23}^* \tilde{\mathcal{P}}.$$

- In [Rot97, Prop. 2.4], the notation $\text{Mod}(X \times X)_{(-1,1)\text{-sp}}$ should be $\text{Mod}(X \times X)_{(1,-1)\text{-sp}}$.

Notation and convention

For a sheaf F on a topological space, let $\text{Supp } F$ be its support. For a (not necessarily commutative) ringed space (X, \mathcal{R}) , let $\text{Mod}(\mathcal{R})$ be the category of left \mathcal{R} -modules. Let $\text{Coh}(\mathcal{R}) \subset \text{Mod}(\mathcal{R})$ be the full subcategory of coherent \mathcal{R} -modules. Given a symbol $*$ $\in \{\emptyset, +, -, b\}$, the notation $D^*(\mathcal{R})$ refers to the unbounded/bounded below/bounded above/bounded derived category of the abelian category $\text{Mod}(\mathcal{R})$ in order. Let $D_c^*(\mathcal{R}) \subset D^*(\mathcal{R})$ be the full subcategory of objects whose cohomologies are coherent \mathcal{R} -modules (in the sense of [Sta24, Tag 01BV]).

Let k be an algebraically closed field. An algebraic variety refers to an integral scheme of finite type and separated over k . For a complex manifold Z and $z \in Z$, let $i_z : (z, \mathbb{C}) \rightarrow (Z, \mathcal{O}_Z)$ be the closed embedding of complex manifolds. Set $\mathbb{C}_z := (i_z)_* \mathbb{C}$, which is a coherent \mathcal{O}_Z -module. Let X, Y be complex tori dual to each other and of dimension g .

6.2 Preliminaries

For the convenience of the reader, we recall the notation of [Rot97, Sec. 2.1].

6.2.1 Categories of splittings

For a complex manifold Z and a (holomorphic) vector bundle $M \rightarrow Z$, by [Har77, III, Prop. 6.3 (c)], one has $H^1(Z, M) = \text{Ext}^1(O_Z, M)$. Thus, every $\alpha \in H^1(Z, M)$ determines a short exact sequence in $\text{Mod}(O_Z)$

$$0 \rightarrow M \rightarrow \mathcal{E}_\alpha \xrightarrow{\mu_\alpha} O_Z \rightarrow 0. \quad (6.3)$$

Since O_Z is a flat O_Z -module, by [Sta24, Tag 05NJ], for every $F \in \text{Mod}(O_Z)$, the sequence (6.3) remains exact after tensored with F :

$$0 \rightarrow M \otimes_{O_Z} F \rightarrow \mathcal{E}_\alpha \otimes_{O_Z} F \xrightarrow{\mu_\alpha \otimes \text{Id}_F} F \rightarrow 0.$$

Definition 6.2.1.1. Define a category $\text{Mod}(O_Z)_{\alpha\text{-sp}}$ as follows: the objects are pairs (F, ψ) , where $F \in \text{Mod}(O_Z)$ and $\psi : F \rightarrow \mathcal{E}_\alpha \otimes_{O_Z} F$ is an α -splitting on F , i.e., an O_Z -linear splitting of $\mu_\alpha \otimes \text{Id}_F$. The morphisms in $\text{Mod}(O_Z)_{\alpha\text{-sp}}$ are required to be compatible with the splittings.

Example 6.2.1.2. When $\alpha = 0$, the sequence (6.3) identifies \mathcal{E}_0 with $M \oplus O_Z$. There is a natural functor $\text{Mod}(O_Z) \rightarrow \text{Mod}(O_Z)_{0\text{-sp}}$ defined by $F \mapsto (F, \psi)$, where $\psi : F \rightarrow \mathcal{E}_0 \otimes F = (M \otimes_{O_Z} F) \oplus F$ is the canonical injection to the second factor. If further $M = \Omega_Z^1$, then an α -splitting ϕ on a vector bundle $E \rightarrow Z$ is exactly a holomorphic 1-form on Z with values in $\text{End}(E)$. The pair (E, ϕ) is a Higgs bundle (in the sense of [Sim92, p.6]) if and only if $[\phi, \phi] = 0$.

Lemma 6.2.1.3. For an O_Z -module F , there is an α -splitting on F if and only if the map $i_* : H^1(Z, M) \rightarrow H^1(Z, M \otimes_{O_Z} \text{End}(F))$ (induced by the natural morphism $O_Z \rightarrow \text{End}(F)$) sends α to 0. In that case, the set of α -splittings on F has a natural simple transitive action of the abelian group $\text{Hom}_{O_Z}(F, M \otimes_{O_Z} F)$.

Proof. The natural morphism $O_Z \rightarrow \text{End}(F)$ induces a morphism $i : M \rightarrow \text{Hom}_{O_Z}(F, M \otimes_{O_Z} F)$, $i(m)(f) = m \otimes f$. There is a canonical evaluation morphism $\text{ev} : \text{Hom}_{O_Z}(F, M \otimes_{O_Z} F) \otimes F \rightarrow M \otimes_{O_Z} F$, $\text{ev}(\phi \otimes f) = \phi(f)$. The five-term exact sequence of the spectral sequence

$$E_2^{i,j} = \text{Ext}^i(O_Z, \text{Ext}^j(F, M \otimes_{O_Z} F)) \Rightarrow \text{Ext}^{i+j}(F, M \otimes_{O_Z} F)$$

gives an injection $\iota : \text{Ext}^1(O_Z, \text{Hom}(F, M \otimes_{O_Z} F)) \rightarrow \text{Ext}^1(F, M \otimes_{O_Z} F)$, which is $\text{Ext}^1(F, \text{ev}) \circ (\cdot \otimes F)$:

$$\begin{array}{ccccc} & & \text{Ext}^1(F, M \otimes_{O_Z} F) & & \\ & \nearrow \cdot \otimes F & \downarrow = & \searrow (i \otimes \text{Id}_F)_* & \\ \text{Ext}^1(O_Z, M) & & \text{Ext}^1(F, M \otimes_{O_Z} F) & \xleftarrow{\text{Ext}^1(F, \text{ev})} & \text{Ext}^1(F, \text{Hom}(F, M \otimes_{O_Z} F) \otimes F) \\ & \searrow i_* & \uparrow \iota & \nearrow \cdot \otimes F & \\ & & \text{Ext}^1(O_Z, \text{Hom}(F, M \otimes_{O_Z} F)) & & \end{array}$$

One has

$$\text{ev} \circ (i \otimes \text{Id}_F)(m \otimes f) = \text{ev}(i(m) \otimes f) = i(m)(f) = m \otimes f,$$

so $\text{ev} \circ (i \otimes \text{Id}_F) = \text{Id}_{M \otimes_{O_Z} F}$ as morphisms $M \otimes_{O_Z} F \rightarrow M \otimes_{O_Z} F$. Therefore, the diagram is commutative. Then F admits an α -splitting if and only if $\alpha \otimes F = 0$ if and only if $i_*(\alpha) = 0$. Any two α -splittings on F differ by a unique element of $\text{Hom}(F, M \otimes_{O_Z} F)$. \square

To each object $(F, \psi) \in \text{Mod}(O_Z)_{\alpha\text{-sp}}$, we assign an element

$$[\psi, \psi] \in \Gamma(Z, (\wedge^2 M) \otimes_{O_Z} \mathcal{E}nd(F)) \quad (6.4)$$

as follows. The sequence (6.3) induces a short exact sequence

$$0 \rightarrow \wedge^2 M \rightarrow \wedge^2 \mathcal{E}_\alpha \xrightarrow{\omega_\alpha} M \rightarrow 0,$$

where

$$\omega_\alpha(\rho_1 \wedge \rho_2) = \mu_\alpha(\rho_1)\rho_2 - \mu_\alpha(\rho_2)\rho_1.$$

The flatness of M ensures the exactness when tensoring with F :

$$0 \rightarrow (\wedge^2 M) \otimes F \rightarrow (\wedge^2 \mathcal{E}_\alpha) \otimes F \xrightarrow{\omega_\alpha \otimes \text{Id}_F} M \otimes_{O_Z} F \rightarrow 0. \quad (6.5)$$

Let $a : \mathcal{E}_\alpha \otimes \mathcal{E}_\alpha \rightarrow \wedge^2 \mathcal{E}_\alpha$ be the morphism defined by $e \otimes e' \mapsto e \wedge e'$. Let ψ^1 be the composition

$$\mathcal{E}_\alpha \otimes F \xrightarrow{\text{Id}_{\mathcal{E}_\alpha} \otimes \psi} \mathcal{E}_\alpha \otimes (\mathcal{E}_\alpha \otimes F) \xrightarrow{\sim} (\mathcal{E}_\alpha \otimes \mathcal{E}_\alpha) \otimes F \xrightarrow{a \otimes \text{Id}_F} (\wedge^2 \mathcal{E}_\alpha) \otimes F,$$

where the isomorphism in the middle is from the associativity of tensor product.

Lemma 6.2.1.4. *One has $(\omega_\alpha \otimes \text{Id}_F)\psi^1\psi = 0$.*

Proof. Locally, the vector bundle \mathcal{E}_α has a (holomorphic) frame $\{e_1, \dots, e_r\}$. For a local section $f \in F$, write $\psi(f) = \sum_{i=1}^r e_i \otimes f_i$, where f_i are local sections of F . For every $1 \leq i \leq r$, write $\psi(f_i) = \sum_{j=1}^r e_j \otimes f_j^{(i)}$, where $f_j^{(i)}$ are local sections of F . As ψ is a section to $\mu_\alpha \otimes \text{Id}_F$, one has

$$f = (\mu_\alpha \otimes \text{Id}_F)\psi(f) = \sum_{i=1}^r \mu_\alpha(e_i)f_i; \quad (6.6)$$

$$f_i = (\mu_\alpha \otimes \text{Id}_F)\psi(f_i) = \sum_{j=1}^r \mu_\alpha(e_j)f_j^{(i)}. \quad (6.7)$$

Thus,

$$\psi(f) \stackrel{(6.6)}{=} \sum_{i=1}^r \mu_\alpha(e_i)\psi(f_i). \quad (6.8)$$

By construction, $\psi^1\psi(f) = \sum_{i,j=1}^r (e_i \wedge e_j) \otimes f_j^{(i)}$. Then

$$\begin{aligned}
(\omega_\alpha \otimes \text{Id}_F)\psi^1\psi(f) &= \sum_{i,j=1}^r [\mu_\alpha(e_i)e_j - \mu_\alpha(e_j)e_i] \otimes f_j^{(i)} \\
&= \sum_{i=1}^r \mu_\alpha(e_i) \sum_{j=1}^r e_j \otimes f_j^{(i)} - \sum_{i=1}^r e_i \otimes [\sum_{j=1}^r \mu_\alpha(e_j)f_j^{(i)}] \\
&\stackrel{(6.7)}{=} \sum_{i=1}^r \mu_\alpha(e_i)\psi(f_i) - \sum_{i=1}^r e_i \otimes f_i \\
&\stackrel{(6.8)}{=} \psi(f) - \psi(f) = 0.
\end{aligned}$$

□

From Lemma 6.2.1.4 and (6.5), one has $\psi^1\psi(F) \subset (\wedge^2 M) \otimes F$. The morphism $\psi^1\psi : F \rightarrow (\wedge^2 M) \otimes F$ gives an element $[\psi, \psi] \in \Gamma(Z, (\wedge^2 M) \otimes_{O_Z} \mathcal{E}nd(F))$.

Example 6.2.1.5. For the complex torus X , set $\mathfrak{g} = H^1(X, O_X)$. Then

$$H^1(X, \mathfrak{g}^* \otimes_{\mathbb{C}} O_X) = \mathfrak{g}^* \otimes_{\mathbb{C}} \mathfrak{g} = \text{End}(\mathfrak{g}).$$

Hence a category $\text{Mod}(O_X)_{T-\text{sp}}$ for each $T \in \text{End}(\mathfrak{g})$. The identity element $1 \in \text{End}(\mathfrak{g})$ corresponds to the tautological exact sequence [Rot96, (1.3)]:

$$0 \rightarrow \mathfrak{g}^* \otimes_{\mathbb{C}} O_X \rightarrow \mathcal{E} \rightarrow O_X \rightarrow 0. \quad (6.9)$$

We also write $\text{Mod}(O_X)_{\text{sp}}$ for $\text{Mod}(O_X)_{1-\text{sp}}$. For $(F, \psi) \in \text{Mod}(O_X)_{\text{sp}}$, the element $[\psi, \psi]$ lies in

$$\Gamma(X, \wedge^2 \mathfrak{g}^* \otimes_{\mathbb{C}} O_X \otimes_{O_X} \mathcal{E}nd(F)) = \wedge^2 \mathfrak{g}^* \otimes_{\mathbb{C}} \text{End}(F),$$

and we recover [Rot96, (4.8)]. Similarly, $H^1(X \times X, \mathfrak{g}^* \otimes_{O_X \times X}) = \text{End}(g) \oplus \text{End}(g)$, so for every pair $T_1, T_2 \in \text{End}(g)$, the category $\text{Mod}(O_{X \times X})_{(T_1, T_2)-\text{sp}}$ is defined.

6.2.2 Categories of twisted connection

We continue to review the twisted (relative) connection introduced in [Rot97, p.206]. Consider a smooth morphism of complex manifolds $f : Z \rightarrow S$, with relative cotangent sheaf Ω_f^1 . As f is smooth, Ω_f^1 is a vector bundle on Z . Let $d_f : O_Z \rightarrow \Omega_f^1$ denote the differential relative to f . An element $\alpha \in H^1(Z, \Omega_f^1)$ determines an extension

$$0 \rightarrow \Omega_f^1 \rightarrow \mathcal{E}_\alpha \xrightarrow{\mu_\alpha} O_Z \rightarrow 0. \quad (6.10)$$

Definition 6.2.2.1. On an O_Z -module F , an α -connection is an $f^{-1}(O_S)$ -linear splitting $\nabla : F \rightarrow \mathcal{E}_\alpha \otimes_{O_Z} F$ to $\mu_\alpha \otimes \text{Id}_F$, satisfying the Leibniz rule

$$\nabla(h\phi) = h\nabla(\phi) + d_f(h) \otimes \phi, \quad (6.11)$$

where h and ϕ are local sections of O_Z and F respectively. Let $\text{Mod}(O_Z)_{f,\alpha-\text{cxn}}$ be the category of pairs (F, ∇) , where $F \in \text{Mod}(O_Z)$ and ∇ is an α -connection on F .

Example 6.2.2.2. If $\alpha = 0$, then α -connection are exactly f -relative connection. Define a sheaf $\tilde{D}_{Z/S}$ of noncommutative O_Z -algebras by gluing the following local data. If $\{\xi_1, \dots, \xi_n\}$ is a local frame of $(\Omega_f^1)^\vee$ (the vector bundle dual to Ω_f^1) on an open subset $U \subset Z$, then a multiplication law on $O_U\{\xi_1, \dots, \xi_n\}$ is introduced by imposing the commutation relation $[\xi_i, h] = \xi_i(h)$ for local sections h of O_Z . Let it be $\tilde{D}_{Z/S}|_U$. Then $\text{Mod}(Z)_{f,0-\text{cxn}} = \text{Mod}(\tilde{D}_{Z/S})$. The category $\text{Mod}(O_Z)_{f,0-\text{cxn}}$ is denoted by $\text{Mod}(O_Z)_{\text{cxn}}$ when f is the structure morphism $Z \rightarrow \text{Specan}(\mathbb{C})$.

Remark 6.2.2.3. In fact, a twisted connection is a particular splitting. Define another extension

$$0 \rightarrow \Omega_f^1 \rightarrow \mathcal{E}_{\alpha'} \rightarrow O_Z \rightarrow 0 \quad (6.12)$$

in $\text{Mod}(O_Z)$ as follows. As an extension of abelian sheaves, (6.12) is same as (6.10). Let h (resp. s') be a local section of O_Z (resp. $\mathcal{E}_{\alpha'}$) and s denote the local section of \mathcal{E}_α induced by s' . The O_Z -module structure on $\mathcal{E}_{\alpha'}$ is defined such that the local section $hs + \mu_\alpha(s)d_f h$ of \mathcal{E}_α induces the local section hs' of $\mathcal{E}_{\alpha'}$.

We claim this indeed defines an O_Z -module structure on $\mathcal{E}_{\alpha'}$. For local sections h_1, h_2 of O_Z , let t be the local section of \mathcal{E}_α induced by $h_2 s'$. Then $t = h_2 s + \mu_\alpha(s)d_f h_2$, so $\mu_\alpha(t) = h_2 \mu_\alpha(s)$. Thus, the local section of \mathcal{E}_α corresponding to $h_1(h_2 s')$ is

$$h_1 t + \mu_\alpha(t)d_f h_1 = h_1 h_2 s + h_1 \mu_\alpha(s)d_f h_2 + h_2 \mu_\alpha(s)d_f h_1 = (h_1 h_2)s + \mu_\alpha(s)d_f(h_1 h_2).$$

Therefore, $h_1(h_2 s') = (h_1 h_2)s'$. The claim is proved.

By construction, the morphisms in (6.12) are O_Z -linear. Then (6.12) is indeed an extension in $\text{Mod}(O_Z)$, hence a new extension class $\alpha' \in \text{Ext}(O_Z, \Omega_f^1)$. An α -connection on $F \in \text{Mod}(O_Z)$ is equivalent to an α' -splitting on F . Hence an equivalence of categories

$$\text{Mod}(O_Z)_{f,\alpha-\text{cxn}} \rightarrow \text{Mod}(O_Z)_{\alpha'-\text{sp}}.$$

There is a notion of integrable α -connection ([Rot97, Remark, p.206]). Let $\text{Mod}(O_Z)_{f,\alpha-\text{cxn},\text{fl}}$ be the full subcategory of $\text{Mod}(O_Z)_{f,\alpha-\text{cxn}}$ comprised of objects whose connection are integrable. Then $\text{Mod}(O_Z)_{f,0-\text{cxn},\text{fl}}$ coincides with $\text{MIC}(f)$ defined in [ABC20, 4.3.7], which is further equivalent to $\text{Mod}(D_{Z/S})$. Here $D_{Z/S}$ is the sheaf of ring of relative differential operators on Z/S defined in [SS94, p.9].

Example 6.2.2.4. For the dual complex tori X, Y , consider the projection $p_X : X \times Y \rightarrow X$. Since $\Omega_{p_X}^1 = p_X^*(\mathfrak{g}^* \otimes_{\mathbb{C}} O_X)$, there is a natural morphism

$$p_X^* : \text{End}(\mathfrak{g}) = H^1(X, \mathfrak{g}^* \otimes_{\mathbb{C}} O_X) \rightarrow H^1(X \times Y, \Omega_{p_X}^1).$$

For every $T \in \text{End}(g)$, the category $\text{Mod}(O_{X \times Y})_{p_X, p_X^* T - \text{cxn}}$ (resp. $\text{Mod}(O_{X \times Y})_{p_X, p_X^* T - \text{cxn}, \text{fl}}$) is also written as $\text{Mod}(O_{X \times Y})_{T - \text{cxn}}$ (resp. $\text{Mod}(O_{X \times Y})_{T - \text{cxn}, \text{fl}}$).

Fact 6.2.2.5 is taken from the two remarks in [Rot97, pp.206–207].

Fact 6.2.2.5. *The Poincaré bundle \mathcal{P} is naturally an object of $\text{Mod}(O_{X \times Y})_{-1 - \text{cxn}, \text{fl}}$.*

In local coordinates, the $p_X^*(-1)$ -connection on \mathcal{P} is explained in [Rot96, (1.10) and p.575ff.] (except that we use a Stein open cover of X instead of Rothstein’s affine open cover).

6.2.3 Functors between them

Recall that the Fourier-Mukai transform (6.1) is the composition of the pullback, the tensor product with \mathcal{P} as well as the derived direct image. Rothstein’s lift to modules with connection keeps an extra track of the splittings and connection.

Remark 6.2.3.1. Combining [Rot97, (2.21)] with the fact that twisted relative connection are kinds of splittings (Remark 6.2.2.3), the categories under consideration ($\text{Mod}(O_X)_{\text{sp}}$, $\text{Mod}(O_{X \times Y})_{T - \text{cxn}}$, etc.) are equivalent to categories of modules over sheaves of certain noncommutative *flat* O -algebras. In particular, each of them is a Grothendieck abelian category. Each has enough K -injectives ([Sta24, Tag 079P]) and enough objects flat over O ([HT07, Lem. 1.5.2 (ii)]), cf. [Rot97, Cor. 2.3]. Thus, all the (left exact) direct image functors involved below admit right derived functors on the unbounded derived categories (see [Sta24, Tag 070K] and [Sta24, Tag 079P]).¹

From splittings to connection

Given $T \in \text{End}(\mathfrak{g})$ and $(F, \psi) \in \text{Mod}(O_X)_{T - \text{sp}}$, the induced morphism

$$p_X^{-1}\psi : p_X^{-1}F \rightarrow p_X^{-1}\mathcal{E} \otimes_{p_X^{-1}O_X} p_X^{-1}F$$

is $p_X^{-1}O_X$ -linear. By Example 6.2.2.4, the sequence (6.9) induces a short exact sequence in $\text{Mod}(O_{X \times Y})$

$$0 \rightarrow \Omega_{p_X}^1 \rightarrow p_X^*\mathcal{E} \rightarrow O_{X \times Y} \rightarrow 0.$$

¹Rothstein [Rot97, Sec. 2.2] uses Čech resolutions for quasi-coherent sheaves, while we are dealing with *all* O -modules.

Its extension class is $p_X^*T \in H^1(X \times Y, \Omega_{p_X}^1)$. Define another $p_X^{-1}O_X$ -linear morphism

$$\begin{aligned} \nabla_\psi : p_X^*F &= (O_{X \times Y} \otimes_{p_X^{-1}O_X} p_X^{-1}F) \rightarrow p_X^*\mathcal{E} \otimes_{O_{X \times Y}} p_X^*F (= \\ p_X^*\mathcal{E} \otimes_{p_X^{-1}O_X} p_X^{-1}F &= O_{X \times Y} \otimes_{p_X^{-1}O_X} p_X^{-1}\mathcal{E} \otimes_{p_X^{-1}O_X} p_X^{-1}F) \end{aligned}$$

by

$$\nabla_\psi(h \otimes s) = d_{p_X}(h) \otimes s + h \otimes [(p_X^{-1}\psi)(s)],$$

where h (resp. s) is a local section of $O_{X \times Y}$ (resp. $p_X^{-1}F$). By construction, ∇_ψ satisfies the Leibniz rule (6.11). So it is a p_X^*T -connection on p_X^*F . Thus, we get the *exact* functor in [Rot97, (2.5)]:

$$p_X^* : \text{Mod}(O_X)_{T\text{-sp}} \rightarrow \text{Mod}(O_{X \times Y})_{T\text{-cxn}}. \quad (6.13)$$

Tensoring with Poincaré bundle

By Fact 6.2.2.5 and [Rot97, (2.10)], the functor

$$\cdot \otimes_{O_{X \times Y}} \mathcal{P} : \text{Mod}(O_{X \times Y})_{1\text{-cxn}} \rightarrow \text{Mod}(O_{X \times Y})_{0\text{-cxn}} \quad (6.14)$$

restricts to a functor $\text{Mod}(O_{X \times Y})_{1\text{-cxn, fl}} \rightarrow \text{Mod}(O_{X \times Y})_{0\text{-cxn, fl}} (\cong \text{Mod}(D_{X \times Y/X}))$. The functor (6.14) is an equivalence of abelian categories, with a quasi-inverse $\cdot \otimes_{O_{X \times Y}} \mathcal{P}^{-1}$.

From connection to splittings

For every $(F, \nabla) \in \text{Mod}(O_{X \times Y})_{1\text{-cxn}}$, the morphism

$$\nabla : F \rightarrow p_X^*\mathcal{E} \otimes_{O_{X \times Y}} F (= p_X^{-1}\mathcal{E} \otimes_{p_X^{-1}O_X} F)$$

is a $p_X^{-1}O_X$ -splitting to $(p_X^{-1}\mu_1) \otimes \text{Id}_F$. By projection formula (see e.g. [KS90, Prop. 2.6.6]), the induced morphism

$$p_{X*}\nabla : p_{X*}F \rightarrow \mathcal{E} \otimes_{O_X} p_{X*}F$$

is an O_X -linear splitting to $\mu_1 \otimes_{O_X} \text{Id}_{p_{X*}F}$. Hence $(p_{X*}F, p_{X*}\nabla) \in \text{Mod}(O_X)_{\text{sp}}$. Thus, we get a left exact functor (a special case of [Rot97, (2.13)]):

$$p_{X*} : \text{Mod}(O_{X \times Y})_{1\text{-cxn}} \rightarrow \text{Mod}(O_X)_{\text{sp}}. \quad (6.15)$$

If (F, ∇) is integrable, then $[p_{X*}\nabla, p_{X*}\nabla]$ defined in (6.4) is zero.

Between connection

We define the inverse image and the direct image of relative connection on changing bases. Consider a cartesian square of complex manifolds

$$\begin{array}{ccc} W & \xrightarrow{g'} & Z \\ \downarrow f' & \square & \downarrow f \\ T & \xrightarrow{g} & S, \end{array} \quad (6.16)$$

where f is smooth. For every $(F, \nabla) \in \text{Mod}(O_Z)_{f,0-\text{c}x\text{n}}$, by [ABC20, Sec. 4.2], the relative connection ∇ is equivalent to an O_Z -linear splitting to the natural projection $P_f^1 \otimes_{O_Z} F \rightarrow F$, where P_\bullet^1 denotes the sheaf of first order jets (defined in [ABC20, Sec. 4.1.2]). Applying g'^* to the induced splitting, we get an O_W -linear splitting to the natural projection $P_{f'}^1 \otimes_{O_W} g'^* F \rightarrow g'^* F$. This is equivalent to an f' -connection on $g'^* F$. Hence an inverse image functor

$$g'^* : \text{Mod}(O_Z)_{f,0-\text{c}x\text{n}} \rightarrow \text{Mod}(O_W)_{f',0-\text{c}x\text{n}}. \quad (6.17)$$

It is right exact. By [ABC20, Sec. 5.1], the connection induced by ∇ is integrable if ∇ is so.

Now for direct image. Fix $\alpha \in F^1(Z, \Omega_f^1)$. For every

$$(F, \nabla) \in \text{Mod}(O_W)_{f',g'^*\alpha-\text{c}x\text{n}},$$

by projection formula (see e.g, [Har77, II, Ex. 5.1 (d)]), one has

$$g'_*(F \otimes_{O_W} g'^* \mathcal{E}_\alpha) = (g'_* F) \otimes_{O_Z} \mathcal{E}_\alpha.$$

Then the induced morphism

$$g'_* \nabla : g'_* F \rightarrow (g'_* F) \otimes_{O_Z} \mathcal{E}_\alpha$$

is $f^{-1}(O_S)$ -linear. Since $d_{f'} : O_W \rightarrow \Omega_{f'}^1$ and $d_f : O_Z \rightarrow \Omega_f^1$ are related by $g'^* d_f = d_{f'}$, the induced map $g'_* \nabla$ satisfies the Leibniz rule (6.11). Hence, the pair $(g'_* F, g'_* \nabla) \in \text{Mod}(O_Z)_{f,\alpha-\text{c}x\text{n}}$. In this manner, we get a left exact functor

$$g'_* : \text{Mod}(O_W)_{f',g'^*\alpha-\text{c}x\text{n}} \rightarrow \text{Mod}(O_Z)_{f,\alpha-\text{c}x\text{n}}. \quad (6.18)$$

When $\alpha = 0$, the functor (6.18) sends $\text{MIC}(f')$ to $\text{MIC}(f)$.

Example 6.2.3.2. Take (6.16) to be

$$\begin{array}{ccc} X \times Y & \xrightarrow{p_Y} & Y \\ \downarrow p_X & \square & \downarrow \\ X & \longrightarrow & \text{Specan}(\mathbb{C}), \end{array}$$

then $p_Y^* : \text{Mod}(O_Y)_{\text{cxd}} \rightarrow \text{Mod}(O_{X \times Y})_{0-\text{cxd}}$ sits on the left of the diagram [Rot97, (2.15)] and

$$p_{Y*} : \text{Mod}(O_{X \times Y})_{0-\text{cxd}} \rightarrow \text{Mod}(Y)_{\text{cxd}} \quad (6.19)$$

is [Rot97, (2.12)]. They restrict respectively to functors

$$p_{Y*} : \text{Mod}(D_{X \times Y/X}) \rightarrow \text{Mod}(D_Y); \quad (6.20)$$

$$p_Y^* : \text{Mod}(D_Y) \rightarrow \text{Mod}(D_{X \times Y/X}). \quad (6.21)$$

Remark 6.2.3.3. Take $\alpha = 0 \in H^1(Z, \Omega_f^1)$. From another point of view, the morphism $O_Z \rightarrow g'_* O_W$ between sheaves of rings extends to a morphism $\tilde{D}_{Z/S} \rightarrow g'_* \tilde{D}_{W/T}$. Then (6.17) and (6.18) are respectively the pullback and the pushout along the induced morphism $(W, \tilde{D}_{W/T}) \rightarrow (Z, \tilde{D}_{Z/S})$ of ringed spaces. By [Sta24, Tag 0096], the functor (6.17) is the left adjoint to (6.18). Then from [Sta24, Tag 09T5], the derived functor

$$Lg'^* : D(\text{Mod}(Z)_{f,0-\text{cxd}}) \rightarrow D(\text{Mod}(W)_{f',0-\text{cxd}})$$

is the left adjoint to

$$Rg'_* : D(\text{Mod}(W)_{f',0-\text{cxd}}) \rightarrow D(\text{Mod}(Z)_{f,0-\text{cxd}}).$$

6.3 Rothstein transform on modules with connection

6.3.1 Construction

Definition 6.3.1.1. Define functors $R\mathfrak{S}_1 : D(\text{Mod}(O_X)_{\text{sp}}) \rightarrow D(\text{Mod}(O_Y)_{\text{cxd}})$ and $R\mathfrak{S}_2 : D(\text{Mod}(O_Y)_{\text{cxd}}) \rightarrow D(\text{Mod}(O_X)_{\text{sp}})$ by

$$\begin{aligned} R\mathfrak{S}_1 &= Rp_{Y*}(\mathcal{P} \otimes_{O_{X \times Y}} p_{X*}^*), \\ R\mathfrak{S}_2 &= Rp_{X*}(\mathcal{P}^{-1} \otimes_{O_{X \times Y}} p_Y^*). \end{aligned}$$

Here Rp_{Y*} (resp. Rp_{X*}) is the right derived functor of (6.19) (resp. (6.15)). The pair $(R\mathfrak{S}_1, R\mathfrak{S}_2)$ is called the *Rothstein transform*.

Let $D_{O-\text{good}}(\text{Mod}(O_Y)_{\text{cxd}}) \subset D(\text{Mod}(O_Y)_{\text{cxd}})$ (resp. $D_{O-\text{good}}(\text{Mod}(O_X)_{\text{sp}}) \subset D(\text{Mod}(O_X)_{\text{sp}})$) be the full subcategory of objects whose cohomologies are good O -modules (in the sense of [Kas03, Def. 4.22]). In view of Proposition 6.3.1.2, Rothstein transform is compatible with Fourier-Mukai transform.

Proposition 6.3.1.2. *There are commutative squares*

$$\begin{array}{ccc} D(\text{Mod}(O_X)_{\text{sp}}) & \xrightarrow{R\mathfrak{S}_1} & D(\text{Mod}(O_Y)_{\text{cxd}}) \\ \downarrow & & \downarrow \\ D(O_X) & \xrightarrow{RS_1} & D(O_Y), \end{array} \quad \begin{array}{ccc} D(\text{Mod}(O_Y)_{\text{cxd}}) & \xrightarrow{R\mathfrak{S}_2} & D(\text{Mod}(O_X)_{\text{sp}}) \\ \downarrow & & \downarrow \\ D(O_Y) & \xrightarrow{RS_2} & D(O_X), \end{array}$$

where the vertical functors are forgetful. In particular, $R\mathfrak{S}_1$ and $R\mathfrak{S}_2$ restrict to functors $D_{O-\text{good}}(\text{Mod}(O_X)_{\text{sp}}) \rightarrow D_{O-\text{good}}(\text{Mod}(O_Y)_{\text{cxd}})$ and $D_{O-\text{good}}(\text{Mod}(O_Y)_{\text{cxd}}) \rightarrow D_{O-\text{good}}(\text{Mod}(O_X)_{\text{sp}})$.

Proof. All the functors $p_X^* : \text{Mod}(O_X) \rightarrow \text{Mod}(O_{X \times Y})$, (6.13), (6.14) and

$$\mathcal{P} \otimes_{O_{X \times Y}} \cdot : \text{Mod}(O_{X \times Y}) \rightarrow \text{Mod}(O_{X \times Y})$$

are exact. To prove the commutativity of the first square, it remains to do so for the square

$$\begin{array}{ccc} D(\text{Mod}(O_{X \times Y})_{0-\text{cxd}}) & \xrightarrow{Rp_{Y*}} & D(\text{Mod}(O_Y)_{\text{cxd}}) \\ \downarrow \text{for}_{X \times Y} & & \downarrow \text{for}_Y \\ D(O_{X \times Y}) & \xrightarrow{Rp_{Y*}} & D(O_Y). \end{array}$$

Since the forgetful functor $\text{for}_Y : \text{Mod}(O_Y)_{\text{cxd}} \rightarrow \text{Mod}(O_Y)$ is exact, the composition $\text{for}_Y Rp_{Y*} : D(\text{Mod}(O_{X \times Y})_{0-\text{cxd}}) \rightarrow D(O_Y)$ is the right derived functor of

$$\text{for}_Y \circ p_{Y*} : \text{Mod}(O_{X \times Y})_{0-\text{cxd}} \rightarrow \text{Mod}(O_Y).$$

From Remark 6.2.3.1, [Sta24, Tag 0096] and [Sta24, Tag 08BJ], the functor $\text{for}_{X \times Y} : \text{Mod}(O_{X \times Y})_{0-\text{cxd}} \rightarrow \text{Mod}(O_{X \times Y})$ preserves K-injective complexes. By Lemma E.1.0.11, the composition $Rp_{Y*} \text{for}_{X \times Y} : D(\text{Mod}(O_{X \times Y})_{0-\text{cxd}}) \rightarrow D(O_Y)$ is the right derived functor of

$$p_{Y*} \text{for}_{X \times Y} : \text{Mod}(O_{X \times Y})_{0-\text{cxd}} \rightarrow \text{Mod}(O_Y).$$

Since $\text{for}_Y \circ p_{Y*} = p_{Y*} \circ \text{for}_{X \times Y}$, the first square is indeed commutative.

By the commutativity of the first square and Corollary 5.3.1.16, the transform $R\mathfrak{S}_1$ preserves O -goodness. The other half about $R\mathfrak{S}_2$ is similar. \square

6.3.2 Rothstein's theorem

Theorem 6.3.2.1 (Rothstein). *There are natural isomorphisms $R\mathfrak{S}_1 R\mathfrak{S}_2 \cong T^{-g}$ on $D_{O-\text{good}}(\text{Mod}(O_Y)_{\text{cxd}})$ and $R\mathfrak{S}_2 R\mathfrak{S}_1 \cong T^{-g}$ on $D_{O-\text{good}}(\text{Mod}(O_X)_{\text{sp}})$.*

We begin the proof of Theorem 6.3.2.1 with Lemma 6.3.2.2, a direct adaption of [Rot97, Prop. 2.4] for complex tori.

Lemma 6.3.2.2. *Let $\Delta \subset X \times X$ be the diagonal. Define a morphism of complex tori $\epsilon_X : X \times X \rightarrow X$, $(x_1, x_2) \mapsto x_2 - x_1$. Then*

$$Rp_{12*}(\epsilon_X \times 1_Y)^* \mathcal{P} \cong O_{\Delta}[-g]$$

in $D^b(\text{Mod}(O_{X \times X})_{(1,-1)-\text{sp}})$, where $p_{12} : X \times X \times Y \rightarrow X \times X$ is the projection.

Proof. The identification $Rp_{X*}\mathcal{P} \cong \mathbb{C}_0[-g]$ in $D^b(O_X)$ from [Kem91, Thm. 3.15] can be lifted to an isomorphism in $D^b(\text{Mod}(O_X)_{-1-\text{sp}})$. As stated in the last sentence of the proof of [Vig21, Prop. 2.1.21], a morphism of modules with splittings (or connection) is an isomorphism whenever the underlying morphism of O -modules is so. Then apply Theorem 5.3.2.3 to the cartesian square

$$\begin{array}{ccc} X \times X \times Y & \xrightarrow{\epsilon_X \times 1_Y} & X \times Y \\ \downarrow p_{12} & & \downarrow p_X \\ X \times X & \xrightarrow{\epsilon_X} & X. \end{array}$$

□

Arguing as in Lemma 6.3.2.2, we can prove the analytic version of [Rot97, Prop. 2.5; Prop. 3.1]. These three results are used in the proof of Theorem 6.3.2.1 below.

Proof of Theorem 6.3.2.1. Repeat the proof of [Rot97, Thm. 3.2], which requires the projection formula and smooth base change theorem for modules with *connection*. For this, we first construct the corresponding comparison morphism that is compatible with the underlying O -module comparison morphism. The construction reduces to the adjunction between derived inverse image and derived direct image of relative connection in Remark 6.2.3.3.

The compatibility with O -module comparison morphism can be proved in a way similar to Proposition 6.3.1.2. On the level of O -modules, the comparison morphism is an isomorphism by Fact 5.3.2.15 and Theorem 5.3.2.3. (This type of arguments can also be found in the proof of [Vig21, Prop. 2.1.21; Thm. 2.1.33].) □

Remark 6.3.2.3. Rothstein's first proof ([Rot96, Thm. 2.2]) is based on a problematic lemma [Rot96, Lem. 2.3]. the problem is explained in [Rot97, Sec. 1]. To save his first proof, one may attempt to replace this "lemma" by its close variant, the bounded way-out lemma (see e.g., [Lip60, Lem. 1.11.3 (i)]). The difficulty is that, *a priori*, there is no canonical choice of a natural transformation between the two functors to be compared, which is required by way-out argument. For instance, in the end of the proof of Proposition 5.4.2.3, there are isomorphism arrows of opposite directions.

6.3.3 Matsushima's theorem

A holomorphic vector bundle $H \rightarrow Y$ is called homogeneous if T_y^*H is isomorphic to H for all $y \in Y$, where $T_y : Y \rightarrow Y$ is the translation by y . The first half of Theorem 6.3.3.1 is a special case of [Mat59, Thm. 1].

Theorem 6.3.3.1 (Matsushima). *Let E be a coherent \mathcal{O}_Y -module with a connection ∇ . Then E is a homogeneous vector bundle and the pair (E, ∇) is translation invariant.*

Proof. By Proposition 6.3.1.2, for every integer i , the coherent \mathcal{O}_X -module $H^i RS_2(E)$ admits a 1-splitting. By Lemma 6.3.3.2, the support of $H^i RS_2(E)$ is finite. Consequently, in $D_c^b(\mathcal{O}_X)$ there is an isomorphism $RS_2(E) \cong \bigoplus_{i \in \mathbb{Z}} T^{-i} H^i RS_2(E)$. From Proposition 5.5.3.2 3 and Fact 6.1.2.1 2, it induces an isomorphism in $D_c^b(\mathcal{O}_Y)$

$$T^{-g}E \rightarrow \bigoplus_{i \in \mathbb{Z}} T^{-i} H^0 RS_1(H^i RS_2(E)),$$

and each $H^0 RS_1(H^i RS_2(E))$ is a homogeneous vector bundle on Y . Then E is isomorphic to $H^0 RS_1(H^g RS_2(E))$, hence a homogeneous vector bundle.

We adopt the argument in [BK09, Footnote (6), p.388]. For every $y \in Y$, $T_y^* \nabla$ is a connection on $T_y^* E \xrightarrow{\sim} E$ and $T_0^* \nabla = \nabla$. The map

$$Y \rightarrow H^0(Y, \Omega_Y^1 \otimes \mathcal{E}nd(E)), \quad y \mapsto T_y^* \nabla - \nabla$$

is holomorphic. It is constantly 0 since Y is compact and $H^0(Y, \Omega_Y^1 \otimes \mathcal{E}nd(E))$ is a finite-dimensional vector space (Cartan-Serre's theorem). Hence $T_y^*(E, \nabla) = (E, \nabla)$ for all $y \in Y$. \square

Lemma 6.3.3.2 ([Rot96, Lem. 3.1]). *Let F be a coherent module with a 1-splitting on the complex torus X , then F is finitely supported.*

Proof. Suppose to the contrary that $\text{Supp}(F)$ is infinite. By [GR84, p.76], $\text{Supp}(F)$ is an analytic set in X . Then $\dim \text{Supp}(F) \geq 1$. Let C be an irreducible component of $\text{Supp}(F)$ of maximal dimension. Write $i : C \rightarrow X$ for the inclusion. Take a morphism $h : Z \rightarrow X$ provided by Lemma 5.5.3.3. Then $h(Z) = C$ and $F'' := F'/T(F')$ is a vector bundle on Z of positive rank r , where $F' = h^* F$ and $T(*)$ denotes the torsion part of a sheaf of modules. In consequence, the morphism of complex tori $h^* : \text{Pic}^0(X) \rightarrow \text{Pic}^0(Z)$ is nonzero. However, we claim that its tangent map at origin $h^* : \mathfrak{g} \rightarrow H^1(Z, \mathcal{O}_Z)$ is zero.

Let $\mathcal{E}' = h^* \mathcal{E}$. Because \mathcal{O}_X is flat over itself, pulling back (6.9) to Y and tensoring with F'' , by [Sta24, Tag 05NJ] we get a short exact sequence

$$0 \rightarrow \mathfrak{g}^* \otimes_{\mathbb{C}} F'' \rightarrow \mathcal{E}' \otimes_{\mathcal{O}_Z} F'' \rightarrow F'' \rightarrow 0. \quad (6.22)$$

Since \mathcal{E}' is a vector bundle on Z , one has

$$\frac{\mathcal{E}' \otimes F'}{T(\mathcal{E}' \otimes F')} = \mathcal{E}' \otimes F''.$$

Then the splitting on F induces a splitting $F'' \xrightarrow{\psi'} \mathcal{E}' \otimes F''$ of (6.22). Let β be the natural morphism $\beta : O_Z \rightarrow \mathcal{E}nd(F'')$. By Lemma 6.2.1.3, the composition

$$\mathrm{End}(g) \xrightarrow{\mathrm{Id}_{\mathfrak{g}^*} \otimes h^*} \mathfrak{g}^* \otimes_{\mathbb{C}} H^1(Z, O_Z) \xrightarrow{\mathrm{Id}_{\mathfrak{g}^*} \otimes H^1(Z, \beta)} \mathfrak{g}^* \otimes_{\mathbb{C}} H^1(Z, \mathcal{E}nd(F''))$$

sends $1 \in \mathrm{End}(\mathfrak{g})$ to 0. Therefore, the map $H^1(Z, \beta)h^* : \mathfrak{g} \rightarrow H^1(Z, \mathcal{E}nd(F''))$ is zero. Taking trace, we get a morphism $\tau : \mathcal{E}nd(F'') \rightarrow O_Z$ with $\tau\beta = r \cdot \mathrm{Id}_{O_Z}$. Then $h^* = \frac{1}{r}\tau_* H^1(Z, \beta)h^* = 0$ as a map $\mathfrak{g} \rightarrow H^1(Z, O_Z)$. The claim follows. The claim gives a contradiction. \square

Corollary 6.3.3.3. *Every local system (of finite dimensional \mathbb{C} -vector spaces) on a complex torus is translation invariant.*

Proof. Let L be a local system on Y . By Theorem 6.3.3.1, the pair $(L \otimes_{\mathbb{C}} O_Y, \mathrm{Id}_L \otimes d)$ is translation invariant. The result follows from the Riemann-Hilbert correspondence [Del70, I, Thm. 2.17]. \square

6.4 Laumon-Rothstein sheaf of algebras

6.4.1 Construction

To lift the Fourier-Mukai transform to D -modules, we recall (in Definition 6.4.1.1) the sheaf \mathcal{A}_X from [Rot96, p.576]. In the notation of (6.9), fix a \mathbb{C} -basis $\{\omega^1, \dots, \omega^g\}$ of the \mathbb{C} -vector space

$$H^0(Y, \Omega_Y^1) = \mathfrak{g}^* = \Gamma(X, \mathfrak{g}^* \otimes_{\mathbb{C}} O_X) \subset \Gamma(X, \mathcal{E}).$$

For each Stein open subset $U \subset X$, by Cartan's Theorem B (see, e.g., [KK83, Sec. 52, Thm. B]) one has $H^1(U, \mathfrak{g}^* \otimes_{\mathbb{C}} O_X) = 0$. Thence (6.9) induces a short exact sequence

$$0 \rightarrow \mathfrak{g}^* \otimes_{\mathbb{C}} O_X(U) \rightarrow \mathcal{E}(U) \xrightarrow{\mu} O_X(U) \rightarrow 0.$$

Whence, there is $\rho \in \mathcal{E}(U)$ with $\mu(\rho) = 1 \in O_X(U)$. For two such pairs (U, ρ) and $(\tilde{U}, \tilde{\rho})$ with $U \cap \tilde{U} \neq \emptyset$, one has $\mu(\tilde{\rho} - \rho) = 0 \in O_X(U \cap \tilde{U})$, so $\tilde{\rho} - \rho \in \mathfrak{g}^* \otimes_{\mathbb{C}} O_X(U \cap \tilde{U})$. There exists a unique tuple $f_1, \dots, f_g \in O_X(U \cap \tilde{U})$ such that

$$\tilde{\rho} - \rho = \sum_{i=1}^g \omega^i \otimes f_i$$

in $\mathcal{E}(U \cap \tilde{U})$.

Definition 6.4.1.1. For each chosen pair (U, ρ) as above, introduce independent variables $x_1^\rho, \dots, x_g^\rho$ and put

$$\mathcal{A}_X|_U = O_U[x_1^\rho, \dots, x_g^\rho].$$

For another choice $(\tilde{U}, \tilde{\rho})$ with the tuple (f_1, \dots, f_g) as above, we glue $\mathcal{A}_X|_U$ and $\mathcal{A}_X|_{\tilde{U}}$ by the rule

$$x_i^\rho - x_i^{\tilde{\rho}}|_{U \cap \tilde{U}} = f_i. \quad (6.23)$$

The resulting sheaf \mathcal{A}_X is a sheaf of commutative O_X -algebra.

Let

$$0 \rightarrow \mathfrak{g}^* \rightarrow X^\natural \xrightarrow{\pi} X \rightarrow 0 \quad (6.24)$$

be the universal vectorial extension² of X constructed in (F.22). In coordinate-free terms, \mathcal{A}_X is the O_X -subalgebra of $\pi_* O_{X^\natural}$ of sections whose restriction to each fiber of π is a polynomial on \mathfrak{g}^* . For every integer $m \geq 0$, let $O_{X^\natural}(m) \subset O_{X^\natural}$ denote the subsheaf of sections whose restriction to the fibers of π are homogeneous polynomials of degree m . Similar to [Bjö93, Def 1.6.1], there exists a sheaf of graded rings $O_{[X^\natural]} := \bigoplus_{m \geq 0} O_{X^\natural}(m) (\subset O_{X^\natural})$ on X^\natural . Then $\mathcal{A}_X = \pi_* O_{[X^\natural]}$ and $\Gamma(X, \mathcal{A}_X) = \mathbb{C}$.

Remark 6.4.1.2. Unlike the analytic case, if X is an abelian variety, then the notation \mathcal{A}_X in [Rot96, p.576] is the *algebraic* direct image $\pi_* O_{X^\natural}$. Morally, such difference also lies between algebraic and analytic D -modules. For a complex manifold or a smooth algebraic variety V , let $p : T^*V \rightarrow V$ be the natural projection of the cotangent bundle. Denote by GD_V the associated graded ring of the degree filtration on D_V . Then $GD_V = p_* O_{T^*V}$ in the algebraic case ([HT07, p.57]). By contrast, in the analytic case, GD_V is the O_V -submodule of $p_* O_{T^*V}$ of sections whose restriction to each fiber of p is a polynomial.

Remark 6.4.1.3. The sheaf of rings \mathcal{A}_X is functorial in X in the following sense. Let $\phi : X' \rightarrow X$ be a morphism of complex tori. Let $\hat{\phi} : Y \rightarrow Y'$ be the morphism dual to ϕ . By Proposition F.5.4.7, it induces a morphism $\phi^\natural : X'^\natural \rightarrow X^\natural$ of complex Lie groups fitting into a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^0(Y', \Omega_{Y'}^1) & \longrightarrow & X'^\natural & \xrightarrow{\pi'} & X' \longrightarrow 0 \\ & & \downarrow \hat{\phi}^* & & \downarrow \phi^\natural & & \downarrow \phi \\ 0 & \longrightarrow & H^0(Y, \Omega_Y^1) & \longrightarrow & X^\natural & \xrightarrow{\pi} & X \longrightarrow 0. \end{array}$$

For each local section of $O_{[X^\natural]}$, its ϕ^\natural -pullback (a local section of $O_{X'^\natural}$) restricts to a polynomial on each fiber of π' . Indeed, this restriction is the $\hat{\phi}^*$ -pullback of a restriction to a fiber of π . Therefore, the natural morphism $O_{X^\natural} \rightarrow \phi_*^\natural O_{X'^\natural}$ restricts to a morphism $O_{[X^\natural]} \rightarrow \phi_*^\natural O_{[X'^\natural]}$. The resulting morphism of ringed spaces $(X'^\natural, O_{[X'^\natural]}) \rightarrow (X^\natural, O_{[X^\natural]})$ descends to another morphism of ringed spaces

$$\tilde{\phi} : (X', \mathcal{A}_{X'}) \rightarrow (X, \mathcal{A}_X), \quad (6.25)$$

²By [Rot96, p.567], it is also the \mathfrak{g}^* -principal bundle associated to the tautological extension (6.9).

which is compatible with ϕ . In particular, the following square

$$\begin{array}{ccc} D(\mathcal{A}_{X'}) & \xrightarrow{R\tilde{\phi}_*} & D(\mathcal{A}_X) \\ \downarrow & & \downarrow \\ D(\mathcal{O}_{X'}) & \xrightarrow{R\phi_*} & D(\mathcal{O}_X) \end{array} \quad (6.26)$$

is commutative, where the vertical functors are forgetful. If M is an \mathcal{O}_X -module, then

$$L\tilde{\phi}^*(\mathcal{A}_X \otimes_{\mathcal{O}_X} M) = \mathcal{A}_{X'} \otimes_{\mathcal{O}_{X'}} L\phi^* M. \quad (6.27)$$

6.4.2 Basic properties

Notice that \mathcal{A}_X has a natural degree filtration $\{\mathcal{A}_X(m)\}_{m \in \mathbb{Z}}$, where

$$\mathcal{A}_X(m) = \pi_*(\oplus_{j=0}^m \mathcal{O}_{X^\sharp}(j))$$

is the \mathcal{O}_X -submodule of \mathcal{A}_X of polynomials of degree at most m . See also [Rot96, Sec. 5.3] and the end of [Lau96, p.10]. Then $\mathcal{A}_X(0) = \mathcal{O}_X$, $\mathcal{A}_X(1) = \mathcal{E}^\vee$ (cf. the start of [Lau96, p.10]), and every $\mathcal{A}_X(m)$ is a locally free \mathcal{O}_X -module of finite rank. Moreover, for any integers $m, n \geq 0$, one has

$$\mathcal{A}_X(n)\mathcal{A}_X(m) = \mathcal{A}_X(n+m). \quad (6.28)$$

Thus, \mathcal{A}_X is a sheaf of positively filtered rings (in the sense of [Bjö93, p.459; p.464]) on the complex torus X .

We review some terminology from [Bjö93, A:III]. A coherent sheaf of rings on a locally compact Hausdorff space is called *noetherian* if every increasing sequence of ideal sheaves is stationary over relatively compact subsets ([Bjö93, 2.24, p.470]). Let R be a commutative filtered ring. If the subring $\oplus_{v \in \mathbb{Z}} R_v T^v$ of $R[T, T^{-1}]$ is a noetherian ring, then R is called a *noetherian filtered ring*.

Definition 6.4.2.1 ([Bjö93, A:III, 1.7; Def. 1.11; 1.19]). A filtration on an R -module M is a family of additive subgroups $\{M_v\}_{v \in \mathbb{Z}}$ such that

$$M_v \subset M_{v+1}; \quad R_k M_v \subset M_{k+v}; \quad \cup_v M_v = M.$$

This filtration is called *separated* if $\cap_{v \in \mathbb{Z}} M_v = 0$, and called *good* if $\oplus_{v \in \mathbb{Z}} M_v T^v$ is a finitely generated $\oplus_{v \in \mathbb{Z}} R_v T^v$ -module.

A *zariskian filtered ring* is a noetherian filtered ring such that all the good filtrations on every finitely generated module are separated. A filtered sheaf of rings is called *stalkwise zariskian* if every stalk is a zariskian filtered ring ([Bjö93, Def. 2.6, p.465]).

Lemma 6.4.2.2. *The sheaf of rings \mathcal{A}_X is coherent and noetherian. The sheaf of filtered rings \mathcal{A}_X is stalkwise zariskian.*

Proof. By (6.23), the graded ring associated to the degree filtration of \mathcal{A}_X is

$$G\mathcal{A}_X := \oplus_{m \geq 0} \mathcal{A}_X(m) / \mathcal{A}_X(m-1) = \text{Sym}(\mathfrak{g}) \otimes_{\mathbb{C}} \mathcal{O}_X = \mathcal{O}_X[x_1, \dots, x_g]. \quad (6.29)$$

Here for each chosen pair (U, ρ) as above, $x_i|_U \in \Gamma(U, \mathcal{A}_X(1)/\mathcal{A}_X(0)) \subset \Gamma(U, G\mathcal{A}_X)$ is the image of $x_i^\rho \in \Gamma(U, \mathcal{A}_X(1))$. From [Bjö79, Thm. 1.26, p.460], \mathcal{A}_X is stalkwise zariskian. The other part follows from [Bjö79, Prop. 1.27, p.460; Thm. 2.7, p.465]. (See also the proof of [Bjö93, Thm. 1.2.5].) \square

In view of the difference mentioned in Remark 6.4.1.2, the statement of [Rot96, Prop. 4.4] is slightly modified as Fact 6.4.2.3. For every \mathcal{A}_X -module F and every chosen pair (U, ρ) as above, define $\psi_U^\rho : F(U) \rightarrow \mathcal{E}(U) \otimes_{\mathcal{O}_X(U)} F(U)$ by

$$\psi_U^\rho(s) = \rho \otimes s + \sum_{i=1}^g \omega^i|_U \otimes (x_i^\rho s).$$

Then $(\mu_1 \otimes \text{Id}_F)(\psi_U^\rho(s)) = s$. In light of (6.23), the family $\{\psi_U^\rho\}_{(U, \rho)}$ glue to a 1-splitting ψ on F . By the commutativity of \mathcal{A}_X and [Rot96, (4.9)], one has $[\psi, \psi] = 0$.

Fact 6.4.2.3. *The resulting functor $\text{Mod}(\mathcal{A}_X) \rightarrow \text{Mod}(\mathcal{O}_X)_{\text{sp}}$, $F \mapsto (F, \psi)$ induces an equivalence from $\text{Mod}(\mathcal{A}_X)$ to the full subcategory of $\text{Mod}(\mathcal{O}_X)_{\text{sp}}$ comprised of objects (F, ψ) with $[\psi, \psi] = 0$.*

From Fact 6.4.2.3 and the proof of [Rot96, Prop. 4.1], the functor (6.13) restricts to an exact functor $p_X^* : \text{Mod}(\mathcal{A}_X) \rightarrow \text{Mod}(\mathcal{O}_{X \times Y})_{1-\text{cxn}, \text{fl}}$. Similarly by [Rot96, Prop. 4.2], the functor (6.15) restricts to a functor

$$p_{X*} : \text{Mod}(\mathcal{O}_{X \times Y})_{1-\text{cxn}, \text{fl}} \rightarrow \text{Mod}(\mathcal{A}_X). \quad (6.30)$$

6.5 Laumon-Rothstein transform

6.5.1 Construction and properties

Definition 6.5.1.1. Define functors

$$RS_1 = R p_{Y*}(\mathcal{P} \otimes_{\mathcal{O}_{X \times Y}}^L p_X^* \cdot) : D(\mathcal{A}_X) \rightarrow D(D_Y); \quad (6.31)$$

$$RS_2 = R p_{X*}(\mathcal{P}^{-1} \otimes_{\mathcal{O}_{X \times Y}}^L p_Y^* \cdot) : D(D_Y) \rightarrow D(\mathcal{A}_X), \quad (6.32)$$

where $R p_{Y*} : D(\text{MIC}(p_X)) \rightarrow D(D_Y)$ (resp. $R p_{X*} : D(\text{Mod}(\mathcal{O}_{X \times Y})_{1-\text{cxn}, \text{fl}}) \rightarrow D(\mathcal{A}_X)$) is the right derived functor of (6.20) (resp. (6.30)), and $p_Y^* :$

$D(D_Y) \rightarrow D(D_{X \times Y/X})$ is from the exact functor (6.21). The pair is called the *Laumon-Rothstein transform*.

The situation is depicted below.

$$\begin{array}{ccc} \mathrm{Mod}(\mathcal{A}_X) & \xrightarrow{H^0 RS_1} & \mathrm{Mod}(D_Y) \\ \downarrow p_X^* & & \uparrow p_Y^* \\ \mathrm{Mod}(O_{X \times Y})_{1-\mathrm{cxn}, \mathrm{fl}} & \xrightarrow{\mathcal{P} \otimes \cdot} & \mathrm{Mod}(O_{X \times Y})_{0-\mathrm{cxn}, \mathrm{fl}}; \end{array} \quad \begin{array}{ccc} \mathrm{Mod}(D_Y) & \xrightarrow{H^0 RS_2} & \mathrm{Mod}(\mathcal{A}_X) \\ \downarrow p_Y^* & & \uparrow p_X^* \\ \mathrm{Mod}(O_{X \times Y})_{0-\mathrm{cxn}, \mathrm{fl}} & \xrightarrow{\mathcal{P}^{-1} \otimes \cdot} & \mathrm{Mod}(O_{X \times Y})_{1-\mathrm{cxn}, \mathrm{fl}}. \end{array}$$

Proposition 6.5.1.2. *There are commutative squares*

$$\begin{array}{ccc} D(\mathcal{A}_X) & \xrightarrow{RS_1} & D(D_Y) \\ \downarrow & & \downarrow \\ D(O_X) & \xrightarrow{RS_1} & D(O_Y); \end{array} \quad \begin{array}{ccc} D(D_Y) & \xrightarrow{RS_2} & D(\mathcal{A}_X) \\ \downarrow & & \downarrow \\ D(O_Y) & \xrightarrow{RS_2} & D(O_X), \end{array}$$

where the vertical functors are forgetful. In particular, RS_1 (resp. RS_2) sends $D_{O-\mathrm{good}}(\mathcal{A}_X)$ (resp. $D_{O-\mathrm{good}}(D_Y)$) to $D_{O-\mathrm{good}}(D_Y)$ (resp. $D_{O-\mathrm{good}}(\mathcal{A}_X)$).

Proof. The proof is similar to that of Proposition 6.3.1.2, as \mathcal{A}_X (resp. D_Y) is flat over O_X (resp. O_Y). \square

With Proposition 6.5.1.2, the proof of Theorem 6.5.1.3 is similar to that of Theorem 6.3.2.1.

Theorem 6.5.1.3 (Laumon, Rothstein). *There are natural isomorphisms of functors $RS_1 RS_2 \cong T^{-g}$ on $D_{O-\mathrm{good}}(D_Y)$ and $RS_2 RS_1 \cong T^{-g}$ on $D_{O-\mathrm{good}}(\mathcal{A}_X)$.*

Proposition 6.5.1.4 follows from Proposition 6.5.1.2, Theorem 6.5.1.3 and Fact 6.1.1.1 1 as in the proof of [Rot96, Thm. 6.1], cf. [Lau96, Prop. 3.1.2; Cor. 3.2.4].

Proposition 6.5.1.4. *There are natural isomorphisms of functors*

$$\begin{aligned} RS_2(D_Y \otimes_{O_Y}^L \cdot) &\cong \mathcal{A}_X \otimes_{O_X}^L RS_2(\cdot) : D_{\mathrm{good}}(O_Y) \rightarrow D_{O-\mathrm{good}}(\mathcal{A}_X); \\ RS_1(\mathcal{A}_X \otimes_{O_X}^L \cdot) &\cong D_Y \otimes_{O_Y}^L RS_1(\cdot) : D_{\mathrm{good}}(O_X) \rightarrow D_{O-\mathrm{good}}(D_Y). \end{aligned}$$

For $x \in X$ (resp. $y \in Y$), let $P_x = \mathcal{P}|_{x \times Y}$ (resp. $P_y = \mathcal{P}|_{X \times y}$) be the pullback line bundle on Y (resp. X). For a closed analytic subset S of a complex manifold Z , [Kas03, (3.30), p.51] defines a D_Z -module $\mathcal{B}_S|_Z$.

Corollary 6.5.1.5. *For any $x \in X$ and $y \in Y$, one has*

$$\begin{aligned} RS_2(D_Y \otimes_{O_Y} \mathbb{C}_y) &= \mathcal{A}_X \otimes_{O_X} P_{-y}; \\ T^g RS_1(\mathcal{A}_X \otimes_{O_X} P_{-y}) &= D_Y \otimes_{O_Y} \mathbb{C}_y = i_{y+} \mathbb{C} = \mathcal{B}_{\{y\}|Y}; \\ RS_1(\mathcal{A}_X \otimes_{O_X} \mathbb{C}_x) &= D_Y \otimes_{O_Y} P_x; \\ T^g RS_2(D_Y \otimes_{O_Y} P_x) &= \mathcal{A}_X \otimes_{O_X} \mathbb{C}_x. \end{aligned}$$

Proof. By [HT07, Example 1.6.4], one has $D_Y \otimes_{O_Y} \mathbb{C}_y = \mathcal{B}_{\{y\}|Y}$. The result follows from Theorem 6.5.1.3, Proposition 6.5.1.4, Fact 6.1.2.1 and Lemma 5.2.0.8. \square

6.5.2 Matsushima-Morimoto theorem

Proposition 6.5.2.1, due to Matsushima [Mat59, Thm. 1] and Morimoto [Mor59, Thm. 2], is a converse to Theorem 6.3.3.1. For abelian varieties, Nakayashiki [Nak94, Prop. 5.9] gives a proof using the Fourier-Mukai transform.

Proposition 6.5.2.1. *A homogeneous vector bundle on a complex torus admits an integrable connection.*

Proof. Let $E \rightarrow Y$ be a homogeneous vector bundle. Set $\hat{E} = H^q RS_2(E)$. According to Proposition 5.5.3.2 and Fact 6.1.1.1, one has $E = H^0 RS_1(\hat{E})$ and $\text{Supp}(\hat{E})$ is finite. By Lemma 6.5.2.2, \hat{E} has an \mathcal{A}_X -module structure. By Proposition 6.5.1.2, the O_Y -module underlying $H^0 RS_1(\hat{E})$ is E . The D_Y -module $H^0 RS_1(\hat{E})$ carries naturally an integrable connection. \square

The proof of Proposition 6.5.2.1 needs Lemma 6.5.2.2, a converse to Lemma 6.3.3.2.

Lemma 6.5.2.2. *If F is an O_X -module with finite support on the complex torus X , then F admits a 1-splitting ψ with $[\psi, \psi] = 0$.*

Proof. There is a decomposition $F = \bigoplus_{i=1}^m F_i$, where $\text{Supp}(F_i)$ is a singleton for each i . Thus, one may assume that $\text{Supp}(F)$ is a singleton. Then there exists an open neighborhood $U \subset X$ of $\text{Supp}(F)$ and a morphism of complex manifolds $s : U \rightarrow X^\natural$ that is a local section to (6.24). Let $\iota : U \rightarrow X$ be the inclusion. Applying π_* to the morphism of sheaves of rings $O_{X^\natural} \rightarrow s_* O_U$, one gets a morphism $\pi_* O_{X^\natural} \rightarrow \iota_* O_U$. As \mathcal{A}_X is an O_X -subalgebra of $\pi_* O_{X^\natural}$, this endows $\iota_* O_U$ an \mathcal{A}_X -module structure.³ Since the canonical O_X -morphism $\text{Id}_F \otimes \iota^\# : F \rightarrow F \otimes_{O_X} \iota_* O_U$ is an isomorphism, F also obtains an \mathcal{A}_X -module structure. This induces such a splitting by Fact 6.4.2.3. \square

Proposition 6.5.2.1, together with Theorem 6.3.3.1, yields (a slight generalization of) Morimoto's theorem [Mor59, Thm. 2, p.91].

Corollary 6.5.2.3 (Morimoto). *A coherent module admitting a connection on a complex torus is a vector bundle admitting an integrable connection.*

³This example shows that Lemma 6.3.3.2 fails without coherent condition.

6.6 Good modules

6.6.1 Definition

We define good \mathcal{A}_X -modules. We also review several definitions of good D -modules in the literature, and show that they are equivalent.

Let Z be a complex manifold.

Definition 6.6.1.1 ([Bjö93, 2.5, p.465]). Let \mathcal{R} be a positively filtered sheaf of rings on Z such that the associated graded ring $G\mathcal{R}$ is coherent. Let M be a coherent left \mathcal{R} -module. A filtration on M is an increasing sequence of subsheaves $\{M_v\}_{v \in \mathbb{Z}}$ satisfying $\cup_{v \in \mathbb{Z}} M_v = M$ and $\mathcal{R}_k M_v \subset M_{k+v}$ for all integers $k \geq 0$ and v . This filtration is called

- *B-good* ([Bjö93, Remark 2.16, p.467]) if for every $x \in Z$, there exists an open neighborhood U , a finite set $\{m_1, \dots, m_s\} \subset \Gamma(U, M)$ and integers k_1, \dots, k_s such that $M_v|_U = \sum_{i=1}^s \mathcal{R}_{v-k_i} m_i$ for all integers v .
- *locally good* ([Meb89, Prop. 2.1.12 (i)]) if every M_v is coherent over \mathcal{O}_Z , and if for every $x \in Z$, there is an open neighborhood U of x and an integer $k_0 \geq 0$ such that $\mathcal{R}_m M_{k_0} = M_{m+k_0}$ on U for all integers $m \geq 0$.

The proof of Lemma 6.6.1.2 is similar to that of [HT07, Prop. 2.1.1; Def. 2.1.2].

Lemma 6.6.1.2. *Let $M. = (M_v)_{v \in \mathbb{Z}}$ be a filtration on a coherent \mathcal{A}_X -module M . Then $M.$ is B-good if and only if $M.$ is locally good. (In that case, we call $M.$ a good filtration on M .)*

Proof. • Assume that $M.$ is B-good. By Lemma 6.4.2.2 and [Bjö93, Thm. 2.17, p.467], the $G\mathcal{A}_X$ -module $\oplus_{v \in \mathbb{Z}} M_v/M_{v-1}$ is coherent. Because of (6.29) and the proof of [Bjö93, Prop. 1.4.5], for every integer v , the \mathcal{O}_X -module M_v/M_{v-1} is coherent. From [Bjö93, Prop. 2.23, p.470], the filtration $M.$ is locally bounded blow. Then by induction on $v \in \mathbb{Z}$, one proves that the \mathcal{O}_X -module M_v is coherent.

For every $x \in X$, by definition, there is an open neighborhood $U \subset X$ of x , sections $m_1, \dots, m_s \in \Gamma(U, M)$ and integers k_1, \dots, k_s such that $M_v|_U = \sum_{i=1}^s \mathcal{A}_X(v - k_i) m_i$ for all integers v . Put $k_0 = \max_{j=1}^s k_j$. For every integer $k \geq 0$, one has $\mathcal{A}_X(k) M_{k_0} \subset M_{k+k_0}$. Moreover,

$$M_{k+k_0}|_U = \sum_{i=1}^s \mathcal{A}_X(k+k_0-k_i) m_i \stackrel{(a)}{\subset} \sum_{i=1}^s \mathcal{A}_X(k) \mathcal{A}_X(k_0-k_i) m_i \subset \mathcal{A}_X(k) M_{k_0},$$

where (a) uses (6.28). Hence $\mathcal{A}_X(k) M_{k_0} = M_{k+k_0}$ on U .

- Conversely, assume that M is locally good. For a fixed $x \in X$, take U and k_0 provided by the definition of local goodness. Since M_{k_0} is coherent over O_X , by shrinking U , one may assume that the O_U -module $M_{k_0}|_U$ is generated by sections $s_1, \dots, s_m \in \Gamma(U, M_{k_0})$. Define a morphism of \mathcal{A}_X -modules $\phi : \mathcal{A}_X^m|_U \rightarrow M|_U$, $(f_1, \dots, f_m) \mapsto \sum_{j=1}^m f_j s_j$. Since M is a filtration, for every integer v , one has $\mathcal{A}_X(v - k_0)M_{k_0} \subset M_v$. Hence $\phi(\mathcal{A}_X(v - k_0)^m) \subset M_v$. By construction, one has $\phi(\mathcal{A}_X(0)^m) = M_{k_0}|_U$. For every integer $k \geq k_0$, on U one has

$$M_k = \mathcal{A}_X(k - k_0)M_{k_0} = \mathcal{A}_X(k - k_0)\phi(\mathcal{A}_X(0)^m) \subset \phi(\mathcal{A}_X(k - k_0)^m).$$

Therefore, the filtration M is B-good. □

From [HT07, Thm. 2.1.3 (i)], a coherent D_V -module on a smooth algebraic variety V admits a globally defined good filtration. By contrast, Malgrange [Mal04, p.405] gives a coherent D -module on the complex manifold $\mathbb{C}^* \times \mathbb{CP}^1$ that does not admit any global good filtration.

Definition 6.6.1.3. An O_Z -module F is called

- *countably quasi-good* ([KS97, p.942]) if every compact subset of Z has an open neighborhood U such that $F|_U$ is the union of an increasing sequence of coherent O_U -submodules.
- *quasi-good* ([KS16, p.12]) if for every relatively compact open subset $U \subset Z$, the restriction $F|_U$ is a sum of coherent O_U -submodules.

A D_Z -module M is called

- *good coherent* if for every relatively compact open subset U of Z , there is a finite filtration $\{M_k\}_{k \in \mathbb{Z}}$ of $M|_U$ such that each quotient M_k/M_{k-1} is a coherent D_U -module admitting a good filtration. ([Sai89, p.369], [SS94, p.10] and [KS96, p.43].)
- *S-quasi-good* ([KS96, p.43]) if for every relatively compact open subset $U \subset Z$, the restriction $M|_U$ admits a filtration $\{M_v\}_{v \in \mathbb{Z}}$ by coherent D_U -submodule such that each quotient M_v/M_{v-1} admits a good filtration and $M_v = 0$ for $v \ll 0$.

Proposition 6.6.1.4. Let M be a coherent D_Z -module. Then the following are equivalent.

1. For every relatively compact open subset U of Z , there is a coherent O_U -submodule $F \subset M|_U$ with $D_U \cdot F = M|_U$.
2. For every relatively compact open subset U of Z , the D_U -module $M|_U$ admits a good filtration.

3. The D_Z -module M is good coherent.
4. The D_Z -module M is S -quasi-good.
5. The O_Z -module M is countably quasi-good.
6. The O_Z -module M is good.
7. The O_Z -module M is quasi-good.

Proof. We follow the circular chain.

1 implies 2 See [Bjö93, 1.4.10].

2 implies 3 For every relatively compact open subset U of Z , define a finite filtration of $M|_U$ by $M_0 = 0$ and $M_1 = M|_U$. Then the graded piece M_1/M_0 admits a good filtration over U .

3 implies 4 For every relatively compact open subset U of Z , consider the filtration $\{M_k\}$ in the definition. By induction on k , one proves that each M_k is D_U -coherent.

4 implies 5 Every quotient M_v/M_{v-1} admits a good filtration, then by [Bjö93, Cor. 1.4.6], it is countably quasi-good. By induction on v and using [KS97, Lem. 2.1.1], one proves that every M_v is countably quasi-good. Therefore, for every integer v , there is an increasing sequence $\{M_v^k\}_{k \geq 1}$ of coherent O_U -submodules of M_v with $M_v = \cup_{k \geq 1} M_v^k$. For every integer $k \geq 1$, let $M^k := \sum_{i \leq k, v \leq k} M_v^i$. By [Sta24, Tag 01BY], M^k is a coherent O_U -submodule of M_k . Then

$$M = \cup_{v \in \mathbb{Z}} M_v = \cup_{v \in \mathbb{Z}} \cup_{i \geq 1} M_v^i = \cup_{k \geq 1} M^k,$$

so M is countably quasi-good.

5 implies 6 An increasing sequence forms a directed family.

6 implies 7 By definition.

7 implies 1 Let U be a relatively compact open subset of Z . Because M is a finite type D_Z -module, for every $x \in \bar{U}$, there is a relatively compact open neighborhood $U(x) \subset Z$ of x , an integer $n(x) \geq 1$ and sections

$$\{s_i^x\}_{1 \leq i \leq n(x)} \subset \Gamma(U(x), M)$$

generating the $D_{U(x)}$ -module $M|_{U(x)}$. By compactness of \bar{U} , the open cover $\{U(x)\}_{x \in \bar{U}}$ of \bar{U} has a finite subcover $\{U(x_j)\}_{1 \leq j \leq r}$. Then $V = \cup_{j=1}^r U(x_j)$ is a relatively compact open subset of Z containing U . By

Condition 7, one may write $M|_V = \sum_{\alpha \in I} G_\alpha$, where I is an index set, and each G_α is a coherent O_V -submodule of $M|_V$.

For every $x \in \bar{U}$, there is an open neighborhood $V(x) \subset U(x)$ of x , such that for each $1 \leq i \leq n(x)$, the restriction $s_i^x|_{V(x)} \in \Gamma(V(x), G_{\alpha(x,i)})$ for some index $\alpha(x,i) \in I$. By compactness of \bar{U} again, the open cover $\{V(x)\}_{x \in \bar{U}}$ has a finite subcover $\{V(x'_k)\}_{1 \leq k \leq m}$. Then

$$F := \sum_{1 \leq k \leq m, 1 \leq i \leq n(x'_k)} G_{\alpha(x'_k, i)}$$

is a finite type O_V -submodule of $M|_V$. By Lemma 6.6.2.7, it is coherent over O_V . Moreover, $D_U \cdot F|_U = M|_U$.

□

The proof of Proposition 6.6.1.5 is similar to that of Proposition 6.6.1.4.

Proposition 6.6.1.5. *Let M be a coherent \mathcal{A}_X -module on the complex torus X . Then the O_X -module M is good if and only if there is a coherent O_X -submodule $F \subset M$ with $\mathcal{A}_X \cdot F = M$.*

Let the sheaf of rings \mathcal{R} be either D_Z or \mathcal{A}_X on the fixed complex torus X .

Definition 6.6.1.6. [Kas03, Def. 4.24] A coherent \mathcal{R} -module is *good* if the underlying O -module is good.

For example, by Lemma 6.4.2.2 and [Bjö93, Thm. 1.2.5], the left \mathcal{R} -module \mathcal{R} is good. Let $\text{Good}(\mathcal{R}) \subset \text{Coh}(\mathcal{R})$ (resp. $D_{\text{good}}^b(\mathcal{R}) \subset D_{O-\text{good}}^b(\mathcal{R})$) be the full subcategory of good \mathcal{R} -modules (resp. objects whose cohomologies are good \mathcal{R} -modules). By Proposition 6.6.1.4, the category $D_{\text{good}}^b(D_Z)$ is what Björk denotes by $D_{\text{coh}}^b(D_Z)_f$ in [Bjö93, p.119].

Fact 6.6.1.7 (GAGA).

- ([HK84, Thm. 1.1 (2)]) *Let V be a smooth proper complex algebraic variety. Then the analytification functor induces an equivalence $\text{Coh}(D_V) \rightarrow \text{Good}(D_{V^{\text{an}}})$.*
- *Let A be a complex abelian variety. Then the analytification functor induces an equivalence $\text{Coh}(\mathcal{A}_A) \rightarrow \text{Good}(\mathcal{A}_{A^{\text{an}}})$.*

A coherent D_Z -module is called *holonomic* if its characteristic variety is of (minimal) dimension $\dim Z$ ([Bjö93, Def. 3.1.1]). Malgrange ([Mal94, p.35], [Mal96, p.367], see also [Sab11, Thm. 4.3.4 (2)]) claims to have proved that every holonomic D_Z -module is generated by a coherent O_Z -submodule, so it is a good D_Z -module. Let $D_h^b(D_Z) \subset D^b(D_Z)$ be the full subcategory of objects with holonomic cohomologies.

6.6.2 Basic properties

Let \mathcal{R} be either D_Z on a complex manifold Z or \mathcal{A}_X on the fixed complex torus X .

Lemma 6.6.2.1 (Induced modules). *The functor $\mathcal{R} \otimes_{O_Z} \cdot : \text{Mod}(O_Z) \rightarrow \text{Mod}(\mathcal{R})$ is exact. It restricts to a functor $\mathcal{R} \otimes_{O_Z} \cdot : \text{Coh}(Z) \rightarrow \text{Good}(\mathcal{R})$, and induces a t-exact functor $\mathcal{R} \otimes_{O_Z}^L \cdot : D_c^b(O_Z) \rightarrow D_{\text{good}}^b(\mathcal{R})$.*

Proof. As \mathcal{R} is flat over O_Z , the functor is exact. Consider the degree filtration $\{\mathcal{R}(m)\}_{m \geq 0}$ of \mathcal{R} , where $\mathcal{R}(m) \subset \mathcal{R}$ is the O_Z -submodule of polynomials of degree at most m . Each $\mathcal{R}(m)$ is vector bundle on Z and $\mathcal{R} = \text{colim}_m \mathcal{R}(m)$. Therefore, the O -module \mathcal{R} is good. By Proposition 5.3.1.5 2, for every coherent O_Z -module F , the O -module $\mathcal{R} \otimes_{O_Z} F$ is good. Because F is an O_Z -module of finite presentation, $\mathcal{R} \otimes_{O_Z} F$ is an \mathcal{R} -module of finite presentation. Then it is \mathcal{R} -coherent by [Bjö93, Thm. 1.2.5] and Lemma 6.4.2.2. The other part follows. \square

Lemma 6.6.2.2. *The category $\text{Good}(\mathcal{R})$ is a weak Serre subcategory of $\text{Mod}(\mathcal{R})$. In particular, $D_{\text{good}}^b(\mathcal{R})$ is a triangulated subcategory of $D^b(\mathcal{R})$.*

Proof. The first half is a combination of [Kas03, Prop. 4.23], [Sta24, Tag 01BY] and [Sta24, Tag 0754]. The second half follows from [Yek19, Prop. 7.4.5]. \square

For a morphism of complex manifolds $f : M \rightarrow N$, the direct image of D -modules $f_+ : D(D_M) \rightarrow D(D_N)$ is constructed in [Bjö93, 2.3.12].

Fact 6.6.2.3 ([Bjö93, Thm. 2.8.1, 2.8.7]). *Let $f : W \rightarrow Z$ be a morphism of complex manifolds. For every $M \in D_{\text{good}}^b(D_W)$, if $f|_{\text{Supp}(M)} : \text{Supp}(M) \rightarrow Z$ is proper, then $f_+ M \in D_{\text{good}}^b(D_Z)$.*

Lemma 6.6.2.4. *Let $f : W \rightarrow Z$ be a proper morphism of complex manifolds. Then the direct image functor $f_+ : D(D_W) \rightarrow D(D_Z)$ restricts to a functor $D_{O\text{-good}}(D_W) \rightarrow D_{O\text{-good}}(D_Z)$.*

Proof. Take $M \in D_{O\text{-good}}(D_W)$. By [Sab11, Remark 3.3.4 (4)], the functor f_+ has finite cohomological dimension. So to prove $f_+ M \in D_{O\text{-good}}(D_Z)$, by [Har66, I, Prop. 7.3 (iii)], one may assume that $M \in \text{Mod}(D_W)$. Define a morphism $i : W \rightarrow W \times Z$, $w \mapsto (w, f(w))$, which is a closed embedding. Let $q : W \times Z \rightarrow Z$ be the projection. By [Sab11, Thm. 3.3.6 (1)], one has $f_+ = q_+ i_+$. The restriction $q|_W : W \rightarrow Z$ is proper. By [Bjö93, Prop. 2.4.8], one has $f_+ M = Rq_* DR_{W \times Z/Z}(i_+ M)[\dim Z]$. As each term of the (relative) de Rham complex $DR_{W \times Z/Z}(i_+ M)$ is $O_{W \times Z}$ -good and supported on W , by Theorem 5.3.1.7, $Rq_*[DR_{W \times Z/Z}(i_+ M)] \in D_{\text{good}}(O_Z)$. \square

For a closed embedding $i : M \rightarrow N$ of complex manifolds, the inverse image $i^* : \text{Mod}(D_N) \rightarrow \text{Mod}(D_M)$ may not preserve D -coherence ([HT07, Rk. 1.5.10]). For smooth morphisms, Fact 6.6.2.5 can be proved by applying [Kas03, Thm. 4.7] or repeating the proof of [HT07, Prop. 1.5.13 (ii)].

Fact 6.6.2.5. *Let $f : M \rightarrow N$ be a smooth morphism of complex manifolds. Then $Lf^* : D^b(D_N) \rightarrow D^b(D_M)$ restricts to functors $D_c^b(D_N) \rightarrow D_c^b(D_M)$ and $D_{\text{good}}^b(D_N) \rightarrow D_{\text{good}}^b(D_M)$.*

Lemma 6.6.2.6 concerns the local existence of good filtrations on coherent \mathcal{A}_X -modules.

Lemma 6.6.2.6. *Let M be a coherent \mathcal{A}_X -module on the complex torus X . For every $x \in X$, there is an open neighborhood U of x and a positive good filtration on $M|_U$.*

Proof. Let $\mathcal{A}_X^q|_U \xrightarrow{\phi} \mathcal{A}_X^p|_U \xrightarrow{\epsilon} M|_U \rightarrow 0$ be a local presentation of M on a relatively compact open neighborhood U of x . For every integer v , set $M_v = \epsilon(\mathcal{A}_X(v)^p)$, which is an O_U -submodule of $M|_U$. Then $M_v = 0$ when $v < 0$. Moreover, $\cup_{v \in \mathbb{Z}} M_v = M|_U$ and for any integers $m, k \geq 0$, one has $\mathcal{A}_X(m)M_k \subset M_{k+m}$. Thus, $\{M_v\}_{v \in \mathbb{Z}}$ is a positive filtration of $M|_U$. For every integer $k \geq 0$, one has $\mathcal{A}_X(k)M_0 = M_k$. It remains to prove that M_k is coherent over O_U .

We claim that $\phi(\mathcal{A}_X(m)^q) \cap \mathcal{A}_X(k)^p$ is coherent over O_U . In fact, for every $y \in U$, there is an integer $s \geq \max(0, k - m)$ such that $\phi(\mathcal{A}_X(m)^q) \subset \mathcal{A}_X(m + s)^p$ near y . In side the coherent O_X -module $\mathcal{A}_X(m + s)^p$, the two O_X -submodules $\phi(\mathcal{A}_X(m)^q)$ and $\mathcal{A}_X(k)^p$ are finite type. By [Sta24, Tag 01BY], their intersection $\phi(\mathcal{A}_X(m)^q) \cap \mathcal{A}_X(k)^p$ is coherent near y . The claim is proved.

Because $\mathcal{A}_X(k)^p$ is a noetherian O_X -module, the increasing sequence of submodules $\{\phi(\mathcal{A}_X(m)^q) \cap \mathcal{A}_X(k)^p\}_{m \geq 0}$ is stationary on U . Therefore, the union $\phi(\mathcal{A}_X^q) \cap \mathcal{A}_X(k)^p = \ker(\epsilon) \cap \mathcal{A}_X(k)^p$ is coherent over O_U . Since the sequence

$$0 \rightarrow \ker(\epsilon) \cap \mathcal{A}_X(k)^p \rightarrow \mathcal{A}_X(k)^p \rightarrow M_k|_U \rightarrow 0$$

is exact in $\text{Mod}(O_U)$, the restriction $M_k|_U$ is O_U -coherent. The constructed filtration is therefore good. \square

When $\mathcal{R} = D_Z$, Lemma 6.6.2.7 is [Sab11, Exercise E.2.4 (4)]. On a complex manifold Z , an O_Z -module F is *pseudo-coherent* if for every open subset U of X , every finite type O_U -submodule of $F|_U$ is of finite presentation ([Kas03, Def. A.5]).

Lemma 6.6.2.7. *If M is a coherent \mathcal{R} -module, then M is pseudo-coherent over O_Z .*

Proof. Let $F \subset M$ be a finite type O -submodule. For every point x , by [Meb89, Prop. 2.1.9] (in the case $\mathcal{R} = D_Z$) and Lemma 6.6.2.6 (in the case $\mathcal{R} = \mathcal{A}_X$), there exists an open neighborhood U of x and a good filtration on $M|_U$. By [Bjö93, Cor. 1.4.6] (in the case $\mathcal{R} = D_Z$) and Lemma 6.6.1.2 (in the case $\mathcal{R} = \mathcal{A}_X$), $M|_U$ is the sum of an increasing sequence of coherent O_U -submodules. Hence $M|_U$ is good over O_U . By Lemma A.1.4.2 1, the O_U -module $M|_U$ is pseudo-coherent. As pseudo-coherence is a local property, M is pseudo-coherent over O_Z . \square

Lemma 6.6.2.8. *Let M be a good \mathcal{R} -module. Let N be a finite type \mathcal{R} -submodule of M . Then N is good over \mathcal{R} .*

Proof. By [Sta24, Tag 01BY (1)], N is coherent over \mathcal{R} . For every relatively compact open subset U of X and every $x \in \bar{U}$, there is an open neighborhood $U(x) \subset X$ of x , an integer $n(x) > 0$ and sections $\{s_i(x)\}_{i=1}^{n(x)} \subset \Gamma(U(x), N)$ generating the $\mathcal{R}|_{U(x)}$ -module $N|_{U(x)}$. The open cover $\{U(x)\}_{x \in \bar{U}}$ of \bar{U} has a finite subcover $\{U(x_j)\}_{j=1}^m$. Let N_0 be the O_U -submodule of $N|_U$ generated by the finitely many local sections

$$\{s_i(x_j)\}_{1 \leq j \leq m, 1 \leq i \leq n(x_j)}.$$

Then N_0 is a finite type O_U -module. Because $M|_U$ is good over $\mathcal{R}|_U$, by Lemma 6.6.2.7, the O_U -module N_0 is coherent. By construction, one has $\mathcal{R}|_U \cdot N_0 = N|_U$. Therefore, the \mathcal{R} -module N is good by Propositions 6.6.1.4 (in the case $\mathcal{R} = D_Z$) and 6.6.1.5 (in the case $\mathcal{R} = \mathcal{A}_X$). \square

6.6.3 Preservation of goodness

Theorem 6.6.3.1. *The functor $RS_1 : D(\mathcal{A}_X) \rightarrow D(D_Y)$ restricts to an equivalence $D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(D_Y)$, with a quasi-inverse $T^g RS_2 : D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_X)$.*

Proof. 1. For every coherent O_Y -module F , one has $RS_2(D_Y \otimes_{O_Y}^L F) \in D_{\text{good}}^b(\mathcal{A}_X)$.

By Proposition 6.5.1.4, one has $RS_2(D_Y \otimes_{O_Y}^L F) = \mathcal{A}_X \otimes_{O_X}^L RS_2(F)$. By Fact 6.1.2.1 2, one has $RS_2(F) \in D_c^b(O_X)$. From Lemma 6.6.2.1, one gets $\mathcal{A}_X \otimes_{O_X}^L RS_2(F) \in D_{\text{good}}^b(\mathcal{A}_X)$.

2. For every $M \in \text{Good}(D_Y)$ and every integer i , the \mathcal{A}_X -module $H^i RS_2(M)$ is good.

Descending induction on $i \in \mathbb{Z}$. The O_X -module underlying $H^i RS_2(M)$ is $H^i RS_2(M)$. By Lemma 6.6.3.2, one has $H^i RS_2(M) = 0$ when $i > 2g$. In particular, $H^i RS_2(M)$ is good over \mathcal{A}_X .

Assume the statement for $i+1$. By Proposition 6.6.1.4, there is a coherent O_Y -submodule $F \subset M$ with $D_Y \cdot F = M$. Let M' be the kernel of the natural epimorphism $D_Y \otimes_{O_Y} F \rightarrow M$. Then

$$0 \rightarrow M' \rightarrow D_Y \otimes_{O_Y} F \rightarrow M \rightarrow 0 \quad (6.33)$$

is a short exact sequence in $\text{Mod}(D_Y)$. By Lemma 6.6.2.1, the D_Y -module $D_Y \otimes_{O_Y} F$ is good. By Lemma 6.6.2.2, so is M' . From (6.33), one gets an exact sequence in $\text{Mod}(\mathcal{A}_X)$

$$H^i RS_2(M') \rightarrow H^i RS_2(D_Y \otimes_{O_Y} F) \rightarrow H^i RS_2(M) \rightarrow H^{i+1} RS_2(M') \rightarrow H^{i+1} RS_2(D_Y \otimes_{O_Y} F). \quad (6.34)$$

By 1, the \mathcal{A}_X -module $H^j RS_2(D_Y \otimes_{O_Y} F)$ is good for $j \in \{i, i+1\}$. By the inductive hypothesis, so is $H^{i+1} RS_2(M')$.

Let $G = \ker[H^{i+1} RS_2(M') \rightarrow H^{i+1} RS_2(D_Y \otimes_{O_Y} F)]$. By Lemma 6.6.2.2, the \mathcal{A}_X -module G is good (hence of finite type). The sequence (6.34) yields an exact sequence

$$H^i RS_2(D_Y \otimes_{O_Y} F) \rightarrow H^i RS_2(M) \rightarrow G \rightarrow 0,$$

so $H^i RS_2(M)$ is a finite type \mathcal{A}_X -module for every coherent D_Y -module M . In particular, $H^i RS_2(M')$ is a finite type \mathcal{A}_X -module.

Let $N = \text{im}(H^i RS_2(M') \rightarrow H^i RS_2(D_Y \otimes_{O_Y} F))$. It is a finite type \mathcal{A}_X -submodule of the good \mathcal{A}_X -module $H^i RS_2(D_Y \otimes_{O_Y} F)$. By Lemma 6.6.2.8, the \mathcal{A}_X -module N is a good. The sequence (6.34) yields an exact sequence

$$0 \rightarrow N \rightarrow H^i RS_2(D_Y \otimes_{O_Y} F) \rightarrow H^i RS_2(M) \rightarrow H^{i+1} RS_2(M') \rightarrow H^{i+1} RS_2(D_Y \otimes_{O_Y} F).$$

By Lemma 6.6.2.2, the \mathcal{A}_X -module $H^i RS_2(M)$ is good. The induction is completed.

From 2, Lemma 6.6.2.2 and [Har66, I, Prop. 7.3 (i)], the functor RS_2 restricts to a functor $D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_X)$. Similarly, using Proposition 6.6.1.5, one can prove that RS_1 restricts to a functor $D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(D_Y)$. By Theorem 6.5.1.3, the restrictions are equivalences. \square

The proof of Theorem 6.6.3.1 needs a cohomological dimension estimation.

Lemma 6.6.3.2. *For an O_X -module F , we have $RS_1(F) \in D^{[0, 2g]}(O_Y)$. Similarly, for an O_Y -module G , we have $RS_2(G) \in D^{[0, 2g]}(O_X)$.*

Proof. By left exactness of the functor $p_{Y*} : \text{Mod}(O_{X \times Y}) \rightarrow \text{Mod}(O_Y)$, one has $R^i S_1(F) = 0$ for every integer $i < 0$. For every $y \in Y$, let M be the restriction (as sheaves) of $\mathcal{P} \otimes_{O_{X \times Y}} p_X^* F$ to $X \times y$. For every integer j , by the proper base change theorem (see e.g., [Mil13, Thm. 17.2]), one has $R^j S_1(F)_y = H^j(X \times y, M)$. When $j > 2g$, by [KS90, Prop. 3.2.2 (iv)], one has $H^j(X \times y, M) = 0$. Therefore, $R^j S_1(F) = 0$. The other part is similar. \square

6.7 Relations with other functors

The properties [Muk81, (3.1), (3.4), (3.8)] of the Fourier-Mukai transform have analogs for the Laumon-Rothstein transform.

6.7.1 Exchange of translation and multiplication

For every $y \in Y$, we view P_y as an object of $\text{Mod}(O_X)_{0-\text{sp}}$ via Example 6.2.1.2. There is a canonical isomorphism $T_{(0,y)}^* \mathcal{P} \cong \mathcal{P} \otimes_{O_{X \times Y}} p_X^* P_y$ in $\text{Mod}(X \times Y)_{-1-\text{cxn}}$, where $p_X^* : \text{Mod}(O_X)_{0-\text{sp}} \rightarrow \text{Mod}(O_{X \times Y})_{0-\text{cxn}}$ is defined in (6.13) and the functor

$$\mathcal{P} \otimes_{O_{X \times Y}} (\cdot) : \text{Mod}(O_{X \times Y})_{0-\text{cxn}} \rightarrow \text{Mod}(O_{X \times Y})_{-1-\text{cxn}}$$

is from [Rot97, (2.10)]. Arguing as in [Muk81, (3.1)], we get Proposition 6.7.1.1 from the projection formula.

Proposition 6.7.1.1.

$$\begin{aligned} RS_2 \circ T_y^* &\cong (\cdot \otimes_{O_X} P_y) \circ RS_2 : D(D_Y) \rightarrow D(\mathcal{A}_X); \\ RS_2 \circ (\cdot \otimes_{O_Y} P_x) &\cong T_{-x}^* \circ RS_2 : D(D_Y) \rightarrow D(\mathcal{A}_X); \\ RS_1 \circ (\cdot \otimes_{O_X} P_y) &\cong T_y^* \circ RS_1 : D(\mathcal{A}_X) \rightarrow D(D_Y); \\ RS_1 \circ T_x^* &\cong (\cdot \otimes_{O_Y} P_{-x}) \circ RS_1 : D(\mathcal{A}_X) \rightarrow D(D_Y). \end{aligned}$$

Similar results hold for $R\mathfrak{S}_1$ and $R\mathfrak{S}_2$.

Remark 6.7.1.2. Goodness over O is not necessary in Proposition 6.7.1.1, as the proof does not use the smooth base change.

6.7.2 Duality

Let Z be a complex manifold. Denote by Δ^{O_Z} the duality (contravariant) functor $R\mathcal{H}om_{O_Z}(\cdot, \omega_Z^{-1})[\dim Z] : D_c^b(O_Z) \rightarrow D_c^b(O_Z)$. The duality functor on D_Z -modules $\Delta^{D_Z} : D(D_Z) \rightarrow D(D_Z)$ is defined by $\Delta^{D_Z} F = G[\dim Z]$, where G is the complex of left D_Z -modules associated with the complex $R\mathcal{H}om_{D_Z}(F, D_Z)$ of right D_Z -modules. By [Bjö93, Def. 2.11.1], Δ^{D_Z} restricts to a functor $D_c^b(D_Z) \rightarrow D_c^b(D_Z)$, and the natural transformation $\text{Id} \rightarrow \Delta^{D_Z} \circ \Delta^{D_Z}$ is an isomorphism of functors $D_c^b(D_Z) \rightarrow D_c^b(D_Z)$.

Lemma 6.7.2.1 ([KS16, p.16]). *The functor $\Delta^{D_Z} : D(D_Z) \rightarrow D(D_Z)$ restricts to a functor $D_{\text{good}}^b(D_Z) \rightarrow D_{\text{good}}^b(D_Z)$.*

Proof. Suppose F is a coherent O_Z -module and $N = D_Z \otimes_{O_Z} F$, then by [Bjö93, (ii), p.122], there is $G \in D_c^b(O_Z)$ with $\Delta^{D_Z} N = D_Z \otimes_{O_Z} G$. By Lemma 6.6.2.1, $\Delta^{D_Z} N \in D_{\text{good}}^b(D_Z)$.

Take $M \in D_{\text{good}}^b(D_Z)$. To prove $\Delta^{D_Z} M \in D_{\text{good}}^b(D_Z)$, by [Har66, I, Prop. 7.3 (i)], one may assume $M \in \text{Good}(D_Z)$. For every relatively compact open subset $U \subset Z$, by [Bjö93, Thm. 1.5.8] and Proposition 6.6.1.4, there is a finite length exact sequence in $\text{Mod}(D_U)$:

$$0 \rightarrow D_U \otimes_{O_U} F^{-n} \rightarrow \cdots \rightarrow D_U \otimes_{O_U} F^0 \rightarrow M|_U \rightarrow 0,$$

where each F^i is a coherent O_U -module. For every i , one has $\Delta^{D_U}(D_U \otimes_{O_U} F^i) \in D_{\text{good}}^b(D_U)$. By Lemma 6.6.2.2, one has $(\Delta^{D_Z} M)|_U = \Delta^{D_U}(M|_U) \in D_{\text{good}}^b(D_U)$. Hence $\Delta^{D_Z} M \in D_{\text{good}}^b(D_Z)$. \square

For algebraic varieties, an analogue of Fact 6.7.2.2 is stated as [HT07, Cor. 2.6.8 (iii), Prop. 3.2.1]. From [HT07, p.101], all the arguments in [HT07, Sec. 2.6] are valid for analytic D -modules.

Fact 6.7.2.2.

1. The contravariant functor $\Delta^{D_Z} : D_h^b(D_Z) \rightarrow D_h^b(D_Z)$ an equivalence.
2. Let M be a coherent D_Z -module. Then M is holonomic if and only if $H^i(\Delta^{D_Z} M) = 0$ for all integers $i \neq 0$.
3. [Bjö93, Thm. 3.2.13 (3)] The bifunctor $-\otimes_{O_Z}^L + : D^b(D_Z) \times D^b(D_Z) \rightarrow D^b(D_Z)$ restricts to a bifunctor $D_h^b(D_Z) \times D_h^b(D_Z) \rightarrow D_h^b(D_Z)$.

Fact 6.7.2.3. Let $f : W \rightarrow Z$ be a morphism of complex manifolds. Then:

1. [Bjö93, Thm. 3.2.13 (1)] The inverse image $Lf^* : D^b(D_Z) \rightarrow D^b(D_W)$ restricts to a functor $D_h^b(D_Z) \rightarrow D_h^b(D_W)$.
2. [Sab11, Thm. 4.4.1] If $F \in D_h^b(D_W)$ is such that $f|_{\text{Supp}(F)}$ is proper, then $f_+ F \in D_h^b(D_Z)$.

Restricted to the complex torus Y , [Bjö93, (ii), p.122] becomes [Rot96, (6.12)]:

$$\Delta^{D_Y}(D_Y \otimes_{O_Y}^L \cdot) \cong D_Y \otimes_{O_Y}^L \Delta^{O_Y} : D_c^b(O_Y) \rightarrow D_c^b(D_Y).$$

Define the duality (contravariant) functor $\Delta^{\mathcal{A}_X} : D^b(\mathcal{A}_X) \rightarrow D^b(\mathcal{A}_X)$ as

$$\Delta^{\mathcal{A}_X} = T^g R\mathcal{H}om_{\mathcal{A}_X}(\cdot, \mathcal{A}_X).$$

It restricts to a functor $D_c^b(\mathcal{A}_X) \rightarrow D_c^b(\mathcal{A}_X)$. Similar to Lemma 6.7.2.1, it restricts to a functor $D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(\mathcal{A}_X)$. Theorem 6.7.2.4 follows from Proposition 6.7.2.5 and Fact 6.7.2.2 2, in the same way how Theorem 6.5 follows from Propositions 6.3 and 6.4 in [Rot96].

Theorem 6.7.2.4 (Rothstein). Let $F \in D_{\text{good}}^b(\mathcal{A}_X)$ be an object such that $RS_1(F)$ is concentrated in a single degree $i \in \mathbb{Z}$. Then $H^i RS_1(F)$ is holonomic if and only if $RS_1 \Delta^{\mathcal{A}_X} F$ is concentrated in degree $g - i$.

Proposition 6.7.2.5 can be deduced from Corollary 6.7.2.8, Proposition 6.5.1.4 and Proposition 5.5.1.8, in the same way that [Rot96, Prop. 6.3] is proved.

Proposition 6.7.2.5. *One has*

$$\begin{aligned} RS_2 \Delta^{D_Y} &= [-1]_X^* T^{-g} \Delta^{\mathcal{A}_X} RS_2 : D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_X), \\ \Delta^{D_Y} RS_1 &= [-1]_Y^* T^g RS_1 \Delta^{\mathcal{A}_X} : D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(D_Y). \end{aligned}$$

Remark 6.7.2.6. Both [Rot96, (6.13), (6.14)] miss a factor $[-1]^*$, due to a missing $[-1]_X^*$ in [Rot96, (6.15)]. Still, this sign does not affect the statement of [Rot96, Thm. 6.5].

Lemma 6.7.2.7 ([Huy06, (3.13)]). *For any objects $K, L \in D(O_Z)$ and $M \in D_c^-(O_Z)$, the natural morphism (provided by [Sta24, Tag 0BYS])*

$$K \otimes_{O_Z}^L R\mathcal{H}om_{O_Z}(M, L) \rightarrow R\mathcal{H}om_{O_Z}(M, K \otimes_{O_Z}^L L) \quad (6.35)$$

is an isomorphism in $D(O_Z)$.

Proof. By [Har66, I, Prop. 7.1 (ii)], one may assume that $M \in \text{Coh}(O_Z)$. By [Sta24, Tag 08DL] and [GH78, p.696], one may shrink Z such that M admits a globally free resolution $F \rightarrow M$, where the complex F is

$$0 \rightarrow O_Z^{k_n} \rightarrow \cdots \rightarrow O_Z^{k_1} \rightarrow O_Z^{k_0} \rightarrow 0$$

with $O_Z^{k_i}$ placed in degree $-i$. The morphism (6.35) becomes

$$K \otimes_{O_Z}^L \mathcal{H}om_{O_Z}(F, L) \rightarrow \mathcal{H}om_{O_Z}(F, K \otimes_{O_Z}^L L),$$

which is an isomorphism. \square

Corollary 6.7.2.8 proves the analytic counterpart of [Rot96, (6.12)].

Corollary 6.7.2.8. *There is a canonical isomorphism $\Delta^{\mathcal{A}_X}(\mathcal{A}_X \otimes_{O_X}^L \cdot) \cong \mathcal{A}_X \otimes_{O_X}^L \Delta^{O_X} \cdot$ of functors $D_c^b(O_X) \rightarrow D_c^b(\mathcal{A}_X)$.*

Proof. By [Rot96, (6.2)], one has

$$\Delta^{\mathcal{A}_X}(\mathcal{A}_X \otimes_{O_X}^L \cdot) = T^g R\mathcal{H}om_{\mathcal{A}_X}(\mathcal{A}_X \otimes_{O_X}^L \cdot, \mathcal{A}_X) = T^g R\mathcal{H}om_{O_X}(\cdot, \mathcal{A}_X).$$

By Lemma 6.7.2.7, it equals $T^g R\mathcal{H}om_{O_X}(\cdot, O_X) \otimes_{O_X}^L \mathcal{A}_X = \mathcal{A}_X \otimes_{O_X}^L \Delta^{O_X} \cdot$. \square

Example 6.7.2.9. Let $F = T^g \mathcal{A}_X \in D_{\text{good}}^b(\mathcal{A}_X)$. By Corollary 6.5.1.5, one has $RS_1(F) = D_Y \otimes_{O_Y} \mathbb{C}_0$. One has $\Delta^{\mathcal{A}_X} F = \mathcal{A}_X$, and $RS_1 \Delta^{\mathcal{A}_X} F$ is concentrated in degree g . Then by Theorem 6.7.2.4, the D_Y -module $D_Y \otimes_{O_Y} \mathbb{C}_0$ is holonomic.

Example 6.7.2.9 leads to a question: When is an induced D -module holonomic? A full answer is in Proposition E.2.0.1.

Remark 6.7.2.10. There is seemingly a paradox. Suppose $g = 1$ and let $i : 0 \rightarrow Y$ be the inclusion. Then the O_Y -modules pullback $i^*\mathbb{C}_0 = \mathbb{C}$ and $i^*(D_Y \otimes_{O_Y} \mathbb{C}_0) = (i^*D_Y) \otimes_{\mathbb{C}} (i^*\mathbb{C}_0) = i^*D_Y$ is the fiber of D_Y at 0, which is nonzero. On the other hand, by [Bjö93, p.87] the derived inverse image $i^+(D_Y \otimes_{O_Y} \mathbb{C}_0)$ is a complex of $D_0 = \mathbb{C}$ -modules concentrated in degree -1 . From [Bjö93, 2.3.7], its underlying complex of $O_0 = \mathbb{C}$ -modules is $Li^*(D_Y \otimes_{O_Y} \mathbb{C}_0)$. Its 0-th cohomology $i^*(D_Y \otimes_{O_Y} \mathbb{C}_0)$ is zero, a contradiction! We suggest catching the mistake.

In fact, D_Z has two different structures of O_Z -modules. Consider local sections h (resp. δ) of O_Z (resp. D_Z). One module structure defines $h \cdot \delta$ as $h\delta$, the product in D_Z . This O_Z -module is denoted by D_Z^l . Then $\text{for}_Z(D_Z) = D_Z^l$.

The other O_Z -module structure defines $h \cdot \delta$ as the reversed product δh in D_Z . Denote this O_Z -module by D_Z^r . Given an O_Z -module F , which one is used in the tensor product defining the induced left D_Z -module $D_Z \otimes_{O_Z} F$? In fact, it relies on the (D_Z, O_Z) -bimodule structure on D_Z^r . But $M := \text{for}_Z(D_Z \otimes_{O_Z} F)$ is NOT the O_Z -module tensor product of F with neither D_Z^r nor D_Z^l .

Return to the special case $F = \mathbb{C}_0 = O_Y/I$ on Y , where $I \subset O_Y$ is the coherent ideal sheaf corresponding to the closed embedding $i : 0 \rightarrow Y$. Take a local coordinate z around $0 \in Y$ such that $O_{Y,0} = \mathbb{C}\{z\}$ and the maximal ideal $m_0 = (z) \subset O_{Y,0}$. Let ∂ be the corresponding local vector field near $0 \in Y$. Let $M = \text{for}_Y(D_Y/D_Y I)$. The \mathbb{C} -vector space $M_0 = \mathbb{C}[\partial]$, and its $O_{Y,0}$ -action is defined by $z \cdot v = zv$ for all $v \in M_0$. The O_0 -module $\text{for}_0(H^0 i^+(D_Y \otimes_{O_Y} \mathbb{C}_0)) = i^*M = M_0/m_0 M_0$.

By $[\partial, z] = 1$, for every integer $k \geq 0$, one has $\partial^k = \frac{1}{k+1}(\partial^{k+1}z - z\partial^{k+1}) \in D_{Y,0}m_0 + m_0 D_{Y,0}$. So, $M_0/m_0 M_0 = 0$, even though the O -module pullback to 0 of both $D_Y^l \otimes_{O_Y} \mathbb{C}_0$ and $D_Y^r \otimes_{O_Y} \mathbb{C}_0$ are nonzero.

6.7.3 Pullback and pushout

Proposition 6.7.3.1 ([Lau96, Prop. 3.3.2]). *Let $f : X' \rightarrow X$ be a morphism of complex tori, with $\dim X' = g'$. Let $\hat{f} : Y \rightarrow Y'$ be the morphism dual to f . Let $\tilde{f} : (X', \mathcal{A}_{X'}) \rightarrow (X, \mathcal{A}_X)$ be the induced morphism (6.25). Then there are canonical isomorphisms of functors*

1.

$$L\hat{f}^*RS'_1 \cong RS_1R\tilde{f}_* : D_{O-\text{good}}(\mathcal{A}_{X'}) \rightarrow D_{O-\text{good}}(D_Y); \quad (6.36)$$

$$R\tilde{f}_*RS'_2 \cong T^{g-g'}RS_2L\hat{f}^* : D_{O-\text{good}}(D_{Y'}) \rightarrow D_{O-\text{good}}(\mathcal{A}_X). \quad (6.37)$$

2.

$$RS'_2 \hat{f}_+ \cong L\tilde{f}^* RS_2 : D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_{X'}); \quad (6.38)$$

$$\hat{f}_+ RS_1 \cong T^{g'-g} RS'_1 L\tilde{f}^* : D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(D_{Y'}). \quad (6.39)$$

Proof. 1. The isomorphism (6.37) follows from (6.36) as follows:

$$\begin{aligned} R\tilde{f}_* RS'_2 &\stackrel{(a)}{\cong} T^g RS_2 RS_1 R\tilde{f}_* RS'_2 \\ &\stackrel{(b)}{\cong} T^g RS_2 L\hat{f}^* RS'_1 RS'_2 \\ &\stackrel{(c)}{\cong} T^{g-g'} RS_2 L\hat{f}^*, \end{aligned}$$

where (6.36) (resp. Theorem 6.5.1.3) is used in (b) (resp. (a) and (c)). Then we prove (6.36).

By (6.26) (resp. the proof of [HT07, Prop. 1.5.8]), the derived direct image (resp. inverse image) functor of \mathcal{A} -modules (resp. D -modules) regards that of the underlying O -modules. From Proposition 5.3.1.2 2, the functor $\mathcal{P}' \otimes_{O_{X' \times Y'}}^L p_{X'}^* \cdot : D(\mathcal{A}_{X'}) \rightarrow D(O_{X' \times Y'})$ restricts to a functor $D_{O-\text{good}}(\mathcal{A}_{X'}) \rightarrow D_{\text{good}}(O_{X' \times Y'})$. An application of Lemma 5.3.2.13 to the cartesian square

$$\begin{array}{ccc} X' \times Y & \xrightarrow{1_{X'} \times \hat{f}} & X' \times Y' \\ p_2 \downarrow & \square & \downarrow p_{Y'} \\ Y & \xrightarrow{\hat{f}} & Y' \end{array}$$

yields a canonical isomorphism of functors

$$L\hat{f}^* R p_{Y'} \rightarrow R p_{2*} L(1_{X'} \times \hat{f})^* : D_{\text{good}}(O_{X' \times Y'}) \rightarrow D_{\text{good}}(O_Y). \quad (6.40)$$

Applying Theorem 5.3.2.3 to the cartesian square

$$\begin{array}{ccc} X' \times Y & \xrightarrow{p_1} & X' \\ f \times 1_Y \downarrow & \square & \downarrow f \\ X \times Y & \xrightarrow{p_X} & X, \end{array}$$

of complex manifolds, one gets a natural isomorphism

$$p_X^* R\tilde{f}_* \rightarrow R(f \times 1_Y)_* p_1^* \quad (6.41)$$

of functors $D_{O-\text{good}}(\mathcal{A}_{X'}) \rightarrow D(\text{Mod}(O_{X \times Y})_{1-\text{cxn}, \text{fl}})$.

Then

$$\begin{aligned}
L\hat{f}^*RS'_1 &= L\hat{f}^*Rp_{Y'}(\mathcal{P}' \otimes_{O_{X' \times Y'}}^L p_{X'}^* \cdot) \\
&\stackrel{(a)}{\cong} Rp_{2*}L(1_{X'} \times \hat{f})^*(\mathcal{P}' \otimes_{O_{X' \times Y'}}^L p_{X'}^* \cdot) \\
&\cong Rp_{2*}[L(1_{X'} \times \hat{f})^*\mathcal{P}' \otimes_{O_{X' \times Y'}}^L L(1_{X'} \times \hat{f})^*p_{X'}^* \cdot] \\
&\cong Rp_{2*}[(1_{X'} \times \hat{f})^*\mathcal{P}' \otimes_{O_{X' \times Y'}}^L p_1^* \cdot] \\
&\stackrel{(b)}{\cong} Rp_{2*}[(f \times 1_Y)^*\mathcal{P} \otimes_{O_{X' \times Y}}^L p_1^* \cdot] \\
&\cong Rp_{Y*}R(f \times 1_Y)_*[(f \times 1_Y)^*\mathcal{P} \otimes_{O_{X' \times Y}}^L p_1^* \cdot] \\
&\stackrel{(c)}{\cong} Rp_{Y*}[\mathcal{P} \otimes_{O_{X \times Y}}^L R(f \times 1_Y)_*p_1^* \cdot] \\
&\stackrel{(d)}{\cong} Rp_{Y*}[\mathcal{P} \otimes_{O_{X \times Y}}^L p_X^* R\tilde{f}_* \cdot] \\
&= RS_1 R\tilde{f}_*,
\end{aligned}$$

where (a), (b), (c) and (d)) use (6.40), (5.26), Fact 5.3.2.15 and (6.41) respectively. This proves (6.36).

2. The isomorphism (6.39) follows from (6.38) as follows:

$$\begin{aligned}
\hat{f}_+ RS_1 &\stackrel{(a)}{\cong} T^{g'} RS'_1 RS'_2 \hat{f}_+ RS_1 \\
&\stackrel{(b)}{\cong} T^{g'} RS'_1 L\tilde{f}^* RS_2 RS_1 \\
&\stackrel{(c)}{\cong} T^{g'-g} RS'_1 L\tilde{f}^*,
\end{aligned}$$

where (a) and (c) use Theorem 6.6.3.1, and (b) uses (6.38). Then we prove (6.38).

Using (6.27), one can prove that $L\tilde{f}^* : D(\mathcal{A}_X) \rightarrow D(\mathcal{A}_{X'})$ restricts to a functor $D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(\mathcal{A}_{X'})$. From Fact 6.6.2.3, the direct image functor $\hat{f}_+ : D^b(D_Y) \rightarrow D^b(D_{Y'})$ restricts to a functor $D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(D_{Y'})$. There are canonical isomorphism of

bifunctors $D_{\text{good}}^b(D_Y)^{\text{op}} \times D_{\text{good}}^b(\mathcal{A}_{X'}) \rightarrow \text{Ab}$:

$$\begin{aligned}
\text{Hom}_{D_{\text{good}}^b(\mathcal{A}_{X'})}(RS'_2\hat{f}_+- , +) &\stackrel{(a)}{\cong} \text{Hom}_{D_{\text{good}}^b(D_{Y'})}(\hat{f}_+- , T^{g'}RS'_1+) \\
&\stackrel{(b)}{\cong} \text{Hom}_{D_{\text{good}}^b(D_Y)}(- , T^g L\hat{f}^* RS'_1+) \\
&\stackrel{(c)}{\cong} \text{Hom}_{D_{\text{good}}^b(D_Y)}(- , T^g RS_1 R\tilde{f}_*+) \\
&\stackrel{(d)}{\cong} \text{Hom}_{D_{\text{good}}^b(\mathcal{A}_X)}(RS_2- , R\tilde{f}_*+) \\
&\cong \text{Hom}_{D_{\text{good}}^b(\mathcal{A}_{X'})}(L\tilde{f}^* RS_2- , +),
\end{aligned}$$

where (a) and (d) use Theorem 6.6.3.1, (b) uses [Bjö93, Thm. 2.11.8], and (c) uses (6.36). From Yoneda's lemma, there is a canonical isomorphism $RS'_2\hat{f}_+ \cong L\tilde{f}^* RS_2$ of functors $D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_{X'})$. \square

6.7.4 External tensor product

For two complex manifolds U, V , recall the (exact) external tensor product bifunctor

$$(\cdot) \boxtimes_O (\cdot) : \text{Mod}(D_U) \times \text{Mod}(D_V) \rightarrow \text{Mod}(D_{U \times V}) \quad (6.42)$$

defined in [Bjö93, 2.4.4]. By exactness, it descends to

$$D(D_U) \times D(D_V) \rightarrow D(D_{U \times V}). \quad (6.43)$$

Remark 6.7.4.1. By [Bjö93, 2.4.13], the bifunctor (6.42) restricts to bifunctors $\text{Coh}(D_U) \times \text{Coh}(D_V) \rightarrow \text{Coh}(D_{U \times V})$ and $\text{Good}(D_U) \times \text{Good}(D_V) \rightarrow \text{Good}(D_{U \times V})$. Then by [Har66, I, Prop. 7.3 (i)], the bifunctor (6.43) restricts to bifunctors $D_c^b(D_U) \times D_c^b(D_V) \rightarrow D_c^b(D_{U \times V})$ and $D_{\text{good}}^b(D_U) \times D_{\text{good}}^b(D_V) \rightarrow D_{\text{good}}^b(D_{U \times V})$. By [Bjö93, p.139], it also restricts to a bifunctor $D_h^b(D_U) \times D_h^b(D_V) \rightarrow D_h^b(D_{U \times V})$.

Using Lemma 5.5.1.6 (at the place of [HT07, Lem. 1.5.31]), Lemma 6.6.2.4 and [Sab11, Thm. 3.3.6 (1)], one can argue as in [HT07, Prop. 1.5.30] to get Fact 6.7.4.2.

Fact 6.7.4.2.

1. Let U, V, Z be complex manifolds. Let $f : U \rightarrow V$ be a proper morphism. Then the natural transformation

$$f_+(-) \boxtimes_O (+) \rightarrow (f \times \text{Id}_Z)_+(- \boxtimes_O +) : D_{O-\text{good}}(D_U) \times D(D_Z) \rightarrow D(D_{V \times Z})$$

is an isomorphism.

2. Let $f_i : U_i \rightarrow V_i$ ($i = 1, 2$) be two proper morphisms of complex manifolds. Then the natural transformation

$$(f_{1+} -) \boxtimes_O (f_{2+} +) \rightarrow (f_1 \times f_2)_+ (- \boxtimes_O +) : D_{O\text{-good}}(DU_1) \times D_{O\text{-good}}(DU_2) \rightarrow D_{O\text{-good}}(DV_1 \times V_2)$$

is an isomorphism.

For a complex torus X , let $\text{for}_X : \text{Mod}(\mathcal{A}_X) \rightarrow \text{Mod}(O_X)$ be the forgetful functor. Let X' be another complex torus. Set $X'' = X \times X'$. Write $u : X'' \rightarrow X$ and $u' : X'' \rightarrow X'$ for the projections. Let Y', Y'' be the dual of X' and X'' respectively. For an \mathcal{A}_X -module F and an $\mathcal{A}_{X'}$ -module G , denote $\tilde{u}^* F \otimes_{\mathcal{A}_{X''}} \tilde{u}'^* G$ by $F \boxtimes_{\mathcal{A}_X} G$. As

$$F \boxtimes_{\mathcal{A}_X} G = u^{-1} F \otimes_{u^{-1} \mathcal{A}_X} \mathcal{A}_{X''} \otimes_{u'^{-1} \mathcal{A}_{X'}} u'^{-1} G,$$

and $\mathcal{A}_{X''}$ is flat over $u^{-1} \mathcal{A}_X$ and over $u'^{-1} \mathcal{A}_{X'}$, the bifunctor

$$- \boxtimes_{\mathcal{A}_X} + : \text{Mod}(\mathcal{A}_X) \times \text{Mod}(\mathcal{A}_{X'}) \rightarrow \text{Mod}(\mathcal{A}_{X''})$$

is exact in both arguments. Consider the diagonal morphism $\delta : X \rightarrow X^2$. There is a canonical isomorphism of bifunctors

$$L\tilde{\delta}^*[- \boxtimes_{\mathcal{A}_X} +] \cong (-) \otimes_{\mathcal{A}_X}^L (+) : D(\mathcal{A}_X) \times D(\mathcal{A}_X) \rightarrow D(\mathcal{A}_X). \quad (6.44)$$

Although the tensor product of two \mathcal{A}_X -modules is different from the tensor product of the underlying O_X -module, Lemma 6.7.4.3 shows that external products do agree. It is used in the proof of Lemma 6.7.4.5.

Lemma 6.7.4.3. *There is a natural isomorphism of bifunctors*

$$\text{for}_{X''}(- \boxtimes_{\mathcal{A}} +) \rightarrow (\text{for}_X -) \boxtimes_O (\text{for}_{X'} +) : \text{Mod}(\mathcal{A}_X) \times \text{Mod}(\mathcal{A}_{X'}) \rightarrow \text{Mod}(O_{X''}).$$

Proof. By construction, one has

$$\mathcal{A}_{X''} = \mathcal{A}_X \boxtimes_O \mathcal{A}_{X'} = u^{-1} \mathcal{A}_X \otimes_{u^{-1} O_X} u'^* \mathcal{A}_{X'}. \quad (6.45)$$

There are natural isomorphisms of functors $\text{Mod}(\mathcal{A}_X) \rightarrow \text{Mod}(O_{X''})$:

$$\begin{aligned} \text{for}_{X''} \tilde{u}^* &:= u^{-1} \cdot \otimes_{u^{-1} \mathcal{A}_X} \mathcal{A}_{X''} \\ &\stackrel{(a)}{=} u^{-1} \cdot \otimes_{u^{-1} \mathcal{A}_X} (u^{-1} \mathcal{A}_X \otimes_{u^{-1} O_X} u'^* \mathcal{A}_{X'}) \\ &\cong u^{-1} \cdot \otimes_{u^{-1} O_X} u'^* \mathcal{A}_{X'} \\ &\cong (u^{-1} \cdot \otimes_{u^{-1} O_X} O_{X''}) \otimes_{O_{X''}} u'^* \mathcal{A}_{X'} \\ &\cong u^* \text{for}_X \cdot \otimes_{O_{X''}} u'^* \mathcal{A}_{X'}, \end{aligned}$$

where (a) uses (6.45). Similarly, there is a natural isomorphism of functors $\text{for}_{X''} \tilde{u}'^* \cong u^* \mathcal{A}_X \otimes_{O_{X''}} u'^* \text{for}_{X'} : \text{Mod}(\mathcal{A}_{X'}) \rightarrow \text{Mod}(O_{X''})$. One has natural isomorphisms of bifunctors

$$\begin{aligned} \text{for}_{X''}(- \boxtimes_{\mathcal{A}_X} +) &:= \tilde{u}^* - \otimes_{\mathcal{A}_{X''}} \tilde{u}'^* + \\ &\cong (u^* \text{for}_X - \otimes_{O_{X''}} u'^* \mathcal{A}_{X'}) \otimes_{u^* \mathcal{A}_X \otimes_{O_{X''}} u'^* \mathcal{A}_{X'}} (u^* \mathcal{A}_X \otimes_{O_{X''}} u'^* \text{for}_{X'} +) \\ &\cong (u^* \text{for}_X -) \otimes_{O_{X''}} (u'^* \text{for}_{X'} +) \\ &:= (\text{for}_X -) \boxtimes_O (\text{for}_{X'} +). \end{aligned}$$

□

Remark 6.7.4.4. We can reprove (6.45) as follows:

$$\begin{aligned} \mathcal{A}_X \boxtimes_O \mathcal{A}_{X'} &\stackrel{(a)}{=} \text{for}(RS_2(D_Y \otimes_{O_Y} \mathbb{C}_0)) \boxtimes_O \text{for}(RS'_2(D_{Y'} \otimes_{O_{Y'}} \mathbb{C}_0)) \\ &\stackrel{(b)}{=} RS_2(D_Y \otimes_{O_Y} \mathbb{C}_0) \boxtimes_O RS'_2(D_{Y'} \otimes_{O_{Y'}} \mathbb{C}_0) \\ &\stackrel{(c)}{=} RS''_2((D_Y \otimes_{O_Y} \mathbb{C}_0) \boxtimes_O (D_{Y'} \otimes_{O_{Y'}} \mathbb{C}_0)) \\ &= RS''_2((D_Y \boxtimes_O D_{Y'}) \otimes_{O_{Y''}} (\mathbb{C}_0 \boxtimes_O \mathbb{C}_0)) \\ &\stackrel{(d)}{=} RS''_2(D_{Y''} \otimes \mathbb{C}_0) \\ &\stackrel{(e)}{=} \text{for}(RS''_2(D_{Y''} \otimes \mathbb{C}_0)) \\ &\stackrel{(f)}{=} \mathcal{A}_{X''}, \end{aligned}$$

where (a) and (f) use Corollary 6.5.1.5, (b) and (e) use Proposition 6.5.1.2, and (c) (resp. (d)) uses Proposition 5.5.1.5 (resp. [Bjö93, 2.4.4, (i)]).

Lemma 6.7.4.5. *There are canonical isomorphisms of bifunctors*

$$RS''_2[- \boxtimes_O +] \cong RS_2 - \boxtimes_{\mathcal{A}} RS'_2 + : D_{O\text{-good}}(D_Y) \times D_{O\text{-good}}(D_{Y'}) \rightarrow D_{O\text{-good}}(\mathcal{A}_{X''}); \quad (6.46)$$

$$RS''_1[- \boxtimes_{\mathcal{A}} +] \cong RS_1 - \boxtimes_O RS'_1 + : D_{O\text{-good}}(\mathcal{A}_X) \times D_{O\text{-good}}(\mathcal{A}_{X'}) \rightarrow D_{O\text{-good}}(D_{Y''}). \quad (6.47)$$

Proof. It follows from Proposition 5.5.1.5, Lemma 6.7.4.3 and Proposition 6.5.1.2. □

6.7.5 Convolution and tensor product

For the dual complex tori X and Y , let $m : X^2 \rightarrow X$ and $\mu : Y^2 \rightarrow Y$ be their respective group law.

Definition 6.7.5.1 (Convolution, [Lau96, p.22]). Define bifunctors

$$\begin{aligned} *_D : D(D_Y) \times D(D_Y) &\rightarrow D(D_Y), & - *_D + &= \mu_+[- \boxtimes_O +], \\ *_\mathcal{A} : D(\mathcal{A}_X) \times D(\mathcal{A}_X) &\rightarrow D(\mathcal{A}_X), & - *_\mathcal{A} + &= R\tilde{m}_*[- \boxtimes_{\mathcal{A}} +]. \end{aligned}$$

As μ is proper, by Fact 6.6.2.3, Lemma 6.6.2.4 and Facts 6.7.2.2 and 6.7.2.3 2, the direct image μ_+ restricts to functors $D_{\text{good}}^b(D_{Y^2}) \rightarrow D_{\text{good}}^b(D_Y)$, $D_{O-\text{good}}(D_{Y^2}) \rightarrow D_{O-\text{good}}(D_Y)$ and $D_h^b(D_{Y^2}) \rightarrow D_h^b(D_Y)$. Together with Remark 6.7.4.1, this implies that the bifunctor $*_D$ restricts to bifunctors $D_{\text{good}}^b(D_Y) \times D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(D_Y)$, $D_{O-\text{good}}(D_Y) \times D_{O-\text{good}}(D_Y) \rightarrow D_{O-\text{good}}(D_Y)$ and $D_h^b(D_Y) \times D_h^b(D_Y) \rightarrow D_h^b(D_Y)$.

Lemma 6.7.5.2. *The pair $(D(D_Y), *_D)$ is a symmetric tensor triangulated category (in the sense of [Bal10, Def. 3]) with unit $D_Y \otimes_{O_Y} \mathbb{C}_0$.*

Proof. Let $i : \text{Specan}(\mathbb{C}) \rightarrow Y$ be the inclusion of $0 \in Y$. Then $D_Y \otimes_{O_Y} \mathbb{C}_0 = i_+ \mathbb{C}$. There are canonical isomorphisms

$$\begin{aligned} (i_+ \mathbb{C}) *_D \cdot &:= \mu_+[(i_+ \mathbb{C}) \boxtimes_O \cdot] \\ &= \mu_+[(i_+ \mathbb{C}) \boxtimes_O (\text{Id}_{Y^+} \cdot)] \\ &\stackrel{(a)}{\cong} \mu_+(i \times \text{Id}_Y)_+(\mathbb{C} \boxtimes_O \cdot) \\ &\stackrel{(b)}{\cong} \text{Id}_{Y^+} = \text{Id}_{D(D_Y)} \end{aligned}$$

of functors $D(D_Y) \rightarrow D(D_Y)$, where (a) and (b) use Fact 6.7.4.2 1 and [Sab11, Thm. 3.3.6 (1)] respectively. Therefore, $D_Y \otimes_{O_Y} \mathbb{C}_0$ is the unit. The other axioms can be verified as in [Wei07, pp. 10-11]. \square

Proposition 6.7.5.3 ([Wei11]). *For every $M \in D_{\text{good}}^b(D_Y)$, the functor $\cdot *_D M : D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(D_Y)$ admits a right adjoint $([-1]_Y^* \Delta^{D_Y} M) *_D \cdot$.*

Proof. Define an automorphism $f : Y^2 \rightarrow Y^2$ of the complex torus Y^2 by $f(a, b) = (a + b, -a)$. Then $p_1 f = \mu$, $p_2 f = [-1]_Y p_1$ and $\mu f = p_2$. One has $Lf^* O_{Y^2} = O_{Y^2}$ in $D^b(D_{Y^2})$.

For any objects $F, G \in D_{\text{good}}^b(D_Y)$, there are canonical bijections

$$\begin{aligned}
& \text{Hom}_{D_{\text{good}}^b(D_Y)}(F *_D M, G) := \text{Hom}_{D_{\text{good}}^b(D_Y)}(\mu_+(F \boxtimes_O M), G) \\
& \stackrel{(a)}{=} \text{Hom}_{D(D_{Y^2})}(F \boxtimes_O M, T^g \mu^* G) \\
& \stackrel{(b)}{=} \text{Hom}_{D(D_{Y^2})}(O_{Y^2}, \Delta^{D_{Y^2}}(F \boxtimes_O M) \otimes_{O_{Y^2}}^L T^g \mu^* G) \\
& \stackrel{(c)}{=} \text{Hom}_{D(D_{Y^2})}(O_{Y^2}, (\Delta^{D_Y} F) \boxtimes_O (\Delta^{D_Y} M) \otimes_{O_{Y^2}}^L T^g \mu^* G) \\
& = \text{Hom}_{D(D_{Y^2})}(O_{Y^2}, p_1^* \Delta^{D_Y} F \otimes_{O_{Y^2}}^L p_2^* \Delta^{D_Y} M \otimes_{O_{Y^2}}^L T^g \mu^* G) \\
& = \text{Hom}_{D(D_{Y^2})}(f^* O_{Y^2}, f^*[p_1^* \Delta^{D_Y} F \otimes_{O_{Y^2}}^L p_2^* \Delta^{D_Y} M \otimes_{O_{Y^2}}^L T^g \mu^* G]) \\
& = \text{Hom}_{D(D_{Y^2})}(O_{Y^2}, \mu^* \Delta^{D_Y} F \otimes_{O_{Y^2}}^L p_1^* [-1]_Y^* \Delta^{D_Y} M \otimes_{O_{Y^2}}^L T^g p_2^* G) \\
& = \text{Hom}_{D(D_{Y^2})}(O_{Y^2}, T^g \mu^* \Delta^{D_Y} F \otimes_{O_{Y^2}}^L ([-1]_Y^* \Delta^{D_Y} M \boxtimes_O G)) \\
& \stackrel{(d)}{=} \text{Hom}_{D(D_{Y^2})}(O_{Y^2}, T^g \Delta^{D_Y} (\mu^* F) \otimes_{O_{Y^2}}^L ([-1]_Y^* \Delta^{D_Y} M \boxtimes_O G)) \\
& \stackrel{(e)}{=} \text{Hom}_{D(D_{Y^2})}(\mu^* F, T^g ([-1]_Y^* \Delta^{D_Y} M \boxtimes_O G)) \\
& \stackrel{(f)}{=} \text{Hom}_{D(D_Y)}(F, \mu_+([-1]_Y^* \Delta^{D_Y} M \boxtimes_O G)) \\
& \stackrel{(g)}{=} \text{Hom}_{D_{\text{good}}^b(D_Y)}(F, ([-1]_Y^* \Delta^{D_Y} M) * G),
\end{aligned}$$

where (a), (c), (d), (f) and (g) use [Bjö93, Thm. 2.11.8], Proposition 6.7.5.4, [Kas03, Thm. 4.12], [Kas03, Thm. 4.40] and Lemma 6.7.2.1 in order, and both (b), (e) use [Kas03, (3.13)]. As the bijections are functorial in F and G , the adjunction follows. \square

The proof of Proposition 6.7.5.3 needs the commutativity of duality with external tensor product for D -modules.

Proposition 6.7.5.4. *Let Z_i ($i = 1, 2$) be two complex manifolds. Then there is a canonical isomorphism*

$$(\Delta^{D_{Z_1}} -) \boxtimes_O (\Delta^{D_{Z_2}} +) \rightarrow \Delta^{D_{Z_1 \times Z_2}}(- \boxtimes_O +) : D_c^b(D_{Z_1}) \times D_c^b(D_{Z_2}) \rightarrow D_c^b(D_{Z_1 \times Z_2})^{\text{op}}.$$

Proof. For a complex manifold Z , the sheaf $D_Z \otimes_{\mathbb{C}_Z} D_Z^{\text{op}}$ is naturally a \mathbb{C}_Z -algebra, and D_Z is naturally a left $D_Z \otimes_{\mathbb{C}_Z} D_Z^{\text{op}}$ -module. For $N_i \in D(D_{Z_i}^{\text{op}})$, by [HT07, p.39], there is a natural isomorphism in $D(D_{Z_1 \times Z_2}^{\text{op}})$:

$$N_1 \boxtimes_O N_2 = (N_1 \boxtimes_{\mathbb{C}} N_2) \otimes_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}} D_{Z_1 \times Z_2}. \quad (6.48)$$

First, we construct the natural transformation. Take $M_i \in D_c^b(D_{Z_i})$.

Claim 6.7.5.5. Then there is a natural morphism in $D^b((D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2})^{\text{op}})$:

$$\begin{aligned} & R\mathcal{H}om_{D_{Z_1}}(M_1, D_{Z_1}) \boxtimes_{\mathbb{C}} R\mathcal{H}om_{D_{Z_2}}(M_2, D_{Z_2}) \\ & \rightarrow R\mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}). \end{aligned} \quad (6.49)$$

Claim 6.7.5.6. There is a natural morphism in $D^b(D_{Z_1 \times Z_2}^{\text{op}})$:

$$\begin{aligned} & R\mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}) \otimes_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}} D_{Z_1 \times Z_2} \\ & \rightarrow R\mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, D_{Z_1 \times Z_2}). \end{aligned} \quad (6.50)$$

Again, there is a natural morphism in $D^b(D_{Z_1 \times Z_2}^{\text{op}})$:

$$R\mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, D_{Z_1 \times Z_2}) \rightarrow R\mathcal{H}om_{D_{Z_1 \times Z_2}}(M_1 \boxtimes_O M_2, D_{Z_1 \times Z_2}), \quad (6.51)$$

which can be defined by taking a $D_{Z_1 \times Z_2} \otimes_{\mathbb{C}} D_{Z_1 \times Z_2}^{\text{op}}$ -injective resolution of $D_{Z_1 \times Z_2}$.

Composing the morphisms (6.48), (6.49), (6.50) and (6.51) in order, one gets a natural morphism in $D^b(D_{Z_1 \times Z_2}^{\text{op}})$:

$$R\mathcal{H}om_{D_{Z_1}}(M_1, D_{Z_1}) \boxtimes_O R\mathcal{H}om_{D_{Z_2}}(M_2, D_{Z_2}) \rightarrow R\mathcal{H}om_{D_{Z_1 \times Z_2}}(M_1 \boxtimes_O M_2, D_{Z_1 \times Z_2}). \quad (6.52)$$

We prove that the constructed natural transformation is an isomorphism. To show (6.52) is an isomorphism, by [Har66, I, Prop. 7.1 (i)], one may assume $M_i \in \text{Coh}(D_{Z_i})$ for $i = 1, 2$. By shrinking Z_i and using [KS90, Prop. 11.2.6], one may find a bounded resolution of M_i by free D_{Z_i} -modules of finite rank. Thus, one may further assume that $M_i = D_{Z_i}$. Since $\omega_{Z_1 \times Z_2} = \omega_{Z_1} \boxtimes_O \omega_{Z_2}$ in $\text{Mod}(D_{Z_1 \times Z_2}^{\text{op}})$, by [HT07, Eg. 2.6.3], in this case (6.52) is an isomorphism. \square

Proof of Claim 6.7.5.5. Take a $D_{Z_i} \otimes_{\mathbb{C}} D_{Z_i}^{\text{op}}$ -injective resolution $D_{Z_i} \rightarrow I_i^*$. Then $I_1^* \boxtimes_{\mathbb{C}} I_2^*$ is a complex of modules over

$$(D_{Z_1} \otimes_{\mathbb{C}} D_{Z_1}^{\text{op}}) \boxtimes_{\mathbb{C}} (D_{Z_2} \otimes_{\mathbb{C}} D_{Z_2}^{\text{op}}) = (D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}) \otimes_{\mathbb{C}} (D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2})^{\text{op}}. \quad (6.53)$$

By [Sta24, Tag 013K (2)], there exists an injective resolution $I_1^* \boxtimes_{\mathbb{C}} I_2^* \rightarrow I^*$ (hence an induced injective resolution $D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2} \rightarrow I^*$) over (6.53). The natural morphism $D_{Z_i} \rightarrow D_{Z_i} \otimes_{\mathbb{C}} D_{Z_i}^{\text{op}}$ is flat, so every injective $D_{Z_i} \otimes_{\mathbb{C}} D_{Z_i}^{\text{op}}$ -module is injective over D_{Z_i} . Similarly, every term of the complex I^* is injective over $D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}$. Then (6.49) is defined to be the composition of the natural morphisms

$$\begin{aligned} & \mathcal{H}om_{D_{Z_1}}(M_1, I_1^*) \boxtimes_{\mathbb{C}} \mathcal{H}om_{D_{Z_2}}(M_2, I_2^*) \rightarrow \mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, I_1^* \boxtimes_{\mathbb{C}} I_2^*) \\ & \rightarrow \mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, I^*). \end{aligned}$$

\square

Proof of Claim 6.7.5.6. Take an injective resolution $D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2} \rightarrow J^*$ over (6.53). By [Sta24, Tag 013K (2)], over $(D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}) \otimes_{\mathbb{C}} D_{Z_1 \times Z_2}^{\text{op}}$ there exists an injective resolution $J^* \otimes_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}} D_{Z_1 \times Z_2} \rightarrow K^*$. Then (6.50) is defined to be the composition of the natural morphisms

$$\begin{aligned} & \mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, J^*) \otimes_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}} D_{Z_1 \times Z_2} \\ & \rightarrow \mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, J^* \otimes_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}} D_{Z_1 \times Z_2}) \\ & \rightarrow \mathcal{H}om_{D_{Z_1} \boxtimes_{\mathbb{C}} D_{Z_2}}(M_1 \boxtimes_{\mathbb{C}} M_2, K^*). \end{aligned}$$

□

Corollary 6.7.5.7 ([Lau96, Cor. 3.3.3]). *The equivalence $RS_2 : (D_{\text{good}}^b(D_Y), *_D) \rightarrow (D_{\text{good}}^b(\mathcal{A}_X), \otimes_{\mathcal{A}_X}^L)$ is a strong monoidal functor. In fact, there are canonical isomorphisms of bifunctors*

$$RS_2(- *_D +) \cong (RS_2-) \otimes_{\mathcal{A}_X}^L (RS_2+) : D_{\text{good}}^b(D_Y) \times D_{\text{good}}^b(D_Y) \rightarrow D_{\text{good}}^b(\mathcal{A}_X); \quad (6.54)$$

$$(RS_1-) *_D (RS_1+) \cong T^{-g} RS_1(- \otimes_{\mathcal{A}_X}^L +) : D_{\text{good}}^b(\mathcal{A}_X) \times D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(D_Y); \quad (6.55)$$

$$RS_1(- *_A +) \cong (RS_1-) \otimes_{O_Y}^L (RS_1+) : D_{O-\text{good}}(\mathcal{A}_X) \times D_{O-\text{good}}(\mathcal{A}_X) \rightarrow D_{O-\text{good}}(D_Y); \quad (6.56)$$

$$(RS_2-) *_A (RS_2+) \cong T^{-g} RS_2(- \otimes_{O_Y}^L +) : D_{O-\text{good}}(D_Y) \times D_{O-\text{good}}(D_Y) \rightarrow D_{O-\text{good}}(\mathcal{A}_X). \quad (6.57)$$

Proof. Let $\delta_X : X \rightarrow X^2 =: X'$ be the diagonal morphism. Its dual morphism is $\mu : Y^2 \rightarrow Y$. There are canonical isomorphisms of bifunctors

$$\begin{aligned} RS_2(- *_D +) &:= RS_2 \mu_+(- \boxtimes_O +) \\ &\stackrel{(a)}{\cong} L\tilde{\delta}_X^* RS_2'(- \boxtimes_O +) \\ &\stackrel{(b)}{\cong} L\tilde{\delta}_X^* (RS_2 - \boxtimes_A RS_2 +) \\ &\stackrel{(c)}{\cong} (RS_2-) \otimes_{\mathcal{A}_X}^L (RS_2+), \end{aligned}$$

where (a), (b) and (c) use (6.38), (6.46) and (6.44) respectively. This shows (6.54).

By Corollary 6.5.1.5, the functor RS_2 preserves units, so it is strong

monoidal. In addition, (6.55) follows:

$$\begin{aligned}
(RS_1 -) *_{\mathcal{D}} (RS_1 +) &\stackrel{(a)}{\cong} T^g RS_1 RS_2 (RS_1 - *_{\mathcal{D}} RS_1 +) \\
&\stackrel{(b)}{\cong} T^g RS_1 (RS_2 RS_1 - \otimes_{\mathcal{A}_X}^L RS_2 RS_1 +) \\
&\stackrel{(c)}{\cong} T^g RS_1 (T^{-g} - \otimes_{\mathcal{A}_X}^L T^{-g} +) \\
&= T^{-g} RS_1 (- \otimes_{\mathcal{A}_X}^L +),
\end{aligned}$$

where (a) and (c) (resp. (b)) use Theorem 6.6.3.1, (resp. (6.54)).

Because the diagonal morphism $\delta_Y : Y \rightarrow Y^2$ is dual to $m : X' = X^2 \rightarrow X$, there are canonical isomorphisms of bifunctors

$$\begin{aligned}
RS_1(- *_{\mathcal{A}} +) &:= RS_1 R\tilde{m}_*(- \boxtimes_{\mathcal{A}} +) \\
&\stackrel{(a)}{\cong} L\delta_Y^* RS_1'(- \boxtimes_{\mathcal{A}} +) \\
&\stackrel{(b)}{\cong} L\delta_Y^* (RS_1 - \boxtimes_O RS_1 +) \\
&\stackrel{(c)}{\cong} (RS_1 -) \otimes_{O_Y}^L (RS_1 +),
\end{aligned}$$

where (a), (b) and (c) use (6.36), (6.47) and [HT07, p.39] respectively. This demonstrates (6.56). Then (6.57) follows:

$$\begin{aligned}
(RS_2 -) *_{\mathcal{A}} (RS_2 +) &\stackrel{(a)}{\cong} T^g RS_2 RS_1 (RS_2 - *_{\mathcal{A}} RS_2 +) \\
&\stackrel{(b)}{\cong} T^g RS_2 (RS_1 RS_2 - \otimes_{O_Y}^L RS_1 RS_2 +) \\
&\stackrel{(c)}{\cong} T^g RS_2 (T^{-g} - \otimes_{O_Y}^L T^{-g} +) \\
&= T^{-g} RS_2 (- \otimes_{O_Y}^L +),
\end{aligned}$$

where (a) and (c) (resp. (b)) use Theorem 6.5.1.3 (resp. (6.56)). \square

Remark 6.7.5.8. We reprove Proposition 6.7.5.3 using the Laumon-Rothstein transform as follows. By [Sta24, Tag 08DJ], for every object $M \in D_{\text{good}}^b(\mathcal{A}_X)$, the functor $\cdot \otimes_{\mathcal{A}_X}^L M : D_{\text{good}}^b(\mathcal{A}_X) \rightarrow D_{\text{good}}^b(\mathcal{A}_X)$ admits a right adjoint $R\mathcal{H}om_{\mathcal{A}_X}(M, \cdot)$. By [Huy06, p.84], the right adjoint is naturally isomorphic to $T^{-g} \Delta^{\mathcal{A}_X}(M) \otimes_{\mathcal{A}_X}^L \cdot$. Then combining Proposition 6.7.2.5 with Corollary 6.7.5.7, one gets Proposition 6.7.5.3.

Appendix A

Sheaves of modules

A.1 Sheaves of modules

We recall some facts about sheaves of modules. Let (X, O_X) be a ringed space.

A.1.1 Generalities

Definition A.1.1.1. An O_X -module F is called

1. ([Sta24, Tag 01B5]) of *finite type* if every $x \in X$ admits an open neighborhood U such that $F|_U$ is generated by finitely many sections;
2. ([Sta24, Tag 01BN]) of *finite presentation* if for every $x \in X$, there is an open neighborhood $U \subset X$, integers $n, m \geq 0$ and an exact sequence of O_U -modules

$$O_U^m \rightarrow O_U^n \rightarrow F|_U \rightarrow 0;$$

3. ([Gro60, 5.1.3]) *quasi-coherent* if for every $x \in X$, there is an open neighborhood $U \subset X$, two sets I, J and a morphism $O_U^{\oplus J} \rightarrow O_U^{\oplus I}$ whose cokernel is isomorphic to $F|_U$;
4. ([Kas03, Def. A.5 (1)]) *pseudo-coherent* if for every open subset $U \subset X$, every finite type O_U -submodule of $F|_U$ is of finite presentation. Let $\text{PCoh}(X) \subset \text{Mod}(O_X)$ be full subcategory of pseudo-coherent modules;
5. ([Kas03, Def. A.5 (2)]) *K-coherent* if F is pseudo-coherent and of finite type;
6. ([Sta24, Tag 01BV]) *coherent* if F is of finite type and for every open subset $U \subset X$ and every finite collection $\{s_i\}_{1 \leq i \leq n}$ in $F(U)$, the kernel of the associated morphism $O_U^n \rightarrow F|_U$ is of finite type over O_U .

Every property in Definition A.1.1.1 is local, in the sense that it restricts to every open subset, and if it holds on each member of an open covering of X , then it holds on X .

Lemma A.1.1.2. *Let $0 \rightarrow F \xrightarrow{i} G \xrightarrow{r} H \rightarrow 0$ be a short exact sequence in $\text{Mod}(O_X)$. If F, H are of finite presentation, then so is G .*

Proof. For every $x \in X$, by [Sta24, Tag 01B8], there is an open neighborhood U of x such that the sequence $G(U) \xrightarrow{r_U} H(U) \rightarrow 0$ is exact. Up to shrinking U , there exist integers $m, n, p, q \geq 0$ and two exact sequences

$$O_U^m \rightarrow O_U^n \xrightarrow{f} F|_U \rightarrow 0, \quad O_U^p \rightarrow O_U^q \xrightarrow{h} H|_U \rightarrow 0.$$

The morphism h is defined by q elements s_1, \dots, s_q of $H(U)$. For each $1 \leq i \leq q$, choose a preimage $t_i \in G(U)$ of s_i . Consider the morphism $\phi : O_U^{n+q} \rightarrow G|_U$ determined by $if(e_1), \dots, if(e_n), t_1, \dots, t_q \in G(U)$. Hence a commutative diagram with two exact middle rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & O_U^m & \longrightarrow & \ker(\phi) & \longrightarrow & O_U^p \\ & & \downarrow & & \downarrow & \nearrow & \downarrow \\ 0 & \longrightarrow & O_U^n & \longrightarrow & O_U^{n+q} & \longrightarrow & O_U^q \longrightarrow 0 \\ & & \downarrow f & & \downarrow \phi & & \downarrow g \\ 0 & \longrightarrow & F|_U & \longrightarrow & G|_U & \longrightarrow & H|_U \longrightarrow 0 \\ & & \downarrow & & \downarrow & \nwarrow & \downarrow \\ & & 0 & \longrightarrow & \text{coker}(\phi) & \longrightarrow & 0. \end{array}$$

By the snake lemma, ϕ is surjective and $\ker(\phi)$ is finite type. Shrinking U again, one may find an integer $k \geq 0$ and a surjection $O_U^k \rightarrow \ker(\phi)$. The induced sequence $O_U^k \rightarrow O_U^{n+q} \rightarrow G|_U \rightarrow 0$ is exact. Therefore, G is of finite presentation. \square

A.1.2 Pseudo-coherent modules

Lemma A.1.2.1.

1. *Let $0 \rightarrow F \xrightarrow{i} G \xrightarrow{r} H \rightarrow 0$ be a short exact sequence in $\text{Mod}(O_X)$. If F, H are pseudo-coherent, then so is G .*
2. *Let I be a directed set. Let (M_i, f_{ij}) be a direct system over I consisting of pseudo-coherent O_X -modules. Then $M := \text{colim}_{i \in I} M_i$ in $\text{Mod}(O_X)$ is pseudo-coherent.*
3. *If $\{M_\alpha\}_{\alpha \in A}$ is a family of pseudo-coherent O_X -modules, then $S := \bigoplus_{\alpha \in A} M_\alpha$ is also pseudo-coherent.*

Proof. Let U be an open subset of X .

1. Let M be a finite type submodule of $G|_U$. Then the kernel of $r|_M : M \rightarrow H|_U$ is $(F|_U) \cap M$. Thus, $r|_M$ induces an injection $M/(F|_U \cap M) \rightarrow H|_U$. As H is pseudo-coherent, the finite type O_U -submodule $M/(F|_U \cap M)$ is of finite presentation. By [Sta24, Tag 01BP (2)], $F|_U \cap M$ is of finite type. As F is pseudo-coherent, $F|_U \cap M$ is of finite presentation. By Lemma A.1.1.2 applied to the exact sequence $0 \rightarrow F|_U \cap M \rightarrow M \rightarrow M/(F|_U \cap M) \rightarrow 0$, the O_U -module M is of finite presentation. Thus, G is pseudo-coherent.
2. Let N be a finite type submodule of $M|_U$. For every $x \in U$, from the first three lines of the proof of [Sta24, Tag 01BB], there is an open neighborhood $V \subset U$ of x and $i \in I$ such that $N|_V \subset F_i|_V$. Since F_i is pseudo-coherent, $N|_V$ is of finite presentation. As finite presentation is a local property, N is of finite presentation. Thus, M is pseudo-coherent.
3. Let I be the set of all finite subsets of A with the inclusion order. Then I is a directed set. For $B \in I$, set $F_B = \bigoplus_{\alpha \in B} M_\alpha$. By Point 1, F_B is pseudo-coherent. For $B \leq B'$ in I , set $f_{B,B'} : F_B \rightarrow F_{B'}$ to be the inclusion. Hence a direct system $(F_B, f_{B,B'})$ over I . By Point 2, the O_X -module $S = \operatorname{colim}_{B \in I} F_B$ is pseudo-coherent.

□

Lemma A.1.2.2 allows one to apply results on K-coherent sheaves in [Kas03] to coherent sheaves.

Lemma A.1.2.2. *An O_X -module is K-coherent if and only if it is coherent.*

Proof. Let $U \subset X$ be an open subset. Assume that F is a K-coherent module. Let $\{s_i\}_{1 \leq i \leq n}$ be a finite collection in $F(U)$, and let $f : O_U^n \rightarrow F|_U$ be the associated morphism. Then $\operatorname{im} f$ is a finite type submodule of $F|_U$. Because F is pseudo-coherent, $\operatorname{im} f$ is of finite presentation over O_U . From [Sta24, Tag 01BP (2)], $\ker f$ is of finite type over O_U . Therefore, F is coherent.

Conversely, assume that F is a coherent O_X -module. Let M be a finite type submodule of $F|_U$. By [Sta24, Tag 01BY (1)], M is coherent over O_U . From [Sta24, Tag 01BW], M is of finite presentation. Thus, F is pseudo-coherent and hence K-coherent. □

The module O_X is quasi-coherent, but in general not pseudo-coherent. If it is pseudo-coherent, then O_X is called a coherent sheaf of rings ([Kas03, p.214], [Bjö93, A:II, Def. 6.29]).

Lemma A.1.2.3. *If X is a locally Noetherian scheme, then every quasi-coherent module is pseudo-coherent.*

Proof. By [Gro60, Cor. 9.4.9], a quasi-coherent module is a directed limit of coherent modules, hence pseudo-coherent by Lemma A.1.2.1 2. \square

Example A.1.2.4. Let $X = \mathbf{A}^1$ be the affine line over a field. Let $U = X \setminus \{0\}$, and let $j : U \rightarrow X$ be the inclusion. By [Har77, II, Example 5.2.3], the \mathcal{O}_X -module $j_! \mathcal{O}_U$ is not quasi-coherent. From [Har77, II, Exercise 1.19 (c)], it is a submodule of the coherent module \mathcal{O}_X . Hence, $j_! \mathcal{O}_U$ is pseudo-coherent.

Definition A.1.2.5 defines a local property. It is weaker than [Bjö93, A:III, 2.24] and [Kas03, Def. A.7].

Definition A.1.2.5. Assume that \mathcal{O}_X is a coherent sheaf of rings. If for every open subset $U \subset X$, every family of coherent ideal sheaves $\{I_i\}_i$ in \mathcal{O}_U , the ideal sheaf $\sum_i I_i$ is \mathcal{O}_U -coherent, then \mathcal{O}_X is called a quasi-Noetherian sheaf of rings.

Example A.1.2.6. 1. If (X, \mathcal{O}_X) is a locally Noetherian scheme, then \mathcal{O}_X is quasi-Noetherian.

2. If (X, \mathcal{O}_X) is a complex analytic space, then by the Oka-Cartan theorem (see, e.g., [Kas03, Thm. A.12]), \mathcal{O}_X is quasi-Noetherian.

A.1.3 Analytic coherent modules

Let X be a complex analytic space. We show that a coherent \mathcal{O}_X -module admits a local free resolution, from which we deduce that coherence is preserved by derived pullbacks and tensor products. An analog of Lemma A.1.3.1 for algebraic varieties is [Har77, III, Example 6.5.1]. By local syzygies [GH78, p.696], on complex manifolds, every coherent module local admits a finite-length, finite free resolution.

Lemma A.1.3.1. *Every $x \in X$ admits an open neighborhood U , such that for every coherent \mathcal{O}_X -module F , there is a (possibly infinite-length) resolution*

$$\cdots \rightarrow \mathcal{O}_U^{n_1} \rightarrow \mathcal{O}_U^{n_0} \rightarrow F|_U \rightarrow 0,$$

where $n_i \geq 0$ are integers.

Proof. Shrinking X to an open neighborhood of x , one may assume that X is Stein. By [GR04, Thm. 8, p.108], there is a compact neighborhood $K \subset X$ of x , such that Theorem B is valid on K in the sense of [GR04, Def. 1, p.100]. Let $U = K^\circ$.

For a coherent \mathcal{O}_X -module F , we construct inductively a sequence of morphisms. From [GR04, Cor. p.101], there is an integer $n_0 \geq 0$, an open neighborhood U_0 of $K \subset X$ and a morphism $f_0 : \mathcal{O}_{U_0}^{n_0} \rightarrow F|_{U_0}$ in $\text{Mod}(\mathcal{O}_{U_0})$ such that $f_0|_U$ is an epimorphism in $\text{Mod}(\mathcal{O}_U)$. Set $\ker(f_{-1})|_{U_0} = F|_{U_0}$.

Given such a morphism $f_j : O_{U_j}^{n_j} \rightarrow \ker(f_{j-1})|_{U_j}$ for an integer $j \geq 0$ and an open neighborhood $U_j \subset X$ of K , by [Sta24, Tag 01BY (3)], the O_{U_j} -module $\ker(f_j)$ is coherent. By [GR04, Cor. p.101], there is an open neighborhood $U_{j+1} \subset U_j$ of K , an integer $n_{j+1} \geq 0$ and a morphism $f_{j+1} : O_{U_{j+1}}^{n_{j+1}} \rightarrow \ker(f_j)|_{U_{j+1}}$ in $\text{Mod}(O_{U_{j+1}})$ such that $f_{j+1}|_U$ is an epimorphism. Thus, one gets a sequence

$$\cdots \rightarrow O_U^{n_2} \xrightarrow{f_2|_U} O_U^{n_1} \xrightarrow{f_1|_U} O_U^{n_0} \xrightarrow{f_0|_U} F|_U \rightarrow 0$$

in $\text{Mod}(O_U)$. By construction, it is exact, hence a resolution of $F|_U$. \square

Example A.1.3.2. Assume that $x \in X$ is a singular point. Then $F := \mathbb{C}_x$ is a coherent O_X -module, but for every open neighborhood $U \subset X$ of x , there is no finite-length resolution of $F|_U$ by finite locally free O_U -modules. (Otherwise, such a resolution induces a finite-length free resolution of the $O_{X,x}$ -module $F_x = \mathbb{C} = O_{X,x}/m_x$. From [Osb12, Ch. 4, Prop. 4.4], the projective dimension $\text{pd}_{O_{X,x}} O_{X,x}/m_x$ is finite. By [Mat87, Lem. 1, p.154] and [Osb12, Prop. 4.9], the global dimension of the ring $O_{X,x}$ is finite. By Serre's theorem (see, e.g., [Osb12, p.332]), the local ring $O_{X,x}$ is regular. From [Ser56, p.6], x is a smooth point of X , a contradiction.)

Therefore, Lemma A.1.3.1 fails if one consider only finite-length resolutions. See also [EP⁺96, Thm. 4.1.2].

Lemma A.1.3.3. *Let $f : X \rightarrow Y$ be a morphism of complex analytic spaces. Then derived pullback $Lf^* : D(Y) \rightarrow D(X)$ restricts to a functor $\text{Coh}(Y) \rightarrow D_c(X)$.*

Proof. Let F be a coherent O_Y -module. For every $x \in X$, by Lemma A.1.3.1, there is an open neighborhood V of $f(x) \in Y$, such that there is a resolution $E_\bullet \rightarrow F|_V \rightarrow 0$ by finite free O_V -modules. Let $g : f^{-1}(V) \rightarrow V$ be the base change of f along the inclusion $V \rightarrow Y$. Then the morphism $g^*E_\bullet \rightarrow (Lf^*F)|_{f^{-1}(V)}$ in $D(f^{-1}(V))$ is an isomorphism. For every integer $j \geq 0$, the $O_{f^{-1}(V)}$ -module g^*E_j is finite free. Thus, the $O_{f^{-1}(V)}$ -module $(H^{-j}Lf^*F)|_{f^{-1}(V)}$ is coherent. Since coherence is a local property, the O_X -module $H^{-j}(Lf^*F)$ is coherent. \square

Lemma A.1.3.4. *For any coherent O_X -modules F and G , one has $F \otimes_{O_X}^L G \in D_c(X)$.*

Proof. For every $x \in X$, by Lemma A.1.3.1, there is an open neighborhood $U \subset X$ of x and a resolution $E_\bullet \rightarrow F|_U \rightarrow 0$ by finite free O_U -modules. The natural morphism $E_\bullet \otimes_{O_U} G|_U \rightarrow F|_U \otimes_{O_U}^L G|_U$ in $D(U)$ is an isomorphism. For every integer n , the O_U -module $H^n(E_\bullet \otimes_{O_U}^L G|_U) = H^n(E_\bullet \otimes_{O_U} G|_U)$ is coherent. Therefore, the O_U -module $H^n(F \otimes_{O_X}^L G)|_U = H^n(F|_U \otimes_{O_U}^L G|_U)$ is coherent. Since coherence is a local property, the O_X -module $H^n(F \otimes_{O_X}^L G)$ is coherent. \square

A.1.4 Good modules

Assume that the ringed space X is locally compact Hausdorff.

Definition A.1.4.1. [Kas03, Def. 4.22] An O_X -module F is called *good* if for every relatively compact open subset $U \subset X$, there exists a directed family $\{G_i\}_{i \in I}$ of coherent O_U -submodules of $F|_U$ such that $F|_U = \sum_{i \in I} G_i$, where $\{G_i\}_{i \in I}$ being a directed family means that for any $i, i' \in I$, there is $i'' \in I$ with $G_i + G_{i'} \subset G_{i''}$ (and hence $F|_U = \operatorname{colim}_{i \in I} G_i$). The full subcategory of $\operatorname{Mod}(O_X)$ consisting of good O_X -modules is denoted by $\operatorname{Good}(X)$.

Lemma A.1.4.2 (Goodness vs. pseudo-coherence).

1. ([Kas03, p.77]) One has $\operatorname{Coh}(X) \subset \operatorname{Good}(X) \subset \operatorname{PCoh}(X)$.
2. Let E be a pseudo-coherent O_X -module. If on every relatively compact open subset $U \subset X$, the O_U -module $E|_U$ is the sum of its finite type submodules, then E is good.

Proof.

1. By definition, every coherent O_X -module is good. Let E be a good O_X -module. Let W be an open subset of X , and let $F \subset E|_W$ be a finite type O_W -submodule. We show that F is of finite presentation over O_W . Replacing (X, E) with $(W, E|_W)$, one may assume that $W = X$. Because X is locally compact, for every $x \in X$, there exists a *relatively compact* open neighborhood $U \subset X$ of x and finitely many sections $s_1, \dots, s_n \in F(U)$ generating $F|_U$. As E is good, $E|_U = \sum_{i \in I} G_i$ is the sum of a directed family of coherent submodules. There exists $i_0 \in I$ and an open neighborhood V of $x \in U$ with $s_i|_V \in G_{i_0}(V)$ for all $1 \leq i \leq n$. Then $F|_V$ is a finite type submodule of $G_{i_0}|_V$. By [Sta24, Tag 01BY (1)], $F|_V$ is O_V -coherent. As coherence is a local property, F is coherent. From [Sta24, Tag 01BW], F is of finite presentation.
2. The family of finite type submodules of $E|_U$ is directed, since the sum of two finite type submodules is of finite type. For every relatively compact open subset $U \subset X$, as E is pseudo-coherent, every finite type submodule of $E|_U$ is pseudo-coherent and hence coherent. Thus, E is good.

□

Basic properties of good modules (similar to those of quasi-coherent modules on algebraic varieties) are recapped in Lemma A.1.4.3. Point 3 should be compared to [Con06, Lemma 2.1.8 (1)].

Lemma A.1.4.3.

1. For every family of objects $\{F_i\}_{i \in I}$ in $\text{Good}(X)$, the direct sum $\bigoplus_{i \in I} F_i$ in $\text{Mod}(O_X)$ is good.
2. The subcategory $D_{\text{good}}(X)$ is closed under direct sums in $D(X)$. Moreover, the inclusion functor $\text{Good}(X) \rightarrow D_{\text{good}}(X)$ commutes with direct sums.

Suppose that O_X is quasi-Noetherian. Then:

3. The subcategory $\text{Good}(X) \subset \text{Mod}(O_X)$ is weak Serre and closed under filtered colimits in $\text{Mod}(O_X)$. In particular, $\text{Good}(X)$ is a locally Noetherian category (in the sense of [Gab62, p.356]).
4. The inclusion functor $D_{\text{good}}(X) \rightarrow D(X)$ is a triangulated subcategory.

Proof.

1. Over every relatively compact open subset U of X , the direct sum $(\bigoplus_{i \in I} F_i)|_U$ is the sum of its coherent O_U -submodules. By Lemma A.1.2.1 3, the O_X -module $\bigoplus_{i \in I} F_i$ is pseudo-coherent. By Lemma A.1.4.2 2, it is good.
2. Since $\text{Mod}(O_X)$ is a Grothendieck abelian category, by [Sta24, Tag 07D9], the category $D(X)$ has arbitrary direct sums and they are computed by taking termwise direct sums of any representative complexes. Then by [Wei95, Exercise 1.2.1], for every integer q , the functor $H^q : D(X) \rightarrow \text{Mod}(O_X)$ commutes with direct sums. The result follows from Point 1.

3. As O_X is quasi-Noetherian, by [Sta24, Tag 0754] and the proof of [Kas03, Prop. 4.23], $\text{Good}(X)$ is a weak Serre subcategory of $\text{Mod}(O_X)$. From [KS06, Thm. 18.1.6 (v)], the category $\text{Mod}(O_X)$ is a Grothendieck abelian category. By Point 1 and [Sta24, Tag 002P], the filtered colimits in $\text{Good}(X)$ exist and agree with the filtered colimits in $\text{Mod}(O_X)$. Thus, filtered colimits in $\text{Good}(X)$ are exact.

Because of [Sta24, Tag 01BC], there is a set of coherent O_X -modules $\{F_i\}_{i \in I}$ such that each coherent O_X -module is isomorphic to exactly one of the F_i . Then $\{F_i\}$ is a family of Noetherian generators of $\text{Good}(X)$. Therefore, the category $\text{Good}(X)$ is locally Noetherian.

4. It follows from [Yek19, Prop. 7.4.5] and Point 3.

□

Lemma A.1.4.4. *A good module on a complex analytic space is quasi-coherent.*

Proof. Let F be a good module on a complex analytic space X . From [Fri67, Thm. I, 9; Rem. I, 10], every $x \in X$ admits a neighborhood K that is a Noetherian Stein compactum. There is a relative compact open subset U of X containing K . As F is good, the \mathcal{O}_U -module $F|_U = \sum_{i \in I} F_i$ is the sum of a directed family of coherent subsheaves. Applying the functor $\Gamma(K, \cdot)$ to the directed family $\{F_i\}_{i \in I}$ in $\text{Coh}(U)$, by [Tay02, Prop. 11.9.2], one gets a directed family of finitely generated $\Gamma(K, \mathcal{O}_K)$ -submodule $\{M_i\}_{i \in I}$ of $\Gamma(K, F)$, whose associated family in $\text{Mod}(\mathcal{O}_K)$ is $\{F_i|_K\}_{i \in I}$. Let M be $\text{colim}_{i \in I} M_i$ in $\text{Mod}(\Gamma(K, \mathcal{O}_K))$. Since the localization functor $\text{Mod}(\Gamma(K, \mathcal{O}_K)) \rightarrow \text{Mod}(\mathcal{O}_K)$ is left adjoint to $\Gamma(K, \cdot) : \text{Mod}(\mathcal{O}_K) \rightarrow \text{Mod}(\Gamma(K, \mathcal{O}_K))$, the localization preserves colimits. Then $F|_K$ is associated to M . By Lemma C.2.0.5, F is quasi-coherent. \square

Remark A.1.4.5. The restriction of a good \mathcal{O}_X -module to an open subset U is a good \mathcal{O}_U -module. Unlike quasi-coherence on schemes, goodness is not a local property. In fact, by Lemma A.1.4.3 3, every free module on a complex manifold is good, while Gabber [Con06, Eg. 2.1.6] gives a locally free (hence quasi-coherent and pseudo-coherent), but not good module on the unit open disk in \mathbb{C} . (In particular, the converse of Lemma A.1.4.4 is wrong for noncompact complex manifolds.) Still, given an \mathcal{O}_X -module F , if for every relatively compact open subset $U \subset X$, the \mathcal{O}_U -module $F|_U$ is good, then F is good.

Definition A.1.4.6 ([KS06, Def. 6.3.3]). In a category \mathcal{C} with small filtered colimits, an object X is of finite presentation, if $\text{Hom}_{\mathcal{C}}(X, \cdot) : \mathcal{C} \rightarrow \text{Set}$ commutes with small filtered colimits.

Remark A.1.4.7. In an additive category with arbitrary direct sums, an object of finite presentation is necessarily compact, but the converse is false. Let M be a finite module but not of finite presentation over a commutative ring R . By [Ren69, no. 2], M is a compact object of the abelian category $\text{Mod}(R)$. From [Sta24, Tag 0G8P], M is not an object of finite presentation.

Lemma A.1.4.8. Let (X, \mathcal{O}_X) be a ringed space. If the topology is Hausdorff compact, then every \mathcal{O}_X -module of finite presentation is an object of finite presentation of $\text{Mod}(\mathcal{O}_X)$.

Proof. Let $G = \text{colim}_{i \in I} G_i$ be a filtered colimit in $\text{Mod}(\mathcal{O}_X)$. Let F be an \mathcal{O}_X -module of finite presentation. By [Sta24, Tag 0GMV], the canonical morphism $\text{colim}_{i \in I} \text{Hom}_{\mathcal{O}_X}(F, G_i) \rightarrow \text{Hom}_{\mathcal{O}_X}(F, G)$ is an isomorphism. By compactness of X and [God58, Thm. 4.12.1], the canonical map $\text{colim}_{i \in I} H^0(X, \text{Hom}_{\mathcal{O}_X}(F, G_i)) \rightarrow H^0(X, \text{colim}_{i \in I} \text{Hom}_{\mathcal{O}_X}(F, G_i))$ is bijective. Then the canonical map $\text{colim}_{i \in I} \text{Hom}_{\text{Mod}(\mathcal{O}_X)}(F, G_i) \rightarrow \text{Hom}_{\text{Mod}(\mathcal{O}_X)}(F, G)$ is bijective. \square

Lemma A.1.4.9. Let X be a compact complex analytic space. Then the objects of finite presentation in $\text{Good}(X)$ are precisely objects of $\text{Coh}(X)$.

Proof. Let $F \in \text{Good}(X)$ be an object of finite presentation. By compactness of X , there is a directed family of coherent submodules $\{F_i\}_{i \in I}$ with $F = \sum_{i \in I} F_i$. Then the canonical morphism $\text{colim}_{i \in I} \text{Hom}(F, F_i) \rightarrow \text{Hom}(F, F)$ of abelian groups is an isomorphism. Thus, there is $i_0 \in I$ and an element $\text{Hom}(F, F_{i_0})$ lying over Id_F . As Id_F factors through F_{i_0} , one has $F = F_{i_0} \in \text{Coh}(X)$.

Conversely, by [Sta24, Tag 01BW], every object of $\text{Coh}(X)$ is an O_X -module of finite presentation. From Lemma A.1.4.8 and compactness of X , it is an object of finite presentation in $\text{Mod}(O_X)$. By Lemma A.1.4.3 3, it is also an object of finite presentation in $\text{Good}(X)$. \square

A.1.5 Sections of direct sum of sheaves

By [Har77, II, Exercise 1.11], on a Noetherian topological space, taking section commutes with (possibly infinite) direct sum of sheaves. This fails on complex manifolds, as Example A.1.5.1 shows.

Example A.1.5.1. Let $X = \mathbb{C}$. Let F be the O_X -module $\oplus_{n \geq 0} \mathbb{C}_n$. There is a section $s \in \Gamma(X, F^{\oplus \mathbb{N}})$, such that for every integer $n \geq 0$, the stalk $s_n \in (F^{\oplus \mathbb{N}})_n = (F_n)^{\oplus \mathbb{N}} = \mathbb{C}^{\oplus \mathbb{N}}$ is $(1, 1, \dots, 1, 0, 0, \dots)$, where the first $n + 1$ entries are 1 and all the other entries are 0. Then s has no preimage under the canonical map $\Gamma(X, F)^{\oplus \mathbb{N}} \rightarrow \Gamma(X, F^{\oplus \mathbb{N}})$. For otherwise, let $(t^n)_{n \geq 0} \in \Gamma(X, F)^{\oplus \mathbb{N}}$ be a preimage of s . Then there are only finitely many integers $n \geq 0$ with $t^n \neq 0$. Every t^n has only finitely many nonzero stalks. However, s has infinitely many nonzero stalks, which is a contradiction.

Let X be a complex manifold. An O_X -module is called *privileged* if for every connected open subset $U \subset X$ and every $x \in U$, the map $\Gamma(U, F) \rightarrow F_x$ taking the stalk at x is injective. By the identity theorem (see, e.g., [GH78, p.7]), O_X is privileged.

Lemma A.1.5.2. Assume that X is connected. Let $\{F_i\}_{i \in I}$ be a family of privileged O_X -modules. Then the canonical map $\oplus_{i \in I} \Gamma(X, F_i) \rightarrow \Gamma(X, \oplus_{i \in I} F_i)$ is bijective.

Proof. Let P be the presheaf direct sum of $\{F_i\}_{i \in I}$. Let $\theta : P \rightarrow \oplus_{i \in I} F_i$ be the sheafification morphism. Then $P(X) = \oplus_{i \in I} \Gamma(X, F_i)$ and $\theta_X : \oplus_{i \in I} \Gamma(X, F_i) \rightarrow \Gamma(X, \oplus_{i \in I} F_i)$ is the colimit of

$$\theta_X^{(J)} : \oplus_{i \in J} \Gamma(X, F_i) \rightarrow \Gamma(X, \oplus_{i \in I} F_i),$$

where J runs through the finite subsets of I . For every such J , by [Sta24, Tag 01AH (4)], the presheaf direct sum of $\{F_i\}_{i \in J}$ is a subsheaf of $\oplus_{i \in I} F_i$, so the map $\theta_X^{(J)}$ is injective. Therefore, their limit map θ_X is also injective. We prove that θ_X is surjective.

By construction of sheafification in [Har77, p.64], for every $s \in \Gamma(X, \oplus_{i \in I} F_i)$, there is a covering $\{U_\alpha\}_{\alpha \in A}$ of X by nonempty *connected* open subsets and an element $t_\alpha \in \Gamma(U_\alpha, P)$ for each $\alpha \in A$ such that $s_x = t_{\alpha,x}$ in $(\oplus_{i \in I} F_i)_x = \oplus_{i \in I} F_{i,x}$ for every $x \in U_\alpha$.

Fix $x_0 \in X$ and $\alpha_0 \in A$ with $x_0 \in U_{\alpha_0}$. Then there is a finite subset $I_0 \subset I$ such that $t_{\alpha_0} \in \Gamma(X, \oplus_{i \in I_0} F_i) \subset \Gamma(X, P)$. Let $B \subset A$ be the subset of indices α with $t_\alpha \notin \Gamma(U_\alpha, \oplus_{i \in I_0} F_i)$. Set $V = \cup_{\alpha \in B} U_\alpha$. Then V is open in X and its complement

$$X \setminus V \subset \cup_{\alpha \in A \setminus B} U_\alpha. \quad (\text{A.1})$$

For every $\alpha \in A \setminus B$, we claim that $U_\alpha \subset X \setminus V$.

In fact, for every $y \in U_\alpha$, every $\beta \in A$ with $y \in U_\beta$ and every $i \in I \setminus I_0$, the stalk $t_{\beta,y}^i = s_y^i = t_{\alpha,y}^i = 0$ in $F_{i,y}$. Since F_i is privileged and U_β is connected, the map $\Gamma(U_\beta, F_i) \rightarrow F_{i,y}$ is injective. Thus, $t_\beta^i = 0$ in $\Gamma(U_\beta, F_i)$. Therefore, $t_\beta \in \Gamma(X, \oplus_{i \in I_0} F_i)$, i.e., $\beta \notin B$. Hence $y \notin V$.

From the claim and (A.1), the subset $X \setminus V = \cup_{\alpha \in A \setminus B} U_\alpha$ is also open in X and contains U_{α_0} . Since X is connected, one has $V = B = \emptyset$. Consequently, $t_\alpha \in \Gamma(X, \oplus_{i \in I_0} F_i)$ for every $\alpha \in A$. Then the family $\{t_\alpha\}_{\alpha \in A}$ glues to a preimage of s in $\Gamma(X, \oplus_{i \in I_0} F_i) \subset \Gamma(X, P)$. Thus, θ_X is surjective and hence a group isomorphism. \square

Corollary A.1.5.3. *If F is a locally free (possibly of infinite rank) \mathcal{O}_X -module, then F is privileged.*

Proof. Let U be a connected open subset of X . Fix $x_0 \in U$. We prove that the map $\Gamma(U, F) \rightarrow F_{x_0}$ is injective. Take $s \in \Gamma(U, F)$ with $s_{x_0} = 0$. By [Har77, II, Exercise 1.14], the set $Z := \{x \in U : s_x = 0\}$ is open in U .

We claim that Z is closed in U . Let $\{x_n\}_{n \geq 1}$ be a sequence of points in Z converging to $y \in U$. Because F is locally free, there is a connected open neighborhood $V \subset U$ of y , a set I and an isomorphism $\phi : F|_V \xrightarrow{\sim} \mathcal{O}_V^{\oplus I}$ of \mathcal{O}_V -modules. There is an integer $N > 0$ with $x_N \in V$. Because \mathcal{O}_V is privileged, from Lemma A.1.5.2, the map on the bottom of the commutative square

$$\begin{array}{ccc} \Gamma(V, F) & \longrightarrow & F_{x_N} \\ \downarrow \phi_V & & \downarrow \phi_{x_N} \\ \Gamma(V, \mathcal{O}_V^{\oplus I}) & \longrightarrow & \mathcal{O}_{V, x_N}^{\oplus I} \end{array}$$

is injective. Then so is the map on the top. Since $s_{x_N} = 0$, one has $s|_V = 0$ and $s_y = 0$. Hence $y \in Z$. The claim is proved.

Because U is connected and $x_0 \in Z$, by claim one has $Z = U$. Therefore, $s = 0$ in $\Gamma(U, F)$. \square

Corollary A.1.5.4. *Let X be a connected complex manifold. Let $\{F_i\}_{i \in I}$ be a family of locally free O_X -modules. Then the canonical map $\oplus_{i \in I} \Gamma(X, F_i) \rightarrow \Gamma(X, \oplus_{i \in I} F_i)$ is bijective.*

Proof. It follows from Lemma A.1.5.2 and Corollary A.1.5.3. \square

A.2 Gabber's example

We present an example of a locally free module on the open unit disc that is not good. It illustrates that goodness on complex manifolds, unlike quasi-coherence on algebraic varieties, is not a local property. This construction is already exhibited in the context of rigid geometry by [Con06, Example 2.1.6], which attributes the originality to Gabber. Furthermore, in [Con06, p.1058] it is mentioned that Gabber's example makes sense in complex-analytic geometry as well. We reproduce this construction with a few extra details.

Lemma A.2.0.1. *Let X be a complex manifold, U be a dense open subset of X . If F is a locally free O_X -module, then the restriction map $r : \Gamma(X, F) \rightarrow \Gamma(U, F)$ is injective.*

Proof. Every $x \in X$ admits a connected open neighborhood V such that $F|_V$ is free O_V -module. Then $\Gamma(V, F)$ is a free $\Gamma(V, O_X)$ -module by Corollary A.1.5.4. By density of U , $V \cap U$ is nonempty. For every $s \in \ker(r)$, $s|_{V \cap U} = 0$. As $F|_V$ is free and the map $\Gamma(V, O_X) \rightarrow \Gamma(V \cap U, O_X)$ is injective, the restriction $s|_V = 0$. By local nature of sheaves, $s = 0$. \square

Lemma A.2.0.2. *Let X be a Hausdorff locally compact space, K be a compact subspace and $j : K \rightarrow X$ be the inclusion. Then for every $q \in \mathbb{Z}$ and every $F \in \text{Ab}(X)$, the canonical morphism $\psi^q : \text{colim}_U H^q(U, F) \rightarrow H^q(K, j^{-1}F)$ is an isomorphism, where U ranges through the family of open neighborhoods of K in X . The two groups are written as $H^q(K, F)$.*

Proof. We prove that both sides are the q -th right derived functor applied to F of a same functor.

Define a category I as follows. The objects are the open subsets of X containing K . For every $U, V \in I$, if $U \supset V$, then $\text{Hom}_I(U, V)$ is a singleton; else $\text{Hom}_I(U, V) = \emptyset$. Thus, I is a small category. Let Ab^I be the category of functors from I to Ab . By [Wei95, Exercise 2.3.7], Ab^I is an abelian category with enough injectives. Recall that Ab is a Grothendieck abelian category, so $\text{colim}_I : \text{Ab}^I \rightarrow \text{Ab}$ is exact. By [KS90, Prop 2.5.1], the composition of the functor $\Phi : \text{Ab}(X) \rightarrow \text{Ab}^I$ defined by $\Phi(F)(U) = \Gamma(U, F)$ with $\text{colim}_I : \text{Ab}^I \rightarrow \text{Ab}$ is $\Gamma(K, j^{-1}\cdot) : \text{Ab}(X) \rightarrow \text{Ab}$. Therefore, the q -th right derived functor of $\Gamma(K, j^{-1}\cdot)$ is $\text{colim}_I \circ R^q\Phi = \text{colim}_U H^q(U, \cdot)$.

The functor $\Gamma(K, j^{-1}\cdot) : \text{Ab}(X) \rightarrow \text{Ab}$ is the composition of $j^{-1} : \text{Ab}(X) \rightarrow \text{Ab}(K)$ with $\Gamma(K, \cdot) : \text{Ab}(K) \rightarrow \text{Ab}(X)$. Every injective object G of $\text{Ab}(X)$ is c-soft, so $j^{-1}G$ is c-soft by [KS90, Proposition 2.5.7 (i)]. By [KS90, Proposition 2.5.10], $j^{-1}G$ is right acyclic for $\Gamma(K, \cdot)$. By [Sta24, Tag 015M], $H^q(K, j^{-1}\cdot)$ is also the q -th right derived functor of $\Gamma(K, j^{-1}\cdot)$. We conclude that ψ^q is an isomorphism. \square

Definition A.2.0.3 (Compact Stein set). [Con06, p.1053] Let K be a compact subset of a complex manifold X . If $H^q(K, F) = 0$ for every open neighborhood U of $K \subset X$, every coherent \mathcal{O}_U -module and every $q \in \mathbb{N}^*$, then K is called a compact Stein set in X .

Lemma A.2.0.4 ([Con06, p.1058]). *Let K be a compact compact Stein set in a complex manifold X , F be a good \mathcal{O}_X -module, then $H^q(K, F) = 0$ for all $q \in \mathbb{N}^*$.*

Proof. There is a relative compact open subset $U \subset X$ containing K . By definition, $F|_U = \text{colim}_i F_i$, where $\{F_i\}$ is a direct family of coherent \mathcal{O}_U -submodules of $F|_U$. By [God58, II, Thm. 4.12.1], $H^q(K, F) = \text{colim}_i H^q(K, F_i) = 0$. \square

Example A.2.0.5 (Gabber). Let Δ be the open unit disc in \mathbb{C} and let $K = \{z \in \mathbb{C} : |z| \leq 1/2\}$. Then $B(0, 2/3)$ is a relatively compact open subset of Δ containing K . By [Dou66, Thm. 3 (B), p.51; (a) p.53], K is a compact Stein set in Δ .

Let x', x'' be two distinct points of the interior of K . Let $U' = \Delta \setminus \{x'\}$, $U'' = \Delta \setminus \{x''\}$ and define $U = U' \cap U''$. Let

$$F' = \bigoplus_{n \in \mathbb{Z}} \mathcal{O}_{U'} e'_n, \quad F'' = \bigoplus_{n \in \mathbb{Z}} \mathcal{O}_{U''} e''_n$$

be two free sheaves with countably infinite rank on U' and U'' respectively.

We glue F' and F'' to define a locally free \mathcal{O}_Δ -module F as follows. Define $h \in \mathcal{O}_\Delta(U)$ by

$$h(z) = e^{\frac{1}{z-x'} + \frac{1}{z-x''}}, \quad \forall z \in U.$$

Then h has essential singularities at x' and x'' . Define F by identifying $F'|_U$ and $F''|_U$ with the free sheaf $\bigoplus_{n \in \mathbb{Z}} \mathcal{O}_U e_n$ via the conditions

$$\begin{aligned} e_{2m} &= e'_{2m}|_U = e''_{2m}|_U + h e''_{2m+1}|_U, \\ e_{2m+1} &= e''_{2m+1}|_U = e'_{2m+1}|_U + h e'_{2m+2}|_U \end{aligned}$$

for every $m \in \mathbb{Z}$ respectively.

We prove that $\Gamma(K, F) = 0$. For every $s \in \Gamma(K, F)$, by Lemma A.2.0.2, there is an open subset W of Δ containing K such that s lifts to an element of $\Gamma(W, F)$. By Corollary A.1.5.4, $\Gamma(U, F) = \bigoplus_{n \in \mathbb{Z}} \Gamma(U, \mathcal{O}_\Delta) e_n$. So, $s|_{U \cap W} =$

$\sum_{n \in \mathbb{Z}} f_n e_n$ with $f_n \in O_\Delta(U \cap W)$ that vanish for all but finitely many n . Note that

$$s|_{U''} = \sum_{n \in \mathbb{Z}} (f_{2n} e_{2n}'' + (f_{2n} h + f_{2n+1}) e_{2n+1}'')|_{U''}.$$

Therefore, f_{2n} and $f_{2n} h + f_{2n+1}$ are holomorphic near x' for all $n \in \mathbb{Z}$. Similarly, f_{2n+1} and $f_{2n+1} h + f_{2n+2}$ are holomorphic near x'' for all $n \in \mathbb{Z}$.

We claim that $s|_{U \cap W} = 0$. Otherwise, let n_0 be the maximum with $f_{n_0} \neq 0$. If n_0 is odd (resp. even), f_{n_0} and $h f_{n_0}$ are holomorphic near x'' (resp. x'). The ratio $h = h f_{n_0} / f_{n_0}$ is meromorphic near x'' (resp. x'). It contradicts the choice of h . The claim is proved.

By Lemma A.2.0.1, the restriction map $\Gamma(W, F) \rightarrow \Gamma(W \cap U, F)$ is injective, so $s = 0$.

We prove that F is not good. Let t be the standard coordinate on Δ , then $0 \rightarrow F \xrightarrow{t} F \rightarrow F/tF \rightarrow 0$ is a short exact sequence in $\text{Mod}(O_\Delta)$. The associated cohomology sequence induces an injection $H^0(K, F/tF) \rightarrow H^1(K, F)$ by Lemma A.2.0.2. As F/tF is the skyscraper supported at the origin, we have $H^0(K, F/tF) \neq 0$ and hence $H^1(K, F) \neq 0$. By Lemma A.2.0.4, the O_Δ -module F is not good. and $F|_K$ is not induced by a $\Gamma(K, O_K)$ -module. In particular, F is not quasi-coherent in the sense of last paragraph of [BBBP07, p.443]. Nevertheless, F is quasi-coherent in the sense of [Gro60, 5.1.3] since it is locally free.

Appendix B

Quasi-coherent GAGA

B.1 Introduction

Let X be a complex algebraic variety. Then the set of complex points $X(\mathbb{C})$ underlies a natural complex analytic space (in the sense of [Ser56, Déf. 1]) structure, denoted by X^{an} . When X is a projective variety, Serre [Ser56, Théorèmes 2 et 3] proves that the abelian category of (algebraic) coherent modules on X is naturally equivalent to that of (analytic) coherent modules on X^{an} . Hall [Hal23] extends the equivalence to the bounded derived category of coherent modules (Fact B.2.0.1).

A natural question is to find analogous equivalences for the larger category of quasi-coherent sheaves on X . We show that good modules (in the sense of Kashiwara, Definition A.1.4.1) is a analytic counterpart of quasi-coherent sheaves on algebraic varieties.

For a ringed space (X, \mathcal{O}_X) , let $\text{Mod}(\mathcal{O}_X)$ be the abelian category of \mathcal{O}_X -modules. Let $D(X)$ be its unbounded derived category.

For an algebraic variety (resp. a complex analytic space) X , let $\text{Qch}(X) \subset \text{Mod}(\mathcal{O}_X)$ (resp. $\text{Good}(X) \subset \text{Mod}(\mathcal{O}_X)$) be the full subcategory of quasi-coherent (resp. good) modules. Let $D_{\text{qc}}(X)$ (resp. $D_{\text{good}}(X)$) be the full subcategory of $D(X)$ comprised of objects with quasi-coherent (resp. good) cohomologies.

Theorem (Proposition B.3.0.2, Theorem B.4.0.2). *If X is proper over \mathbb{C} , then the analytification functor $D_{\text{qc}}(X) \rightarrow D_{\text{good}}(X^{\text{an}})$ is an equivalence of triangulated categories.*

B.2 Review

We recall the work of Serre [Ser56] (known as “GAGA”), which gives an equivalence of algebraic coherent modules and analytic coherent modules

on complex, projective varieties. The theory is extended to complex, proper algebraic varieties in [GR71, Exp. XII].

Let X be a complex algebraic variety. Let \mathbf{An} (resp. \mathbf{Set}) be the category of complex analytic spaces (resp. sets). Let Ψ_X be the functor $\mathbf{An} \rightarrow \mathbf{Set}$ sending a complex analytic space Y to the set $\mathrm{Hom}_{\mathbb{C}}(Y, X)$ of morphisms of spaces with a sheaf of \mathbb{C} -algebras. By [GR71, Exp. XII, Thm. 1.1], the functor Ψ_X is represented by a complex analytic space¹ X^{an} (called the analytification of X) and a *flat* morphism $\psi_X \in \mathrm{Hom}_{\mathbb{C}}(X^{\mathrm{an}}, X)$. Because X is of finite type over \mathbb{C} , from [GR71, Exp. XII, Prop. 2.1 (viii)], the dimension of X^{an} is finite.

By [GR71, Exp. XII, 1.2], for every morphism $f : X \rightarrow Y$ of complex algebraic varieties, there is a commutative square

$$\begin{array}{ccc} X^{\mathrm{an}} & \xrightarrow{\psi_X} & X \\ \downarrow f^{\mathrm{an}} & & \downarrow f \\ Y^{\mathrm{an}} & \xrightarrow{\psi_Y} & Y \end{array} \quad (\text{B.1})$$

in the category of ringed spaces. In other words, the analytification induces a functor $(\cdot)^{\mathrm{an}}$ from the category of complex algebraic varieties to \mathbf{An} .

For a ringed space (Y, \mathcal{O}_Y) , let $\mathrm{Coh}(Y) \subset \mathrm{Mod}(\mathcal{O}_Y)$ be the full subcategory comprised of coherent modules (in the sense of [Sta24, Tag 01BV]). Let $D_c(Y) \subset D(Y)$ be the full subcategory consisting of objects with coherent cohomologies. The pullback functor

$$\psi_X^* : \mathrm{Mod}(\mathcal{O}_X) \rightarrow \mathrm{Mod}(\mathcal{O}_{X^{\mathrm{an}}}), \quad F \mapsto F^{\mathrm{an}} \quad (\text{B.2})$$

is exact and admits a right adjoint, so it commutes with colimits. It extends to a functor $D(X) \rightarrow D(X^{\mathrm{an}})$, which is t-exact relative to the standard t-structures. From [GR71, Exp. XII, 1.3], it restricts to a functor $D_c^b(X) \rightarrow D_c^b(X^{\mathrm{an}})$ and $\mathrm{Coh}(X) \rightarrow \mathrm{Coh}(X^{\mathrm{an}})$.

Fact B.2.0.1 can be retracted from [Hal23, Remark 1.1 and the proof of Theorem A]. Neeman [Nee21, Example A.2] modifies Hall's proof to some extent.

Fact B.2.0.1. *Assume that the complex algebraic variety X is proper. Then the functor (B.2) induces an equivalence $D_c^b(X) \rightarrow D_c^b(X^{\mathrm{an}})$ of triangulated categories. In particular, it restricts to an equivalence $\mathrm{Coh}(X) \rightarrow \mathrm{Coh}(X^{\mathrm{an}})$ of abelian categories.*

By Lemma A.1.4.3, $\mathrm{Good}(X)$ is a weak Serre subcategory of $\mathrm{Mod}(\mathcal{O}_X)$, and $D_{\mathrm{good}}(X)$ is a triangulated subcategory of $D(X)$.

¹Strictly speaking, complex analytic spaces are allowed to be non-Hausdorff in [GR71, Exp. XII]. In our case, the algebraic variety X is assumed to be separated over \mathbb{C} , by [GR71, Exp. XII, Prop. 3.1 (viii)], the topology of X^{an} is Hausdorff.

Lemma B.2.0.2. *For the complex algebraic variety X , the functor (B.2) restricts to a functor*

$$\mathrm{Qch}(X) \rightarrow \mathrm{Good}(X^{\mathrm{an}}) \quad (\mathrm{B.3})$$

and induces a functor

$$D_{\mathrm{qc}}(X) \rightarrow D_{\mathrm{good}}(X^{\mathrm{an}}). \quad (\mathrm{B.4})$$

Proof. For every quasi-coherent O_X -module F , by Fact B.2.0.3,

$$F = \sum_{i \in I} F_i \quad (\mathrm{B.5})$$

is the sum of a direct family of coherent O_X -submodules. As ψ_X^* commutes with colimits, one has

$$\psi_X^* F = \mathrm{colim}_{i \in I} \psi_X^* F_i \quad (\mathrm{B.6})$$

in the category $\mathrm{Mod}(O_{X^{\mathrm{an}}})$. Since ψ_X^* is exact, each $\psi_X^* F_i$ is a coherent $O_{X^{\mathrm{an}}}$ -submodule of $\psi_X^* F$. Therefore, the $O_{X^{\mathrm{an}}}$ -module $\psi_X^* F$ is good.

For every $G \in D_{\mathrm{qc}}(X)$ and every integer n , because (B.2) is an exact functor, the $O_{X^{\mathrm{an}}}$ -module $H^n(\psi_X^* G) = \psi_X^*(H^n G)$ is good by last paragraph. Hence $\psi_X^* G \in D_{\mathrm{good}}(X^{\mathrm{an}})$. \square

Fact B.2.0.3 ([Gro60, Cor. 9.4.9], [Sta24, Tag 01PG]). *On a Noetherian scheme, every quasi-coherent sheaf is the sum of the directed family of all coherent submodules.*

B.3 GAGA for quasi-coherent modules

Using Fact B.2.0.3 and that ψ_X^* commutes with colimits, we extend GAGA from coherent O_X -modules to quasi-coherent O_X -modules. When $Y = \mathrm{Spec} \mathbb{C}$, Proposition B.3.0.1 generalizes [Ser56, Thm. 1].

Proposition B.3.0.1. *Let $f : X \rightarrow Y$ be a proper morphism of complex algebraic varieties. Then the base change natural transformation $(Rf_*)^{\mathrm{an}} \rightarrow Rf_*^{\mathrm{an}}(\cdot)^{\mathrm{an}}$ (induced by the commutative square (B.1)) induces an isomorphism of functors $D_{\mathrm{qc}}(X) \rightarrow D_{\mathrm{good}}(Y^{\mathrm{an}})$.*

Proof. For every $F \in D_{\mathrm{qc}}(X)$, by [Lip60, Prop. 3.9.2], one has $Rf_* F \in D_{\mathrm{qc}}(Y)$. By Lemma B.2.0.2, one has $F^{\mathrm{an}} \in D_{\mathrm{good}}(X^{\mathrm{an}})$ and $(Rf_* F)^{\mathrm{an}} \in D_{\mathrm{good}}(Y^{\mathrm{an}})$. Since f is proper, from [GR71, Exp. XII, Prop. 3.2 (v)], the morphism $f^{\mathrm{an}} : X^{\mathrm{an}} \rightarrow Y^{\mathrm{an}}$ is proper. As X^{an} has finite dimension, by Theorem 5.3.1.7, one has $Rf_*^{\mathrm{an}} F^{\mathrm{an}} \in D_{\mathrm{good}}(Y^{\mathrm{an}})$. Therefore, both functors $(Rf_*)^{\mathrm{an}}$ and $Rf_*^{\mathrm{an}}(\cdot)^{\mathrm{an}}$ restrict to functors $D_{\mathrm{qc}}(X) \rightarrow D_{\mathrm{good}}(Y^{\mathrm{an}})$.

We prove that the morphism $(Rf_* F)^{\mathrm{an}} \rightarrow Rf_*^{\mathrm{an}} F^{\mathrm{an}}$ is an isomorphism. By Lemma 5.3.1.11 (resp. [Lip60, Prop. 3.9.2]), the functor $Rf_*^{\mathrm{an}} : D(X^{\mathrm{an}}) \rightarrow D(Y^{\mathrm{an}})$ (resp. $Rf_* : D_{\mathrm{qc}}(X) \rightarrow D_{\mathrm{qc}}(Y)$) is bounded. From

[Sta24, Tag 06YZ], the inclusion functor $\mathrm{Qch}(X) \rightarrow \mathrm{Mod}(O_X)$ exhibits $\mathrm{Qch}(X)$ as a weak Serre subcategory (in the sense of [Sta24, Tag 02MO]) of $\mathrm{Mod}(O_X)$. Then by (way-out argument) [Har66, I, Prop. 7.1 (iii)], one may assume $F \in \mathrm{Qch}(X)$. By [KS06, Prop. 13.1.5 (ii), p.320], it suffices to check that for every integer $n \geq 0$, the natural morphism $(R^n f_* F)^{\mathrm{an}} \rightarrow R^n f_*^{\mathrm{an}}(F^{\mathrm{an}})$ in $\mathrm{Mod}(O_{Y^{\mathrm{an}}})$ is an isomorphism.

By Fact B.2.0.3, one can write $F = \sum_{i \in I} F_i$ as the sum of a direct family of coherent O_X -submodules of F . By [Sta24, Tag 07TB], one has

$$\mathrm{colim}_{i \in I} R^n f_* F_i \xrightarrow{\sim} R^n f_* F.$$

The analytification commutes with colimits, so

$$\mathrm{colim}_{i \in I} (R^n f_* F_i)^{\mathrm{an}} \xrightarrow{\sim} (R^n f_* F)^{\mathrm{an}}.$$

By [GR71, XII, Thm. 4.2], the natural morphisms $(R^n f_* F_i)^{\mathrm{an}} \rightarrow R^n f_*^{\mathrm{an}}(F_i^{\mathrm{an}})$ are isomorphisms. By Lemma 5.3.1.9, the natural morphism

$$\mathrm{colim}_{i \in I} R^n f_*^{\mathrm{an}}(F_i^{\mathrm{an}}) \rightarrow R^n f_*^{\mathrm{an}}(F^{\mathrm{an}})$$

is an isomorphism. □

Proposition B.3.0.2 shows that goodness on complex analytic spaces is an analytic counterpart of quasi-coherence on complex algebraic varieties.

Proposition B.3.0.2. *Suppose that the complex algebraic variety X is proper. Then (B.3) is an equivalence of abelian categories.*

Proof. • The functor (B.3) is essentially surjective: Indeed, because X is proper over \mathbb{C} , by [GR71, Exp. XII, Prop. 3.2 (v)], the complex analytic space X^{an} is compact. Then for every good $O_{X^{\mathrm{an}}}$ -module G , one can write $G = \sum_{i \in I} G_i$ as the sum of a directed family of coherent $O_{X^{\mathrm{an}}}$ -submodules. From the equivalence $\psi_X^* : \mathrm{Coh}(X) \rightarrow \mathrm{Coh}(X^{\mathrm{an}})$ ([GR71, XII, Thm. 4.4]), there is a filtered inductive system $\{H_i\}_{i \in I}$ in $\mathrm{Coh}(X)$ whose analytification is the filtered inductive system $\{G_i\}_{i \in I}$. By [Sta24, Tag 01LA (4)], the colimit H of $\{H_i\}$ in $\mathrm{Mod}(O_X)$ exists and lies in $\mathrm{Qch}(X)$. Because ψ_X^* commutes with colimits, one has $H^{\mathrm{an}} = \mathrm{colim}_{i \in I} G_i$. In particular, H^{an} is isomorphic to G in $\mathrm{Good}(X^{\mathrm{an}})$.

- The functor (B.3) is fully faithful: For any quasi-coherent O_X -modules F and G , we have to show that the canonical morphism

$$\mathrm{Hom}_{O_X}(F, G) \rightarrow \mathrm{Hom}_{O_{X^{\mathrm{an}}}}(F^{\mathrm{an}}, G^{\mathrm{an}}) \tag{B.7}$$

is an isomorphism. Assume first that F is coherent.

- From [GW20, Exercise 7.20 (b)], one has

$$[\mathcal{H}om_{O_X}(F, G)]^{\mathrm{an}} = \mathcal{H}om_{O_{X^{\mathrm{an}}}}(F^{\mathrm{an}}, G^{\mathrm{an}}).$$

- As F is of finite presentation, the O_X -module $\mathcal{H}om_{O_X}(F, G)$ is quasi-coherent.

Therefore, by Proposition B.3.0.1, the canonical morphism

$$H^0(X, \mathcal{H}om_{O_X}(F, G)) \rightarrow H^0(X^{\text{an}}, \mathcal{H}om_{O_{X^{\text{an}}}}(F^{\text{an}}, G^{\text{an}}))$$

is an isomorphism, which is exactly (B.7).

By (B.5) and (B.6), the general case follows. □

B.4 Derived category of quasi-coherent sheaves

By [Sta24, Tag 0BKN], for every ringed space Y , the derived category $D(Y)$ has products and derived limits. This plays an essential role in step 4 of the proof of Theorem B.4.0.2.

Definition B.4.0.1. [Sta24, Tag 07LS] Let \mathcal{A} be an additive category with arbitrary direct sums. An object $K \in \mathcal{A}$ is called compact, if $\text{Hom}_{\mathcal{A}}(K, \cdot) : \mathcal{A} \rightarrow \text{Ab}$ preserves direct sums.

Theorem B.4.0.2. *If the complex algebraic variety X is proper, then the functor (B.4) is an equivalence of triangulated categories.*

Proof. Since X^{an} is compact, by Lemma B.4.0.6, the perfect complex $O_{X^{\text{an}}}$ is a compact object of $D(X^{\text{an}})$. Then from the proof of [Hal23, Lem. 4.3], the functor $\psi_X^* : D_{\text{qc}}(X) \rightarrow D(X^{\text{an}})$ admits a right adjoint functor $R\psi_{\text{qc},*} : D(X^{\text{an}}) \rightarrow D_{\text{qc}}(X)$ which preserves small coproducts.

1. The functor $\psi_X^* : D_{\text{qc}}(X) \rightarrow D(X^{\text{an}})$ is fully faithful.

From Fact B.2.0.1, the unit of the adjunction $\eta : \text{Id} \rightarrow R\psi_{\text{qc},*}\psi_X^*$ (a natural transformation of functors $D_{\text{qc}}(X) \rightarrow D_{\text{qc}}(X)$) restricts to an isomorphism of functors $D_c^b(X) \rightarrow D_c^b(X)$. By [BB03, Thm. 3.1.1 1], the compact objects of $D_{\text{qc}}(X)$ are precisely the perfect complexes. From [Nee96, Prop. 2.5], $D_{\text{qc}}(X)$ is generated by a family of perfect complexes $\{E_i\}_{i \in I}$. By [Sta24, Tag 0FXU (1)], every perfect complex in $D(X)$ belongs to $D_c^b(X)$, so the η_{E_i} are isomorphisms. From Lemma B.4.0.5, η is an isomorphism of functors $D_{\text{qc}}(X) \rightarrow D_{\text{qc}}(X)$. Thus, 1 is proved.

2. The functor (B.4) restricts to an equivalence $D_{\text{qc}}^b(X) \rightarrow D_{\text{good}}^b(X^{\text{an}})$.

We prove that every $F \in D_{\text{good}}^b(X^{\text{an}})$ is in the essential image of $D_{\text{qc}}^b(X) \rightarrow D_{\text{good}}^b(X^{\text{an}})$. Induction on the cohomological length of F . By Proposition B.3.0.2, it holds when F has length zero. Suppose that it is true for objects

of length $\leq n$ and F has length $n + 1$. There is an integer i such that $\tau^{\leq i} F, \tau^{> i} F$ have length $\leq n$. There is a canonical exact triangle

$$\tau^{\leq i} F \rightarrow F \rightarrow \tau^{> i} F \xrightarrow{+1} \tau^{\leq i} F[1]$$

in $D_{\text{good}}^b(X^{\text{an}})$. By 1 and the inductive hypothesis, the morphism $+1 : \tau^{> i} F \rightarrow \tau^{\leq i} F[1]$ is in the essential image of $D_{\text{qc}}^b(X) \rightarrow D_{\text{good}}^b(X^{\text{an}})$. Then so is F . The essential surjectivity together with 1 proves 2.

3. The functor $\psi_X^* : D_{\text{qc}}^+(X) \rightarrow D_{\text{good}}^+(X^{\text{an}})$ is an equivalence.

For every $F \in D_{\text{good}}^+(X^{\text{an}})$, by Lemma B.4.0.3, one has $\text{hocolim}_{n>0} \tau^{\leq n} F \xrightarrow{\sim} F$. Every $\tau^{\leq n} F$ is in $D_{\text{good}}^b(X^{\text{an}})$. From 2, there is a system $(K_n)_{n>0}$ of objects of $D_{\text{qc}}^b(X)$, whose image under ψ_X^* is isomorphic to the system $(\tau^{\leq n} F)_{n>0}$. Since (B.4) respects coproducts, it respects homotopy colimits. Since $\text{Qch}(X)$ is closed under filtered colimits in $\text{Mod}(O_X)$, the subcategory $D_{\text{qc}}(X)$ is closed under homotopy colimits in $D(X)$.

Then F is isomorphic to the image of $K := \text{hocolim}_{n>0} K_n \in D_{\text{qc}}(X)$ under ψ_X^* . There is an integer q , such that $H^i(F) = 0$ for every integer $i < q$. Then $\psi_X^* H^i(K) = H^i(\psi_X^* K) = 0$. By Proposition B.3.0.2, one has $H^i(K) = 0$. Hence $K \in D_{\text{qc}}^+(X)$. Thus, the functor $\psi_X^* : D_{\text{qc}}^+(X) \rightarrow D_{\text{good}}^+(X^{\text{an}})$ is essentially surjective. By 1, it is an equivalence.

4. Every $Z \in D_{\text{good}}(X^{\text{an}})$ is in the essential image of (B.4).

By Lemma B.4.0.4, the canonical morphism $Z \rightarrow \text{Rlim}_{n>0} \tau^{\geq -n} Z$ is an isomorphism in $D(X^{\text{an}})$. By 3, there is an inverse system (Y^{-n}) of objects of $D_{\text{qc}}^+(X)$, whose image is isomorphic to the inverse system $(\tau^{\geq -n} Z)_{n>0}$. Let Y be $\text{Rlim}_{n>0} Y^{-n}$ in $D(X)$. For any integers $n \geq 1$ and q , the functor ψ_X^* transforms the morphism $H^q(Y^{-n-1}) \rightarrow H^q(Y^{-n})$ in $\text{Qch}(X)$ to $H^q(\tau^{\geq -n-1} Z) \rightarrow H^q(\tau^{\geq -n} Z)$ in $\text{Good}(X^{\text{an}})$.

The morphism $H^q(\tau^{\geq -n-1} Z) \rightarrow H^q(\tau^{\geq -n} Z)$ is surjective, and when $n \geq -q$, it is an isomorphism. By Proposition B.3.0.2, the morphism $H^q(Y^{-n-1}) \rightarrow H^q(Y^{-n})$ is surjective, and when $n \geq -q$, it is an isomorphism. By [Sta24, Tag 0A0J (1)], the canonical morphism $H^q(Y) \rightarrow H^q(Y^{\min(q, -1)})$ is an isomorphism. In particular, the O_X -module $H^q(Y)$ is quasi-coherent. Hence $Y \in D_{\text{qc}}(X)$.

For every integer $m > 0$, the functor ψ_X^* transforms $\prod_{n>0} Y^{-n} \rightarrow Y^{-m}$ to $\psi_X^*(\prod_{n>0} Y^{-n}) \rightarrow \tau^{\geq -m} Z$. Hence a morphism $\psi_X^*(\prod_{n>0} Y^{-n}) \rightarrow \prod_{n>0} \tau^{\geq -n} Z$ in $D(X^{\text{an}})$. It fits to a commutative diagram

$$\begin{array}{ccccccc} \psi_X^*(\prod_{n>0} Y^{-n})[-1] & \longrightarrow & \psi_X^*(\prod_{n>0} Y^{-n})[-1] & \longrightarrow & \psi_X^* Y & \longrightarrow & \psi_X^*(\prod_{n>0} Y^{-n}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \prod_{n>0} \tau^{\geq -n} Z[-1] & \longrightarrow & \prod_{n>0} \tau^{\geq -n} Z[-1] & \longrightarrow & Z & \longrightarrow & \prod_{n>0} \tau^{\geq -n} Z \end{array}$$

in $D(X^{\text{an}})$, where the rows are exact triangles. By TR3, it induces a morphism of triangles. Hence a commutative square

$$\begin{array}{ccc} H^q(\psi_X^* Y) & \xrightarrow{\sim} & H^q(\psi_X^* Y^{\min(q, -1)}) \\ \downarrow & & \downarrow \sim \\ H^q(Z) & \xrightarrow{\sim} & H^q(\tau^{\geq \min(q, -1)} Z) \end{array}$$

in $\text{Mod}(O_{X^{\text{an}}})$. Therefore, for every integer q , the induced morphism $H^q(\psi_X^* Y) \rightarrow H^q(Z)$ is an isomorphism. Therefore, the morphism $\psi_X^* Y \rightarrow Z$ is an isomorphism in $D_{\text{good}}(X^{\text{an}})$. Thus, 4 is proved. By 4 and 1, the functor (B.4) is an equivalence. \square

Lemma B.4.0.3. *Let \mathcal{A} be an abelian category, where colimits over \mathbb{N} exist and are exact. Then the natural transformation $\text{hocolim}_{n>0} \tau^{\leq n} \cdot \rightarrow \text{Id}$ is an isomorphism of functors $\mathcal{A} \rightarrow \mathcal{A}$.*

Proof. It follows from [Sta24, Tag 0949] and the construction of canonical truncations. \square

Lemma B.4.0.4. *Let X be a complex analytic space. Then the natural transformation $\text{Id} \rightarrow \text{Rlim}_{n>0} \tau^{\geq -n} \cdot$ is an isomorphism of functors $D(X) \rightarrow D(X)$.*

Proof. For every $x \in X$, there is an integer $d_x \geq 0$, and a fundamental system \mathcal{U}_x of open neighborhoods of x , such that every $U \in \mathcal{U}_x$ is a closed complex subspace of a domain in \mathbb{C}^{d_x} . By Fact 5.3.1.10, for every $E \in D(X)$, any integers $p > 2d_x$ and q , one has $H^p(U, H^q(E)) = 0$. By [Sta24, Tag 0D63], the canonical morphism $E \rightarrow \text{Rlim}_{n>0} \tau^{\geq -n} E$ is an isomorphism in $D(X)$. \square

Lemma B.4.0.5. *Let \mathcal{C}, \mathcal{D} be triangulated categories. Assume that \mathcal{C} has direct sums. Let $\{E_i\}_{i \in I}$ be a family of compact objects of \mathcal{C} such that $\oplus_{i \in I} E_i$ generates \mathcal{C} . Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be triangulated functors preserving direct sums. Let $\eta : F \rightarrow G$ be a natural transformation. If for every $i \in I$, the morphism $\eta_{E_i} : F(E_i) \rightarrow G(E_i)$ is an isomorphism in \mathcal{D} , then η is an isomorphism.*

Proof. From [Sta24, Tag 09SN], every object $X \in \mathcal{C}$ can be written as $X = \text{hocolim}_{n>0} X_n$, where

- X_1 is a direct sum of shifts of the E_i ,
- each transition morphism $X_n \rightarrow X_{n+1}$ fits into an exact triangle $Y_n \rightarrow X_n \rightarrow X_{n+1} \rightarrow Y_n[1]$,
- and Y_n is a direct sum of shifts of the E_i .

Since F, G preserve direct sums, and the η_{E_i} are isomorphisms, so are the $\{\eta_{X_n}\}_{n>0}$ and η_{X_1} . By [Sta24, Tag 014A] and induction on $n > 0$, one proves that the η_{X_n} are isomorphisms. By [BN93, Lem. 4.1], $F, G : \mathcal{C} \rightarrow \mathcal{D}$ preserve homotopy colimits. Therefore, η_X is an isomorphism. \square

Lemma B.4.0.6. *Let X be a compact complex analytic space. Then every perfect object of $D(X)$ belongs to $D_c^b(X)$. It is a compact object of $D(X)$ and of $D_{\text{good}}(X)$.*

Proof. Let $E \in D(X)$ be a perfect object. By definition, there is an open covering $X = \cup_{i \in I} U_i$, such that for each $i \in I$, there is a morphism of complexes $E_i^\bullet \rightarrow E|_{U_i}$ which is a quasi-isomorphism, with $E_i^j = 0$ for all but finite many integers j , and every E_i^j is a direct summand of a finite free \mathcal{O}_X -module. Since X is compact, one has $E \in D^b(X)$. By [Sta24, Tag 01BY (1)], every E_i^j is coherent. Therefore, every $H^j(E)|_{U_i}$ is coherent over \mathcal{O}_{U_i} . Thus, $H^j(E)$ is coherent over \mathcal{O}_X for all j . Hence $E \in D_c^b(X)$. In particular, E is in $D_{\text{good}}(X)$.

Let $E^\vee := R\mathcal{H}om(E, \mathcal{O}_X) \in D(X)$. From [Sta24, Tag 08DQ], there is a natural isomorphism of functors $\text{Hom}_{D(X)}(E, \cdot) \rightarrow H^0(X, E^\vee \otimes_{\mathcal{O}_X}^L \cdot) : D(X) \rightarrow \text{Ab}$. The functor $E^\vee \otimes_{\mathcal{O}_X}^L \cdot : D(X) \rightarrow D(X)$ commutes with direct sums. Since X is compact, $\dim X$ is finite. Then by Lemma B.4.0.7, the functor $H^0(X, \cdot) : D(X) \rightarrow \text{Ab}$ also commutes with direct sums. Therefore, E is a compact object of $D(X)$. By Lemma A.1.4.3 2, $D_{\text{good}}(X)$ is closed under direct sums in $D(X)$. Then E is also a compact object of $D_{\text{good}}(X)$. \square

Lemma B.4.0.7. *Let $f : X \rightarrow Y$ be a proper morphism of complex analytic spaces. If $\dim X$ is finite, then the functor $Rf_* : D(X) \rightarrow D(Y)$ commutes with direct sums.*

Proof. First, we prove that for every integer q , there is a natural isomorphism

$$R^q f_* \xrightarrow{\sim} R^q f_* \tau_{\geq q-2 \dim X} : D(X) \rightarrow \text{Mod}(\mathcal{O}_Y). \quad (\text{B.8})$$

Indeed, by [Sta24, Tag 08J5], for every object $E \in D(X)$, there is an exact triangle $\tau_{\leq q-2 \dim X-1} E \rightarrow E \rightarrow \tau_{\geq q-2 \dim X} E \rightarrow (\tau_{\leq q-2 \dim X-1} E)[1]$. It induces an exact sequence

$$R^q f_* \tau_{\leq q-2 \dim X-1} E \rightarrow R^q f_* E \rightarrow R^q f_* \tau_{\geq q-2 \dim X} E \rightarrow R^{q+1} f_* \tau_{\leq q-2 \dim X-1} E$$

in $\text{Mod}(\mathcal{O}_Y)$. From Lemma 5.3.1.11, one has

$$R^q f_* \tau_{\leq q-2 \dim X-1} E = R^{q+1} f_* \tau_{\leq q-2 \dim X-1} E = 0.$$

Hence an isomorphism $R^q f_* E \rightarrow R^q f_* \tau_{\geq -q-2 \dim X} E$ functorial in E .

Let $\{E_i : i \in I\}$ be a family of objects of $D(X)$. Set $E = \oplus_{i \in I} E_i$. To prove that the canonical morphism $\oplus_{i \in I} Rf_* E_i \rightarrow Rf_* E$ in $D(Y)$ is

an isomorphism, it suffices to show that for every integer q , the induced morphism $\oplus_{i \in I} R^q f_* E_i \rightarrow R^q f_* E$ in $\text{Mod}(O_Y)$ is an isomorphism. Since $\tau_{\geq q-2 \dim X} E = \oplus_{i \in I} \tau_{\geq q-2 \dim X} E_i$, by (B.8), one may assume that E and all the E_i are in $D^{\geq q-2 \dim X}(X)$. Then from [Sta24, Tag 015J], one has canonical spectral sequences

$$R^s f_* H^t(E) \Rightarrow R^{s+t} f_* E, \quad R^s f_* H^t(E_i) \Rightarrow R^{s+t} f_* E_i.$$

By Lemma 5.3.1.9, for any integers s and t , the canonical morphism $\oplus_{i \in I} R^s f_* H^t(E_i) \rightarrow R^s f_* H^t(E)$ in $\text{Mod}(O_Y)$ is an isomorphism. Consequently, the canonical morphism $\oplus_{i \in I} R^q f_* E_i \rightarrow R^q f_* E$ is an isomorphism. \square

Corollary B.4.0.8. *If the complex algebraic variety X is proper, then the functor $\psi_X^* : D_c(X) \rightarrow D_c(X^{\text{an}})$ is an equivalence of triangulated categories.*

Proof. For every $F \in D_c(X)$ and every integer i , the $O_{X^{\text{an}}}$ -module $H^i(\psi_X^* F) = \psi_X^* H^i(F)$ is coherent. Thus, the functors $\psi_X^* : D_c(X) \rightarrow D_c(X^{\text{an}})$ is well-defined. By Theorem B.4.0.2, the functor $\psi_X^* : D_c(X) \rightarrow D_c(X^{\text{an}})$ is fully faithful. For every $F \in D_c(X^{\text{an}})$, by Theorem B.4.0.2, there is $G \in D_{\text{qc}}(X)$ with $\psi_X^* G$ isomorphic to F . Then $\psi_X^* H^i(G) = H^i(\psi_X^* G) \xrightarrow{\sim} H^i(F)$ is coherent over $O_{X^{\text{an}}}$. By Fact B.2.0.1 and Proposition B.3.0.2, the O_X -module $H^i(G)$ is coherent. Hence $G \in D_c(X)$. Therefore, $\psi_X^* : D_c(X) \rightarrow D_c(X^{\text{an}})$ is essential surjective and hence an equivalence. \square

B.5 Compact objects

Corollary B.5.0.1. *Suppose that the complex algebraic variety X is proper. Then the compact objects of $D_{\text{good}}(X^{\text{an}})$ are precisely the perfect complexes in $D(X^{\text{an}})$.*

Proof. By compactness of X^{an} and Lemma B.4.0.6, perfect complexes are compact objects of $D_{\text{good}}(X^{\text{an}})$. Conversely, let F be a compact object of $D_{\text{good}}(X^{\text{an}})$. By Theorem B.4.0.2, there is a compact object $G \in D_{\text{qc}}(X)$ with $\psi_X^* G$ isomorphic to F . By [Sta24, Tag 09M1], G is a perfect complex in $D(X)$. By definition, F is a perfect complex in $D(X^{\text{an}})$. \square

Let X be a compact complex manifold.

Question B.5.0.2. Does the full subcategory of $D_{\text{good}}(X)$ of compact objects coincide with $D_c^b(X)$?

Question B.5.0.3. Is the category $D_{\text{good}}(X)$ compactly generated?

When X is the analytification of a smooth proper complex algebraic variety, Corollary B.5.0.1 (resp. Theorem B.4.0.2) answers Questions B.5.0.2 (resp. B.5.0.3) affirmatively.

Appendix C

Quasi-coherent sheaves on complex analytic spaces

C.1 Introduction

Let (X, O_X) be a ringed space. The category of O_X -modules is denoted by $\text{Mod}(O_X)$.

Definition C.1.0.1. An O_X -module F is called *quasi-coherent* if for every $x \in X$, there is an open neighborhood $U \subset X$ of x , two sets I, J and a morphism $O_U^{\oplus J} \rightarrow O_U^{\oplus I}$ with cokernel isomorphic to $F|_U$. The full subcategory of $\text{Mod}(O_X)$ of quasi-coherent modules is denoted by $\text{Qch}(X)$.

According to [Sta24, Tag 01BD], in general $\text{Qch}(X)$ is not an abelian category. By [Sta24, Tag 06YZ], if X is a scheme, then $\text{Qch}(X)$ is a weak Serre subcategory (in the sense of [Sta24, Tag 02MO (2)]) of $\text{Mod}(O_X)$. We show a complex analytic analog of this result, contrary to a guess made in [?].

Theorem C.1.0.2. *If X is a complex analytic space, then $\text{Qch}(X) \subset \text{Mod}(O_X)$ is a weak Serre subcategory. In particular, it is an abelian subcategory.*

Acknowledgment

I thank my supervisor, Anna Cadoret, for her constant support.

C.2 Preliminaries

Let (X, O_X) be a ringed space. Every $O_X(X)$ induces naturally a quasi-coherent module.

Example C.2.0.1 ([Sta24, Tag 01BI]). Let $f : (X, O_X) \rightarrow (\{*\}, O_X(X))$ be the morphism of ringed spaces, with $f : X \rightarrow \{*\}$ the unique map and $f_*^\sharp : O_X(X) \rightarrow O_X(X)$ the identity. Then f is flat. For an $O_X(X)$ -module M , its pullback f^*M is called the sheaf associated with M . This O_X -module is quasi-coherent. The functor $f^* : \text{Mod}(O_X(X)) \rightarrow \text{Mod}(O_X)$ is called *localization* and denoted by \sim .

From [Gro60, 4.1.1], on a scheme the direct sum of any family of quasi-coherent modules is quasi-coherent. It fails for complex manifolds, shown by Example C.2.0.2.

Example C.2.0.2. [hs] Let $X \subset \mathbb{C}$ be the unit open disk. For every integer $n \geq 2$, Gabber ([Con06, Eg. 2.1.6], see also Example A.2.0.5) constructs a locally free (hence quasi-coherent) O_X -module F_n of infinite rank, such that for every open subset $U \subset X$ containing $\{\pm 1/n\}$, one has $\Gamma(U, F_n) = 0$. We prove that $F := \bigoplus_{n \geq 2} F_n$ is not quasi-coherent.

Assume the contrary. Then there is an open neighborhood V of $0 \in X$, a set I and a quotient morphism $q : O_V^{\oplus I} \rightarrow F|_V$. There is an integer $N \geq 2$ with $\{\pm 1/N\} \subset V$. Let $p : F|_V \rightarrow F_N|_V$ be the quotient morphism. Because $\text{Hom}_{\text{Mod}(O_V)}(O_V, F_N|_V) = \Gamma(V, F_N) = 0$, the morphism $pq = 0$. However, it contradicts $F_N|_V \neq 0$.

Let X be a complex analytic space in the sense of [GR04, p.18]. For an inclusion $i : K \rightarrow X$ of a compact subset, let $O_K = i^{-1}O_X$. Then O_K is naturally a sheaf of rings on K .

Definition C.2.0.3. A compact subset $K \subset X$ is a *Stein compactum*, if K has a fundamental system of open neighborhoods that are Stein subspaces of X . A Stein compactum K is *Noetherian* if $O_K(K)$ is a Noetherian ring.

Fact C.2.0.4 ([Fri67, Thm. I, 9; Rem. I, 10]). Every $x \in X$ admits a neighborhood which is a Noetherian Stein compactum in X .

Lemma C.2.0.5. Let F be an O_X -module. Then the following conditions are equivalent:

1. ([BBBP07, Def. 5.1]) Every $x \in X$ admits a neighborhood K which is a Noetherian Stein compactum, such that $F|_K$ is associated with a $O_K(K)$ -module.
2. The O_X -module F is quasi-coherent.

Proof.

- Assume Condition 1. For every $x \in X$, take such a K and suppose that $F|_K$ is associated with a $O_K(K)$ -module M . There are sets I, J and an

exact sequence $O_K(K)^{\oplus I} \rightarrow O_K(K)^{\oplus J} \rightarrow M \rightarrow 0$ in the category of $O_K(K)$ -modules. By [Sta24, Tag 01BH], it induces an exact sequence $O_K^{\oplus I} \rightarrow O_K^{\oplus J} \rightarrow F|_K \rightarrow 0$ in $\text{Mod}(O_K)$. Then the O_{K° -module $F|_{K^\circ}$ is quasi-coherent. Thus, Condition 2 is proved.

- Assume Condition 2. Because X is locally compact Hausdorff, for every $x \in X$, by [Sta24, Tag 01BK], there is an open neighborhood $U \subset X$ of x such that $F|_U$ is associated with a $\Gamma(U, O_X)$ -module. From Fact C.2.0.4, there is a neighborhood K of $x \in U$ which is a Noetherian Stein compactum. By [Sta24, Tag 01BJ] applied to the morphism $(K, O_K) \rightarrow (U, O_U)$ of ringed spaces, $F|_K$ is associated with a $O_K(K)$ -module. Thus, Condition 1 is proved.

□

Lemma C.2.0.6. *Let K be a Noetherian Stein compactum in X .*

1. *The natural transformation $\text{Id} \rightarrow \Gamma(K, \cdot)$ of functors $\text{Mod}(O_K(K)) \rightarrow \text{Mod}(O_K(K))$ is an isomorphism.*
2. *The localization functor $\cdot^\sim : \text{Mod}(O_K(K)) \rightarrow \text{Mod}(O_K)$ is exact, fully faithful.*
3. *For every $O_K(K)$ -module M and every integer $q > 0$, one has $H^q(K, \tilde{M}) = 0$.*

Proof.

1. Let M be a $O_K(K)$ -module. We prove that the morphism $M \rightarrow \Gamma(K, \tilde{M})$ is an isomorphism. When M is finitely generated, the result follows from [Tay02, p.299]. Assume now that M is arbitrary. Let $\{M_i\}_{i \in I}$ be the family of all finitely generated submodules of M . This family is directed in the inclusion relation and

$$M = \sum_{i \in I} M_i. \quad (\text{C.1})$$

By [Sta24, Tag 01BH (4)], the localization functor preserves colimits. Therefore,

$$\tilde{M} = \text{colim}_{i \in I} \tilde{M}_i. \quad (\text{C.2})$$

By [God58, Thm. 4.12.1], one has

$$\Gamma(K, \tilde{M}) = \text{colim}_{i \in I} \Gamma(K, \tilde{M}_i) = \text{colim}_{i \in I} M_i = M.$$

2. The exactness is proved in [Tay02, Prop. 11.9.3 (ii)]. For any $M, N \in \text{Mod}(O_K(K))$, we prove that the natural morphism

$$\text{Hom}_{O_K(K)}(M, N) \rightarrow \text{Hom}_{O_K}(\tilde{M}, \tilde{N}) \quad (\text{C.3})$$

is an isomorphism.

Assume first that M is finitely generated. As the ring $O_K(K)$ is Noetherian, the $O_K(K)$ -module M is of finite presentation. Then by [GW20, Exercise 7.20 (b), p.205], one has $\widetilde{\text{Hom}_{O_K(K)}(M, N)} = \text{Hom}_{O_K}(\tilde{M}, \tilde{N})$. By Point 1, the morphism (C.3) is an isomorphism. Assume now that M is arbitrary. By (C.1) and (C.2), the morphism (C.3) is the inverse limit of the morphisms $\text{Hom}_{O_K(K)}(M_i, N) \rightarrow \text{Hom}_{O_K}(\tilde{M}_i, \tilde{N})$, each of which is an isomorphism.

3. When M is finitely generated, it follows from [Tay02, Prop. 11.9.2] and [Car57, Thm. 1 (B)]. Assume now that M is arbitrary. By (C.2) and [God58, Thm. 4.12.1], one has $H^q(K, \tilde{M}) = \text{colim}_i H^q(K, \tilde{M}_i) = 0$.

□

C.3 Proof of Theorem C.1.0.2

- For every morphism $f : F \rightarrow G$ in $\text{Qch}(X)$, we prove that $\ker(f), \text{coker}(f)$ in $\text{Mod}(O_X)$ lie in $\text{Qch}(X)$.

For every $x \in X$, by Lemma C.2.0.5, there is a neighborhood A (resp. B) of $x \in X$ which is a Noetherian Stein compactum and an $O_A(A)$ -module M (resp. $O_B(B)$ -module N), such that $F|_A$ (resp. $G|_B$) is associated with M (resp. N). By Fact C.2.0.4, there is a neighborhood C of $x \in A^\circ \cap B^\circ$ which is a Noetherian Stein compactum. From [Sta24, Tag 01BJ], $F|_C$ (resp. $G|_C$) is associated with $M \otimes_{O_A(A)} O_C(C)$ (resp. $N \otimes_{O_B(B)} O_C(C)$). By Lemma C.2.0.6 2, there is a morphism $\phi : M \otimes_{O_A(A)} O_C(C) \rightarrow N \otimes_{O_B(B)} O_C(C)$ in $\text{Mod}(O_C(C))$ whose localization is $f|_C : F|_C \rightarrow G|_C$. The restriction functor $\text{Mod}(O_X) \rightarrow \text{Mod}(O_{C^\circ})$ is exact, so $\ker(f)|_{C^\circ}$ (resp. $\text{coker}(f)|_{C^\circ}$) is the localization of $\ker(\phi \otimes_{O_C(C)} \text{Id}_{O_X(C^\circ)})$ (resp. $\text{coker}(\phi \otimes_{O_C(C)} \text{Id}_{O_X(C^\circ)})$) in $\text{Mod}(O_X(C^\circ))$. Therefore, the O_X -modules $\ker(f), \text{coker}(f)$ are quasi-coherent.

- Let

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0 \quad (\text{C.4})$$

be a short exact sequence in $\text{Mod}(O_X)$, with F', F'' quasi-coherent. We prove that F is quasi-coherent.

By Lemma C.2.0.5, for every $x \in X$, there is a neighborhood K' (resp. K'') of x which is a Noetherian Stein compactum, and an $O_{K'}(K')$ -module M' (resp. $O_{K''}(K'')$ -module M'') whose localization is $F'|_{K'}$ (resp. $F''|_{K''}$). By Fact C.2.0.4, there is a neighborhood K of $x \in K'^\circ \cap K''^\circ$ that is a Noetherian

Stein compactum. From [Sta24, Tag 01BJ], $F'|_K$ (resp. $F''|_K$) is associated with the $O_K(K)$ -module $M' \otimes_{O_{K'}(K')} O_K(K)$ (resp. $M'' \otimes_{O_{K''}(K'')} O_K(K)$).

Let $P = \Gamma(K, F)$. By Lemma C.2.0.6 1 and 3, the sequence (C.4) induces a short exact sequence in $\text{Mod}(O_K(K))$:

$$0 \rightarrow M' \otimes_{O_{K'}(K')} O_K(K) \rightarrow P \rightarrow M'' \otimes_{O_{K''}(K'')} O_K(K) \rightarrow 0.$$

From Lemma C.2.0.6 2, its localization induces a short exact sequence in $\text{Mod}(O_K)$:

$$0 \rightarrow M' \otimes_{O_{K'}(K')} \widetilde{O_K(K)} \rightarrow \tilde{P} \rightarrow M'' \otimes_{O_{K''}(K'')} \widetilde{O_K(K)} \rightarrow 0.$$

By restricting to K° and [Sta24, Tag 01BJ], one has a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' \otimes_{O_{K'}(K')} \widetilde{O_X(K^\circ)} & \longrightarrow & P \otimes_{O_K(K)} \widetilde{O_X(K^\circ)} & \longrightarrow & M'' \otimes_{O_{K''}(K'')} \widetilde{O_X(K^\circ)} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F'|_{K^\circ} & \longrightarrow & F|_{K^\circ} & \longrightarrow & F''|_{K^\circ} \longrightarrow 0 \end{array}$$

in $\text{Mod}(O_{K^\circ})$, where the vertical morphisms are given by the adjunction of $\tilde{\cdot} : \text{Mod}(O_X(K^\circ)) \rightarrow \text{Mod}(O_{K^\circ})$ and $\Gamma(K^\circ, \cdot) : \text{Mod}(O_{K^\circ}) \rightarrow \text{Mod}(O_X(K^\circ))$. The rows are exact, and the two outside vertical arrows are isomorphisms. By the five lemma, the middle vertical morphism is an isomorphism. By Example C.2.0.1, the O_{K° -module $F|_{K^\circ}$ is quasi-coherent. Consequently, F is quasi-coherent.

By [Sta24, Tag 0754], $\text{Qch}(X)$ is a weak Serre subcategory of $\text{Mod}(O_X)$.

Appendix D

Complex analytic geometry

D.1 Dimension of the fiber product

Section D.1 aims at understanding the dimension of the fiber product of algebraic varieties/complex analytic spaces. The proof in the analytic case is inspired by that in the algebraic case. Therefore, we begin with the algebraic situation.

D.1.1 Algebraic case

Fix a field k . Under flatness condition, the dimension of the fiber product behaves well.

Lemma D.1.1.1. *Let X, Y, Z be three schemes of finite type over k and $f : X \rightarrow Z$, $g : Y \rightarrow Z$ be k -morphisms. Assume that the schemes X, Z are irreducible, Y is equidimensional, and g is flat. Put $W = X \times_Z Y$. If W is nonempty, then W is equidimensional of dimension $\dim X + \dim Y - \dim Z$.*

Proof. Applying [Har77, Ch. III, Corollary 9.6] to the flat morphism g , we find that g is of relative dimension $\dim Y - \dim Z$. By virtue of [Sta24, Tag 02NK], its base change $W \rightarrow X$ is also flat of relative dimension $\dim Y - \dim Z$. Then the reverse direction of the cited [Har77, Ch. III, Corollary 9.6] shows that W is equidimensional of dimension $\dim Y - \dim Z + \dim X$. \square

In the proof of Proposition D.1.1.2, the general case is reduced to the case of a flat morphism.

Proposition D.1.1.2. *Let $X, Y, Z/k$ be three finite type schemes, $f : X \rightarrow Z$, $g : Y \rightarrow Z$ be dominant k -morphisms. Assume that X, Z are irreducible and Y is equidimensional, and put $W = X \times_Z Y$, then $\dim W + \dim Z \geq \dim X + \dim Y$.*

Proof. Since the reduction $Z_{\text{red}} \rightarrow Z$ is a universal homeomorphism, we may assume that Z is an integral scheme. By generic flatness [Sta24, Tag 052A],

there is a nonempty affine open subset $U \subset Z$ such that the restriction $g^{-1}(U) \rightarrow U$ is flat. By [Sta24, Tag 01UA], the morphism $g^{-1}(U) \rightarrow U$ is open. By shrinking U , we may assume further that $g^{-1}(U) \rightarrow U$ is surjective.

Because f is dominant, $f^{-1}(U)$ is a nonempty open subset of X . Therefore, by [Har77, Ch. II, Exercise 3.20 (e)] we have $\dim U = \dim Z$, $\dim f^{-1}(U) = \dim X$ and $g^{-1}(U)$ is equidimensional of dimension $\dim Y$. Hence, we may base change everything along $U \rightarrow Z$ which does not increase $\dim W$. In particular, we can assume that g is flat surjective. Then $W \rightarrow X$ is also flat surjective. In particular, $W \neq \emptyset$. We conclude by Lemma D.1.1.1. \square

Example D.1.1.3 shows that the inequality in Proposition D.1.1.2 can be strict.

Example D.1.1.3. If $f : X \rightarrow P_k^3$ is the blow up at a point $p \in P^3(k)$, then the morphism f is projective surjective, $\dim X = 3$, $\dim X \times_{P_k^3} X = 4$ and the defect of semismallness $r(f) = 1$.

Corollary D.1.1.4. Let $X, Y/k$ be two finite type schemes and $f : X \rightarrow Y$ be a k -morphism. If the scheme X is irreducible, then $\dim X \times_Y X \geq 2 \dim X - \dim \overline{f(X)}$, where $\overline{f(X)}$ is the Zariski closure of $f(X)$ in Y .

Proof. Because the reduction $X_{\text{red}} \rightarrow X$ is a universal homeomorphism, we may assume that X is reduced. Let $Z \rightarrow Y$ be the scheme theoretic image of f . By [Har77, Ch. II, Exercise 3.11 (d)], the induced morphism $X \rightarrow Z$ is dominant and the underlying topological space of Z is $\overline{f(X)}$. Therefore, Z is also irreducible. By magic square [Vak23, 1.3.S], the natural morphism $X \times_Z X \rightarrow X \times_Y X$ is the base change of the diagonal isomorphism $Z \rightarrow Z \times_Y Z$, hence also an isomorphism. By Proposition D.1.1.2, $\dim X \times_Y X = \dim X \times_Z X \geq 2 \dim X - \dim Z$. \square

D.1.2 Analytic case

The contents of this section is parallel to those of Section D.1.1. Lemma D.1.2.1 is an analogue of [Har77, III, Corollary 9.6], whose proof is also a direct adaptation. A complex analytic space is called equidimensional if every irreducible component is of same dimension.

Lemma D.1.2.1. Let $f : X \rightarrow Y$ be a flat morphism of complex analytic spaces, and assume that Y is irreducible. Then the following conditions are equivalent:

1. X is equidimensional of dimension $n + \dim Y$;
2. for every $y \in f(X)$, the fiber X_y is equidimensional of dimension n .

In that case, we say f is flat of relative dimension n .

Proof. Assume 1. Given $y \in f(X)$, let Z be an irreducible component of X_y . Because the set of irreducible components of a complex analytic space is locally finite, there is $x \in Z$ which is not in any other irreducible component of X_y . Applying [CD94, Proposition 2.11, p.113], we have $\dim_x Z + \dim_y Y = \dim_x X$. As Y, Z are irreducible hence pure dimensional, we have $\dim_y Y = \dim Y$ and $\dim_x Z = \dim Z$. Now that $\dim_x X = \dim Y + n$, we have $\dim Z = n$.

Conversely, assume 2. Let W be an irreducible component of X . Let $x \in W$ be a point which is not contained in any other irreducible component of X and $y = f(x)$. Then we have $\dim_x X = \dim W$ and $\dim_y Y = \dim Y$. Applying [CD94, Proposition 2.11, p.113], we obtain

$$\dim_x(X_y) + \dim_y Y = \dim_x X.$$

By assumption, $\dim_x(X_y) = n$. Thus $\dim W = \dim Y + n$ as required. \square

Lemma D.1.2.2 is similar to Lemma D.1.1.1.

Lemma D.1.2.2. *Let $f : X \rightarrow Z$, $g : Y \rightarrow Z$ be complex analytic space morphisms. Assume that X, Z are irreducible, Y is equidimensional, and g is flat. Put $W = X \times_Z Y$. If W is nonempty, then W is equidimensional of dimension $\dim X + \dim Y - \dim Z$.*

Proposition D.1.2.3 is the main result of Section D.1.

Proposition D.1.2.3. *Let X, Y, Z be irreducible complex analytic spaces. Let $f : X \rightarrow Z$, $g : Y \rightarrow Z$ be morphisms and put $W = X \times_Z Y$. If f is surjective and the (Euclidean) topology of X is second-countable, then $\dim W + \dim Z \geq \dim X + \dim Y$.*

Proof. Because reduction does not change the dimension [GR84, p.96], we may assume that X, Y, Z are reduced. Let $A = \{x \in X : f \text{ is not flat at } x\}$. By Frisch's theorem [CD94, Theorem 2.8, p.112], A is an analytic subset of X and $f(A) \neq Z$. Then $X \setminus f^{-1}(f(A)) \rightarrow Z \setminus f(A)$ is a surjective flat morphism. By shrinking X, Y, Z suitably, we may assume further that f is flat surjective. Then W is nonempty and we conclude by Lemma D.1.2.2. \square

The invariant $\dim X \times_Y X$ considered in Corollary D.1.2.4 appears in the definition of defect of semismallness (Definition 4.5.2.1).

Corollary D.1.2.4. *Let $f : X \rightarrow Y$ be a proper morphism of irreducible complex analytic spaces. If the (Euclidean) topology of X is second-countable, then $\dim X \times_Y X \geq 2 \dim X - \dim f(X)$.*

Proof. The image $Z := f(X)$ is an analytic subset of Y . Endow Z with the reduced structure of complex analytic space. Then Z is also irreducible and the morphism $f : X \rightarrow Z$ is surjective. The natural morphism $X \times_Z X \rightarrow X \times_Y X$ is an isomorphism. Then we conclude by Proposition D.1.2.3. \square

D.2 Connection on line bundles

The purpose of Section D.2 is to show Lemma D.2.0.4. For one thing, it is closely related to Corollary 4.4.2.2. For another, it implies the possibility to extend the Donaldson-Uhlenbeck-Yau theorem and nonabelian Hodge theory to manifolds more general than Kähler ones (Remarks D.2.0.6 and D.2.0.7). For work towards this direction, see [BD23], which extends nonabelian Hodge theory to Fujiki class \mathcal{C} manifolds.

We begin the proof with a variation of the classical maximum principle.

Proposition D.2.0.1. *Let $U \subset \mathbb{R}^n$ be a nonempty connected open subset, $f : U \rightarrow \mathbb{C}$ be a harmonic function. If $|f|$ attains its maximum in U , then f is constant.*

Lemma D.2.0.2 concerns the uniqueness of solution to $\bar{\partial}\partial$ -equation.

Lemma D.2.0.2. *Let $f : X^n \rightarrow \mathbb{C}$ be a smooth function on a compact connected complex manifold X with $\bar{\partial}\partial f = 0$, then f is constant.*

Proof. Since X is compact, the subset $A = \{x \in X : |f(x)| = \max_{t \in X} |f(t)|\}$ is nonempty closed in X . For any $p \in A$, there exists a local holomorphic coordinate $(U; z_1, \dots, z_n)$, where U is a connected open neighborhood of p in X . With this chart, we identify U as an open subset of \mathbb{C}^n . Since $\bar{\partial}\partial f = 0$, we have $\frac{\partial^2 f}{\partial \bar{z}_j \partial z_l} = 0$ for all $1 \leq j, l \leq n$. In particular, $\sum_{j=1}^n \frac{\partial^2 f}{\partial \bar{z}_j \partial z_j} = 0$, or equivalently, f is a harmonic function on U . By Proposition D.2.0.1, f is constant on U and so $U \subset A$. Therefore, A is open in X . By connectedness of X , $A = X$. So for any $p \in X$, f is locally constant near p . By connectedness of X again, f is constant. \square

Let X be a regular manifold for the rest of Section D.2.

We need a comparison between the Atiyah class ([Huy05, Def. 4.2.18]) and the first Chern class. For Kähler manifolds, it is [Ati57a, Prop. 12].

Lemma D.2.0.3. *Let X be a regular manifold. Let $L \rightarrow X$ be a holomorphic line bundle. Let $A(L) \in H^1(X, \Omega_X^1)$ be the Atiyah class of L . Then*

$$\frac{i}{2\pi} A(L) = c_1^{\mathbb{R}}(L)$$

in $H^2(X, \mathbb{R})$. In particular, L admits a holomorphic connection if and only if $L \in \text{Pic}^{\tau}(X)$.

Proof. By Corollary 4.3.1.4, we have a commutative diagram

$$\begin{array}{ccc} Z^{1,1}(X) & \hookrightarrow & Z^2(X) \\ \downarrow \phi & & \downarrow \psi \\ H^1(X, \Omega_X^1) & \xrightarrow{\iota^{1,1}} & H_{dR}^2(X; \mathbb{C}), \end{array}$$

where ϕ is taking Dolbeault cohomology class and ψ is taking de Rham cohomology class. Take a hermitian metric h on L . Let R be the corresponding Chern curvature form. By [Huy05, Corollary 4.4.5], $R \in Z^{1,1}(X)$. Then by [Huy05, Proposition 4.3.10], $A(L) = \phi(R)$ and $c_1^{\mathbb{R}}(L) = \frac{i}{2\pi}\psi(R)$. The equality follows. The second part follows from [Huy05, Proposition 4.2.19]. \square

Lemma D.2.0.4. *Let X be a regular manifold, $L \in \text{Pic}^{\tau}(X)$, then:*

1. *L admits a unique (up to a positive scalar) hermitian metric whose Chern connection is flat;*
2. *Every holomorphic connection on L is flat.*

Proof. 1. We begin with the existence. By Corollary 4.4.2.2 2, there is a unitary local system $\mathcal{L} \in \text{Loc}^{u,1}(X)$ on X with $\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_X = L$. Applying Theorem 4.2.3.1 the existence of such metric follows.

Now for uniqueness. Let h, h' be two hermitian metrics whose respective Chern connections ∇, ∇' are flat holomorphic connections. By Theorem 4.2.3.1, $\ker(\nabla), \ker(\nabla') \in \text{Loc}^{u,1}(X)$ have the same induced line bundle. By Corollary 4.4.2.2 2, $\ker(\nabla) = \ker(\nabla')$ in $\text{Loc}^{u,1}(X)$. The hermitian metrics h, h' restrict to two monodromy invariant hermitian forms on the common local system $\ker(\nabla)$. Moreover, by Theorem 4.2.3.1 one can recover the hermitian metric on the line bundle L from the restricted hermitian form on the local system. Since this local system is of rank 1, at one stalk these two hermitian forms differ by a scalar. Globally they differ by this scalar as they are monodromy invariant. Then the metrics h, h' also differ by a scalar.

2. By Lemma D.2.0.3 and [Huy05, Prop. 4.2.19], L admits a holomorphic connection. We show that the curvature forms (which are global holomorphic 2 forms) of different holomorphic connections on L are the same. In fact, for two such connections D, D' on L , by [Huy05, p.179], $D' - D \in H^0(X, \Omega_X^1)$. This form is d -closed by [Uen06, Corollary 9.5, p.101]. By [Huy05, Lemma 4.3.4], the curvature of D' equals that of D .

We adopt the argument in [BK09, Footnote (6), p.388]. By Cartan-Serre theorem [Car53, Théorème], the complex vector space $H^0(X, \Omega_X^2)$ is finite dimensional. Taking the curvature form of one (hence every) holomorphic connection on elements of $\text{Pic}^0(X)$, we get a holomorphic map $\text{Pic}^0(X) \rightarrow H^0(X, \Omega_X^2)$. As the complex torus $\text{Pic}^0(X)$ is compact connected, this map is constant. The canonical connection on the trivial line bundle $\mathcal{O}_X(\in \text{Pic}^0(X))$ is flat, so this map is constantly zero. In other words, for every $K \in \text{Pic}^0(X)$, any holomorphic connection on K is flat.

As $L \in \text{Pic}^\tau(X)$, there is an integer $n \geq 1$ such that $L^{\otimes n} \in \text{Pic}^0(X)$. Take a holomorphic connection on L of curvature form R , then it induces a holomorphic connection on $L^{\otimes n}$ of curvature form nR . As $nR = 0$, it holds that $R = 0$. The flatness follows from the first paragraph and the existence in Point 1. \square

Remark D.2.0.5. Here is a second proof of Lemma D.2.0.4 1. Take a hermitian metric h on L . Locally its Chern curvature is given by $\nabla = d + h^{-1}\partial h$. More precisely, let s be a local holomorphic frame for L , and by abuse of notation let h be the local (smooth positive) function $h(s, s)$. Then $\nabla(s) = (h^{-1}\partial h) \otimes s$ and the Chern curvature form $R = \bar{\partial}(h^{-1}\partial h)$ is a d -closed smooth $(1, 1)$ -form whose de Rham class is 0. Moreover iR is a real form. (This is part of Chern-Weil theory, see [Huy05, Proposition 4.3.8 (iii); 4.3.10 and p.196].) Therefore, by Fact 4.3.1.2, there is a smooth function $f : X \rightarrow \mathbb{R}$ with

$$R + \bar{\partial}\partial f = 0. \quad (\text{D.1})$$

Define a new hermitian metric h' by

$$h'(s, s) = e^f h(s, s). \quad (\text{D.2})$$

Then the new Chern connection is given by $\nabla'(s) = \nabla(s) + (\partial f) \otimes s$. The new curvature form $R' = R + \bar{\partial}\partial f = 0$, i.e., the new Chern connection is flat and compatible with the holomorphic structure, hence a holomorphic connection.

So far we have established the existence of such metric. As for uniqueness, any hermitian metric h' with flat Chern connection is in the form of (D.2) where f is a solution to (D.1). Lemma D.2.0.2 shows that such a solution f is unique up to addition by constant. So such metric h' is unique up to a positive scalar.

Remark D.2.0.6. When X is a compact Kähler manifold, Lemma D.2.0.4 1 is a consequence of known results. In fact [Kob87, Proposition 5.7.7 (a)] shows a holomorphic line bundle is slope stable. By Donaldson-Uhlenbeck-Yau theorem [UY86, Corollary 8.1, p.292], there is $\mathcal{L} \in \text{Loc}^{u,1}(X)$ such that $L = \mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_X$, and \mathcal{L} induces such a metric via Theorem 4.2.3.1. For any such hermitian metric, its Chern connection is a Hermitian-Yang-Mills connection. The uniqueness of such metric is mentioned in [Bea92, (3.2) c)] and follows from [UY86, Theorem, p.262] and [Che22, Corollary 2.18].

Remark D.2.0.7. Lemma D.2.0.4 can be viewed as a step toward nonabelian Hodge theory on regular manifolds. In fact, a semisimple local system on a compact Kähler manifold is unitary if and only if the associated Higgs bundle (E, θ) has $\theta = 0$ ([Sim92, Example p.21]). The metric given by Lemma D.2.0.4 is exactly the harmonic metric provided by Corlette Theorem [GRR15, Theorem 1, p.151].

D.3 Jacobi inversion theorem

In this section, we give a refinement of Proposition 4.4.1.2 3.

Lemma D.3.0.1. *For a pointed regular manifold (X, x_0) , for every $n \geq h^{1,0}(X)$, the holomorphic map $f_n : X^n \rightarrow \text{Alb}(X)$ defined by $(x_1, \dots, x_n) \mapsto \sum_{i=1}^n \alpha_{x_0}(x_i)$ is surjective.*

When X is a compact Riemann surface, then Lemma D.3.0.1 reduces to (part of) Jacobi inversion theorem in [GH78, p.235].

Two proofs are provided. They are inspired by [Voi02, Lemma 12.11] and [BL04, Proposition 11.11.8] respectively, but with an extra attention to the feasible range of n . The first proof is shorter, while the second proof provides a stronger result, Lemma D.3.2.1.

D.3.1 First proof

Lemma D.3.1.1. *Let X be a compact complex manifold. Then there is a subset $S \subset X$ with $\#S \leq h^{1,0}(X)$ such that, for any $\eta \in H^0(X, \Omega_X^1)$ with $\eta(x) = 0$ in the (holomorphic) cotangent space $(T_x^h X)^\vee$, we have $\eta = 0$.*

Proof. For every $x \in X$, let V_x be the subspace $\{\eta \in H^0(X, \Omega_X^1) : \eta(x) = 0\}$ of $H^0(X, \Omega_X^1)$. Then $\cap_{x \in X} V_x = \{0\}$. Hence, there is a subset $S \subset X$ with $\#S \leq h^{1,0}(X)$ and $\cap_{x \in S} V_x = \{0\}$. \square

Here is the first proof.

First proof of Lemma D.3.0.1. Consider the cotangent map $(d_p f_n)^* : (T_{f_n(p)}^h \text{Alb}(X))^\vee \rightarrow (T_p^h X^n)^\vee$ at $p = (p_1, \dots, p_n) \in X^n$. Since the cotangent bundle $\Omega_{\text{Alb}(X)}^1$ is trivial, this map is identified with the composition

$$H^0(\text{Alb}(X), \Omega_{\text{Alb}(X)}^1) \rightarrow (T_{f_n(p)}^h \text{Alb}(X))^\vee \rightarrow \prod_{i=1}^n (T_{p_i}^h X)^\vee.$$

By Proposition 4.4.1.2 4, it is further identified with the natural map

$$H^0(X, \Omega_X^1) \rightarrow \prod_{i=1}^n (T_{p_i}^h X)^\vee. \quad (\text{D.3})$$

By Lemma D.3.1.1, there exist $n_0 \leq h^{1,0}(X)$ and $x = (x_1, \dots, x_{n_0}) \in X^{n_0}$ such that for any $\eta \in H^0(X, \Omega_X^1)$ with $\eta(x_i) = 0$ for all i , we have $\eta = 0$. Then for every $n \geq n_0$, the map (D.3) is injective when $p = (x, x_0, \dots, x_0)$. Or equivalently, f_n is a submersion of smooth manifolds near p . From local normal form theorem, the image $f_n(X^n)$ contains a nonempty open subset of $\text{Alb}(X)$. By Remmert theorem [Whi72, Theorem 4A, p.150], $f_n(X^n)$ is an analytic subset of $\text{Alb}(X)$. By [GR84, Theorem, p.168], $f_n(X^n) = \text{Alb}(X)$, i.e., f_n is surjective. \square

D.3.2 Second proof

To certain extent, Lemma D.3.2.1 shows that a generating subset of a complex torus generates the complex torus “uniformly”.

Lemma D.3.2.1. *Let A be a g -dimensional commutative complex Lie group. Let M be a compact irreducible analytic subset of A containing 0. If the complex Lie subgroup of A generated by M is A , then for every integer $n \geq g$, the map $f_n : M^n \rightarrow A$ defined by $(x_1, \dots, x_n) \mapsto \sum_{i=1}^n x_i$ is surjective. In particular, A is a complex torus.*

Proof. Since M is connected, the identity component of A contains M . Therefore, A is connected.

The statement is true when $g = 0$. So we assume $g > 0$, then $M \neq \{0\}$ and hence $\dim M \geq 1$. For every $n \geq 1$, let $A_n = f_n(M^n)$, which is an analytic subset of A by Remmert theorem [Whi72, Theorem 4A, p.150]. Since $f_1 : M \rightarrow A$ is the inclusion, we find $A_1 = M \ni 0$. For every $x \in M^n$, $f_{n+1}(x, 0) = f_n(x)$, so $A_n \subset A_{n+1}$, hence an increasing sequence of analytic subsets of A :

$$A_1 \subset A_2 \subset \dots$$

Consider the integer sequence of analytic dimensions $\{\dim_0 A_n\}_{n \geq 1}$. By [GR84, p.96], this sequence is non-decreasing and bounded above by $\dim_0 A = g$. Therefore, there is $n_0 \leq g$ such that $\dim_0 A_{n_0} = \dim_0 A_{n_0+1}$.

By assumption, M^n is an irreducible complex analytic space. By [CD94, (14.14), p.89], the complex analytic space A_n is irreducible and pure dimensional for every $n \geq 1$ and $A_{n_0} = A_{n_0+1}$.

We claim that for every $m > n_0$, $A_{n_0} = A_m$.

We prove the claim by induction on m . It holds when $m = n_0 + 1$. If it is true for $m - 1$ with $m \geq n_0 + 2$, then for every $(x_1, \dots, x_m) \in M^m$, $\sum_{i=1}^{m-1} x_i \in A_{m-1} = A_{n_0}$, so there is $(p_1, \dots, p_{n_0}) \in M^{n_0}$ with $\sum_{j=1}^{n_0} p_j = \sum_{i=1}^{m-1} x_i$. Then

$$\sum_{i=1}^m x_i = x_m + \sum_{j=1}^{n_0} p_j \in A_{n_0+1} = A_{n_0}.$$

Therefore, $A_m = A_{n_0}$. The induction is completed.

For every $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in M^n$, we have $f_n(x) + f_n(y) = f_{2n}(x, y)$, so $A_n + A_n \subset A_{2n}$. In particular, $A_{n_0} + A_{n_0} \subset A_{2n_0} = A_{n_0}$. This shows A_{n_0} is closed under addition.

We are going to show that $-A_{n_0} = A_{n_0}$ and then A_{n_0} would be a subgroup of A .

As A is commutative connected, by [AK01, Proposition 1.1.2], its universal covering is in the form of $\pi : \mathbb{C}^g \rightarrow A$ and the lattice $\ker(\pi)$ is identified with the fundamental group $\pi_1(A, 0)$. As π is locally

biholomorphic and every A_n is irreducible, the preimage $\pi^{-1}(A_n)$ is an analytic subset of \mathbb{C}^g , every irreducible component of whom is of dimension $\dim A_n$. Any two different irreducible components are disjoint and differ by a translation by an element of $\ker(\pi)$.

Let V_n be the unique irreducible component of $\pi^{-1}(A_n)$ containing 0, then $\pi(V_n) = A_n$. Fix an integer $k \geq 1$ and let $[k] : A \rightarrow A$ be the multiplication by k . As A_{n_0} is closed under addition, we get $[k]A_{n_0} \subset A_{n_0}$. As π is a group morphism, we have $k \cdot \pi^{-1}(A_{n_0}) \subset \pi^{-1}(A_{n_0})$. As $k : \mathbb{C}^g \rightarrow \mathbb{C}^g$ is biholomorphic, kV_{n_0} is an irreducible analytic subset of \mathbb{C}^g isomorphic to V_{n_0} . As $0 \in kV_n$, we have $kV_{n_0} \subset V_{n_0}$. As $\dim kV_{n_0} = \dim V_{n_0}$, by [CD94, (14.14), p.89] again, we get $kV_{n_0} = V_{n_0}$, i.e., the morphism $k : V_{n_0} \rightarrow V_{n_0}$ is biholomorphic.

For every $x(\neq 0) \in A_{n_0}$, we check that $-x \in A_{n_0}$. In fact, take $v \in V_{n_0} \cap \pi^{-1}(x)$. Then $v \neq 0$. By last paragraph, $v/k \in V_{n_0}$ for every $k \geq 1$. Let l be the complex line in \mathbb{C}^g spanned by v . By the identity theorem for holomorphic functions on l , the smallest analytic subset of l containing $\{v/k\}_{k \geq 1}$ is l . Now that $V_{n_0} \cap l$ is an analytic subset of l containing $\{v/k\}_{k \geq 1}$, we get $l = V_{n_0} \cap l \subset V_{n_0}$. In particular, $-v \in V_{n_0}$ and then $-x \in A_{n_0}$ as desired.

So far we have shown that A_{n_0} is a subgroup of A that is a complex analytic subset. By Corollary F.2.0.5, A_{n_0} is an embedded complex Lie subgroup of A . By assumption, $A_{n_0} = A$. From the claim we get the surjectivity of f_n for every $n \geq n_0$. In particular, A is compact, hence a complex torus. \square

Example D.3.2.2. In Lemma D.3.2.1, we cannot remove the condition that $0 \in M$. For example, consider $A = \mathbb{C}^*$ and $M = \{2\}$. The irreducibility of M is also necessary. For instance, take A to be the elliptic curve $\mathbb{C}/\mathbb{Z}[i]$, $M = \{0, x\}$, where $x \in A \setminus A_{\text{tor}}$. Then f_n is not surjective for all integers $n \geq 1$.

Second proof of Lemma D.3.0.1. Let $M = \alpha_{x_0}(X)$, which is an irreducible analytic subset of $\text{Alb}(X)$ by Remmert theorem [Whi72, Theorem 4A, p.150] and [CD94, (14.14), p.89]. In addition, $0 \in M$. The proof is completed by citing Proposition 4.4.1.2 3 and Lemma D.3.2.1. \square

Appendix E

D -modules

E.1 Unbounded Bernstein's equivalence

In Section E.1, let X be a smooth algebraic variety over an algebraically closed field k of characteristic 0. Let $\mathrm{Qch}(O_X) \subset \mathrm{Mod}(O_X)$ and $\mathrm{Mod}_{\mathrm{qc}}(D_X) \subset \mathrm{Mod}(D_X)$ be the full subcategories of objects quasi-coherent over O_X . They are weak Serre subcategories.

Fact E.1.0.1 (Bernstein, [B⁺87, VI, Thm. 2.10]). *The natural functor*

$$\iota'_X : D^b(\mathrm{Mod}_{\mathrm{qc}}(D_X)) \rightarrow D_{\mathrm{qc}}^b(D_X)$$

is an equivalence.

Remark E.1.0.2. The first sentence of the proof in [B⁺87] needs (implicitly) [Mur07, Remark 64] and Fact E.1.0.3.

Fact E.1.0.3 can be proved as [B⁺87, I, Prop. 12.8; VI, Prop. 1.14].

Fact E.1.0.3. *Let B be a weak Serre subcategory of an abelian category A . Then the full class $\mathrm{Ob}(B)$ of objects in B is a generating class of $D_B^b(A)$ (defined in [Sta24, Tag 06UP]) in the sense of [B⁺87, I, Def. 12.4].*

Theorem E.1.0.4 is an unbounded generalization of Fact E.1.0.1. It is left “to the reader to state and prove” in [Nee96, p.207]. We follow the strategy pointed out in [gh], and do not claim originality here.

Theorem E.1.0.4. *The functor*

$$\iota'_X : D(\mathrm{Mod}_{\mathrm{qc}}(D_X)) \rightarrow D_{\mathrm{qc}}(D_X) \tag{E.1}$$

induced by the inclusion $\mathrm{Mod}_{\mathrm{qc}}(D_X) \rightarrow \mathrm{Mod}(D_X)$ is an equivalence of categories.

We need a series of lemmas for the proof of Theorem E.1.0.4. For an open immersion $j : U \rightarrow X$, we have a natural morphism of ringed spaces $j : (U, D_U) \rightarrow (X, D_X)$. From [B⁺87, VI, 5.2] and [HT07, Prop. 1.5.29], the functor $j_+ : D(D_U) \rightarrow D(D_X)$ is the right derived functor of the corresponding (left exact) direct image $j_* : \text{Mod}(D_U) \rightarrow \text{Mod}(D_X)$. By [Ber83, 2, p.12] and [Sta24, Tag 0096], the inverse image $j^* : \text{Mod}(D_X) \rightarrow \text{Mod}(D_U)$ is left adjoint to j_* . Lemma E.1.0.5 2 helps to construct a quasi-inverse to (E.1).

Lemma E.1.0.5.

1. *The category $\text{Mod}_{\text{qc}}(D_X)$ is locally noetherian.*
2. *The inclusion functor $\iota' : \text{Mod}_{\text{qc}}(D_X) \rightarrow \text{Mod}(D_X)$ admits a right adjoint $Q' = Q'_X : \text{Mod}(D_X) \rightarrow \text{Mod}_{\text{qc}}(D_X)$. The unit natural transform $\eta' : \text{Id}_{\text{Mod}_{\text{qc}}(D_X)} \rightarrow Q'\iota'$ is an isomorphism.*

Proof. By [Sta24, Tag 01LA (4)], $\text{Qch}(O_X) \subset \text{Mod}(O_X)$ is an abelian subcategory closed under colimits. Then so is $\text{Mod}_{\text{qc}}(D_X) \subset \text{Mod}(D_X)$.

1. When X is affine, by [HT07, Prop. 1.4.4 (ii)], the functor $\Gamma(X, \cdot) : \text{Mod}_{\text{qc}}(D_X) \rightarrow \text{Mod}(D_X(X))$ is an equivalence of abelian categories. As the ring $D_X(X)$ is left noetherian, the category $\text{Mod}(D_X(X))$ is locally noetherian by the last paragraph of [Gab62, p.402].

For a general X , one may assume that there exists an open covering $X = U \cup V$, such that the statement holds for U and V . Arguing as in [Gab62, Prop. 2, p.441], one can prove that $\text{Mod}_{\text{qc}}(D_X)$ is the gluing of $\text{Mod}_{\text{qc}}(D_U)$ and $\text{Mod}_{\text{qc}}(D_V)$ along $\text{Mod}_{\text{qc}}(D_{U \cap V})$ in the sense of [Gab62, VI. 1]. Let $j : U \rightarrow X$ be the inclusion. Then

$$j^* : \text{Mod}_{\text{qc}}(D_X) \rightarrow \text{Mod}_{\text{qc}}(D_U)$$

is exact and left adjoint to

$$j_* : \text{Mod}_{\text{qc}}(D_U) \rightarrow \text{Mod}_{\text{qc}}(D_X).$$

The (counit) natural transformation $\epsilon : j^*j_* \rightarrow \text{Id}_{\text{Mod}_{\text{qc}}(D_U)}$ is an isomorphism. From [Gab62, Prop. 5, p.374], the subcategory $\ker(j^*)$ is localizing in $\text{Mod}_{\text{qc}}(D_X)$ (in the sense of [Gab62, p372]) and j^* induces an equivalence

$$\text{Mod}_{\text{qc}}(D_X) / \ker(j^*) \rightarrow \text{Mod}_{\text{qc}}(D_U).$$

A similar result holds for V . Then by [Gab62, Lem. 2, p.442], the gluing category $\text{Mod}_{\text{qc}}(D_X)$ is locally noetherian.

2. It follows from 1 and Lemma E.1.0.7.

□

Remark E.1.0.6. For an affine (possibly singular) variety V , by [GR14, 4.7.1; 5.5], the abelian category $\text{Mod}_{\text{qc}}(D_V)$ is still Grothendieck.

Lemma E.1.0.7. *Let \mathcal{A} be a Grothendieck abelian category. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a functor preserving all colimits.*

1. *Then F admits a right adjoint $G : \mathcal{B} \rightarrow \mathcal{A}$.*
2. *If further F is fully faithful, then the unit natural transformation $\eta : \text{Id}_{\mathcal{A}} \rightarrow GF$ is an isomorphism.*

Proof. 1. Let Set be the category of sets. For each object $Y \in \mathcal{B}$, consider the functor

$$\text{Hom}_{\mathcal{B}}(F(\cdot), Y) : \mathcal{A}^{\text{op}} \rightarrow \text{Set}.$$

It transforms colimits into limits. Then by [Sta24, Tag 07D7], it is representable. From [ML78, Cor. 2, p.85], the functor F admits a right adjoint.

2. If follows from Yoneda's lemma.

□

By [Sta24, Tag 077P (2)], the inclusion $\iota = \iota_X : \text{Qch}(O_X) \rightarrow \text{Mod}(O_X)$ admits a right adjoint $Q_X = Q : \text{Mod}(O_X) \rightarrow \text{Qch}(O_X)$, called the *coherator* of X . To reduce the problem to the study of O_X -modules, consider the square

$$\begin{array}{ccc} \text{Mod}(D_X) & \xrightarrow{Q'_X} & \text{Mod}_{\text{qc}}(D_X) \\ \downarrow \text{for}_X & & \downarrow \text{for}_X \\ \text{Mod}(O_X) & \xrightarrow{Q_X} & \text{Qch}(O_X), \end{array} \quad (\text{E.2})$$

where the vertical functors are forgetful.

Lemma E.1.0.8. *Suppose that X is affine. Write $R = \Gamma(X, D_X)$. Then:*

1. *The functor $\tilde{\cdot} := D_X \otimes_R \cdot : \text{Mod}(R) \rightarrow \text{Mod}(D_X)$ is left adjoint to the global section functor $\Gamma(X, \cdot) : \text{Mod}(D_X) \rightarrow \text{Mod}(R)$;*
2. *The square (E.2) is commutative.*

Proof.

1. Let $(\sigma, \sigma^\#) : (X, D_X) \rightarrow (\{*\}, R)$ be the morphism of ringed spaces, with $\sigma : X \rightarrow \{*\}$ the unique map and $\sigma^\#$ given by Id_R . Then $\Gamma(X, \cdot) = \sigma_* : \text{Mod}(D_X) \rightarrow \text{Mod}(R)$. By [Sta24, Tag 01BH], the functor $\tilde{\cdot} = \sigma^*$. The adjunction follows from [Sta24, Tag 0096].

2. From 1 and [HT07, Prop. 1.4.4 (ii)], the functor $Q' : \text{Mod}(D_X) \rightarrow \text{Mod}_{\text{qc}}(D_X)$ is the composition of $\Gamma(X, \cdot) : \text{Mod}(D_X) \rightarrow \text{Mod}(R)$ with $\tilde{\cdot} : \text{Mod}(R) \rightarrow \text{Mod}_{\text{qc}}(D_X)$. The largest rectangle in the following diagram

$$\begin{array}{ccccccc}
 & & Q' & & & & \\
 & \swarrow & & \searrow & & & \\
 \text{Mod}(D_X) & \xrightarrow{\Gamma(X, \cdot)} & \text{Mod}(R) & \xrightarrow{D_X \otimes_R \cdot} & \text{Mod}_{\text{qc}}(D_X) & \xrightarrow{\Gamma(X, \cdot)} & \text{Mod}(R) \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \text{Mod}(O_X) & \xrightarrow{\Gamma(X, \cdot)} & \text{Mod}(O_X(X)) & \xrightarrow{O_X \otimes_{O_X(X)} \cdot} & \text{Qch}(O_X) & \xrightarrow{\Gamma(X, \cdot)} & \text{Mod}(O_X(X)) \\
 & \searrow & & \swarrow & & & \\
 & & Q & & & &
 \end{array}$$

is same as the small square on the left, hence commutative. Moreover, the two horizontal functors $\Gamma(X, \cdot)$ on the right are equivalences, so Q' is compatible with Q .

□

The abelian categories $\text{Mod}(D_X)$ and $\text{Mod}(O_X)$ are Grothendieck. By [Sta24, Tag 079P] and [Sta24, Tag 070K], the functor $Q' : \text{Mod}(D_X) \rightarrow \text{Mod}_{\text{qc}}(D_X)$ and $Q : \text{Mod}(O_X) \rightarrow \text{Qch}(O_X)$ admit right derived functors $RQ' : D(D_X) \rightarrow D(\text{Mod}_{\text{qc}}(D_X))$ and $RQ : D(O_X) \rightarrow D(\text{Qch}(O_X))$.

Lemma E.1.0.9. 1. The square (E.2) is commutative.

2. The square

$$\begin{array}{ccc}
 D(D_X) & \xrightarrow{RQ'_X} & D(\text{Mod}_{\text{qc}}(D_X)) \\
 \downarrow \text{for}_X & & \downarrow \text{for}_X \\
 D(O_X) & \xrightarrow{RQ_X} & D(\text{Qch}(O_X)),
 \end{array}$$

is commutative.

Proof.

1. We deduce a formula for Q'_X . Since X is quasi-compact, there is a finite cover $\{U_\alpha\}_{\alpha \in I}$ of X by affine opens. For any $\alpha \neq \beta$ in I , since X is separated over k , the scheme $U_{\alpha\beta} := U_\alpha \cap U_\beta$ is affine. Denote all the various open immersions $U_{\alpha\beta} \rightarrow X$ and $U_\alpha \rightarrow X$ as j . For

every D_X -module F , the sheaf axiom gives an equalizer diagram in $\text{Mod}(D_X)$:

$$0 \rightarrow F \rightarrow \oplus_{\alpha} j_*(F|_{U_{\alpha}}) \rightrightarrows \oplus_{(\alpha, \beta)} j_*(F|_{U_{\alpha\beta}}),$$

where the two right morphisms are induced by the inclusions $U_{\alpha\beta} \rightarrow U_{\alpha}$ and $U_{\alpha\beta} \rightarrow U_{\beta}$. By Lemma E.1.0.10, it induces another equalizer diagram in $\text{Mod}_{\text{qc}}(D_X)$:

$$0 \rightarrow Q'_X F \rightarrow \oplus_{\alpha} j_* Q'_{U_{\alpha}}(F|_{U_{\alpha}}) \rightrightarrows \oplus_{(\alpha, \beta)} j_* Q'_{U_{\alpha\beta}}(F|_{U_{\alpha\beta}}). \quad (\text{E.3})$$

There is a natural transformation $\iota' Q'_X \rightarrow \text{Id}_{\text{Mod}(D_X)} : \text{Mod}(D_X) \rightarrow \text{Mod}(D_X)$. Applying $\text{for}_X : \text{Mod}(D_X) \rightarrow \text{Mod}(O_X)$, one gets a natural transformation $\text{for}_X \circ \iota' \circ Q'_X \rightarrow \text{for}_X : \text{Mod}(D_X) \rightarrow \text{Mod}(O_X)$. Since $\text{for}_X \circ \iota' = \iota \circ \text{for}_X : \text{Mod}_{\text{qc}}(D_X) \rightarrow \text{Mod}(O_X)$ and Q_X is right adjoint to ι , there is a natural transformation

$$\mu_X : \text{for}_X \circ Q'_X \rightarrow Q_X \circ \text{for}_X$$

of functors $\text{Mod}(D_X) \rightarrow \text{Qch}(O_X)$. By Lemma E.1.0.8 2, it is an isomorphism when X is affine.

For a general X , by (E.3) and [TT90, (B.14.2)], there is a commutative diagram of functors $\text{Mod}(D_X) \rightarrow \text{Qch}(O_X)$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{for}_X Q'_X & \longrightarrow & \oplus_{\alpha} j_* \text{for}_{U_{\alpha}} Q'_{U_{\alpha}}(\cdot|_{U_{\alpha}}) & \rightrightarrows & \oplus_{(\alpha, \beta)} j_* \text{for}_{U_{\alpha\beta}} Q'_{U_{\alpha\beta}}(\cdot|_{U_{\alpha\beta}}) \\ & & \downarrow \mu_X & & \downarrow & & \downarrow \\ 0 & \longrightarrow & Q_X \text{for}_X & \longrightarrow & \oplus_{\alpha} j_* Q_{U_{\alpha}} \text{for}_{U_{\alpha}}(\cdot|_{U_{\alpha}}) & \rightrightarrows & \oplus_{(\alpha, \beta)} j_* Q_{U_{\alpha\beta}} \text{for}_{U_{\alpha\beta}}(\cdot|_{U_{\alpha\beta}}), \end{array}$$

where the two vertical arrows on the right are isomorphisms. Therefore, μ_X is an isomorphism.

2. The morphism $(X, D_X) \rightarrow (X, O_X)$ of ringed spaces is flat, and the direct image functor is the forgetful functor $\text{for}_X : \text{Mod}(D_X) \rightarrow \text{Mod}(O_X)$. By [Sta24, Tag 08BJ], it preserves K-injective complexes. The conclusion follows from Point 1, Lemma E.1.0.11 and [Sta24, Tag 070K].

□

Lemma E.1.0.10. *Let $j : U \rightarrow X$ be an open immersion. Then the natural transformation $j_* \circ Q'_U \rightarrow Q'_X \circ j_* : \text{Mod}(D_U) \rightarrow \text{Mod}_{\text{qc}}(D_X)$ is an isomorphism.*

Proof. As $j^* : \text{Mod}(D_X) \rightarrow \text{Mod}(D_U)$ restricts to a functor $\text{Mod}_{\text{qc}}(D_X) \rightarrow \text{Mod}_{\text{qc}}(D_U)$, one has $\iota'_U j^* = j^* \iota'_X$ as functors $\text{Mod}_{\text{qc}}(D_X) \rightarrow \text{Mod}(D_U)$. The functor $j_* : \text{Mod}(D_U) \rightarrow \text{Mod}(D_X)$ regards the direct image $j_* : \text{Mod}(O_U) \rightarrow \text{Mod}(O_X)$, so it also restricts to a functor $\text{Mod}_{\text{qc}}(D_U) \rightarrow \text{Mod}_{\text{qc}}(D_X)$. As Q' is right adjoint to ι' and j_* is right adjoint to j^* , the isomorphism follows. \square

Lemma E.1.0.11. *Let $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{C}$ be left exact functors of abelian categories. Assume that \mathcal{A}, \mathcal{B} are Grothendieck. If for ever K -injective complex I over \mathcal{A} , the natural morphism $GF(I) \rightarrow RG(F(I))$ in $D(\mathcal{C})$ is an isomorphism,¹ then the canonical natural transformation (constructed in [Sta24, Tag 05T2 (1)]) $t : R(G \circ F) \rightarrow RG \circ RF$ is an isomorphism of functors from $D(\mathcal{A}) \rightarrow D(\mathcal{C})$.*

Proof. Let A be a complex over \mathcal{A} . As \mathcal{A} is Grothendieck, by [Sta24, Tag 079P], there is a quasi-isomorphism $A \rightarrow I$ such that I is a K -injective complex. By [Sta24, Tag 070K], the morphism t_A is the composition of isomorphisms

$$R(G \circ F)(A) = GF(I) \rightarrow RG(F(I)) = RG(RF(A)).$$

\square

Proof of Theorem E.1.0.4. By [Sta24, Tag 09T5], $RQ' : D(D_X) \rightarrow D(\text{Mod}_{\text{qc}}(D_X))$ is right adjoint to $L\iota' = \iota' : D(\text{Mod}_{\text{qc}}(D_X)) \rightarrow D(D_X)$. Let $\Psi' : D_{\text{qc}}(D_X) \rightarrow D(\text{Mod}_{\text{qc}}(D_X))$ (resp. $\Psi : D_{\text{qc}}(O_X) \rightarrow D(\text{Qch}(O_X))$) be the restriction of RQ' (resp. RQ). By Lemma E.1.0.9 2, there are natural commutative squares

$$\begin{array}{ccc} D(\text{Mod}_{\text{qc}}(D_X)) & \xrightarrow{L\iota'} & D_{\text{qc}}(D_X) & \xrightarrow{\Psi'} & D(\text{Mod}_{\text{qc}}(D_X)) \\ \downarrow \text{for} & & \downarrow \text{for} & & \downarrow \text{for} \\ D(\text{Qch}(O_X)) & \xrightarrow{L\iota} & D_{\text{qc}}(O_X) & \xrightarrow{\Psi} & D(\text{Qch}(O_X)), \end{array}$$

where $L\iota$ is induced by the inclusion $\iota : \text{Qch}(O_X) \rightarrow \text{Mod}(O_X)$.

Since Ψ is right adjoint to ι , the counit $\epsilon' : \iota' \Psi' \rightarrow \text{Id}_{D_{\text{qc}}(D_X)}$ (resp. unit $\eta' : \text{Id}_{D(\text{Mod}_{\text{qc}}(D_X))} \rightarrow \Psi' \iota'$) is compatible with the counit $\epsilon : \iota \Psi \rightarrow \text{Id}_{D_{\text{qc}}(O_X)}$ (resp. unit $\eta : \text{Id}_{D(\text{Qch}(O_X))} \rightarrow \Psi \iota$). The functor $\text{for} : D(D_X) \rightarrow D(O_X)$ is conservative. By [Sta24, Tag 09T4], the counit ϵ and the unit η are isomorphisms, so are the counit ϵ' and the unit η' . In particular, the functor (E.1) is an equivalence with a quasi-inverse Ψ' . \square

¹i.e., $F(I)$ computes RG in the sense of [Sta24, Tag 05SX (1)]

E.2 When is an induced D -module holonomic?

Proposition E.2.0.1. *Let X be a complex manifold. Let F be an O_X -module. Then the following conditions are equivalent:*

1. *the induced module $D_X \otimes_{O_X} F$ is holonomic;*
2. *F is coherent with $\text{Supp}(F)$ discrete.*

Lemma E.2.0.2 and Lemma E.2.0.3 are needed for the proof of Proposition E.2.0.1.

Lemma E.2.0.2. *Let A be a Gorenstein local ring (in the sense of [Sta24, Tag 0DW7 (1)]) of Krull dimension n . Let M be a finite A -module. Then the following conditions are equivalent:*

1. *For all integers $i \neq n$, one has $\text{Ext}^i(M, A) = 0$;*
2. *the length of M is finite.*

Proof. Let k be the residue field of A .

- Assume Condition 1. To prove 2, one may assume $M \neq 0$. As A is Gorenstein, $A[0]$ is a dualizing complex of A . By [Mat87, Thm. 18.1, p.141], one has $R\mathcal{H}om_A(k, A[n]) = k[0]$, so $A[n]$ is the normalized dualizing complex of A (in the sense of [Sta24, Tag 0A7M]). Let d be the depth of M . By [Sta24, Tag 0B5A], the module M is Cohen-Macaulay and

$$M = \text{Ext}_A^{n-d}(\text{Ext}_A^{n-d}(M, A), A).$$

Thus, $\text{Ext}_A^{n-d}(M, A) \neq 0$. By Condition 1, one has $n - d = n$. Hence $\dim \text{Supp}(M) = d = 0$. By [Ati69, Exercise 19 v), p.46], one has $\dim A/\text{Ann}(M) = 0$. Then $A/\text{Ann}(M)$ is an artinian ring. From [Eis95, Cor. 2.17], the length of M is finite.

- Assume Condition 2. Induction on the length $l(M)$ of M . When $l(M) = 0$, one has $M = 0$ and Condition 1 holds. Now assume $l(M) > 0$ and the statement holds for all modules of length less than $l(M)$. There is a submodule N of M such that M/N is a simple module and $l(N) < l(M)$. By [Sta24, Tag 00J2], the module M/N is isomorphic to k . For every integer $i \neq n$, the short exact sequence $0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$ induces an exact sequence $\text{Ext}^i(M/N, A) \rightarrow \text{Ext}^i(M, A) \rightarrow \text{Ext}^i(N, A)$. By the inductive hypothesis, $\text{Ext}^i(N, A) = 0$. By [Mat87, Thm. 18.1, p.141], one has $\text{Ext}^i(M/N, A) = 0$. Hence $\text{Ext}^i(M, A) = 0$.

□

Lemma E.2.0.3. *Let X be a complex analytic space. Let F be a coherent O_X -module. Then the length of the $O_{X,x}$ -module F_x is finite for all $x \in X$ if and only if the subspace $\text{Supp}(F) \subset X$ is discrete.*

Proof. The “if” part follows from Lemma 5.5.2.8 1. We prove the “only if” part. By coherence of F and [GR84, p.76], $\text{Supp}(F)$ is a closed analytic set of X . Assume to the contrary that $\text{Supp}(F)$ is not discrete. Then $\dim \text{Supp}(F) > 0$. Let C be an irreducible component of $\text{Supp}(F)$ of maximal dimension. Endow C with the reduced induced closed subspace structure. Let $i : C \rightarrow X$ be the closed embedding of complex analytic spaces.

For every $x \in C$, the morphism $O_{X,x} \rightarrow O_{C,x}$ is surjective. Then by [Sta24, Tag 00IX], one has $l_{O_{C,x}}(i^*F)_x = l_{O_{X,x}}(i^*F)_x$. The morphism $F_x \rightarrow (i^*F)_x$ of $O_{X,x}$ -modules is surjective, so $l_{O_{X,x}}(i^*F)_x \leq l_{O_{X,x}}F_x$. In particular, the length of $(i^*F)_x$ over $O_{C,x}$ is finite. By [GD71, Cor. 5.2.4.1], the support of i^*F is C . Replacing (X, F) by (C, i^*F) , one may assume further that X is irreducible with $\dim X > 0$.

By the generic freeness [Ros68, Prop. 3.1], there is $x_0 \in X$ such that F_{x_0} is a free O_{X,x_0} -module. As the support of F is X , from [RS17, p.238], F is not a torsion sheaf. Then by irreducibility of X and [Ros68, p.69], the O_{X,x_0} -module F_{x_0} has positive rank. Thus, O_{X,x_0} has finite length over itself, hence an artinian ring. The dimension formula in [GR84, p.96] and [CD94, (14.14), p.89] yield $\dim X = \dim_{x_0} X = \dim O_{X,x_0} = 0$, a contradiction. \square

Proof of Proposition E.2.0.1. Let $M = D_X \otimes_{O_X} F$ and $\hat{F} = R\mathcal{H}om_{O_X}(F, O_X)$. By [Sta24, Tag 08DJ], one has

$$\mathcal{H}om_{O_X}(\omega_X, \hat{F}) = R\mathcal{H}om_{O_X}(\omega_X \otimes_{O_X} F, O_X). \quad (\text{E.4})$$

Provided that F is coherent, [Bjö93, (ii) p.122] gives

$$\Delta^{D_X} M = D_X \otimes_{O_X} \mathcal{H}om_{O_X}(\omega_X, \hat{F})[\dim X]. \quad (\text{E.5})$$

Plugging (E.4) into (E.5), one gets

$$\Delta^{D_X} M = D_X \otimes_{O_X} R\mathcal{H}om_{O_X}(\omega_X \otimes_{O_X} F, O_X)[\dim X].$$

For every nonzero integer i , one has

$$H^i(\Delta^{D_X} M) = D_X \otimes_{O_X} \mathcal{E}xt_{O_X}^{i+\dim X}(\omega_X \otimes_{O_X} F, O_X).$$

By [Sta24, Tag 01CB] and [GH78, 1. p.700], its stalk at $x \in X$ is isomorphic to

$$D_{X,x} \otimes_{O_{X,x}} \mathcal{E}xt_{O_{X,x}}^{i+\dim_x X}(F_x, O_{X,x})$$

.

- Assume Condition 2. By [Bjö93, 1.5.1], the D_X -module M is coherent. By Lemma E.2.0.3, the $O_{X,x}$ -module F_x has finite length. As $O_{X,x}$ is a noetherian regular local ring of Krull dimension $\dim_x X$, by Lemma E.2.0.2, one has $\text{Ext}_{O_{X,x}}^{i+\dim_x X}(F_x, O_{X,x}) = 0$ for all $x \in X$. Hence $H^i(\Delta^{D_X} M) = 0$. From Fact 6.7.2.2 2, the D_X -module M is holonomic.
- Assume Condition 1. From [SS94, p.55], the O_X -module F is coherent. From Fact 6.7.2.2 2, for every nonzero integer i , one has $H^i(\Delta^{D_X} M) = 0$. As $D_{X,x}$ is a nonzero free $O_{X,x}$ -module, one gets $\text{Ext}_{O_{X,x}}^{i+\dim_x X}(F_x, O_{X,x}) = 0$. By Lemma E.2.0.2, the $O_{X,x}$ -module F_x has finite length for every $x \in X$. From Lemma E.2.0.3, the support of F is discrete.

□

The proof of Proposition E.2.0.4 (an algebraic analog of Proposition E.2.0.1) is similar.

Proposition E.2.0.4. *Let X be a smooth algebraic variety over an algebraically closed field of characteristic 0. Let F be an O_X -module. Then the following conditions are equivalent:*

1. *the induced module $D_X \otimes_{O_X} F$ is holonomic;*
2. *F is coherent with $\text{Supp}(F)$ finite.*

Appendix F

Group extensions of complex Lie groups

F.1 Introduction

In the history of cohomology theory of abelian varieties over positive characteristic fields, the study of group extension problem played an important role. For instance, Rosenlicht obtains Fact F.1.0.1 through considering vectorial extensions of abelian varieties. Let k be an algebraically closed field and A/k be an abelian variety with $\dim A = g$. The dual abelian variety of A is denoted by A^\vee .

Fact F.1.0.1 ([Ros58, Theorems 1 and 2]). *The dimension of the k -vector space $H^1(A, O_A)$ is g .*

A notable byproduct of Rosenlicht's work is the existence of the following object, the so-called *universal vectorial extension*.

Fact F.1.0.2 ([Ros58, Prop. 11]). *There is a short exact sequence¹ of commutative algebraic groups over k : $0 \rightarrow \mathbb{G}_a^g \rightarrow A^\natural \rightarrow A \rightarrow 0$, where A^\natural is the moduli space of line bundles equipped with an integrable connection on A^\vee .*

In [Rot96, (1.17)] and [Lau96, Thm. 3.2.1], it is proved that the Fourier-Mukai transform $D^b(\mathrm{Qch}(O_A)) \rightarrow D^b(\mathrm{Qch}(O_{A^\vee}))$ lifts to an equivalence $D^b(\mathrm{Qch}(O_{A^\natural})) \rightarrow D^b(\mathrm{Qch}(D_{A^\vee}))$, where for a smooth algebraic variety M/k , $\mathrm{Qch}(O_M)$ (resp. $\mathrm{Qch}(D_M)$) refers to the category of O_M (resp. left D_M) modules that are O_M -quasi-coherent.

The cohomology theory of complex analytic analogue of abelian varieties, namely complex tori, is elementary. By contrast, as far as we know, the existence of universal vectorial extension in the analytic setting is not

¹in the sense of [Ros58, Sec. 2, p.691]

covered in the literature, though admittedly easier and should be known. The main results are summarized as follows.

Proposition (Proposition F.4.3.1). *For two commutative complex Lie groups A, B , the commutative extensions of A by B are classified by the abelian group*

$$\mathrm{Ext}_{\mathbb{Z}}^1(\pi_0(A), \pi_0(B)) \oplus \mathrm{Hom}_{\mathrm{Ab}}(\pi_1(A_0), \pi_0(B)) \oplus \mathrm{coker}(s).$$

Here s is the restriction morphism $\mathrm{Hom}_{\mathrm{Vec}}(L(A), L(B_0)) \rightarrow \mathrm{Hom}_{\mathrm{Ab}}(\pi_1(A_0), B_0)$, A_0 (resp. B_0) signifies the identity component of A (resp. B), the notation $\pi_1(*)$ refers to the fundamental group, and $A/A_0 = \pi_0(A)$ denotes the 0-th homotopy group of A and similar for B .

Theorem. *Let A be a complex torus of dimension g . Then:*

- (Theorem F.5.2.4 (resp. F.5.3.2)) *The dual torus $\mathrm{Pic}^0(A)$ (resp. tangent space $T_0A = H^1(A, O_A)$) naturally classifies the extensions of A by the multiplicative group \mathbb{C}^* (resp. additive group \mathbb{C}).*
- (Propositions F.5.4.5 1 and F.5.4.7) *There is an extension*

$$0 \rightarrow H^0(A^\vee, \Omega_{A^\vee}^1) \rightarrow (\mathbb{C}^*)^{2g} \rightarrow A \rightarrow 0$$

that is universal among all vectorial extensions of A .

We emphasize some differences between the analytic case and the algebraic case. For a complex torus A , let $\mathrm{Div}(A)$ be the group of analytic divisors on A modulo linear equivalence. Let $\mathrm{Pic}(A)$ be the group of isomorphic classes of line bundles on A . The natural map $\mathrm{Div}(A) \rightarrow \mathrm{Pic}(A)$ is surjective if and only if A is an abelian variety ([Deb05, Sec. 4.3, Cor. 4]). This is why the Picard group is used in Theorem F.5.2.4 while divisor group appears in its algebraic analogue ([Wei49, no. 2], [Ser88, Thm. 6]). Discrete groups like \mathbb{Z} are not (finite type) algebraic groups, but there is no reason to exclude them as complex Lie groups. Plenty of important analytic morphisms are not algebraic, like the universal covering (exponential map) $\exp : \mathbb{C} \rightarrow \mathbb{C}^*$.

The organization is as follows. The main goal of this text is to classify extensions of complex Lie groups. Section F.2 contains preliminaries about complex Lie groups. In Section F.3 we define complex Lie group extensions and give several first results about the classification. Then we focus on commutative extensions in Section F.4. Commutative extensions of complex tori deserve extra attention, and they are discussed in Section F.5. Some extensions with complex-tori base are automatically commutative, as Section F.6 shows. Noncommutative extensions are treated superficially in Section F.7.

Convention and notation

A statement about Lie groups is understood to hold for both real and complex Lie groups. The topology underlying a Lie group is always assumed to be second countable.²

For every Lie group G , the identity component of G is denoted by G_0 . The Lie algebra of G is written as $L(G)$. And $Z(G)$ denotes the center of G . The automorphism group of G is denoted by $\text{Aut}(G)$. Let $\text{Inn} : G \rightarrow \text{Aut}(G)$ be the group morphism defined by taking conjugation $g \mapsto g \bullet g^{-1}$. Then the subgroup $\text{Inn}(G)$ of inner automorphisms is normal in $\text{Aut}(G)$. Let $\text{Out}(G) = \text{Aut}(G)/\text{Inn}(G)$ be the group of outer automorphisms. Let G^{op} be the Lie group opposite to G . (If G is complex, then so is G^{op} .) There is a natural identification of real/complex manifolds $G \rightarrow G^{\text{op}}$ denoted by $g \mapsto g^*$. If G is connected, then the universal covering group of G is denoted by \tilde{G} and the fundamental group of G with the identity e_G as the base point is denoted by $\pi_1(G)$.

Complex Lie subgroups refer to embedded closed complex Lie subgroups. If G is a complex Lie group and $S \subset G$ is a subset, by [HN11, Exercise 15.1.3 (b)] there is a smallest complex Lie subgroup of G containing S , called the complex Lie subgroup generated by S .

Let Vec (resp. Ab , resp. \mathcal{C} , resp. Set) be the category of finite dimensional complex vector spaces (resp. abelian groups, resp. commutative complex Lie groups, resp. sets). For a complex manifold X and a commutative complex Lie group B , let \mathcal{B}_X be the abelian sheaf on X of germs of holomorphic maps from X to B .

F.2 Generalities on complex Lie groups

Two fundamental facts about complex Lie groups are recalled.

Fact F.2.0.1 ([Bou72, Ch. III, §3, no. 8, Prop. 28]). *Let $f : G \rightarrow H$ be a morphism of complex Lie groups. Then:*

1. *$\ker(f)$ is a normal complex Lie subgroup of G and $L(\ker(f)) = \ker(d_e f : L(G) \rightarrow L(H))$.*
2. *If $f(G)$ is closed in H , then $f(G)$ is a complex Lie subgroup of H , and f induces a complex Lie group isomorphism $G/\ker(f) \rightarrow f(G)$. In particular, if f is surjective, then $d_e f : L(G) \rightarrow L(H)$ is surjective. If f is bijective, then f is an isomorphism.*

Remark F.2.0.2. Fact F.2.0.1 2 fails if the topology of G is not assumed to be second countable. For example, let τ (resp. τ') be the discrete topology

²A partial reason for such restriction is that, in this case, Condition (2) of [Hoc51b, Definition 1.1] is implied by Condition (1), showed in p.542 *loc.cit.*

(resp. the Euclidean topology) of \mathbb{C} , then $\text{Id} : (\mathbb{C}, \tau) \rightarrow (\mathbb{C}, \tau')$ is a bijective morphism but not open.

Right principal bundle is defined in [Bou07, 6.2.1]. Left principal bundle can be defined similarly.

Fact F.2.0.3 ([HBS66, Thm. 3.4.3], [Bou72, Ch. III, §1, Propositions 10 and 11]). *Suppose G is a complex Lie group and K is a normal complex Lie subgroup of G . Then the group G/K has a unique structure of complex manifold, such that the quotient map $\pi : G \rightarrow G/K$ is a submersion.³ With this structure, G/K is a complex Lie group and p is a left principal K -bundle under the natural left group action $K \times G \rightarrow G$ defined by $(k, g) \mapsto kg$. In particular, every surjective morphism of complex Lie groups is open.*

We recall that principal bundles are classified by the first sheaf cohomology, in the following way. Let X (resp. B) be a complex manifold (resp. commutative complex Lie group). Let S be the set of isomorphism classes of principal B -bundles⁴ over X . Define a map

$$\Psi : S \rightarrow H^1(X, \mathcal{B}_X) \quad (\text{F.1})$$

as follows. For every $[p : P \rightarrow X] \in S$, there exists an open cover $\{U_i\}_{i \in I}$ of X and a family of local trivializations $f_i : U_i \times B \rightarrow p^{-1}(U_i)$ for every $i \in I$. For any indices $i, j \in I$ and every $x \in U_i \cap U_j$, there exists a unique element $b_{ij}(x) \in B$ such that $b_{ij}(x) \cdot f_i(y) = f_j(y)$ for all $y \in p^{-1}(x)$. Hence a morphism $b_{ij} : U_i \cap U_j \rightarrow B$ of complex manifolds. Moreover, for any indices $i, j, k \in I$ and every $x \in U_i \cap U_j \cap U_k$, they satisfy the 1-cocycle relation $b_{ij}(x) + b_{jk}(x) + b_{ki}(x) = 0$. Thus, the family $\{b_{ij}\}_{i,j \in I}$ defines an element $\Psi(p)$ of $H^1(X, \mathcal{B}_X)$.

As per [HBS66, 3.2 b), p.41], the map Ψ is bijective. The structure of abelian group on $H^1(X, \mathcal{B}_X)$ is translated to S via Ψ . The zero element of S is the class of the trivial principal B -bundle. For every pair $[p_1 : P_1 \rightarrow X]$ and $[p_2 : P_2 \rightarrow X]$ in S , by taking a family of trivialization for each p_i , we can define a morphism $\phi : P_1 \times_X P_2 \rightarrow P_1 + P_2$ of principal B -bundles on X such that for every $b, b' \in B$, $u \in P_1$, $v \in P_2$ with $p_1(u) = p_2(v)$, one has

$$\phi(b \cdot u, b' \cdot v) = (b + b') \cdot \phi(u, v). \quad (\text{F.2})$$

In particular, ϕ is surjective. Restricted to the fiber at some $x \in X$, ϕ is induced by the group law of B and the chosen trivializations.

We need a complex version of Cartan's subgroup theorem. Notice that a real analytic closed subgroup of a complex Lie group may not be a complex analytic subset.

³in the sense of [Bou07, 5.9.1]

⁴Here B is commutative, so it is unnecessary to specify the principal bundle to be left or right.

Lemma F.2.0.4 ([Bjö93, p.513]). *Let X be a complex manifold, $Y \subset X$ be a complex analytic subset. If $p \in Y$ is a smooth point of Y , then near p , the subset Y is an embedded complex submanifold of X .*

Proof. As the problem is local, we may assume that X is an open subset \mathbb{C}^n , there exist $f_1, \dots, f_m \in O_X(X)$ with $O_{X,p}/(f_1, \dots, f_m) = O_{Y,p}$ and $Y = Z(f_1, \dots, f_m)$. Let $r = \text{rank}_p(f_1, \dots, f_m)$. By reordering subscripts, one may assume

$$\det\left(\frac{\partial f_i}{\partial z_j}\right)_{1 \leq i, j \leq r} \neq 0.$$

Then $(f_1, \dots, f_r) : X \rightarrow \mathbb{C}^r$ is a holomorphic submersion near p . Therefore, near p , the subset $Z(f_1, \dots, f_r)$ is an embedded complex submanifold of X of dimension $n - r$. By the Jacobian criterion (see, e.g., [GR84, p.114]), one has $\text{emb}_p Y = n - r$. By the criterion of smoothness ([GR84, p.116]), one has $\dim_p Y = n - r$. Now that $Y \subset Z(f_1, \dots, f_r)$, near p the subset Y is an irreducible component of $Z(f_1, \dots, f_r)$, hence also an embedded complex submanifold of X . \square

Corollary F.2.0.5 contains [Lee01, Prop. 1.23] as a special case.

Corollary F.2.0.5 (Complex Cartan subgroup theorem). *Let G be a complex Lie group, and let H be a subgroup that is a complex analytic subset of G . Then H is a complex Lie subgroup of G .*

Proof. Endow H with the induced structure of reduced complex analytic space. By [GR84, p.117], the complex analytic space H has a smooth point p . For every $q \in H$, the left multiplication by qp^{-1} gives a biholomorphic map $G \rightarrow G$, which sends H to H and maps p to q . Therefore, q is also a smooth point of H . By Lemma F.2.0.4, H is a complex submanifold of G near q for all $q \in H$. Thus, H is a complex submanifold of G and hence a complex Lie subgroup. \square

In Lemma F.2.0.6, if G is furthermore connected, then the result of is contained in [Bou72, Ch.III, Sec. 6, no. 4, Cor. 4].

Lemma F.2.0.6. *Let G be a complex Lie group. Then the center $Z(G)$ is a complex Lie subgroup of G .*

Proof. The holomorphic map $G \times G \rightarrow G$ defined by $(x, y) \mapsto yxy^{-1}$ is a group action of G on itself. By [Bou72, Ch. III, Sec. 1, no. 7, Prop. 14], for every $x \in G$, the stabilizer $C_G(x)$ of $x \in G$ is a complex Lie subgroup of G . Therefore, so is $Z(G) = \cap_{x \in G} C_G(x)$ by [HN11, Exercise 15.1.3 (a)]. \square

A complex Lie group isomorphic to a complex Lie subgroup of $\text{GL}_n(\mathbb{C})$ for some integer $n \geq 1$ is called *linear*. Proposition F.2.0.7, due to Matsushima and Morimoto, is a characterization of commutative linear complex Lie groups.

Proposition F.2.0.7. *Let B be a connected commutative complex Lie group. Then the following conditions are equivalent:*

1. B is isomorphic to $\mathbb{C}^m \times (\mathbb{C}^*)^n$ for some integers $m, n \geq 0$;
2. the complex Lie group B is linear;
3. B is a Stein group (i.e., the underlying complex manifold is a Stein manifold).

In that case, the pair (m, n) is unique.

Proof. See [HN11, Exercise 15.3.1] for the fact that 1 implies 2. Since $\mathrm{GL}_n(\mathbb{C})$ is a Stein manifold, 2 implies 3. As per [MM60, Proposition 4], 3 implies 1. The uniqueness is contained in the Remmert-Morimoto decomposition (see, e.g., [AK01, Thm. 1.1.5]). \square

Remark F.2.0.8. The commutativity of B in Proposition F.2.0.7 is important. In fact, there is a connected Stein group that is not linear ([Ari19, Sec. 1]). This differs from the algebraic case where every algebraic group that is an affine variety is linear ([Mil17a, Cor. 4.10]).

In some sense, Definition F.2.0.9 is an antipode to Stein groups.

Definition F.2.0.9. A connected complex Lie group on which every holomorphic function is constant is called a toroidal group.⁵

Complex tori are toroidal groups, but there exist toroidal groups that are not compact ([AK01, p.1]). Every toroidal group is a semi-torus in the sense of [NW13, Def. 5.1.5].

By [AK01, 1.1.5], every connected commutative complex Lie group G is uniquely isomorphic to $\mathbb{C}^l \times (\mathbb{C}^*)^m \times X$ with a toroidal group X . In particular, G can be presented as an extension of a complex torus by a connected linear group. (From [NW13, pp.169-170], a semi-torus can admit nonequivalent presentations, while semiabelian varieties admit exactly one algebraic presentation.)

F.3 Group extensions

Given a surjective Lie group morphism $p : E \rightarrow Q$, by Fact F.2.0.1, $K := \ker(p)$ is a normal Lie subgroup of E and the induced morphism $E/K \rightarrow Q$ is an isomorphism. We write it as

$$1 \rightarrow K \xrightarrow{i} E \xrightarrow{p} Q \rightarrow 1 \tag{F.3}$$

⁵also known as a Cousin group

and call it a short exact sequence. In that case, E is called an *extension* of the base Q by the extension kernel K . Moreover, $d_{ep} : L(E) \rightarrow L(Q)$ is surjective of kernel $L(K)$, hence an extension of Lie algebras

$$0 \rightarrow L(K) \rightarrow L(E) \xrightarrow{d_{ep}} L(Q) \rightarrow 0.$$

When $K \subset Z(E)$, such an extension is called *central*. If (F.3) is a central extension with Q commutative, as in [MRM74, p.222], using Fact F.2.0.3 one can construct a skew-symmetric bimorphism

$$e : Q \times Q \rightarrow K, \quad (\text{F.4})$$

to measure the deviation of E from commutativity. Indeed, the group E is commutative if and only if e is constant.

Several topological properties of Lie groups are preserved by extensions.

Fact F.3.0.1. *If K, Q in (F.3) are compact (resp. connected, resp. discrete), then so is E .*

Proof. The statement concerning connectedness is in [Che46, Prop. 2, p.36]. The others are consequences of Fact F.2.0.3. \square

Fact F.3.0.2 ([HN11, Cor. 16.3.9]). *If (F.3) is a central extension of complex Lie groups, where K is finite and E is connected, then Q is linear if and only if E is linear.*

The finiteness of K in Fact F.3.0.2 is necessary. Consider the exact sequence $0 \rightarrow \mathbb{Z}^2 \rightarrow \mathbb{C} \rightarrow A \rightarrow 0$ defining a complex torus A . Here \mathbb{Z}^2 and \mathbb{C} are linear, while A is not.

Similarly, an extension E of a finite group Q by a linear group K is linear. Indeed, let $\rho : K \rightarrow \text{GL}_n(\mathbb{C})$ be a faithful representation, then the induced representation $\text{Ind}_K^E \rho : E \rightarrow \text{GL}_{mn}(\mathbb{C})$ is also faithful, where $m = \#Q$. Again, the finiteness of Q is essential here. Example F.3.0.3 shows the statement fails when Q is only discrete and linear but infinite.

Example F.3.0.3. Work of Deligne [Del78] (see also [KRW20, p.470]) shows that for any integers $g \geq 2, n \geq 3$, there is a central extension $1 \rightarrow \mathbb{Z}/n \rightarrow G \rightarrow \text{Sp}_{2g}(\mathbb{Z}) \rightarrow 1$ for which G is not residually finite. By Malcev's theorem ([Mal40, Thm. VII]; see also [Nic13, p.1]), the discrete complex Lie group G is not linear, even though \mathbb{Z}/n and $\text{Sp}_{2g}(\mathbb{Z})$ are linear.

We turn to the classification of extensions. Two Lie group extensions C and C' of B by A are called equivalent if there exists a morphism $f : C \rightarrow C'$ making a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \longrightarrow & C & \longrightarrow & A \longrightarrow 0 \\ & & \downarrow \text{Id} & & \downarrow f & & \downarrow \text{Id} \\ 0 & \longrightarrow & B & \longrightarrow & C' & \longrightarrow & A \longrightarrow 0. \end{array}$$

In this case, f is bijective, hence an isomorphism by Fact F.2.0.1. The *trivial* extension of Q by K refers to the equivalence class of the obvious sequence

$$1 \rightarrow K \rightarrow K \times Q \rightarrow Q \rightarrow 1.$$

Fact F.3.0.4 ([Bou72, Ch.III, no.4, Prop. 8]). *The Lie group extension (F.3) is trivial if and only if there is a morphism $r : E \rightarrow K$ with $ri = \text{Id}_K$. The extension is a semidirect product if and only if there is a morphism $s : Q \rightarrow E$ with $ps = \text{Id}_Q$.*

The extension (F.3) defines a group morphism $\psi : Q \rightarrow \text{Out}(K)$, called the *outer action* corresponding to the extension. We call (K, ψ) the extension kernel of (F.3). Equivalent extensions induce the same outer action. For two complex Lie groups Q, K and a group morphism $\psi : Q \rightarrow \text{Out}(K)$, denote by $\text{Ext}(Q, K, \psi)$ the set of equivalence classes of extensions of Q by K with outer action ψ .

Since the center $Z(K)$ is a characteristic complex Lie subgroup of K by Lemma F.2.0.6, there is a canonical group morphism $\text{Aut}(K) \rightarrow \text{Aut}(Z(K))$ which passes to another group morphism $\text{Out}(K) \rightarrow \text{Aut}(Z(K))$. Hence a group morphism

$$\psi_0 : Q \rightarrow \text{Aut}(Z(K)) \tag{F.5}$$

induced by ψ . When K is *commutative*, $\psi = \psi_0$ and the construction of Baer sum ((F.41) and [FLA19, p.444]) makes $\text{Ext}(Q, K, \psi)$ an abelian group.

F.3.1 Pullback and pushout

Extensions can be pulled back.

Example F.3.1.1 (Pullback). Given a morphism $g : Q' \rightarrow Q$ of complex Lie groups, pulling (F.3) back along g gives an extension of Q' by K as follows.

The map $E \times Q' \rightarrow Q$ defined by $(x, h') \mapsto p(x)^{-1}g(h')$ is holomorphic, so the preimage E' of the identity element $e_Q \in Q$ is an analytic subset of $E \times Q'$. As $E' = \{(x, h') \in E \times Q' : p(x) = g(h')\}$ is a subgroup of $E \times Q'$, by Corollary F.2.0.5, E' is a complex Lie subgroup of $E \times Q'$ (which is the extension group). Let $p' : E' \rightarrow Q'$ and $\epsilon : E' \rightarrow E$ be the projections. Then the triple (E', ϵ, p') is the fiber product $E \times_Q Q'$ in the category of complex Lie groups.

For every $h' \in Q'$, by surjectivity of p , there is $x \in E$ with $p(x) = g(h')$. Then $(x, h') \in E'$ with $p'(x, h') = h'$. Hence p' is surjective.

Define a morphism $i' : K \rightarrow E'$ by $i'(k) = (k, e_{Q'})$. Then i' is injective. Since $p'i'$ is trivial, $i'(K) \subset \ker(p')$. Conversely, for every $(x, h') \in \ker(p')$, $h' = e_{Q'}$ and $p(x) = g(e_{Q'}) = e_Q$. Thus, $x \in K$ and $(x, h') = i'(x) \in i'(K)$. Hence a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 1 & \longrightarrow & K & \xrightarrow{i'} & E' & \xrightarrow{p'} & Q' & \longrightarrow & 1 \\ & & \downarrow \text{Id} & & \downarrow \epsilon & & \downarrow g & & \\ 1 & \longrightarrow & K & \xrightarrow{i} & E & \xrightarrow{p} & Q & \longrightarrow & 1 \end{array}$$

The first row is called the pullback extension of (F.3) along g . Its outer action is $\psi g : Q' \rightarrow \text{Out}(K)$. Hence a map $\text{Ext}(Q, K, \psi) \rightarrow \text{Ext}(Q', K, \psi g)$. It is a group morphism when K is commutative ([Hoc51a, p.99]).

The universal property of pullback shows that the first row of every such commutative diagram is determined by the second row and $g : Q' \rightarrow Q$. By construction, the pullback of a central extension is also central.

A pushout extension along a morphism $f : K \rightarrow K'$ of complex Lie groups may not exist. When it exists, it satisfies a universal property.

Lemma F.3.1.2. *Consider a commutative diagram of complex Lie groups, where each row is exact*

$$\begin{array}{ccccccc}
1 & \longrightarrow & K & \longrightarrow & E & \xrightarrow{p} & Q \longrightarrow \cdot \\
& & \downarrow f & & \downarrow m & & \downarrow \text{Id} \\
1 & \longrightarrow & K' & \xrightarrow{\iota} & E' & \xrightarrow{\pi} & Q \longrightarrow 1.
\end{array} \tag{F.6}$$

Then the triple (E', m, ι) has the following universal property: For every commutative diagram of complex Lie groups

$$\begin{array}{ccc}
K & \xrightarrow{i} & E \\
\downarrow f & & \downarrow m \\
K' & \xrightarrow{\iota} & E'
\end{array}
\begin{array}{c}
\searrow \phi \\
\searrow \eta \\
\searrow \psi
\end{array}
\begin{array}{c}
\\
\\
H
\end{array}$$

with $\psi(m(c)^{-1}bm(c)) = \phi(c)^{-1}\psi(b)\phi(c)$ for every $c \in E$ and $b \in K'$, there exists a unique morphism $\eta : E' \rightarrow H$ keeping the diagram commutative.

In particular, up to a unique equivalence, the second row of (F.6) has at most one choice when the first row and $f : K \rightarrow K'$ are given.

Proof. We construct a map $\eta : E' \rightarrow H$ as follows. For every $c' \in E'$, there exists $c \in E$ with $p(c) = \pi(c')$. Let $b' = m(c)^{-1}c'$. Then $\pi(b') = p(c)^{-1}\pi(c') = e_Q$, so $b' \in K'$. Define $\eta(c') = \phi(c)\psi(b')$.

To show that η is well-defined, we claim that $\eta(c')$ is independent of the choice of c . Indeed, take another $c_1 \in E$ with $p(c_1) = \pi(c')$, then $p(c^{-1}c_1) = e_Q$, hence $c^{-1}c_1 \in K$. This time the element in K' is $b'_1 = m(c_1)^{-1}c'$, so $b' = f(c^{-1}c_1)b'_1$ in K' and hence $\psi(b') = \phi(c^{-1}c_1)\psi(b'_1)$. Therefore, $\phi(c)\psi(b') = \phi(c_1)\psi(b'_1)$ in H as claimed.

We check that η is holomorphic near $c' \in E'$. Indeed, by Fact F.2.0.3, there is an open neighborhood U of $\pi(c') \in Q$, and a holomorphic map $s : U \rightarrow E$ with $ps = \text{Id}_U$. The map

$$\pi^{-1}(U) \rightarrow U \times K', \quad x \mapsto (\pi(x), [ms\pi(x)]^{-1}x)$$

is biholomorphic. The map

$$U \times K' \rightarrow H, \quad (u, b') \mapsto \phi(s(u))\psi(b')$$

is holomorphic. The composition is exactly $\eta|_{\pi^{-1}(U)}$.

We check that η is a group morphism. For $c'_i \in E'$ ($i = 1, 2$), choose $c_i \in E$ with $p(c_i) = \pi(c'_i)$. Then for $c'_1 c'_2$ we can choose $c_1 c_2$. Let $b'_1 = m(c_1)^{-1}c'_1$ and $b'_2 = m(c_2)^{-1}c'_2$. Then

$$b' := m(c_1 c_2)^{-1}c'_1 c'_2 = m(c_2)^{-1}b'_1 m(c_2)b'_2.$$

By the construction of η , one has

$$\begin{aligned} \eta(c'_1 c'_2) &= \phi(c_1 c_2)\psi(b') \\ &= \phi(c_1)\phi(c_2)\psi[m(c_2)^{-1}b'_1 m(c_2)]\psi(b'_2) \\ &= \phi(c_1)\psi(b'_1)\phi(c_2)\psi(b'_2) = \eta(c'_1)\eta(c'_2) \end{aligned}$$

Then η is a morphism of complex Lie groups. By construction, η is the unique group morphism keeping the diagram commutative. \square

Example F.3.1.3. Assume that Q is connected. As the map $p : E \rightarrow Q$ in (F.3) is open by Fact F.2.0.3, $p(E_0)$ is a nonempty open subgroup of Q and hence $p(E_0) = Q$ by the connectedness of Q . Then the following diagram is commutative and each row is exact

$$\begin{array}{ccccccc} 1 & \longrightarrow & K \cap E_0 & \longrightarrow & E_0 & \longrightarrow & Q \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \text{Id} \\ 1 & \longrightarrow & K & \longrightarrow & E & \longrightarrow & Q \longrightarrow 1 \end{array}$$

By Lemma F.3.1.2, the second row is determined by the inclusion $K \cap E_0 \rightarrow K$ (an open normal subgroup) and the first row.

F.3.2 Rudimentary classification

Let K, Q be complex Lie groups, where Q is discrete. Consider an abstract group extension $1 \rightarrow K \rightarrow E \rightarrow Q \rightarrow 1$. Then as a set $E = \sqcup_x xK$, where x runs through a set of left representatives of E/K . Thus E admits a unique complex manifold structure making the maps holomorphic. However, the group law of E needs not to be holomorphic in this complex structure. The

semidirect product sequence $1 \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rtimes \mathbb{Z}/2 \rightarrow \mathbb{Z}/2 \rightarrow 1$ serves as an example, where $\mathbb{Z}/2$ acts on \mathbb{C} by complex conjugation. But when the base is discrete and the outer action is trivial, the Lie group extension problem reduces to the abstract group extension problem.

Proposition F.3.2.1. *Let K, Q be complex Lie groups. If Q is discrete, then the natural forgetful map $\phi : \text{Ext}(Q, K, 1) \rightarrow \text{Ext}_{\text{Abs}}(Q, K, 1)$ is bijective, where $\text{Ext}_{\text{Abs}}(Q, K, 1)$ denotes the set of isomorphism classes of abstract group extensions of Q by K with trivial outer action. In fact, for every abstract group extension $1 \rightarrow K \rightarrow E \rightarrow Q \rightarrow 1$, E admits a unique complex manifold structure making the sequence an extension of complex Lie groups.*

Proof. We prove that ϕ is injective. Consider $E_1, E_2 \in \text{Ext}(Q, K, 1)$ with $\phi(E_1) = \phi(E_2)$. Then there is an abstract group isomorphism $f : E_1 \rightarrow E_2$ making a commutative diagram

$$\begin{array}{ccc} K & \hookrightarrow & E_1 \\ & \searrow & \downarrow f \\ & & E_2. \end{array}$$

For every $x \in E_1$, the restriction $xK \rightarrow f(x)K$ of f is holomorphic, since the left multiplication $K \rightarrow xK$ (resp. $K \rightarrow f(x)K$) by x (resp. $f(x)$) in E_1 (resp. E_2) is biholomorphic. Thus, f is holomorphic and hence an equivalence of complex Lie group extensions.

We prove that ϕ is surjective. Given an abstract group extension $1 \rightarrow K \rightarrow E \rightarrow Q \rightarrow 1$ in $\text{Ext}_{\text{Abs}}(Q, K, 1)$, we endow E with the complex structure making the maps holomorphic. We show the group law $m : E \times E \rightarrow E$ is holomorphic. Choose a set-theoretic section $s : Q \rightarrow E$. Then the map $K \times Q \rightarrow E$ defined by $(a, b) \mapsto as(b)$ is biholomorphic. With this identification, m becomes the map

$$\mu : K \times Q \times K \times Q \rightarrow K \times Q, \quad (a, b, a', b') \mapsto (as(b)a's(b')s(bb')^{-1}, bb') = (a\rho(a')s(b)s(b'))s(bb')^{-1}, b$$

where $\rho : K \rightarrow K$ is $x \mapsto s(b)xs(b)^{-1}$. Since the outer action is trivial, $\rho \in \text{Inn}(K)$. Therefore, the map $K \times K \rightarrow K$ defined by $(a, a') \mapsto a\rho(a')$ is holomorphic. Because Q is discrete, μ (and hence m) is holomorphic. Then E is a complex Lie group and the abstract extension lifts to $\text{Ext}(Q, K, 1)$. \square

Corollary F.7.2.6 below is a result about discrete base with nontrivial outer action. We turn to two other simple cases.

Proposition F.3.2.2. *Every extension of \mathbb{C} is a semidirect product. In particular, every central extension of \mathbb{C} is trivial.*

Proof. Let $0 \rightarrow B \rightarrow C \xrightarrow{p} \mathbb{C} \rightarrow 0$ be an extension. Then $0 \rightarrow L(B) \rightarrow L(C) \xrightarrow{dep} L(\mathbb{C}) \rightarrow 0$ is an exact sequence of Lie algebras. Take a \mathbb{C} -linear map

$ds : L(\mathbb{C}) \rightarrow L(C)$ with $d_e p \circ ds = \text{Id}_{L(\mathbb{C})}$. Because $\dim_{\mathbb{C}} L(\mathbb{C}) = 1$, ds is a Lie algebra morphism. As \mathbb{C} is simply connected, there is a unique morphism $s : \mathbb{C} \rightarrow C$ with $d_e s = ds$. Since $d_e(ps) = \text{Id}_{L(\mathbb{C})}$, one has $ps = \text{Id}_{\mathbb{C}}$. Therefore, this extension is a semidirect product by Fact F.3.0.4. \square

Proposition F.3.2.3. *Let B be a connected commutative complex Lie group. Then every central extension of \mathbb{C}^* by B is trivial.*

Proof. Let C be a central extension of \mathbb{C}^* by B . Consider the pullback extension along $\exp(2\pi i \bullet) : \mathbb{C} \rightarrow \mathbb{C}^*$. By Proposition F.3.2.2, there is a morphism $\rho : \mathbb{C} \rightarrow C'$ with $p'\rho = \text{Id}_{\mathbb{C}}$. Then $p\epsilon\rho(1) = \exp(2\pi i) = 1$, so $\epsilon\rho(1) \in B$. As B is connected commutative, its exponential map $\exp_B : L(B) \rightarrow B$ is surjective. Take $v \in L(B)$ with $\exp_B(-v) = \epsilon\rho(1)$.

$$\begin{array}{ccccccc}
 & & & & \mathbb{Z} & & \\
 & & & & \downarrow & & \\
 1 & \longrightarrow & B & \longrightarrow & C' & \xleftarrow{\rho} & \mathbb{C} \longrightarrow 0 \\
 & & \downarrow \text{Id} & & \downarrow \epsilon & \nearrow p' & \downarrow \exp(2\pi i \bullet) \\
 1 & \longrightarrow & B & \longrightarrow & C & \xrightarrow{p} & \mathbb{C}^* \longrightarrow 1
 \end{array}$$

Define a holomorphic map

$$\rho' : \mathbb{C} \rightarrow C', \quad \rho'(z) = \exp_B(zv)\rho(z).$$

We check that ρ' is a group morphism. For every $z, w \in \mathbb{C}$,

$$\begin{aligned}
 \rho'(z+w) &= \exp_B((z+w)v)\rho(z+w) = \exp_B(zv)\exp_B(wv)\rho(z)\rho(w) \\
 &= \exp_B(zv)\rho(z)\exp_B(wv)\rho(w) = \rho'(z)\rho'(w),
 \end{aligned}$$

where the last but one equality uses $B \subset Z(C)$.

Therefore, ρ' is a complex Lie group morphism. Moreover, $\rho'(1) = \exp_B(v)\rho(1) = \epsilon\rho(-1)\rho(1)$. Then $\epsilon\rho'(1) = e_C$. Therefore, $\rho'(\mathbb{Z}) \subset \ker(\epsilon)$. Thus, ρ' induces a morphism $s : \mathbb{C}^* \rightarrow C$ making a commutative diagram

$$\begin{array}{ccccc}
 \mathbb{C} & \xrightarrow{\rho'} & C' & \xrightarrow{p'} & C \\
 \downarrow \exp(2\pi i \bullet) & & \downarrow \epsilon & & \downarrow \exp(2\pi i \bullet) \\
 \mathbb{C}^* & \xrightarrow{s} & C & \xrightarrow{p} & \mathbb{C}^*
 \end{array}$$

Since $p'\rho' = \text{Id}_{\mathbb{C}}$ and $\exp(2\pi i \bullet) : \mathbb{C} \rightarrow \mathbb{C}^*$ is surjective, $ps = \text{Id}_{\mathbb{C}^*}$. From Fact F.3.0.4, the extension C is trivial. \square

Example F.7.1.7 gives a result about non-central extensions of \mathbb{C}^* .

Now assume that the Lie group K is discrete and *commutative*. We recall results⁶ from [Hoc51b, Sec. 3].

⁶They are stated for real Lie groups, but the proofs extend to the complex setting.

Fact F.3.2.4 ([Hoc51b, p.545]). *Let K, Q be Lie groups. If K is discrete commutative and Q is connected, then the extension (F.3) of Lie groups is central.*

Corollary F.3.2.5. *Let K, Q be commutative Lie groups. If Q is connected and K is discrete, then every extension of Q by K is commutative.*

Proof. Let (F.3) be such an extension. By Fact F.3.2.4, this extension is central. Then consider the induced continuous map (F.4). Since Q is connected and K is discrete, this map is constant, or equivalently, E is commutative. \square

Let Ab_c be the abelian category of abelian groups that are at most countable. Fact F.3.2.6 shows that the universal cover of a connected Lie group is “universal” among all the extensions with discrete commutative kernels.

Fact F.3.2.6 (Hochschild, [Hoc51b, Thm. 3.2 and Cor.]). *Let Q be a connected Lie group. Then the functor $\text{Ext}(Q, \cdot, 1) : \text{Ab}_c \rightarrow \text{Ab}$ is represented by $\pi_1(Q)$ and the class of the universal cover sequence $1 \rightarrow \pi_1(Q) \rightarrow \tilde{Q} \rightarrow Q \rightarrow 1$ in $\text{Ext}(Q, \pi_1(Q), 1)$. Hence an isomorphism $\Gamma_K : \text{Ext}(Q, K, 1) \rightarrow \text{Hom}_{\text{Ab}}(\pi_1(Q), K)$ functorial in $K \in \text{Ab}_c$. Moreover, $E \in \text{Ext}(Q, K, 1)$ is connected if and only if $\Gamma_K(E)$ is surjective.*

F.4 Commutative Extensions

F.4.1 Generalities

Lemma F.4.1.1. *The category \mathcal{C} is naturally additive with finite direct products.*

Proof. The Hom sets are commutative groups, and composition of morphisms is bilinear. Moreover, the product $G_1 \times G_2$ of two commutative complex Lie groups is both a product and a coproduct of G_1 and G_2 in \mathcal{C} . \square

Although the category Alg of commutative complex algebraic groups is an abelian category ([Mil17a, Thm. 5.62]), as Example F.4.1.2 and Example F.4.1.3 show, \mathcal{C} is NOT an abelian category.

Example F.4.1.2. The map $i : \mathbb{Z}^2 \rightarrow \mathbb{C}$ defined by $(a, b) \mapsto a + b\sqrt{2}$ is injective. The image is not closed in \mathbb{C} as it is dense in \mathbb{R} . For every morphism $f : \mathbb{C} \rightarrow X$ in \mathcal{C} , with $fi = 0$, we have $f = 0$ by identity theorem for holomorphic maps. Thus i is a monomorphism and epimorphism in \mathcal{C} , but not an isomorphism.

Example F.4.1.3. Let $p : \mathbb{C}^2 \rightarrow \mathbb{C}^2/\mathbb{Z}^4$ be the natural projection. Let $i : \mathbb{C} \rightarrow \mathbb{C}^2$ be the closed embedding defined by $z \mapsto (z, \sqrt{2}z)$. Then the composition $pi : \mathbb{C} \rightarrow \mathbb{C}^2/\mathbb{Z}^4$ is an injective morphism (hence a monomorphism) in \mathcal{C} . By [Lee13, Example 7.19], $pi(C)$ is a connected dense subset of $\mathbb{C}^2/\mathbb{Z}^4$. In particular, pi is an epimorphism in \mathcal{C} . The cokernel of pi is the zero morphism $\mathbb{C}^2/\mathbb{Z}^4 \rightarrow 0$. However, pi is not an isomorphism in \mathcal{C} .

Proposition F.4.1.4 3 is a special case of [Con14, Prop. D.2.1]. An elementary proof is given.

Proposition F.4.1.4.

1. $\text{Hom}_{\mathcal{C}}(\mathbb{C}^*, \mathbb{C}) = 0$.
2. For $A \in \mathcal{C}$, the map

$$\text{Hom}_{\mathcal{C}}(\mathbb{C}^n, A) \rightarrow \text{Hom}_{\text{Vec}}(L(\mathbb{C}^n), L(A)), \quad f \mapsto d_e f$$

is a group isomorphism.

3. Let $f : \mathbb{C}^* \rightarrow \mathbb{C}^*$ be a morphism in \mathcal{C} . Then there is an integer k such that $f(z) = z^k$ for every $z \in \mathbb{C}^*$. Hence an isomorphism $\mathbb{Z} = \text{Hom}_{\mathcal{C}}(\mathbb{C}^*, \mathbb{C}^*)$.

Proof. The Lie algebra of \mathbb{C}^* is \mathbb{C} . The exponential map $\exp : \mathbb{C} \rightarrow \mathbb{C}^*$ is normalized as $w \mapsto e^{2\pi i w}$.

1. Let $f : \mathbb{C}^* \rightarrow \mathbb{C}^*$ be a morphism. Then $d_e f : \mathbb{C} \rightarrow \mathbb{C}$ is linear. There is $a \in \mathbb{C}$ with $d_e f(v) = av$ for all $v \in \mathbb{C}$. Since $1 \in \mathbb{C} = L(\mathbb{C}^*)$ is mapped to $1 \in \mathbb{C}^*$ under the exponential map $\exp(2\pi i \bullet)$, one has $0 = f(1) = d_e f(1) = a$. Then $d_e f = 0$ and $f = 0$.
2. It follows from the fact that \mathbb{C}^n is simply connected and both groups are commutative.
3. Consider the induced linear map on Lie algebras $df : \mathbb{C} \rightarrow \mathbb{C}$. There is a unique complex number k such that $df(w) = kw$ for all $w \in \mathbb{C}$. Then

$$e^{2\pi i k} = \exp(df(1)) = f \exp(1) = f(1) = 1.$$

Therefore, k is an integer. For every $z \in \mathbb{C}^*$, there is $w \in \mathbb{C}$ with $\exp(w) = z$. Then $f(z) = f(\exp(w)) = \exp df(w) = \exp(kw) = z^k$.

□

For $A, B \in \mathcal{C}$, the set of isomorphism classes of *commutative* extensions of A by B is denoted by $\text{Ext}(A, B)$.

Proposition F.4.1.5.

1. $\text{Ext}(\bullet, \bullet) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \text{Set}$ is a covariant functor.
2. Let \mathcal{E} be the collection of extensions in \mathcal{C} . Then the pair $(\mathcal{C}, \mathcal{E})$ is an exact category.⁷

Proof. 1. Fix $A, B \in \mathcal{C}$ and an element of $\text{Ext}(A, B)$: $0 \rightarrow B \xrightarrow{i} C \xrightarrow{p} A \rightarrow 0$.

(a) If $f : B \rightarrow B'$ is a morphism in \mathcal{C} , then

$$g : B \rightarrow C \times B', \quad b \mapsto (-b, f(b))$$

is a morphism in \mathcal{C} . It is injective and the (set-theoretic) image is closed in $C \times B'$. By Fact F.2.0.1 2, g identifies B as a complex Lie subgroup of $C \times B'$. Let f_*C be the quotient $(C \times B')/B$ provided by Fact F.2.0.3. The canonical map $B' \rightarrow C \times B'$ induces an injective morphism $f_*i : B' \rightarrow f_*C$ since $B \cap (\{0\} \times B') = \{0\}$. Moreover, B is in the kernel of the composition $C \times B' \rightarrow A$ by $(c, \beta) \mapsto p(c)$, hence a surjective morphism $f_*p : f_*C \rightarrow A$.

Note that $f_*p \circ f_*i = 0$, so $f_*i(B') \subset \ker(f_*p)$. For every element x of $\ker(f_*p)$, take a representative $(c, \beta) \in C \times B'$. As $p(c) = 0$, $c \in B$. Then $(0, \beta + f(c)) - (c, \beta) = g(c)$. Therefore,

$$x = [(0, \beta + f(c))] = f_*i(\beta + f(c)) \in f_*(B').$$

Thus, $f_*i(B') = \ker(f_*p)$

Therefore, the sequence

$$0 \rightarrow B' \xrightarrow{f_*i} f_*C \xrightarrow{f_*p} A \rightarrow 0$$

is exact and $f_*C \in \text{Ext}(A, B')$. Hence a morphism $f_* : \text{Ext}(A, B) \rightarrow \text{Ext}(A, B')$ in the category Set .

Let F be the canonical morphism $C \rightarrow f_*C$. By construction, the extension $f_*C \in \text{Ext}(A, B')$ has the following universal property: for every morphism $h : A \rightarrow A'$ in \mathcal{C} , every $C' \in \text{Ext}(A', B')$, every morphism $G : C \rightarrow C'$ making the diagram commutative

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \longrightarrow & C & \longrightarrow & A \longrightarrow 0 \\ & & \downarrow f & & F \downarrow \searrow & & \downarrow \text{Id} \\ 0 & \longrightarrow & B' & \longrightarrow & f_*C & \xrightarrow{G} & A \longrightarrow 0 \\ & & \downarrow \text{Id} & & u \downarrow \swarrow & & \downarrow h \\ 0 & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & A' \longrightarrow 0, \end{array} \quad (\text{F.7})$$

⁷see [Sta24, Tag 05SF]

there exists a unique morphism $u : f_*C \rightarrow C'$ keeping the diagram commutative.

- (b) If $h : A' \rightarrow A$ is a morphism in \mathcal{C} , by Example F.3.1.1, we get a morphism $h^* : \text{Ext}(A, B) \rightarrow \text{Ext}(A', B)$ in the category Set . Let F be the canonical projection $h^*C \rightarrow C$. By construction, the extension g^*C has the following universal property: for every morphism $g : B' \rightarrow B$, every extension $C' \in \text{Ext}(A', B')$, every morphism $G : C' \rightarrow C$ making the following diagram commutative

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & A' & \longrightarrow & 0 \\
 & & \downarrow g & & \downarrow v & & \downarrow \text{Id} & & \\
 0 & \longrightarrow & B & \longrightarrow & h^*C & \xrightarrow{G} & A' & \longrightarrow & 0 \\
 & & \downarrow \text{Id} & & \downarrow F & & \downarrow h & & \\
 0 & \longrightarrow & B & \longrightarrow & C & \longrightarrow & A & \longrightarrow & 0,
 \end{array} \quad (\text{F.8})$$

there exists a unique morphism $v : C' \rightarrow h^*C$ keeping the diagram commutative.

- (c) Let $f : B \rightarrow B'$, $g : A \rightarrow A'$ be morphisms in \mathcal{C} , $C \in \text{Ext}(A, B)$, and $C' \in \text{Ext}(A', B')$. Then the relation $f_*C = g^*C'$ in $\text{Ext}(A, B')$ is equivalent to the existence of a morphism $F : C \rightarrow C'$ making a commutative diagram

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & B & \longrightarrow & C & \longrightarrow & A & \longrightarrow & 0 \\
 & & \downarrow f & & \downarrow F & & \downarrow g & & \\
 0 & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & A' & \longrightarrow & 0.
 \end{array}$$

Indeed, it follows from the universal properties in Points (1a) and (1b). For every $X \in \text{Ext}(A', B)$, in view of the diagram

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & B & \longrightarrow & g^*X & \longrightarrow & A & \longrightarrow & 0 \\
 & & \downarrow \text{Id} & & \downarrow & & \downarrow g & & \\
 0 & \longrightarrow & B & \longrightarrow & X & \longrightarrow & A' & \longrightarrow & 0 \\
 & & \downarrow f & & \downarrow & & \downarrow \text{Id} & & \\
 0 & \longrightarrow & B' & \longrightarrow & f_*X & \longrightarrow & A' & \longrightarrow & 0,
 \end{array}$$

one has $f_*g^*X = g^*f_*X$. This completes the proof.

2. It follows from Point 1 and Lemma F.4.1.1. □

Example F.4.1.6. If A is a complex torus with $\dim A = g$, B is the discrete group \mathbb{Q}/\mathbb{Z} , then $\text{Ext}(A, B)$ is isomorphic to B^{2g} by Fact F.3.2.6. Even

though B is an injective object of \mathbf{Ab} , the functor $\mathrm{Ext}(\cdot, B) : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Ab}$ is nonzero.

Example F.4.1.7. For an extension $0 \rightarrow B \xrightarrow{i} C \xrightarrow{p} A \rightarrow 0$ in \mathcal{C} , the pushout $i_*C \in \mathrm{Ext}(A, C)$ is the trivial extension. In fact, $i_*C = C \times C/B$ with the embedding

$$B \rightarrow C \times C, \quad b \mapsto (-b, b).$$

The group law $C \times C \rightarrow C$ descends to a morphism $r : i_*C \rightarrow C$. Then $r \circ i_*(i) = \mathrm{Id}_C$. By Fact F.3.0.4, i_*C is trivial.

Similarly, as the diagonal inclusion $C \rightarrow C \times C$ factors through a morphism $s : C \rightarrow p^*C$ and $p^*(p) \circ s = \mathrm{Id}_C$, the pullback $p^*C \in \mathrm{Ext}(C, B)$ is also trivial.

Fact F.4.1.8 follows from Proposition F.4.1.5.

Fact F.4.1.8 ([Ros58, Prop. 5], [Ser88, Prop. 1, p.163]). *1. For every $A, B \in \mathcal{C}$, under the Baer sum $\mathrm{Ext}(A, B)$ is an abelian subgroup of $\mathrm{Ext}(A, B, 1)$.*

2. The functor $\mathrm{Ext}(\bullet, \bullet) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Ab}$ is an additive bifunctor.

3. For any $C, C' \in \mathrm{Ext}(A, B)$, the product $C \times C'$ is naturally an element of $\mathrm{Ext}(A \times A, B \times B)$.

*4. Let $d : A \rightarrow A \times A$ the diagonal map of A and $s : B \times B \rightarrow B$ the group law of B . Then $C + C' = d^*s_*(C \times C')$ in $\mathrm{Ext}(A, B)$.*

Corollary F.4.1.9. *For every commutative complex Lie group A , the restriction $\mathrm{Ext}(A, \cdot) : \mathrm{Vec} \rightarrow \mathbf{Ab}$ factors through a functor from Vec to the category of all complex vector spaces.*

By Example F.4.3.3 below, for every $V \in \mathrm{Vec}$, $\dim_{\mathbb{C}} \mathrm{Ext}(A, V)$ is finite. Hence an additive functor $\mathrm{Ext}(A, \cdot) : \mathrm{Vec} \rightarrow \mathrm{Vec}$.

Example F.4.1.10. Endowing each object of \mathbf{Ab}_c the discrete topology gives a functor $\mathbf{Ab}_c \rightarrow \mathcal{C}$, identifying \mathbf{Ab}_c as a full subcategory of \mathcal{C} . The subcategory \mathbf{Ab}_c is closed under extension by Fact F.3.0.1. From Proposition F.3.2.1, the forgetful natural transformation $\mathrm{Ext}(-, +) \rightarrow \mathrm{Ext}_{\mathbb{Z}}^1(-, +)$ is an isomorphism of functors $\mathbf{Ab}_c^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathbf{Ab}$.

Example F.4.1.11. Analytification functor $(\bullet)^{\mathrm{an}} : \mathbf{Alg} \rightarrow \mathcal{C}$ identifies \mathbf{Alg} as a subcategory of \mathcal{C} (which is not full). The extension problem within the subcategory \mathbf{Alg} is discussed by Rosenlicht [Ros58] and Serre [Ser88, Ch. VII]. They define a similar additive functor $\mathrm{Ext}_{\mathbf{Alg}} : \mathbf{Alg}^{\mathrm{op}} \times \mathbf{Alg} \rightarrow \mathbf{Ab}$. For every $A, B \in \mathbf{Alg}$, there is a natural morphism $\mathrm{Ext}_{\mathbf{Alg}}(A, B) \rightarrow \mathrm{Ext}(A^{\mathrm{an}}, B^{\mathrm{an}})$. In general, this morphism is neither injective nor surjective.

By [MM66, Introduction, 1.], when A is a complex abelian variety, one has $\mathrm{Ext}_{\mathbf{Alg}}(\mathbb{G}_a, A) = 0$ while $\mathrm{Ext}_{\mathbf{Alg}}(\mathbb{G}_m, A)$ is (non-canonically) isomorphic

to the torsion subgroup A_{tor} of A . But $\text{Ext}(\mathbb{C}^*, A^{\text{an}}) = 0$ by Proposition F.3.2.3, so the natural morphism $\text{Ext}_{\text{Alg}}(\mathbb{G}_m, A) \rightarrow \text{Ext}(\mathbb{C}^*, A^{\text{an}})$ is not injective.

For any two complex abelian varieties X_i ($i = 1, 2$) of positive dimension, $\text{Ext}_{\text{Alg}}(X_2, X_1)$ is countable while $\text{Ext}(X_2^{\text{an}}, X_1^{\text{an}})$ is uncountable. In fact, the natural morphism $\text{Ext}_{\text{Alg}}(X_2, X_1) \rightarrow \text{Ext}(X_2^{\text{an}}, X_1^{\text{an}})$ is an embedding onto the torsion subgroup of $\text{Ext}(X_2^{\text{an}}, X_1^{\text{an}})$ ([BL99, Ch. 1; Prop. 6.1, Cor. 6.3]).

Lemma F.4.1.12 is mentioned at the bottom of [Hoc51b, p.546].

Lemma F.4.1.12. *If G is a commutative connected Lie group, then G is a divisible \mathbb{Z} -module.*

Proof. The exponential map $\exp : L(G) \rightarrow G$ is surjective. For every $x \in G$, there is $v \in L(G)$ with $\exp(v) = x$. For every integer $n \geq 1$, $\exp(v/n) \in G$ and $n(\exp(v/n)) = x$. \square

Corollary F.4.1.13. *An extension $0 \rightarrow B \rightarrow C \rightarrow A \rightarrow 0$ in \mathcal{C} with B connected and A discrete is trivial. In particular, for every $G \in \mathcal{C}$, the natural exact sequence*

$$0 \rightarrow G_0 \rightarrow G \rightarrow G/G_0 \rightarrow 0$$

is a trivial extension, hence a non-canonical isomorphism $G \rightarrow G_0 \times G/G_0$ in \mathcal{C} .

Proof. By Lemma F.4.1.12, the \mathbb{Z} -module B is divisible, so the functor $\text{Ext}_{\mathbb{Z}}^1(\cdot, B) : \text{Ab} \rightarrow \text{Ab}$ is zero. Since A is discrete, the result follows from Example F.4.1.10. \square

Example F.4.1.14. The abelian group underlying a complex torus B is divisible by Lemma F.4.1.12, hence an injective object of Ab and $\text{Ext}_{\mathbb{Z}}^1(\bullet, B) : \text{Ab} \rightarrow \text{Ab}$ is zero. However, $\text{Ext}(\bullet, B) : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}$ can be nonzero. In fact, [BL04, (8) b), p.68] gives an example of a nontrivial exact sequence of complex tori

$$0 \rightarrow B \rightarrow C \rightarrow A \rightarrow 0$$

with $\dim A = \dim B = 1$.

F.4.2 Exact sequences of Ext

Let $0 \rightarrow A' \xrightarrow{i} A \xrightarrow{p} A'' \rightarrow 0$ be an exact sequence in \mathcal{C} , i.e., $A \in \text{Ext}(A'', A')$. For $f \in \text{Hom}(A', B)$, there is $f_* A \in \text{Ext}(A'', B)$. Hence a map

$$d : \text{Hom}(A', B) \rightarrow \text{Ext}(A'', B), \quad d(f) = f_* A.$$

Then d is a group morphism. The formation of d is functorial in B .

Proposition F.4.2.1. *Let $B \in \mathcal{C}$. The sequence in \mathbf{Ab} with obvious morphisms*

$0 \rightarrow \operatorname{Hom}_{\mathcal{C}}(A'', B) \rightarrow \operatorname{Hom}_{\mathcal{C}}(A, B) \rightarrow \operatorname{Hom}_{\mathcal{C}}(A', B) \xrightarrow{d} \operatorname{Ext}(A'', B) \xrightarrow{p^*} \operatorname{Ext}(A, B) \rightarrow \operatorname{Ext}(A', B)$
is exact and functorial in B .

Proof.

- Exactness at $\operatorname{Hom}(A, B)$ follows from Fact F.2.0.1.
- Exactness at $\operatorname{Hom}(A', B)$: By Example F.4.1.7, the composition

$$\operatorname{Hom}(A, B) \xrightarrow{i_*} \operatorname{Hom}(A', B) \rightarrow \operatorname{Ext}(A'', B)$$

is zero. Now take $\phi \in \ker(d)$. By Fact F.3.0.4, there is a morphism $r : \phi_* A \rightarrow B$ with $r\phi_*(i) = \operatorname{Id}_B$. Let $F : A \rightarrow \phi_* A$ be the canonical morphism. Then $rFi = r\phi_*(i)\phi = \phi$. Hence $\phi \in \operatorname{im}(i_*)$.

- Exactness at $\operatorname{Ext}(A'', B)$: By Example F.4.1.7, for every $\phi \in \operatorname{Hom}(A', B)$, $p^*d\phi = p^*\phi_* A = \phi_* p^* A = 0$, i.e., the composition

$$\operatorname{Hom}(A', B) \xrightarrow{d} \operatorname{Ext}(A'', B) \xrightarrow{p^*} \operatorname{Ext}(A, B)$$

is zero.

Now take $C \in \ker(p^*) \subset \operatorname{Ext}(A'', B)$ with connecting morphisms $f : B \rightarrow C$ and $g : C \rightarrow A''$. By Fact F.3.0.4, there is a morphism $s : A \rightarrow p^*C$ with $p^*(p) \circ s = \operatorname{Id}_A$. For every $a' \in A'$, the image of $s(a')$ in A'' is $p(a') = 0$, so the image of $s(a')$ in C lies in B . Thus, the restriction of s to A' is a morphism $\phi : A' \rightarrow B$. By construction, the extension group of $d(\phi) = \phi_* A \in \operatorname{Ext}(A'', B)$ is $A \times B/D$, where $D = \{(-a', \phi(a')) : a' \in A'\}$.

Define $\psi : A \rightarrow C$ by $\psi = F \circ s$. Define

$$A \times B \rightarrow C, \quad (a, b) \mapsto \psi(a) + f(b).$$

For every $a' \in A'$, $\psi(-a') + f(s(a')) = 0$, hence a factorization $\phi_* A \rightarrow C$ in the middle keeping the diagram commutative:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & A' & \xrightarrow{i} & A & \xrightarrow{p} & A'' & \longrightarrow & 0 \\
 & & \downarrow \phi & & \downarrow & & \downarrow \operatorname{Id} & & \\
 0 & \longrightarrow & B & \xrightarrow{\phi_* i} & \phi_* A & \longrightarrow & A'' & \longrightarrow & 0 \\
 & & \downarrow \operatorname{Id} & & \downarrow & & \downarrow \operatorname{Id} & & \\
 0 & \longrightarrow & B & \xrightarrow{f} & C & \xrightarrow{g} & A'' & \longrightarrow & 0 \\
 & & \uparrow \operatorname{Id} & & \uparrow F & & \uparrow p & & \\
 0 & \longrightarrow & B & \longrightarrow & p^* C & \xrightarrow{p^* p} & A & \longrightarrow & 0.
 \end{array}$$

$\swarrow \text{---} s \text{---} \nwarrow$

Then $C = \phi_* A = d\phi$ in $\text{Ext}(A'', B)$. Therefore, $\ker(p^*) = \text{im}(d)$.

- **Exactness at $\text{Ext}(A, B)$:** As the composition $A' \rightarrow A \rightarrow A''$ is zero and $\text{Ext}(\bullet, B) : C^{\text{op}} \rightarrow \text{Ab}$ is an additive functor, the composition $\text{Ext}(A'', B) \rightarrow \text{Ext}(A, B) \rightarrow \text{Ext}(A', B)$ is zero.

Conversely, if $C_1 \in \text{Ext}(A, B)$ with $i^* C_1 = 0$ in $\text{Ext}(A', B)$, then there is a morphism $s : A' \rightarrow i^* C_1$ with $i^* g \circ s = \text{Id}_{A'}$. The composition $\phi : A' \rightarrow C_1$ is injective. Indeed, if $a' \in \ker(\phi)$, then $s(a') = (a', 0)$ in $A' \times C_1$. Thus, $i(a') = 0$ by the construction of pullback extension. Since i is injective, $a' = 0$.

Let $C_1 \rightarrow C = C_1/\phi(A')$ be the quotient morphism. Let $f_0 : B \rightarrow C$ be the induced morphism. Then f_0 is injective. Indeed, if $b \in \ker(f_0)$, then $f(b) = \phi(a')$ for some $a' \in A'$. Then $(a', f(b)) \in i^* C_1$, so $i(a') = gf(b) = 0$. Hence $a' = 0$ and $f(b) = 0$. Therefore, $b = 0$.

Because $pg\phi = p \circ i = 0$, the morphism $pg : C_1 \rightarrow A''$ descends to a surjective morphism $g_0 : C \rightarrow A''$. We prove that the bottom row of the following diagram is exact:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & B & \longrightarrow & i^* C_1 & \xrightarrow{\quad \quad} & A' \longrightarrow 0 \\
 & & \downarrow \text{Id} & & \downarrow & \swarrow i^* g & \downarrow i \\
 0 & \longrightarrow & B & \xrightarrow{\quad f \quad} & C_1 & \xrightarrow{\quad g \quad} & A \longrightarrow 0 \\
 & & \downarrow \text{Id} & & \downarrow & \swarrow \phi & \downarrow p \\
 0 & \longrightarrow & B & \xrightarrow{\quad f_0 \quad} & C & \xrightarrow{\quad g_0 \quad} & A'' \longrightarrow 0.
 \end{array}$$

Since $gf = 0$, one has $g_0 f_0 = 0$. Therefore, $f_0(B) \subset \ker(g_0)$. Conversely, for $c \in \ker(g_0)$, there is $c_1 \in C_1$ with $[c_1] = c$. Since $pg(c_1) = g_0(c) = 0$, one gets $g(c_1) \in A'$. Then $g\phi g(c_1) = gc_1$. So $c_1 - \phi g(c_1) \in \ker(g) = B$ and

$$f_0(c_1 - \phi g(c_1)) = [c_1 - \phi g(c_1)] = c.$$

Therefore, $\ker(g_0) = f_0(B)$. In particular, the bottom row is exact, i.e., $C \in \text{Ext}(A'', B)$. By the universal property showed in the diagram (F.8), $C_1 = p^* C$.

□

Example F.4.2.2. Let A be a complex torus, and let B be a finite abelian group. Then $\text{Hom}_{\mathcal{C}}(A, B) = 0$. Let integer $n(\geq 1)$ be a multiple of $\#B$. Applying Proposition F.4.2.1 to the exact sequence in \mathcal{C}

$$0 \rightarrow A[n] \rightarrow A \xrightarrow{[n]_A} A \rightarrow 0,$$

one gets an exact sequence in Ab :

$$0 \rightarrow \text{Hom}(A[n], B) \rightarrow \text{Ext}(A, B) \xrightarrow{f} \text{Ext}(A, B).$$

Since the morphism $[n]_B : B \rightarrow B$ is zero in \mathcal{C} , by Fact F.4.1.8, $f = ([n]_B)_* = 0$. Hence an isomorphism $\text{Hom}(A[n], B) \rightarrow \text{Ext}(A, B)$ that is functorial in B , which is also confirmed by Fact F.3.2.6.

Let $0 \rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0$ be an exact sequence in \mathcal{C} . If $A \in \mathcal{C}$ and $\phi \in \text{Hom}(A, B'')$, then $\phi^*B \in \text{Ext}(A', B)$. Define a map $d : \text{Hom}(A, B'') \rightarrow \text{Ext}(A, B')$ by $d(\phi) = \phi^*B$.

Proposition F.4.2.3. *Let $0 \rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0$ be an exact sequence in \mathcal{C} and $A \in \mathcal{C}$. Then the sequence*

$$0 \rightarrow \text{Hom}(A, B') \rightarrow \text{Hom}(A, B) \rightarrow \text{Hom}(A, B'') \xrightarrow{d} \text{Ext}(A, B') \rightarrow \text{Ext}(A, B) \rightarrow \text{Ext}(A, B'')$$

in Ab is exact and functorial in A .

The proof is analogous to that of Proposition F.4.2.1 and is thereby omitted.

Consider the extension problem with connected bases. Corollary F.4.2.4 should be compared to [Sha49, Thm. 1]: for two compact connected real Lie groups G, H , the cokernel of the restriction morphism $\text{Hom}(\tilde{H}, Z(G)) \rightarrow \text{Hom}(\pi_1(H), Z(G))$ is isomorphic to the group of extensions of H by G .

Corollary F.4.2.4. *Let A, B be commutative complex Lie groups. Assume that A is connected with universal cover $\omega : \tilde{A} \rightarrow A$. Then there is a canonical exact sequence in Ab :*

$$0 \rightarrow \text{Hom}_{\mathcal{C}}(A, B) \xrightarrow{\omega^*} \text{Hom}_{\mathcal{C}}(\tilde{A}, B) \xrightarrow{r} \text{Hom}_{\text{Ab}}(\pi_1(A), B) \rightarrow \text{Ext}(A, B) \rightarrow 0, \quad (\text{F.9})$$

where r is induced by restriction.

Proof. By Proposition F.3.2.2, Fact F.3.2.6 and Corollary F.4.1.13, the functor $\text{Ext}(\mathbb{C}, \bullet) : \mathcal{C} \rightarrow \text{Ab}$ is zero. By Fact F.4.1.8,

$$\text{Ext}(\mathbb{C}^n, \bullet) = 0. \quad (\text{F.10})$$

The proof is concluded by Proposition F.4.2.1. \square

Example F.4.2.5. In Corollary F.4.2.4, if B discrete, then $\text{Hom}_{\mathcal{C}}(\tilde{A}, B) = 0$ and the natural morphism $\text{Hom}(\pi_1(A), B) \rightarrow \text{Ext}(A, B)$ is an isomorphism, which agrees with Fact F.3.2.6.

F.4.3 Determination of commutative extension group

The commutative extension problem of complex Lie groups is answered by Proposition F.4.3.1. Fix two commutative complex Lie groups A, B .

Proposition F.4.3.1. *There is a non-canonical isomorphism in Ab :*

$$\text{Ext}(A, B) \rightarrow \text{Ext}_{\mathbb{Z}}^1(A/A_0, B/B_0) \oplus \text{Hom}_{\text{Ab}}(\pi_1(A_0), B/B_0) \oplus \text{Ext}(A_0, B_0),$$

and $\text{Ext}(A_0, B_0)$ is the cokernel of the natural restriction morphism

$$s : \text{Hom}_{\text{Vec}}(L(A), L(B)) \rightarrow \text{Hom}_{\text{Ab}}(\pi_1(A_0), B_0).$$

Proof. By Corollary F.4.1.13, there are non-canonical isomorphisms in \mathcal{C} : $A \rightarrow A/A_0 \times A_0$ and $B \rightarrow B/B_0 \times B_0$. Using Fact F.4.1.8, one gets an isomorphism in Ab :

$$\text{Ext}(A, B) \rightarrow \text{Ext}(A/A_0, B_0) \oplus \text{Ext}(A/A_0, B/B_0) \oplus \text{Ext}(A_0, B/B_0) \oplus \text{Ext}(A_0, B_0).$$

The first factor $\text{Ext}(A/A_0, B_0) = 0$ by Corollary F.4.1.13. By Example F.4.1.10, the natural morphism $\text{Ext}(A/A_0, B/B_0) \rightarrow \text{Ext}_{\mathbb{Z}}^1(A/A_0, B/B_0)$ is an isomorphism. Fact F.3.2.6 gives a natural isomorphism $\text{Hom}_{\text{Ab}}(\pi_1(A_0), B/B_0) \rightarrow \text{Ext}(A_0, B/B_0)$. Corollary F.4.2.4 identifies $\text{Ext}(A_0, B_0)$ with the cokernel of the restriction map $r : \text{Hom}_{\mathcal{C}}(\tilde{A}_0, B_0) \rightarrow \text{Hom}_{\text{Ab}}(\pi_1(A_0), B_0)$. By Proposition F.4.1.4 2, the group morphism

$$t : \text{Hom}_{\mathcal{C}}(\tilde{A}_0, B_0) \rightarrow \text{Hom}_{\text{Vec}}(L(A), L(B)), \quad \phi \mapsto d_e \phi$$

is an isomorphism. The proof is finished by setting $s = rt^{-1}$. \square

For every $C \in \text{Ext}(A, B)$, by Fact F.2.0.3, the morphism $C \rightarrow A$ is a principal B -bundle. The bijection (F.1) gives rise to a canonical map

$$\pi : \text{Ext}(A, B) \rightarrow H^1(A, \mathcal{B}_A). \quad (\text{F.11})$$

Fact F.4.3.2 is taken from [Ros58, pp.698-699] and the proof of [Ser88, Ch. VII, no. 5, Prop. 5].

Fact F.4.3.2. *The map (F.11) is a group morphism and the formation of π is functorial, in the sense that it commutes with the morphisms $f_* : \text{Ext}(A, B) \rightarrow \text{Ext}(A, B')$ defined by $f : B \rightarrow B'$ and $g^* : \text{Ext}(A, B) \rightarrow \text{Ext}(A', B)$ defined by $g : A' \rightarrow A$. When B is a vector group, the map π is \mathbb{C} -linear.*

Example F.4.3.3. Let X be a toroidal group, and let $\omega : \tilde{X} \rightarrow X$ be the universal covering of kernel F . Then F is a discrete subgroup of the vector space \tilde{X} . By Proposition F.4.2.1,

$$\text{Hom}_{\mathcal{C}}(X, \mathbb{C}) \rightarrow \text{Hom}_{\mathcal{C}}(\tilde{X}, \mathbb{C}) \rightarrow \text{Hom}_{\mathcal{C}}(F, \mathbb{C}) \rightarrow \text{Ext}(X, \mathbb{C}) \rightarrow \text{Ext}(\tilde{X}, \mathbb{C})$$

is an exact sequence in Ab. From Definition F.2.0.9, $\text{Hom}_{\mathcal{C}}(X, \mathbb{C}) = 0$. By Proposition F.10, $\text{Ext}(\tilde{X}, \mathbb{C}) = 0$. Hence the first exact row of Diagram (F.12).

According to [AK01, p.48], there is a \mathbb{C} -linear isomorphism $\text{Hom}_{\mathcal{C}}(\tilde{X}, \mathbb{C}) \rightarrow H^0(X, \Omega_X^1)$ and every global holomorphic 1-form on X is d -closed. So taking de Rham cohomology class results in a linear map $H^0(X, \Omega_X^1) \rightarrow H^1(X, \mathbb{C})$. The inclusion $\mathbb{C}_X \rightarrow \mathcal{O}_X$ induces a linear map $H^1(X, \mathbb{C}) \rightarrow H^1(X, \mathcal{O}_X)$. By universal coefficient theorem (see, e.g., [Hat05, Thm. 3.2]), the natural morphism $\text{Hom}_{\mathcal{C}}(F, \mathbb{C}) \rightarrow H^1(X, \mathbb{C})$ is an isomorphism. Hence a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}_{\mathcal{C}}(\tilde{X}, \mathbb{C}) & \longrightarrow & \text{Hom}_{\mathcal{C}}(F, \mathbb{C}) & \longrightarrow & \text{Ext}(X, \mathbb{C}) \longrightarrow 0 \\ & & \downarrow \simeq & & \downarrow \simeq & & \downarrow (F.11) \\ 0 & \longrightarrow & H^0(X, \Omega_X^1) & \longrightarrow & H^1(X, \mathbb{C}) & \longrightarrow & H^1(X, \mathcal{O}_X). \end{array} \quad (\text{F.12})$$

Let $b_1(X) := \dim_{\mathbb{C}} H^1(X, \mathbb{C})$ be the first Betti number of X , i.e., the \mathbb{Z} -rank of F . From [AK01, p.48], as a \mathbb{C} -vector space

$$\text{Ext}(X, \mathbb{C}) = \frac{H^1(X, \mathbb{C})}{H^0(X, \Omega_X^1)} \quad (\text{F.13})$$

is of dimension $b_1(X) - \dim X$.

If X is a toroidal theta group,⁸ then $\pi : \text{Ext}(X, \mathbb{C}) \rightarrow H^1(X, \mathcal{O}_X)$ is a \mathbb{C} -linear isomorphism by [AK01, Thm. 2.2.6 b)]. Otherwise, X is a toroidal wild group⁸ and $H^1(X, \mathcal{O}_X)$ is infinite dimensional by [AK01, Prop. 2.2.7].

A seemingly different way to compute the last factor in Proposition F.4.3.1, i.e., the group of commutative extensions of two connected commutative complex Lie groups, is given in Example F.4.3.4.

Example F.4.3.4. Start by the special case that X is a toroidal group and B is a connected commutative complex Lie group. Denote the kernel of the universal cover of B (resp. X) by $\iota : K \rightarrow \tilde{B}$ (resp. $F \rightarrow \tilde{X}$). By (F.10) and Proposition F.4.1.4 2, the sequence

$$0 \rightarrow \text{Hom}_{\mathcal{C}}(\tilde{X}, K) \rightarrow \text{Hom}_{\mathcal{C}}(\tilde{X}, \tilde{B}) \rightarrow \text{Hom}_{\mathcal{C}}(\tilde{X}, B) \rightarrow 0$$

is exact in Ab. As F is a free \mathbb{Z} -module,

$$0 \rightarrow \text{Hom}_{\mathcal{C}}(F, K) \rightarrow \text{Hom}_{\mathcal{C}}(F, \tilde{B}) \rightarrow \text{Hom}_{\mathcal{C}}(F, B) \rightarrow 0$$

in Ab is also exact. Applying Proposition F.4.2.1 and the snake lemma to the commutative diagram

⁸in the sense of [AK01, Def. 2.2.1]

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(\tilde{X}, K) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(\tilde{X}, \tilde{B}) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(\tilde{X}, B) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(F, K) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(F, \tilde{B}) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(F, B) \longrightarrow 0,
\end{array}$$

one gets an exact sequence in Ab :

$$0 \rightarrow \mathrm{Hom}_{\mathcal{C}}(X, B) \xrightarrow{j} \mathrm{Ext}(X, K) \xrightarrow{\iota_*} \mathrm{Ext}(X, \tilde{B}) \rightarrow \mathrm{Ext}(X, B) \rightarrow 0. \quad (\mathrm{F.14})$$

Since K is a free \mathbb{Z} -module, by Fact F.3.2.6, $\mathrm{Ext}(X, K) = H^1(X, \mathbb{Z}) \otimes_{\mathbb{Z}} K$. By Fact F.4.1.8 and (F.13), one has

$$\mathrm{Ext}(X, \tilde{B}) = \frac{H^1(X, \mathbb{C})}{H^0(X, \Omega_X^1)} \otimes_{\mathbb{C}} \tilde{B}.$$

The group morphism ι_* is induced by the \mathbb{Z} -bilinear map

$$H^1(X, \mathbb{Z}) \times K \rightarrow \left(\frac{H^1(X, \mathbb{C})}{H^0(X, \Omega_X^1)} \right) \otimes_{\mathbb{C}} \tilde{B}, \quad (\eta, x) \mapsto [\eta] \otimes \iota(x).$$

Thus we can compute $\mathrm{Ext}(X, B)$ from (F.14).

For a general connected commutative complex Lie group A , by [AK01, 1.1.5], $A = \mathbb{C}^l \times (\mathbb{C}^*)^m \times X_0$ for some integers $l, m \geq 0$ and a toroidal group X_0 . By Fact F.4.1.8, Proposition F.3.2.2 and Proposition F.3.2.3, $\mathrm{Ext}(A, B) = \mathrm{Ext}(X_0, B)$, reducing to the previous case.

F.5 Commutative extensions of complex tori

F.5.1 Primitive cohomology classes

Every central extension of a compact real Lie group by a vector group is trivial, shown by Fact F.5.1.1.

Fact F.5.1.1 (Iwasawa, [Iwa49, Lem. 3.7], [Hoc51a, Footnote 10, p.107]). *Let (F.3) be an exact sequence of real Lie groups. If K is a vector group and Q is compact, then this extension is a semidirect product. In particular, if this extension is central, then it is trivial.*

Contrary to the real case, Example F.5.1.2 shows a commutative extension of a complex torus by a vector group can be nontrivial.

Example F.5.1.2 ([MM60, p.145, Exemple], [LH76, Sec. I.3]). Set $C = \mathbb{C}^* \times \mathbb{C}^*$. Then $B = \{(e^z, e^{iz}) : z \in \mathbb{C}\}$ is a complex Lie subgroup of C (but not an algebraic subgroup of $\mathbb{G}_m \times \mathbb{G}_m$) isomorphic to \mathbb{C} . The quotient $A = C/B$ is an elliptic curve. The exact sequence $0 \rightarrow B \rightarrow C \rightarrow A \rightarrow 0$ is a nontrivial extension, as C is not biholomorphic to $B \times A$.

In the remainder of Section F.5, unless otherwise specified, let A be a complex torus of dimension g and B be a commutative complex Lie group. Let $s_A : A \times A \rightarrow A$ be the group law of A . The dual of A is $A^\vee = \text{Pic}^0(A)$.

The analogue of Proposition F.5.1.3 for abelian varieties is [Ros58, Prop. 9].

Proposition F.5.1.3. *The morphism (F.11) is injective.*

Proof. Let $C \in \ker(\pi)$. The principal bundle $C \rightarrow A$ is trivial, so there is a morphism $s : A \rightarrow C$ of complex manifolds with $ps = \text{Id}_A$. Then there exists a unique $b \in B$ with $b \cdot s(e_A) = e_C$, where dot signifies the action of B on the fiber $p^{-1}(e_A)$. Define

$$s' : A \rightarrow C, \quad s'(a) = b \cdot s(a).$$

Then s' is a complex manifold morphism with $ps' = \text{Id}_A$. Replacing s by s' , we may suppose that $s(e_A) = e_C$. By [NW13, Thm. 5.1.36], s is a morphism in \mathcal{C} . By Fact F.3.0.4, $C = 0$ in $\text{Ext}(A, B)$. Therefore, π is injective. \square

We propose to determine the image of (F.11). Let Mfd be the category of complex manifolds. Define a functor

$$T : \text{Mfd}^{\text{op}} \rightarrow \text{Ab}, \quad T(X) = H^1(X, \mathcal{B}_X).$$

When X is a point, $T(X) = 0$. Let $X_1, X_2 \in \text{Mfd}$, and let $p_i : X_1 \times X_2 \rightarrow X_i$ ($i = 1, 2$) be the projection to the i -th factor. There is a morphism $p_1^* \oplus p_2^* : T(X_1) \times T(X_2) \rightarrow T(X_1 \times X_2)$.

Definition F.5.1.4. [Ser88, (29), no.14, Ch. VII] For $A \in \mathcal{C}$, an element $x \in T(A) = H^1(A, \mathcal{B}_A)$ is called primitive if $s_A^*(x) = p_1^*(x) + p_2^*(x)$ in $T(A \times A)$. Denote by $\text{PT}(A)$ the subgroup of $T(A)$ formed by the primitive elements.

Fact F.5.1.5 ([Ser88, Lem. 8, p.181]). *The functor $\text{PT} : \mathcal{C}^{\text{op}} \rightarrow \text{Ab}$ is additive.*

Theorem F.5.1.6 is an analytic analog of [Ser88, Thm. 5, p.181].

Theorem F.5.1.6. *Assume that B_0 is linear. Then the image of the morphism (F.11) is the set of primitive elements of $H^1(A, \mathcal{B}_A)$.*

Proof. Take $C \in \text{Ext}(A, B)$ and put $x = \pi(C)$. By Facts F.4.1.8 and F.4.3.2,

$$s_A^*(x) = s_A^*\pi(C) = \pi s_A^*(C) = \pi(p_1^*C + p_2^*C) = p_1^*\pi(C) + p_2^*\pi(C) = p_1^*x + p_2^*x,$$

so x is primitive.

Conversely, let $x \in H^1(A, \mathcal{B}_A)$ be a primitive element and let $p : C \rightarrow A$ be the corresponding principal B -bundle. We show that there exists a structure of commutative complex Lie group on C which makes it an extension of A by B .

By Corollary F.4.1.13, every morphism of complex manifolds $A \rightarrow B$ is constant. Let $C' \rightarrow A \times A$ be the pull-back of $C \rightarrow A$ along $s_A : A \times A \rightarrow A$. As x is primitive, $C' = p_1^*C + p_2^*C$ in $T(A \times A)$. Choose a surjection $p_1^*C \times_{A \times A} p_2^*C \rightarrow C'$ satisfying (F.2). Since $p_1^*C = C \times A$ and $p_2^*C = A \times C$, as a complex manifold $p_1^*C \times_{A \times A} p_2^*C$ is isomorphic to $C \times C$. Hence a morphism $g : C \times C \rightarrow C$ of complex manifolds:

$$\begin{array}{ccc}
 & & g \\
 & \searrow & \nearrow \\
 C \times C = p_1^*C \times_{A \times A} p_2^*C & \longrightarrow & p_1^*C + p_2^*C = C' \longrightarrow C \\
 & \downarrow & \downarrow p \\
 & A \times A & \xrightarrow{s_A} A
 \end{array} \quad (\text{F.15})$$

By construction, it satisfies

$$g(b \cdot c, b' \cdot c') = (b + b') \cdot g(c, c') \quad (\text{F.16})$$

for every $c, c' \in C$ and $b, b' \in B$.

Choose a point $e \in p^{-1}(e_A)$. Since $p(g(e, e)) = s_A(e_A, e_A) = e_A$, there exists a unique $b \in B$ with $b \cdot g(e, e) = e$. Replacing e by $b \cdot e$, we can suppose that

$$g(e, e) = e. \quad (\text{F.17})$$

We verify that (C, e, g) is a group.

Identity According to (F.15), there is a morphism $h : C \rightarrow B$ of complex manifolds with $g(c, e) = h(c) \cdot c$ for all $c \in C$. By (F.17), $h(e) = e_B$. Furthermore, (F.16) shows that $h(b \cdot c) = h(c)$ for all $b \in B$. Therefore, h factors as $C \xrightarrow{p} A \xrightarrow{\bar{h}} B$. The morphism \bar{h} of complex manifolds is constant, so $g(c, e) = c$ for all $c \in C$. The formula $g(e, c) = c$ is proved similarly.

Associativity According to (F.15), there is a complex manifold morphism $u : C \times C \times C \rightarrow B$ with

$$g(c, g(c', c'')) = u(c, c', c'') \cdot g(g(c, c'), c'')$$

for all $c, c', c'' \in C$. Then $u(e, e, e) = e_B$. Equation (F.16) shows that u factors through a morphism $\bar{u} : A \times A \times A \rightarrow B$ of complex manifolds. Then \bar{u} is of constant value e_B . Therefore, $g(c, g(c', c'')) = g(g(c, c'), c'')$ for all $c, c', c'' \in C$.

Inverse Denote by $i_A : A \rightarrow A$ (resp. $i_B : B \rightarrow B$) the inverse map of A (resp. B). Let $C^- \rightarrow A$ be the principal B -bundle corresponding to

$-x \in H^1(A, \mathcal{B}_A)$. There is a morphism $f : C \rightarrow C^-$ of principal B -bundles over A , such that for every $b \in B, c \in C, f(b \cdot c) = (-b) \cdot c$. Since $0_A = i_A + \text{Id}_A$, by Fact F.5.1.5, $0 = 0_A^*x = i_A^*x + x$, hence $i_A^*x = -x$. In other words, the pullback of $p : C \rightarrow A$ along i_A is $C^- \rightarrow A$.

$$\begin{array}{ccccc} & & i & & \\ & \nearrow & & \searrow & \\ C & \xrightarrow{f} & i_A^*C & \longrightarrow & C \\ & \downarrow & & & \downarrow p \\ & A & \xrightarrow{i_A} & & A \end{array}$$

The induced morphism $i : C \rightarrow C$ of complex manifolds is such that for every $c \in C, b \in B$,

$$i(b \cdot c) = (-b) \cdot i(c). \quad (\text{F.18})$$

Since $i(e) \in p^{-1}(e_A)$, there is $b \in B$ with $b \cdot i(e) = e$. Define $i' : C \rightarrow C$ by $i'(x) = b \cdot i(x)$ and replace i by i' . Then we may further assume that $i(e) = e$. Because

$$p(g(c, i(c))) = s_A(p(c), pi(c)) = s_A(p(c), i_A(p(c))) = e_A,$$

there exists a morphism $v : C \rightarrow B$ of complex manifolds such that $g(c, i(c)) = v(c) \cdot e$ and $v(e) = e_B$. By (F.16) and (F.18), v factors through $\bar{v} : A \rightarrow B$, which is of constant value e_B . Therefore, $g(c, i(c)) = e$ for all $c \in C$.

In conclusion, (C, e, g, i) is a complex Lie group and (F.15) shows that $p : C \rightarrow A$ is a morphism. Define an injective map $\iota : B \rightarrow C$ by $b \mapsto b \cdot e$. By (F.16), then ι is a morphism. Since $\iota(B) = p^{-1}(e)$, the sequence

$$0 \rightarrow B \xrightarrow{\iota} C \xrightarrow{p} A \rightarrow 0$$

is exact. By Proposition F.6.0.2 2 below, C is commutative and hence $C \in \text{Ext}(A, B)$. (The commutativity of C can also be proved using an argument of similar type.) Therefore, $x = \pi(C)$ is in the image of π . \square

F.5.2 The case $B = \mathbb{C}^*$

We review some basics about (holomorphic) line bundles on complex tori.

Definition F.5.2.1. [Wei48, Ch.VIII, n.58] Let $L \rightarrow A$ be a line bundle on a complex torus. If for every $a \in A$, the pullback line bundle T_a^*L is isomorphic to L , then we write $L \equiv O_A$. Here $T_a : A \rightarrow A$ is defined by $T_a(x) = x + a$.

By [BL04, p.36], L induces a morphism

$$\phi_L : A \rightarrow A^\vee, \quad a \mapsto T_a^* L \otimes L^{-1}.$$

Then $L \equiv O_A$ is equivalent to $\phi_L = 0$. Then [BL04, Prop. 2.5.3] becomes Fact F.5.2.2.

Fact F.5.2.2. *Let $L \rightarrow A$ be a line bundle on a complex torus. The following conditions are equivalent:*

1. L is analytically equivalent to O_A ;
2. $L \in \text{Pic}^0(A)$;
3. $L \equiv O_A$.

Proposition F.5.2.3. *Let $L \rightarrow A$ be a line bundle on complex torus. Then $L \equiv O_A$ if and only if $s_A^* L = p_1^* L \otimes p_2^* L$.*

Proof. If $s_A^* L = p_1^* L \otimes p_2^* L$, then for every $a \in A$, the line bundle $T_a^* L = (s_A^* L)|_{A \times a} = (p_1^* L \otimes p_2^* L)|_{A \times a} = L$, i.e., $L \equiv O_A$.

Conversely, if $L \equiv O_A$, then for every $a \in A$, $(s_A^* L)|_{A \times a} = T_a^* L = L = (p_1^* L)|_{A \times a}$. Therefore, $s^* L \otimes p_1^* L^{-1} \rightarrow A \times A$ is a line bundle, whose restriction to $A \times a$ is trivial for all $a \in A$. By seesaw theorem [BL04, A.8], there is a line bundle $M \rightarrow A$ such that $s^* L \otimes p_1^* L^{-1} = p_2^* M$. Then $s^* L = p_1^* L \otimes p_2^* M$. Hence, $L = s^* L|_{0 \times A} = (p_1^* L \otimes p_2^* M)|_{0 \times A} = M$. Therefore, $s^* L = p_1^* L \otimes p_2^* L$. \square

Theorem F.5.2.4 is mentioned without proof in [KKN08, Sec. 1.2]. The analogue for abelian varieties is in [Wei49, no. 2].

Theorem F.5.2.4 (Weil). *If A is a complex torus, then $\pi : \text{Ext}(A, \mathbb{C}^*) \rightarrow \text{Pic}^0(A)$ is an isomorphism.*

Proof. For $B = \mathbb{C}^*$, the sheaf $\mathcal{B}_A = O_A^*$ and $H^1(A, \mathcal{B}_A) = \text{Pic}(A)$. The class of a line bundle $L \rightarrow A$ is primitive means the line bundle $s_A^* L$ is isomorphic to $p_1^* L \otimes p_2^* L$ on $A \times A$. By Proposition F.5.2.3 and Fact F.5.2.2, it is equivalent to $[L] \in \text{Pic}^0(A)$. Then Proposition F.5.1.3 and Theorem F.5.1.6 complete the proof. \square

With the identifications provided by Theorem F.5.2.4 and Proposition F.4.1.4 3, [AK01, Remark 1.1.16] can be rephrased in a coordinate-free way as follows. It is a criterion telling whether a semi-torus is a toroidal group.

Fact F.5.2.5. *Let $r \geq 1$ be an integer, and let $0 \rightarrow (\mathbb{C}^*)^r \rightarrow X \rightarrow A \rightarrow 0$ be an extension in \mathcal{C} . Denote by $(L_1, \dots, L_r) \in (A^\vee)^r$ the point corresponding to the equivalent class $[X] \in \text{Ext}(A, (\mathbb{C}^*)^r)$. Then the following are equivalent:*

- X is a toroidal group;

- for all $\sigma \in \mathbb{Z}^r \setminus \{0\}$, $\sum_{i=1}^r \sigma_i L_i \neq 0$ in A^\vee ;
- for every nontrivial morphism $f : (\mathbb{C}^*)^r \rightarrow \mathbb{C}^*$, the pushout extension $f_* X$ of A by \mathbb{C}^* is nontrivial.

F.5.3 The case $B = \mathbb{C}$

When $B = \mathbb{C}$, the sheaf $\mathcal{B}_A = \mathcal{O}_A$. Fact F.5.3.1 can be found in, e.g., [Men20, (3.1)].

Fact F.5.3.1 (Künneth formula). *Let X, Y be connected complex manifolds. Assume that Y is compact. Then there is a canonical decomposition $H^1(X \times Y, \mathcal{O}_{X \times Y}) = H^1(X, \mathcal{O}_X) \oplus H^1(Y, \mathcal{O}_Y)$.*

The analogue of Theorem F.5.3.2 for abelian varieties is [Ros58, Theorem 1].

Theorem F.5.3.2 (Rosenlicht, Serre). *If A is a complex torus, then the canonical morphism $\pi : \text{Ext}(A, \mathbb{C}) \rightarrow H^1(A, \mathcal{O}_A)$ is a \mathbb{C} -linear isomorphism. In particular, $\dim_{\mathbb{C}} \text{Ext}(A, \mathbb{C}) = \dim A$.*

Proof. Let m_1 (resp. m_2) be the injection $A \rightarrow A \times A$ defined by $a \mapsto (a, 0)$ (resp. $a \mapsto (0, a)$). Let $p_i : A \times A \rightarrow A$ ($i = 1, 2$) be the two projections. By Fact F.5.3.1, p_1^* and p_2^* identify $T(A \times A)$ as the direct sum $T(A) \oplus T(A)$. The projection to i th factor is m_i^* . Because $s_A \circ m_i = \text{Id}_A$, one has $s_A^*(x) = p_1^*x + p_2^*x$ for every $x \in T(A)$, i.e., x is primitive. Then Proposition F.5.1.3 and Theorem F.5.1.6 conclude the proof. \square

Remark F.5.3.3. Another way to prove Theorem F.5.3.2 is to use (F.13). In this case, the diagram (F.12) can be completed into a commutative diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & \text{Hom}_{\mathbb{C}}(\tilde{A}, \mathbb{C}) & \longrightarrow & \text{Hom}(\pi_1(A), \mathbb{C}) & \longrightarrow & \text{Ext}(A, \mathbb{C}) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \pi \\
0 & \longrightarrow & H^0(A, \Omega_A^1) & \longrightarrow & H^1(A, \mathbb{C}) & \longrightarrow & H^1(A, \mathcal{O}_A) \longrightarrow 0.
\end{array}
\tag{F.19}$$

The bottom row comes from the Hodge structure on $H^1(A, \mathbb{C})$ ([Huy05, Lem. 3.3.1]).

Corollary F.5.3.4. *Let A be a complex abelian variety, and let $n(\geq 0)$ be an integer. Then the natural morphism $\text{Ext}_{\text{Alg}}(A, \mathbb{G}_a^n) \rightarrow \text{Ext}(A^{\text{an}}, \mathbb{C}^n)$ is an isomorphism.*

Proof. It is a combination of [Ser88, Thm. 7, p.185], Theorem F.5.3.2 and [Ser56, Thm. 1]. \square

F.5.4 Universal vectorial extension

Definition F.5.4.1. [Ros58, p.705] Let H be a vector group. An extension

$$0 \rightarrow H \rightarrow G \rightarrow A \rightarrow 0 \quad (\text{F.20})$$

in \mathcal{C} is called decomposable if there exists an extension

$$0 \rightarrow H_1 \rightarrow G_1 \rightarrow A \rightarrow 0$$

in \mathcal{C} of A by a vector subgroup H_1 of H , and H' is a vector subgroup of H of positive dimension with an isomorphism $f : G_1 \oplus H' \rightarrow G$ such that the maps $H_1 \rightarrow H \rightarrow G$ and $H_1 \rightarrow G_1 \xrightarrow{f|_{G_1}} G$ coincide. Otherwise, the extension G is called indecomposable.

Proposition F.5.4.2. *The extension (F.20) is decomposable if and only if there is a strict vector subgroup H_1 of H and an extension $0 \rightarrow H_1 \rightarrow G_1 \xrightarrow{p_1} A \rightarrow 0$ with $\iota_* G_1 = G$, where $\iota : H_1 \rightarrow H$ is the inclusion.*

Proof. If G is decomposable, by definition, we can write $G = G_1 \oplus H'$, where $H' \subset H$ is a positive-dimensional vector subgroup and $0 \rightarrow H_1 \rightarrow G_1 \rightarrow A \rightarrow 0$ is an extension in \mathcal{C} of A by a vector subgroup $H_1 \subset H$ making a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H_1 & \longrightarrow & G_1 & \longrightarrow & A & \longrightarrow & 0 \\ & & \downarrow \iota & & \downarrow & & \downarrow \text{Id} & & \\ 0 & \longrightarrow & H & \longrightarrow & G & \longrightarrow & A & \longrightarrow & 0 \end{array}$$

By the universal property (F.7), $G = \iota_* G_1$. Moreover,

$$\dim H_1 = \dim G_1 - \dim A = \dim G - \dim H' - \dim A = \dim H - \dim H' < \dim H.$$

Conversely, assume that $\iota_* G_1 = G$. Choose a vector subspace H' of H with $H = H' \oplus H_1$, then $\dim H' = \dim H - \dim H_1 > 0$. The composed morphism $G_1 \oplus H' \xrightarrow{p_1} G_1 \xrightarrow{p_1} A$ is surjective of kernel $H_1 \oplus H' = H$, hence a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H_1 & \longrightarrow & G_1 & \longrightarrow & A & \longrightarrow & 0 \\ & & \downarrow \iota & & \downarrow & & \downarrow \text{Id} & & \\ 0 & \longrightarrow & H & \longrightarrow & G_1 \oplus H' & \longrightarrow & A & \longrightarrow & 0 \end{array}$$

with exact rows. By the universal property (F.7), $G = \iota_* G_1 = G_1 \oplus H'$. This identification makes the maps $H_1 \rightarrow H \rightarrow G$ and $H_1 \rightarrow G_1 \rightarrow G$ coincide. Therefore, G is decomposable. \square

Proposition F.5.4.3. *Let $0 \rightarrow \mathbb{C}^n \rightarrow G \rightarrow A \rightarrow 0$ be an extension in \mathcal{C} . Let $q_i : \mathbb{C}^n \rightarrow \mathbb{C}$ be the i -th coordinate function. Then G is indecomposable if and only if the family $\{q_{i,*}G\}_{1 \leq i \leq n}$ of vectors in $\text{Ext}(A, \mathbb{C})$ is linearly independent.*

Proof. Assume that $\{q_{i,*}G\}$ is linearly dependent. By changing of coordinate, one may assume that $q_{n,*}G = 0$ in $\text{Ext}(A, \mathbb{C})$. By Fact F.3.0.4, there is a morphism $r : q_{n,*}G \rightarrow \mathbb{C}$ with $i_n r = \text{Id}$ on $q_{n,*}G$.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{C}^n & \xrightarrow{i} & G & \longrightarrow & A \longrightarrow 0 \\ & & \downarrow q_n & \swarrow r & \downarrow \alpha & & \downarrow \text{Id} \\ 0 & \longrightarrow & \mathbb{C} & \xrightarrow{i_n} & q_{n,*}G & \longrightarrow & A \longrightarrow 0 \end{array}$$

Then $i_n r \alpha i = \alpha i = i_n q_n$. Since i_n is injective, one has

$$r \alpha i = q_n. \quad (\text{F.21})$$

Let $q : \mathbb{C}^n \rightarrow \mathbb{C}^{n-1}$ be the projection to the first $(n-1)$ coordinates. Let $\beta : G \rightarrow q_*G$ be the canonical morphism. Define a morphism

$$\epsilon : G \rightarrow q_*G \oplus \mathbb{C}, \quad g \mapsto (\beta(g), r\alpha(g)).$$

Then the right square of the following diagram is commutative.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{C}^n & \xrightarrow{i} & G & \longrightarrow & A \longrightarrow 0 \\ & & \downarrow q \oplus q_n & & \downarrow \epsilon & & \downarrow \text{Id} \\ 0 & \longrightarrow & \mathbb{C}^{n-1} \oplus \mathbb{C} & \longrightarrow & q_*G \oplus \mathbb{C} & \longrightarrow & A \longrightarrow 0 \end{array}$$

By (F.21), the left square of the above diagram is commutative. Therefore, ϵ is an equivalence of extensions and $G = q_*G \oplus \mathbb{C}$ is decomposable.

Conversely, assume that G is decomposable. By Proposition F.5.4.2, there is a vector subgroup $\iota : H_1 \rightarrow \mathbb{C}^n$ with $\dim H_1 < n$ and an extension $0 \rightarrow H_1 \rightarrow G_1 \rightarrow A \rightarrow 0$ with $\iota_*G_1 = G$. There is a linear combination $f = \sum_{i=1}^n a_i q_i : \mathbb{C}^n \rightarrow \mathbb{C}$, where $a_1, \dots, a_n \in \mathbb{C}$ are not all zero, such that $f\iota = 0$. Then $\sum_{i=1}^n a_i q_{i,*}G = f_*G = (f\iota)_*G_1 = 0$. Thus, the family $\{q_{i,*}G\}_i$ is linearly dependent. \square

Corollary F.5.4.4 follows from Proposition F.5.4.3 and Theorem F.5.3.2.

Corollary F.5.4.4. *Let $0 \rightarrow V \rightarrow G \rightarrow A \rightarrow 0$ be an extension in \mathcal{C} by a vector group V . If $\dim_{\mathbb{C}} V > g$, then G is decomposable.*

Proposition F.5.4.5 is an analytic analogue of [Ros58, Prop. 11].

Proposition F.5.4.5.

1. There is a \mathbb{C} -vector group H with $\dim_{\mathbb{C}} H = g$ and an indecomposable extension

$$0 \rightarrow H \rightarrow G \rightarrow A \rightarrow 0 \quad (\text{F.22})$$

such that for every $V \in \text{Vec}$, the map

$$\phi_V : \text{Hom}_{\text{Vec}}(H, V) \rightarrow \text{Ext}(A, V), \quad l \mapsto l_* G \quad (\text{F.23})$$

is a linear isomorphism. In other words, H together with the extension (F.22) represents the functor $\text{Ext}(A, \bullet) : \text{Vec} \rightarrow \text{Vec}$.

2. A $G' \in \text{Ext}(A, V)$ is indecomposable if and only if the corresponding linear map $\phi_V^{-1}(G') : H \rightarrow V$ is surjective.

Proof.

1. By Theorem F.5.3.2, $\dim_{\mathbb{C}} \text{Ext}(A, \mathbb{C}) = g$. Take a \mathbb{C} -basis $\{G_1, \dots, G_g\}$ of $\text{Ext}(A, \mathbb{C})$. By Fact F.4.1.8, $\text{Ext}(A, \mathbb{C}^g) = \bigoplus_{i=1}^g \text{Ext}(A, \mathbb{C})$, so there is an element $G \in \text{Ext}(A, \mathbb{C}^g)$ corresponding to $(G_1, \dots, G_g) \in \bigoplus_{i=1}^g \text{Ext}(A, \mathbb{C})$. Hence an extension $0 \rightarrow H \rightarrow G \rightarrow A \rightarrow 0$, where $H = \mathbb{C}^g$. By Proposition F.5.4.3, G is indecomposable.

When $l \in H^\vee$ is taking the i -th coordinate of $H = \mathbb{C}^g$, $l_* G = G_i$. Therefore, the image of the linear map $\phi_{\mathbb{C}}$ contains a basis of $\text{Ext}(A, \mathbb{C})$. Thus, $\phi_{\mathbb{C}}$ is surjective. Since $\dim_{\mathbb{C}} H^\vee = \dim_{\mathbb{C}} \text{Ext}(A, \mathbb{C})$, $\phi_{\mathbb{C}}$ is a linear isomorphism. Since every $V \in \text{Vec}$ is the direct sum of finitely many copies of \mathbb{C} and the formation of ϕ_V is functorial in V , ϕ_V is also a linear isomorphism.

2. By Proposition F.5.4.2, G' is decomposable iff there is a proper linear subspace $\iota : V_1 \rightarrow V$ with G' in the image of the map $\iota_* : \text{Ext}(A, V_1) \rightarrow \text{Ext}(A, V)$ iff there is a proper linear subspace $\iota : V_1 \rightarrow V$ with $\phi_V^{-1}(G')$ in the image of the map $\iota_* : \text{Hom}_{\text{Vec}}(H, V_1) \rightarrow \text{Hom}_{\text{Vec}}(H, V)$ iff $\phi_V^{-1}(G') : H \rightarrow V$ factors through a proper linear subspace $\iota : V_1 \rightarrow V$ iff $\phi_V^{-1}(G') : H \rightarrow V$ is not surjective.

□

The extension (F.22) is called the *universal vectorial extension* of A . (As a representing object, such an extension is unique up to equivalence.) By (F.23) and Theorem F.5.3.2, $H = H^0(A^\vee, \Omega_{A^\vee}^1)$.

EXAMPLE F.5.1.2 (CONTINUED). Since $\dim \text{Ext}(A, C) = 1$, this nontrivial extension is equivalent to the universal vectorial extension.

We proceed to give an explicit construction of the universal vectorial extension.

Proposition F.5.4.6. *Let $B^{\natural 1}$ be the group of isomorphic classes of rank 1 local systems on A . Let B^{\natural} be the group of isomorphic classes of pairs (L, ∇) , where $L \rightarrow A$ is a holomorphic line bundle and ∇ is a flat holomorphic connection on L . Then there exist natural identifications of groups*

$$B^{\natural} = B^{\natural 1} = \text{Hom}_{\text{Ab}}(\pi_1(A), \mathbb{C}^*) = H^1(A, \mathbb{C}^*) = \frac{H^1(A, \mathbb{C})}{H^1(A, \mathbb{Z})}.$$

They are isomorphic to $(\mathbb{C}^)^{2g}$.*

Proof. By the Riemann-Hilbert correspondence [Del70, Théorème 2.17, p.12], the map $B^{\natural} \rightarrow B^{\natural 1}$ defined by $(L, \nabla) \mapsto \ker(\nabla)$ is a group isomorphism. By [Del70, Corollaire 1.4, p.4], there is an isomorphism $B^{\natural 1} \rightarrow \text{Hom}_{\text{Ab}}(\pi_1(A), \mathbb{C}^*)$. By the universal coefficient theorem [Hat05, Thm. 3.2], there is a natural isomorphism $H^1(A, \mathbb{C}^*) \rightarrow \text{Hom}(\pi_1(A), \mathbb{C}^*)$. The exact sequences $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \xrightarrow{\exp(2\pi i \bullet)} \mathbb{C}^* \rightarrow 0$ of constant sheaves on A gives rise to an exact sequence

$$H^0(A, \mathbb{C}) \rightarrow H^0(A, \mathbb{C}^*) \rightarrow H^1(A, \mathbb{Z}) \rightarrow H^1(A, \mathbb{C}) \rightarrow H^1(A, \mathbb{C}^*) \rightarrow H^2(A, \mathbb{Z}) \rightarrow H^2(A, \mathbb{C}).$$

Since the first map is surjective and the last map is injective, it breaks into a short exact sequence

$$0 \rightarrow H^1(A, \mathbb{Z}) \rightarrow H^1(A, \mathbb{C}) \rightarrow H^1(A, \mathbb{C}^*) \rightarrow 0$$

and hence an isomorphism $H^1(A, \mathbb{C})/H^1(A, \mathbb{Z}) \rightarrow H^1(A, \mathbb{C}^*)$ functorial in A . Moreover, there is a non-canonical isomorphism $H^1(A, \mathbb{C}^*) \rightarrow (\mathbb{C}^*)^{2g}$. \square

For every $(L, \nabla) \in B^{\natural}$, the line bundle $L \in \text{Pic}^0(A) = A^{\vee}$ by [Dem12, Ch. V, §9]. The bottom row of (F.19) induces an exact sequence in \mathcal{C} :

$$0 \rightarrow H^0(A, \Omega_A^1) \rightarrow \frac{H^1(A, \mathbb{C})}{H^1(A, \mathbb{Z})} \rightarrow \frac{H^1(A, \mathcal{O}_A)}{H^1(A, \mathbb{Z})} \rightarrow 0. \quad (\text{F.24})$$

Using the identifications $B^{\natural} \cong \frac{H^1(A, \mathbb{C})}{H^1(A, \mathbb{Z})}$ from Proposition F.5.4.6 and $A^{\vee} = \text{Pic}^0(A) = H^1(A, \mathcal{O}_A)/H^1(A, \mathbb{Z})$, (F.24) is an extension of A^{\vee} by $H^0(A, \Omega_A^1)$ and gives a morphism $B^{\natural} \rightarrow \text{Pic}^0(A)$, which sends (L, ∇) to L . Hence a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}_{\mathcal{C}}(\tilde{A}, \mathbb{C}^*) & \longrightarrow & \text{Hom}_{\text{Ab}}(\pi_1(A), \mathbb{C}^*) & \longrightarrow & \text{Ext}(A, \mathbb{C}^*) \longrightarrow 0 \\ & & \downarrow & & \uparrow & & \downarrow \pi \\ 0 & \longrightarrow & H^0(A, \Omega_A^1) & \longrightarrow & B^{\natural} & \xrightarrow{u} & \text{Pic}^0(A) \longrightarrow 0, \end{array}$$

where the first exact row is (F.9) and the second comes from (F.24). The left vertical isomorphism uses Proposition F.4.1.4 2 and the isomorphism $L(A)^\vee \rightarrow H^0(A, \Omega_A^1)$ given by [BL04, Thm. 1.4.1 b)]. The middle vertical isomorphism is contained in Proposition F.5.4.6.

When A is an abelian variety, it is proved in [Mes73, p.260] that (F.24) is the universal vectorial extension of A^\vee . The proof is based on [Ros58, Thm. 1]. In a similar manner, Proposition F.5.4.7 follows from Theorem F.5.3.2.

Proposition F.5.4.7. *The extension (F.24) is the universal vectorial extension of $A^\vee = \text{Pic}^0(A)$. In particular, the extension group is isomorphic to $(\mathbb{C}^*)^{2g}$ (as a complex Lie group).*

Proof. Let $U = H^0(A, \Omega_A^1)$. Pushing out the extension (F.24) defines a natural transformation $\psi : \text{Hom}_{\text{Vec}}(U, \bullet) \rightarrow \text{Ext}(A^\vee, \bullet)$ between two functors on Vec .

We claim that $\psi_{\mathbb{C}}$ is an isomorphism. Choose $u \in \ker(\psi_{\mathbb{C}}) \subset \text{Hom}_{\text{Vec}}(U, \mathbb{C})$. As the push-out along u is trivial, by Fact F.3.0.4, there is a morphism $r : E \rightarrow \mathbb{C}$ with $ir = \text{Id}_E$. Let $u' : H^1(A, \mathbb{C}) \rightarrow \mathbb{C}$ be the morphism in \mathcal{C} induced by r . Then $u' = d_e u'$ is \mathbb{C} -linear. Now that $u'(H^1(A, \mathbb{Z})) = 0$ and $H^1(A, \mathbb{Z})$ contains a \mathbb{C} -basis of $H^1(A, \mathbb{C})$, one has $u' = 0$. As the diagram commutes, $u = 0$.

$$\begin{array}{ccccccc}
 & & & H^1(A, \mathbb{C}) & & & \\
 & & & \downarrow & & & \\
 0 & \longrightarrow & U & \xrightarrow{u'} & \frac{H^1(A, \mathbb{C})}{H^1(A, \mathbb{Z})} & \longrightarrow & A^\vee \longrightarrow 0 \\
 & & \downarrow u & \swarrow & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathbb{C} & \xrightarrow{i} & E & \longrightarrow & A^\vee \longrightarrow 0
 \end{array}$$

(Note: A dashed arrow labeled r points from E to \mathbb{C} , and a dashed arrow labeled u' points from $H^1(A, \mathbb{C})$ to \mathbb{C} .)

Therefore, $\psi_{\mathbb{C}}$ is injective. By Theorem F.5.3.2, $\dim_{\mathbb{C}} \text{Ext}(A^\vee, \mathbb{C}) = \dim_{\mathbb{C}} \text{Hom}_{\text{Vec}}(U, \mathbb{C})$. Therefore, $\psi_{\mathbb{C}}$ is a linear isomorphism. Similar to the proof of Proposition F.5.4.5 1, ψ is a natural isomorphism of the two functors. \square

Another construction of the universal vectorial extension is in [Nak94, Prop. 2.4]⁹.

Remark F.5.4.8. The real Lie group extension underlying (F.24) is trivial by Fact F.5.1.1. Indeed, consider the real analytic group morphism $A^\vee \rightarrow B^\natural$ defined by $L \mapsto (L, \nabla^L)$, where ∇^L is the unique flat Chern connection on L given by Lemma D.2.0.4 1. This map is a real Lie group section to (F.24), but not holomorphic.

⁹stated for complex abelian varieties but the proof extends to complex tori.

Remark F.5.4.9. Let A be a complex abelian variety of dimension g . By Corollary F.5.3.4, the extension (F.22) is equivalent to an algebraic one. Thus, the analytification of the algebraic universal vectorial extension $0 \rightarrow \mathbb{G}_a^g \rightarrow E \rightarrow A \rightarrow 0$ is exactly the analytic universal vectorial extension. From [Bri09, Prop. 2.3 (i)] and the footnote in [MRM74, p.34], the algebraic variety E is anti-affine, i.e., every morphism $E \rightarrow A_{\mathbb{C}}^1$ of algebraic varieties is constant. On the other hand, by Proposition F.5.4.7, E^{an} is isomorphic to $(\mathbb{C}^*)^{2g}$ as a complex Lie group, so E^{an} is not a toroidal group. Although E is not an affine variety, E^{an} is a Stein manifold. See also Serre's example [Har70, Exampe 3.2, p.232].

Remark F.5.4.10. Universal vectorial extensions can be defined for not only complex tori but also toroidal groups. Consider a toroidal group X of dimension n . Similar to Proposition F.5.4.5, the functor $\text{Ext}(X, \cdot) : \text{Vec} \rightarrow \text{Vec}$ is represented by $\text{Ext}(X, \mathbb{C})^{\vee}$, which is the kernel of the natural linear map $H_1(X, \mathbb{C}) \rightarrow H^0(X, \Omega_X^1)^{\vee}$ by (F.13).

An extrinsic description is possible. Choose a presentation

$$0 \rightarrow (\mathbb{C}^*)^{n-q} \rightarrow X \rightarrow T \rightarrow 0 \quad (\text{F.25})$$

according to [AK01, 1.1.14], where T is a complex torus of dimension q . For every $V \in \text{Vec}$, by Proposition F.4.2.1, the induced sequence

$$\text{Hom}_{\mathbb{C}}((\mathbb{C}^*)^{n-q}, V) \rightarrow \text{Ext}(T, V) \rightarrow \text{Ext}(X, V) \rightarrow \text{Ext}((\mathbb{C}^*)^{n-q}, V)$$

is exact in Vec . By Proposition F.4.1.4 1, $\text{Hom}_{\mathbb{C}}((\mathbb{C}^*)^{n-q}, V) = 0$. By Proposition F.3.2.3, $\text{Ext}((\mathbb{C}^*)^{n-q}, V) = 0$. Thus, the morphism $\text{Ext}(T, V) \rightarrow \text{Ext}(X, V)$ is a \mathbb{C} -linear isomorphism. In other words, the natural transformation $\text{Ext}(T, \cdot) \rightarrow \text{Ext}(X, \cdot)$ between the two functors on Vec is an isomorphism. In this way, the case of toroidal groups is reduced to the case of complex tori.

F.5.5 Application to the functor $\text{Ext}(A, \bullet)$

Analogue of Proposition F.5.5.1 for abelian varieties is [Ros58, Cor., p.711].

Proposition F.5.5.1. *If B is a complex Lie subgroup (not necessarily connected) of A , then there is a natural exact sequence in Ab :*

$$0 \rightarrow \text{Ext}(A/B, \mathbb{C}) \rightarrow \text{Ext}(A, \mathbb{C}) \rightarrow \text{Ext}(B, \mathbb{C}) \rightarrow 0.$$

Proof. By Corollary F.4.1.13, there is an isomorphism $B \rightarrow B_0 \times B/B_0$ in \mathcal{C} and $\text{Ext}(B/B_0, \mathbb{C}) = 0$. By Fact F.4.1.8, $\text{Ext}(B, \mathbb{C}) = \text{Ext}(B_0, \mathbb{C})$. Since B is compact and B_0 is open in B , the quotient B/B_0 is finite, thus $\text{Hom}_{\text{Ab}}(B/B_0, \mathbb{C}) = 0$. By the compactness of B_0 , $\text{Hom}_{\mathbb{C}}(B_0, \mathbb{C}) = 0$. Then $\text{Hom}(B, \mathbb{C}) = 0$. Now that $A, B_0, A/B$ are complex tori, Theorem F.5.3.2 implies $\dim \text{Ext}(A, \mathbb{C}) = \dim \text{Ext}(A/B, \mathbb{C}) + \dim \text{Ext}(B, \mathbb{C})$. This together with Proposition F.4.2.1 proves the stated exactness. \square

The proof of Theorem F.5.5.2 is shorter than that of its algebraic analogue [Ser88, Thm. 12, p.195].

Theorem F.5.5.2. *If $0 \rightarrow B' \rightarrow B \xrightarrow{\phi} B'' \rightarrow 0$ is an exact sequence in \mathcal{C} , then the sequence¹⁰ in \mathbf{Ab}*

$$\mathrm{Ext}(A, B') \rightarrow \mathrm{Ext}(A, B) \xrightarrow{\phi_*} \mathrm{Ext}(A, B'') \rightarrow 0 \quad (\text{F.26})$$

is exact. If B''_0 is linear, then the first map in (F.26) is injective.

Proof. By Proposition F.4.2.3, it suffices to prove that $\phi_* : \mathrm{Ext}(A, B) \rightarrow \mathrm{Ext}(A, B'')$ is surjective. From (F.10) and Proposition F.4.2.1, one obtains a commutative square

$$\begin{array}{ccc} \mathrm{Hom}(\pi_1(A), B) & \longrightarrow & \mathrm{Hom}(\pi_1(A), B'') \\ \downarrow & & \downarrow \\ \mathrm{Ext}(A, B) & \xrightarrow{\phi_*} & \mathrm{Ext}(A, B''), \end{array}$$

where the vertical maps are surjective. Since $\pi_1(A)$ is a free \mathbb{Z} -module, the top row is surjective, then so is the bottom.

Now assume that B''_0 is linear, then $\mathrm{Hom}_{\mathcal{C}}(A, B'') = 0$. By Proposition F.4.2.3, the first map is injective. \square

Remark F.5.5.3. The linearity of B''_0 is necessary to guarantee the injectivity in Theorem F.5.5.2. For instance, let $0 \rightarrow \mathbb{C}^g \rightarrow (\mathbb{C}^*)^{2g} \rightarrow A \rightarrow 0$ be the universal vectorial extension of A and assume $g \geq 1$. By Proposition F.4.2.3, the natural sequence $0 \rightarrow \mathrm{Hom}_{\mathcal{C}}(A, A) \rightarrow \mathrm{Ext}(A, \mathbb{C}^g) \rightarrow \mathrm{Ext}(A, (\mathbb{C}^*)^{2g})$ is exact. Thus, Id_A is a nonzero element in the kernel of the first map of (F.26).

Example F.5.5.4. Applying Theorem F.5.5.2 to the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \xrightarrow{\exp(2\pi i \bullet)} \mathbb{C}^* \rightarrow 1$, and using Fact F.3.2.6, Theorems F.5.2.4 and F.5.3.2, one gets an exact sequence

$$0 \rightarrow \mathrm{Hom}(\pi_1(A), \mathbb{Z}) \rightarrow H^1(A, \mathcal{O}_A) \rightarrow \mathrm{Pic}^0(A) \rightarrow 0. \quad (\text{F.27})$$

In particular, $\mathrm{Ext}(A, \cdot)$ turns the exponential map to the universal cover of the complex torus A^\vee . Identifying $\mathrm{Hom}(\pi_1(A), \mathbb{Z})$ with the sheaf cohomology $H^1(A, \mathbb{Z})$, the sequence (F.27) is also induced by the exponential sequence of sheaves on A :

$$0 \rightarrow \mathbb{Z}_A \rightarrow \mathcal{O}_A \xrightarrow{\exp(2\pi i)} \mathcal{O}_A^* \rightarrow 1.$$

Theorem F.5.5.5 is an analytic version of [Ser88, Thm. 13, p.196]

¹⁰induced by Proposition F.4.2.3

Theorem F.5.5.5. *If $0 \rightarrow L \xrightarrow{i} C \rightarrow A \rightarrow 0$ is an exact sequence in \mathcal{C} with L connected and $G \in \text{Ab}_c$. Then there is a natural exact sequence*

$$0 \rightarrow \text{Ext}(A, G) \rightarrow \text{Ext}(C, G) \xrightarrow{i^*} \text{Ext}(L, G) \rightarrow 0.$$

Proof. As L is connected and G is discrete, $\text{Hom}_{\mathcal{C}}(L, G) = 0$. By Proposition F.4.2.1, it suffices to show that $i^* : \text{Ext}(C, G) \rightarrow \text{Ext}(L, G)$ is surjective. For every $L' \in \text{Ext}(L, G)$, by Theorem F.5.5.2, the map $\text{Ext}(A, L') \rightarrow \text{Ext}(A, L)$ is surjective. Thus, there exists $C' \in \text{Ext}(A, L')$ having image $C \in \text{Ext}(A, L)$.

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & G & \xrightarrow{\beta} & \ker(\alpha) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & L' & \longrightarrow & C' & \longrightarrow & A \longrightarrow 0 \\
& & \downarrow & & \downarrow \alpha & & \downarrow \\
0 & \longrightarrow & L & \xrightarrow{i} & C & \longrightarrow & A \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & \longrightarrow & 0 & \longrightarrow & 0
\end{array}$$

(Note: A dashed arrow points from C' to C in the original diagram.)

By the snake lemma, α is surjective and β is an isomorphism. Therefore, $C' \in \text{Ext}(C, G)$ and $i^*C' = L'$ in $\text{Ext}(L, G)$. \square

In Example F.5.5.6, we give another proof of [BL99, Prop. 5.7, p.21], which computes the extension group of two complex tori.

Example F.5.5.6. Let $X_i = \mathbb{C}^{g_i} / \Pi_i \mathbb{Z}^{2g_i}$ ($i = 1, 2$) be two complex tori, where the chosen period matrix is of the form $\Pi_i = (\tau_i, I_{g_i})$ with $\tau_i \in M_{g_i}(\mathbb{C})$ and $\det(\text{Im}(\tau_i)) \neq 0$. Define $\xi : M(2g_1 \times 2g_2, \mathbb{Z}) \rightarrow M(g_1 \times g_2, \mathbb{C})$ by $\xi(P) = \Pi_1 P \begin{pmatrix} I_{g_2} \\ \tau_2 \end{pmatrix}$.

Define a map $\rho : M(g_1 \times g_2, \mathbb{C}) \rightarrow \text{Ext}(X_2, \tilde{X}_1)$ as follows. For every $\alpha \in M(g_1 \times g_2, \mathbb{C})$, let $\alpha' = (\alpha, 0) \in M(g_1 \times 2g_2, \mathbb{C})$. Consider the sequence

$$0 \rightarrow \mathbb{C}^{g_1} \xrightarrow{i} \frac{\mathbb{C}^{g_1+g_2}}{\{(\alpha'v, \Pi_2 v) : v \in \mathbb{Z}^{2g_2}\}} \xrightarrow{p} X_2 \rightarrow 0,$$

where i is induced by $\mathbb{C}^{g_1} \rightarrow \mathbb{C}^{g_1+g_2}$ defined by $x \mapsto (x, 0)$ and p is induced by the second projection $\mathbb{C}^{g_1+g_2} \rightarrow \mathbb{C}^{g_2}$. It is an exact sequence. Denote its class by $\rho(M) \in \text{Ext}(X_2, \tilde{X}_1)$. This sequence fits into a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \tilde{X}_1 & \longrightarrow & \frac{\mathbb{C}^{g_1+g_2}}{\{(\alpha'v, \Pi_2v) : v \in \mathbb{Z}^{2g_2}\}} & \longrightarrow & X_2 \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & X_1 & \longrightarrow & X & \longrightarrow & X_2 \longrightarrow 0,
\end{array}$$

where the second row is $\psi_{\Pi_1, \Pi_2}(\alpha') \in \text{Ext}(X_2, X_1)$ defined in [BL99, p.20], and

$$X = \frac{\mathbb{C}^{g_1+g_2}}{\{(\Pi_1u + \alpha'v, \Pi_2v) : u \in \mathbb{Z}^{2g_1}, v \in \mathbb{Z}^{2g_2}\}}.$$

Then ρ is a linear isomorphism by Theorem F.5.3.2.

Define a map $\phi : M(2g_1 \times 2g_2, \mathbb{Z}) \rightarrow \text{Ext}(X_2, \pi_1(X_1))$ as follows. Given $P = \begin{pmatrix} P_1 & P_2 \\ P_3 & P_4 \end{pmatrix} \in M(2g_1 \times 2g_2, \mathbb{Z})$, with each $P_i \in M(g_1 \times g_2, \mathbb{Z})$, we set $A = \tau_1 P_2 + P_4 \in M(g_1 \times g_2, \mathbb{C})$ and $\alpha = \xi(P)$. The linear map $\mathbb{C}^{g_1+g_2} \xrightarrow{(I, -A)} \mathbb{C}^{g_1}$ sends $(u, 0)$ to u for all $u \in \mathbb{C}^{g_1}$ and sends $(\alpha'v, \Pi_2v)$ to $\Pi_1 \begin{pmatrix} P_1 & -P_2 \\ P_3 & -P_4 \end{pmatrix} v \in \Pi_1 \mathbb{Z}^{2g_1}$ for all $v \in \mathbb{Z}^{2g_2}$. Thus it descends to the vertical morphism in the middle of the following commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathbb{C}^{g_1} & \longrightarrow & \frac{\mathbb{C}^{g_1+g_2}}{\{(\alpha'v, \Pi_2v) : v \in \mathbb{Z}^{2g_2}\}} & \longrightarrow & X_2 \longrightarrow 0 \\
& & \downarrow & & \downarrow (I_{g_1}, -A) & & \downarrow \\
0 & \longrightarrow & X_1 & \longrightarrow & X_1 & \longrightarrow & 0 \longrightarrow 0,
\end{array} \tag{F.28}$$

where the first row is of class $\rho(\alpha) = \rho(\xi(P))$. The snake lemma gives an extension of X_2 by $\pi_1(X_1)$, whose class is denoted by $\phi(P)$.

The image of $\phi(P)$ under the pushout map $\text{Ext}(X_2, \pi_1(X_1)) \rightarrow \text{Ext}(X_2, \tilde{X}_1)$ is exactly the first row of (F.28), i.e., $\rho(\xi(P))$. Then ϕ is a group isomorphism by Fact F.3.2.6. And there is a commutative diagram

$$\begin{array}{ccccccc}
& & & & & & M(g_1 \times 2g_2, \mathbb{C}) \\
& & & & & \nearrow \alpha \mapsto \alpha' & \\
M(2g_1 \times 2g_2, \mathbb{Z}) & \xrightarrow{\xi} & M(g_1 \times g_2, \mathbb{C}) & \longrightarrow & \frac{M(g_1 \times g_2, \mathbb{C})}{\text{Im}(\xi)} & & \\
\downarrow \phi & & \downarrow \rho & & \downarrow \text{dotted} & \nwarrow \psi_{\Pi_1, \Pi_2} & \\
\text{Ext}(X_2, \pi_1(X_1)) & \longrightarrow & \text{Ext}(X_2, \tilde{X}_1) & \longrightarrow & \text{Ext}(X_2, X_1) & \longrightarrow & 0
\end{array}$$

where the second row is from (F.14) and the induced dotted isomorphism is exactly the content of [BL99, Proposition 5.7, p.21].

To conclude Section F.5.5, we show that the groups of commutative extensions of complex tori by linear groups are naturally complex Lie

groups. Let \mathcal{T} (resp. \mathcal{S}) be the full subcategory of \mathcal{C} comprised of complex tori (resp. objects whose identity component is linear). Then $\text{Ext} : \mathcal{T}^{\text{op}} \times \mathcal{S} \rightarrow \text{Ab}$ is an additive functor by Fact F.4.1.8. Theorem F.5.5.7, an analytic analogue of [Wu86, Theorem 5], lifts this functor.

Theorem F.5.5.7 (Wu). *There is a natural way to lift $\text{Ext} : \mathcal{T}^{\text{op}} \times \mathcal{S} \rightarrow \text{Ab}$ to an additive functor $\text{Ext} : \mathcal{T}^{\text{op}} \times \mathcal{S} \rightarrow \mathcal{C}$.*

Proof. First we define a complex Lie group structure on $\text{Ext}(A, H)$, where $A \in \mathcal{T}$ and $H \in \mathcal{S}$. Let $g = \dim A$.

If there is an isomorphism $f : H \rightarrow (\mathbb{C}^*)^n$ in \mathcal{S} , then by Theorem F.5.2.4, f gives rise to an isomorphism $\text{Ext}(A, H) \rightarrow (A^\vee)^n$ making $\text{Ext}(A, H)$ a complex torus. The complex structure on $\text{Ext}(A, H)$ is independent of the choice of the isomorphism f .

If H is connected, by Proposition F.2.0.7, there is an isomorphism $u : H \rightarrow V \times H_m$, where $V \in \text{Vec}$ and H_m is a power of \mathbb{C}^* . Then $u_* : \text{Ext}(A, H) \rightarrow \text{Ext}(A, V) \times \text{Ext}(A, H_m)$ is an isomorphism. By Theorem F.5.3.2, the vector space $\text{Ext}(A, V)$ is *finite dimensional*. Together with last paragraph, $\text{Ext}(A, H)$ inherits a complex Lie group structure, which is independent of the choice of u .

For a general object $H \in \mathcal{S}$, the natural exact sequence $0 \rightarrow H_0 \rightarrow H \rightarrow H/H_0 \rightarrow 0$ in \mathcal{C} is trivial by Corollary F.4.1.13. Thus, the resulting exact sequence $0 \rightarrow \text{Ext}(A, H_0) \rightarrow \text{Ext}(A, H) \rightarrow \text{Ext}(A, H/H_0) \rightarrow 0$ in Ab is also trivial. Now that $\text{Ext}(A, H/H_0) = \text{Hom}_{\text{Ab}}(\pi_1(A), H/H_0)$ by Fact F.3.2.6, one regards it as a discrete group. From the complex structure on $\text{Ext}(A, H_0)$, the group $\text{Ext}(A, H)$ has a unique complex Lie group structure, such that the identity component is $\text{Ext}(A, H_0)$.

It remains to show:

1. If $A \in \mathcal{T}$ is fixed, then $\text{Ext}(A, \cdot)$ sends morphisms in \mathcal{S} to morphisms in \mathcal{C} .
2. If $H \in \mathcal{S}$ is fixed, then $\text{Ext}(\cdot, H)$ sends morphisms in \mathcal{T} to morphisms in \mathcal{C} .

To show 1, let $h : H \rightarrow H'$ be a morphism in \mathcal{S} . By decomposing H, H' according to Corollary F.4.1.13 and Proposition F.2.0.7, one may assume that each of H and H' is either discrete, \mathbb{C} or \mathbb{C}^* .

- If H is discrete, then so is $\text{Ext}(A, H)$, hence $\text{Ext}(A, h)$ is a morphism in \mathcal{C} .
- If $H = H' = \mathbb{C}$, by Proposition F.4.1.4 2, h is a linear map. By Corollary F.4.1.9, so is $\text{Ext}(A, h)$.
- If $H = \mathbb{C}, H' = \mathbb{C}^*$. By Proposition F.4.1.4 2, h is the composition of a linear map $\mathbb{C} \rightarrow \mathbb{C}$ followed by the exponential map $\exp(2\pi i \cdot) :$

$\mathbb{C} \rightarrow \mathbb{C}^*$. By Example F.5.5.4, $\text{Ext}(A, h)$ is the composition of a linear map $H^1(A, O_A) \rightarrow H^1(A, O_A)$ followed by the universal cover $H^1(A, O_A) \rightarrow A^\vee$. Thus, $\text{Ext}(A, h)$ is a morphism in \mathcal{C} .

- If H' is discrete and H is connected, then h is trivial and so is $\text{Ext}(A, h)$.
- If $H = \mathbb{C}^*$ and $H' = \mathbb{C}$, then h is trivial by Proposition F.4.1.4 1 and so is $\text{Ext}(A, h)$.
- If $H = H' = \mathbb{C}^*$, then h is a power map by Proposition F.4.1.4 3. Then $\text{Ext}(A, h)$ is a power map of A^\vee , hence a morphism in \mathcal{C} .

This proves 1.

To show 2, let $g : A \rightarrow A'$ be a morphism in \mathcal{T} . By decomposing H again, we may divide the proof into three cases.

- $H = \mathbb{C}^*$. By pulling back line bundles, g induces the dual morphism $g^* : \text{Pic}^0(A') \rightarrow \text{Pic}^0(A)$. It is identified with $\text{Ext}(g, H)$ by Fact F.4.3.2 and Theorem F.5.2.4.
- H is discrete. Then so is $\text{Ext}(A', H)$ and thus $\text{Ext}(g, H)$ is a morphism in \mathcal{C} .
- $H = \mathbb{C}$. By pulling back, g induces a \mathbb{C} -linear map $H^1(A', O_{A'}) \rightarrow H^1(A, O_A)$. It is identified with $\text{Ext}(g, H)$ by Fact F.4.3.2 and Theorem F.5.3.2.

This proves 2. □

Remark F.5.5.8. In Theorem F.5.5.7, we cannot generalize from complex tori to toroidal groups, nor remove the linear restriction.

Let X be a toroidal group. Then $\text{Hom}_{\mathcal{C}}(X, \mathbb{C}^*) = 0$, hence (F.14) specializes to

$$0 \rightarrow \text{Ext}(X, \mathbb{Z}) \xrightarrow{i} \text{Ext}(X, \mathbb{C}) \rightarrow \text{Ext}(X, \mathbb{C}^*) \rightarrow 0. \quad (\text{F.29})$$

Note that $\text{Ext}(X, \mathbb{Z}) = H^1(X, \mathbb{Z})$ (Fact F.3.2.6), and by (F.13) the injection i is the composition of the inclusion $H^1(X, \mathbb{Z}) \rightarrow H^1(X, \mathbb{C})$ with the projection $H^1(X, \mathbb{C}) \rightarrow \frac{H^1(X, \mathbb{C})}{H^0(X, \Omega_X^1)}$.

When X is compact, the sequence (F.29) lifts to an exact sequence in \mathcal{C} by Theorem F.5.5.7. As opposed to the compact case, when X is not compact and consider the presentation (F.25), one has $1 \leq q < n$, so

$$\text{rank}_{\mathbb{Z}} \text{Ext}(X, \mathbb{Z}) = n + q > 2q = \dim_{\mathbb{R}} \text{Ext}(X, \mathbb{C}).$$

Therefore, the image of i is not closed in the vector space $\text{Ext}(X, \mathbb{C})$ (a phenomenon seen in Example F.4.1.2). In particular, the sequence (F.29) has no lift to an exact sequence in \mathcal{C} .

Let A, B be two complex tori, $g = \dim A$, $g' = \dim B$ and reconsider (F.14):

$$0 \rightarrow \operatorname{Hom}_{\mathcal{C}}(A, B) \xrightarrow{j} \operatorname{Ext}(A, \pi_1(B)) \rightarrow \operatorname{Ext}(A, \tilde{B}) \rightarrow \operatorname{Ext}(A, B) \rightarrow 0.$$

Here, $\operatorname{Ext}(A, \tilde{B})$ is a \mathbb{C} -vector space of dimension gg' by Theorem F.5.3.2. Identifying $\operatorname{Ext}(A, \pi_1(B))$ with $\operatorname{Hom}_{\operatorname{Ab}}(\pi_1(A), \pi_1(B))$ via Fact F.3.2.6, j is the map ρ_r in [BL04, p.10]. The quotient $\frac{\operatorname{Ext}(A, \pi_1(B))}{\operatorname{Hom}_{\mathcal{C}}(A, B)}$ is a free abelian group of rank $4gg' - \operatorname{rank}_{\mathbb{Z}} \operatorname{Hom}_{\mathcal{C}}(A, B)$. As long as $\operatorname{rank}_{\mathbb{Z}} \operatorname{Hom}_{\mathcal{C}}(A, B) < 2gg'$ (say, when $A = B$ is an elliptic curve without complex multiplication, then $\mathbb{Z} = \operatorname{Hom}_{\mathcal{C}}(A, B)$), the image of the induced injection $\frac{\operatorname{Ext}(A, \pi_1(B))}{\operatorname{Hom}_{\mathcal{C}}(A, B)} \rightarrow \operatorname{Ext}(A, \tilde{B})$ is not closed. In particular, $\operatorname{Ext}(A, B)$ has no structure of complex Lie group making this sequence exact in \mathcal{C} .

F.6 Extensions of complex tori are often commutative

In Section F.6, we prove that under suitable hypotheses, an extension of a complex torus is commutative.

Proposition F.6.0.1. *If $1 \rightarrow B \rightarrow C \xrightarrow{p} A \rightarrow 1$ is a central extension of complex Lie groups, where A is a toroidal group, then C is commutative. Or equivalently, for every $B \in \mathcal{C}$, the natural injection $\operatorname{Ext}(A, B) \rightarrow \operatorname{Ext}(A, B, 1)$ is an isomorphism.*

Proof. Consider the holomorphic map $A \times A \rightarrow B$ given by (F.4). By [NW13, Thm. 5.1.36], it is a group morphism, so constant. Thus, C is commutative. \square

An algebraic analogue of Proposition F.6.0.2 is [Wu86, Cor. 2, p.370].

Proposition F.6.0.2. *Let $1 \rightarrow K \rightarrow E \rightarrow A \rightarrow 1$ be an extension of complex Lie groups, where A is a complex torus.*

1. *If $Z(K)_0$ is Stein, then $Z(K) = Z(E) \cap K$.*
2. *If K is commutative and K_0 is Stein, then E is commutative.*

Proof.

1. Since $Z(E) \cap K \subset Z(K)$, it suffices to prove that $Z(K) \subset Z(E)$. Consider the group morphism (F.5): $\theta : A \rightarrow \operatorname{Aut}(Z(K))$. For every $x \in Z(K)$, the map

$$\phi : A \rightarrow Z(K), \quad a \mapsto \theta_a(x)x^{-1}$$

is continuous. Moreover, $\phi(0) = e_K$. By the connectedness of A , $\phi(A) \subset Z(K)_0$. As $Z(K)_0$ is Stein and A is compact, $\phi(A)$ is the singleton $\{e_K\}$. Therefore, $\theta_a(x) = x$ for every $x \in Z(K)$, which proves $Z(K) \subset Z(E)$.

2. By 1, $K \subset Z(E)$. By Proposition F.6.0.1, E is commutative. □

In Proposition F.6.0.3, when B is isomorphic to \mathbb{C}^n for some integer $n \geq 0$ or to \mathbb{C}^* , we recover [BZ23a, Lem. 2.10].

Proposition F.6.0.3. *Let $1 \rightarrow B \rightarrow C \xrightarrow{p} A \rightarrow 1$ be an exact sequence of complex Lie groups, where A is a complex torus and B is commutative. If the group B/B_0 is torsion (i.e., every element of B/B_0 has finite order), then C is commutative.*

Proof. Let Z be the center of C . By Proposition F.6.0.1, it suffices to check $B \subset Z$.

The outer action induces a morphism $A \rightarrow \text{Aut}(B_0)(\leq \text{GL}(L(B)))$. It is trivial by the compactness of A , i.e., $B_0 \leq Z$. By Corollary F.4.1.13, one may assume $B = B_0 \times D$, where D is a discrete subgroup of B isomorphic to B/B_0 and $D \cap B_0 = \{e_B\}$. Let $q : B \rightarrow D$ and $r : B \rightarrow B_0$ be the corresponding projections.

It remains to show that $0 \times D(\leq B)$ is contained in Z . Fix $d \in D$ and put $b = (0, d) \in B$. The map

$$\nu : C \rightarrow C, \quad c \mapsto cb c^{-1}$$

is holomorphic and $\nu(e) = b$. For every $b' \in B$, one has

$$\nu(cb') = cb'bb'^{-1}c^{-1} = cb c^{-1} = \nu(c).$$

The right multiplication action of B on the complex manifold C has quotient A by Fact F.2.0.3, so ν factors through a morphism $u : A \rightarrow B$ of complex manifolds. Then $qu : A \rightarrow D$ is continuous. Since A is connected, qu is constant. Since $qu(e_A) = d$, one gets $qu \equiv d$.

On the other hand, the map $ru : A \rightarrow B_0$ is holomorphic. By assumption, there is an integer $n \geq 1$ (depending on d) such that $d^n = e_D$ in D . Thus, $b^n = e_B$. For every $c \in C$, one has $\nu(c)^n = (cb c^{-1})^n = cb^n c^{-1} = e_B$. Therefore, $ru(A)$ is contained in the torsion subgroup $B_{0,\text{tor}}$ of B_0 . In view of [AK01, Prop. 1.1.2], $B_{0,\text{tor}}$ is totally disconnected. Since A is connected, ru is constant.

Since $ru(e_A) = 0$, one has $ru \equiv 0$. Therefore, $u \equiv b$, i.e., $b \in Z$. Therefore, $0 \times D \subset Z$ and the proof is completed. □

Corollary F.6.0.4 follows immediately from Proposition F.6.0.3.

Corollary F.6.0.4. *Given an extension*

$$0 \rightarrow (\mathbb{C}^*)^n \rightarrow G \rightarrow A \rightarrow 0 \tag{F.30}$$

of complex Lie groups, where A is a complex torus and $n(\geq 1)$ is an integer, then G is a semi-torus.

Corollary F.6.0.5. *In Corollary F.6.0.4, if A is algebraic, then G admits a unique structure of semiabelian variety such that (F.30) defines a commutative extension of algebraic groups.*

Proof. From Corollary F.6.0.4, (F.30) defines an element of $\text{Ext}(A^{\text{an}}, (\mathbb{C}^*)^n)$. By [Ser88, Thm. 6, p.184] and Theorem F.5.2.4, the natural map $\text{Ext}_{\text{Alg}}(A, \mathbb{G}_m^n) \rightarrow \text{Ext}(A^{\text{an}}, (\mathbb{C}^*)^n)$ is identified with the analytification map $[\text{Pic}^0(A)]^n \rightarrow [\text{Pic}^0(A^{\text{an}})]^n$, hence a group isomorphism. In particular, there is a unique exact sequence $0 \rightarrow \mathbb{G}_m^n \rightarrow C \rightarrow A \rightarrow 0$ in Alg whose analytification is equivalent to (F.30). \square

Lemma F.6.0.6 is used in the proof of Proposition F.6.0.7.

Lemma F.6.0.6. *Let G be a real Lie group with Lie algebra \mathfrak{g} .*

1. *If $X, Y \in \mathfrak{g}$ are such that $[X, [X, Y]] = 0$ and $[Y, [X, Y]] = 0$, then $\exp(X) \exp(Y) \exp(-X) \exp(-Y) = \exp([X, Y])$.*
2. *If $X \in \mathfrak{g}$ satisfies that $\exp(X)$ commutes with every element of G_0 and $[X, \mathfrak{g}] \subset Z(\mathfrak{g})$, then $X \in Z(\mathfrak{g})$.*

Proof.

1. According to Baker-Campbell-Hausdorff formula (see, e.g., [Far08, Cor. 3.4.5]), there is a symmetric open neighborhood U of $0 \in \mathfrak{g}$ such that for every $A, B \in U$, $\exp(A) \exp(B) = \exp(Z)$, where

$$Z = Z(A, B) = A + B + [A, B]/2 + \dots$$

and "..." indicates terms involving higher commutators of A and B . There is a symmetric open neighborhood V of $0 \in U$ such that $Z(A, B) \in U$ for every $A, B \in V$.

Define $f : \mathbb{R} \rightarrow G$ by

$$f(t) = \exp(tX) \exp(tY) \exp(-tX) \exp(-tY) \exp(-t^2[X, Y]).$$

Then f is real analytic. There is $\epsilon > 0$ such that $tX, tY \in V$ for all $t \in (-\epsilon, \epsilon)$. By assumption, $[Z(tX, tY), Z(-tX, -tY)] = 0$ and $Z(tX, tY) + Z(-tX, -tY) = t^2[X, Y]$. Then

$$f(t) = \exp(Z(tX, tY)) \exp(Z(-tX, -tY)) \exp(-t^2[X, Y]) = e_G$$

for all $t \in (-\epsilon, \epsilon)$ (see [Laz54, p.144]). By [ADGK23, Cor. A.5], $f(1) = e_G$.

2. Let $D = \exp^{-1}(e_G)$. There is an open neighborhood W of $0 \in \mathfrak{g}$ such that $\exp(W)$ is open in G and $\exp : W \rightarrow \exp(W)$ is a diffeomorphism. Then $D \cap W = \{0\}$. For every $Y \in \mathfrak{g}$, there is $k > 0$ with

$[X, Y/k] \in W$. By assumption, $[X, Y/k] \in Z(\mathfrak{g})$, so $[X, [X, Y/k]] = 0$ and $[Y/k, [X, Y/k]] = 0$. Since $\exp(Y/k) \in G_0$, it commutes with $\exp(X)$. By 1, $\exp([X, Y/k]) = e_G$. Then $[X, Y/k] \in D \cap W$. Therefore, $[X, Y] = 0$. Thus, $X \in Z(\mathfrak{g})$.

□

An algebraic analogue of Proposition F.6.0.7 is [Ros56, Cor. 2, p.433].

Proposition F.6.0.7. *Let $1 \rightarrow B \rightarrow C \xrightarrow{p} A \rightarrow 1$ be an exact sequence of complex Lie groups, with A complex torus and B commutative. Then C_0 is commutative.*

Proof. We may assume that C is connected by replacing C (resp. B) with C_0 (resp. $B \cap C_0$). Let $\omega : \mathbb{C}^g \rightarrow A$ be the universal covering of A . Denote by \mathfrak{b} (resp. \mathfrak{c}) the Lie algebra of B (resp. C). Let $\eta : A \rightarrow \text{Aut}(B)$ be the outer action. Then η induces a holomorphic morphism $\eta_0 : A \rightarrow \text{Aut}(B_0)$. Because $\text{Aut}(B_0)$ is complex Lie subgroup of $\text{GL}(\mathfrak{b})$, η_0 is trivial.

Consider the pullback extension along ω .

$$\begin{array}{ccccccc}
 & 0 & \longrightarrow & \ker(\epsilon) & \xrightarrow{\pi|_{\ker(\epsilon)}} & \ker(\omega) & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 1 & \longrightarrow & B & \longrightarrow & E & \xrightarrow{\pi} & \mathbb{C}^g \longrightarrow 1 \\
 & & \downarrow \text{Id} & & \downarrow \epsilon & \nearrow \pi & \downarrow \omega \\
 1 & \longrightarrow & B & \longrightarrow & C & \xrightarrow{p} & A \longrightarrow 1 \\
 & & \downarrow & & \downarrow & \nwarrow & \\
 & & 0 & & & &
 \end{array}$$

By the snake lemma, ϵ is surjective and π restricts to an isomorphism $\ker(\pi) \rightarrow \ker(\omega)$. In particular, $d_{e\epsilon} : L(E) \rightarrow L(C)$ is an isomorphism. By Fact F.2.0.3, the morphism ϵ is open. Since E_0 is open in E , $\epsilon(E_0)$ is an open subgroup of C . By connectedness of C , $\epsilon(E_0) = C$. Similarly, $\pi(E_0) = \mathbb{C}^g$. By Fact F.7.2.7 1 below, $B \cap E_0$ is connected. Therefore, $B \cap E_0 \subset B_0$. Since $B_0 \subset B \cap E_0$, one has $B_0 = B \cap E_0$. Hence an extension $1 \rightarrow B_0 \rightarrow E_0 \rightarrow \mathbb{C}^g \rightarrow 1$. The outer action is $\eta_0 \omega : \mathbb{C}^g \rightarrow \text{Aut}(B_0)$, so it is a central extension. Then

$$0 \rightarrow \mathfrak{b} \rightarrow \mathfrak{c} \rightarrow \mathbb{C}^g \rightarrow 0 \quad (\text{F.31})$$

is a central extension of Lie algebras. In particular, $\mathfrak{b} \subset Z(\mathfrak{c})$. We shall prove the extension (F.31) is trivial.

We show that $\exp_E : \mathfrak{c} \rightarrow E_0$ is surjective. Indeed, for every $x \in E_0$, there is $v \in \mathfrak{c}$ with $d_{e\epsilon}p(v) = \pi(x)$. Then $\pi(\exp_E(v)) = \pi(x)$, so $\pi(x \exp_E(-v)) = 0$ and hence $x \exp_E(-v) \in B_0$. As B_0 is connected commutative, there

is $u \in \mathfrak{b}$ with $\exp_B(u) = x \exp_E(-v)$. Since $u \in Z(\mathfrak{c})$, one gets $x = \exp_E(u) \exp_E(v) = \exp_E(u + v)$.

By Corollary F.4.1.13, there is a decomposition $B = B_0 \times D$, where $D \in \text{Ab}_c$ is discrete. The natural morphism $E_0 \times D \rightarrow E_0 \rightarrow \mathbb{C}^g$ is surjective of kernel $B_0 \times D$, hence the first row of the diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & B & \longrightarrow & E_0 \times D & \longrightarrow & \mathbb{C}^g \longrightarrow 1 \\
 & & \uparrow & & \uparrow & & \uparrow \text{Id} \\
 1 & \longrightarrow & B_0 & \longrightarrow & E_0 & \xrightarrow{\phi} & \mathbb{C}^g \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \text{Id} \\
 1 & \longrightarrow & B & \longrightarrow & E & \longrightarrow & \mathbb{C}^g \longrightarrow 1
 \end{array}$$

By Lemma F.3.1.2, there is an equivalence of extensions $\phi : E \rightarrow E_0 \times D$.

Fix $x \in \ker(\epsilon)$, let $\phi(x) = (\phi_1(x), \phi_2(x)) \in E_0 \times D$. For every $y \in E_0$,

$$(y, 1)\phi(x)(y, 1)^{-1} = (y\phi_1(x)y^{-1}, \phi_2(x)) \in \phi(\ker(\epsilon)).$$

Hence, $\phi^{-1}((y\phi_1(x)y^{-1}, \phi_2(x))) \in \ker(\epsilon)$. The map

$$E_0 \rightarrow \ker(\epsilon), \quad y \mapsto \phi^{-1}((y\phi_1(x)y^{-1}, \phi_2(x)))$$

is continuous. As E_0 is connected and $\ker(\epsilon)$ is discrete, this map is constantly x . Thus, $y\phi_1(x)y^{-1} = \phi_1(x)$. Therefore, $\phi_1(x)$ commutes with every element of E_0 . As $\exp_E : \mathfrak{c} \rightarrow E_0$ is surjective, there is $X \in \mathfrak{c}$ with $\exp_E(X) = \phi_1(x)$. Since \mathbb{C}^g is an abelian Lie algebra, $[\mathfrak{c}, \mathfrak{c}]$ is contained in the kernel of $d_{ep} : \mathfrak{c} \rightarrow \mathbb{C}^g$, which is \mathfrak{b} . Then $[\mathfrak{c}, \mathfrak{c}] \subset Z(\mathfrak{c})$, i.e., $[\mathfrak{c}, [\mathfrak{c}, \mathfrak{c}]] = 0$. By Lemma F.6.0.6 2, $X \in Z(\mathfrak{c})$.

Consider the commutative diagram

$$\begin{array}{ccc}
 \mathfrak{c} & \xrightarrow{d_{ep}} & \mathbb{C}^g \\
 \downarrow \exp_E & & \downarrow \text{Id} \\
 E & \xrightarrow{\pi} & \mathbb{C}^g
 \end{array}$$

Then $\pi(x) = \pi(\phi_1(x)) = d_{ep}(X) \in d_{ep}(Z(\mathfrak{c}))$. Therefore, $\ker(\omega) = \pi(\ker(\epsilon)) \subset d_{ep}(Z(\mathfrak{c}))$. Since d_{ep} is \mathbb{C} -linear and $\ker(\omega)$ contains a \mathbb{C} -basis of \mathbb{C}^g , one has $d_{ep}(Z(\mathfrak{c})) = \mathbb{C}^g$. Consequently, there is a \mathbb{C} -linear map $s : \mathbb{C}^g \rightarrow Z(\mathfrak{c})$ with $d_{ep} \circ s = \text{Id}_{\mathbb{C}^g}$. As $s : \mathbb{C}^g \rightarrow \mathfrak{c}$ is a Lie algebra morphism, the central extension (F.31) is trivial and \mathfrak{c} is the direct sum of \mathfrak{b} and \mathbb{C}^g . In particular, \mathfrak{c} is abelian. As C is connected and its Lie algebra is abelian, C is commutative. \square

Example F.6.0.8 shows that the condition that B/B_0 is torsion (resp. K_0 is Stein) in Proposition F.6.0.3 (resp. Proposition F.6.0.2 2) is necessary. Moreover, in Proposition F.6.0.7, the commutativity of C fails in general.

Example F.6.0.8. Let A be a complex torus and $B = A \times \mathbb{Z}$ be the product group. Consider the complex manifold morphism $A \times B \rightarrow B$ defined by $(a, a', k) \mapsto (a' + ka, k)$. It is a non trivial group action of A on B . Let C be the corresponding semidirect product (see [Bou72, Ch.III, no. 4, Prop. 7]), then the resulting complex Lie group extension $1 \rightarrow B \rightarrow C \rightarrow A \rightarrow 1$ is not central.

F.7 Noncommutative extensions

F.7.1 Lifted extensions

The real Lie group extension problem is studied by G. Hochschild in [Hoc51a] and [Hoc51b]. As Example F.7.1.1 shows, the case of real Lie groups is different from the case of complex Lie groups.

Example F.7.1.1. Let $G = \mathbb{C}$. The morphism of real Lie groups $\rho : \mathbb{C} \rightarrow \mathbb{C}^* = \text{Aut}(G)$ defined by $z \mapsto e^{\bar{z}}$ is an action of G on itself which is real analytic but not holomorphic. Hence an exact sequence of real Lie groups $1 \rightarrow G \rightarrow G \rtimes_{\rho} G \rightarrow G \rightarrow 1$ by [Bou72, Ch. III, no. 4, Prop. 7]. However, the middle term has no structure of complex Lie group making the maps holomorphic. Therefore, [Iwa49, Theorem 7] fails for complex Lie groups. Besides, this shows that the real Lie group extension problem and the complex one are different.

In Section F.7, we review Hochschild's work, but in the context of complex Lie groups. References to the original statement are given when the proofs are similar modulo slight modifications. All results in the sequel are essentially known.

In Section F.7.1, the goal is to derive Corollary F.7.1.6, a result about the extensions of a commutative group by a connected group.

Let L be a complex Lie group and $K \in \mathcal{C}$. For a fixed holomorphic group action $L \times K \rightarrow K$, let $\phi : L \rightarrow \text{Aut}(K)$ denote the induced group morphism. Let $Z(L, K, \phi)$ denote the set of crossed morphisms, i.e., morphisms $\rho : L \rightarrow K$ of complex manifolds such that $\rho(l_1 l_2) = \rho(l_1) \phi_{l_1}(\rho(l_2))$ for all $l_1, l_2 \in L$. Then $Z(L, K, \phi)$ is an abelian group under addition. (When ϕ is trivial, $Z(L, K, \phi) = \text{Hom}(L, K)$.)

For a normal complex Lie subgroup H of L , define

$$\text{Ophom}_L(H, K, \phi) = \{\psi \in \text{Hom}(H, K) : \psi(lhl^{-1}) = \phi_l(\psi(h)), \forall l \in L, h \in H\}.$$

Then $\text{Ophom}_L(H, K, \phi)$ is a subgroup of $\text{Hom}(H, K)$. When $H \subset Z(L)$, one has

$$\text{Ophom}_L(H, K, \phi) = \text{Hom}_{\mathcal{C}}(H, K^{\phi(L)}), \quad (\text{F.32})$$

where $K^{\phi(L)} = \cap_{l \in L} \{x \in K : \phi_l(x) = x\}$ is the set of elements fixed by $\phi(L) (\leq \text{Aut}(K))$. Here $K^{\phi(L)}$ is indeed a complex Lie subgroup of K by Corollary F.2.0.5. When ϕ is trivial, $\text{Ophom}_L(H, K, \phi)$ is the set of morphisms $H \rightarrow K$ invariant under the conjugation action of L .

Proposition F.7.1.2. *Assume that H is a normal complex Lie subgroup of L contained in $\ker(\phi)$. For every $\rho \in Z(L, K, \phi)$, $\rho|_H \in \text{Ophom}_L(H, K, \phi)$, hence a group morphism $Z(L, K, \phi) \rightarrow \text{Ophom}_L(H, K, \phi)$, whose image is denoted by $Z_H(L, K, \phi)$.*

Proof. For every $h, h' \in H$, $\rho(hh') = \rho(h)\phi_h(\rho(h')) = \rho(h)\rho(h')$ since $h \in \ker(\phi)$. Thus $\rho|_H \in \text{Hom}(H, K)$. In particular, $\rho(e_L) = e_K$. For every $l \in L$,

$$e_K = \rho(e_L) = \rho(ll^{-1}) = \rho(l)\phi_l(\rho(l^{-1})),$$

so $\rho(l)^{-1} = \phi_l(\rho(l^{-1}))$. Then

$$\begin{aligned} \rho(lhl^{-1}) &= \rho(lh)\phi_{lh}(\rho(l^{-1})) \\ &= \rho(lh)\phi_l(\rho(l^{-1})) = \rho(lh)\rho(l)^{-1} \\ &= \rho(l)\phi_l(\rho(h))\rho(l)^{-1} = \phi_l(\rho(h)). \end{aligned}$$

The last equality uses the commutativity of K . Therefore, $\rho|_H \in \text{Ophom}_L(H, K, \phi)$. \square

Let $\omega : Q' \rightarrow Q$ be a surjective morphism of connected complex Lie groups with kernel F . Let $\eta : Q \rightarrow \text{Aut}(K)$ be a group morphism such that the induced group action $Q \times K \rightarrow K$ is *holomorphic*. As K is commutative, the pulling back map $\omega^* : \text{Ext}(Q, K, \eta) \rightarrow \text{Ext}(Q', K, \eta\omega)$ is a group morphism. Fact F.7.1.3 gives a description of $\ker(\omega^*)$.

Define a map $\sigma : \text{Ophom}_{Q'}(F, K, \eta\omega) \rightarrow \text{Ext}(K, Q, \eta\omega)$ as follows. As the group action defined by η is holomorphic, the semidirect complex Lie group $K \rtimes_{\eta\omega} Q'$ exists by [Bou72, Ch.III, no.4, Prop. 7]. For $\psi \in \text{Ophom}_{Q'}(F, K, \eta\omega)$, the morphism $F \rightarrow K \rtimes_{\eta\omega} Q'$ defined by $k \mapsto (\psi(k), k)$ identifies F as a normal complex Lie subgroup of $K \rtimes_{\eta\omega} Q'$. Let $E = K \rtimes_{\eta\omega} Q' / F$. The projection $K \rtimes_{\eta\omega} Q' \rightarrow Q'$ descends to a morphism $E \rightarrow Q$. The injection $K \rightarrow K \rtimes_{\eta\omega} Q'$ induces a morphism $K \rightarrow E$. Then the resulting sequence $1 \rightarrow K \rightarrow E \rightarrow Q \rightarrow 1$ is exact with outer action $\eta\omega$, whose equivalence class is denoted by $\sigma(\psi)$.

Fact F.7.1.3 ([Hoc51a, Thm. 1.1]). *The map σ is a group morphism and the sequence*

$$Z(Q', K, \eta\omega) \rightarrow \text{Ophom}_{Q'}(F, K, \eta\omega) \xrightarrow{\sigma} \text{Ext}(Q, K, \eta) \xrightarrow{\omega^*} \text{Ext}(Q', K, \eta\omega)$$

is exact.

The use of Fact F.7.1.3 is based on the existence of $\omega : Q' \rightarrow Q$ such that every extension in $\text{Ext}(Q, K, \eta)$ becomes a semidirect product when pulled back to $\text{Ext}(Q', K, \eta\omega)$ along ω .

Fact F.7.1.4 ([Hoc51a, Thm. 2.1]). *Let Q be a connected complex Lie group. Assume that $\eta : Q \rightarrow \text{Aut}(K)$ is a group morphism such that the induced group action is holomorphic. Then there exists a simply connected complex Lie group Q' and a surjective morphism $\omega : Q' \rightarrow Q$ such that the pullback morphism $\omega^* : \text{Ext}(Q, K, \eta) \rightarrow \text{Ext}(Q', K, \eta\omega)$ is zero.*

Remark F.7.1.5. The connectedness condition of the extension kernel in [Hoc51a, Theorems 1.1 and 2.1] is in fact unnecessary.

Corollary F.7.1.6 follows from Fact F.7.1.3 and Fact F.7.1.4.

Corollary F.7.1.6 ([Hoc51a, Cor. 2.1]). *In the notation of Fact F.7.1.4, $\text{Ext}(Q, K, \eta) = \text{Ophom}_{Q'}(F, K, \eta\omega)/Z_F(Q', K, \eta\omega)$, where $F = \ker(\omega)$.*

Example F.7.1.7. Let $Q = \mathbb{C}^*$, $L = \mathbb{C}$ and $\omega : L \rightarrow \mathbb{Q}$ be defined by $\omega(z) = e^{2\pi iz}$. Then $F = \ker(\omega) = \mathbb{Z}$. Let $\mathbb{C}^* \times K \rightarrow K$ be a holomorphic group action and $\eta : \mathbb{C}^* \rightarrow \text{Aut}(K)$ be the induced group morphism. Then $\text{Ophom}_L(F, K, \eta\omega) = \text{Hom}(\mathbb{Z}, K^{\eta(\mathbb{C}^*)}) = K^{\eta(\mathbb{C}^*)}$. By Proposition F.3.2.2 and Corollary F.7.1.6, one has $\text{Ext}(\mathbb{C}^*K, \eta) = K^{\eta(\mathbb{C}^*)}/Z_{\mathbb{Z}}(\mathbb{C}, K, \eta\omega)$.

F.7.2 Factor systems

It is well-known that extensions of abstract groups can be classified in terms of factor systems, see [CE99, Ch. XIV, Sec. 4]. This description relies on the existence of set-theoretical cross sections. In general, nevertheless, it is not possible to find a continuous cross section to a surjective morphism of topological groups.

Consider the extension (F.3) of complex Lie groups with outer action $\psi : Q \rightarrow \text{Out}(K)$.

Example F.7.2.1. Assume that there is a cross section to (F.3), i.e., a morphism $s : Q \rightarrow E$ of complex manifolds with $ps = \text{Id}_Q$. Replacing s by $s(e_Q)^{-1}s$ when necessary, one may assume that s is normalized as $s(e_Q) = e_E$. Define

$$f : Q \times Q \rightarrow E, \quad f(g, h) = s(g)s(h)s(gh)^{-1}.$$

Then f is holomorphic. Since $p(f(g, h)) = e_Q$, $f(g, h) \in K$, so f factors through K . The map f measures the failure of s to be a morphism. If E is commutative, then additionally f is symmetric in the sense of [Ser88, (16), p.166]:

$$f(x, y) = f(y, x) \quad \forall x, y \in Q. \quad (\text{F.33})$$

Define $\phi : Q \rightarrow \text{Aut}(K)$ by $\phi_g = \text{Inn}_{s(g)}|_K$. Then ϕ is a map (but not necessarily a group morphism) lifting ψ , and the induced map

$$Q \times K \rightarrow K, \quad (g, x) \mapsto \phi_g(x) \quad (\text{F.34})$$

is holomorphic. When K is commutative, $\phi = \psi$ is a group morphism independent of the choice of s . When (F.3) is a central extension, ϕ is constantly Id_K .

Moreover, f and ϕ satisfy the following relations:

$$\begin{aligned} f(e_Q, h) &= f(g, e_Q) = e_K; \\ \phi_e &= \text{Id}_K; \\ \phi_g \phi_h &= \text{Inn}_{f(g, h)} \phi_{gh}; \\ f(g, h) f(gh, k) &= \phi_g(f(h, k)) f(g, hk). \end{aligned} \quad (\text{F.35})$$

Example F.7.2.1 motivates Definition F.7.2.2.

Definition F.7.2.2 (Factor system). If a morphism $f : Q \times Q \rightarrow K$ of complex manifolds and a map $\phi : Q \rightarrow \text{Aut}(K)$ making (F.34) holomorphic satisfy the relations (F.35), then f is called a ϕ -factor system (and simply a factor system when ϕ is trivial, in which case the last relation in (F.35) is $f(g, h) f(gh, k) = f(h, k) f(g, hk)$.) A factor system f is called symmetric if (F.33) holds.

When K is commutative, the set of ϕ -factor systems is an abelian group under addition.

We examine how the ϕ -factor system f induced by s in Example F.7.2.1 depends on the choice of the cross section s .

Example F.7.2.3. Let $s' : Q \rightarrow E$ be another normalized cross section still inducing ϕ . Define

$$g : Q \rightarrow E, \quad g(x) = s(x)^{-1} s'(x).$$

Then $g(e_Q) = e_E$ as s, s' are normalized and g is holomorphic. For every $x \in Q$, $p(g(x)) = e_Q$, so $g(x) \in K$. For every $k \in K$, $\text{Inn}_{s(x)} k = \phi_x(k) = \text{Inn}_{s'(x)} k$, so $g(x) \in Z(K)$, i.e., g factors through $Z(K)$. Then $s'(x) = s(x)g(x)$. Let f' be the factor system induced by s' . Then

$$\begin{aligned} f'(x, y) &= s'(x) s'(y) s'(xy)^{-1} \\ &= s(x) g(x) s(y) g(y) [s(xy) g(xy)]^{-1} \\ &= \phi_x(g(x)) s(x) s(y) g(y) g(xy)^{-1} s(xy)^{-1} \\ &= \phi_x(g(x)) f(x, y) s(xy) g(y) g(xy)^{-1} s(xy)^{-1} \\ &= \phi_x(g(x)) f(x, y) \phi_{xy}(g(y) g(xy)^{-1}) \\ &= g^\phi(x, y) f(x, y), \end{aligned}$$

where $g^\phi : Q \times Q \rightarrow K$ is a morphism of complex manifolds defined by

$$g^\phi(x, y) = \phi_x(g(x))\phi_{xy}(g(y)g(xy)^{-1}). \quad (\text{F.36})$$

When (F.3) is a central extension, ϕ is trivial, then (F.36) reduces to [Ser88, (15), p.166]: $g^\phi(x, y) = g(x)g(y)g(xy)^{-1}$.

Example F.7.2.3 motivates Definition F.7.2.4.

Definition F.7.2.4. Let f, f' be two ϕ -factor systems. If there is a holomorphic map $g : Q \rightarrow Z(K)$ with $g(e_Q) = e_E$ such that $f' = g^\phi f$ with g^ϕ defined by (F.36), then f and f' are called ϕ -equivalent, denoted by $f \sim_\phi f'$.

In Definition F.7.2.4, \sim_ϕ is an equivalent relation on the set of ϕ -factor systems. When K is commutative, inside the group of all ϕ -factor systems, the elements ϕ -equivalent to the zero form a subgroup. A result similar to Proposition F.7.2.5 for algebraic groups is in [Ser88, Ch. VII, Sec. 1, no.4].

Proposition F.7.2.5. Let K, Q be complex Lie groups with a map $\phi : Q \rightarrow \text{Aut}(K)$ such that (F.34) is holomorphic and the induced map $\psi : Q \rightarrow \text{Out}(K)$ is a group morphism. Then:

1. The set \mathcal{F} of \sim_ϕ -equivalence classes of ϕ -factor systems is canonically identified with the subset $\mathcal{E} \subset \text{Ext}(Q, K, \psi)$ of equivalence classes of extensions of Q by K which admit at least one normalized cross section inducing ϕ .
2. When K is commutative, the identification in 1 is a group isomorphism.
3. If further Q is also commutative and $\phi = \psi = 1$ is trivial, then the subgroup of equivalence classes of symmetric factor systems corresponds to the subgroup of equivalence classes of commutative extensions.

Proof. We only prove 1. Examples F.7.2.1 and F.7.2.3 construct a map $\Phi : \mathcal{E} \rightarrow \mathcal{F}$. (Note that equivalent extensions induces the same ϕ -equivalence class.)

Conversely, we define a map $\Psi : \mathcal{F} \rightarrow \mathcal{E}$ by the following construction. Given a ϕ -factor system f , one can construct an exact sequence $1 \rightarrow K \rightarrow E_{f,\phi} \rightarrow Q \rightarrow 1$ of complex Lie groups with a (holomorphic) normalized cross section $s : Q \rightarrow E_{f,\phi}$ as follows. Let $E_{f,\phi} = K \times Q$ as a complex manifold. Define a map

$$g : E_{f,\phi} \times E_{f,\phi} \rightarrow E_{f,\phi}, \quad g((k, x), (l, y)) = (k\phi_x(l)f(x, y), xy).$$

As f and the map (F.34) are holomorphic, so is g . Moreover, (F.35) shows g defines an associative multiplication. The pair $(1, 1) \in E_{f,\phi}$ is the identity, and the inverse of (k, x) is

$$(\phi_x^{-1}[k^{-1}f(x, x^{-1})^{-1}], x^{-1}).$$

Hence $(E_{f,\phi}, g)$ is a complex Lie group. The projection $p : E_{f,\phi} \rightarrow Q$ is a surjective morphism. The map $i : K \rightarrow E_{f,\phi}$ by $k \mapsto (k, 1)$ is the kernel of p . Moreover, define $s : Q \rightarrow E_{f,\phi}$ by $s(g) = (1, g)$, then s is normalized cross section. Put $\Psi(f) = E_{f,\phi}$.

We check that $\Psi\Phi = \text{Id}_{\mathcal{E}}$. Indeed, the map $E_{f,\phi} \rightarrow E$ defined by $(k, x) \mapsto ks(x)$ is an equivalence of extensions. We check that $\Phi\Psi = \text{Id}_{\mathcal{F}}$, or equivalently s induces f and ϕ . In fact, for every $x \in Q$, $k \in K$, one has

$$\phi_x(k)s(x) = (\phi_x(k), 1)(1, x) = (\phi_x(k), x) = (1, x)(k, 1) = s(x)k,$$

so $\phi_x = \text{Inn}_{s(x)}|_K$, i.e., s induces ϕ . For every $y \in Q$,

$$\begin{aligned} s(x)s(y)s(xy)^{-1} &= (1, x)(1, y)(1, xy)^{-1} \\ &= (f(x, y), xy)(\phi_{xy}^{-1}[f(xy, y^{-1}x^{-1})^{-1}], y^{-1}x^{-1}) \\ &= (f(x, y)\phi_{xy}\phi_{xy}^{-1}(f(xy, y^{-1}x^{-1})^{-1})f(xy, y^{-1}x^{-1}), 1) \\ &= (f(x, y), 1). \end{aligned}$$

Therefore, s induces f . □

When the base Q of (F.3) is discrete, then a set-theoretic cross section is automatically holomorphic.

Corollary F.7.2.6. *Let Q be a discrete complex Lie groups, and let $\eta : Q \rightarrow \text{Aut}(K)$ be a group morphism. Then the group $\text{Ext}(Q, K, \eta)$ is isomorphic to the group of \sim_η -equivalence classes of η -factor systems. Furthermore, if Q is also commutative, then $\text{Ext}(Q, K)$ is isomorphic to the group of \sim -equivalence classes of symmetric factor systems.*

Proof. Since Q is discrete, the group action $Q \times K \rightarrow K$ induced by η is holomorphic. The first (resp. second) half follows from Proposition F.7.2.5 2 (resp. 3). □

Another important case where a cross section exists is with simply connected bases. For this, we need a holomorphic version of Malcev's theorem ([Mal42, (E), p.12], [Hoc51a, Lemma 3.1], [Mac60, Theorem 3.2]).

Fact F.7.2.7 (Malcev, [Bou72, Ch. III, § 6, no. 6, Prop. 14; Cor. 2]). *Let L be a connected complex Lie group, N be a normal immersed complex Lie subgroup of L .*

1. *If N is closed in L and L/N is simply connected, then N is connected.*
2. *If L is simply connected, N is connected, then N is closed in L and there exists a biholomorphic map $f : L \rightarrow N \times L/N$ making a commutative diagram*

$$\begin{array}{ccc}
L & \xrightarrow{f} & N \times L/N \\
& \searrow q & \downarrow p_2 \\
& & L/N,
\end{array}$$

where p_2 is the projection to the second factor and $q : L \rightarrow L/N$ is the quotient morphism.

In the same way that [Hoc51a, Theorem 3.1] follows from [Hoc51a, Lemma 3.1], Fact F.7.2.8 can be deduced from Fact F.7.2.7.

Fact F.7.2.8. *Let (F.3) be an exact sequence of complex Lie groups, where E is connected and Q is simply connected. Then there exists a cross section, i.e., a holomorphic map $s : Q \rightarrow E$ with $ps = \text{Id}_Q$. In particular, the principal K -bundle $p : E \rightarrow Q$ is trivial.*

Example F.7.2.9. Let A be a complex elliptic curve. Take a nonzero element of A^\vee , which induces a nontrivial extension E of A by \mathbb{C}^* via Theorem F.5.2.4. By Proposition F.5.1.3, the principal \mathbb{C}^* -bundle $E \rightarrow A$ is nontrivial. Therefore, Fact F.7.2.8 fails if the base is not simply connected.

Corollary F.7.2.10 follows immediately from Fact F.7.2.8 and Proposition F.7.2.5.

Corollary F.7.2.10. *Let K, Q be complex Lie groups, where K is connected commutative and Q is simply connected. Let $\eta : Q \rightarrow \text{Aut}(K)$ be a complex Lie group morphism¹¹. Then $\text{Ext}(Q, K, \eta)$ is isomorphic to the group of \sim_η -equivalence classes of η -factor systems.*

Similar to [Hoc51a, Theorem 3.2], Fact F.7.2.11 can be proved using Fact F.7.2.7 and Fact F.7.2.8,

Fact F.7.2.11. *Let K, Q be complex Lie groups, where K is connected and Q is simply connected. Then the map (on the set of equivalence classes) which associates with each extension of Q by K the induced extension of $L(Q)$ by $L(K)$ is injective. The image is the set of classes of those extensions $0 \rightarrow L(K) \rightarrow \mathfrak{E} \rightarrow L(Q) \rightarrow 0$ in which the derivation*

$$[x, \bullet]_{\mathfrak{E}|_{L(K)}} \in \text{Der}(L(K)) = L(\text{Aut}(L(K)))$$

belongs to $L(\text{Aut}(K))$ for every $x \in \mathfrak{E}$. Furthermore, if K is commutative and $\eta : Q \rightarrow \text{Aut}(K)$ is a morphism, then the resulting map

$$\text{Ext}(Q, K, \eta) \rightarrow \text{Ext}(L(Q), L(K), d_\eta \eta)$$

is a group isomorphism.

¹¹Here $\text{Aut}(K)$ is a complex Lie subgroup of $\text{GL}(L(K))$ by [Lee01, Propositions 1.26 and 1.27].

A connected Lie group is called semisimple if its Lie algebra is semisimple. Analogue of Fact F.7.2.12 for semisimple real Lie groups H and real vector groups G is contained in the proof of [Hoc51b, Theorem 5.1]. Fact F.7.2.12 can be proved in a similar way.

Fact F.7.2.12. *Let G, H be connected complex Lie groups, where G is commutative and H is semisimple. Let $\eta : H \rightarrow \text{Aut}(G)$ be a morphism of complex Lie groups. If $\phi \in Z(H, G, \eta)$ is a crossed morphism, then there exists $g \in G$ such that $\phi(x) = \eta_x(g)g^{-1}$ for all $x \in H$. In particular, $\phi \equiv e_G$ on $\ker(\eta)$.*

Theorem F.7.2.13 is a complex version of [Hoc51a, Theorem 4.4].

Theorem F.7.2.13. *In Fact F.7.2.12, $\text{Ext}(H, G, \eta)$ is canonically isomorphic to $\text{Hom}_{\text{Ab}}(\pi_1(H), G^{\eta(H)})$.*

Proof. Let $\omega : \tilde{H} \rightarrow H$ be the universal covering of H . Then $\ker(\omega) = \pi_1(H)$ is a discrete subgroup of \tilde{H} . By Fact F.3.2.4, $\pi_1(H) \subset Z(\tilde{H})$. Then (F.32) gives

$$\text{Ophom}_{\tilde{H}}(\ker(\omega), G, \eta\omega) = \text{Hom}(\pi_1(H), G^{\eta(H)}).$$

By Fact F.7.2.12, for every $\rho \in Z(\tilde{H}, G, \eta\omega)$, $\rho|_{\ker \omega} = 1$, i.e., $Z_{\ker(\omega)}(\tilde{H}, G, \eta\omega) = 0$. By Fact F.7.2.11, the natural map $\text{Ext}(\tilde{H}, G, \eta\omega) \rightarrow \text{Ext}(L(H), L(G), d_e\eta)$ is a group isomorphism. Since $L(H)$ is a semisimple complex Lie algebra, Levi's theorem [Ser64, Theorem 4.1, p.48] affirms that $\text{Ext}(L(H), L(G), d_e\eta) = 0$. By Fact F.7.1.3, $\text{Ext}(H, G, \eta) = \text{Hom}(\pi_1(H), G^{\eta(H)})$. \square

F.7.3 Non-abelian kernels and extensions of the center

For two complex Lie groups K, Q and a group morphism $\theta : Q \rightarrow \text{Out}(K)$, if θ is induced by some extension of Q by K , then the extension kernel (K, θ) is called *extendible*. The problem to determine the extendibility of a given extension kernel is more difficult than that for abstract groups treated in [EM47, Theorem 8.1], because of the obstruction to the existence of a cross section. For extendible kernels, Corollary F.7.3.8 shows that the problem for extensions by K can be reduced to that with an *abelian* kernel, namely $Z(K)$.

Let $1 \rightarrow K \rightarrow E \xrightarrow{p} Q \rightarrow 1$ and $1 \rightarrow K' \rightarrow E' \xrightarrow{p'} Q \rightarrow 1$ be two extension of complex Lie groups. Denote their outer action by $\theta : Q \rightarrow \text{Out}(K)$ and $\theta' : Q \rightarrow \text{Out}(K')$ respectively. Assume that $Z(K) = Z(K') := C$ and θ, θ' induce a common center action¹² $\theta_0 : Q \rightarrow \text{Aut}(C)$. Hence a commutative

¹²see (F.5)

diagram

$$\begin{array}{ccc}
 & \text{Out}(K) & \\
 \theta \nearrow & & \searrow \\
 Q & \xrightarrow{\theta_0} & \text{Aut}(C) \\
 \theta' \searrow & & \nearrow \\
 & \text{Out}(K') &
 \end{array} \tag{F.37}$$

We recall the multiplication of kernels defined in [EM47, Sec. 4]. The group law $C \times C \rightarrow C$ is holomorphic, so the subset

$$C^* := \{(x, x^{-1}) : x \in C\} \tag{F.38}$$

is analytic in $C \times C$. By Lemma F.2.0.6, $C \times C$ is an analytic subset of $K \times K'$. As C^* is a central subgroup of $K \times K'$, it is also a complex Lie subgroup of $K \times K'$ by Corollary F.2.0.5. Let $K'' = K \times K' / C^*$. From [EM47, p.328], the morphism $C \rightarrow K''$ by $g \mapsto [(g, 1)]$ identifies C as the center of K'' .

For every $x \in Q$, select automorphisms $\alpha \in \theta(x)(\subset \text{Aut}(K))$ and $\alpha' \in \theta'(x)(\subset \text{Aut}(K'))$. Because the diagram (F.37) is commutative, $\alpha \times \alpha'$ is an automorphism of $K \times K'$ sending C^* into itself. It thus determines an automorphism α'' of K'' . The class $[\alpha''] \in \text{Out}(K'')$ depends only on θ, θ' , but not the choices of α, α' . Hence a group morphism

$$\theta'' : Q \rightarrow \text{Out}(K'') \tag{F.39}$$

that also induces $\theta_0 : Q \rightarrow \text{Aut}(C)$.

Definition F.7.3.1. The pair (K'', θ'') constructed above is called the C -product of the two given extension kernels (K, θ) and (K', θ') .

Example F.7.3.2. If $K' = C$ is commutative, it is asserted in [EM47, (4.4)] that K' acts as an identity for the C -product. To make it explicit, we define a surjective morphism $\phi : K \times C \rightarrow K$ of complex manifolds by $\phi(k, k') = k'k$. Then ϕ is a morphism and $C^* = \ker(\phi)$. Thus, ϕ induces an isomorphism $\sigma : K'' \rightarrow K$ satisfying [EM47, (4.2), (4.3)].

Then we review the multiplication of the given two extensions, contained the proof of [EM47, Lem. 5.1].

As the map $E \times E' \rightarrow Q$ by $(x, x') \mapsto p'(x')p(x)^{-1}$ is holomorphic, the preimage of e_Q

$$D = D_{p,p'}(E, E') = \{(x, x') \in E \times E' : p(x) = p'(x')\}, \tag{F.40}$$

is analytic in $E \times E'$. Since D is a subgroup of $E \times E'$, by Corollary F.2.0.5, D is a complex Lie subgroup of $E \times E'$.

For every $(x, x') \in D$ with $y = p(x) = p(x')$, every $g \in C$, the element

$$(x, x')(g, g^{-1})(x^{-1}, x'^{-1}) = (\theta_0(y)(g), \theta_0(y)(g)^{-1})$$

is in C^* . Therefore, C^* defined by (F.38) is normal in D .

As C^* is a normal complex Lie subgroup of D , we can set $E'' = D/C^*$. The inclusion $K \times K' \rightarrow D$ descends to an injective morphism $K'' \rightarrow E''$. The map $D \rightarrow Q$ defined by $(x, x') \mapsto p(x)$ induces a surjective morphism $p'' : E'' \rightarrow Q$ whose kernel is K'' . Hence an extension $1 \rightarrow K'' \rightarrow E'' \rightarrow Q \rightarrow 1$. The induced outer action $Q \rightarrow \text{Out}(K'')$ is (F.39). We call (E'', p'') the C -product of the two given extensions (E, p) and (E', p') , written as $(E'', p'') = (E, p) \otimes (E', p')$. Thus, [EM47, Lemmas 5.1 and 5.2] hold for complex Lie groups.

Fact F.7.3.3. *The C -product of two extendible kernels is extendible. The kernel of the C -product $(E, p) \otimes (E', p')$ of two extensions is the C -product of the two kernels.*

Proposition F.7.3.4. *When $K' = C$, (E', p') is the semidirect product $C \rtimes_{\theta_0} Q$, then (E'', p'') is naturally equivalent to (E, p) .*

Proof. Consider the subgroup $D \leq E \times E' = E \times (C \rtimes_{\theta_0} Q)$ defined in (F.40). Define a map $\psi : D \rightarrow E$ by $(x, c, q) \mapsto cx$ for $x \in E$ and $(c, q) \in C \rtimes_{\theta_0} Q$. Then ψ is holomorphic.

We check that ψ is a group morphism. Take another $(x, c', q') \in D$. Since $\theta_{0,q}(c') = \theta_{p(x)}(c') = xc'x^{-1}$, one has

$$\begin{aligned} \psi((x, c, q)(x', c', q')) &= \psi(xx', c\theta_{0,q}(c'), qq') \\ &= c\theta_{0,q}(c')xx' = cxc'x' = \psi(x, c, q)\psi(x', c', q'). \end{aligned}$$

For every $g \in C$, $\psi(g, g^{-1}) = e_E$, so $C^* \subset \ker \psi$. Thus, ψ induces a morphism $\epsilon : E'' \rightarrow E$. Together with σ defined in Example F.7.3.2, ϵ fits into a commutative diagram.

$$\begin{array}{ccccccc} 1 & \longrightarrow & K'' & \longrightarrow & E'' & \longrightarrow & Q \longrightarrow 1 \\ & & \downarrow \sigma & & \downarrow \epsilon & & \downarrow \text{Id} \\ 1 & \longrightarrow & K & \longrightarrow & E & \longrightarrow & Q \longrightarrow 1 \end{array}$$

Therefore, ϵ is an equivalence of extensions. \square

By construction, C -product defines a map $\text{Ext}(Q, K, \theta) \times \text{Ext}(Q, K', \theta') \rightarrow \text{Ext}(Q, K'', \theta'')$. When $K' = C$, it specializes to

$$\text{Ext}(Q, K, \theta) \times \text{Ext}(Q, C, \theta_0) \rightarrow \text{Ext}(Q, K, \theta), \quad (\text{F.41})$$

which defines an action of the abelian group $\text{Ext}(Q, C, \theta_0)$ on the set $\text{Ext}(Q, K, \theta)$. If further K is also commutative, by [Hoc51a, p.97], (F.41) is exactly the group law defined by the Baer sum on $\text{Ext}(Q, C, \theta_0)$.

Definition F.7.3.5. [EM47, p.329] For every extension kernel (K, θ) , let θ^* be the composition of $\theta : Q \rightarrow \text{Out}(K)$ with the natural group isomorphism $\text{Out}(K) \rightarrow \text{Out}(K^{\text{op}})$. Then the extension kernel $(K^{\text{op}}, \theta^*)$ is called the inverse of (K, θ) .

For every $(E, p) \in \text{Ext}(Q, K, \theta)$, define $p^* : E^{\text{op}} \rightarrow Q$ by $p^*(x^*) = p(x^{-1})$, then it is a surjective morphism. Since $\ker(p^*) = K^{\text{op}}$, $1 \rightarrow K^{\text{op}} \rightarrow E^{\text{op}} \xrightarrow{p^*} Q \rightarrow 1$ is an extension. The associated outer action is θ^* . Thus, we get an element $(E^{\text{op}}, p^*) \in \text{Ext}(Q, K^{\text{op}}, \theta^*)$ of (E, p) . It is called the inverse of (E, p) and its extension kernel is the inverse of (K, θ) .

It is a classical result that the group action (F.41) is simple transitive. For abstract groups, see [EM47, Lem. 11.2 and 11.3]. For algebraic groups, see [FLA19, Thm. 1.1]. It remains true for complex Lie groups. The first half, Fact F.7.3.6, can be proved in the same way as in [Hoc51b, Thm. 1.1], using the inverse in the group $\text{Ext}(Q, C, \theta_0)$ and Proposition F.7.3.4.

Fact F.7.3.6. *Let K, Q be complex Lie groups, $C = Z(K)$. Let $\theta : Q \rightarrow \text{Out}(K)$ be a group morphism that induces $\theta_0 : Q \rightarrow \text{Aut}(C)$. Then the action of $\text{Ext}(Q, C, \theta_0)$ on $\text{Ext}(Q, K, \theta)$ defined by (F.41) is free.*

Theorem F.7.3.7 is analogue to [EM47, Lemma 11.2].

Theorem F.7.3.7. *In the notation of Fact F.7.3.6, if $\text{Ext}(Q, K, \theta)$ is nonempty (i.e., the extension kernel (K, θ) is extendible), then its $\text{Ext}(Q, C, \theta_0)$ -action defined by (F.41) is transitive. Equivalently, for every $(E, p), (E_1, p_1) \in \text{Ext}(Q, K, \theta)$, there exists $F \in \text{Ext}(Q, C, \theta_0)$ with $F \otimes E$ equivalent to E_1 .*

Proof. Define $D_{p_1, p^*}(E_1, E^{\text{op}})$ like (F.40). Set

$$S = \{(x_1^{-1}, x^*) \in D_{p_1, p^*}(E_1, E^{\text{op}}) : x_1 k x_1^{-1} = x k x^{-1}, \forall k \in K\}.$$

Then S is a subgroup of $E_1 \times E^{\text{op}}$. For every $k \in K$, the map

$$\phi_k : E_1 \times E^{\text{op}} \rightarrow K \quad (x_1, x^*) \mapsto x_1^{-1} k x_1 x k^{-1} x^{-1}$$

is holomorphic, so $\phi_k^{-1}(e_K)$ is analytic in $E_1 \times E^{\text{op}}$. Then $S = D_{p_1, p^*}(E_1, E^{\text{op}}) \cap \bigcap_{k \in K} \phi_k^{-1}(e_K)$ is analytic in $E_1 \times E^{\text{op}}$, by [Whi72, Theorem 9C, p.100]. By Corollary F.2.0.5, S is a complex Lie subgroup of $E_1 \times E^{\text{op}}$.

The map $K \times K^{\text{op}} \rightarrow K$ by $(k, k'^*) \mapsto k k'$ is holomorphic, so $K^* = \{(k^{-1}, k^*) : k \in K\}$ is an analytic subset of $K \times K^{\text{op}}$. It is a subgroup of S , hence a complex Lie subgroup of S by Corollary F.2.0.5.

For every $(x_1^{-1}, x^*) \in S$, $k \in K$, one has

$$\begin{aligned} (x_1^{-1}, x^*)(k^{-1}, k^*)(x_1, (x^*)^{-1}) &= (x_1^{-1} k^{-1} x_1, x^* k^* (x^{-1})^*) \\ &= (x^{-1} k^{-1} x, (x^{-1} k x)^*) \in K^*, \end{aligned}$$

so K^* is a normal subgroup of S . Let $F = S/K^*$ and $\nu : S \rightarrow F$ be the quotient morphism. The map $i : C \rightarrow F$ defined by $c \mapsto [(c, 1)]$ is an injective morphism.

The map $\bar{\phi} : S \rightarrow Q$ defined by $\bar{\phi}(x_1^{-1}, x^*) = p(x^{-1})$ is a morphism with K^* contained in the kernel. We check that $\bar{\phi}$ is surjective. For every $h \in Q$, there exist $x \in E$ and $x_1 \in E_1$ with $p(x) = p_1(x_1) = h^{-1}$. Since the two automorphisms of K , $\text{Inn}_x|_K$ and $\text{Inn}_{x_1}|_K$ have the same class $\theta_{h^{-1}}$ in $\text{Out}(K)$, there exists $k_0 \in K$ such that $\text{Inn}_{x_1}|_K = \text{Inn}_x|_K \text{Inn}_{k_0}$. Then $(x_1^{-1}, (xk_0)^*) \in S$ and $\bar{\phi}(x_1^{-1}, (xk_0)^*) = h$.

If $(x_1^{-1}, x^*) \in \ker \bar{\phi}$, then $p_1(x_1) = p(x_1) = e_Q$, so $x_1, x \in K$. Moreover, $x_1 k x_1^{-1} = x k x^{-1}$ for all $k \in K$. Then $x_1^{-1} x \in C$, so $(x_1^{-1}, x^*) = (x_1^{-1} x, 1^*)(x^{-1}, x^*)$. Thus, $[(x_1^{-1}, x^*)] = i(x_1^{-1} x) \in i(C)$.

Thus $\bar{\phi}$ induces a surjective morphism $\phi : F \rightarrow Q$ with $i(C) \supset \ker \phi$. In addition, ϕi is trivial, so $i(C) \subset \ker(\phi)$. Hence an extension $1 \rightarrow C \xrightarrow{i} F \xrightarrow{\phi} Q \rightarrow 1$ with the induced action $Q \rightarrow \text{Aut}(C)$ coinciding with θ_0 .

It remains to show that the C -product extension $F \otimes E$ is equivalent to E_1 . By construction, $F \otimes E$ is represented by $G = D_{\phi,p}(F, E)/C^*$, where $C^* = \{(c, c^{-1}) \in F \times E : c \in C\}$. The pullback of $D_{\phi,p}(F, E)$ along the natural surjection $S \times E \rightarrow F \times E$ is $D_{\phi\nu,p}(S, E)$.

For every $(a, b^*, x) \in D_{\phi\nu,p}(S, E) \subset E_1 \times E^{\text{op}} \times E$, one has $p_1(a) = p(b^{-1}) = p(x)$, whence $bx \in K$ and $a \cdot (bx) \in E_1$. Define a holomorphic map $\tau : D_{\phi\nu,p}(S, E) \rightarrow E_1$ by $\tau(a, b^*, x) = a \cdot (bx)$.

$$\begin{array}{ccccc}
 & E_1 & & & \\
 & \nwarrow \tau & & \nearrow \nu^* & \\
 D_{\phi\nu,p}(S, E) & \longrightarrow & D_{\phi,p}(F, E) & \longrightarrow & G \\
 \downarrow & & \downarrow & & \\
 S \times E & \xrightarrow{\nu \times \text{Id}_E} & F \times E & & \\
 \downarrow & & & & \\
 E_1 \times E^{\text{op}} \times E & & & &
 \end{array}$$

We check that τ is a group morphism. For every $(a, b^*, x), (a', b'^*, x') \in D_{\phi\nu,p}(S, E)$, since $(a', b'^*) \in S$ and $bx \in K$, one has $a'^{-1}(bx)a' = b'(bx)b'^{-1}$. Hence,

$$\begin{aligned}
 \tau(a, b^*, x) \tau(a', b'^*, x') &= [a(bx)][a'(b'x')] \\
 &= aa'[a'^{-1}(bx)a'](b'x') = aa'[b'(bx)b'^{-1}](b'x') \\
 &= aa'(b'bxx') = \tau(aa', (b'b)^*, xx') = \tau(aa', b^*b'^*, xx').
 \end{aligned}$$

We check that τ is surjective. For every $x_1 \in E_1$, $p_1(x_1) \in Q$. As $\phi\nu : S \rightarrow Q$ is surjective, there is $(a, b^*) \in S$ with $\phi\nu(a, b^*) = p_1(x_1)$. Then $p_1(a) = p_1(x_1)$. Thus, $a^{-1}x_1 \in K$. Let $x = b^{-1}(a^{-1}x_1) \in E$. Then $p(x) =$

$p(b^{-1}) = \phi\nu(a, b^*)$, so $(a, b^*, x) \in D_{\phi\nu, p}(S, E)$ and $\tau(a, b^*, x) = a(bx) = a(a^{-1}x_1) = x_1$.

We check that $\ker(\nu^*) \subset \ker(\tau)$. For every $(x_1, x^*, y) \in \ker(\nu^*) \subset E_1 \times E^{\text{op}} \times E$, there is $c \in C$ with $[(x_1, x^*)], y = (c, c^{-1})$ in $F \times E$. Equivalently, $y = c^{-1}$ in E and $[(x_1, x^*)] = [(c, 1^*)]$ in $F = S/K^*$. Whence, $(x_1 c^{-1}, x^*) \in K^*$, i.e., $x \in K$ and $x_1 = x^{-1}c$. Therefore, $(x_1, x^*, y) = (x^{-1}c, x^*, c^{-1})$ with $x \in K, c \in C$. Thus, $\tau(x_1, x^*, y) = x^{-1}c(xc^{-1}) = e_{E_1}$ and $(x_1, x^*, y) \in \ker(\tau)$.

Conversely, we check $\ker(\tau) \subset \ker(\nu^*)$. For every $(a, b^*, x) \in \ker(\tau)$, one has $a(bx) = e_{E_1}$, so $a \in K$. Because $(a, b^*) \in D_{p_1, p^*}(E_1, E^{\text{op}})$, we obtain $p(b^{-1}) = p(a) = e_Q$ and hence $b \in K$. Since $\text{Inn}_{a^{-1}} = \text{Inn}_b \in \text{Aut}(K)$, one has $ab \in C$. Therefore, $[(a, b^*)] = [(ab, 1^*)] = i(ab)$ in $F = S/K^*$ and $(a, b^*, x) = (ab, (ab)^{-1}) \in C^* \leq F \times E$. Then $(a, b^*, x) \in \ker(\nu^*)$.

Therefore, $\ker(\tau) = \ker(\nu^*)$, so τ induces an isomorphism $G \rightarrow E_1$ that establishes an equivalence between the two elements of $\text{Ext}(Q, K, \theta)$. \square

Fact F.7.3.6 and Theorem F.7.3.7 yield Corollary F.7.3.8.

Corollary F.7.3.8. *Let K, Q be complex Lie groups, $C = Z(K)$, $\theta : Q \rightarrow \text{Out}(K)$ be a group morphism. Let $\theta_0 : Q \rightarrow \text{Aut}(C)$ be the induced group morphism. If $\text{Ext}(Q, K, \theta)$ is nonempty, then $\text{Ext}(Q, K, \theta)$ is in (non-canonical) bijection with $\text{Ext}(Q, C, \theta_0)$.*

F.8 Maximal morphisms

A result stronger than Proposition F.5.1.3 holds.

Definition F.8.0.1. [Ser88, Definition 1, p.125]. Let X be a complex manifold, A be a complex torus. A morphism $f : X \rightarrow A$ is called maximal if whenever f factors as $X \xrightarrow{g} A' \xrightarrow{h} A$, where $A' \in \mathcal{C}$ is connected and $h - h(0) : A' \rightarrow A$ is a finite morphism, it holds that $h - h(0)$ is an isomorphism.

Proposition F.8.0.2. *If X is a regular manifold¹³, then the Albanese morphism $f : X \rightarrow \text{Alb}(X)$ associated to some base point $x \in X$ is maximal.*

Proof. Assume that f factors as $X \xrightarrow{g} A' \xrightarrow{h} \text{Alb}(X)$, where $A' \in \mathcal{C}$ is a connected and $h - h(0)$ is a finite morphism. Then A' is compact, hence a complex torus. Choosing $g(x)$ as the new zero element of A' , we get a new structure of complex torus on A' , to which we stick from now on. Then h is a finite morphism. By Proposition 4.4.1.2 3, there is a morphism $\phi : \text{Alb}(X) \rightarrow A'$ with $\phi f = g$ and the complex Lie subgroup of $\text{Alb}(X)$ generated by $f(X)$ is $\text{Alb}(X)$ itself. Then $h\phi f = f$ and hence

¹³in the sense of [Var86, p.233]

$h\phi = \text{Id}_{\text{Alb}(X)}$. In particular, h is surjective. By Fact F.3.0.4, the exact sequence $0 \rightarrow \ker(h) \rightarrow A' \xrightarrow{h} A \rightarrow 0$ defines a trivial extension, so A' is isomorphic to $\ker(h) \times A$. By connectedness of A' , $\ker(h) = 0$ and h is an isomorphism. \square

When $f = \text{Id}_A$, Proposition F.8.0.3 reduces to Proposition F.5.1.3.

Proposition F.8.0.3 ([Ser88, Prop. 14, p.188]). *Let X be a connected compact complex manifold, A be a complex torus, $B \in \mathcal{C}$. Let $f : X \rightarrow A$ be a maximal morphism. If B_0 is linear, then the composed morphism*

$$\text{Ext}(A, B) \xrightarrow{\pi} H^1(A, \mathcal{B}_A) \xrightarrow{f^*} H^1(X, \mathcal{B}_X) \quad (\text{F.42})$$

is injective.

Proof. Let $C \in \ker(f^* \circ \pi)$. Then the principal fiber bundle $f^*p : f^*C \rightarrow X$ is trivial. Fix a point $c \in f^*C$ lying over $0 \in C$. Then there is a morphism $s : X \rightarrow f^*C$ with $f^*p \circ s = \text{Id}_X$ and $s(f^*p(c)) = c$. Let $t : X \rightarrow C$ be the morphism induced by s .

$$\begin{array}{ccccccc} & & & & f^*C & \xrightarrow{f^*p} & X \\ & & & & \downarrow & \swarrow t & \downarrow f \\ 0 & \longrightarrow & B & \longrightarrow & C & \xrightarrow{p} & A \longrightarrow 0 \\ & & \uparrow & & \uparrow & \nwarrow h^{-1} & \uparrow \text{Id} \\ 0 & \longrightarrow & B \cap A' & \longrightarrow & A' & \xrightarrow{h} & A \longrightarrow 0 \end{array}$$

By Remmert's theorem [Whi72, Theorem 4A, p.150], $t(X)$ is an analytic subset of C . By [CD94, (14.14), p.89], the analytic space $t(X)$ is irreducible. Moreover, $t(X)$ is compact and $0 = t(f^*p(c)) \in t(X)$. Let A' be the complex Lie subgroup of C generated by $t(X)$. By Lemma D.3.2.1, A' is a complex torus. Then $(A' \cap B)_0$ is a compact. As a closed complex submanifold of B_0 , $(A' \cap B)_0$ is also a Stein manifold, hence a point. Thus, $A' \cap B$ is discrete and compact, hence finite. Therefore, $h : A' \rightarrow A$ is a finite morphism. As the maximal morphism f factors as $X \xrightarrow{t} A' \xrightarrow{h} A$, h is an isomorphism. Then $h^{-1} : A \rightarrow C$ is a morphism and $ph^{-1} = \text{Id}_A$. By Fact F.3.0.4, $C = 0$ in $\text{Ext}(A, B)$. \square

Example F.8.0.4. Let X be a regular manifold, $f : X \rightarrow A$ be the Albanese morphism associated to some base point $x \in X$. When $B = \mathbb{C}$, the composed morphism (F.42) is a linear isomorphism $f^* : H^1(A, \mathcal{O}_A) \rightarrow H^1(X, \mathcal{O}_X)$. When $B = \mathbb{C}^*$, it is the inclusion of the identity component $\text{Pic}^0(A) \rightarrow \text{Pic}(X)$.

F.9 Commutative extensions of real Lie groups

Let \mathcal{R} be the category of commutative real Lie groups. The solution to the extension problem within \mathcal{R} is summarized in Proposition F.9.0.2. Similar to Lemma F.4.1.1, the category \mathcal{R} is additive but not abelian. Parallel to the construction in Section F.4, we can define an additive functor $\text{Ext}_{\mathcal{R}} : \mathcal{R}^{\text{op}} \times \mathcal{R} \rightarrow \text{Ab}$ by considering commutative extensions.

Proposition F.9.0.1 generalizes [IH76, Proposition 5, p.110] (which says that C is isomorphic to $A \times B$) and [HN11, Lemma 15.3.2] (which is for real tori). The similar statement for complex tori is false, shown by Example F.4.1.14.

Proposition F.9.0.1. *Let $0 \rightarrow B \rightarrow C \rightarrow A \rightarrow 0$ be an extension of commutative real Lie groups. If A, B are connected, this extension is trivial.*

Proof. Similar to Proposition F.3.2.2, every extension of \mathbb{R} is a semidirect product, hence $\text{Ext}_{\mathcal{R}}(\mathbb{R}, \bullet) = 0$ on \mathcal{R} . Similar to Proposition F.3.2.3, $\text{Ext}_{\mathcal{R}}(S^1, B) = 0$. According to [IH76, Proposition 4, p.109], A is isomorphic to $(S^1)^n \times \mathbb{R}^m$ for some $m, n \in \mathbb{N}$. As the functor $\text{Ext}_{\mathcal{R}}(\bullet, B) : \mathcal{R} \rightarrow \text{Ab}$ is additive, we get $\text{Ext}_{\mathcal{R}}(A, B) = 0$. \square

Proposition F.9.0.2. *For every $A, B \in \mathcal{R}$, there is a non canonical isomorphism in Ab :*

$$\text{Ext}_{\mathcal{R}}(A, B) \rightarrow \text{Ext}_{\mathbb{Z}}^1(A/A_0, B/B_0) \oplus \text{Hom}_{\text{Ab}}(\pi_1(A_0), B/B_0).$$

Proof. By a real version of Corollary F.4.1.13, there are non canonical isomorphisms in \mathcal{R} : $A \rightarrow A/A_0 \times A_0$ and $B \rightarrow B/B_0 \times B_0$. By additivity of the bifunctor $\text{Ext}_{\mathcal{R}}$, we get an isomorphism in Ab :

$$\text{Ext}_{\mathcal{R}}(A, B) \rightarrow \text{Ext}_{\mathcal{R}}(A/A_0, B_0) \oplus \text{Ext}_{\mathcal{R}}(A/A_0, B/B_0) \oplus \text{Ext}_{\mathcal{R}}(A_0, B/B_0) \oplus \text{Ext}_{\mathcal{R}}(A_0, B_0).$$

Using Lemma F.4.1.12, one can prove that $\text{Ext}_{\mathcal{R}}(A/A_0, B_0) = 0$. Identical to Example F.4.1.10, $\text{Ext}_{\mathcal{R}}(A/A_0, B/B_0) = \text{Ext}_{\mathbb{Z}}^1(A/A_0, B/B_0)$. Similar to Corollary F.3.2.5 and [Hoc51b, Thm. 3.2], $\text{Ext}_{\mathcal{R}}(A_0, B/B_0) = \text{Hom}_{\text{Ab}}(\pi_1(A_0), B/B_0)$. By Proposition F.9.0.1, $\text{Ext}_{\mathcal{R}}(A_0, B_0) = 0$. The proof is completed. \square

Bibliography

- [ABC20] Yves André, Francesco Baldassarri, and Maurizio Cailotto. *De Rham cohomology of differential modules on algebraic varieties*. Springer, 2nd edition, 2020.
- [Ach21] Pramod N Achar. *Perverse sheaves and applications to representation theory*, volume 258. American Mathematical Soc., 2021.
- [ADGK23] Perla Azzi, Rodrigue Desmorat, Julien Grivaux, and Boris Kolev. Rationality of normal forms of isotropy strata of a representation of a compact Lie group. <https://arxiv.org/pdf/2301.08599v1.pdf>, 2023.
- [AHR20] Pramod N Achar, William D Hardesty, and Simon Riche. Representation theory of disconnected reductive groups. *Documenta Mathematica*, 25:2149–2177, 2020.
- [AK01] Yukitaka Abe and Klaus Kopfermann. *Toroidal groups: line bundles, cohomology and quasi-Abelian varieties*. Springer, 2001.
- [AKO02] Yves André, Bruno Kahn, and Peter O’Sullivan. Nilpotence, radicaux et structures monoïdales. *Rendiconti del Seminario Matematico della Università di Padova*, 108:107–291, 2002.
- [And92] Yves André. Mumford-Tate groups of mixed Hodge structures and the theorem of the fixed part. *Compositio Mathematica*, 82(1):1–24, 1992.
- [And21] Yves André. Normality criteria, and monodromy of variations of mixed Hodge structures. *Tunisian Journal of Mathematics*, 3(3):645–656, 2021. Appendix C to Kahn, B. *Albanese kernels and Griffiths groups*.
- [Ara90] Donu Arapura. Hodge theory with local coefficients on compact varieties. *Duke Mathematical Journal*, 61(2):531–543, 1990.

- [Ari19] Oleg Aristov. Holomorphic functions of exponential type on connected complex Lie groups. *Journal of Lie Theory*, 29(4):1045–1070, 2019.
- [AT13] Daniele Angella and Adriano Tomassini. On the $\partial\bar{\partial}$ -lemma and Bott-Chern cohomology. *Inventiones mathematicae*, 192(1):71–81, 2013.
- [Ati57a] Michael Francis Atiyah. Complex analytic connections in fibre bundles. *Transactions of the American Mathematical Society*, 85(1):181–207, 1957.
- [Ati57b] Michael Francis Atiyah. Vector bundles over an elliptic curve. *Proceedings of the London Mathematical Society*, 3(1):414–452, 1957.
- [Ati69] Michael Atiyah. *Introduction to commutative algebra*. CRC Press, 1969.
- [B⁺87] Armand Borel et al. *Algebraic D-Modules*, volume 2 of *Perspectives in Mathematics*. Academic Press, 1987.
- [Bal10] Paul Balmer. Tensor triangular geometry. In *Proceedings of the International Congress of Mathematicians*, volume 2, pages 85–112, 2010.
- [Bar23] Owen Barrett. The singular support of an ℓ -adic sheaf. <https://arxiv.org/pdf/2309.02587v2.pdf>, 2023.
- [BB03] Alexei Bondal and Michel Van den Bergh. Generators and representability of functors in commutative and noncommutative geometry. *Moscow Mathematical Journal*, 3(1):1–36, 2003. doi:10.17323/1609-4514-2003-3-1-1-36.
- [BBBP07] Oren Ben-Bassat, Jonathan Block, and Tony Pantev. Non-commutative tori and Fourier–Mukai duality. *Compositio Mathematica*, 143(2):423–475, 2007.
- [BBDG82] Alexander Beilinson, Joseph Bernstein, Pierre Deligne, and Ofer Gabber. *Faisceaux pervers*. Société mathématique de France, 1982.
- [BBHM63] A Bialynicki-Birula, G Hochschild, and GD Mostow. Extensions of representations of algebraic linear groups. *American Journal of Mathematics*, 85(1):131–144, 1963.
- [BBR94] C Bartocci, U Bruzzo, and D Hernández Ruipérez. Fourier-Mukai transform and index theory. *manuscripta mathematica*, 85(1):141–163, 1994.

- [BC09] Olivier Brinon and Brian Conrad. CMI summer school notes on p -adic Hodge theory. <https://math.stanford.edu/~conrad/papers/notes.pdf>, 2009.
- [BC18] Patrick Brosnan and Timothy Y. Chow. Unit interval orders and the dot action on the cohomology of regular semisimple Hessenberg varieties. *Advances in Mathematics*, 329:955–1001, 2018.
- [BD21] Sébastien Boucksom and Simone Diverio. A note on Lang’s conjecture for quotients of bounded domains. *Épjournal de Géométrie Algébrique*, 5, 2021. doi:10.46298/epiga.2021.volume5.6050.
- [BD23] Indranil Biswas and Sorin Dumitrescu. Nonabelian Hodge theory for Fujiki class \mathcal{C} manifolds. *Proceedings American Mathematical Society*, 151(5):1937–1948, 2023.
- [BD24] Indranil Biswas and Sorin Dumitrescu. Principal bundles with holomorphic connections over a Kähler Calabi-Yau manifold, 2024. doi:10.1016/j.difgeo.2023.102093.
- [Bea92] Arnaud Beauville. Annulation du H^1 pour les fibrés en droites plats. In *Complex algebraic varieties*, pages 1–15. Springer, 1992.
- [Ber83] J Bernstein. Algebraic theory of D -modules. https://gauss.math.yale.edu/~il282/Bernstein_D_mod.pdf, 1983.
- [BF84] Jürgen Bingenier and Hubert Flenner. Constructible and quasiconstructible sheaves on analytic spaces. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, 54(1):119–139, 1984.
- [BGR84] Siegfried Bosch, Ulrich Güntzer, and Reinhold Remmert. *Non-archimedean analysis*, volume 261. Springer Berlin, 1984.
- [Bjö79] Jan-Erik Björk. *Rings of differential operators*, volume 21. North-Holland Mathematical Library, 1979.
- [Bjö93] Jan-Erik Björk. *Analytic \mathcal{D} -modules and applications*, volume 247. Springer Science & Business Media, 1993.
- [BK06] Gebhard Böckle and Chandrashekhara Khare. Mod ℓ representations of arithmetic fundamental groups II: A conjecture of A. J. de Jong. *Compositio Mathematica*, 142(2):271–294, 2006.

- [BK09] Jean-Benoît Bost and Klaus Künnemann. Hermitian vector bundles and extension groups on arithmetic schemes II. The arithmetic Atiyah extension. In Dai Xianzhe, Léandre Rémi, Xiaonan Ma, and Zhang Weiping, editors, *From probability to geometry (I) - Volume in honor of the 60th birthday of Jean-Michel Bismut*, number 327 in Astérisque. Société mathématique de France, 2009. URL: http://www.numdam.org/item/AST_2009__327__361_0/.
- [BL99] Christina Birkenhake and Herbert Lange. *Complex tori*. Springer, 1999.
- [BL04] Christina Birkenhake and Herbert Lange. *Complex abelian varieties*, volume 6. Springer, 2004.
- [Bla56] André Blanchard. Sur les variétés analytiques complexes. *Annales scientifiques de l'École Normale Supérieure*, 73(2):157–202, 1956.
- [Blo84] Spencer Bloch. The proof of the Mordell conjecture. *The mathematical intelligencer*, 6(2):41–47, 1984.
- [Blo10] J Block. Duality and equivalence of module categories in noncommutative geometry. In *A celebration of the mathematical legacy of Raoul Bott*, volume 50 of *CRM Proceedings & Lecture notes*, pages 311–339. Amer. Math. Soc., 2010.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*, volume 21. Springer Science & Business Media, 1990. doi:10.1007/978-3-642-51438-8.
- [BLSW17] Nero Budur, Yongqiang Liu, Luis Saumell, and Botong Wang. Cohomology support loci of local systems. *Michigan Mathematical Journal*, 66(2):295–307, 2017.
- [BN93] Marcel Bökstedt and Amnon Neeman. Homotopy limits in triangulated categories. *Compositio Mathematica*, 86(2):209–234, 1993.
- [Bon06] Juhani Bonsdorff. A Fourier transformation for Higgs bundles. *Journal für die reine und angewandte Mathematik*, 2006(591):21–48, 2006. doi:10.1515/CRELLE.2006.013.
- [Bon10] Juhani Bonsdorff. Autodual connection in the Fourier transform of a Higgs bundle. *Asian Journal of Mathematics*, 14(2):153–174, 2010.

- [Bor91] Armand Borel. *Linear algebraic groups*, volume 126. Springer Science & Business Media, 2nd edition, 1991.
- [Bou72] Nicolas Bourbaki. *Gropes et Algèbres de Lie: Chapitres 2 et 3*. Springer, 1972.
- [Bou07] Nicolas Bourbaki. *Variétés différentielles et analytiques: fascicule de résultats*, volume 8. Springer Science & Business Media, 2007.
- [BP01] Alexander Beilinson and Alexander Polishchuk. Torelli theorem via Fourier-Mukai transform. In *Moduli of Abelian Varieties*, pages 127–132. Springer, 2001.
- [Bre12] Glen E Bredon. *Sheaf theory*, volume 170. Springer Science & Business Media, 2012.
- [Bri98] Tom Bridgeland. Fourier-Mukai transforms for elliptic surfaces. *Journal für die reine und angewandte Mathematik*, 1998(498):115–133, 1998. doi:doi:10.1515/crll.1998.046.
- [Bri09] Michel Brion. Anti-affine algebraic groups. *Journal of Algebra*, 321(3):934–952, 2009.
- [Bro78] Robert Brody. Compact manifolds in hyperbolicity. *Transactions of the American Mathematical Society*, 235:213–219, 1978. doi:10.2307/1998216.
- [BS17] Indranil Biswas and Georg Schumacher. Line bundles and flat connections. *Proceedings-Mathematical Sciences*, 127(3):547–549, 2017.
- [BSS18] Bhargav Bhatt, Christian Schnell, and Peter Scholze. Vanishing theorems for perverse sheaves on abelian varieties, revisited. *Selecta Mathematica*, 24(1):63–84, 2018.
- [BT19] Benjamin Bakker and Jacob Tsimerman. The Ax-Schanuel conjecture for variations of Hodge structures. *Inventiones mathematicae*, 217:77–94, 2019.
- [BZ23a] Tatiana Bandman and Yuri G Zarhin. Automorphism groups of \mathbb{P}^1 -bundles over a non-uniruled base. *Russian Mathematical Surveys*, 78(1):1–64, 2023. doi:10.4213/rm10093e.
- [BZ23b] Tatiana Bandman and Yuri G. Zarhin. Simple complex tori of algebraic dimension 0. *Proc. Steklov Inst. Math.*, 320:27–45, 2023. doi:10.4213/tm4259.

- [Cam91] Frédéric Campana. The class \mathcal{C} is not stable by small deformations. *Mathematische Annalen*, 290(1):19–30, 1991.
- [Cam01] Frédéric Campana. Ensembles de Green-Lazarsfeld et quotients résolubles des groupes de Kähler. *Journal of Algebraic Geometry*, 10(4):599–622, 2001.
- [Car53] Henri Cartan. Un théoreme de finitude. *Séminaire Henri Cartan*, 6:1–11, 1953.
- [Car57] Henri Cartan. Variétés analytiques réelles et variétés analytiques complexes. *Bulletin de la Société Mathématique de France*, 85:77–99, 1957.
- [Cat83] Fabrizio Catanese. Moduli of surfaces of general type. In *Algebraic geometry—open problems*, pages 90–112. Springer, 1983.
- [Cat91] Fabrizio Catanese. Moduli and classification of irregular Kaehler manifolds (and algebraic varieties) with Albanese general type fibrations. *Inventiones mathematicae*, 104(1):263–289, 1991.
- [CD94] Frédéric Campana and G Dethloff. *Several complex variables VII: sheaf-theoretical methods in complex analysis*, volume 74. Springer Science & Business Media, 1994.
- [CE99] Henry Cartan and Samuel Eilenberg. *Homological algebra*, volume 19 of *Princeton Mathematical Series*. Princeton university press, 1999.
- [Che46] Claude Chevalley. *Theory of Lie groups*. Princeton University Press, 1946.
- [Che09] Ke Chen. *Special subvarieties of mixed Shimura varieties*. PhD thesis, Université Paris-Sud, 2009. http://www.numdam.org/item/BJHTUP11_2009__0773__A1_0/.
- [Che22] Xuemiao Chen. Admissible Hermitian-Yang-Mills connections over normal varieties, 2022. <https://arxiv.org/abs/2205.12305v3>.
- [CHM97] Lucia Caporaso, Joe Harris, and Barry Mazur. Uniformity of rational points. *Journal of the American Mathematical Society*, 10(1):1–35, 1997.
- [CK16] Anna Cadoret and Arno Kret. Galois-generic points on Shimura varieties. *Algebra & Number Theory*, 10(9):1893–1934, 2016.

- [CLZ16] Ke Chen, Xin Lu, and Kang Zuo. On the Oort conjecture for Shimura varieties of unitary and orthogonal types. *Compositio Mathematica*, 152(5):889–917, 2016.
- [Con00] Brian Conrad. *Grothendieck duality and base change*. Springer, 2000.
- [Con06] Brian Conrad. Relative ampleness in rigid geometry. *Annales de l’institut Fourier*, 56(4):1049–1126, 2006.
- [Con14] Brian Conrad. Reductive group schemes. In *Autour des schémas en groupes I*, volume 42-43 of *Panoramas et Synthèses*, pages 93–444. Société Mathématique de France Paris, 2014.
- [Cos05] Izzet Coskun. The arithmetic and geometry of Kobayashi hyperbolicity. In *Snowbird Lectures in Algebraic Geometry*, volume 388 of *Contemporary Mathematics*, pages 77–88. American Mathematical Society, 2005.
- [COUV16] Manuel Ceballos, Antonio Otal, Luis Ugarte, and Raquel Villacampa. Invariant complex structures on 6-nilmanifolds: classification, Frölicher spectral sequence and special Hermitian metrics. *The Journal of Geometric Analysis*, 26(1):252–286, 2016.
- [Day23] Mathieu Daylies. Base change for coherent cohomology in Berkovich geometry. <https://arxiv.org/pdf/2211.17001v2.pdf>, 2023.
- [dCM03] Mark Andrea A. de Cataldo and Luca Migliorini. The Hodge theory of algebraic maps. <https://arxiv.org/pdf/math/0306030v1.pdf>, 2003.
- [dCM05] Mark Andrea A. de Cataldo and Luca Migliorini. The Hodge theory of algebraic maps. *Annales scientifiques de l’Ecole normale supérieure*, 38(5):693–750, 2005.
- [dCM09] Mark Andrea A. de Cataldo and Luca Migliorini. The decomposition theorem, perverse sheaves and the topology of algebraic maps. *Bulletin of the American Mathematical Society*, 46(4):535–633, 2009.
- [Deb05] Olivier Debarre. *Complex tori and abelian varieties*, volume 11 of *SMF/AMS Texts and Monographs*. American Mathematical Soc., 2005.
- [Del68] Pierre Deligne. Théorème de Lefschetz et critères de dégénérescence de suites spectrales. *Publications*

- Mathématiques de l'Institut des Hautes Études Scientifiques*, 35(1):107–126, 1968.
- [Del70] Pierre Deligne. *Équations différentielles à points singuliers réguliers*, volume 163. Springer, 1970.
- [Del71a] Pierre Deligne. Théorie de Hodge, II. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques*, 40(1):5–57, 1971.
- [Del71b] Pierre Deligne. Travaux de Shimura. In *Séminaire Bourbaki vol. 1970/71 Exposés 382–399*, pages 123–165. Springer, 1971.
- [Del74] Pierre Deligne. La conjecture de Weil. I. *Publications Mathématiques de l'Institut des Hautes Études Scientifiques*, 43(1):273–307, 1974.
- [Del77] P Deligne. *Séminaire de Géométrie Algébrique du Bois Marie-Cohomologie étale (SGA 4 1/2)*. Springer, Berlin; New York, 1977. doi:10.1007/BFb0091516.
- [Del78] Pierre Deligne. Extensions centrales non résiduellement finies de groupes arithmétiques. *Comptes rendus mathématiques de l'Académie des sciences*, 287:203–208, 1978.
- [Del90] Pierre Deligne. Catégories tannakiennes. In *The Grothendieck Festschrift II*, pages 111–195. Springer, 1990.
- [Del02] Pierre Deligne. Catégories tensorielles. *Mosc. Math. J*, 2(2):227–248, 2002. doi:10.17323/1609-4514-2002-2-2-227-248.
- [Dem12] Jean-Pierre Demailly. Complex analytic and differential geometry. <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>, 2012.
- [DGMS75] Pierre Deligne, Phillip Griffiths, John Morgan, and Dennis Sullivan. Real homotopy theory of Kähler manifolds. *Inventiones mathematicae*, 29(3):245–274, 1975.
- [Dim04] Alexandru Dimca. *Sheaves in topology*. Springer Science & Business Media, 2004.
- [DJK20] Christopher Daw, Ariyan Javanpeykar, and Lars Kühne. Effective estimates for the degrees of maximal special subvarieties. *Selecta Mathematica*, 26:1–31, 2020.
- [DM22] Pierre Deligne and JS Milne. Tannakian categories. <https://www.jmilne.org/math/>, 2022.

- [Dou66] Adrien Douady. Le problème des modules pour les sous-espaces analytiques compacts d'un espace analytique donné. *Annales de l'Institut Fourier*, 16(1):1–95, 1966. doi:10.5802/aif.226.
- [DR10] Michael Dettweiler and Stefan Reiter. Rigid local systems and motives of type G_2 . with an appendix by Michale Dettweiler and Nicholas M. Katz. *Compositio Mathematica*, 146(4):929–963, 2010. doi:10.1112/S0010437X10004641.
- [DR16] Simone Diverio and Erwan Rousseau. The exceptional set and the Green-Griffiths locus do not always coincide. *L'Enseignement Mathématique*, 61(3):417–452, 2016. doi:10.4171/LEM/61-3/4-6.
- [Dri01] Vladimir Drinfeld. On a conjecture of Kashiwara. *Mathematical Research Letters*, 8(6):713–728, 2001.
- [Eis95] David Eisenbud. *Commutative algebra: with a view toward algebraic geometry*, volume 150. Springer Science & Business Media, 1995. doi:10.1007/978-1-4612-5350-1.
- [EM47] Samuel Eilenberg and Saunders MacLane. Cohomology theory in abstract groups II: Group extensions with a non-abelian kernel. *Annals of Mathematics*, 48(2):326–341, 1947. doi:10.2307/1969174.
- [Eng95] Ryszard Engelking. Theory of dimensions, finite and infinite. *Sigma Series in Pure Mathematics*, 10, 1995.
- [Enr39] Federigo Enriques. Des courbes paracanonicques appartenant à une surface algébrique irrégulière. *Bulletin de la Société Royale des Sciences de Liège*, 8:422–423, 1939. <http://dspace.bcu-iasi.ro/handle/123456789/44377>.
- [Enr49] Federigo Enriques. *Le superficie algebriche*. N. Zanichelli, 1949.
- [EP⁺96] Jörg Eschmeier, Mihai Putinar, et al. *Spectral decompositions and analytic sheaves*. Number 10 in London Mathematical Society monographs. Clarendon Press Press, 1996.
- [EV89] Hélène Esnault and Eckart Viehweg. Vanishing and non-vanishing theorems. In Barlet D., Esnault H., Elzein F., Verdier Jean-Louis, and Viehweg E., editors, *Théorie de Hodge - Luminy, Juin 1987*, number 179-180 in Astérisque. Société mathématique de France, 1989. URL: http://www.numdam.org/item/AST_1989__179-180__97_0/.

- [EY03] Bas Edixhoven and Andrei Yafaev. Subvarieties of Shimura varieties. *Annals of Mathematics*, 157(2):621–645, 2003.
- [Fal82] G Faltings. Arithmetic varieties and rigidity. In *Seminar on number theory, Paris*, pages 63–77, 1982.
- [Fal83] Gerd Faltings. Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. *Inventiones mathematicae*, 73(3):349–366, 1983.
- [Fal88] Gerd Faltings. Crystalline cohomology and p -adic Galois representations. In *Algebraic analysis, geometry, and number theory*, pages 25–80. Johns Hopkins University Press, 1988.
- [Fal91] Gerd Faltings. Diophantine approximation on abelian varieties. *Annals of Mathematics*, 133(3):549–576, 1991. doi:10.2307/2944319.
- [Fal94] Gerd Faltings. The general case of S. Lang’s conjecture. In *Barsotti Symposium in Algebraic Geometry*, pages 175–182, 1994.
- [Far08] Jaques Faraut. *Analysis of Lie Groups*. Cambridge University Press Cambridge, 2008. doi:10.1017/CB09780511755170.
- [Fav12] David Favero. Reconstruction and finiteness results for Fourier-Mukai partners. *Advances in Mathematics*, 230(4-6):1955–1971, 2012.
- [Fis76] Gerd Fischer. *Complex analytic geometry*, volume 538. Springer, 1976. doi:10.1007/BFb0080338.
- [FJJ⁺71] D Ferrand, Jean-Pierre Jouanolou, O Jussila, S Kleiman, M Raynaud, and Jean-Pierre Serre. *Théorie des Intersections et Théorème de Riemann-Roch: Séminaire de Géométrie Algébrique du Bois Marie 1966/67 (SGA 6)*, volume 225. Springer, 1971. doi:10.1007/BFb0066283.
- [FK88] Eberhard Freitag and Reinhardt Kiehl. *Etale cohomology and the Weil conjecture*, volume 13. Springer Science & Business Media, 1988. doi:10.1007/978-3-662-02541-3.
- [FL14] Carlos Florentino and Thomas Ludsteck. Unipotent Schottky bundles on Riemann surfaces and complex tori. *International Journal of Mathematics*, 25(06):1450056, 2014. doi:10.1142/S0129167X14500566.
- [FLA19] Mathieu Florence and Giancarlo Lucchini Arteche. On extensions of algebraic groups. *L’Enseignement Mathématique*, 65(3):441–455, 2019.

- [Fri67] Jacques Frisch. Points de platitude d'un morphisme d'espaces analytiques complexes. *Inventiones mathematicae*, 4:118–138, 1967.
- [FS13] Teresa Monteiro Fernandes and Claude Sabbah. On the de Rham complex of mixed twistor \mathcal{D} -modules. *International Mathematics Research Notices*, 2013(21):4961–4984, 2013.
- [Fuj78] Akira Fujiki. Closedness of the Douady spaces of compact Kähler spaces. *Publications of the Research Institute for Mathematical Sciences*, 14(1):1–52, 1978.
- [Gab62] P. Gabriel. Des catégories abéliennes. *Bulletin de la Société Mathématique de France*, 90:323–448, 1962. doi:10.24033/bsmf.1583.
- [Gai07] Dennis Gaitsgory. On de Jong's conjecture. *Israel Journal of Mathematics*, 157(1):155–191, 2007.
- [GD71] A. Grothendieck and J.A. Dieudonné. *Eléments de Géométrie Algébrique I*, volume 166 of *Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete*. Springer Berlin Heidelberg, 1971.
- [gh] gdb (<https://mathoverflow.net/users/115211/gdb>). Compare bounded (unbounded) derived categories of D -modules. <https://mathoverflow.net/q/361206>.
- [GH78] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. John Wiley & Sons, 1978.
- [Gil11] William D Gillam. Localization of ringed spaces. *Advances in Pure Mathematics*, 1(5):250–263, 2011. doi:10.4236/apm.2011.15045.
- [GL87] Mark Green and Robert Lazarsfeld. Deformation theory, generic vanishing theorems, and some conjectures of Enriques, Catanese and Beauville. *Inventiones mathematicae*, 90(2):389–407, 1987.
- [GL91] Mark Green and Robert Lazarsfeld. Higher obstructions to deforming cohomology groups of line bundles. *Journal of the American Mathematical Society*, 4(1):87–103, 1991.
- [GL96] Ofer Gabber and François Loeser. Faisceaux pervers ℓ -adiques sur un tore. *Duke Mathematical Journal*, 83(3):501–606, 1996.

- [GM88] Mark Goresky and Robert MacPherson. *Stratified Morse theory*, volume 14 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, 1988. doi:10.1007/978-3-642-71714-7.
- [GN20] A Genestier and BC Ngô. Lectures on Shimura varieties. In *Shimura varieties*, pages 22–71. Cambridge University Press, 2020.
- [God58] Roger Godement. Topologie algébrique et théorie des faisceaux. *Publications de l'Institut de Mathématique de L'Université de Strasbourg XIII*, 1, 1958.
- [GR71] Alexander Grothendieck and Michele Raynaud. *Revêtements étales et groupe fondamental (SGA 1)*. Springer-Verlag, 1971. doi:10.1007/BFb0058656.
- [GR84] Hans Grauert and Reinhold Remmert. *Coherent analytic sheaves*, volume 265. Springer Science & Business Media, 1984.
- [GR04] Hans Grauert and Reinhold Remmert. *Theory of Stein spaces*, volume 236. Springer Science & Business Media, 2004. doi:10.1007/978-3-642-18921-0.
- [GR14] Dennis Gaitsgory and Nick Rozenblyum. Crystals and D -modules. *Pure and Applied Mathematics Quarterly*, 10(1):57–154, 2014.
- [Gro55] A Grothendieck. *Produits tensoriels topologiques et espaces nucléaires*. Number 16 in *Memoirs of the American Mathematical Society*. American Mathematical Society, 1955.
- [Gro57a] Alexander Grothendieck. Sur la classification des fibrés holomorphes sur la sphere de Riemann. *American Journal of Mathematics*, 79(1):121–138, 1957.
- [Gro57b] Alexandre Grothendieck. Sur quelques points d'algèbre homologique. *Tohoku Mathematical Journal, Second Series*, 9(2):119–183, 1957.
- [Gro60] Alexander Grothendieck. Éléments de géométrie algébrique : I. Le langage des schémas (EGA I). *Publications Mathématiques de l'IHÉS*, 4:5–228, 1960. URL: http://www.numdam.org/item/PMIHES_1960__4__5_0.
- [Gro61a] Alexander Grothendieck. Techniques de construction en géométrie analytique. II. Généralités sur les espaces annelés et les espaces analytiques. *Séminaire Henri Cartan*, 13(1), 1960-1961. talk:9. URL: http://www.numdam.org/item/SHC_1960-1961__13_1_A5_0/.

- [Gro61b] Alexander Grothendieck. Techniques de construction en géométrie analytique. VI. étude locale des morphismes : germes d'espaces analytiques, platitude, morphismes simples. *Séminaire Henri Cartan*, 13(1), 1960-1961. talk:13. URL: http://www.numdam.org/item/SHC_1960-1961__13_1_A9_0/.
- [Gro61c] Alexander Grothendieck. Éléments de géométrie algébrique : II. Étude globale élémentaire de quelques classes de morphismes. *Publications Mathématiques de l'IHÉS*, 8:5–222, 1961. URL: http://www.numdam.org/item/PMIHES_1961__8__5_0/.
- [Gro65] Alexander Grothendieck. Éléments de géométrie algébrique : IV. Étude locale des schémas et des morphismes de schémas, Seconde partie. *Publications Mathématiques de l'IHÉS*, 24:5–231, 1965. URL: http://www.numdam.org/item/PMIHES_1965__24__5_0/.
- [Gro66] Alexander Grothendieck. Éléments de géométrie algébrique : IV. étude locale des schémas et des morphismes de schémas, Troisième partie. *Publications Mathématiques de l'IHÉS*, 28:5–255, 1966. URL: http://www.numdam.org/item/PMIHES_1966__28__5_0/.
- [Gro06] Frank D Grosshans. *Algebraic homogeneous spaces and invariant theory*. Springer, 2006.
- [GRR15] Alberto García-Raboso and Steven Rayan. Introduction to nonabelian Hodge theory. In *Calabi-Yau varieties: arithmetic, geometry and physics*, pages 131–171. Springer, 2015.
- [GSK09] Jon González-Sánchez and Benjamin Klopsch. Analytic pro- p groups of small dimensions. *Journal of Group Theory*, 12(5):711–734, 2009. doi:10.1515/JGT.2009.006.
- [GW20] Ulrich Görtz and Torsten Wedhorn. *Algebraic Geometry I: Schemes*. Springer, 2nd edition, 2020.
- [Hac04] Christopher D Hacon. A derived category approach to generic vanishing. *Journal für die reine und angewandte Mathematik*, 575:173–187, 2004. doi:10.1515/crll.2004.078.
- [Hal23] Jack Hall. GAGA theorems. *Journal de Mathématiques Pures et Appliquées*, 175:109–142, 2023. doi:10.1016/j.matpur.2023.05.004.
- [Har66] Robin Hartshorne. *Residues and duality*, volume 20. Springer, 1966. doi:10.1007/BFb0080482.

- [Har70] Robin Hartshorne. *Ample subvarieties of algebraic varieties*, volume 156. Springer, 1970. doi:10.1007/BFb0067839.
- [Har77] Robin Hartshorne. *Algebraic geometry*, volume 52. Springer Science & Business Media, 1977.
- [Hat05] Allen Hatcher. *Algebraic topology*. Cambridge University Press, 2005.
- [HBS66] Friedrich Hirzebruch, Armand Borel, and RLE Schwarzenberger. *Topological methods in algebraic geometry*, volume 175. Springer Berlin-Heidelberg-New York, 1966.
- [Hin98] Marc Hindry. Introduction to abelian varieties and the Mordell-Lang conjecture. In *Model Theory and Algebraic Geometry: An introduction to E. Hrushovski's proof of the geometric Mordell-Lang conjecture*, pages 85–100. Springer, 1998.
- [Hir60] Heisuke Hironaka. *On the theory of birational blowing-up*. PhD thesis, Harvard University, 1960.
- [Hir77] Heisuke Hironaka. Stratification and flatness. In *Real and complex singularities*, pages 199–265, 1977.
- [HK84] Ryoshi Hotta and Masaki Kashiwara. The invariant holonomic system on a semisimple Lie algebra. *Inventiones mathematicae*, 75(2):327–358, 1984.
- [HN11] Joachim Hilgert and Karl-Hermann Neeb. *Structure and geometry of Lie groups*. Springer Science & Business Media, 2011.
- [Hoc51a] G. Hochschild. Group extensions of Lie groups. *Annals of Mathematics*, 54(1):96–109, 1951. doi:10.2307/1969314.
- [Hoc51b] G Hochschild. Group extensions of Lie groups II. *Annals of Mathematics*, 54(3):537–551, 1951. doi:10.2307/1969488.
- [Hov99] Mark Hovey. Model category structures on chain complexes of sheaves, 1999. <https://arxiv.org/abs/math/9909024v1>. arXiv:math/9909024.
- [HRS23] Tamir Hemo, Timo Richarz, and Jakob Scholbach. Constructible sheaves on schemes. *Advances in Mathematics*, 429:109179, 2023. doi:10.1016/j.aim.2023.109179.
- [hs] Jason Starr (<https://mathoverflow.net/users/13265/jason-starr>). Is an infinite direct sum of quasi-coherent

- \mathcal{O}_X -modules quasi-coherent on a complex manifold?
<https://mathoverflow.net/q/453047> (version: 2023-08-18).
- [HS00] Marc Hindry and Joseph H Silverman. *Diophantine geometry: an introduction*, volume 201. Springer Science & Business Media, 2000.
 - [HS23] David Hansen and Peter Scholze. Relative perversity. *Comm. Amer. Math. Soc.*, 3:631–668, 2023. doi:10.1090/cams/21.
 - [HT07] Ryoshi Hotta and Toshiyuki Tanisaki. *D-modules, perverse sheaves, and representation theory*, volume 236. Springer Science & Business Media, 2007.
 - [Huy05] Daniel Huybrechts. *Complex geometry: an introduction*, volume 78. Springer, 2005.
 - [Huy06] Daniel Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Clarendon Press, 2006.
 - [Ill06] Luc Illusie. Vanishing cycles over general bases after P. Deligne, O. Gabber, G. Laumon and F. Orgogozo. 数理解析研究所研究, 1521:35–53, 2006. <http://hdl.handle.net/2433/58796>.
 - [Iwa49] Kenkichi Iwasawa. On some types of topological groups. *Annals of Mathematics*, 50(3):507–558, 1949. doi:10.2307/1969548.
 - [JKLM23] Ariyan Javanpeykar, Thomas Krämer, Christian Lehn, and Marco Maculan. The monodromy of families of subvarieties on abelian varieties, 2023. arXiv:2210.05166v2.
 - [JM22] Jia Jia and Sheng Meng. Moishezon manifolds with no nef and big classes. <https://arxiv.org/pdf/2208.12013v1.pdf>, 2022.
 - [Kas98] Masaki Kashiwara. Semisimple holonomic \mathcal{D} -modules. In *Topological field theory, primitive forms and related topics*, pages 267–271. Springer, 1998.
 - [Kas03] Masaki Kashiwara. *D-modules and microlocal calculus*, volume 217. American Mathematical Soc., 2003.
 - [Kas04] M Kashiwara. t -structures on the derived categories of holonomic \mathcal{D} -modules and coherent \mathcal{O} -modules. *Moscow Mathematical Journal*, 4(4):847–868, 2004. doi:10.17323/1609-4514-2004-4-4-847-868.

- [Kat88] Nicholas M. Katz. *Gauss Sums, Kloosterman Sums, and Monodromy Groups*. Princeton University Press, 1988. URL: <http://www.jstor.org/stable/j.ctt1b7x80x>.
- [Kat90] Nicholas M Katz. *Exponential sums and differential equations*, volume 124 of *Annals of Mathematics Studies*. Princeton University Press, 1990.
- [Kem91] George R Kempf. *Complex abelian varieties and theta functions*. Springer-Verlag Berlin, 1991.
- [KK83] Ludger Kaup and Burchard Kaup. *Holomorphic functions of several variables: an introduction to the fundamental theory*, volume 3. Walter de Gruyter, 1983. doi:10.1515/9783110838350.
- [KKN08] Takeshi Kajiwarra, Kazuya Kato, and Chikara Nakayama. Analytic log Picard varieties. *Nagoya Mathematical Journal*, 191:149–180, 2008.
- [KL85] Nicholas M Katz and Gérard Laumon. Transformation de Fourier et majoration de sommes exponentielles. *Publications Mathématiques de l’IHÉS*, 62:145–202, 1985.
- [KM23] Thomas Krämer and Marco Maculan. Arithmetic finiteness of very irregular varieties, 2023. <https://arxiv.org/abs/2310.08485v2>.
- [Kob87] Shoshichi Kobayashi. *Differential geometry of complex vector bundles*, volume 793. Princeton University Press, 1987.
- [Kob98] Shoshichi Kobayashi. *Hyperbolic complex spaces*, volume 318. Springer Science & Business Media, 1998.
- [Kod64] Kunihiko Kodaira. On the structure of compact complex analytic surfaces, I. *American Journal of Mathematics*, 86(4):751–798, 1964.
- [Krä21] Thomas Krämer. Characteristic cycles and the microlocal geometry of the Gauss map, II. *Journal für die reine und angewandte Mathematik*, 2021(774):53–92, 2021.
- [Krä22] Thomas Krämer. Characteristic cycles and the microlocal geometry of the Gauss map, I. *Annales Scientifiques de l’École Normale Supérieure*, 55(6):1475–1527, 2022.
- [KRL19] Nicholas M Katz and Antonio Rojas-León. A rigid local system with monodromy group $2.J_2$. *Finite Fields and Their Applications*, 57:276–286, 2019.

- [KRW20] Manuel Krannich and Oscar Randal-Williams. Mapping class groups of simply connected high-dimensional manifolds need not be arithmetic. *Comptes Rendus. Mathématique*, 358(4):469–473, 2020.
- [KS90] Masaki Kashiwara and Pierre Schapira. *Sheaves on Manifolds*, volume 292. Springer Science & Business Media, 1990. doi:10.1007/978-3-662-02661-8.
- [KS96] Masaki Kashiwara and Pierre Schapira. *Moderate and formal cohomology associated with constructible sheaves*. Number 64 in Mémoires de la Société Mathématique de France. Société mathématique de France, 1996. URL: http://www.numdam.org/item/MSMF_1996_2_64__1_0/, doi:10.24033/msmf.378.
- [KS97] Masaki Kashiwara and Pierre Schapira. Integral transforms with exponential kernels and Laplace transform. *Journal of the American Mathematical Society*, 10(4):939–972, 1997.
- [KS06] Masaki Kashiwara and Pierre Schapira. *Categories and Sheaves*, volume 332. Springer Berlin, Heidelberg, 2006. doi:10.1007/3-540-27950-4.
- [KS16] Masaki Kashiwara and Pierre Schapira. *Regular and irregular holonomic D-modules*, volume 433. Cambridge University Press, 2016.
- [KT19a] Nicholas M Katz and Pham Huu Tiep. Local systems and finite unitary and symplectic groups. *Advances in Mathematics*, 358:106859, 2019. doi:10.1016/j.aim.2019.106859.
- [KT19b] Nicholas M Katz and Pham Huu Tiep. Rigid local systems and finite symplectic groups. *Finite Fields and Their Applications*, 59:134–174, 2019.
- [KUY18] Bruno Klingler, Emmanuel Ullmo, and Andrei Yafaev. Bi-algebraic geometry and the André-Oort conjecture. In *Algebraic geometry: Salt Lake City 2015*, volume 97, pages 319–359. American Mathematical Society Providence, 2018.
- [KW01] Reinhardt Kiehl and Rainer Weissauer. *Weil conjectures, perverse sheaves and l'adic Fourier transform*, volume 42. Springer Science & Business Media, 2001.
- [KW15a] Thomas Krämer and Rainer Weissauer. On the Tannaka group attached to the theta divisor of a generic principally polarized abelian variety. *Mathematische Zeitschrift*, 281(3):723–745, 2015.

- [KW15b] Thomas Krämer and Rainer Weissauer. Vanishing theorems for constructible sheaves on abelian varieties. *Journal of Algebraic Geometry*, 24(3):531–568, 2015. doi:10.1090/jag/645.
- [KY14] Bruno Klingler and Andrei Yafaev. The André-Oort conjecture. *Annals of mathematics*, 180(3):867–925, 2014.
- [Lan74] Serge Lang. Higher dimensional Diophantine problems. *Bulletin of the American Mathematical Society*, 80(5):779 – 787, 1974.
- [Lan86] Serge Lang. Hyperbolic and Diophantine analysis. *Bulletin of the American Mathematical Society*, 14(2):159–205, 1986.
- [Lan91] Serge Lang. *Number theory III: Diophantine geometry*, volume 60 of *Encyclopaedia of Mathematical Sciences*. Springer Science & Business Media, 1991. doi:10.1007/978-3-642-58227-1.
- [Lan02] Serge Lang. *Algebra*. Springer, 3rd edition, 2002.
- [Lau96] Gérard Laumon. Transformation de Fourier généralisée. <https://arxiv.org/pdf/alg-geom/9603004v1.pdf>, 1996.
- [Laz54] Michel Lazard. Sur les groupes nilpotents et les anneaux de Lie. *Annales scientifiques de l'École Normale Supérieure*, 71(2):101–190, 1954.
- [Laz04] Robert K Lazarsfeld. *Positivity in algebraic geometry I: Classical setting: line bundles and linear series*, volume 48. Springer, 2004. doi:10.1007/978-3-642-18808-4.
- [Lee01] Dong Hoon Lee. *The structure of complex Lie groups*. Chapman and Hall/CRC, 2001.
- [Lee13] John M Lee. *Introduction to smooth manifolds*. Springer, 2nd edition, 2013.
- [Len68] Helmut Lenzing. Direkte Produkte von freien Moduln. *Mathematische Zeitschrift*, 106:206–212, 1968.
- [IH76] P de la Harpe. Introduction to complex tori. In *Complex analysis and its applications Vol. II*, pages 101–144. International Atomic Energy Agency, Vienna, 1976.
- [Lip60] Joseph Lipman. *Notes on derived functors and Grothendieck duality*, pages 1–259. Springer, 1960.
- [Liu06] Qing Liu. *Algebraic geometry and arithmetic curves*. Oxford University Press, 2nd edition, 2006.

- [LS20] Brian Lawrence and Will Sawin. The Shafarevich conjecture for hypersurfaces in abelian varieties, 2020. <https://arxiv.org/abs/2004.09046v2>.
- [LTZZ10] Steven Lu, Yuping Tu, Qi Zhang, and Quan Zheng. On semistability of Albanese maps. *Manuscripta Mathematica*, 131(3):531–535, 2010.
- [LV20] Brian Lawrence and Akshay Venkatesh. Diophantine problems and p -adic period mappings. *Inventiones mathematicae*, 221(3):893–999, 2020. doi:10.1007/s00222-020-00966-7.
- [LZ19] Xin Lu and Kang Zuo. The Oort conjecture on Shimura curves in the Torelli locus of curves. *Journal de Mathématiques Pures et Appliquées*, 123:41–77, 2019. doi:10.1016/j.matpur.2019.01.001.
- [Mac60] RA Macauley. Analytic group kernels and Lie algebra kernels. *Transactions of the American Mathematical Society*, 95(3):530–553, 1960.
- [Mal40] Anatolii Malcev. On isomorphic matrix representations of infinite groups. *Matematicheskii Sbornik*, 50(3):405–422, 1940.
- [Mal42] A Malcev. On the simple connectedness of invariant subgroups of Lie groups. *Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS, Nouvelle Série*, 34:10–13, 1942.
- [Mal94] Bernard Malgrange. Filtration des modules holonomes. In *Analyse algébrique des perturbations singulières*. Hermann, 1994.
- [Mal96] Bernard Malgrange. Connexions méromorphes 2 Le réseau canonique. *Inventiones mathematicae*, 124(1):367–387, 1996.
- [Mal04] Bernard Malgrange. On irregular holonomic D -modules. In *Éléments de la théorie des systèmes différentiels géométriques*, volume 8, pages 391–410. Société mathématique de France, 2004.
- [Mat59] Yozô Matsushima. Fibres holomorphes sur un tore complexe. *Nagoya Mathematical Journal*, 14:1–24, 1959.
- [Mat87] Hideyuki Matsumura. *Commutative Ring Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1987. doi:10.1017/CB09781139171762.
- [Mat12] John N. Mather. Notes on topological stability. *Bulletin of the American Mathematical Society*, 49:475–506, 2012.

- [Max19] Laurențiu Maxim. *Intersection Homology & Perverse Sheaves: With Applications to Singularities*. Springer, 2019.
- [MD91] Jacob Murre and Ch Deninger. Motivic decomposition of abelian schemes and the Fourier transform. *Journal für die reine und angewandte Mathematik*, 1991(422):201–219, 1991.
- [Meb89] Zoghman Mebkhout. *Le formalisme des six opérations de Grothendieck pour les D_X -modules cohérents*. Hermann, Paris, 1989.
- [Men20] Lingxu Meng. Leray-Hirsch theorem and blow-up formula for Dolbeault cohomology. *Annali di Matematica Pura ed Applicata (1923-)*, 199(5):1997–2014, 2020.
- [Mes73] William Messing. The universal extension of an abelian variety by a vector group. In *Symposia Mathematica*, volume 11, pages 359–372, 1973.
- [Mil80] James S Milne. *Étale Cohomology*. Princeton university press, 1980. doi:10.1515/9781400883981.
- [Mil99] JS Milne. Descent for Shimura varieties. *Michigan Mathematical Journal*, 46(1):203–208, 1999.
- [Mil07] James S Milne. Quotients of Tannakian categories. *Theory and Applications of Categories*, 18(21):654–664, 2007.
- [Mil13] James S. Milne. Lectures on Étale cohomology (v2.21). <https://www.jmilne.org/math/CourseNotes/LEC.pdf>, 2013.
- [Mil17a] James S Milne. *Algebraic groups: the theory of group schemes of finite type over a field*, volume 170. Cambridge University Press, 1st edition, 2017.
- [Mil17b] James S Milne. Introduction to Shimura varieties (revised version). <https://www.jmilne.org/math/>, 2017.
- [Mil20] Joseph S Milne. Complex multiplication, July 2020. Available at <https://www.jmilne.org/math/CourseNotes/>.
- [ML78] Saunders Mac Lane. *Categories for the working mathematician*, volume 5. Springer Science & Business Media, 1978. doi:10.1007/978-1-4757-4721-8.
- [MM60] Yozō Matsushima and Akihiko Morimoto. Sur certains espaces fibrés holomorphes sur une variété de Stein. *Bulletin de la Société Mathématique de France*, 88:137–155, 1960.

- [MM66] Hideyuki Matsumura and Masayoshi Miyanishi. On some dualities concerning abelian varieties. *Nagoya Mathematical Journal*, 27(2):447–462, 1966.
- [MM07] Xiaonan Ma and George Marinescu. *Holomorphic Morse inequalities and Bergman kernels*, volume 254. Springer Science & Business Media, 2007.
- [Moc12] Shinichi Mochizuki. Topics in absolute anabelian geometry I: generalities. *J. Math. Sci. Univ. Tokyo*, 19(2):139–242, 2012.
- [Moo98a] Ben Moonen. Linearity properties of Shimura varieties, I. *Journal of algebraic geometry*, 7(3):539–568, 1998.
- [Moo98b] Ben Moonen. Models of Shimura varieties in mixed characteristics. In *Galois representations in arithmetic algebraic geometry*, pages 267–350. Cambridge University Press, 1998.
- [Mor22] L. J. Mordell. On the rational solutions of the indeterminate equation of the third and fourth degree. *Proc. Camb. Phil. Soc.*, 21:179–192, 1922. <https://ia800202.us.archive.org/23/items/proceedingscambr21camb/proceedingscambr21camb.pdf>.
- [Mor59] Akihiko Morimoto. Sur la classification des espaces fibrés vectoriels holomorphes sur un tore complexe admettant des connexions holomorphes. *Nagoya Mathematical Journal*, 15:83–154, 1959.
- [Mor95] Atsushi Moriwaki. Remarks on rational points of varieties whose cotangent bundles are generated by global sections. *Mathematical Research Letters*, 2(1):113–118, 1995. doi:10.4310/MRL.1995.v2.n1.a10.
- [MRM74] David Mumford, Chidambaram Padmanabhan Ramanujam, and Jurij Ivanovič Manin. *Abelian varieties*, volume 5. Oxford university press Oxford, 2nd edition, 1974.
- [MS08] Emanuele Macrì and Paolo Stellari. Automorphisms and autoequivalences of generic analytic K3 surfaces. *Journal of Geometry and Physics*, 58(1):133–164, 2008.
- [Muk78] Shigeru Mukai. Semi-homogeneous vector bundles on an abelian variety. *Journal of Mathematics of Kyoto University*, 18(2):239–272, 1978.
- [Muk81] Shigeru Mukai. Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves. *Nagoya Mathematical Journal*, 81:153–175, 1981.

- [Mun00] James R Munkres. *Topology*, volume 2. Prentice Hall Upper Saddle River, 2nd edition, 2000.
- [Mur06] Daniel Murfet. Derived categories of sheaves, 2006. <http://therisingsea.org/notes/DerivedCategoriesOfSheaves.pdf>.
- [Mur07] Daniel Murfet. Triangulated categories part I. <http://therisingsea.org/notes/TriangulatedCategories.pdf>, 2007.
- [Nad89] Alan Michael Nadel. The nonexistence of certain level structures on abelian varieties over complex function fields. *Annals of Mathematics*, 129(1):161–178, 1989.
- [Nak94] Atsushi Nakayashiki. Commuting partial differential operators and vector bundles over abelian varieties. *American Journal of Mathematics*, 116(1):65–100, 1994.
- [Nee96] Amnon Neeman. The Grothendieck duality theorem via Bousfield’s techniques and Brown representability. *Journal of the American Mathematical Society*, 9(1):205–236, 1996.
- [Nee21] Amnon Neeman. Triangulated categories with a single compact generator and a Brown representability theorem, 2021. arXiv: 1804.02240v4.
- [Nic13] Bogdan Nica. Linear groups-Malcev’s theorem and Selberg’s lemma, 2013. <https://arxiv.org/abs/1306.2385v1>.
- [Noo06] Rutger Noot. Correspondances de Hecke, action de Galois et la conjecture d’André-Oort. In *Séminaire Bourbaki : volume 2004/2005, exposés 938-951*, number 307 in Astérisque, pages 165–197. Société mathématique de France, 2006. talk:942. URL: http://www.numdam.org/item/SB_2004-2005_47__165_0/.
- [NW13] Junjirō Noguchi and Jörg Winkelmann. *Nevanlinna theory in several complex variables and Diophantine approximation*, volume 350. Springer Science & Business Media, 2013.
- [Ogu94] Keiji Ogus. Two remarks on Calabi-Yau Moishezon threefolds. *Journal für die reine und angewandte Mathematik*, 452:153–161, 1994. doi:10.1515/crll.1994.452.153.
- [Osb12] M Scott Osborne. *Basic homological algebra*, volume 196. Springer Science & Business Media, 2012.

- [Par68] Aleksey Nikolaevich Parshin. Algebraic curves over function fields. I. *Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya*, 32(5):1191–1219, 1968. doi:10.1070/IM1968v002n05ABEH000723.
- [Pil00] Anand Pillay. Some model theory of compact complex spaces. In *Hilbert’s Tenth Problem: Relations with Arithmetic and Algebraic Geometry*, volume 270 of *Contemporary Mathematics*, pages 323–338. American Mathematical Society, 2000.
- [Pin90] Richard Pink. *Arithmetical compactification of mixed Shimura varieties*. PhD thesis, Rheinischen Friedrich-Wilhelms-Universität Bonn, 1990. <https://people.math.ethz.ch/~pink/ftp/phd/PinkDissertation.pdf>.
- [Poo17] Bjorn Poonen. *Rational points on varieties*, volume 186. American Mathematical Soc., 2017.
- [PPS17] Giuseppe Pareschi, Mihnea Popa, and Christian Schnell. Hodge modules on complex tori and generic vanishing for compact Kähler manifolds. *Geometry & Topology*, 21(4):2419–2460, 2017.
- [PS13] Mihnea Popa and Christian Schnell. Generic vanishing theory via mixed Hodge modules. *Forum of Mathematics, Sigma*, 1:e1, 2013. doi:10.1017/fms.2013.1.
- [Rei64] Hans-Jörg Reiffen. Prolongement de Riemann concernant les classes de cohomologie à supports compacts. *Comptes rendus hebdomadaires des séances de l’Académie des sciences*, 259(15):2333–2335, 1964.
- [Ren69] Rudolf Rentschler. Sur les modules M tels que $\text{Hom}(M, -)$ commute avec les sommes directes. *CR Acad. Sci. Paris*, 268:930–933, 1969. Série A.
- [Ric14] Timo Richarz. A new approach to the geometric Satake equivalence. *Documenta Mathematica*, 19:209–246, 2014.
- [Rie71] Oswald Riemenschneider. Characterizing Moisëzon spaces by almost positive coherent analytic sheaves. *Mathematische Zeitschrift*, 123(3):263–284, 1971.
- [Rie17] Emily Riehl. *Category theory in context*. Courier Dover Publications, 2017.
- [Rob00] Alain M Robert. *A course in p -adic analysis*, volume 198. Springer Science & Business Media, 2000.

- [Ros56] Maxwell Rosenlicht. Some basic theorems on algebraic groups. *American Journal of Mathematics*, 78(2):401–443, 1956.
- [Ros58] Maxwell Rosenlicht. Extensions of vector groups by abelian varieties. *American Journal of Mathematics*, 80(3):685–714, 1958.
- [Ros68] Hugo Rossi. Picard variety of an isolated singular point. *Rice Institute Pamphlet-Rice University Studies*, 54(4), 1968.
- [Rot96] Mitchell Rothstein. Sheaves with connection on abelian varieties. *Duke Mathematical Journal*, 84(3):565–598, 1996.
- [Rot97] Mitchell Rothstein. Correction to “Sheaves with connection on abelian varieties”. *Duke Mathematical Journal*, 87(1):205–211, 1997.
- [RR70] Jean-Pierre Ramis and Gabriel Ruget. Complexe dualisant et théorèmes de dualité en géométrie analytique complexe. *Publications Mathématiques de l’IHÉS*, 38:77–91, 1970.
- [RR74] Jean-Pierre Ramis and Gabriel Ruget. Résidus et dualité. *Inventiones mathematicae*, 26:89–131, 1974. doi:10.1007/BF01435691.
- [RRV71] JP Ramis, G Ruget, and JL Verdier. Dualité relative en géométrie analytique complexe. *Invent. Math.*, 13:261–283, 1971.
- [RS17] Jean Ruppenthal and Martin Sera. Modifications of torsion-free coherent analytic sheaves. *Annales de l’Institut Fourier*, 67(1):237–265, 2017.
- [RZ10] Luis Ribes and Pavel Zalesskii. *Profinite groups*. Springer, 2nd edition, 2010. doi:10.1007/978-3-642-01642-4.
- [Sab07] Claude Sabbah. De Rham and Dolbeault cohomology of \mathcal{D} -modules, 2007. https://perso.pages.math.cnrs.fr/users/claude.sabbah/livres/sabbah_oberwolfach07.pdf.
- [Sab11] Claude Sabbah. Introduction to the theory of \mathcal{D} -modules. https://perso.pages.math.cnrs.fr/users/claude.sabbah/livres/sabbah_nankai110705.pdf, 2011.
- [Sai89] Morihiko Saito. Induced D -modules and differential complexes. *Bulletin de la Société Mathématique de France*, 117(3):361–387, 1989.
- [SBW89] Jean-Pierre Serre, Martin Brown, and Michel Waldschmidt. *Lectures on the Mordell-Weil theorem*. Springer, 1989.

- [Sch14] René Scheider. *The de Rham realization of the elliptic polylogarithm in families*. PhD thesis, Universität Regensburg, 2014. <https://arxiv.org/pdf/1408.3819v1.pdf>.
- [Sch15] Christian Schnell. Holonomic D -modules on abelian varieties. *Publications mathématiques de l’IHÉS*, 121(1):1–55, 2015. doi:10.1007/s10240-014-0061-x.
- [Sch19] Peter Scholze. Lectures on analytic geometry, 2019. <https://www.math.uni-bonn.de/people/scholze/Analytic.pdf>.
- [Sch22] Peter Scholze. Six-functor formalisms, 2022. <https://people.mpim-bonn.mpg.de/scholze/SixFunctors.pdf>.
- [Ser56] Jean-Pierre Serre. Géométrie algébrique et géométrie analytique. *Annales de l’institut Fourier*, 6:1–42, 1956. doi:10.5802/aif.59.
- [Ser58] Jean-Pierre Serre. Morphismes universels et variété d’Albanese. *Séminaire Claude Chevalley*, 4:1–22, 1958.
- [Ser64] Jean-Pierre Serre. *Lie algebras and Lie groups: 1964 lectures given at Harvard University*. Springer, 1964.
- [Ser88] Jean-Pierre Serre. *Algebraic groups and class fields*, volume 117. Springer Science & Business Media, 1988. doi:10.1007/978-1-4612-1035-1.
- [Sev42] Francesco Severi. La teoria generale dei sistemi continui di curve sopra una superficie algebrica. *Memorie della R. Acc. d’Italia*, XII:337–430, 1942.
- [Sha49] Arnold Shapiro. Group extensions of compact Lie groups. *Annals of Mathematics*, pages 581–586, 1949.
- [Sha18] Jafar Shaffaf. A solution of the Erdős–Ulam problem on rational distance sets assuming the Bombieri–Lang conjecture. *Discrete & Computational Geometry*, 60:283–293, 2018. doi:10.1007/s00454-018-0003-3.
- [Sie29] Carl Ludwig Siegel. Über einige Anwendungen diophantischer Approximationen. *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, 1929(1):209–266, 1929.
- [Sim92] Carlos Simpson. Higgs bundles and local systems. *Publications Mathématiques de l’IHÉS*, 75:5–95, 1992.

- [Sim93] Carlos Simpson. Subspaces of moduli spaces of rank one local systems. *Annales scientifiques de l'Ecole normale supérieure*, 26(3):361–401, 1993.
- [Sim97] C Simpson. The Hodge filtration on nonabelian cohomology. In *Algebraic geometry–Santa Cruz 1995*, volume 62 of *Proceedings of Symposia in Pure Mathematics*, pages 217–281, 1997. Part 2.
- [Spa88] N. Spaltenstein. Resolutions of unbounded complexes. *Compositio Mathematica*, 65(2):121–154, 1988.
- [SS94] Pierre Schapira and Jean-Pierre Schneiders. Index theorem for elliptic pairs. *Astérisque*, 224:1–4, 1994.
- [Sta24] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2024.
- [Tay02] Joseph L Taylor. *Several complex variables with connections to algebraic geometry and Lie groups*, volume 46. American Mathematical Society Providence, 2002.
- [th16] t3suji (<https://mathoverflow.net/users/2653/t3suji>). Fourier Mukai transform for non-quasi coherent sheaves. MathOverflow, 2016.
- [Tim87] Klaus Timmerscheidt. Mixed Hodge theory for unitary local systems. *Journal für die reine und angewandte Mathematik*, 379:152–171, 1987. doi:10.1515/crll.1987.379.152.
- [TT90] Robert W Thomason and Thomas Trobaugh. Higher algebraic K-theory of schemes and of derived categories. In *The Grothendieck Festschrift*, volume III, pages 247–435. Springer, 1990.
- [Tu97] Lihuang Tu. A proposition of the convergent power series ring. *Acta Mathematica Sinica, Chinese Edition*, 40(3):365–368, 1997. https://actamath.cjoe.ac.cn/Jwk_sxxb_cn/EN/10.12386/A1997sxxb0059. doi:10.12386/A1997sxxb0059.
- [Uen80] Kenji Ueno. On three-dimensional compact complex manifolds with non-positive Kodaira dimension. *Proceedings of the Japan Academy, Series A, Mathematical Sciences*, 56(10):479–483, 1980.
- [Uen83] Kenji Ueno, editor. *Classification of algebraic and analytic manifolds: Katata symposium proceedings 1982*, volume 39. Birkhäuser, 1983.

- [Uen06] Kenji Ueno. *Classification theory of algebraic varieties and compact complex spaces*, volume 439. Springer, 2006.
- [Ull04] Emmanuel Ullmo. Points rationnels des variétés de Shimura. *International Mathematics Research Notices*, 2004(76):4109–4125, 2004.
- [UY86] Karen Uhlenbeck and Shing-Tung Yau. On the existence of Hermitian-Yang-Mills connections in stable vector bundles. *Communications on Pure and Applied Mathematics*, 39(S1):S257–S293, 1986.
- [UY10] Emmanuel Ullmo and Andrei Yafaev. Points rationnels des variétés de Shimura: un principe du “tout ou rien”. *Mathematische Annalen*, 348(3):689–705, 2010.
- [UY14] Emmanuel Ullmo and Andrei Yafaev. Galois orbits and equidistribution of special subvarieties: towards the André-Oort conjecture. *Annals of mathematics*, 180(3):823–865, 2014.
- [Vak23] Ravi Vakil. The rising sea: Foundations of algebraic geometry. <http://math.stanford.edu/~vakil/216blog/FOAGapr0123public.pdf>, 2023.
- [Var86] Jean Varouchas. Propriétés cohomologiques d’une classe de variétés analytiques complexes compactes. In *Séminaire d’Analyse*, pages 233–243. Springer, 1986. doi:10.1007/BFb0077057.
- [Var89] Jean Varouchas. Kähler spaces and proper open morphisms. *Mathematische Annalen*, 283:13–52, 1989.
- [Ver66] Jean-Louis Verdier. Dualité dans la cohomologie des espaces localement compacts. In *Séminaire Bourbaki : années 1964/65 1965/66, exposés 277-312*, number 9 in Séminaire Bourbaki. Société mathématique de France, 1966. talk:300. URL: http://www.numdam.org/item/SB_1964-1966__9__337_0/.
- [Vig21] Florian Viguier. *D-modules arithmétiques et transformation de Fourier-Mukai*. PhD thesis, Université de Strasbourg, 2021.
- [Voi02] Claire Voisin. *Hodge theory and complex algebraic geometry I*. Cambridge University Press Cambridge, 2002.
- [Wan16] Botong Wang. Torsion points on the cohomology jump loci of compact Kähler manifolds. *Mathematical Research Letters*, 23(2):545–563, 2016.

- [Wei48] André Weil. *Variétés abéliennes et courbes algébriques*, volume 121. Paris Hermann, 1948.
- [Wei49] A Weil. Variétés abéliennes. In *Colloque d’algebre et théorie des nombres*. CNRS, 1949.
- [Wei95] Charles A Weibel. *An introduction to homological algebra*. Number 38 in Cambridge Studies in Advanced Mathematics. Cambridge university press, 1995.
- [Wei07] Rainer Weissauer. Brill-Noether sheaves, 2007. <https://arxiv.org/pdf/math/0610923v4.pdf>.
- [Wei11] R. Weissauer. A remark on rigidity of BN-sheaves, 2011. <https://arxiv.org/pdf/1111.6095v1.pdf>.
- [Wei16] Rainer Weissauer. Vanishing theorems for constructible sheaves on abelian varieties over finite fields. *Mathematische Annalen*, 365(1):559–578, 2016.
- [Whi72] Hassler Whitney. *Complex analytic varieties*, volume 131. Addison-Wesley Reading, 1972.
- [Wib22] Michael Wibmer. A remark on torsors under affine group schemes. *Transformation Groups*, pages 1–8, 2022. doi:10.1007/s00031-022-09767-z.
- [Wil70] Stephen Willard. *General topology*. Addison-Wesley publishing company, 1970.
- [Wil16] Geordie Williamson. The Hodge theory of the decomposition theorem (after de Cataldo and Migliorini), 2016.
- [Wu86] Xiaolong Wu. On the extensions of abelian varieties by affine group schemes. In *Group Theory, Beijing 1984: Proceedings of an International Symposium held in Beijing, Aug. 27–Sep. 8, 1984*, pages 361–387. Springer, 1986. doi:10.1007/BFb0076184.
- [Yam19] Katsutoshi Yamanoi. Pseudo Kobayashi hyperbolicity of subvarieties of general type on abelian varieties. *Journal of the Mathematical Society of Japan*, 71(1):259–298, 2019.
- [Yek19] Amnon Yekutieli. *Derived categories*, volume 183. Cambridge University Press, 2019. doi:10.1017/9781108292825.
- [Yun14] Zhiwei Yun. Motives with exceptional Galois groups and the inverse Galois problem. *Inventiones mathematicae*, 196(2):267–337, 2014. doi:10.1007/s00222-013-0469-9.

- [Zhu17] Xinwen Zhu. An introduction to affine Grassmannians and the geometric Satake equivalence. In *Geometry of moduli spaces and representation theory*, volume 24 of *IAS/Park City Math. Ser.*, pages 59–154. Amer. Math. Soc., Providence, RI, 2017.

Index

- D_X -module, 30
- $\mathrm{Mod}_{\mathrm{qc}}(D_X)$, 245
- $\mathrm{Mon}(K, \psi)$ monodromy group, 80
- λ -connection, 32
- Albanese, 102
- Berstein equivalence theorem, 245
- Brody hyperbolic, 16
- coherator, 247
- coherent sheaves, 209
- convolution, 22
- cotorus, 70
- defect of semismallness, 109
- Fourier-Mukai transform, 28
- Fujiki class \mathcal{C} , 117
- general type, 15
- generic Mumford-Tate group, 44
- geometric Lang conjecture, 16
- good sheaves, 214
- Green-Lazarsfeld generic vanishing theorem, 26
- Hecke correspondence, 43
- Higgs bundle, 32
- holonomic D -modules, 190
- homogeneous vector bundle, 30
- integral Lang locus, 51
- Jacobian, 101
- Kodaira dimension, 15
- Kr mer-Weissauer generic vanishing theorem, 26
- Lang locus, 38
- Laumon-Rothstein theorem
 - algebraic, 165
 - analytic, 185
- Laumon-Rothstein transform, 185
- Lawrence-Sawin theorem, 21
- Moishezon manifold, 119
- Mordell conjecture, 13
- Mukai duality, 29
- neat subgroup, 41
- neutral Tannakian category, 22
- quasi-coherent sheaves, 209
- regular manifold, 97
- relative perverse sheaves, 64
- Rothstein transform, 177
- semisimple constructible complex of sheaves, 58
- Shafarevich conjecture, 14
- special subvariety, 44
- Spectrum of perverse sheaf, 75
- Tannakian monodromy group, 78
- unipotent vector bundle, 30
- universally locally acyclic, 61