

ON THE CONSTRUCTION OF VALUATIONS AND GENERATING SEQUENCES ON HYPERSURFACE SINGULARITIES

STEVEN DALE CUTKOSKY, HUSSEIN MOURTADA AND BERNARD TEISSIER

ABSTRACT. Suppose that (K, ν) is a valued field, $f(z) \in K[z]$ is a unitary and irreducible polynomial and (L, ω) is an extension of valued fields, where $L = K[z]/(f(z))$. Let A be a local domain with quotient field K dominated by the valuation ring of ν and such that $f(z)$ is in $A[z]$.

This paper is devoted to the problem of describing the structure of the associated graded ring $\text{gr}_\omega A[z]/(f(z))$ of $A[z]/(f(z))$ for the filtration defined by ω as an extension of the associated graded ring of A for the filtration defined by ν . In particular we give an algorithm which in many cases produces a finite set of elements of $A[z]/(f(z))$ whose images in $\text{gr}_\omega A[z]/(f(z))$ generate it as a $\text{gr}_\nu A$ -algebra as well as the relations between these images. We also work out the interactions of our method with phenomena which complicate the study of ramification and local uniformization in positive characteristic, such as the non tameness and the defect of an extension. For a valuation ν of rank one and a separable extension of valued fields $(K, \nu) \subset (L, \omega)$ as above our algorithm produces a generating sequence in a local birational extension A_1 of A dominated by ν if and only if there is no defect. In this case, $\text{gr}_\omega A_1[z]/(f(z))$ is a finitely presented $\text{gr}_\nu A_1$ -module.

(French) Soient (K, ν) un corps valué, $f(z) \in K[z]$ un polynôme unitaire irréductible et (L, ω) une extension de corps valués où $L = K[z]/(f(z))$. Soit encore A un anneau local dont le corps de fractions est K dominé par l'anneau de valuation de ν , et tel que $f(z) \in A[z]$. Cet article est consacré à la description de de l'anneau gradué $\text{gr}_\omega A[z]/(f(z))$ associé à la filtration de $A[z]/(f(z))$ déterminée par la valuation ω en tant qu'extension de l'anneau gradué associé à la filtration de A déterminée par ν . En particulier nous définissons un algorithme qui dans de nombreux cas produit un ensemble fini d'éléments de $A[z]/(f(z))$ donts les images dans $\text{gr}_\omega A[z]/(f(z))$ l'engendrent en tant que $\text{gr}_\nu A$ -algèbre ainsi que les relations entre ces images. Nous décrivons aussi les interactions entre notre méthode et des phénomènes qui compliquent l'étude de la ramification et de l'uniformisation locale en caractéristique positive, tels que la non modération et le défaut. Pour une valuation ν de rang un et une extension séparable de corps valués comme ci-dessus notre algorithme produit une suite génératrice dans une extension birationnelle A_1 de A dominée par ν si et seulement si l'extension est sans défaut. Dans ce cas $\text{gr}_\omega A_1[z]/(f(z))$ est un $\text{gr}_\nu A_1$ -module de présentation finie.

1. INTRODUCTION

Given a field K and an extension L of K , the study of the problem of extending a valuation from K to L has a long history motivated in part by its close relation with ramification theory, whether in number theory or in algebraic geometry. It has an incarnation in logic,

2000 *Mathematics Subject Classification.* 12J20, 16W60, 14B25.

Key words and phrases. Valuations, ramification, defect, generating sequences, key polynomials.

The first author was partially supported by NSF grant DMS-1700046. He also thanks the Université Paris Diderot and the Fondation Sciences Mathématiques de Paris. The second author was partially supported by The Miller scholar in residence program of the University of Missouri at Columbia.

the model theory of valued fields which provides another viewpoint on ramification theory. After fundamental work by E. Artin, H. Hasse, A. Ostrowski and others, S. MacLane created a method for describing all extensions of a discrete rank one valuation on a field K to a primitive extension $K(z)$, be it algebraic or transcendental. The method is based on the existence of *key polynomials* in $K[z]$ which provide successive approximations of a given extension of the valuation and, by the behavior of their degrees, a measure of its complexity.

On the side of algebraic geometry, Zariski's approach to resolution of singularities of algebraic varieties using local uniformization of valuations provides a strong motivation for the study of valuations on local domains essentially of finite type over a field, which waned after Hironaka's proof of resolution in characteristic zero but later revived as an approach to resolution in positive characteristic.

In the 1970's and 1980's appeared (see [33], [32], [15]) the idea that the associated graded ring $\text{gr}_\nu A$ of a local domain A with respect to the filtration of A associated to a valuation ν of its field of fractions centered in A (non negative on A and positive on its maximal ideal) encoded in a geometric way essential characters of the valuation and could be used at least in special cases to obtain local uniformization. For example, representatives in A of the generators of the graded algebra associated to a one dimensional integral complex analytic algebra can be used to embed the corresponding curve in an affine space where a single birational toric modification provides an embedded resolution of singularities (see [15]). It also became apparent that some of MacLane's essential definitions are better understood using associated graded rings.

Somewhat later, MacLane's theory was generalized by Vaquié who extended to all Krull valuations the construction of sequences of key polynomials, now indexed by ordinals (see [36], [37], [38] and section 7 below). He also described the extension $\text{gr}_\nu K \subset \text{gr}_\omega K[z]$ of graded rings corresponding to an extension of valuations from ν on K to ω on $K[z]$, for z algebraic or transcendental over K . It appeared that the images of MacLane's and Vaquié's key polynomials in the graded algebra $\text{gr}_\omega K[z]$ were related to its generation as a $\text{gr}_\nu K$ -algebra.

In the last three decades or so the problem of describing a *generating sequence* for a valuation, which is a set of elements of a ring A whose images in $\text{gr}_\nu A$ provide a presentation by generators and relations has become of major interest for the ramification theory of extensions of valued fields as well as for local uniformization in positive characteristic, which is still an open problem.

In fact it has become apparent that given an extension $(A, \nu) \subset (B, \omega)$ of valued rings the extension $\text{gr}_\nu A \subset \text{gr}_\omega B$ of the associated graded algebras, as well as the similar extensions obtained after birational extensions of A and B encodes in a comparatively simple language, such as the condition of being finitely generated, essential information about the ramification of the original extension. This concerns especially the defect and the possibility to uniformize ω on B if we can uniformize ν on A . But we can access this information only if we have descriptions by generators and relations of $\text{gr}_\nu A$ and $\text{gr}_\omega B$, or of $\text{gr}_\omega B$ as a $\text{gr}_\nu A$ -algebra. This is the main motivation for this work.

The problem of constructing generating sequences in a Noetherian local domain A which is dominated by a valuation ω is extremely difficult, and little is known about this problem in general. The difficulty reflects the fact that the structure of the semigroup of values $S^A(\omega) = \omega(A \setminus \{0\})$ can be extremely complicated. It is well understood in the case that A has dimension one (see [33], [15]), and for regular local rings of dimension two ([32], [9],

[25]). It is known for certain valuations dominating two dimensional quotient singularities [12] and for certain valuations dominating three dimensional regular local rings [18].

Here we consider the case where the essence of the difficulty resides: suppose that (K, ν) is a valued field, $f(z) \in K[z]$ is a unitary and irreducible polynomial and (L, ω) is a finite field extension, where $L = K[z]/(f(z))$. Further suppose that A is a local domain with quotient field K such that ν dominates A and that $f(z)$ is in $A[z]$. We provide an algorithm producing the first significant part of a generating sequence for extensions of a valuation ν to $A[z]/(f(z))$.

The associated graded ring of $K[z]/(f(z))$ along ω as an extension of the associated graded ring of K along ν has been constructed implicitly, in the papers [23], [24] of MacLane for discrete rank one valuations, and for general valuations by Vaquié in [36], [37], [38]. Further papers on this topic, and comparison with the method of pseudo convergent sequences (introduced by Ostrowski in [28, Teil III, §11] and developed by Kaplansky in [17]) are [2], [27], [31], [16] and [10]. Finding generating sequences for $A[z]/(f(z))$ in the case where A is an arbitrary noetherian subring dominated by R_ν and with the same field of fractions is much more difficult.

This paper is devoted to this problem. We describe the relationship of our method with the key polynomials of MacLane and Vaquié. We also work out the interactions of our method of computation with phenomena which complicate the study of ramification in positive characteristic, such as the lack of tameness and the defect of an extension.

We now give more details about the content of this paper:

Let G_ν be the value group of ν and R_ν be the valuation ring of ν , with maximal ideal m_ν . Given a subring A of the field of fractions of R_ν , the associated graded ring of A along ν is defined as

$$\text{gr}_\nu(A) = \bigoplus_{\gamma \in G_\nu} \mathcal{P}_\gamma(A) / \mathcal{P}_\gamma^+(A)$$

where

$$\mathcal{P}_\gamma(A) = \{g \in A \setminus \{0\} \mid \nu(g) \geq \gamma\} \text{ and } \mathcal{P}_\gamma^+(A) = \{g \in A \setminus \{0\} \mid \nu(g) > \gamma\}.$$

The ring $\text{gr}_\nu(A)$ is an algebra over its degree zero subring. It is a domain which is generally not Noetherian. In this text we shall consider subrings of R_ν so that the semigroup $S^A(\nu)$ of values of elements of $A \setminus \{0\}$ which indexes the homogeneous components of $\text{gr}_\nu(A)$ is contained in the positive part of G_ν . We shall see more about this semigroup below.

Important invariants of a finite extension $(K, \nu) \subset (L, \omega)$ of valued fields are the reduced ramification index and residue degree of ω over ν , which are

$$e(\omega/\nu) = [G_\omega : G_\nu] \text{ and } f(\omega/\nu) = [R_\omega/m_\omega : R_\nu/m_\nu].$$

Another, very subtle invariant is the defect $\delta(\omega/\nu)$ of the extension, which is a power of the characteristic p of the residue field R_ν/m_ν . The defect and its role in local uniformization is explained in [19]. We give the definition of the defect in (43) below. In the case where ω is the unique extension of ν to L we have that

$$(1) \quad [L : K] = e(\omega/\nu) f(\omega/\nu) \delta(\omega/\nu).$$

If A and B are local domains with quotient fields K and L such that ω dominates B and B dominates A , we have a graded inclusion of graded domains

$$\text{gr}_\nu(A) \rightarrow \text{gr}_\omega(B).$$

The index of quotient fields

$$[\mathrm{QF}(\mathrm{gr}_\omega(B)) : \mathrm{QF}(\mathrm{gr}_\nu(A))] = e(\omega/\nu)f(\omega/\nu)$$

by Proposition 3.3 of [3]. The defect seems to disappear, but it manifests itself in mysterious behavior in the extensions of associated graded rings of injections $A' \rightarrow B'$ of birational extensions of Noetherian local domains A, B . For instance, if ν has rational rank 1 but is not discrete, the defect $\delta(\omega/\nu)$ is larger than 1 and A and B are two dimensional excellent local domains, then $\mathrm{gr}_\omega(B')$ is not a finitely generated $\mathrm{gr}_\nu(A')$ -algebra for any regular local rings $A' \rightarrow B'$ which are dominated by ω and dominate A and B as shown in [6].

The construction of generating sequences is closely related to the problem of local uniformization. In Theorem 7.1 [7], it is shown how reduction of multiplicity along a rank 1 valuation can be achieved in a defectless extension $A \rightarrow A[z]/(f(z))$. The statement “defectless” means that the rank 1 valuations ν and ω satisfy $\delta(\omega/\nu) = 1$. From this assumption, it follows that either $\omega(z - K)$ has a largest element, or the limsup of this set is ∞ . If the limsup of this set is ∞ , then in an appropriate extension, the valuation ω corresponds to a linear factor of $f(z)$, and it is not difficult to realize a reduction of multiplicity by blowing up. So assume that $\omega(z - K)$ has a largest element $\gamma \in G_\omega$. We then have $\gamma \notin G_\nu$. After a birational extension A_1 of A and a change of variables of z in $A_1[z]$, we obtain that $\omega(z) = \gamma$ and then after a Cremona transformation involving z , we obtain a reduction of the multiplicity of the strict transform of f .

In [34] and [35], it is shown how associated graded rings along a valuation can be used to prove local uniformization, at least when the associated graded rings are finitely generated algebras over A/m_A . A suitable toric resolution of singularities of the associated graded ring induces a local uniformization of the given valuation.

The subring of degree zero elements of the graded ring $\mathrm{gr}_\nu(A)$ is $(\mathrm{gr}_\nu(A))_0 = A/Q$ where Q is the prime ideal in A of elements of positive value. A generating sequence for ν on A is an ordered set of elements of A whose classes in $\mathrm{gr}_\nu(A)$ generate $\mathrm{gr}_\nu(A)$ as a graded $(\mathrm{gr}_\nu(A))_0$ -algebra. To be meaningful, a generating sequence should come with a formula for computing the values of elements of A , and their relations in $\mathrm{gr}_\nu(A)$. In particular, a generating sequence should give the structure of $\mathrm{gr}_\nu(A)$ as a graded $(\mathrm{gr}_\nu(A))_0$ -algebra.

In the case of an inclusion $A \subset B$ of domains, and an extension ω of ν to the quotient field of B such that ω has nonnegative value on B , a generating sequence of the extension is an ordered sequence of elements of B whose classes in $\mathrm{gr}_\omega(B)$ generate $\mathrm{gr}_\omega(B)$ as a $\mathrm{gr}_\nu(A)$ -algebra. A generating sequence for an extension should come with a formula for computing the values of elements of B , relative to the values of elements of A , and give their relations in $\mathrm{gr}_\nu(B)$. That is, a generating sequence should give the structure of $\mathrm{gr}_\omega(B)$ as a graded $\mathrm{gr}_\nu(A)$ -algebra.

In this paper, we give a very simple algorithm which allows us to compute a generating sequence and the structure of $\mathrm{gr}_\omega(A[z]/(f(z)))$ in many situations. Throughout this paper, we have the assumption that A is a local domain which contains an algebraically closed field k such that its residue field $A/m_A = k$, ν dominates A and the residue field of the valuation ring R_ν of ν is $R_\nu/m_\nu = k$ (ν is a “rational valuation”). This algorithm is derived in Section 4. The algorithm is valid for an arbitrary extension ω of an arbitrary valuation ν dominating A ($m_\nu \cap A = m_A$).

A realization of our algorithm produces a subring of $\mathrm{gr}_\omega(R_\nu[z]/(f(z)))$ which is the quotient of a graded polynomial ring C over $\mathrm{gr}_\nu(R_\nu)$ in either finitely many or countably many variables, and a set of generators of the graded prime ideal I of C such that C/I is isomorphic to the subring of $\mathrm{gr}_\omega(R_\nu[z]/(f(z)))$. Our algorithm gives an explicit representation of

the subring as

$$\mathrm{gr}_\nu(R_\nu)[\bar{\varphi}_1, \dots, \bar{\varphi}_k, \dots]/I$$

where

$$I = (\bar{\varphi}_1^{n_1} - \bar{c}_1, \bar{\varphi}_2^{n_2} - \bar{c}_2 \bar{\varphi}_1^{j_1(2)}, \dots, \bar{\varphi}_k^{n_k} - \bar{c}_k \bar{\varphi}_1^{j_1(k)} \bar{\varphi}_2^{j_2(k)} \dots \bar{\varphi}_{k-1}^{j_{k-1}(k)}, \dots)$$

with $\bar{c}_1, \dots, \bar{c}_k, \dots \in \mathrm{gr}_\nu(R_\nu)$ homogeneous elements. The elements $\bar{\varphi}_i$ are homogeneous with strictly increasing values. If our algorithm terminates in a finite number of steps k , then elements $\varphi_1, \dots, \varphi_k \in R_\nu[z]$ whose classes are $\bar{\varphi}_1, \dots, \bar{\varphi}_k$ form a generating sequence of $R_\nu[z]/(f(z))$ over R_ν and we have built up the entire associated graded ring

$$\mathrm{gr}_\nu(R_\nu[z]/(f(z))) = \mathrm{gr}_\nu(R_\nu)[\bar{\varphi}_1, \dots, \bar{\varphi}_k]/I$$

where

$$I = (\bar{\varphi}_1^{n_1} - \bar{c}_1, \bar{\varphi}_2^{n_2} - \bar{c}_2 \bar{\varphi}_1^{j_1(2)}, \dots, \bar{\varphi}_k^{n_k} - \bar{c}_k \bar{\varphi}_1^{j_1(k)} \bar{\varphi}_2^{j_2(k)} \dots \bar{\varphi}_{k-1}^{j_{k-1}(k)}).$$

In this case, we have that $\mathrm{gr}_\nu(R_\nu[z]/(f(z)))$ is a finitely generated and presented $\mathrm{gr}_\nu(R_\nu)$ -module.

When we compare our algorithm to the theory of Vaquié ([36], [37], [38]) in Subsection 7.1, we conclude in Proposition 7.1 that a realization of our algorithm produces the “first simple admissible family” $\mathcal{S}^{(1)}$ of an “admissible family” \mathcal{S} determining the valuation ω .

Our algorithm produces in many situations a finite sequence of elements of $A[z]$ whose images generate the $\mathrm{gr}_\nu A$ -algebra $\mathrm{gr}_\omega A[z]$. It does this even in cases where there are infinitely many key polynomials.

More precisely, if the characteristic p of k does not divide the degree of f , A is a domain as above and ω is the unique extension of ν to a valuation of the quotient field L of $A[z]/(f(z))$, then we show in Theorem 5.1 that our algorithm produces a finite generating sequence in $A[z]/(f(z))$. The associated graded ring of $A[z]/(f(z))$ along ω is then a finitely generated and presented module over the associated graded ring of A along ν .

Since the defect $\delta(\omega/\nu)$ is always a power of p , the assumption that p does not divide the degree of f in Theorem 5.1 and the assumption that ω is the unique extension of ν forces the defect $\delta(\omega/\nu)$ to be 1 by (1).

We show that if any of the above assumptions are removed, then the conclusions of Theorem 5.1 do not hold (Examples of Section 4 and Section 11). For instance, the assumption that $R_\nu[z]/(f(z))$ is a “hypersurface singularity” is shown to be necessary for finite generation to hold in Example 11.3.

To illustrate the power of Theorem 5.1, we compute in Example 5.2 the associated graded ring when $f(z)$ is a quadratic polynomial, k has characteristic not equal to 2 and ω is the unique extension of ν . It has the simple form

$$\mathrm{gr}_\omega(A[z]/(f(z))) \cong \mathrm{gr}_\nu(A)[\bar{\varphi}]/(\bar{\varphi}^2 - \bar{c})$$

for some homogeneous $\bar{c} \in \mathrm{gr}_\nu(A)$. From the classification of associated graded rings of valuations dominating a two dimensional regular local ring A ([32] and [9]) we see that we are able to completely calculate the associated graded ring along an extended valuation in the local rings of two dimensional rational double points, when the extension ω is unique. In contrast, if ω is not the unique extension of ν , then $\mathrm{gr}_\omega(A[z]/(f(z)))$ might not be a finitely generated $\mathrm{gr}_\nu(A)$ -module, as shown in Examples 5.2 and 11.4.

In Theorem 8.2, we consider an arbitrary separable extension (with no assumption on the degree) and assume that A is an excellent local domain. We show that an extension of a rank one valuation ν is without defect if and only if there exists a realization of our algorithm with coefficients in a birational extension A_1 of A which constructs ω , either

as a valuation or a limit valuation. A birational extension A_1 of A is a localization of a finitely generated A -algebra whose quotient field is K and which is dominated by ν .

An example showing that the conclusions of Theorem 8.2 may not hold if ν has rank larger than one is given in Section 10. In Example 8.3, it is shown that the conclusions of Theorem 8.2 may not hold if $f(z)$ is not separable over K .

In Section 9 we analyze our algorithm in a rank 1 example with defect from [11] to motivate the necessary condition of Theorem 8.2. We explicitly show that a generating sequence does not exist in $A_1[z]$ for any birational extension A_1 of A which is dominated by ν , and the valuation ω is not realizable as a limit valuation.

In the final section, Section 11, we give examples showing that the finite generation of extensions of associated graded rings and valuation semigroups ensured by Theorem 5.1 may fail if any of the assumptions of the theorem are removed. The semigroup $S^A(\nu)$ of values of ν on A is

$$S^A(\nu) = \{\nu(g) \mid g \in A \setminus \{0\}\}.$$

In Example 11.3, it is shown that there exists an extension L of the quotient field K of A of degree prime to p , a valuation ν of K which dominates A and has a unique extension to L such that if B is the integral closure of A in L , then $\text{gr}_\omega(B)$ is not a finitely generated $\text{gr}_\nu(A)$ -module and the semigroup $S^B(\omega)$ is not a finitely generated $S^A(\nu)$ -module. In particular, the conclusions of Theorem 5.1 do not hold for this extension. This example shows that we must have the condition that $B = A[z]/(f(z))$ is a ‘‘hypersurface singularity’’ for the conclusions of Theorem 5.1 to be true.

We make use of the theory of MacLane, [23], [24], which he developed to construct the extensions of a (rank 1) discrete valuation ν of K to a discrete valuation ω of $k[z]$ or of $K[z]/(f(z))$ for some irreducible unitary polynomial $f(z) \in K[z]$. Our algorithm can be viewed as a realization of MacLane’s method in the context of a general valuation, in a specific, nice form. MacLane’s theory is surveyed in Section 3.

We also make use of Vaquié’s generalization of MacLane’s method in [36], [37], [38] to construct extensions of general valuations in $K[z]$ and $K[z]/(f(z))$ in our proof of Theorem 8.2. The essential new concept in Vaquié’s work is that of a ‘‘limit key polynomial’’. He gave in [38, Exemple 4.1] an example of infinite sequences of key polynomials due to the non uniqueness of valuation extension. Vaquié’s method is surveyed in Section 7, as well as a study of its relationship to our algorithm. In the situation of this paper we shall meet only finite sequences of limit key polynomials since the number of limit key polynomials is bounded by the degree of $f(z)$. In Section 6 we collect and derive some results about Henselizations of rings and valued fields which we need for the proof of Theorem 8.2.

In this paper, a local ring is a commutative ring with a unique maximal ideal. In particular, we do not require a local ring to be Noetherian. We will denote the maximal ideal of a local ring A by m_A . The quotient field of a domain A will be denoted by $\text{QF}(A)$. We will say that a local ring B dominates a local ring A if $A \subset B$ and $m_B \cap A = m_A$.

We will denote the natural numbers by \mathbf{N} and the positive integers by \mathbf{Z}_+ .

2. VALUATIONS AND PSEUDO VALUATIONS

We shall in the sequel consider sequences of valuations which approximate ω . For that reason we change notations and denote these sequences by V_0, V_1, \dots as in [23] and [24]. A general valuation will be denoted by V and the reader may think of ν as V_0 .

Suppose that V is a valuation on a field K . We will denote the valuation ring of V by R_V and its maximal ideal by m_V . The value group of V will be denoted by G_V .

Suppose that A is a Noetherian local domain with quotient field K and $A \rightarrow A_1$ is an extension of local domains such that A_1 is a domain whose quotient field is K and A_1 is essentially of finite type over A (A_1 is a localization of a finitely generated A -algebra). Then we will say that $A \rightarrow A_1$ is a birational extension.

If A is a domain which is contained in R_V , then the associated graded ring of A along V is $\text{gr}_V(A)$ as defined in the introduction, The initial form $\text{In}_V(g)$ of $g \in A$ is the class of g in $\mathcal{P}_{V(g)}(A)/\mathcal{P}_{V(g)}^+(A)$. The semigroup of V on A has also been defined in the introduction. A pseudo valuation V on a domain A is a surjective map $V : A \rightarrow G_V \cup \{\infty\}$ where G_V is a totally ordered Abelian group and a prime ideal

$$I(V)_\infty = I^A(V)_\infty = \{g \in A \mid V(g) = \infty\}$$

of A_1 such that $V : \text{QF}(A/I(V)_\infty) \setminus \{0\} \rightarrow G_V$ is a valuation.

3. THE MACLANE THEORY OF KEY POLYNOMIALS

Suppose that V is a valuation or a pseudo valuation on a domain A . Following MacLane in [23] in the case $A = K[z]$, we can define an equivalence \sim on A defined for $g, h \in A$ by $g \sim h$ in V if $V(g - h) > \min\{V(g), V(h)\}$ or $V(g) = V(h) = \infty$. We say that $g \in A$ is equivalence divisible by h in V , written $h|g$ in V , if there exists $a \in A$ such that $g \sim ah$ in V . An element g is said to be equivalence irreducible in V if $g|ab$ in V implies $g|a$ or $g|b$ in V .

These conditions can be expressed respectively as the statement that $\text{In}_V(h) = \text{In}_V(g)$ in $\text{gr}_V(A)$, that $\text{In}_V(h)$ divides $\text{In}_V(g)$ in $\text{gr}_V(A)$ and that the ideal generated by $\text{In}_V(g)$ in $\text{gr}_V(A)$ is prime.

3.1. MacLane's algorithm. We review MacLane's algorithm [23] to construct the extensions of a valuation V_0 of a field K to a valuation or pseudo-valuation of the polynomial ring $K[z]$. MacLane applied his method to construct extensions of rank 1 discrete valuations of K to $K[z]$. This algorithm has been extended to general valuations by Vaquié [36]. MacLane constructs "augmented sequences of inductive valuations"

$$(2) \quad V_1, \dots, V_k, \dots$$

which extend V_0 to $K[z]$. An augmented sequence (2) is constructed from successive inductive valuations

$$(3) \quad V_k = [V_{k-1}; V_k(\varphi_k) = \mu_k] \text{ for } 1 \leq k$$

of $K[z]$, where φ_k is a "key polynomial" over V_{k-1} and μ_k is a "key value" of φ_k over V_{k-1} . We always take $\varphi_1 = z$.

We say that $\varphi(z) \in K[z]$ is a key polynomial with key value μ over V_{k-1} if

- 1) $\varphi(z)$ is equivalence irreducible in V_{k-1} .
- 2) $\varphi(z)$ is minimal in V_{k-1} ; that is, if $\varphi(z)$ divides $g(z)$ in V_{k-1} , then $\deg_z \varphi(z) \leq \deg_z g(z)$.
- 3) $\varphi(z)$ is unitary and $\deg_z \varphi(z) > 0$.
- 4) $\mu > V_{k-1}(\varphi(z))$.

Following MacLane (Definition 6.1 [23]) we also assume

- 5) $\deg_z \varphi_i(z) \geq \deg_z \varphi_{i-1}(z)$ for $i \geq 2$.
- 6) $\varphi_i(z) \sim \varphi_{i-1}(z)$ in V_{i-1} is false. Here the equivalence is to be understood for polynomials in $K[z]$.

It follows from [23, Theorem 9.3] that

$$(4) \quad \text{if } \varphi(z) \text{ is a key polynomial over } V_{k-1} \text{ then } \deg_z \varphi_{k-1}(z) \text{ divides } \deg_z \varphi(z).$$

The key polynomials $\varphi_k(z)$ can further be assumed to be homogeneous in V_{k-1} , which will be defined after (7).

MacLane shows that if V_0 is discrete of rank 1, then the extensions of V_0 to a valuation or pseudo valuation of $K[z]$ are the V_k arising from augmented sequences of finite length (2) and the limit sequences of augmented sequences of infinite length (2) which determine a limit value V_∞ on $K[z]$ defined by

$$V_\infty(g(z)) = \lim_{k \rightarrow \infty} V_k(g(z)) \text{ for } g(z) \in K[z].$$

We have that $V_\infty(g(z))$ is well defined whenever V_0 has rank 1, and is a valuation or pseudo-valuation by the argument of page 10 [23].

MacLane's method has been extended by Vaquié [36], to eventually construct all extensions of an arbitrary valuation V_0 of K to a valuation or pseudo valuation of $K[z]$. We will discuss Vaquié's method in Section 7.

To compute the " k -th stage" value $V_k(g(z))$ for $g(z) \in K[z]$ by MacLane's method, we consider the unique expansion

$$(5) \quad g(z) = g_m(z)\varphi_k^m(z) + g_{m-1}\varphi_k^{m-1}(z) + \cdots + g_0$$

with $g_i(z) \in K[z]$, $\deg_z g_i(z) < \deg_z \varphi_k(z)$ for all i and $g_m(z) \neq 0$. Then

$$V_k(g(z)) = \min\{V_{k-1}(g_m(z)) + m\mu_k, V_{k-1}(g_{m-1}(z)) + (m-1)\mu_k, \dots, V_{k-1}(g_0(z))\}.$$

This expression suffices to prove by induction, assuming the existence of a unique expansion of the coefficients $g_i(z)$ in terms of the polynomials $\varphi_j(z)$ with $j < k$, that every $g(z) \in K[z]$ has a unique expansion

$$(6) \quad g(z) = \sum_j a_j(z) \varphi_1^{m_{1,j}}(z) \varphi_2^{m_{2,j}}(z) \cdots \varphi_k^{m_{k,j}}(z)$$

with $a_j \in K$ and $0 \leq m_{i,j} < \deg_z \varphi_{i+1} / \deg_z \varphi_i$ for $i = 1, \dots, k-1$. Recall that $\deg_z \varphi_{i+1} / \deg_z \varphi_i$ is a positive integer by (4). Then

$$(7) \quad V_k(g) = \min_j V_k(a_j \varphi_1^{m_{1,j}} \varphi_2^{m_{2,j}} \cdots \varphi_k^{m_{k,j}}).$$

If all terms in (6) have the same values in V_k then g is said to be homogeneous in V_k .

We shall often, as we just did, simplify notations by writing g for $g(z)$, etc. when there is no fear of confusion.

Remark 3.1. *If A is a subring of K such that $\varphi_i \in A[z]$ for $1 \leq i \leq k$ and $g \in A[z]$, then the coefficients a_j in (6) are all in A .*

The polynomial g , with expansion (5), is minimal in V_k if and only if $g_m \in K$ and

$$(8) \quad V_k(g) = V_k(g_m \varphi_k^m)$$

by 2.3 [24] or Theorem 9.3 [23].

By 3.13 of [24] or Theorem 6.5 [23], for $k > i$,

$$(9) \quad V_k(\varphi_i) = V_i(\varphi_i) \text{ and } V_k(g) = V_i(g) \text{ whenever } \deg_z g < \deg_z \varphi_{i+1}.$$

Further, by Theorems 5.1 and 6.4 [23], or 3.11 and 3.12 of [24],

$$(10) \quad \text{For all } g \in K[z], V_k(g) \geq V_{k-1}(g) \text{ with equality if and only if } \varphi_k \nmid g \text{ in } V_{k-1}.$$

3.2. MacLane's algorithm in a finite primitive extension. Suppose $f(z) \in K[z]$ is unitary and irreducible. The extensions of V_0 to valuations of $K[z]/(f(z))$ are the extensions of V_0 to pseudo valuations V of $K[z]$ such that $I(V)_\infty = (f(z))$. MacLane [24] gives an explicit explanation of how his algorithm can be applied to construct the pseudo valuations V of $K[z]$ which satisfy $I(V)_\infty = (f(z))$ in Section 5 of [24] (when V_0 is discrete of rank 1). Vaquié shows in [37] and [38] how this algorithm can be extended to arbitrary valuations V_0 of K .

Suppose V_1, \dots, V_k is an augmented sequence of inductive valuations in $K[z]$. Expand

$$f = f_m \varphi_k^m + \dots + f_0$$

as in (5). Define the projection of V_k by $\text{proj}(V_k) = \alpha - \beta$ where α is the largest and β is the smallest amongst the exponents j for which $V_k(f(z)) = V_k(f_j \varphi_k^j)$. A k -th approximant V_k to $f(z)$ over V_0 is a k -th stage homogeneous (the key polynomial φ_i is homogeneous in V_{i-1} for $i \leq k$) inductive valuation which is an extension of V_0 and which has a positive projection (Definition 3.3 [24]).

First approximants V_1 to f are defined as $V_1 = [V_0; V_1(\varphi_1) = \mu_1]$, where $\varphi_1 = z$ and μ_1 is chosen so that $\text{proj}(V_1) > 0$. MacLane shows in Lemma 3.4 [24] that if V_k is a k -th approximant to $f(z)$, then so is V_i for $i = 1, \dots, k-1$. Further, $\varphi_k | f$ in V_{k-1} and $V_k(f(z)) > V_{k-1}(f(z)) > \dots > V_1(f(z))$. In Theorem 10.1 [24], MacLane shows that if V_0 is a discrete valuation of rank 1 then every extension of V_0 to a valuation of $K[z]/(f(z))$ is an augmented sequence of finite length of approximants V_1, \dots, V_k such that $V_k(f(z)) = \infty$ or a limit of an augmented sequence of approximants of infinite length such that $V_\infty(f(z)) = \infty$. If V_0 is not discrete of rank 1, then there is the possibility that the algorithm will have to be continued to construct a pseudo valuation W of $K[z]$ with $W(f(z)) = \infty$. If this last case occurs, then the situation becomes quite complicated, as we must then extend the family $\{V_k \mid k \in \mathbf{Z}_+\}$ to a "simple admissible family" and possibly make some jumps. This is shown by Vaquié in Theorem 2.5 [36] and is explained in Section 7. An essential point is that for every construction V_1, \dots, V_k of approximants to f over V_0 by MacLane's algorithm, there exists an extension W of V_0 to a pseudo valuation of $K[x]$ such that $I(W)_\infty = (f(z))$ and $W(\varphi_k) = V_k(\varphi_k)$ for all k (This will be deduced from Theorem 1 [38] in Theorem 3.4).

We will assume now that V_0 has rank 1, so we may assume that G_{V_0} is an ordered subgroup of \mathbf{R} . We will now look a little more at the case where we have an infinite sequence of approximants, leading to a limit valuation V_∞ . In this case, there exists k_0 such that $\varphi_k = \varphi_{k_0} + h_k$ with $\deg_z h_k < \deg_z \varphi_{k_0}$ for $k \geq k_0$. Thus for $k > k_0$,

$$V_k(\varphi_k) > V_{k-1}(\varphi_k) \geq V_{k-1}(\varphi_{k-1}).$$

Thus $\lim_{k \rightarrow \infty} V_k(\varphi_k)$ exists, and is either equal to ∞ or an element of \mathbf{R} .

Lemma 3.2. *Suppose that V_0 has rank 1 and V_1, \dots, V_k, \dots is an infinite sequence of approximants to f over V_0 . Then the following are equivalent:*

- 1) $V_\infty = \lim_{k \rightarrow \infty} V_k$ is a pseudo valuation on $K[z]$ (but not a valuation).
- 2) $I^{K[z]}(V_\infty)_\infty = (f(z))$.
- 3) $\lim_{k \rightarrow \infty} V_k(\varphi_k) = \infty$.

Proof. We first prove 1) implies 3). By assumption, there exists $0 \neq h \in I(V_\infty)_\infty$. There exists k_0 such that for $k \geq k_0$, $\deg_z \varphi_k = \deg_z \varphi_{k_0}$. Expand

$$h = h_m \varphi_{k_0}^m + h_{m-1} \varphi_{k_0}^{m-1} + \dots + h_0$$

with $\deg_z h_i < \deg_z \varphi_{k_0}$ for all i and $h_m \neq 0$. There exists $\lambda \in \mathbf{Z}_+$, $1 \leq \lambda \leq \deg \varphi_{k_0}$ such that $\deg_z z^\lambda h_m = \deg_z \varphi_{k_0}$ and so there exists $0 \neq \alpha \in K$ such that $\alpha z^\lambda h_m = \varphi_{k_0} + \eta_m$ with $\deg \eta_m < \deg \varphi_{k_0}$. This implies that $\alpha z^\lambda h$ has an expansion

$$\alpha z^\lambda h = \varphi_{k_0}^{m+1} + \eta_m \varphi_{k_0}^m + \alpha z^\lambda h_{m-1} \varphi_{k_0}^{m-1} + \cdots + \alpha z^\lambda h_{m-j} \varphi_{k_0}^{m-j} + \cdots + \alpha z^\lambda h_0$$

with $\deg_z z^\lambda h_{m-j} < 2 \deg_z \varphi_{k_0}$ for all j . Now we can expand each $\alpha z^\lambda h_{m-j} = \eta_{m-j} \varphi_{k_0} + \theta_{m-j}$, with $\deg_z \eta_{m-j}$ and $\deg_z \theta_{m-j}$ less than $\deg_z \varphi_{k_0}$, so that finally we can expand

$$\alpha z^\lambda h = \varphi_{k_0}^{m+1} + h'_m \varphi_{k_0}^m + \cdots + h'_{m+1-j} \varphi_{k_0}^{m+1-j} + \cdots + h'_0$$

with $\deg_z h'_{m+1-j} < \deg_z \varphi_{k_0}$ for all j . Thus, substituting $\alpha z^\lambda h \in I(V_\infty)_\infty$ for h and continuing to denote by m the degree of its expansion in φ_{k_0} , we may assume that $h_m = 1$. The same argument shows that for $k \geq k_0$ there exist $h_i(k) \in K[z]$ for $i < m$ such that

$$h = \varphi_k^m + h_{m-1}(k) \varphi_k^{m-1} + \cdots + h_0(k)$$

with $\deg_z h_j(k) < \deg_z \varphi_k$. Now by definition of V_k we have

$$V_k(h) \leq m V_k(\varphi_k)$$

for $k \geq k_0$, so $\lim_{k \rightarrow \infty} V_k(\varphi_k) = \infty$.

We now prove that 3) implies 2). In the expansion

$$f = f_m \varphi_k^m + \cdots + f_0$$

with $\deg_z f_i < \deg_z \varphi_k$, we have that at least two distinct terms have the same value

$$V_k(f(z)) = \min_i \{V_{k-1}(f_i) + i V_k(\varphi_k)\}.$$

Thus $V_k(f(z)) \geq V_k(\varphi_k)$ for all k , which implies

$$\lim_{k \rightarrow \infty} V_k(f(z)) = \infty$$

so that $f \in I(V_\infty)_\infty$. Now $I(V_\infty)_\infty$ is a proper principal ideal in $K[z]$ and f is irreducible in $K[z]$ so $I(V_\infty)_\infty = (f(z))$.

Finally, 2) implies 1) follows since $I(V_\infty)_\infty \neq (0)$. \square

We observe that if the equivalent conditions of Lemma 3.2 hold and $g \in K[z]$ is such that $f \nmid g$, then there exists k such that $V_k(g) = V_\infty(g)$. This follows since we can find a φ_k such that $V_k(\varphi_k) = V_\infty(\varphi_k) > V_k(g)$. Then, expanding $g = g_m \varphi_k^m + \cdots + g_0$ with $\deg_z g_i < \deg_z \varphi_k$, we have that $V_\infty(g) = V_k(g) = V_k(g_0)$.

For the rest of this section, we will assume that V_0 has arbitrary rank. MacLane gives the following explanation of how to find all of the extensions of a $(k-1)$ -st stage approximant V_{k-1} to f over V_0 to a k -th stage approximant V_k to f over V_0 .

We say that $e \in K[z]$ is an ‘‘equivalence unit’’ for V_k if there exists an ‘‘equivalence-reciprocal’’ $h \in K[z]$ such that $eh \sim 1$ in V_k . It is shown in Section 4 of [24] that e is an equivalence unit if and only if e is equivalent in V_k to a polynomial g such that $\deg_z g < \deg_z \varphi_k$.

By Theorem 4.2 [24], f has an essentially unique (unique up to equivalence in V_{k-1}) expression

$$(11) \quad f \sim e \varphi_{k-1}^{m_0} \psi_1^{m_1} \cdots \psi_t^{m_t}$$

in V_{k-1} , with $m_0 \in \mathbf{N}$ and $m_1, \dots, m_t > 0$. Here e is an equivalence unit for V_{k-1} and ψ_1, \dots, ψ_t are homogeneous key polynomials over V_{k-1} all not equivalent to φ_{k-1} in V_{k-1}

and not equivalent in V_{k-1} to each other. We have that $t > 0$ since $\text{proj}(V_{k-1}) > 0$. We have that φ_{k-1} is a homogeneous key polynomial in V_{k-1} by Lemma 4.3 [24].

If f is a homogeneous key polynomial for V_{k-1} , then $V_k = [V_{k-1}; V_k(f(z)) = \infty]$ is a pseudo valuation of $K[z]$ with $I(V)_\infty = (f(z))$.

If f is not a homogeneous key polynomial for V_{k-1} , then none of the ψ_i are equal to f , and we may define a k -th stage approximant to f over V_0 which is an inductive valuation of V_{k-1} by $V_k = [V_{k-1}; V_k(\varphi_k) = \mu_k]$ where φ_k is one of the ψ_i . In the expansion (5) of f ,

$$f = f_m \varphi_k^m + \cdots + f_0$$

μ_k must be chosen so that $\text{proj}(V_k) > 0$. All k -th stage approximants V_k to f extending V_{k-1} are found by the above procedure.

Let $T = \mathbf{R} \times G_{V_0}$. Given $\alpha, \beta \in G_{V_0}$ and $q \in \mathbf{R}$, we have the line

$$D = \{(x, \gamma) \in T \mid q\gamma + \alpha x + \beta = 0\}$$

in T . When $q \neq 0$, we define the slope of D to be $-\frac{\alpha}{q} \in G_{V_0} \otimes_{\mathbf{Z}} \mathbf{R}$. Associated to D are two half spaces of T ,

$$H_{\geq}^D = \{(x, \gamma) \in T \mid q\gamma + \alpha x + \beta \geq 0\}$$

and

$$H_{\leq}^D = \{(x, \gamma) \in T \mid q\gamma + \alpha x + \beta \leq 0\}.$$

Given a subset A of T , the convex closure of A is $\text{Conv}(A) = \cap H$ where H runs over the half spaces of T which contain A .

The Newton polygon is constructed as on page 500 [24] and page 2510 [38]. These constructions are equivalent but slightly different. We use the convention of [24]. The possible values μ_k can be conveniently found from the Newton polygon $N(V_{k-1}, \varphi_k)$. This is constructed by taking the convex closure in T of

$$A = \{(m - i, \delta) \mid \delta \geq V_{k-1}(f_i), 0 \leq i \leq m\},$$

where the union is over i such that $f_i \neq 0$. A segment F of $\text{Conv}(A)$ is a subset F of $\text{Conv}(A)$ which is defined by $F = \text{Conv}(A) \cap D$ where D is a line of T such that $\text{Conv}(A)$ is contained in one of the half spaces H_{\geq}^D or H_{\leq}^D defined by D and $F = \text{Conv}(A) \cap D$ contains at least two distinct points.

The slopes of the segments of $N(V_{k-1}, \varphi_k)$ whose slope μ satisfies $\mu > V_{k-1}(\varphi_k)$ are the possible values of φ_k . The polygon composed of those segments of slope μ with $\mu > V_{k-1}(\varphi_k)$ is called the principal part of the Newton polygon $N(V_{k-1}, \varphi_k)$.

In the proof of Theorem 5.1 of [24], it is shown that for $1 \leq i \leq t$, the principal polygon of $N(V_{k-1}, \psi_i)$ (from(11)) is

$$(12) \quad \{(x, y) \in N(V_{k-1}, \psi_i) \mid x \geq m - m_i\}.$$

Further, m_0 is the smallest exponent i such that in the expansion $f = \sum f_i \varphi_{k-1}^i$ with $\deg_z f_i < \deg_z \varphi_{k-1}$, we have that $V_{k-1}(f_i \varphi_{k-1}^i) = V_{k-1}(f(z))$.

Remark 3.3. *If the coefficients of $f(z)$ are all in the valuation ring R_{V_0} of V_0 , then the coefficients of all key polynomials φ_k are also in R_{V_0} , as is established in Theorem 7.1 [24].*

The following theorem follows from a criterion of [38].

Theorem 3.4. *Suppose that V_k is a k -th approximant to f over V_0 . Then there exists a pseudo valuation W of $K[z]$ such that $W|K = V_0$, $I(W)_\infty = (f(z))$, $W(g) \geq V_k(g)$ for all $g \in K[z]$ and $W(\varphi_i) = V_i(\varphi_i)$ for $1 \leq i \leq k$.*

Proof. As explained in the above construction of V_k , we have that $\varphi_k|f$ in V_{k-1} , and there exists a key polynomial ψ for V_k with ψ not equivalent to φ_k in V_k and such that $\psi|f$ in V_k . The theorem now follows from Theorem 1 [38]. \square

4. AN ALGORITHM TO CONSTRUCT GENERATING SEQUENCES

Let V_0 be a valuation of a field K . Suppose that there exists an algebraically closed field k such that $k \subset R_{V_0}$ and $R_{V_0}/m_{V_0} \cong k$. Let $f(z) \in R_{V_0}[z]$ be an irreducible unitary polynomial.

In this section we give an inductive construction of a sequence of approximants to f over V_0 , so that the key polynomials constructed have a particularly nice form. We will call the sequence of approximants “a realization of the algorithm of Section 4”. We will prove the following theorem by induction on k .

Theorem 4.1. *Suppose that W is a pseudo valuation of $K[z]$ extending V_0 such that $I(W)_\infty = (f(z))$. Then we can construct a sequence of approximants to f over V_0*

$$(13) \quad V_1, \dots, V_k, \dots,$$

where

$$(14) \quad V_i = [V_{i-1}; V_i(\varphi_i) = W(\varphi_i)]$$

for all i such that the key polynomials φ_i satisfy $\varphi_1 = z$ in V_0 and

$$(15) \quad \varphi_i = \varphi_{i-1}^{n_{i-1}} - c_{i-1} \varphi_1^{j_1(i-1)} \dots \varphi_{i-2}^{j_{i-2}(i-1)}$$

in V_{i-1} for $2 \leq i \leq k$ with $c_{i-1} \in R_{V_0}$, $n_i = [G_{V_{i-1}} : G_{V_{i-2}}]$ and $0 \leq j_l(m) < n_l$ for all l and m . The sequence (13) is either of finite length k with $\varphi_k = f$ and $V_k(f(z)) = \infty$ or the sequence is infinite.

Observe that we have that

$$\varphi_i^{n_i} \sim c_i \varphi_1^{j_1(i)} \dots \varphi_{i-1}^{j_{i-1}(i)}$$

in V_i for $1 \leq i \leq k-1$, since φ_{i+1} is a key polynomial over V_i .

The proof of the theorem will be given after we have established Lemmas 4.2 and 4.3 and Theorem 4.4.

Lemma 4.2. *Suppose that V_1, \dots, V_k satisfy the conclusions (14) and (15) of Theorem 4.1 and we have an equality*

$$n_k V_k(\varphi_k) = V_k(c_k \varphi_1^{j_1(k)} \dots \varphi_{k-1}^{j_{k-1}(k)})$$

in V_k with $c_k \in K$, $n_k = [G_{V_k} : G_{V_{k-1}}]$ and $0 \leq j_l(k) < n_l$ for all l . Then $c_k \in R_{V_0}$.

Proof. In the case that $k = 1$, we have that $W(z) \geq 0$ since f is unitary and the coefficients of f are in R_{V_0} . Thus $V_0(c_1) \geq 0$.

Now suppose that $k \geq 2$. Since n_i is the smallest positive integer m such that $mV_i(\varphi_i) \in G_{V_{i-1}}$, we have by repeated Euclidean division that every element $\gamma \in G_{V_k}$ has a unique decomposition as

$$(16) \quad \gamma = \gamma_0 + j_1 \mu_1 + \dots + j_k \mu_k$$

where $\gamma_0 \in G_{V_0}$, $\mu_i = W(\varphi_i)$ for $1 \leq i \leq k$ and $0 \leq j_i < n_i$ for $1 \leq i \leq k$. We have from (15) that

$$(17) \quad n_i \mu_i < \mu_{i+1} \text{ for all } 1 \leq i < k.$$

There is a unique representation

$$n_l \mu_l = \gamma_0 + j_1 \mu_1 + \cdots + j_{l-1} \mu_{l-1}$$

of the form of (16). It follows from (17) that

$$j_1 \mu_1 + \cdots + j_{l-1} \mu_{l-1} < n_l \mu_l.$$

Thus $V_0(c_k) = \gamma_0 > 0$. □

Lemma 4.3. *Suppose that V_1, \dots, V_k satisfy the conclusions (14) and (15) of Theorem 4.1 with $I(V_i)_\infty = (0)$ for all $i \leq k$. Let A be a local domain whose quotient field is K and suppose that $f(z) \in A[z]$. Further suppose that A is dominated by V_0 and that A contains k (so that $A/m_A \cong k$). Suppose that $c_i \in A$ for $i \leq k-1$. Then we have a graded k -algebra isomorphism of $\text{gr}_{V_k}(A[z])$ with the quotient $\text{gr}_{V_0}(A)[\bar{\varphi}_1, \dots, \bar{\varphi}_k]/I$ of the graded polynomial ring $\text{gr}_{V_0}(A)[\bar{\varphi}_1, \dots, \bar{\varphi}_k]$ over $\text{gr}_{V_0}(A)$, where*

$$I = (\bar{\varphi}_1^{n_1} - \bar{c}_1, \bar{\varphi}_2^{n_2} - \bar{c}_2 \bar{\varphi}_1^{j_1(2)}, \dots, \bar{\varphi}_{k-1}^{n_{k-1}} - \bar{c}_{k-1} \bar{\varphi}_1^{j_1(k-1)} \cdots \bar{\varphi}_{k-2}^{j_{k-2}(k-1)}).$$

Here $\bar{c}_1, \dots, \bar{c}_{k-1}$ are the initial forms of c_1, \dots, c_{k-1} in $\text{gr}_{V_0}(A)$ and $\bar{\varphi}_i$ has the weight $V_k(\varphi_i)$ for all i .

Suppose there exists $c \in A$ and $j_i \in \mathbf{N}$ for $1 \leq i \leq k-1$ with $0 \leq j_i < n_i$ such that $V_k(\varphi_k^{n_k}) = V_k(c \varphi_1^{j_1} \cdots \varphi_{k-1}^{j_{k-1}})$. Then $(\bar{\varphi}_k^{n_k} - \bar{c} \bar{\varphi}_1^{j_1} \cdots \bar{\varphi}_{k-1}^{j_{k-1}})$ is a prime ideal in $\text{gr}_{V_k}(A[z])$.

Proof. Every $g \in A[z]$ has the unique decomposition of (6) and Remark 3.1,

$$g = \sum_j a_j \varphi_1^{m_{1,j}} \varphi_2^{m_{2,j}} \cdots \varphi_k^{m_{k,j}}$$

with $a_j \in A$, $m_{1,j}, \dots, m_{k,j} \in \mathbf{N}$ and $0 \leq m_{i,j} < n_i$ for $i < k$ and

$$\begin{aligned} V_k(g) &= \min_j \{V_0(a_j) + m_{1,j} V_1(\varphi_1) + \cdots + m_{k,j} V_k(\varphi_k)\} \\ &= \min_j \{V_k(a_j) + m_{1,j} V_k(\varphi_1) + \cdots + m_{k,j} V_k(\varphi_k)\} \end{aligned}$$

by (9).

Since $\text{gr}_{V_k}(A[z])$ is generated by the initial forms of elements of $A[z]$, the natural graded $\text{gr}_{V_0}(A)$ -algebra map

$$\Psi : \text{gr}_{V_0}(A)[\bar{\varphi}_1, \dots, \bar{\varphi}_k] \rightarrow \text{gr}_{V_k}(A[z])$$

is a surjection and I is contained in the kernel. A homogeneous element G of $\text{gr}_{V_0}[\bar{\varphi}_1, \dots, \bar{\varphi}_k]$ has a unique representation

$$G \equiv \bar{c} \bar{\varphi}_1^{j_1} \cdots \varphi_{k-1}^{j_{k-1}} \varphi_k^{j_k} \pmod{I}$$

with $c \in A$, $j_1, \dots, j_k \in \mathbf{N}$ and $0 \leq j_i < n_i$ for $i < k$. Now $\Psi(G) = 0$ implies that $c = 0$ which implies that $G \equiv 0 \pmod{I}$. Thus Ψ is an isomorphism, and the first statement of the lemma follows.

We now prove the second statement. Let

$$\psi = \bar{\varphi}_k^{n_k} - \bar{c} \bar{\varphi}_1^{j_1} \cdots \bar{\varphi}_{k-1}^{j_{k-1}}.$$

We have that $\text{gr}_{V_k}(A[z]) \cong B[\bar{\varphi}_k]$ is a graded polynomial ring over the domain

$$B = \text{gr}_{V_0}(A)[\bar{\varphi}_1, \dots, \bar{\varphi}_{k-1}] / (\bar{\varphi}_1^{n_1} - \bar{c}_1, \dots, \bar{\varphi}_{k-1}^{n_{k-1}} - \bar{c}_{k-1} \bar{\varphi}_1^{j_1(k-1)} \cdots \bar{\varphi}_{k-2}^{j_{k-2}(k-1)}).$$

Let L be an algebraic closure of the quotient field of B . Choose $t \in L$ such that $t^n = \bar{c} \bar{\varphi}_1^{j_1} \cdots \bar{\varphi}_{k-1}^{j_{k-1}}$. Then giving t the weight $V_k(\varphi_k)$, we have that $B[t]$ is a graded domain

which is a free B -module of rank n_k , since $V_k(\varphi_k)$ has order n_k in G_k/G_{k-1} , and so $1, t, \dots, t^{n_k-1}$ is a B -basis of $B[t]$. We have a natural surjection of graded B -modules

$$(18) \quad B[\overline{\varphi}_k]/(\psi) \rightarrow B[t].$$

Now $B[\overline{\varphi}_k]/(\psi)$ is also a free B -module of rank n_k , as $1, \overline{\varphi}_k, \dots, \overline{\varphi}_k^{n_k-1}$ is a B -basis. Thus (18) is an isomorphism, and so $B[\overline{\varphi}_k]/(\psi)$ is a domain. \square

Suppose that G is a totally ordered Abelian group. Let $U = G \otimes_{\mathbf{Z}} \mathbf{R}$, $d \in \mathbf{Z}_+$ and $\gamma \in G$. Since \mathbf{Z} is a principal ideal domain, we have that

$$(19) \quad \left(\frac{1}{d}\mathbf{Z}\gamma\right) \cap G = \frac{1}{\overline{m}}\mathbf{Z}\gamma \quad \text{for some } \overline{m} \in \mathbf{Z}_+.$$

Indeed, we must have $\left(\frac{1}{d}\mathbf{Z}\gamma\right) \cap G = \frac{a}{d}\mathbf{Z}\gamma$ for some $a \in \mathbf{Z}_+$. Now $\gamma \in \frac{a}{d}\mathbf{Z}\gamma$ implies $a|d$, and so there exists $\overline{m} \in \mathbf{Z}_+$ such that $\frac{1}{\overline{m}} = \frac{a}{d}$.

This implies:

$$(20) \quad \frac{1}{\overline{m}}(d, \gamma) \in \mathbf{Z} \oplus G.$$

We shall need the following fact:

$$(21)$$

For $n, q \in \mathbf{Z}_+$, $\frac{q}{n}(d, \gamma) \in \mathbf{Z} \oplus G$ if and only if n divides qd and $\frac{q}{n} = \frac{e}{\overline{m}}$ for some $e \in \mathbf{Z}_+$.

For the reader's convenience, we give a proof of (21). Suppose that $\frac{q}{n}(d, \gamma) \in \mathbf{Z} \oplus G$. Then n divides qd and writing $qd = rn$ we see that $\frac{r}{d}\mathbf{Z}\gamma \subset G$ so that it follows from (19) that $\frac{r}{d} = \frac{q}{n}$ is an integral multiple of $\frac{1}{\overline{m}}$. The converse follows from (20).

Theorem 4.4. *Suppose that we have constructed approximants $V_i = [V_{i-1}, V_i(\varphi_i) = W(\varphi_i)]$ for $1 \leq i \leq k-1$ to f over V_0 satisfying the conclusions of Theorem 4.1, $V_{k-1}(\varphi_{k-1}) < \infty$ and we have an equivalence in V_{k-1}*

$$(22) \quad f \sim e\varphi_{k-1}^{m_0}\overline{\psi}_1^{m_1} \cdots \overline{\psi}_t^{m_t}$$

of the form of (11) with $m_0 \in \mathbf{N}$ and $m_1, \dots, m_t \in \mathbf{Z}_+$ such that e is an equivalence unit for V_{k-1} , $\overline{\psi}_1, \dots, \overline{\psi}_t$ are homogeneous key polynomials to f over V_{k-1} such that there are expressions

$$\overline{\psi}_i = \varphi_{k-1}^{n_{k-1}} - \varepsilon_{k-1,i}c_{k-1}\varphi_1^{j_1(k-1)} \cdots \varphi_{k-2}^{j_{k-2}(k-1)}$$

with $c_{k-1} \in R_{V_0}$ non zero, $\varepsilon_{k-1,i} \in k$ distinct and nonzero, and $0 \leq j_i(k-1) < n_i$ for all i .

Then there exists a unique $\overline{\psi}_i$ such that $W(\overline{\psi}_i) > V_{k-1}(\overline{\psi}_i)$ and setting $\varphi_k = \overline{\psi}_i$, there exists a unique segment S of the principal part of the Newton polygon $N(V_{k-1}, \varphi_k)$ which has slope $s = W(\varphi_k)$.

Defining $V_k = [V_{k-1}, V_k(\varphi_k) = W(\varphi_k)]$, we have that V_k is an approximate to f over V_0 , such that the approximants V_1, \dots, V_k satisfy the conclusions of Theorem 4.1.

Now suppose that $V_k(\varphi_k) < \infty$. The Newton polygon $N(V_{k-1}, \varphi_k)$ is computed from the expansion

$$(23) \quad f = \sum f_i \varphi_k^i$$

with $\deg_z f_i < \deg_z \varphi_k$. Let $(m - i_1, a_1)$ be the lowest point on the segment S and let $(m - i_0, a_0)$ be the highest point. Let

$$(24) \quad F_{k,s}(\varphi_k) = \sum f_i \varphi_k^i,$$

where the sum is restricted to i such that $(m - i, V_{k-1}(f_i))$ is on S . Then there exists a polynomial in φ_k

$$(25) \quad G_{k,s}(\varphi_k) = \sum g_i \varphi_k^i$$

with $g_i \in K[z]$ such that the i such that g_i is not zero are exactly the i such that f_i is a coefficient of $F_{k,s}$ and $g_i \sim f_i$ in V_{k-1} for all such i . Further, factoring the right side of (25) as a polynomial in φ_k ,

$$(26) \quad G_{k,s}(\varphi_k) = f_{i_1}^{\bar{m}+1} \varphi_k^{i_0} \psi_1^{a_1} \dots \psi_t^{a_t}$$

where

$$(27) \quad \psi_i = \varphi_k^{n_k} - \varepsilon_{k,i} c_k \varphi_1^{j_1(k)} \dots \varphi_{k-1}^{j_{k-1}(k)}$$

with $c_k \in R_{V_0}$ nonzero, $\varepsilon_{k,i} \in k$ are distinct and nonzero, $0 \leq j_i(k) < n_i$ for all i . Further, we have that $f_{i_1}^{\bar{m}+1}$ is an equivalence unit in V_k ,

$$n_k = [G_k : G_{k-1}]$$

and the ψ_i are homogeneous key polynomials in V_k . Also, there is a V_k equivalence

$$(28) \quad f \sim G_{k,s}(\varphi_k)$$

in V_k .

Proof. The fact that there exists a $\bar{\psi}_i$ such that $W(\bar{\psi}_i) > V_{k-1}(\bar{\psi}_i)$ follows from the equivalence relation (22), since $W(f(z)) = \infty$ and $W(e\varphi_{k-1}^{m_0}) = V_{k-1}(e\varphi_{k-1}^{m_0})$. Uniqueness of $\bar{\psi}_i$ follows since the $\varepsilon_{k-1,i}$ are distinct. The existence of a segment S of the principal part of the Newton polygon $N(V_{k-1}, \varphi_k)$ with slope $s = W(\varphi_k)$, follows from Theorem 3.4 and the discussion of Subsection 3.2. The fact that upon setting $\varphi_k = \bar{\psi}_i$, we have that $V_k = [V_{k-1}, V_k(\varphi_k) = W(\varphi_k)]$ is an approximate to f over V_0 then follows since $\text{proj}(V_k)$ is positive, as $W(f(z)) = \infty$, and the fact that the approximants V_1, \dots, V_k satisfy the conclusions of Theorem 4.1 follows from our assumptions on the φ_i for $i \leq k$.

Let $y = sx + r$ be the equation of the line containing the segment S , so that

$$s = \frac{a_0 - a_1}{i_1 - i_0}.$$

Let \bar{m} be the largest positive integer such that

$$(29) \quad \frac{1}{\bar{m}}(i_1 - i_0, a_0 - a_1) \in \mathbf{Z} \oplus G_{k-1}.$$

Here \bar{m} is as defined before (21), with $d = i_1 - i_0$, $\gamma = a_0 - a_1$ and $G = G_{k-1}$. Let $(\bar{b}, \bar{c}) = \frac{1}{\bar{m}}(i_1 - i_0, a_0 - a_1)$. If $V_{k-1}(f_i) - s(m - i) = r$, then

$$(m - i, V_{k-1}(f_i)) = (m - i_1, a_1) + \lambda(\bar{b}, \bar{c}) = \left(m - i_1 + \lambda\bar{b}, a_1 + \lambda \left(\frac{a_0 - a_1}{\bar{m}} \right) \right)$$

for some $\lambda \in \mathbf{N}$ with $0 \leq \lambda \leq \bar{m}$ (this follows from (21)). Using the relations (15) for $2 \leq i \leq k$, there exists

$$h = c_k \varphi_1^{j_1(k)} \dots \varphi_{k-1}^{j_{k-1}(k)} \in K[z]$$

with $c_k \in K$ and $0 \leq j_l(k) < n_l$ for $1 \leq l < k$ such that $V_{k-1}(h) = \frac{a_0 - a_1}{\bar{m}}$. We have that

$$\begin{aligned} F_{k,s} &= \sum_{\tau=0}^{\bar{m}} f_{i_0+\tau\bar{b}} \varphi_k^{i_0+\tau\bar{b}} \\ &= \varphi_k^{i_0} \left(\sum_{\tau=0}^{\bar{m}} f_{i_0+\tau\bar{b}} \varphi_k^{\tau\bar{b}} \right) \end{aligned}$$

where

$$(30) \quad \begin{aligned} V_{k-1}(f_{i_0+\tau\bar{b}}) &= s(m - (i_0 + \tau\bar{b})) + r = -\tau \left(\frac{a_0 - a_1}{\bar{m}} \right) + a_0 \\ &= (\bar{m} - \tau) \left(\frac{a_0 - a_1}{\bar{m}} \right) + a_1 = V_{k-1}(h^{\bar{m}-\tau}) + V_{k-1}(f_{i_1}). \end{aligned}$$

By (30), and since V_0 is rational ($R_{V_0}/m_{V_0} = k$), there exist $\gamma_\tau \in k$ such that

$$(31) \quad \gamma_\tau \text{In}(h^{\bar{m}-\tau}) \text{In}(f_{i_1}) = \text{In}(f_{i_0+\tau\bar{b}})$$

in $\text{gr}_{V_{k-1}}(K[z])$. Define $G_{k,s}(\varphi_k)$ by

$$\begin{aligned} G_{k,s}(\varphi_k) &= \varphi_k^{i_0} \left(\sum_{\tau=0}^{\bar{m}} \gamma_\tau h^{\bar{m}-\tau} f_{i_1} \varphi_k^{\tau\bar{b}} \right) \\ &= f_{i_1}^{\bar{m}+1} \varphi_k^{i_0} h^{\bar{m}} \left(\sum_{\tau=0}^{\bar{m}} \gamma_\tau (h^{-1} \varphi_k^{\bar{b}})^\tau \right) \\ &= f_{i_1}^{\bar{m}+1} \varphi_k^{i_0} h^{\bar{m}} \prod_{j=1}^{\bar{m}} ((h^{-1} \varphi_k^{\bar{b}}) - \alpha_j) \\ &= f_{i_1}^{\bar{m}+1} \varphi_k^{i_0} \prod_{j=1}^{\bar{m}} (\varphi_k^{\bar{b}} - \alpha_j h) \end{aligned}$$

for suitable nonzero $\alpha_j \in k$.

We will compute the order

$$[G_k : G_{k-1}] = [(G_{k-1} + s\mathbf{Z}) : G_{k-1}].$$

We will show that the order n_k is $n_k = \bar{b} = \frac{i_1 - i_0}{\bar{m}}$. Since $s = \frac{a_0 - a_1}{i_1 - i_0}$,

$$\bar{b}s = \frac{a_0 - a_1}{\bar{m}} \in G_{k-1}.$$

Now with a as defined in the analysis preceding this proposition, with $d = i_1 - i_0$, $\gamma = a_0 - a_1$ and $G = G_{k-1}$, we have that $a = \frac{d}{\bar{m}} = \bar{b}$.

Suppose $n \in \mathbf{Z}_+$ and $ns \in G_{k-1}$. Now

$$ns = n \left(\frac{a_0 - a_1}{i_1 - i_0} \right) = \frac{n}{d} \gamma \in G_{k-1}.$$

which implies that $a = \bar{b} \mid n$.

Thus we have that

$$n_k = \bar{b} = [G_k : G_{k-1}].$$

We now have that $c_k \in R_{V_0}$ by Lemma 4.2.

The fact that $f \sim G_{k,s}$ in V_k follows since $\gamma_\tau h^{\bar{m}-\tau} f_{i_1} \sim f_{i_0+\tau\bar{b}}$ in V_k , which follows from (31), the facts that by (9),

$$V_k(h) = V_k(c_k) + \sum_{i=1}^{k-1} j_i(k) V_k(\varphi_i) = V_0(c_k) + \sum_{i=1}^{k-1} j_i(k) V_i(\varphi_i) = V_{k-1}(h)$$

and

$$V_k(f_i) = V_{k-1}(f_i)$$

for all i since $\deg_z f_i < \deg_z \varphi_k$.

We know that φ_k is a key polynomial in V_k as discussed after (11). Finally, we verify that each $\psi_i = \varphi_k^{n_k} - \varepsilon_{k,i} c_k \varphi_1^{j_1(k)} \cdots \varphi_{k-1}^{j_{k-1}(k)}$ is a key polynomial in V_k . By Lemma 4.3, the ideal

$$(\text{In}(\psi_i)) = (\bar{\varphi}_k^{n_k} - \varepsilon_{k,i} \bar{c}_k \bar{\varphi}_1^{j_1(k)} \cdots \bar{\varphi}_{k-1}^{j_{k-1}(k)})$$

is a prime ideal in $\text{gr}_{V_k}(R_{V_0}[z])$, where $\bar{c}_k = \text{In}(c_k)$, and $\bar{\varphi}_i = \text{In}(\varphi_i)$. Thus ψ_i is equivalence irreducible in V_k as a polynomial in $R_{V_0}[z]$. Since every non zero element of $R_{V_0}[z]$ is a unit in K this implies that it is equivalence irreducible in V_k as a polynomial in $K[z]$. We have that ψ_i is minimal in V_k by (8). Since ψ_i has the leading coefficient 1 and $\deg_z \psi_i > 0$, we have that ψ_i is a key polynomial over V_k . \square

We now give the proof of Theorem 4.1. Set $\varphi_1 = z$ and $V_1 = [V_0; V_1(\varphi_1) = W(\varphi_1)]$, which is an approximant to f over V_0 since $W(f(z)) = \infty$. By a simplification of the proof of Theorem 4.4, we have that $f \sim ez^{m_0}\overline{\psi}_1^{m_1} \cdots \overline{\psi}_t^{m_t}$ in V_1 , where e is an equivalence unit in V_1 and $\overline{\psi}_i = z^{n_i} - \varepsilon_{1,i}c_1$ with $c_1 \in R_{V_0}$ and $\varepsilon_{1,i} \in k$ are nonzero and distinct.

Now the conclusions of the theorem follow from induction using Theorem 4.4.

Proposition 4.5. *Suppose that there is a unique extension of V_0 to a pseudo valuation W of $K[z]$ with $I(W)_\infty = (f(z))$ and we have constructed a finite or infinite sequence of approximants V_1, \dots, V_k, \dots to f over V_0 satisfying the conclusions of Theorem 4.1. Then we have that for $k \geq 2$, with notation as in (26), setting $e_k = i_0$,*

$$(32) \quad f \sim \varphi_k^{e_k} \text{ in } V_{k-1}$$

where

$$(33) \quad \varphi_{k+1} = \varphi_k^{n_k} - c_k \varphi_1^{j_1(k)} \cdots \varphi_{k-1}^{j_{k-1}(k)}$$

with $c_k \in R_{V_0}$ nonzero, $0 \leq j_i(k) < n_i$ for all i and

$$(34) \quad \deg_z f = e_k \deg_z \varphi_k.$$

Proof. We use the notation of the statement and proof of Theorem 4.4. By Theorem 3.4, every realization of the algorithm to construct a k -th stage approximant V_k to f over V_0 extends to the construction of a pseudo valuation U extending V_0 with $I(U)_\infty = (f(z))$. Since W is unique, every realization of the algorithm must extend to the construction of $U = W$.

We will prove that the following equations,

$$(35) \quad f \sim \varphi_k^{e_k} \text{ in } V_{k-1} \text{ with } \deg_z f = e_k \deg_z \varphi_k$$

and for all $k \geq 2$, (25) of Theorem 4.4 satisfies

$$(36) \quad f \sim \psi_1^{a_1} \cdots \psi_t^{a_t} \text{ in } V_k \text{ with } \deg_z f = a_1 \deg_z \psi_1 + \cdots + a_t \deg_z \psi_t.$$

We will establish (35) and (36) for $k = 2$. Since the extension is unique, every realization of the algorithm must extend to the construction of W , so $N(V_0; \varphi_1 = z)$ has a unique segment. Let $\mu_1 = s = W(z)$ be the slope of this segment, so that

$$V_1 = [V_0; V_1(\varphi_1) = \mu_1].$$

Expand

$$f = z^d + f_{d-1}z^{d-1} + \cdots + f_0$$

with $f_i \in K$. Since $N(V_0, \varphi_1)$ has a unique segment, $i_0 = 0$, $i_1 = d$ and $f_{i_1} = 1$ in (26) for $k = 1$, so by (28) and (26) for $k = 1$,

$$(37) \quad f \sim G_{1,s}(\varphi_1) = \psi_1^{a_1} \cdots \psi_t^{a_t}$$

in V_1 , where

$$(38) \quad \psi_i = \varphi_1^{n_i} - \varepsilon_{1,i}c_1$$

from (27). Suppose that $t > 1$. Any choice of ψ_i is a key polynomial for V_1 , and if $W_2 = [V_1; V_2(\psi_i) = \mu_2]$ is an approximant extending V_1 , then since every realization of the algorithm must extend to the construction of W , we have that

as observed in the first part of the proof,

$$W(\psi_i) = W_2(\psi_i) = \mu_2 > n_1 V_1(\varphi_1) = V_0(c_1).$$

For $j \neq i$,

$$\psi_j = \psi_i + (\varepsilon_{1,i} - \varepsilon_{1,j})c_1$$

so for $j \neq i$,

$$W(\psi_j) = W(\psi_i + (\varepsilon_{1,i} - \varepsilon_{1,j})c_1) = V_0(c_1).$$

This contradiction shows that $t = 1$ in (37) and so $f \sim \varphi_2^{e_2}$ in V_1 with $\deg_z f = e_2 \deg_z \varphi_2$, establishing (35) for $k = 2$.

From (35) for $k = 2$, we have that there is an expression

$$f = \varphi_2^{e_2} + f_{e_2-1} \varphi_2^{e_2-1} + \cdots + f_0$$

with $\deg_z f_i < \deg_z \varphi_2$ for all i . From (12), we then have that the principal part of the Newton polygon $N(V_1, \varphi_2)$ is the entirety of $N(V_1, \varphi_2)$. Further, by uniqueness of the extension of V_0 , we have that $N(V_1, \varphi_2)$ has a unique segment, so $i_0 = 0$, $i_1 = e_2$ and $f_{i_1} = 1$ in (26) for $k = 2$, so

$$f \sim G_{2,s}(\varphi_2) = \psi_1^{a_1} \cdots \psi_t^{a_t}$$

in V_2 with the ψ_i given by (27) for $k = 2$, establishing (36) in V_2 for $k = 2$, with $\deg_z f = a_1 \deg_z \psi_1 + \cdots + a_t \deg_z \psi_t$.

Now by induction on k , repeating the argument for the case $k = 2$ with the application of Theorem 4.4, we obtain the conclusions of Proposition 4.5. \square

Formulas (32) and (34) also follow from Theorem 3.1 [38], and then formula (33) follows from Theorem 4.4.

5. WHEN THE DEGREE IS PRIME TO p AND THE EXTENSION IS UNIQUE

Theorem 5.1. *Suppose that A is a local domain which contains an algebraically closed field k such that $A/m_A \cong k$. Let K be the quotient field of A and suppose that V_0 is a valuation of K which dominates A , such that the residue field of the valuation ring of V_0 is k . Suppose that $f(z) \in A[z]$ is unitary and irreducible, there is a unique extension of V_0 to a valuation ω of $K[z]/(f(z))$ and the characteristic p of k does not divide $\deg_z f$. Let W be the associated pseudo valuation of $K[z]$ such that $I(W)_\infty = (f(z))$ in $K[z]$. Then there exists a realization of the algorithm of Section 4 constructing approximants V_1, \dots, V_k to f over V_0 satisfying equations (14) and (15) for all $i \leq k$ such that $W = V_k$. We have that*

$$\deg_z f = [G_\omega : G_{V_0}] = [G_{V_k} : G_{V_0}].$$

Further, with the notation of (15), $c_i \in A$ for all $1 \leq i \leq k$, and

$$\text{gr}_\omega(A[z]/(f(z))) \cong \text{gr}_{V_0}(A)[\bar{\varphi}_1, \dots, \bar{\varphi}_{k-1}]/I$$

where

$$I = (\bar{\varphi}_1^{n_1} - \bar{c}_1, \bar{\varphi}_2^{n_2} - \bar{c}_2 \bar{\varphi}_1^{j_1(2)}, \dots, \bar{\varphi}_{k-1}^{n_{k-1}} - \bar{c}_{k-1} \bar{\varphi}_1^{j_1(k-1)} \bar{\varphi}_2^{j_2(k-1)} \cdots \bar{\varphi}_{k-2}^{j_{k-2}(k-1)})$$

is a finitely generated and presented $\text{gr}_{V_0}(A)$ -module.

Proof. Suppose by induction on i that we have constructed approximants V_1, \dots, V_i to f over V_0 satisfying equations (14) and (15) with $c_1, \dots, c_{i-1} \in A$ and that φ_i is not equal to f . By Theorem 4.4 and Proposition 4.5, $f \sim G_i = \varphi_{i+1}^{e_{i+1}}$ in V_i , with φ_{i+1} a key polynomial over V_i such that

$$(39) \quad \varphi_{i+1} = \varphi_i^{n_i} - c_i \varphi_1^{j_1(i)} \cdots \varphi_{i-1}^{j_{i-1}(i)} \text{ and } \deg_z f = e_{i+1} \deg_z \varphi_{i+1}$$

for some nonzero $c_i \in R_{V_0}$. Expanding

$$(40) \quad f = \sum f_j \varphi_i^j$$

in $K[z]$, with $\deg_z f_j < \deg_z \varphi_i$, let $F = \sum f_j \varphi_i^j$ where the sum is restricted to f_j such that $V_{i-1}(f_j) + j\mu_i$ (with $\mu_i = W(\varphi_i)$) is minimal, and expanding G_i as a polynomial in φ_i , we see that the coefficients of $G_i = \varphi_i^{n_i e_{i+1}} - e_{i+1} c_i \varphi_1^{j_1(i)} \cdots \varphi_{i-1}^{j_{i-1}(i)} \varphi_i^{n_i(e_{i+1}-1)} + \cdots$ as a polynomial in φ_i and of the coefficients f_j in the expansion $F = \sum f_j \varphi_i^j$ must be equivalent in V_{i-1} by Theorem 4.4.

Now $e_{i+1} n_i \deg_z \varphi_i = \deg_z f$, so since we assume that p does not divide $\deg_z f$, we have that p does not divide e_{i+1} . Comparing the expansions of F and G_i , we see that

$$0 \neq f_{n_i(e_{i+1}-1)} \sim g_{n_i(e_{i+1}-1)} = -e_{i+1} c_i \varphi_1^{j_1(i)} \cdots \varphi_{i-1}^{j_{i-1}(i)}$$

in V_{i-1} . Since $\deg_z f_{n_i(e_{i+1}-1)} < \deg_z \varphi_i$ and $c_1, \dots, c_{i-1} \in A$ by induction, by (6) and Remark 3.1, $f_{n_i(e_{i+1}-1)}$ has a unique expansion (with only finitely many terms)

$$(41) \quad f_{n_i(e_{i+1}-1)} = \sum_{\alpha \geq 1} a_{\sigma_1(\alpha), \dots, \sigma_{i-1}(\alpha)} \varphi_1^{\sigma_1(\alpha)} \cdots \varphi_{i-1}^{\sigma_{i-1}(\alpha)}$$

with

$$W(a_{\sigma_1(\alpha), \dots, \sigma_{i-1}(\alpha)} \varphi_1^{\sigma_1(\alpha)} \cdots \varphi_{i-1}^{\sigma_{i-1}(\alpha)}) < W(a_{\sigma_1(\alpha+1), \dots, \sigma_{i-1}(\alpha+1)} \varphi_1^{\sigma_1(\alpha+1)} \cdots \varphi_{i-1}^{\sigma_{i-1}(\alpha+1)})$$

for all α , $0 \leq \sigma_l(\alpha) < n_l$ for $1 \leq l \leq i-1$ and $a_{\sigma_1(\alpha), \dots, \sigma_{i-1}(\alpha)} \in A$. Thus the minimum value term in V_{i-1} in this expansion is

$$a_{\sigma_1(1), \dots, \sigma_{i-1}(1)} \varphi_1^{\sigma_1(1)} \cdots \varphi_{i-1}^{\sigma_{i-1}(1)}$$

and so

$$j_l(i) = \sigma_l(1) \text{ for } 1 \leq l \leq i-1$$

and

$$-e_{i+1} c_i \sim a_{\sigma_1(1), \dots, \sigma_{i-1}(1)}$$

in V_0 . Replacing c_i with $-\frac{1}{e_{i+1}} a_{\sigma_1(1), \dots, \sigma_{i-1}(1)}$ in (39), we have that $c_i \in A$.

Suppose $n_i = 1$, so that $e_{i+1} = e_i$. Then substituting (39) and (41) into (40), we obtain

$$f = \varphi_{i+1}^{e_i} + \left(\sum_{k \geq 2} a_{\sigma_1(k), \dots, \sigma_{i-1}(k)} \varphi_1^{\sigma_1(k)} \cdots \varphi_{i-1}^{\sigma_{i-1}(k)} \right) \varphi_{i+1}^{e_i-1} + \sum_{j=2}^{e_i-2} f'_j \varphi_{i+1}^j$$

where $\deg_z f'_j < \deg_z \varphi_{i+1} = \deg_z \varphi_i$ for all j . Since (41) is a finite sum, we can only have $n_i = 1$ for finitely many consecutive i .

Since $\deg_z f = e_i n_1 \cdots n_{i-1}$ for all i , we must have that the algorithm terminates in a finite number of iterations k . We then have that $\varphi_k = f$ and $W = V_k$.

The final statement on the structure of $\text{gr}_\omega(A[z]/(f(z)))$ now follows from Lemma 4.3. \square

As an immediate consequence of Theorem 5.1, we have the following example, which allows us to easily compute the associated graded rings and valuation semigroups of many examples, including the rational double point singularities in dimension two, since the semigroups of valuations dominating two dimensional regular local rings are completely known ([32]. [9]).

Example 5.2. *Let k be an algebraically closed field of characteristic $p \neq 2$, and $A = k[[x_1, \dots, x_n]]$ be a power series ring over k . Let $f(z) = z^2 + az + b$ with $a, b \in m_A$ be irreducible and let $B = A[z]/(f(z))$. Suppose that ν is a valuation of the quotient field of A which dominates A and such that $R_\nu/m_\nu = k$.*

Suppose that ν has a unique extension ω to the quotient field of B which dominates B . Then there exists $g \in m_A$ such that setting $\bar{z} = z - g$, we have that

- 1) $\omega(\bar{z})$ is a generator of $G_\omega/G_\nu \cong \mathbf{Z}_2$ and
- 2) $\text{gr}_\omega(B) = \text{gr}_\nu(A)[\text{in}(\bar{z})] \cong \text{gr}_\nu(A)[\bar{\varphi}]/(\bar{\varphi}^2 - \bar{c})$ for some $\bar{c} \in \text{gr}_\nu(A)$.

In contrast, if ν does not have a unique extension to the quotient field of B which dominates B , then it can happen that $\text{gr}_\omega(B)$ is not a finitely generated $\text{gr}_\nu(A)$ -module (as will follow from Example 11.4).

The good conclusions of Theorem 5.1 may fail if either the extension is not unique or p divides $\deg_z f$. In Example 8.1 [35], an example of Guillaume Rond is presented which shows that the conclusions of Theorem 5.1 may fail if the extension of V_0 to a valuation of $K[z]/(f(z))$ is not unique and $p \nmid \deg_z f$.

Example 5.3. *The conclusions of Theorem 5.1 may fail if the characteristic p of the field k divides the degree of $f(z)$. In our example, $f(z)$ is separable and V_0 has a unique extension to $K[z]/(f(z))$.*

We now give the construction of the example. Let k be an algebraically closed field of characteristic 2 and let $A = k[x_1, x_2]_{(x_1, x_2)}$ be a localization of a two dimensional polynomial ring over k . Let K be the quotient field of A . Let V_0 be the rank 1 valuation on K defined by $V_0(x_1) = 1$ and $V_0(x_2) = \sqrt{37}$, so that $G_{V_0} = \mathbf{Z} + \sqrt{37}\mathbf{Z}$. Let

$$f(z) = z^4 + x_1^{317}z + x_1^4x_2^2 + x_2^{31}.$$

We have that $f(z)$ is an irreducible, separable polynomial in $K[z]$.

Setting $\varphi_1 = z$, we have that the Newton polygon $N(V_0, \varphi_1)$ has only one segment, from $(0, 0)$ to $(4, 4 + 2\sqrt{37})$. The slope of this segment is $1 + \frac{1}{2}\sqrt{37}$, giving the first step approximant to f over V_0 , $V_1 = [V_0; V_1(\varphi_1) = 1 + \frac{1}{2}\sqrt{37}]$. We have that $G_{V_1} = \mathbf{Z} + \frac{\sqrt{37}}{2}\mathbf{Z}$.

Now $f \sim (z^2 + x_1^2x_2)^2$ in V_1 and $V_1(z) \notin G_{V_0}$ so $\varphi_2 = z^2 + x_1^2x_2$ is a key polynomial over V_1 . We have that

$$f = \varphi_2^2 + x_1^{317}z + x_2^{31}$$

so the principal part of $N(V_1, \varphi_2)$ is equal to $N(V_1, \varphi_2)$, which has only one segment, from $(0, 0)$ to $(2, 31\sqrt{37})$. The slope is $\frac{31}{2}\sqrt{37}$, giving the 2-nd step approximant to f over V_0 , $V_2 = [V_1; V_2(\varphi_2) = \frac{31}{2}\sqrt{37}]$, with $G_{V_2} = G_{V_1}$. We have that

$$f = (\varphi_2 + zx_1^{-1}x_2^{15})^2 + \varphi_2x_1^{-2}x_2^{30} + x_1^{317}z$$

so that $f \sim (\varphi_2 + zx_1^{-1}x_2^{15})^2$ in V_2 . Thus

$$(42) \quad \varphi_3 = \varphi_2 + zx_1^{-1}x_2^{15}$$

is a key polynomial for V_2 . We have that

$$f = \varphi_3^2 + x_1^{-2}x_2^{30}\varphi_3 + x_1^{-3}x_2^{45}z + x_1^{317}z$$

so the principal part of $N(V_2, \varphi_3)$ is equal to $N(V_2, \varphi_3)$ which has only one segment, from $(0, 0)$ to $(2, \frac{91}{2}\sqrt{37} - 2)$. The slope is $\frac{91}{4}\sqrt{37} - 1$, giving the 3-rd stage approximant to f over V_0 , $V_3 = [V_2; V_3(\varphi_3) = \frac{91}{4}\sqrt{37} - 1]$, with

$$G_{V_3} = G_{V_1} + \left(\frac{91}{4}\sqrt{37} - 1 \right) \mathbf{Z} = \mathbf{Z} + \frac{\sqrt{37}}{4}\mathbf{Z}.$$

Now $f \sim \varphi_3^2 + x_1^{-3}x_2^{45}z$ in V_3 and $V_3(\varphi_3) \notin G_{V_1}$, so $\varphi_4 = \varphi_3^2 + x_1^{-3}x_2^{45}z$ is a key polynomial over V_3 . We have that

$$f = \varphi_4 + x_1^{-2}x_2^{30}\varphi_3 + x_1^{317}z$$

so the principal part of $N(V_3, \varphi_4)$ is $N(V_3, \varphi_4)$, which has only one segment, from $(0, 0)$ to $(1, -3 + \frac{131}{4}\sqrt{37})$. The slope is $-3 + \frac{131}{4}\sqrt{37}$, giving the 4-th stage approximant to f over V_0 , $V_4 = [V_3; V_4(\varphi_4) = -3 + \frac{131}{4}\sqrt{37}]$. We have that $G_{V_4} = G_{V_3}$. Now $f \sim \varphi_4 + x_1^{-2}x_2^{30}\varphi_3$ in V_4 so

$$\varphi_5 = \varphi_4 + x_1^{-2}x_2^{30}\varphi_3$$

is a key polynomial over V_4 . We have that $f = \varphi_5 + x_1^{317}z$ so the principal part of $N(V_4, \varphi_5)$ is $N(V_4, \varphi_5)$, which has only one segment, from $(0, 0)$ to $(1, 318 + \frac{1}{2}\sqrt{37})$. The slope is $318 + \frac{1}{2}\sqrt{37}$, giving the 5-th stage approximant to f over V_4 , $V_5 = [V_4; V_5(\varphi_5) = 318 + \frac{1}{2}\sqrt{37}]$. We have that $G_{V_5} = G_{V_3}$.

Now $f = \varphi_5 + x_1^{317}z$ is a key polynomial for V_5 , so $V_6 = [V_5; V_6(f(z)) = \infty]$ is a pseudo valuation with $I(V_6)_\infty = (f(z))$.

Let ω be the induced extension of V_0 to $K[z]/(f(z))$. We have that $G_\omega = G_{V_3}$ and thus

$$[G_\omega : G_{V_0}] = 4 = \deg_z f = [L : K]$$

showing that ω is the unique extension of V_0 to a valuation of L , and that $\delta(\omega/V_0) = 1$, so that the extension is defectless (Section 8). Observe that we cannot avoid substitutions like (42), leaving the ring A in any realization of the algorithm. Notice that the conclusions of Theorem 5.1 are verified, if we take A_1 to be a birational extension of A containing $x_1^{-1}x_2^{15}$.

Remark 5.4. *In the example, the valuation V_0 is an Abhyankar valuation, which means that there is equality in the fundamental inequality of Abhyankar ([1, Theorem 1]),*

$$\dim_{\mathbf{Q}} G_{V_0} \otimes_{\mathbf{Z}} \mathbf{Q} + \text{trdeg}_{A/m_A} R_{V_0}/m_{V_0} = \dim A.$$

It is known ([20, Theorem 1]) that Abhyankar valuations have “no defect”, a fact which plays a role in this example. We will come back to the study of the effect of defect in Sections 6, 7, 8 and 9 below.

6. HENSELIZATION AND COMPLETION

A valued field (K, ν) is Henselian if for all algebraic extensions L of K , there exists a unique valuation ω of L which extends ν . Some references on the theory of Henselian fields are [19], [14], [30] and [37].

An extension (K^h, ν^h) of a valued field (K, ν) is called a Henselization of (K, ν) if (K^h, ν^h) is Henselian and for all Henselian valued fields (L, ω) and all embeddings $\lambda : (K, \nu) \rightarrow (L, \omega)$, there exists a unique embedding $\tilde{\lambda} : (K^h, \nu^h) \rightarrow (L, \omega)$ which extends λ .

A Henselization (K^h, ν^h) of (K, ν) can be constructed by choosing an extension ν^s of ν to a separable closure K^{sep} of K and letting K^h be the fixed field of the decomposition group

$$\{\sigma \in G(K^{\text{sep}}/K) \mid \nu^s \circ \sigma = \nu^s\}$$

of ν^s , and defining ν^h to be the restriction of ν^s to K^h ([14, Theorem 17.11]).

Lemma 6.1. *Suppose that (K, ν) is a valued field and let (K^h, ν^h) be a Henselization of (K, ν) . Suppose that $f(z) \in K[z]$ is unitary, irreducible and separable, so that $f(z)$ is reduced in $K^h[z]$. Let $f(z) = f_1(z)f_2(z) \cdots f_r(z)$ be the factorization of $f(z)$ into irreducible unitary factors in $K^h[z]$. If the coefficients of $f(z)$ are in R_ν then the coefficients of the $f_i(z)$ are in R_{ν^h} .*

Let ν_i^h be the (unique) extension of ν^h to $K^h[z]/(f_i)$. Then the distinct extensions of ν to $K[z]/(f(z))$ are the r restrictions ν_i of ν_i^h to $K[z]/(f(z))$, under the natural inclusions $K[z]/(f(z)) \rightarrow K^h[z]/(f_i(z))$.

Proof. The polynomial $f(z)$ is reduced in $K^h[z]$ since the separable polynomial $f(z)$ is reduced in $K^{\text{sep}}[z]$ where K^{sep} is a separable closure of K .

Let \bar{z} be a root of $f_i(z)$ in K^{sep} . Then $f(z)$ is the minimal polynomial of \bar{z} in $K[z]$, and $K[z]/(f(z)) \cong K[\bar{z}]$. If \bar{z} is integral over R_ν , then \bar{z} is integral over R_{ν^h} . Thus the coefficients of f_i are in R_{ν^h} since R_{ν^h} is normal (Theorem 5, page 260 [39]).

If L is a finite separable extension of K , then we have two associated sets,

$$\text{Mon}(L, K) = K\text{-embeddings of } L \text{ in } K^{\text{sep}}$$

and

$$\mathcal{E}(L, \nu) = \text{Extensions of } \nu \text{ to a valuation of } L.$$

By Lemma 1.4 [37] or Section 17 [14], the map $\Phi : \text{Mon}(L, K) \rightarrow \mathcal{E}(L, \nu)$, defined by $\Phi(\lambda) = \nu^s \circ \lambda$ is surjective, with $\Phi(\lambda) = \Phi(\lambda')$ if and only if $\lambda \sim_{K^h} \lambda'$. The equivalence \sim_{K^h} is defined by $\lambda \sim_{K^h} \lambda'$ if and only if there exists a K^h -isomorphism $\sigma : K^{\text{sep}} \rightarrow K^{\text{sep}}$ such that $\lambda' = \sigma \circ \lambda$.

The valuation $\nu^s \circ \lambda$ is obtained from the embedding

$$L \cong \lambda(L) \rightarrow \lambda(L) \cdot K^h$$

into the join of $\lambda(L)$ and K^h in K^{sep} , and the restriction of the valuation $\nu^s|_{\lambda(L) \cdot K^h}$ to L .

Let $L = K[z]/(f(z))$. The elements $\lambda \in \text{Mon}(L, K)$ are in one to one correspondence with the distinct roots α_λ of $f(z)$ in K^{sep} . We have $\lambda(L) \cdot K^h = K^h[\alpha_\lambda]$. Thus $\lambda(L) \cdot K^h \cong K^h[z]/(f_i)$ for some i . Further, $\lambda \sim_{K^h} \lambda'$ if and only if α_λ and $\alpha_{\lambda'}$ have the same minimal polynomial f_i in $K^h[z]$.

Since K^h is Henselian, for each i there is a unique extension of ν^h to $K^h[z]/(f_i)$, and so the last assertion of the lemma follows. \square

Suppose that A is a local ring and $g(z) \in A[z]$ is a polynomial. Let $\bar{g}(z) \in A/m_A[z]$ be the polynomial obtained by reducing the coefficients of $g(z)$ mod m_A .

A local ring A is a Henselian local ring if it has the following property. Let $f(z) \in A[z]$ be a unitary polynomial of degree n . If $\alpha(z)$ and $\alpha'(z)$ are relatively prime unitary polynomials in $A/m_A[z]$ of degrees r and $n-r$ respectively such that $\bar{f}(z) = \alpha(z)\alpha'(z)$, then there exist unitary polynomials $g(z)$ and $g'(z)$ in $A[z]$ of degrees r and $n-r$ respectively such that $\bar{g}(z) = \alpha(z)$, $\bar{g}'(z) = \alpha'(z)$ and $f(z) = g(z)g'(z)$.

If A is a local ring, a local ring A^h which dominates A is called a Henselization of A if any local homomorphism from A to a Henselian local ring can be uniquely extended to A^h . A Henselization always exists (Theorem 43.5 [26]). The construction is particularly nice when A is a normal local ring. Let K be the quotient field of A and Let K^{sep} be a separable closure of A . Let \bar{A} be the integral closure of A in K^{sep} and let \bar{m} be a maximal ideal of \bar{A} .

Let H be the decomposition group

$$H = G^s(\bar{A}_{\bar{m}}/A) = \{\sigma \in G(K^{\text{sep}}/K) \mid \sigma(\bar{A}_{\bar{m}}) = \bar{A}_{\bar{m}}\}.$$

Then $A^h = (\bar{A}_{\bar{m}})^H$ is the fixed ring of the action of H on $\bar{A}_{\bar{m}}$. We have

$$A^h = (\bar{A} \cap K^H)_{\bar{m} \cap (\bar{A} \cap K^H)} = \bar{A}_{\bar{m}} \cap K^H = (\tilde{A})_{\bar{m} \cap \tilde{A}}$$

where \tilde{A} is the integral closure of A in K^H .

We remark that if A is an excellent local domain with quotient field K and ν is a valuation of K which dominates A , then there exists a directed system of normal birational extensions A_i of A such that $\cup A_i = R_\nu$.

Lemma 6.2. *Continuing the assumptions of Lemma 6.1, suppose that A is an excellent local domain with quotient field K such that ν dominates A , and that A_i is a directed system of birational extensions of A such that the A_i are normal local domains which are dominated by ν and $\cup A_i = R_\nu$. Then there are natural equalities*

$$R_{\nu^h} = (R_\nu)^h = \cup A_i^h.$$

Proof. Let ν^s be an extension of ν to K^{sep} and

$$H = \{\sigma \in \text{Gal}(K^{\text{sep}}/K) \mid \nu^s \circ \sigma = \nu^s\},$$

so that $K^h = (K^{\text{sep}})^H$. Let \bar{V} be the integral closure of R_ν in K^{sep} , and let $m = \bar{V} \cap m_{\nu^s}$, a maximal ideal in \bar{V} . Since K^{sep} is algebraic over K , we have that $R_{\nu^s} = \bar{V}_m$ by Theorem 12, page 27 [40]. Now, as is shown on the bottom of page 68 of [40], H is the decomposition group

$$H = G^s(R_{\nu^s}/R_\nu) = \{\sigma \in G(K^{\text{sep}}/K) \mid \sigma(R_{\nu^s}) = R_{\nu^s}\},$$

so that

$$(R_\nu)^h = \bar{V}_m \cap K^h = R_{\nu^s} \cap K^h = R_{\nu^h},$$

establishing the first assertion of the lemma.

Suppose that A is a normal local ring with quotient field K . Let \tilde{A} be the integral closure of A in K^h . If A is dominated by $V = R_\nu$, then $\tilde{A}_{m_{\nu^s} \cap \tilde{A}}$ is dominated by $\tilde{V}_{m_{\nu^s} \cap \tilde{V}}$ (where \tilde{V} is the integral closure of V in K^h). Suppose $g, h \in \tilde{V}$ with $h \notin m_{\nu^s} \cap \tilde{V}$. Since \tilde{A}_i is a directed system, there exists i such that $g, h \in \tilde{A}_i$, so $h \notin m_{\nu^s} \cap \tilde{A}_i$ and $\frac{g}{h} \in (\tilde{A}_i)_{m_{\nu^s} \cap \tilde{A}_i}$. Thus

$$\cup_i (\tilde{A}_i)_{m_{\nu^s} \cap \tilde{A}_i} = R_\nu^h.$$

Let \bar{A}_i be the integral closure of A_i in K^{sep} . By Lemma 3.3 [4], we have inclusions of decomposition groups

$$G^s(R_{\nu^s}/R_\nu) \subset G^s((\bar{A}_i)_{m_{\nu^s} \cap \bar{A}_i}/A_i)$$

for all i , and by Lemma 3.4 [4], there exists i_0 such that

$$G^s(R_{\nu^s}/R_\nu) = G^s((\bar{A}_i)_{m_{\nu^s} \cap \bar{A}_i}/A_i)$$

for $i \geq i_0$. Thus $A_i^h \subset (\tilde{A}_i)_{m_{\nu^s} \cap \tilde{A}_i}$ for all i and $A_i^h = (\tilde{A}_i)_{m_{\nu^s} \cap \tilde{A}_i}$ for $i \gg 0$. The last assertion of the lemma now follows. \square

Let (K, ν) be a valued field such that ν has rank 1. The completion $(\hat{K}, \hat{\nu})$ (when ν has rank 1) is defined in Section 2 of [14]. The completion \hat{K} is defined to be the ring of ν -Cauchy sequences in K modulo the maximal ideal of ν -null sequences (ν -Cauchy sequences whose limit is ∞). The extension $\hat{\nu}$ of ν is defined by $\hat{\nu}(h) = \lim_{i \rightarrow \infty} \nu(h_i)$ if (h_i) is a ν -Cauchy sequence in K which converges to h . We have that \hat{K} is a Henselian field (Lemma 16.7 [14]). The following lemma is proven in Theorem 2.12 [14].

Lemma 6.3. *Suppose that (K, ν) is a rank 1 valued field and $(\hat{K}, \hat{\nu})$ is a completion of (K, ν) . Suppose that $f(z) \in K[z]$ is unitary, irreducible and separable, so that $f(z)$ is reduced in $\hat{K}[z]$. Let $f(z) = f_1(z)f_2(z) \cdots f_r(z)$ be the factorization of f into irreducible unitary factors in $\hat{K}[z]$.*

Let $\hat{\nu}_i$ be the (unique) extension of $\hat{\nu}$ to $\hat{K}[z]/(f_i)$. Then the distinct extensions of ν to $K[z]/(f(z))$ are the r restrictions ν_i of $\hat{\nu}_i$ to $K[z]/(f(z))$, under the natural inclusions $K[z]/(f(z)) \rightarrow \hat{K}[z]/(f_i)$.

Lemma 6.4. *Let notation be as in the statement of Lemma 6.3. We then have a factorization $K \rightarrow K^h \rightarrow \hat{K}$ of valued fields. Further, the factorizations of $f(z)$ into products of unitary irreducible polynomials in $K^h[z]$ and $\hat{K}[z]$ are the same.*

Proof. We have a natural inclusion of K^h into \hat{K} since \hat{K} is a Henselian field. The irreducible factors of $f(z)$ in $K^h(z)$ remain irreducible in $\hat{K}[z]$ since there is a 1-1 correspondence of the irreducible factors of $f(z)$ in $\hat{K}[z]$ with the distinct extensions of ν to $L = K[z]/(f(z))$ by Lemma 6.3 and there is a 1-1 correspondence of the irreducible factors of $f(z)$ in $K^h[z]$ with the distinct extensions of ν to L by Lemma 6.1. \square

Some references on the defect of a finite field extension are [19], [14], [30] and [37].

Suppose $(K, \nu) \rightarrow (L, \omega)$ is a finite separable extension of valued fields. Let K^{sep} be a separable closure of K with an embedding of L in K^{sep} . Let ν^s be an extension of ω to a valuation of K^{sep} . As discussed above, we can use ν^s to define the Henselization K^h of (K, ν) , with valuation $\nu^h = \nu^s|_{K^h}$, and then $L^h = L \cdot K^h$, the join of L and K^h in K^{sep} , is a Henselization of (L, ω) with valuation $\omega^h = \nu^s|_{L^h}$ (Lemma 1.3 [37], [19], [14, (17.16)]). The defect of ω over ν is defined as

$$(43) \quad \delta(\omega/\nu) = [L^h : K^h]/e(\omega^h/\nu^h)f(\omega^h/\nu^h) = [L^h : K^h]/e(\omega/\nu)f(\omega/\nu).$$

The defect is a power of the residue characteristic p of the valuation ring of ν by Ostrowski's lemma ([19, Theorem 8.2]).

7. VAQUIÉ'S ALGORITHM

Suppose that K is a field, $f(z) \in K[z]$ is unitary and irreducible, ν is a valuation of K and μ is a pseudo valuation of $K[z]$ which extends ν such that $I(\mu)_\infty = (f(z))$. Vaquié shows in Theorem 2.5 [36] that there exists a “finite admissible family of valuations” \mathcal{S} which determines μ . We will take the last element of \mathcal{S} to be the pseudo valuation μ . This result follows from Proposition 2.3 [36], which gives an algorithm for constructing such a family.

We summarize the definition of an “admissible family of valuations” approximating μ (from Section 2.1 [36]), which takes the following form since $I(\mu)_\infty = (f(z)) \neq 0$. A family \mathcal{S} of iterated augmented valuations is called a “simple admissible family” if it is of the form $\mathcal{S} = (\mu_i)_{i \in I}$ where the set of indices I is the disjoint union $I = B \amalg A$ with B a finite set and A a totally ordered set, where all elements of A are larger than all elements of B and A does not have a largest element.

A family of valuations $\mathcal{A} = (\mu_i)_{i \in I}$ is called an “admissible family” for μ (defined on page 3473 [36]) if it is a finite or countable union of simple admissible families $\mathcal{S}^{(t)} = (\mu_i^{(t)})_{i \in I^{(t)}}$. The first valuation of $\mathcal{S}^{(1)}$ is an inductive valuation of the form $\mu_1^{(1)} = [\mu_0; \mu_1^{(1)}(\varphi_1^{(1)}) = \gamma_1]$ where $\mu_0 = \nu$ is the given valuation of K and $\varphi_1^{(1)}$ is a polynomial of degree 1. For $t \geq 2$, the first valuation $\mu_1^{(t)}$ of $\mathcal{S}^{(t)}$ is a “limit augmented valuation” for the family $(\mu_{\alpha^{(t-1)}})_{\alpha \in A^{(t-1)}}$. The construction of limit augmented valuations will be explained below.

Write $I^{9t} = B^{(t)} \amalg A^{(t)}$ as above and write $B^{(t)} = \{1, \dots, n^{(t)}\}$. Then for $i \geq 2$ in $B^{(t)}$, $\mu_i^{(t)} = [\mu_{i-1}^{(t)}; \mu_i^{(t)}(\varphi_i^{(t)}) = \gamma_i^{(t)}]$ is an inductive valuation (Section 3). For $\alpha \in A^{(t)}$, we have that $\mu_\alpha^{(t)} = [\mu_{n^{(t)}}^{(t)}; \mu_\alpha^{(t)}(\varphi_\alpha^{(t)}) = \gamma_\alpha^{(t)}]$ is an inductive valuation, where $\deg_z \varphi_\alpha^{(t)} = \deg_z \varphi_{n^{(t)}}^{(t)}$.

Vaquié requires that $\deg_z \varphi_{i-1}^{(t)} < \deg_z \varphi_i^{(t)}$ for $i \geq 2$ in $B^{(t)}$ but we do not assume this. By the definition of an inductive value, we do have that $\deg_z \varphi_{i-1}^{(t)} \leq \deg_z \varphi_i^{(t)}$ for $i \geq 2$ in $B^{(t)}$. By the construction of limit key polynomials, we have that $\deg_z \varphi_{n^{(t)}}^{(t)} < \deg_z \varphi_1^{(t+1)}$ for all t .

We require that for $g \in K[z]$ and $i < j \in I$,

$$(44) \quad \mu_i(g) \leq \mu_j(g) \leq \mu(g).$$

Further, $\mu_i(\varphi_i) = \mu(\varphi_i)$ for all i .

We now discuss the construction of limit augmented valuations.

Suppose that $\mathcal{A} = (\mu_\alpha)_{\alpha \in A}$ is an admissible family of valuations for μ . Define ([36, page 3473])

$$\tilde{\Sigma}(\mathcal{A}) = \{g \in K[z] \mid \mu_\alpha(g) < \mu(g) \text{ for all } \mu_i \in \mathcal{A}\}.$$

Define $d(\mathcal{A}) = \infty$ if $\tilde{\Sigma}(\mathcal{A}) = \emptyset$ and

$$d(\mathcal{A}) = \inf\{\deg_z \varphi \mid \varphi \in \tilde{\Sigma}(\mathcal{A})\}$$

if $\tilde{\Sigma}(\mathcal{A}) \neq \emptyset$. Now define

$$\Sigma(\mathcal{A}) = \{\varphi \in \tilde{\Sigma}(\mathcal{A}) \text{ such that } \varphi \text{ is unitary and } \deg_z \varphi = d(\mathcal{A})\}$$

and

$$(45) \quad \Lambda(\mathcal{A}) = \{\mu(\varphi) \mid \varphi \in \Sigma(\mathcal{A})\}.$$

Suppose that $\Lambda(\mathcal{A})$ does not have a largest element. We then define a totally ordered index set C , which does not have a largest element, so that

$$\Lambda(\mathcal{A}) = \{\gamma_\alpha \mid \alpha \in C\},$$

where $\alpha < \beta$ if and only if $\gamma_\alpha < \gamma_\beta$.

A “limit key polynomial” φ for \mathcal{A} is defined on page 3465 of [37]. It satisfies the three properties that φ is \mathcal{A} -minimal, φ is \mathcal{A} -irreducible and φ is unitary. The elements of $\Sigma(\mathcal{A})$ are limit key polynomials for \mathcal{A} by Proposition 1.21 [36]. Choose $\varphi_\alpha \in \Sigma(\mathcal{A})$ for each $\alpha \in C$ so that $\mu(\varphi_\alpha) = \gamma_\alpha$. We then have a limit augmented valuation $\mu_\alpha = [\mathcal{A}; \mu_\alpha(\varphi_\alpha) = \gamma_\alpha]$ (Proposition 1.22 [36]), which is defined by

$$(46) \quad \mu_\alpha(g) = \max_{j \in A} \left\{ \min_i \{ \mu_j(g_i) + i \mu(\varphi_\alpha) \} \right\}$$

for $g \in K[z]$, where

$$g = \sum g_i \varphi_\alpha^i$$

with $\deg_z g_i < \deg_z \varphi_\alpha$.

The “associated family of iterated augmented valuations” to \mathcal{A} is

$$(47) \quad (\mu_\alpha)_{\alpha \in C}.$$

We will explain here how the algorithm proceeds if we are given a discrete simple admissible family $\mathcal{S} = \{\mu_1, \dots, \mu_n\}$ such that $\Sigma(\mu_n)$ is nonempty. We will produce an admissible family of valuations \mathcal{B} such that $d(\mathcal{B}) > d(\mu_n)$.

All elements of $\Sigma(\mu_n)$ are key polynomials for μ_n by Theorem 8.1 [23] or Theorem 1.15 page 3453 [37].

First suppose that the set of values $\Lambda(\mu_n)$ has a largest element γ' (which could be ∞). Then we can define $\mu' = [\mu_n; \nu'(\varphi') = \gamma']$ where $\varphi' \in \Sigma(\mu_n)$ satisfies $\mu(\varphi') = \gamma'$. We then have two cases, depending on if $\deg_z \varphi' > \deg_z \varphi_n$ or if $\deg_z \varphi' = \deg_z \varphi_n$.

Assume that $\deg_z \varphi' > \deg_z \varphi_n$. Set $\varphi_{n+1} = \varphi'$, $\gamma_{n+1} = \gamma'$ and

$$\mu_{n+1} = \mu' = [\mu_n; \mu_{n+1}(\varphi_{n+1}) = \gamma_{n+1}].$$

Then define $\mathcal{B} = \{\mu_1, \dots, \mu_n, \mu_{n+1}\}$ which is a discrete simple admissible family, with $d(\mathcal{B}) \geq \deg_z(\varphi_{n+1}) > \deg_z \varphi_n$.

Now assume that $\deg_z \varphi' = \deg_z \varphi_n$. Then define $\mathcal{B} = \{\mu_1, \dots, \mu_n, \mu'\}$ which is again a discrete simple admissible family with $d(\mathcal{B}) > d(\mu')$ (by Lemma 15.1 [23] or Corollary, page 3448 [36]).

The last case is when $\Lambda(\mu_n)$ does not have a largest element. Define the associated family of iterated augmented valuations $(\mu_\alpha)_{\alpha \in C}$ of (47) for μ_n . For all $\gamma_\alpha \in \Lambda(\mu_n)$, define $\mu_\alpha = [\mu_n; \mu_\alpha(\varphi_\alpha) = \gamma_\alpha]$. Define $\mathcal{S}^{(1)}$ by adding to \mathcal{S} the family $\mathcal{C} = (\mu_\alpha)_{\alpha \in C}$, so $\mathcal{S}^{(1)}$ is indexed by $I' = \{1, \dots, n\} \amalg C$ (which does not have a largest element). We have that $\mathcal{S}^{(1)}$ is a simple admissible family. The family \mathcal{C} is an “exhaustive, continuous family of iterated augmented valuations” with the property that $\deg_z \varphi_\alpha = d(\mu_n)$ for all $\alpha \in C$. We have that $f \notin \Sigma(\mu_n)$ since C does not have a largest element. Thus $\tilde{\Sigma}(\mathcal{C}) \neq \emptyset$. By Proposition 1.21 [36], all polynomials of $\Sigma(\mathcal{C})$ are limit key polynomials for the family \mathcal{C} . We now choose a polynomial $\varphi_1^{(2)} \in \Sigma(\mathcal{C})$, and define the “limit augmented valuation” $\mu_1^{(2)} = [(\mu_\alpha)_{\alpha \in C}; \nu_1^{(2)}(\varphi_1^{(2)}) = \mu(\varphi_1^{(2)})]$ (by the definition on page 2465 [36] and Proposition 1.22 [36] and as explained in (46)) and the discrete, simple admissible family $\mathcal{S}^{(2)} = \{\mu_1^{(2)}\}$. By Proposition 1.27 [36], $\deg_z \varphi_1^{(2)}$ is greater than the degree of the polynomials in $\Sigma(\mu_n)$. Define the admissible family $\mathcal{B} = \mathcal{S}^{(1)} \cup \mathcal{S}^{(2)}$, which is indexed by $I'' = I' \amalg \{1^{(2)}\}$ (where $1^{(2)}$ is larger than every element of I'').

7.1. Comparison of the algorithms of Section 4 and Vaquié. Suppose that W is a pseudo valuation of $K[z]$ which extends a valuation V_0 of K , such that $I(W)_\infty = (f(z))$ where f is unitary and $f(z) \in R_{V_0}[z]$. Let

$$(48) \quad V_1, \dots, V_k, \dots$$

be a sequence of approximants to f over V_0 constructed by the algorithm of Section 4 which satisfy (44) (with $\mu_j = V_j$ and $\mu = W$).

We then either have that $\varphi_k = f$ or $V_0, V_1, \dots, V_k, \dots$ is infinite with $\deg_z \varphi_k = \deg_z \varphi_{k_0}$ for $k \geq k_0$. In the first case, we have that $\mathcal{S} = \{V_1, \dots, V_k\}$ is a discrete simple admissible family of valuations which determines W .

Suppose that V_1, \dots, V_k, \dots is infinite. Then $\varphi_k \in \Sigma(V_{k_0})$ for $k > k_0$, and so $d(V_{k_0}) = \deg_z \varphi_{k_0}$.

If $\Lambda(V_{k_0})$ has a maximal element γ , $\varphi' \in \Sigma(V_{k_0})$ is a key polynomial with $W(\varphi') = \gamma$ and corresponding valuation $\mu' = [V_{k_0}; \mu'(\varphi') = W(\varphi')]$, then $\{V_1, \dots, V_{k_0}, \mu'\}$ is the first part of the discrete part of $\mathcal{S}^{(1)}$ constructed by Vaquié’s algorithm. If $W(\varphi') = \infty$, then $\mathcal{S} = \mathcal{S}^{(1)} = \{V_1, \dots, V_{k_0}, \mu' = W\}$ is an admissible family of valuations which determines W .

Suppose that $\Lambda(V_{k_0})$ does not have a largest element. Let $\mathcal{C} = (\mu_\alpha)_{\alpha \in C}$ be the associated family of iterated augmented valuations associated to V_{k_0} of (47). Choose a limit key polynomial $\varphi_1^{(2)}$ for \mathcal{C} . The next step in Vaquié’s algorithm is to construct $\mathcal{S} = \mathcal{S}^{(1)} \cup \mathcal{S}^{(2)}$ where $\mathcal{S}^{(1)} = \{V_1, \dots, V_{k_0}\} \cup \mathcal{C}$ and $\mathcal{S}^{(2)} = \{V_1^{(2)} = [C; V_1^{(2)}(\varphi_1^{(2)}) = W(\varphi_1^{(2)})]\}$.

Looking again at the case where $\Lambda(V_{k_0})$ has a maximal element γ and $\varphi' \in \Sigma(V_{k_0})$ is the corresponding key polynomial, we have an expression $\varphi' = \varphi_{k_0} + h$ where $h \in K[z]$ has $\deg_z h < \deg_z \varphi_{k_0}$. We further have that $h \in R_{V_0}[z]$ by Remark 3.3. We have an expression

(for some r)

$$h = \sum_{j=1}^r a_j \varphi_1^{\sigma_1(j)} \cdots \varphi_{k_0-1}^{\sigma_{k_0-1}(j)}$$

with $a_j \in R_{V_0}$, $0 \leq \sigma_i(j) < n_i = [G_{V_i} : G_{V_{i-1}}]$ for all i and j and

$$W(a_i \varphi_1^{\sigma_1(i)} \cdots \varphi_{k_0-1}^{\sigma_{k_0-1}(i)}) < W(a_j \varphi_1^{\sigma_1(j)} \cdots \varphi_{k_0-1}^{\sigma_{k_0-1}(j)})$$

if $i < j$. Let

$$(49) \quad \psi_i = \varphi_{k_0} + a_1 \varphi_1^{\sigma_1(1)} \cdots \varphi_{k_0-1}^{\sigma_{k_0-1}(1)} + \cdots + a_i \varphi_1^{\sigma_1(i)} \cdots \varphi_{k_0-1}^{\sigma_{k_0-1}(i)}$$

for $1 \leq i \leq r$. We then have (for instance by the criterion of Proposition 1.9 [36]) that

$$(50) \quad V_1, \dots, V_{k_0}, V'_{k_0+1}, \dots, V'_{k_0+r}$$

is a $(k_0 + r)$ -th stage approximant to f over V_0 , where

$$V'_{k_0+1} = [V_{k_0}; V'_{k_0+1}(\psi_1) = W(\psi_1)] \text{ and } V'_{k_0+i} = [V'_{k_0+i-1}; V'_{k_0+i}(\psi_i) = W(\psi_i)] \text{ for } 2 \leq i \leq r.$$

Further, either $W(\varphi') < \infty$ and

$$d(\{V_1, \dots, V_{k_0}, V'_{k_0+1}, \dots, V'_{k_0+r}\}) > \deg_z \varphi_{k_0},$$

or $W(\varphi') = \infty$, in which case $f = \varphi'$ (since f and φ' are unitary in z of the same degree) and $\psi_r = f$.

We may now continue the algorithm of Section 4 to construct higher stage approximants, starting from V'_{k_0+r} . After a finite number of iterations of this procedure, we construct a sequence of approximants to f ,

$$(51) \quad V_1, \dots, V_{k_1}, \dots$$

so that $\deg_z \varphi_i \leq \deg_z \varphi_{i+1}$ if $i < k_1$ and $\deg_z \varphi_i = \deg_z \varphi_i$ for $i \geq k_1$. which is either of finite length k_1 , so that $V_{k_1} = W$, or there is a jump ($t > 1$) in the construction of the admissible family $\mathcal{S} = \mathcal{S}^{(1)} \cup \dots \cup \mathcal{S}^{(t)}$ determining W .

Suppose that (51) is infinite and the equivalent conditions of Lemma 3.2 hold for (51). Let $\mathcal{C} = (\mu_\alpha)_{\alpha \in \mathcal{C}}$ be the associated family to V_{k_1} of (47). Suppose $g \in K[z]$ and $W(g) < \infty$ and k is so large that $W(\varphi_k) > W(g)$. Write $g = g_m \varphi_k^m + \cdots + g_0$ with $\deg_z g_i < \deg_z \varphi_k$ for all i . We have that

$$V_k(g) = V_k(g_0) = V_{k_0-1}(g_0) = W(g_0) = W(g).$$

Thus $g \notin \tilde{\Sigma}(\mathcal{C})$ and so $\deg_z f$ is the smallest degree of an element of $\tilde{\Sigma}(\mathcal{C})$. Thus $\mathcal{S} = \mathcal{S}^{(1)} \cup \mathcal{S}^{(2)}$ where $\mathcal{S}^{(1)} = \{V_1, \dots, V_{k_1}\} \cup \mathcal{C}$ and $\mathcal{S}^{(2)} = \{V_1^{(2)}\}$ where $V_1^{(2)} = [\mathcal{C}; V_1^{(2)}(f(z)) = \infty]$.

The following proposition follows from our analysis.

Proposition 7.1. *Suppose that V_0 has finite rank. Then there exists a realization of the algorithm of Section 4 which produces the first simple admissible family $\mathcal{S}^{(1)}$ of an admissible family $\mathcal{S} = \mathcal{S}^{(1)} \cup \dots \cup \mathcal{S}^{(t)}$ determining W , where all key polynomials are in $R_\nu[z]$.*

7.2. Invariants of ramification and jumps. Suppose that W is an extension of a valuation $V = V_0$ of K to a pseudo valuation of $K[z]$ with $I(W)_\infty = (f(z))$ in $K[z]$ with f unitary. Let ω be the induced valuation on $L = K[z]/(f(z))$.

The jumps $s^{(j-1)}(\mathcal{S})$ in a family $\mathcal{S} = \mathcal{S}^{(1)} \cup \dots \cup \mathcal{S}^{(t)}$ realizing W are defined by the equations

$$(52) \quad \deg_z \varphi_1^{(j)} = s^{(j-1)}(\mathcal{S}) \deg_z \varphi_\alpha^{(j-1)}$$

where $\varphi_\alpha^{(j-1)}$ is a key polynomial of a member of the continuous family $\mathcal{C}^{(j-1)}$ associated to $\mathcal{S}^{(j-1)}$. The total jump of the family \mathcal{S} is

$$s^{\text{tot}}(\mathcal{S}) = \prod_{j=2}^t s^{(j-1)}(\mathcal{S}).$$

We have by Lemma 2.11 and Corollary 2.10 [37] that

$$(53) \quad \deg_z f = [L : K] = e(\omega/V) f(\omega/V) s^{\text{tot}}(\mathcal{S}).$$

We have that $s^{\text{tot}}(\mathcal{S}) = 1$ if and only if there are no jumps in the construction of approximants. Here $e(\omega/V) = [G_\omega : G_V]$ where G_ω and G_V are the respective value groups of ω and V , and $f(\omega/V)$ is the index of the respective residue fields of the valuation rings of ω and V .

In the case where ω is the unique extension of V to a valuation of L , we have by Ostrowski's lemma that

$$(54) \quad [L : K] = e(\omega/V) f(\omega/V) \delta(\omega/V)$$

where the defect $\delta(\omega/V)$ is a power of the residue characteristic p of V . Comparing with (53), we have that $s^{\text{tot}}(\mathcal{S}) = \delta(\omega/V)$ in this case. Thus (assuming ω is the unique extension of V) there is no jump if and only if there is no defect and in this case,

$$(55) \quad [L : K] = e(\omega/V) f(\omega/V).$$

In contrast to the good property of key polynomials of (4), we have examples of the following type for limit key polynomials.

Example 7.2. *The jumps $s^{(i)}$ and total jump s^{tot} can be rational numbers which are not integers.*

We now construct such an example. Let k be an algebraically closed field and $K = k(x)$ be a rational function field in one variable over k . Let ν be the valuation of K with valuation ring $R_\nu = k[x]_{(x)}$ and such that $\nu(x) = 1$. Let $L = K[z]/(z^3 - z^2 - x) \cong k(z)$. Let ω be the extension of ν to L with valuation ring $R_\omega = k[z]_{(z)}$ and $\omega(z) = \frac{1}{2}$. Then $e(\omega/\nu) = 2$ and $f(\omega/\nu) = 1$. Thus by (53),

$$s^{\text{tot}} = \frac{\deg_z(z^3 - z^2 - x)}{e(\omega/\nu) f(\omega/\nu)} = \frac{3}{2}.$$

8. DEFECTLESS EXTENSIONS

Lemma 8.1. *Suppose that (K, ν) is a valued field containing an algebraically closed field k such that $R_\nu/m_\nu \cong k$ and $f(z) \in R_\nu[z]$ is unitary, irreducible and separable. Let $L = K[z]/(f(z))$ and let ω be an extension of ν to L . Let W be the induced pseudo valuation on $K[z]$. Let $\bar{f}(z) \in K^h[z]$ be the irreducible factor of $f(z)$ which induces ω (by Lemma 6.1) and let ω^h be the (unique) extension of ν^h to $K^h[z]/(\bar{f}(z))$. Let \bar{W} be the induced pseudo valuation on $K^h[z]$. Let $V_0 = \nu$ and $W_0 = \nu^h$. Then the following hold:*

- 1) $\text{gr}_{\nu^h}(R_{\nu^h}) = \text{gr}_{\nu}(R_{\nu})$.
- 2) Set $\varphi_1 = z$, let $V_1 = [V_0; V_1(\varphi_1) = W(\varphi_1)]$ and let $W_1 = [\nu^h; W_1(\varphi_1) = W(\varphi_1)]$. Then $\text{gr}_{W_1}(R_{\nu^h}[z]) = \text{gr}_{V_1}(R_{\nu}[z])$.
- 3) Suppose that $V_i = [V_{i-1}; V_i(\varphi_i) = W(\varphi_i)]$ for $1 \leq i \leq k$ is a realization of the algorithm of Section 4 in $R_{\nu}[z]$ such that $W_i = [W_{i-1}; W_i(\varphi_i) = W(\varphi_i)]$ for $1 \leq i \leq k$ is a realization of the the algorithm of Section 4 in $R_{\nu^h}[z]$ and

$$(56) \quad \text{gr}_{W_i}(R_{\nu^h}[z]) = \text{gr}_{V_i}(R_{\nu}[z]) \text{ for } 1 \leq i \leq k.$$

Suppose that $\varphi_{k+1} \in R_{\nu}[z]$ has an expression $\varphi_{k+1} = \varphi_k^n - c_k \varphi_1^{j_1} \cdots \varphi_{k-1}^{j_{k-1}}$ of the form of (15), φ_{k+1} is a key polynomial for W_k and $W_{k+1} = [W_k; W_{k+1}(\varphi_{k+1}) = W(\varphi_{k+1})]$ is a $(k+1)$ -st approximant of \bar{f} over W_0 . Then φ_{k+1} is a key polynomial for V_k and $V_{k+1} = [V_k; V_{k+1}(\varphi_{k+1}) = W(\varphi_{k+1})]$ is a $(k+1)$ -st approximant of f over V_0 . Further, $\text{gr}_{W_{k+1}}(R_{\nu^h}[z]) = \text{gr}_{V_{k+1}}(R_{\nu}[z])$.

Proof. Statement 1) follows since $G_{\nu^h} = G_{\nu}$ and $R_{\nu^h}/m_{\nu^h} = R_{\nu}/m_{\nu}$ by [14, Theorem 17.19]. Statement 2) follows since

$$\text{gr}_{W_1}(R_{\nu^h}[z]) = \text{gr}_{\nu^h}(R_{\nu^h})[\text{in}_{W_1}(\varphi_1)] = \text{gr}_{\nu}(R_{\nu})[\text{in}_{V_1}(\varphi_1)] = \text{gr}_{V_1}(R_{\nu}[z]).$$

Now we will prove statement 3). To show that φ_{k+1} is a key polynomial over V_k , we must verify that 1) - 6) of the definition of a key polynomial, given after (3) hold for φ_{k+1} over V_k . This follows since these conditions hold for φ_{k+1} over W_k . The fact that W_{k+1} is a $(k+1)$ -st approximant to \bar{f} over W_k implies that φ_{k+1} equivalence divides \bar{f} in W_k . Thus $\text{in}_{W_k}(\varphi_{k+1})$ divides $\text{in}_{W_k}(\bar{f})$ in $\text{gr}_{W_k}(R_{\nu^h}[z])$. Now $\text{in}_{W_k}(\bar{f})$ divides $\text{in}_{W_k}(f(z))$ in $\text{gr}_{W_k}(R_{\nu^h}[z])$. So $\text{in}_{W_k}(\varphi_{k+1})$ divides $\text{in}_{W_k}(f(z))$ in $\text{gr}_{W_k}(R_{\nu^h}[z]) = \text{gr}_{V_k}(R_{\nu}[z])$. Thus φ_{k+1} equivalence divides $f(z)$ in V_k and so V_{k+1} is a $(k+1)$ -st approximant to $f(z)$ over V_k . We have that $n = [G_{W_k} : G_{W_{k-1}}] = [G_{V_k} : G_{V_{k-1}}]$ as $G_{W_{k-1}} = G_{V_{k-1}}$. Finally, we have that $\text{gr}_{V_{k+1}}(R_{\nu}[z]) = \text{gr}_{W_{k+1}}(R_{\nu^h}[z])$ by Lemma 4.3. \square

Theorem 8.2. *Suppose that A is an excellent local domain which contains an algebraically closed field k such that $A/m_A \cong k$. Let K be the quotient field of A and suppose that $V_0 = \nu$ is a rank 1 valuation of K which dominates A and such that the residue field of the valuation ring of V_0 is k . Suppose that $f(z) \in A[z]$ is unitary, irreducible and separable and W is a pseudo valuation of $K[z]$ such that $I(W)_{\infty} = (f(z))$ in $K[z]$ which extends V_0 . Let ω be the induced valuation of $L = K[z]/(f(z))$. Then ω is defectless over ν ($\delta(\omega/\nu) = 1$) if and only if there exists a normal birational extension A_1 of A which is dominated by ν such that there exists a realization*

$$V_1, \dots, V_k, \dots$$

of the algorithm of Section 4 in $A_1[z]$, satisfying (14) and (15) for all k with $c_k \in A_1$ for all $k \geq 1$, such that $W = V_k$ for some finite k or $W = \lim_{k \rightarrow \infty} V_k$.

If these equivalent conditions hold, then there exists a positive integer k such that

$$\text{gr}_{\omega}(A_1[z]/(f(z))) \cong \text{gr}_{\nu}(A_1)[\bar{\varphi}_1, \dots, \bar{\varphi}_k]/I$$

where

$$I = (\bar{\varphi}_1^{n_1} - \bar{c}_1, \bar{\varphi}_2^{n_2} - \bar{c}_2 \bar{\varphi}_1^{j_1(2)}, \dots, \bar{\varphi}_k^{n_k} - \bar{c}_k \bar{\varphi}_1^{j_1(k)} \bar{\varphi}_2^{j_2(k)} \cdots \bar{\varphi}_{k-1}^{j_{k-1}(k)})$$

is a finitely generated and presented $\text{gr}_{\nu}(A_1)$ -module.

An example showing that the conclusions of Theorem 8.2 may not hold if ν has rank larger than one will be given in Section 10. In Example 8.3, it will be shown that the conclusions of Theorem 8.2 may not hold if $f(z)$ is not separable over K .

Proof. First suppose that $\delta(\omega/\nu) = 1$. Let notation be as in Section 6. By Lemma 6.1, there exists an extension \overline{W} of ν^h to a pseudo valuation of $K^h[z]$, such that $I(\overline{W})_\infty = (\overline{f})$ where $\overline{f}(z)$ is an irreducible factor of $f(z)$ in $K^h[z]$, and \overline{W} is an extension of W .

We will construct a special sequence of approximants W_1, \dots, W_{k_0} to \overline{f} over ν^h such that $\overline{W} = W_{k_0}$. In particular,

$$W_{k_0} = [W_{k_0-1}; W_{k_0}(\varphi_{k_0}) = \infty]$$

where $\varphi_{k_0} = \overline{f}$.

Set $\varphi_1 = z$ and let $W_1 = [\nu^h; W_1(\varphi_1) = W(\varphi_1)]$. Suppose by induction on k that we have constructed a sequence of approximants to \overline{f} over ν^h ,

$$W_1, \dots, W_k$$

giving a realization of the algorithm of Section 4, such that expressions

$$\varphi_i = \varphi_{i-1}^{n_{i-1}} - c_{i-1} \varphi_1^{j_1(i-1)} \dots \varphi_{i-2}^{j_{i-2}(i-1)}$$

of the form of 15) hold for $i \leq k$ with $c_i \in R_\nu$ for $i \leq k-1$. After replacing A with a birational extension A_1 of A , we may suppose that $c_i \in A$ for $i \leq k-1$.

If $\Lambda(W_k)$ does not have a largest element, then we have a jump $s^{(1)} > 1$ by (52) and the analysis of this case in Subsection 7.1. But by (53) and (54), there cannot be a jump, and we have a contradiction, showing that $\Lambda(W_k)$ has a largest element.

Suppose we are in the case where $\Lambda(W_k)$ has a maximal element $\gamma \neq \infty$ and $\varphi' \in \Sigma(W_k)$ is a corresponding key polynomial. We will modify the resulting sequence (50) of the analysis in Subsection 7.1, which we will write as

$$(57) \quad W_1, \dots, W_k, W'_{k+1}, \dots, W'_{k+r}$$

by modifying the ψ_i of (49), replacing the a_i with suitable $b_i \in R_\nu$ for $1 \leq i \leq r$. With the notation of Lemma 6.2, since $a_1, \dots, a_r \in R_{\nu^h}$, there exists A_l such that $a_i \in A_l^h$ for $1 \leq i \leq r$ and $\varphi_1, \dots, \varphi_{k_0} \in A_l[z]$. Thus, since \overline{W} induces a rank 1 valuation on $K^h[z]/(\overline{f}(z))$, there exists $n \in \mathbf{Z}_+$ such that $nV_0(m_{A_l^h}) > \overline{W}(\varphi') = \gamma$. Now $A_l \rightarrow A_l^h$ is unramified with no residue field extension, so there exists $b_i \in A_l$ such that $a_i - b_i \in m_{A_l^h}^n$ for $1 \leq i \leq r$. Thus $V_0(b_i) - \nu^h(a_i) > \overline{w}(\varphi')$ for $1 \leq i \leq r$ and we can replace ψ_i with $\psi_{i-1} + b_i \varphi_1^{j_1(i)} \dots \varphi_{k-1}^{j_{k-1}(i)}$ in (49) for $1 \leq i \leq r$, to produce a sequence (57) with $\psi_i \in R_\nu[z]$ for all i . We then have a corresponding sequence to (57),

$$V_1, \dots, V_k, V'_{k+1}, \dots, V'_{k+r}$$

of approximants to f over V_0 by Lemma 8.1.

Now we can continue, using the algorithm of Section 4, applying the above argument as necessary until we reach W_k such that the maximal element of $\Lambda(W_k)$ is ∞ , so that $\overline{f} \in \Sigma(W_k)$.

With this assumption, there exists l (with the notation of Lemma 6.2) such that the coefficients of \overline{f} are in A_l^h and the coefficients of $\varphi_1, \dots, \varphi_k$ are in A_l . We have $\overline{f} = \varphi_k + h$ where $h \in A_l^h[z]$ and $\deg_z h < \deg_z \varphi_k$. Set $\psi_0 = \varphi_k$. By induction, we may construct a sequence $\psi_i \in A_l[z]$ of monic polynomials with $\deg_z \psi_i = \deg_z \varphi_k$, such that for all i , $\overline{f} = \psi_i + h_i$ with $h_i \in (A_l)^h[z]$ a polynomial of degree $< \deg_z \varphi_k$ and

$$\psi_{i+1} = \psi_i + b_i \varphi_1^{\sigma_1(i)} \dots \varphi_{k_0-1}^{\sigma_{k_0-1}(i)}$$

with $b_i \in A_l$ and $0 \leq \sigma_j(i) < n_j$ for $1 \leq j \leq k_0-1$ such that $\overline{W}(\psi_{i+1}) > \overline{W}(\psi_i)$ for all i . Since A_l is Noetherian, and \overline{W} induces a rank 1 valuation on $K^h[z]/(\overline{f}(z))$, we have that

\overline{W} takes on $A_l[z]$ only a finite number of values which are less than or equal to a given finite upper bound. Thus we either obtain that $\psi_i = \overline{f}(z)$ for some i , or that

$$\lim_{i \rightarrow \infty} W(\psi_i) = \lim_{i \rightarrow \infty} \overline{W}(\psi_i) = \infty.$$

By Lemma 8.1, inductively defining $V_i = [V_{i-1}; V_i(\varphi_i) = W(\varphi_i)]$ for $1 \leq i \leq k$ and $V_{i+k} = [V_{i+k-1}; V_i(\psi_i) = W(\psi_i)]$ for $k < i$, we construct a sequence

$$V_1, \dots, V_k, \dots$$

of approximants to $f(z)$ over V_0 such that $\lim_{i \rightarrow \infty} V_i(\varphi_i) = \infty$, so that $W = \lim_{i \rightarrow \infty} V_i$ by Lemma 3.2.

Now suppose there exists a normal birational extension A_1 of A and a realization V_1, \dots, V_k, \dots of the algorithm of Section 4 as in the statement of the theorem. We will show that the defect $\delta(\omega/\nu) = 1$.

First suppose that the sequence is of finite length, terminating with $V_k = W$, so that the last key polynomial is $\varphi_k = f$ (with $V_k(\varphi_k) = \infty$). We have that $\deg_z \varphi_1 = 1$ and $\deg_z \varphi_i = n_{i-1} \deg_z \varphi_{i-1}$ for $i \geq 2$. Thus

$$[G_\omega : G_{V_0}] \delta(\omega/V_0) \leq \deg_z f = n_1 n_2 \cdots n_{k-1} = [G_\omega : G_{V_0}]$$

so that $\delta(\omega/\nu) = 1$.

Now suppose that V_1, \dots, V_k, \dots is of infinite length. We have (by Lemma 6.4) natural extensions of valued fields

$$(K, \nu) \rightarrow (K^h, \nu^h) \rightarrow (\hat{K}, \hat{\nu}).$$

Let $\overline{f}(z)$ be the irreducible factor of $f(z)$ in $K^h[z]$ which induces ω (from Lemma 6.1). Then $\overline{f}(z)$ is irreducible in $\hat{K}[z]$ (by Lemma 6.4) and so is the irreducible factor of $f(z)$ in $\hat{K}[z]$ which induces ω (by Lemma 6.3). Thus the pseudo valuation W extends to a pseudo valuation W^h of $K^h[z]$ and to a pseudo valuation \hat{W} of $\hat{K}[z]$ such that $I(W^h)_\infty = (\overline{f}(z))$ in $K^h[z]$ and $I(\hat{W})_\infty = (\overline{f}(z))$ in $\hat{K}[z]$. By (43),

$$(58) \quad \delta(\omega/\nu) = [L^h : K^h]/[G_\omega : G_\nu] = \deg_z \overline{f}/[G_\omega : G_\nu].$$

There exists k_0 such that $\deg_z \varphi_k = \deg_z \varphi_{k_0}$ for $k \geq k_0$. There exist $a_i \in A_1$ and $j_1(i), \dots, j_{k_0-1}(i)$ with $0 \leq j_l(i) < n_l$ for $1 \leq l \leq k_0 - 1$ such that

$$\varphi_{k_0+i+1} = \varphi_{k_0+i} - a_i \varphi_1^{j_1(i)} \cdots \varphi_{k_0-1}^{j_{k_0-1}(i)}$$

for $i \geq 0$. Now

$$W(\varphi_{k_0+i}) = W(a_i \varphi_1^{j_1(i)} \cdots \varphi_{k_0-1}^{j_{k_0-1}(i)})$$

for $i > 0$ and

$$(59) \quad W(\varphi_{k_0+i}) \mapsto \infty \text{ as } i \mapsto \infty$$

by Lemma 3.2. Thus $\nu(a_i) \mapsto \infty$ as $i \mapsto \infty$.

For fixed (b_1, \dots, b_{k_0-1}) such that $0 \leq b_l < n_l$ for $1 \leq l \leq k_0 - 1$, define

$$c_l(b_1, \dots, b_{k_0-1}) = \sum a_i,$$

where the sum is over $i < l$ such that $(j_1(i), \dots, j_{k_0-1}(i)) = (b_1, \dots, b_{k_0-1})$. Let

$$\tau_i = \sum_{b_1, \dots, b_{k_0-1}} c_i(b_1, \dots, b_{k_0-1}) \varphi_1^{b_1} \cdots \varphi_{k_0-1}^{b_{k_0-1}}$$

where the sum is over b_1, \dots, b_{k_0-1} such that $0 \leq b_j < n_j$ for $1 \leq j \leq k_0 - 1$. We have that $\varphi_{k_0+i} = \varphi_{k_0} - \tau_i$. Thus

$$W(\tau_j - \tau_i) = W(\varphi_{k_0+i} - \varphi_{k_0+j}) \geq \min\{W(\varphi_{k_0+i}), W(\varphi_{k_0+j})\}$$

so $W(\tau_j - \tau_i) \mapsto \infty$ as $j \geq i \mapsto \infty$. We have that

$$W(\tau_j - \tau_i) = \min\{\nu(c_i(b_1, \dots, b_{k_0-1}) - c_j(b_1, \dots, b_{k_0-1})) + W(\varphi_1^{b_1} \cdots \varphi_{k_0-1}^{b_{k_0-1}})\}$$

where the minimum is over b_1, \dots, b_{k_0-1} with $0 \leq b_j < n_j$ for $1 \leq j \leq k_0 - 1$. So for all b_1, \dots, b_{k_0-1} , $\nu(c_i(b_1, \dots, b_{k_0-1}) - c_j(b_1, \dots, b_{k_0-1})) \mapsto \infty$ as $j \geq i \mapsto \infty$. Thus for each b_1, \dots, b_{k_0-1} , $(c_i(b_1, \dots, b_{k_0-1}))$ is a ν -Cauchy sequence.

Thus these sequences have limits in \hat{K} , and so (φ_{k_0+i}) is a ν -Cauchy sequence in $K[z]$ which has a non zero limit φ_∞ in $\hat{K}[z]$ (φ_∞ is necessarily unitary of degree equal to $\deg_z \varphi_{k_0}$). Now $\varphi_\infty \in I(\hat{W})_\infty = (\bar{f})$ by (59). Thus $\deg_z \varphi_\infty \geq \deg_z \bar{f}$. Now

$$\deg_z \bar{f} \leq \deg_z \varphi_\infty = \deg_z \varphi_{k_0} = [G_{V_{k_0}} : G_{V_0}] = [G_\omega : G_\nu].$$

Thus $\deg_z \bar{f} = [G_\omega : G_\nu]$ and $\delta(\omega/\nu) = 1$ by (58). \square

Example 8.3. *The conclusions of Theorem 8.2 may fail if $f(z)$ is not separable over K .*

An example of F.K. Schmidt of a discrete valuation ring (with value group \mathbf{Z}) and an inseparable extension of its quotient field which has defect is explained in [22, Example 3.1]. The example is as follows. Let k be an algebraically closed field of characteristic $p > 0$, $A = k[x, y]_{(x, y)}$ be the localization of a polynomial ring in two variables and K be the quotient field of A . Let $k[[t]]$ be a power series ring and let $s \in k[[t]]$ be transcendental over $k(t)$ and such that $\text{ord}_t(s) > 0$. The k -algebra embedding $K \rightarrow k[[t]]$ defined by $x \mapsto t$ and $y \mapsto s^p$ induces a valuation ν on K which dominates A by $\nu(g(x, y)) = \text{ord}_t(g(t, s^p))$. We have that $G_\nu = \mathbf{Z}$ and $R_\nu/m_\nu = k$. Let $f(z) = z^p - y \in K[z]$. There is a unique extension of ν to a valuation ω of $L = K[z]/(f(z))$ (since L is purely inseparable over K) which is an immediate extension of ν ($G_\omega = G_\nu$ and $R_\omega/m_\omega = R_\nu/m_\nu$). Thus the defect $\delta(\omega/\nu) = \deg_z f = p$ by Ostrowski's lemma (1). Since ν is a rank 1 discrete valuation, by MacLane's theorem (Section 3), ω is a limit valuation which is realized by his algorithm. We will give an explicit construction.

Let W be the pseudo valuation induced by ω on $K[z]$, and let $V_0 = \nu$. We will construct a sequence of approximants V_1, \dots, V_i, \dots to f over V_0 which realize W .

Expand $s = \sum_{i=1}^{\infty} a_i t^i$ with $a_i \in k$. We have that $s^p = \sum_{i=1}^{\infty} a_i^p t^{ip}$. Define

$$\sigma(1) = \text{ord}_t(s) = \min\{i \mid a_i \neq 0\}$$

and for $j > 1$,

$$\sigma(j) = \min\{i \mid \sigma(j-1) < i \text{ and } a_i \neq 0\}.$$

The first approximant is $V_1 = [V_0; V_1(\varphi_1) = \sigma(1)]$ where $\varphi_1 = z$. For $i \geq 1$, V_{i+1} is defined by $V_{i+1} = [V_i; V_{i+1}(\varphi_{i+1}) = \sigma(i+1)]$, where $\varphi_{i+1} = \varphi_i - a_{\sigma(i+1)} x^{\sigma(i+1)}$. Then $\lim_{i \rightarrow \infty} V_i(\varphi_i) = \infty$ and so W is the limit valuation $W = \lim_{i \rightarrow \infty} V_i$ by Lemma 3.2.

9. A RANK 1 SEPARABLE EXAMPLE WITH DEFECT

We consider an example from Theorem 7.38 [11], with regard to the algorithm of Section 4. Let k be an algebraically closed field of characteristic $p > 0$. Let $K = k(u, v)$ be a two

dimensional rational function field over k , and, using the method of [32] and [9], define a valuation ν of K by the following generating sequence:

$$P_0 = u, P_1 = v, P_2 = v^{p^2} - u \text{ and } P_{i+1} = P_i^{p^2} - u^{p^{2i-2}} P_{i-1} \text{ for } i \geq 2.$$

We normalize the valuation ν so that $\nu(u) = 1$. We have the defining relations $v^{p^2} \sim u$ in ν and $P_i^{p^2} \sim u^{p^{2i-2}} P_{i-1}$ in ν for $i \geq 2$. As shown in [11], the value group $G_\nu = \frac{1}{p^\infty} \mathbf{Z} = \cup_{i \geq 1} \frac{1}{p^i} \mathbf{Z}$. Let $f = x^p + ux^{p-1} - u \in K[x]$. By Theorem 7.38 [11], ν has a unique extension to a valuation ω of $L = K[x]/(f(z))$. Further, ω is an immediate extension, so it is a defect extension with $[L : K] = \delta(\omega/\nu) = [L : K] = p$.

Let W be the pseudo valuation induced by ω on $K[x]$. We will construct a realization of the algorithm of Section 4, giving an infinite sequence of approximants to f over $V_0 = \nu$,

$$V_1, \dots, V_k, \dots$$

satisfying (14) and (15) with $c_i \in R_\nu$ for all i .

Setting $\varphi_1 = x$, we have that $N(V_0, \varphi_1)$ has a single segment, which has the slope $\frac{V_0(u)}{p} = \frac{1}{p}$. Thus the first approximant to f over V_0 is $V_1 = [V_0; V_1(\varphi_1) = \frac{1}{p}]$.

We will make use of the following observation when constructing our sequence of approximants. Suppose we have constructed the sequence V_1, \dots, V_k of approximants, where $\deg_x \varphi_i = 1$ for all i . Then for $i \leq k$, $\varphi_i = \varphi_{i-1} + a_{i-1}$ with $a_{i-1} \in R_\nu$ and

$$W(\varphi_i) > W(\varphi_{i-1}) = \nu(a_{i-1}).$$

Expanding

$$ux^{p-1} = g_{p-1} \varphi_k^{p-1} + g_{p-2} \varphi_k^{p-2} + \dots + g_0$$

with $g_i \in R_\nu$, we have that

$$(60) \quad V_0(g_0) \geq \min\{W(g_i \varphi_k^i)\} = V_k(ux^{p-1}) = W(ux^{p-1}) = 1 + \frac{p-1}{p} > \frac{p^4}{p^4-1}.$$

Now $f \sim \varphi_1^p - u$ in V_1 and $u = v^{p^2} - P_2 \sim v^{p^2}$ in V_0 . Thus $f \sim (\varphi_1 - v^p)^p$ in V_1 , and we take our second key polynomial to be $\varphi_2 = \varphi_1 - v^p = \varphi_1 - P_1^p$. We thus have that the second approximant is $V_2 = [V_1; V_2(\varphi_2) = W(\varphi_2)]$. Expanding

$$f = \varphi_2^p + f_{p-1} \varphi_2^{p-1} + \dots + f_1 \varphi_2 + f_0$$

with $f_i \in R_\nu$, by (60) with $k = 2$, we have that

$$f_0 = v^{p^2} - u + \text{terms of value} > \frac{p^4}{p^4-1}$$

and $v^{p^2} - u = P_2$. Now since ω is the unique extension of ν , we have that the principal part of $N(V_1; \varphi_2)$ is $N(V_1, \varphi_2)$ and $N(V_1, \varphi_2)$ has a single segment, which has slope $\frac{V_0(f_0)}{p} = \frac{V_0(P_2)}{p} = \frac{1}{p} + \frac{1}{p^5}$, which is less than $\frac{p^4}{p^4-1}$. The second approximant is $V_2 = [V_1; V_2(\varphi_2) = \frac{1}{p} + \frac{1}{p^5}]$.

By (60), we have that $f \sim \varphi_2^p + P_2$ in V_2 . Now $P_2 \sim \frac{P_3^{p^2}}{u^{p^4}}$ in V_0 , so

$$f \sim \left(\varphi_2 + \frac{P_3^p}{u^{p^3}} \right)^p \text{ in } V_2,$$

and so $\varphi_3 = \varphi_2 + \frac{P_3^p}{u^{p^3}}$ is a key polynomial for V_2 . We thus have that the third approximant is $V_3 = [V_2; V_3(\varphi_3) = W(\varphi_3)]$. We expand

$$f = \varphi_3^p + f_{p-1}\varphi_3^{p-1} + \cdots + f_1\varphi_3 + f_0$$

with $f_i \in R_\nu$. By (60), we have that

$$f_0 = -\frac{P_3^{p^2}}{u^{p^4}} + P_2 + \text{terms of value} > \frac{p^4}{p^4 - 1}.$$

Also,

$$-\frac{P_3^{p^2}}{u^{p^4}} + P_2 = -\frac{P_4}{u^{p^4}}.$$

Now since ω is the unique extension of ν , we have that the principal part of $N(V_2, \varphi_3)$ is $N(V_2, \varphi_3)$ and $N(V_2, \varphi_3)$ has a single segment, which has slope

$$\frac{V_0(f_0)}{p} = \frac{1}{p}V_0\left(-\frac{P_4}{u^{p^4}}\right) = \frac{1}{p} + \frac{1}{p^5} + \frac{1}{p^9} < \frac{p^4}{p^4 - 1}.$$

The third approximant is $V_3 = [V_2; V_3(\varphi_3) = \frac{1}{p} + \frac{1}{p^5} + \frac{1}{p^9}]$.

Continuing in this way, we construct an infinite sequence of approximants V_1, \dots, V_k, \dots to f over V_0 with $V_k = [V_{k-1}; V_k(\varphi_k) = \mu_k]$ and for $k \geq 3$,

$$\varphi_k = \varphi_{k-1} + (-1)^{k-1} \frac{P_{2(k-2)+1}^p}{u^{p^3+p^7+\dots+p^{4(k-3)+3}}}$$

with

$$\mu_k = \frac{1}{p} + \frac{1}{p^5} + \frac{1}{p^9} + \cdots + \frac{1}{p^{4(k-1)+1}}.$$

We have that

$$(61) \quad \lim_{k \rightarrow \infty} \mu_k = \frac{p^4}{p(p^4 - 1)}.$$

In particular, we have by Lemma 3.2, that the limit valuation $V_\infty = \lim_{k \rightarrow \infty} V_k$ is a valuation, and thus is not equal to W .

We observe that there does not exist a birational extension A_1 of $k[u, v]_{(u, v)}$ which is dominated by V_0 such that $\varphi_i \in A_1[x]$ for all i , as there can only be finitely many values of elements in a Noetherian local ring which is dominated by a rank 1 valuation that are less than a fixed finite bound.

We now analyze the extension W of ν in the context of Vaquié's algorithm. We will construct an admissible family of valuations \mathcal{S} which determines W .

In the above realization of the algorithm of Section 4, we started by defining $\varphi_1 = x$, and $V_1 = [V_0; V_1(\varphi_1) = \frac{1}{p}]$. With the notation of Section 7, we have

$$\Sigma(V_1) = \{x - g \mid g \in K \text{ and } W(x - g) > W(x)\}$$

and

$$\Lambda(V_1) = \{W(\varphi) \mid \varphi \in \Sigma(V_1)\}.$$

Let $(\mu_\alpha)_{\alpha \in C}$ be the associated family of iterated augmented valuations to $\mathcal{A} = \{V_1\}$ of (47).

The concept of distance of an element of L from K and the concepts of dependent and independent Artin-Schreier extensions are introduced in [21]. In [13], our extension ω of ν is analyzed, and it is shown that it is a dependent Artin-Schreier extension. We will make use of a calculation in their proof, to determine $\limsup\{\Lambda(V_1)\}$. Suppose that $g \in \Sigma(V_1)$.

Then $W(x - g) > W(x)$ so $\nu(g) = W(x) = \frac{1}{p}$. Thus $g \in R_\nu$, and by 2) of Theorem 4.4 [13], we have that

$$W(g^p - x^p) \leq 1 + \frac{1}{p^4} + \cdots + \frac{1}{p^{4(k+1)}}$$

for some $k \geq 0$. Thus

$$W(x - g) = \frac{1}{p}W(g^p - x^p) \leq \frac{1}{p} + \frac{1}{p^5} + \cdots + \frac{1}{p^{4(k+1)+1}} < \frac{p^4}{p(p^4 - 1)}.$$

By (61), we have that $\limsup\{\Lambda(V_1)\} = \frac{p^4}{p(p^4-1)}$ and $\frac{p^4}{p(p^4-1)} \notin \Lambda(V_1)$. In particular, $\Lambda(V_1)$ does not have a largest element. Thus the first simple admissible family associated to W is

$$\mathcal{S}^{(1)} = \{V_1\} \cup \{(\mu_\alpha)_{\alpha \in C}\}$$

and \mathcal{S} is the union of $t > 1$ simple admissible families. Since ω is an immediate extension of ν , we have by (53) that

$$p = \deg_x f = s^{\text{tot}}(\mathcal{S}) = \prod_{j=2}^t s^{(j-1)}(\mathcal{S}).$$

Since the $s^{(j-1)}(\mathcal{S})$ are positive rational numbers with $s^{(j-1)}(\mathcal{S}) > 1$ for $2 \leq j \leq t$, we have that $t \geq 2$.

Thus an admissible family of valuations \mathcal{S} which determine W starts with $\mathcal{S}^{(1)}$ and has at least one jump.

10. A DEFECTLESS EXTENSION OF A RANK TWO VALUATION WITH MANY JUMPS

In this section we construct the following example, which shows that the conclusions of Theorem 8.2 may not hold if ν has rank larger than one.

Example 10.1. *Let k be an algebraically closed field of characteristic not equal to 2, and let $k[x, y]$ be a polynomial ring in two variables over k . Let $K = k(x, y)$ and let ν be the rank two valuation on K defined by $\nu(x) = (0, 1), \nu(y) = (1, 0) \in (\mathbf{Z}_2)_{\text{lex}}$ and $\nu|(k \setminus 0) = 0$. Let*

$$(62) \quad f = ((z^2 - x^2 - x^3)^2 - y^2(x^2 + 2x^3))^2 - (y^6 + y^7)$$

and let ω be an extension of ν to $K[z]/(f(z))$. Let W be the induced pseudo valuation of $K[z]$. Then a family $\mathcal{S} = \mathcal{S}^{(1)} \cup \cdots \cup \mathcal{S}^{(t)}$ (with notation of Section 7) realizing W has at least three jumps; that is, $t \geq 3$.

We first establish that f is irreducible in $K[z]$. Setting $x = 0$ in f , we obtain the reduction $\tilde{f} = z^8 - (y^6 + y^7) \in k(y)[z]$. We have that

$$\tilde{f} = \prod_{j=0}^7 (z - \tau^j y^{\frac{3}{4}}(1+y)^{\frac{1}{8}})$$

over an algebraic closure of $k(y)$, where τ is a primitive 8-th root of unity in k . A unitary factor of \tilde{f} of degree r must have the constant term $\tau^s (y^{\frac{3}{4}}(1+y)^{\frac{1}{8}})^r$ for some $s \in \mathbf{N}$. But $(y^{\frac{3}{4}}(1+y)^{\frac{1}{8}})^r \in k(y)$ only if $r = 8$, so \tilde{f} is irreducible in $k(y)[z]$, and thus f is irreducible in $K[z]$.

Henselization is discussed in Section 6.

Lemma 10.2. *The polynomial f factors into a product of linear unitary polynomials in $K^h[z]$, where (K^h, ν^h) is a Henselization of (K, ν) .*

Proof. We will solve the equation $f(z) = 0$ in R_{ν^h} . Let

$$Q = z^2 - (x^2 + x^3), U = Q^2 - y^2(x^2 + 2x^3).$$

With these substitutions, the equation $f(z) = 0$ becomes $U^2 = (y^6 + y^7)$. Let $(1 + y)^{\frac{1}{2}}$ be a square root of $1 + y$ in the Henselization A^h of $A = k[x, y]_{(x, y)}$. Then $U = y^3(1 + y)^{\frac{1}{2}}$ in A^h . Thus we have that

$$Q^2 = y^3(1 + y)^{\frac{1}{2}} + y^2(x^2 + 2x^3) = y^2(x^2 + 2x^3 + y(1 + y)^{\frac{1}{2}}).$$

Set $x = x_1$ and $y = x_1^2 y_1$. We have that $x_1, y_1 \in R_{\nu}$. Then

$$Q^2 = x_1^4 y_1^2 (x_1^2 + 2x_1^3 + x_1^2 y_1 (1 + x_1^2 y_1)^{\frac{1}{2}}) = x_1^6 y_1^2 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}}).$$

Let $(1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}}$ be a square root of $1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}}$ in the Henselization A_1^h of $A_1 = k[x_1, y_1]_{(x_1, y_1)}$. Then

$$Q = x_1^3 y_1 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}}$$

in A_1^h . We now have that

$$\begin{aligned} z^2 &= Q + x^2 + x^3 \\ &= x_1^3 y_1 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}} + x_1^2 + x_1^3 \\ &= x_1^2 (1 + x_1 + x_1 y_1 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}}). \end{aligned}$$

Let $(1 + x_1 + x_1 y_1 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}})^{\frac{1}{2}}$ be a square root of

$$1 + x_1 + x_1 y_1 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}}$$

in A_1^h . Then

$$z = x_1 (1 + x_1 + x_1 y_1 (1 + 2x_1 + y_1 (1 + x_1^2 y_1)^{\frac{1}{2}})^{\frac{1}{2}})^{\frac{1}{2}} \in A_1^h \subset R_{\nu^h}$$

by Lemma 6.2.

Since all eight roots of $f(z)$ can be found this way, by making different choices of square roots, we have the desired factorization of $f(z)$ in $K^h[z]$ into a product of linear polynomials. \square

By Lemma 6.1, ω is the restriction to $K[z]/(f(z))$ of the extension of ν^h to a valuation ω^h of $K^h[z]/(\bar{f})$ for some factor \bar{f} of f in $K^h[z]$. Since \bar{f} is a linear polynomial by Lemma 10.2, we have that

$$(63) \quad [G_{\omega} : G_{\nu}] = [G_{\omega^h} : G_{\nu^h}] = \deg_z \bar{f} = 1$$

by (53).

We will require the following remark.

Remark 10.3. *An element $g \in k(z)$ is a square of an element of $k(z)$ if and only if all zeros and poles of $g(z)$ in \mathbf{A}_k^1 have even order.*

The remark follows since every element $g(z)$ of $k(z)$ has a unique factorization

$$g(z) = c(z - a_1)^{n_1} \cdots (z - a_t)^{n_t}$$

with $c \in k$, a_1, \dots, a_t distinct elements of k and n_1, \dots, n_t nonzero integers.

We now turn to the construction of the family \mathcal{S} . We will use the notation of Section 7. To begin with, we observe that the total jump $s^{\text{tot}}(\mathcal{S})$ of \mathcal{S} satisfies

$$(64) \quad s^{\text{tot}}(\mathcal{S}) = \deg_z f(z) = 8$$

by (53) and (63).

Let $V_0 = \nu$. Since $W(f(z)) = \infty$, we have that $W(z) = (0, 1)$ and so the first approximant is $V_1 = [V_0; V_1(z) = (0, 1)]$. As above, let $Q = z^2 - (x^2 + x^3)$. Since $W(f(z)) = \infty$, we have that $W(Q) = (1, 1)$. Let $\sum_{i=1}^{\infty} \alpha_i x^i$ with $\alpha_i \in k$ be a square root of $x^2 + x^3 = x^2(1+x)$ in $k[[x]]$. Let $\bar{z} = z - (\alpha_1 x + \cdots + \alpha_n x^n)$ for some $n \in \mathbf{Z}_+$. Then

$$\begin{aligned} Q &= (\bar{z} + \alpha_1 x + \cdots + \alpha_n x^n)^2 - (x^2 + x^3) \\ &= \bar{z}^2 + 2(\alpha_1 x + \cdots + \alpha_n x^n)\bar{z} + (\alpha_1 x + \cdots + \alpha_n x^n)^2 - (x^2 + x^3) \end{aligned}$$

so that $W(\bar{z}(\bar{z} + 2(\alpha_1 x + \cdots + \alpha_n x^n))) > (0, n)$. Thus

$$(65) \quad W(z - \alpha_1 x - \cdots - \alpha_n x^n) > (0, \frac{n}{2}) \text{ or } W(z + \alpha_1 x + \cdots + \alpha_n x^n) > (0, \frac{n}{2}).$$

Thus $d(V_1) = 1$ and so

$$\Sigma(V_1) = \{\varphi = z + h \mid h \in K \text{ and } V_1(\varphi) < W(\varphi)\}.$$

We will show that

$$(66) \quad \Lambda(V_1) = \{W(\varphi) \mid \varphi \in \Sigma(V_1)\} \subset \{0\} \times \mathbf{Z}_+.$$

We now prove equation (66). Suppose there exists $h \in K$ such that setting $\varphi = z + h$, we have that $W(\varphi) \geq (1, 0)$. Then

$$(67) \quad W(h) = (0, 1).$$

Substituting into Q , we have that $Q = \varphi^2 - 2h\varphi + h^2 - (x^2 + x^3)$. Now $W(Q) = (1, 1)$ implies

$$(68) \quad W(h^2 - (x^2 + x^3)) \geq (1, 0).$$

By (67), we have an expression

$$h = \frac{\alpha_0(x) + y\Omega_1}{\beta_0(x) + y\Omega_2}$$

with $\alpha(x), \beta(x) \in k[x]$ nonzero and $\Omega_1, \Omega_2 \in k[x, y]$. Now substituting into (68), we have that

$$W((\alpha_0(x) + y\Omega_1)^2 + (x^2 + x^3)(\beta_0(x) + y\Omega_2)^2) \geq (1, 0)$$

which implies

$$W\left(\left(\frac{\alpha_0(x)}{\beta_0(x)}\right)^2 - (x^2 + x^3)\right) \geq (1, 0)$$

so that

$$\left(\frac{\alpha_0(x)}{\beta_0(x)}\right)^2 = x^2 + x^3,$$

a contradiction by Remark 10.3. Thus (66) holds.

Let

$$\mathcal{A} = \{\mu_\alpha = [V_1; \mu_\alpha(\varphi_\alpha) = W(\varphi_\alpha) \mid \varphi_\alpha \in \Sigma(V_1)]\}.$$

By (65) and (66), we have that $(1, 0)$ is the least upper bound of $\Lambda(\mathcal{A})$ in $(\mathbf{Z}^2)_{\text{lex}}$ but $(1, 0) \notin \Lambda(\mathcal{A})$. Thus \mathcal{A} does not have a maximal element.

Suppose that $\mu_\alpha \in \mathcal{A}$. Then $\mu_\alpha = [V_1; \mu_\alpha(\varphi_\alpha) = W(\varphi_\alpha)]$ with $\varphi_\alpha = z + h$ for some $h \in K$. Expand

$$Q = \varphi_\alpha^2 - 2h\varphi_\alpha + (h^2 - (x^2 + x^3)),$$

so that

$$\mu_\alpha(Q) \leq 2\mu_\alpha(\varphi_\alpha),$$

and $\mu_\alpha(Q) < (1, 0)$ by (66). Thus $Q \in \tilde{\Sigma}(\mathcal{A})$, and since Q has the smallest possible degree that a polynomial in $\tilde{\Sigma}(\mathcal{A})$ can have (it must have degree greater than $1 = d(V_1)$) we have that $d(\mathcal{A}) = 2$ and $Q \in \Sigma(\mathcal{A})$, and so Q is a limit key polynomial for \mathcal{A} . Let $V_2 = [\mathcal{A}; V_2(Q) = (1, 1)]$. Then the first simple admissible family in \mathcal{S} is $\mathcal{S}^{(1)} = \{V_1\} \cup \{\mathcal{A}\}$, and the second admissible family $\mathcal{S}^{(2)}$ begins with V_2 . Thus the first jump in \mathcal{S} is

$$s^{(1)}(\mathcal{S}) = \frac{\deg_z Q}{\deg_z z} = 2.$$

We have that $f = (Q^2 - y^2(x^2 + 2x^3))^2 - (y^6 + y^7)$. Let

$$U = Q^2 - y^2(x^2 + 2x^3)$$

as above. We have that $W(U) = (3, 0)$ since $W(f(z)) = \infty$. Let $\sum_{i=1}^{\infty} \beta_i x^i$ with $\beta_i \in k$ be a square root of $x^2 + 2x^3 = x^2(1+x)$ in $k[[x]]$. For $n \in \mathbf{Z}_+$, let $\overline{Q} = Q - y(\beta_1 x + \cdots + \beta_n x^n)$. Then

$$U = \overline{Q}^2 + 2y(\beta_1 x + \cdots + \beta_n x^n)\overline{Q} + y^2(\beta_1 x + \cdots + \beta_n x^n)^2 - y^2(x^2 + 2x^3),$$

so that

$$W(\overline{Q}(\overline{Q} + 2y(\beta_1 x + \cdots + \beta_n x^n))) > (2, n).$$

Thus

$$(69) \quad W(Q - y(\beta_1 x + \cdots + \beta_n x^n)) > (1, \frac{n}{2}) \text{ or } W(Q + y(\beta_1 x + \cdots + \beta_n x^n)) > (1, \frac{n}{2}).$$

Thus $d(V_2) = 2$ and so

$$\Sigma(V_2) = \{\varphi = Q + Az + B \mid A, B \in K \text{ and } V_2(\varphi) < W(\varphi)\}.$$

We will show that

$$(70) \quad \Lambda(V_2) = \{W(\varphi) \mid \varphi \in \Sigma(V_2)\} \subset \{1\} \times \mathbf{Z}_+.$$

We now prove equation (70). Suppose there exist $A, B \in K$ such that setting

$$\varphi = Q + Az + B,$$

we have that $W(\varphi) \geq (2, 0)$. We have that $W(Q) = W(Az + B)$. Expand

$$U = \varphi^2 - 2(Az + B)\varphi + (Az + B)^2 - y^2(x^2 + 2x^3).$$

Now $W(\varphi^2) \geq (4, 0)$ and $W((Az + B)\varphi) > (3, 0)$. Since $W(U) = (3, 0)$, we have that

$$W((Az + B)^2 - y^2(x^2 + 2x^3)) \geq (3, 0).$$

Thus

$$(1, 1) = W((Az + B)) = \min\{W(A) + (0, 1), W(B)\}.$$

We can thus write

$$A = y \left(\frac{\alpha_0(x) + y\Omega_1}{\gamma_0(x) + y\Omega_3} \right), B = y \left(\frac{\beta_0(x) + y\Omega_2}{\gamma_0(x) + y\Omega_3} \right)$$

with $\Omega_1, \Omega_2, \Omega_3 \in k[x, y]$, $\gamma_0(x) \neq 0$ and at least one of $\alpha_0(x), \beta_0(x) \neq 0$. Thus

$$W([(\alpha_0(x) + y\Omega_1)z + (\beta_0(x) + y\Omega_2)]^2 - (\gamma_0(x) + y\Omega_3)^2(x^2 + 2x^3)) \geq (1, 0),$$

and so

$$\begin{aligned} (1, 0) &\leq W((\alpha_0(x)z + \beta_0(x))^2 - \gamma_0(x)^2(x^2 + 2x^3)) \\ &= W(\alpha_0(x)^2 z^2 + 2\alpha_0(x)\beta_0(x)z + \beta_0(x)^2 - \gamma_0(x)^2(x^2 + 2x^3)) \\ &= W(\alpha_0(x)^2 Q + 2\alpha_0(x)\beta_0(x)z + (\alpha_0(x)^2(x^2 + x^3) + \beta_0(x)^2 - \gamma_0(x)^2(x^2 + 2x^3))). \end{aligned}$$

Thus

$$W(2\alpha_0(x)\beta_0(x)z + (\alpha_0(x)^2(x^2 + x^3) + \beta_0(x)^2 - \gamma_0(x)^2(x^2 + 2x^3))) \geq (1, 0).$$

But this implies that

$$(71) \quad \alpha_0(x)\beta_0(x) = 0$$

by (66) and thus

$$(72) \quad \alpha_0(x)^2(x^2 + x^3) + \beta_0(x)^2 - \gamma_0(x)^2(x^2 + 2x^3) = 0.$$

We have that $\alpha_0(x) = 0$ or $\beta_0(x) = 0$ by (71). If $\alpha_0(x) = 0$, then (72) becomes

$$\left(\frac{\beta_0(x)}{\gamma_0(x)}\right)^2 = x^2 + 2x^3$$

which is not a square in $k(x)$ by Remark 10.3, giving a contradiction. If $\beta_0(x) = 0$, then (72) becomes

$$\left(\frac{\alpha_0(x)}{\gamma_0(x)}\right)^2 = \frac{x+2}{x+1},$$

again giving a contradiction by Remark 10.3. Thus (70) holds.

Set

$$\mathcal{B} = \{\nu_\beta = [V_2; \nu_\beta(\varphi_\beta) = W(\varphi_\beta)] \mid \varphi_\beta \in \Sigma(V_2)\}.$$

Suppose $\nu_\beta \in \mathcal{B}$. Then $\nu_\beta = [V_2; \nu_\beta(\varphi_\beta) = W(\varphi_\beta)]$ with $\varphi_\beta = Q + Az + B$ for some $A, B \in K$. Expand

$$U = \varphi_\beta^2 - 2(Az + B)\varphi_\beta + (Az + B)^2 - y^2(x^2 + 2x^3)$$

to see that $\nu_\beta(U) \leq 2\nu_\beta(\varphi_\beta)$, and thus $\nu_\beta(U) < (3, 0)$ by (70). Thus $U \in \tilde{\Sigma}(\mathcal{B})$. We thus have that $d(\mathcal{B}) = 4$ or $d(\mathcal{B}) = 3$.

Let $\psi \in \Sigma(\mathcal{B})$, and define $V_3 = [\mathcal{B}; V_3(\psi) = W(\psi)]$. Then the second admissible family in \mathcal{S} begins with V_3 . Thus the second jump is

$$s^{(2)}(\mathcal{S}) = \frac{\deg_z \psi}{\deg_z Q} = \frac{3}{2} \text{ or } 2.$$

Thus

$$s^{(1)}(\mathcal{S})s^{(2)}(\mathcal{S}) \leq 4 < 8 = s^{\text{tot}}(\mathcal{S})$$

so there must be at least one more jump in the construction of \mathcal{S} so that $t \geq 3$.

11. EXTENSIONS OF ASSOCIATED GRADED RINGS AND SEMIGROUPS

We will consider in this section the conditions of finite generation of extensions of associated graded rings along a valuation and relative finite generation of extensions of valuation semigroups.

In this section, we will have the following assumptions. Suppose that A is a Noetherian local domain which contains an algebraically closed field k such that $A/m_A \cong k$. Let K be the quotient field of A and suppose that ν is a rank 1 valuation of K which dominates A , such that the residue field of the valuation ring of ν is k .

Suppose that S is a sub semigroup of a semigroup T . We say that T is a finitely generated module over S if there exists a finite number of elements t_1, \dots, t_r of T such that

$$T = (t_1 + S) \cup \dots \cup (t_r + S).$$

With our assumptions, $\text{gr}_\nu(A)$ is isomorphic to the semigroup algebra $k[t^{S^A(\nu)}]$. Thus if $A \rightarrow B$ is an inclusion of domains and ω is an extension of ν to the quotient field of B which is nonnegative on B such that the residue field of ω is k , then $\text{gr}_\omega(B)$ is a finitely generated $\text{gr}_\nu(A)$ -module if and only if $S^B(\omega)$ is a finitely generated module over $S^A(\nu)$.

We have the following immediate corollary of Theorem 5.1.

Corollary 11.1. *Suppose that $f(z) \in A[z]$ is unitary and irreducible and there is a unique extension of ν to a valuation ω of $K[z]/(f(z))$ and the characteristic p of k does not divide $\deg_z f(z)$. Then $\text{gr}_\omega(A[z]/(f(z)))$ is a finitely generated $\text{gr}_\nu(A)$ -module and $S^{A[z]/(f(z))}(\omega)$ is a finitely generated module over the semigroup $S^A(\nu)$.*

The following corollary addresses the case when the extension of valuations is not unique. It is an immediate corollary of Theorem 8.2.

Corollary 11.2. *Further suppose that A is excellent. Suppose that $f(z) \in A[z]$ is unitary, irreducible and separable and ω is a valuation of $K[z]/(f(z))$ which extends ν and there is no defect in the extension ($\delta(\omega/\nu) = 1$). Then there exists a birational extension A_1 of A which is dominated by ν such that $\text{gr}_\omega(A_1[z]/(f(z)))$ is a finitely generated $\text{gr}_\nu(A_1)$ -module and $S^{A_1[z]/(f(z))}(\omega)$ is a finitely generated module over the semigroup $S^{A_1}(\nu)$.*

If we remove any of the assumptions of Corollary 11.1, then the conclusions of the corollary are false, as is shown in the following three examples. We consider finite extensions $A \rightarrow B$ where A and B are excellent, B is a domain with quotient field L and ω is an extension of ν to L which dominates B .

Example 11.3. *There exists a finite extension $A \rightarrow B$ such that ω is the unique extension of ν to $L = \text{QF}(B)$, p does not divide $[L : K]$ but $\text{gr}_\omega(B)$ is not a finitely generated $\text{gr}_\nu(A)$ -module and $S^B(\omega)$ is not a finitely generated module over the semigroup $S^A(\nu)$.*

In particular, the representation of B as a ‘‘hypersurface singularity’’ over A is essential to the conclusions of Theorem 5.1 and Corollary 11.1.

Example 11.4. *There exists an extension $A \rightarrow B = A[z]/(f(z))$ where $f(z)$ is unitary and irreducible, such that p does not divide $\deg_z f(z)$ but the extension ω of ν to a valuation of $L = \text{QF}(B)$ is not unique such that $\text{gr}_\omega(B)$ is not a finitely generated $\text{gr}_\nu(A)$ -module and $S^B(\omega)$ is not a finitely generated module over the semigroup $S^A(\nu)$.*

Example 11.4 shows that the condition that ω is the unique extension of ν is necessary in Theorem 5.1 and Corollary 11.1, and that the birational extension $A \rightarrow A_1$ in the conclusions of Corollary 11.2 is necessary.

Example 11.5. *There exists an extension $A \rightarrow B = A[z]/(f(z))$ where $f(z)$ is unitary and irreducible, such that the extension ω of ν to a valuation of $L = \text{QF}(B)$ is unique but p divides $\deg_z f(z)$ such that $\text{gr}_\omega(B)$ is not a finitely generated $\text{gr}_\nu(A)$ -module and $S^B(\omega)$ is not a finitely generated module over $S^A(\nu)$. In the example, $\delta(\omega/\nu) = 1$.*

Example 11.5 shows that the condition that $p \nmid \deg_z f(z)$ is necessary in Corollary 11.1. In the remainder of this section, we will construct these three examples.

Examples 11.3 and 11.4 will be obtained from Example 9.3 of [9]. In Example 9.3 [9], k is an arbitrary field. We will make the further restriction that k is an algebraically closed field of characteristic $p > 2$. Let $T = k[x, y]_{(x, y)}$, a localization of a polynomial ring in two variables, and R be the subring $R = k[x^2, xy, y^2]_{(x^2, xy, y^2)}$. Let ω be the rational rank 1 valuation dominating T which is determined by the generating sequence

$$P_0 = x, P_1 = y, P_2 = y^3 - x^5$$

and

$$P_{i+1} = P_i^3 - x^{a_i} P_{i-1} \text{ for } i \geq 2$$

where a_i is even, and chosen so that $S^T(\omega)$ is not a finitely generated module over $S^R(\nu)$, where ν is the restriction of ω to the quotient field M of R . Let N be the quotient field of T .

Since the characteristic of k is not equal to 2, N is Galois over M , and the Galois group is generated by the involution σ defined by $\sigma(x) = -x$ and $\sigma(y) = -y$. Given $0 \neq g \in T$, we expand

$$g = \sum \alpha_{i_0, i_1, \dots, i_r} P_0^{i_0} P_1^{i_1} \dots P_r^{i_r}$$

with $\alpha_{i_0, i_1, \dots, i_r} \in k$, $i_0 \in \mathbf{N}$ and $0 \leq i_j < 3$ for $1 \leq j$, so that

$$\omega(g) = \min\{i_0\omega(P_0) + i_1\omega(P_1) + \dots + i_r\omega(P_r) \mid \alpha_{i_0, i_1, \dots, i_r} \neq 0\}.$$

Then

$$\sigma(g) = \sum \alpha_{i_0, i_1, \dots, i_r} (-1)^{i_0 + i_1 + \dots + i_r} P_0^{i_0} P_1^{i_1} \dots P_r^{i_r}$$

and thus $\omega(\sigma(g)) = \omega(g)$. Since the extensions of a valuation in a finite Galois extension are conjugate (Corollary 3 to Theorem 12, page 66 [40]), we have that ω is the unique extension of ν to N .

We now give a direct verification that T is not isomorphic to $R[z]/(f(z))$ for some $f(z) \in R[z]$. This follows since for a maximal ideal m in $R[z]/(f(z))$, we have that

$$\dim_k m/m^2 \geq 3 > 2 = \dim_k m_T/m_T^2.$$

We thus have that $R \rightarrow T$ gives Example 11.3.

In Example 9.4 [9], it is shown that in the natural extension $S \rightarrow T$, where $S = k[u, v]_{(u, v)}$ and $u = x^2, v = y^2$, with valuation μ obtained by restricting ω to the quotient field of S , that $S^T(\omega)$ is not a finitely generated $S^S(\mu)$ -module. Now we have a factorization of our extension $S \rightarrow U \rightarrow T$ where $U = k[x, v]_{(x, v)}$. Now $U \cong S[z]/(z^2 - u)$ and $T \cong U[z]/(z^2 - v)$. Let τ be the restriction of ω to the quotient field L of U .

Now we must have that $S^U(\tau)$ is not a finitely generated $S^S(\mu)$ -module or $S^T(\omega)$ is not a finitely generated $S^U(\tau)$ -module since $S^T(\omega)$ is not a finitely generated $S^S(\mu)$ -module.

We necessarily have by Corollary 11.1 that either τ is not the unique extension of μ to L or ω is not the unique extension of τ to N , giving Example 11.4.

In [12], a general theory of eigenfunctions for a valuation is developed for two dimensional quotient singularities, and a complete characterization is given of when the resulting extension of associated graded rings along the valuation is finite.

We now construct Example 11.5. Let $A = k[u, v]_{(u, v)}$ with quotient field K and let ν be the valuation of K which dominates A constructed in Theorem 7.38 [11] and analyzed in Section 9. Let $f(x) = x^p + ux^{p-1} - u$. It is shown in Theorem 7.38 [11] that there is a unique extension of ν to a valuation ω of $L = K[x]/(f(x))$. The extension is immediate, with defect $\delta(\omega/\nu) = p$. Let $B = A[x]/(f(x))$.

We see from the generating sequence P_0, \dots, P_i, \dots recalled in the beginning of Section 9 that $\text{gr}_\nu(A) \cong k[\overline{P}_0, \overline{P}_1, \dots]/I$ where

$$I = (\overline{P}_1^{p^2} - \overline{P}_0, \overline{P}_i^{p^2} - \overline{P}_0^{p^{2i-2}} \overline{P}_{i-1} \text{ for } i \geq 2).$$

It is shown in formulas (35) and (36) of [5] that

$$U_0 = x, U_1 = v, U_2 = v^p - x$$

and for $j \geq 2$,

$$U_{j+1} = U_j^p - x^{p^{2j-2}} U_{j-1} \text{ if } j \text{ is odd,}$$

$$U_{j+1} = U_j^{p^3} - x^{p^{2j-1}} U_{j-1} \text{ if } j \text{ is even}$$

is a generating sequence for ω in B . Thus $\text{gr}_\nu(B) \cong k[\overline{U}_0, \overline{U}_1, \dots]/J$ where

$$J = (\overline{U}_1^p - \overline{U}_0, \overline{U}_i^p - \overline{U}_0^{p^{2i-2}} \overline{U}_{i-1} \text{ for } i \geq 2 \text{ odd, } \overline{U}_i^{p^3} - \overline{U}_0^{p^{2i-1}} \overline{U}_{i-1} \text{ for } i \geq 2 \text{ even}).$$

Thus $\overline{U}_n^p = \overline{P}_n$ if n is even and $\overline{U}_n = \overline{P}_n$ if n is odd, and so $\text{gr}_\omega(B)$ is not a finitely generated $\text{gr}_\nu(A)$ -module and $S^\omega(B)$ is not a finitely generated $S^\nu(A)$ -module.

REFERENCES

- [1] S. Abhyankar, On the valuations centered in a local domain, *American Journal of Mathematics* 78 (1956), 321 - 348.
- [2] R. Brown, Valuations, primes and irreducibility in polynomial rings and rational function fields, *Trans. Amer. Math. Soc.* 174 (1972), 451 - 488.
- [3] S.D. Cutkosky, A generalization of the Abhyankar Jung theorem to associated graded rings of valuations, *Math. Proc. Cambridge Phil. Soc.* 160 (2016), 233 - 253.
- [4] S.D. Cutkosky, Finite generation of extensions of associated graded rings along a valuation, *Journal London Math. Soc.* 98 (2018), 177-203.
- [5] S.D. Cutkosky, Ramification of valuations and local rings in positive characteristic, *Communications in Algebra* 44 (2016), 2828-2866.
- [6] S.D. Cutkosky, The role of defect and splitting in finite generation of extensions of associated graded rings along a valuation, *Algebra Number Theory* 11 (2017), 1461 - 1488.
- [7] S.D. Cutkosky and H. Mourtada, Defect and Local Uniformization, arXiv:1711.02726.
- [8] S.D. Cutkosky, H. Mourtada and B. Teissier, On uniqueness of finite extensions of monomial valuations and their uniformization.
- [9] S.D. Cutkosky and Pham An Vinh, Valuation semigroups of two dimensional local rings, *Proceedings of the London Mathematical Society* 108 (2014), 350 - 384.
- [10] J. Decaup, W. Mahboub and M. Spivakovsky, Abstract key polynomials and comparison theorems with the key polynomials of MacLane - Vaquié, arXiv:1611.006392v1
- [11] S.D. Cutkosky and O. Piltant, Ramification of Valuations, *Advances in Math* 183 (2004), 1 - 79.
- [12] A. Dutta, Generating sequences and semigroups of valuations and 2-dimensional normal local rings, ArXiv:1804.04704v1.
- [13] S. El Hitti and L. Ghezzi, Dependent Artin-Schreier defect extensions and strong monomialization, *J. Pure and Applied Alg.* 220 (2010), 1331 - 1342.
- [14] O. Endler, *Valuation Theory*, Springer Verlag, New York, Heidelberg, Berlin, 1972.
- [15] R. Goldin and B. Teissier, Resolving singularities of plane analytic branches with one toric morphism, in: *Resolution of singularities, a research textbook in honor of O. Zariski*, edited by H. Hauser, J. Lipman, F. Oort and A. Quirós, *Progress in Math.*, Vol. 181, Birkhäuser, Basel, 200, 315-340.
- [16] F.J. Herrera Govantes, M.A. Olalla Acosta, W. Mahboub and M. Spivakovsky, Key Polynomials for simple extensions of valued fields, arXiv:1406.0657.
- [17] I. Kaplansky, Maximal fields with valuations, *Duke Math. J.* 9 (1942), 303 - 321.
- [18] O. Kashcheyeva, Constructing examples of semigroups of valuations, *J. Pure Appl. Algebra* 220 (2016), 3826-3860.

- [19] F.-V. Kuhlmann, Valuation theoretic and model theoretic aspects of local uniformization, in Resolution of Singularities - A Research Textbook in Tribute to Oscar Zariski, H. Hauser, J. Lipman, F. Oort, A. Quiros (eds.), Progress in Math. 181, Birkhäuser (2000), 4559 - 4600.
- [20] F.-V. Kuhlmann, Elimination of ramification I: The generalized stability theorem, Trans. AMS 362 (2010), 5697-5727.
- [21] F.-V. Kuhlmann, A classification of Artin-Schreier defect extensions and a characterization of defectless fields, Ill. J. Math 54 (2010), 397 - 445.
- [22] F.-V. Kuhlmann, The defect, in Commutative Algebra - Noetherian and non-Noetherian perspectives. Marco Fontana, Salah-Eddine Kabbaj, Bruce Olberding and Irena Swanson eds. Springer, 2011.
- [23] S. MacLane, A construction for absolute values in polynomial rings, Trans. Amer. Math. Soc. 40 (1936), 363 - 395.
- [24] S. MacLane, A construction for prime ideals as absolute values of an algebraic field, Duke Math. J, (1936), 492 - 510.
- [25] H. Mourtada, Jet schemes and generating sequences of divisorial valuations in dimension two, Michigan Math. J., Volume 66, Issue 1 (2017), 155-174.
- [26] M. Nagata, Local rings, Interscience publishers, New York, London, 1962.
- [27] J. Novacoski and M. Spivakovsky, Key polynomials and pseudo-convergent sequences, J. Algebra 495 (2018), 199 -219.
- [28] A. Ostrowski, Untersuchungen zur arithmetischen Theorie der Körper, Mathematische Zeitschrift, 39 (1935), 269-404.
- [29] M. Raynaud, Anneaux Locaux Henséliens, LNM 169 Springer Verlag, Berlin-Heidelberg, New York, 1970.
- [30] P. Ribenboim, Théorie des valuations, Les Presses Universitaires de Montréal, 1964.
- [31] J.-C. San Saturnino, Defect of an extension, key polynomials and local uniformization, J. Algebra 431 (2017), 91 - 119.
- [32] M. Spivakovsky, Valuations in function fields of surfaces, Amer. J. Math. 112 (1990), 107 - 156.
- [33] B. Teissier, Appendix to: The moduli problem for plane branches, by O. Zariski, translated by Ben Lichtin, University Lecture Series, Vol. 39, A.M.S., 2006. In French: Le problème des modules pour les branches planes, Hermann, Paris 1986, et Ecole Polytechnique, Centre de Mathématiques, 1973.
- [34] B. Teissier, Valuations, deformations, and toric geometry, in Valuation theory in interaction, Proceedings of the second international conference on valuation theory, Vol. II, Fields Institute Communications 33, A.M.S. Providence, R.I., 2013 , 351-459.
- [35] B. Teissier, Overweight deformations of affine toric varieties and local uniformization, in Valuation theory in interaction, Proceedings of the second international conference on valuation theory, Segovia, El Escorial, 2011. Edited by A. Campillo, F.-V. Kuhlmann and B. Teissier. European Math. Soc. Publishing House, Congress Reports Series, Sept. 2014, 474-565.
- [36] M. Vaquié, Extension d'une valuation, Trans. AMS 359 (2007), 3439 - 3481.
- [37] M. Vaquié, Famille admissible de valuations et défaut d'une extension, J. Algebra 311 (2007), 859 - 876.
- [38] M. Vaquié, Extensions de valuation et polygone de Newton, Annales de L'Institut Fourier 58 (2008), 2503 - 2541.
- [39] O. Zariski and P. Samuel, Commutative Algebra Volume I, Van Nostrand, 1958.
- [40] O. Zariski and P. Samuel, Commutative Algebra Volume II, Van Nostrand, 1960.

SDC: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA
Email address: cutkoskys@missouri.edu

HM: IMJ-PRG, CNRS, UNIVERSITÉ PARIS DIDEROT, INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, BÂT. SOPHIE GERMAIN, PLACE AURÉLIE NEMOURS, F-75013, PARIS, FRANCE.
Email address: hussein.mourtada@imj-prg.fr

BT: IMJ-PRG, CNRS, UNIVERSITÉ PARIS DIDEROT, INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, BÂT. SOPHIE GERMAIN, PLACE AURÉLIE NEMOURS, F-75013, PARIS, FRANCE.
Email address: bernard.teissier@imj-prg.fr