

A computation of Kontsevich Integral of torus knots *

Julien Marché †

April 2004

Abstract

We study the rational Kontsevich integral of torus knots. We construct explicitly a series of diagrams made of circles joined together in a tree-like fashion and colored by some special rational functions. We show that this series codes exactly the unwheeled rational Kontsevich integral of torus knots, and that it behaves very simply under branched coverings. Our proof is combinatorial. It uses the results of Wheels and Wheeling and various spaces of diagrams.

1 Introduction and notations

In 1998, in [BNGRT00], a conjecture was formulated about a precise expression for the Kontsevich integral of the unknot (later it was proved, see for instance [BNLT03]). Until now, we do not know any complete formula for this powerful invariant for non-trivial knots. Let \mathcal{B} be the usual space of uni-trivalent diagrams, which is the target space of the Kontsevich integral. We define a localization of \mathcal{B} called \mathcal{B}_s . Unfortunately, the localization map is not injective in high loop degrees. Then, we give a formula for the family of torus knots which takes its values in \mathcal{B}_s and which we state in a “rational” form.

The starting point of our proof is a well-known formula: it has been used by Christine Lescop (see [Les99]) and Dror Bar-Natan in unpublished work. From this expression, we construct a sequence of series of diagrams which are all obtained by “gluing wheels” and which converges to the unwheeled Kontsevich integral of torus knots. (See Section 1.1 for the precise meaning of “unwheeled” here.) Hence, we present the result in a compact way by using “gluing graphs”, which we will define along the way.

Then, in the third part, we compute a rational form of the preceding expression and show that only tree-like gluing diagrams appear as it is suggested by figure 1. More precisely, we show the following theorem:

Theorem. *Let D be the operator on $\mathbb{Q}(t)$ defined by $Dg(t) = tg'(t)$ and let $h(t) = \frac{t+1}{t-1}$ (The operator D acts as $\frac{d}{dx}$ on $g(\exp(x))$).*

There is a series of diagrams $Y_{p,q}^{rat}$ obtained by inserting circles in vertices of tree graphs, such that circles corresponding to vertices of valence k are colored by $D^{k-1}h(t^p)$, $D^{k-1}h(t^q)$ or $D^{k-1}h(t^{pq})$.

Applying the “hair map” (in other words, substituting t with the exponential of a small leg attached to the circle), we obtain a series in \mathcal{B}_s which is equal to the logarithm of the unwheeled Kontsevich integral of the torus knot of parameters p, q plus a fixed series $\log\langle\Omega, \Omega\rangle$. (see 1.1 for precise definitions)

As a consequence of this computation, we show that the operator Lift_r which corresponds to cyclic branched coverings of S^3 along the knot simply acts on $Y_{p,q}^{rat}$ by multiplying a diagram D by $r^{-\chi(D)}$ where χ is the Euler characteristic.

**keywords:* finite type invariants, Kontsevich integral, torus knots, Wheels and Wheeling, rationality. 2000 *Mathematics Subject Classification:* 57M27.

†Institut de Mathématiques de Jussieu, Équipe “Topologie et Géométries Algébriques” Case 7012, Université Paris VII, 75251 Paris CEDEX 05, France. e-mail: marche@math.jussieu.fr




Figure 1: Diagrams appearing in the unwheeled Kontsevich integral of torus knots

This article extends the result of our previous article [Mar] where we computed the unwheeled Kontsevich integral of torus knots up to degree 3. Here, we give a formula for all degrees using quite different methods.

Lev Rozansky also computed formulas for the loop expansion of torus knots in the weight system associated to \mathfrak{sl}_2 , see [Roz97]. The computation of the 2-loop part of torus knots has been done independently by Tomotada Ohtsuki in [Oht] who computed more generally a formula for 2-loop part of knots cabled by torus knots.

We would like to thank Stavros Garoufalidis, Pierre Vogel, and Marcos Marino for useful remarks and Gregor Masbaum for his remarks and his careful reading.

1.1 Normalizations of the Kontsevich integral

Let K be a knot in S^3 and suppose that K has a banded structure with self-linking 0. We will note $Z(K)$ the Kontsevich integral of K in the algebra \mathcal{A} of trivalent diagrams lying on a circle.

Let \mathcal{B} be the algebra of uni-trivalent diagrams. It is well known that the Poincaré-Birkhoff-Witt map $\chi : \mathcal{B} \rightarrow \mathcal{A}$ is an isomorphism but not an algebra isomorphism. We will note σ its inverse.

If U is the trivial knot, we define $\Omega = \sigma Z(U)$. The map $\Upsilon = \chi \circ \partial_\Omega : \mathcal{B} \rightarrow \mathcal{A}$ defined by instance in [Thu00] is known to be an algebra isomorphism. The quantity $Z^\boxtimes(K) = \Upsilon^{-1}Z(K)$ will be called unwheeled Kontsevich integral. It behaves better than $\sigma Z(K)$ under connected sum and cyclic branched coverings.

For each knot K , the quantities $Z(K)$, $\sigma Z(K)$ and $Z^\boxtimes(K)$ are group-like, which means that they are exponentials of a series of connected diagrams. We will note respectively $z(K)$, $\sigma z(K)$ and $z^\boxtimes(K)$ the logarithm of these quantities.

1.2 Loop degree and rationality

If D is a connected diagram of \mathcal{B} , its first Betti number defines a degree called loop degree. The loop degree 1 part of $\sigma Z(K)$ or $Z^\boxtimes(K)$ is well-known: it only depends on the Alexander polynomial of K . For the higher degrees, very little is known. There are formulas for the 2-loop part of small knots in Rozansky's table (see [Roz03]), and we can find in [Gar] a formula for the 2-loop part of untwisted whitehead doubles. In the sequel, we give a formula for the full Kontsevich integral of torus knots.

In order to make precise computations, we will need the following formalism which is yet another version of “diagrams with beads” used in [Roz03],[Kri],[GKb] for instance, and generalized by P. Vogel in [Vog00].

Definition 1. *Let \mathcal{C} be the category whose objects are free abelian groups of finite rank and morphisms are linear isomorphisms. We call \mathcal{C} -module a functor from \mathcal{C} to the category of \mathbb{Q} -vector spaces. In the sequel, we will associate to any \mathcal{C} -module F a \mathbb{Q} -vector space $\mathcal{D}(F)$ which will be called the space of diagrams decorated by F . This construction will be a functor from the category of \mathcal{C} -modules to the category of \mathbb{Q} -vector spaces.*

Let F be a \mathcal{C} -module and Γ be a finite trivalent graph with local orientations at vertices (we allow Γ to have connected components which are circles).

We define $\mathcal{D}(F)$ as the quotient of $\bigoplus_{\Gamma} F(H^1(\Gamma, \mathbb{Z}))$ by the following relations:

- If Γ is isomorphic to Γ' via a map ϕ , then we identify $x \in F(\Gamma')$ and $F(\phi^*)(x) \in F(\Gamma)$ for any x in $F(\Gamma')$.
- If Γ and Γ' just differ by the orientation of a vertex, then we identify x in $F(H^1(\Gamma, \mathbb{Z}))$ with $-x$ in $F(H^1(\Gamma', \mathbb{Z}))$.
- If Γ is a graph with one four-valent vertex, we note $\Gamma_I, \Gamma_H, \Gamma_X$ the three standard resolutions of this vertex. We have canonical identifications between $H^1(\Gamma, \mathbb{Z}), H^1(\Gamma_I, \mathbb{Z}), H^1(\Gamma_H, \mathbb{Z})$ and $H^1(\Gamma_X, \mathbb{Z})$. For every $x \in F(H^1(\Gamma, \mathbb{Z}))$, we add the relation $x_I = x_H - x_X$.

Let us give the main example: we define $F(H) = \mathbb{Q}[[H]] = \prod_{n \geq 0} S^n(H \otimes \mathbb{Q})$. The space $\mathcal{D}(F)$ is obtained by coloring graphs with 1-cohomology classes which can be materialized by small legs attached to the edges. It is not hard to see that $\mathcal{D}(F) \simeq \mathcal{B}$. This isomorphism is a convenient way to express series of diagrams.

Let $f(x)$ be the power series defined by $\frac{1}{2} \log \frac{\sinh x/2}{x/2}$. The famous wheel formula (see [Thu00]) states that $\sigma z(U) = f(x)$ (we suppose that the variable x is a generator of $H^1(\bigcirc, \mathbb{Z})$). Further, it was shown in [Kri] that the loop degree 1 part of $\sigma z(K)$ is $f(x) + Wh_K(x)$ where $Wh_K(x) = -\frac{1}{2} \log \Delta(e^x)$ and Δ is the Alexander polynomial of K . As we are interested in the higher loop degree part, we need to recall the rationality theorem which was proved in [GKb]. For that, we define a new space of diagrams with the help of a new functor.

Let $\mathbb{Q}[\exp(H)]$ be the polynomial algebra on elements on the form $\exp(\lambda), \lambda \in H$ with the relation $\exp(\lambda + \mu) = \exp(\lambda)\exp(\mu)$ for all λ and μ in H . Then, $F(H) = \mathbb{Q}[\exp(H)]_{loc}$ is the localization of the preceding space by expressions such as $P_1(\exp(h_1)) \cdots P_k(\exp(h_k))$ with $h_1, \dots, h_k \in H$ and $P_1, \dots, P_k \in \mathbb{Q}[h]$ such that $P_1(1) \cdots P_k(1) \neq 0$. It defines a new space of diagrams $\mathcal{B}^{rat} = \mathcal{D}(F)$ with a map $\text{Hair} : \mathcal{B}^{rat} \rightarrow \mathcal{B}$ induced by the Taylor expansion: $\mathbb{Q}[\exp(H)]_{loc} \rightarrow \mathbb{Q}[[H]]$ which is a map of \mathcal{C} -modules.

The rationality theorem tells us that there is an element $Z^{rat}(K) \in \mathcal{B}^{rat}$ such that

$$\sigma Z(K) = \exp(f(x) + Wh_K(x)) \text{Hair } Z^{rat}(K).$$

Respectively, there is such an element for the unwheeled invariant which will be recalled later in this article. A construction, developed in [GKb] gives $Z^{rat}(K)$ as an invariant of K ; this is not automatic because the Hair map is not injective, although it is so in small degrees (see [PM]).

2 Diagrammatic expressions of the integral

2.1 First diagrammatic expression

Let p and q be two coprime integers such that $p > 0$. We note $K_{p,q}$ the torus banded knot with parameters p and q and self-linking 0, and $L_{p,q}$ the torus banded knot with banding parallel to the torus on which it lies. This knot has self-linking pq , and its Kontsevich integral is a bit easier to compute.

The method of computation is inspired from [Les99]: we first compute the Kontsevich integral of the following braid.

Let p points be lying on the vertices of a regular p -gon. We note γ the braid obtained by rotating the whole picture by an angle $2\pi \frac{q}{p}$.

Let us associate to any one dimensional manifold Γ the space $\mathcal{A}(\Gamma)$ of trivalent diagrams lying on Γ . This defines a contravariant functor with respect to continuous maps relative to boundaries. Let ϕ_p^* be the map induced by the projection onto the first factor $\phi_p : [0, 1] \times \{1, \dots, p\} \rightarrow [0, 1]$ and \sphericalangle be the only

degree 1 diagram in $\mathcal{A}([0,1])$. Then a direct computation of monodromy of the K-Z connection shows that γ has Kontsevich integral equal to $\phi_p^*(\exp_{\#}(\frac{q}{2p}\curvearrowright))$.

The banded knot $L_{p,q}$ is obtained by closing the previous braid: this translates diagrammatically to the following: let ψ_p be the map from S^1 to itself defined by $\psi_p(z) = z^p$. Then, $Z(L_{p,q}) = \psi_p^*(\nu_{\#} \exp_{\#}(\frac{q}{2p}\curvearrowright))$, where $\nu = Z(U)$.

By lemma 4.10 of [Thu00], the map ψ_p^* viewed in \mathcal{B} has the following form: if $D \in \mathcal{B}$ has k legs (i.e. univalent vertices) then $\sigma\psi_p^*\chi D = p^k D$. We will note more simply D_p the result of this operation which looks like a change of variable.

Then, to compute $Z(K_{p,q})$ from $Z(L_{p,q})$, we only need to change the framing, that is $Z(K_{p,q}) = \exp_{\#}(-\frac{pq}{2}\curvearrowright)\#Z(L_{p,q})$. We will transform this product into the usual one by applying the unwheeling map Υ^{-1} . As a result, we will have a formula for $Z^{\curvearrowright}(K_{p,q})$.

We now sum up the steps of the computation:

1. Computation of $\sigma(\nu_{\#} \exp_{\#}(\frac{q}{2p}\curvearrowright))$
2. Change of variables $x \mapsto px$
3. Unwheeling

We recall that $\Upsilon = \chi \circ \partial_{\Omega}$ is an algebra isomorphism and that $\Upsilon^{-1}\nu = \frac{\Omega}{\langle \Omega, \Omega \rangle}$ and $\Upsilon^{-1}\curvearrowright = \curvearrowright - \Theta/24$. Here $\langle A, B \rangle$ is the sum over all ways of gluing all legs of A to all legs of B .

Then, to achieve the first step, we have to compute

$$\sigma(\nu_{\#} \exp_{\#}(\frac{q}{2p}\curvearrowright)) = \frac{\partial_{\Omega}(\Omega \exp(\frac{q}{2p}\curvearrowright))}{\langle \Omega, \Omega \rangle \exp(\frac{q}{48p}\Theta)}. \quad (1)$$

We explain below what will be a diagram colored by some parameters. In this setting, we will recall how usual operations on diagrams are obtained. Finally we will describe a new operation.

Definition 2. • Let P be a set of parameters. We denote $\mathcal{B}(P)$ the space of couples (D, f) where D is a uni-trivalent graph and f is a map from legs of D to P . We will say that legs of D are labeled or colored by elements of P . If $D \in \mathcal{B}$ and $x \in P$, we will write D_x the diagram D whose legs are colored by x .

- If $D \in \mathcal{B}(x)$, we define the diagram $D_{x+y} \in \mathcal{B}(x, y)$ by replacing legs of D by the same legs colored by x or y in all possible ways.
- If $D, E \in \mathcal{B}(P)$ and $x \in P$ we define $\langle D, E \rangle_x$ as the sum over all gluings of all x -legs of D on all x -legs of E . We also define $\partial_D E$ as the sum over all gluings of all x -legs of D on some x -legs of E . This operator satisfies $\partial_{D_x} E_x = \langle D_y, E_{x+y} \rangle_y$ and for $F \in \mathcal{B}(P)$, $\langle D_x, E_x F_x \rangle_x = \langle \partial_{E_x} D_x, F_x \rangle_x$.
- Let A_x and B_x be two series of diagrams in $\mathcal{B}(P)$ where P contains at least three colors: $\{x, y, z\}$. We define $A \cdot B = \langle A_{y+x}, B_{x+z} \rangle_x$. The y -legs of $A \cdot B$ will be called left legs and z -legs of $A \cdot B$ will be called right legs for obvious reasons. For rational numbers r and r' , the diagram ${}_r A \cdot {}_{r'} B$ will be the result of multiplying all left legs by r and all right legs by r' .

Moreover, if a_x and b_x are two series of connected diagrams, then $\exp(a)$, $\exp(b)$ and $\exp(a) \cdot \exp(b)$ are group-like, hence, we define ${}_y a \times {}_z b = \log({}_y \exp(a) \cdot \exp(b)_z)$.

Using the notation of definition 2, we compute

$$\partial_{\Omega}(\Omega \exp(\frac{q}{2p}\curvearrowright)) = \langle \Omega_x, \Omega_{x+y} \exp(\frac{q}{2p}{}_{x+y}\curvearrowright_{x+y}) \rangle_x.$$

We use the fact that ${}_{x+y}\curvearrowright_{x+y} = {}_x\curvearrowright_x + 2{}_x\curvearrowright_y + {}_y\curvearrowright_y$ and properties recalled in the preceding definition to obtain that this is equal to $\langle \partial_{\exp(\frac{q}{2p}{}_x\curvearrowright_x)} \Omega_x, \Omega_{x+y} \exp(\frac{q}{p}{}_x\curvearrowright_y) \rangle_x \exp(\frac{q}{2p}{}_y\curvearrowright_y)$.

Now we use the fundamental formula $\partial_{D_x} \Omega_x = \langle D_x, \Omega_x \rangle_x \Omega_x$ which is true if D contains only x -legs. The preceding expression reduces to $\langle \exp(\frac{q}{2p} \curvearrowright), \Omega \rangle \left(\frac{q}{p} \Omega \cdot \Omega \right) \exp(\frac{q}{2p} \curvearrowright)$.

Thanks to the identity $\langle \exp(\frac{q}{2p} \curvearrowright), \Omega \rangle = \exp(\langle \frac{q}{2p} \curvearrowright, \Omega \rangle) = \exp(\frac{q}{48p} \Theta)$, we may cancel this factor in the expression (1) above.

Multiplying by p to the power the number of legs, we obtain a formula for $Z(L_{p,q})$:

$$Z(L_{p,q}) = \frac{q \Omega \cdot \Omega_p \exp(\frac{qp}{2} \curvearrowright)}{\langle \Omega, \Omega \rangle}.$$

We obtain a formula for $Z^\boxtimes(K_{p,q})$ by unwheeling $Z(L_{p,q})$ and multiplying by $\Upsilon^{-1} \exp_{\#}(-\frac{pq}{2} \curvearrowright) = \exp(-\frac{pq}{2} \curvearrowright + \frac{pq}{48} \Theta)$.

$$Z^\boxtimes(K_{p,q}) = \partial_{\Omega}^{-1}(q \Omega \cdot \Omega_p \exp(\frac{qp}{2} \curvearrowright)) \frac{\exp(-\frac{pq}{2} \curvearrowright + \frac{pq}{48} \Theta)}{\langle \Omega, \Omega \rangle}. \quad (2)$$

2.2 A sequence converging to the integral of torus knots

In this part, we will start from the formula (2) to deduce new formulas by an induction process.

We define $\mathcal{B}^c = \mathcal{B}(\{\text{active, inert}\})$. There is a forgetful map $\mathcal{B}^c \rightarrow \mathcal{B}$. Let us put $\omega^{-1} = \frac{1}{p} \omega \times \omega_{\frac{1}{q}}$ and $\omega^0 = \omega$. These are elements of \mathcal{B}^c if we declare that all their legs are active.

Let us define three operators analogous to those of definition 2 but for diagrams in \mathcal{B}^c in the following way:

Definition 3. • For $A \in \mathcal{B}^c$ and $r \in \mathbb{Q}$, we set A_r to be the diagram A whose active legs are multiplied by r , viewed as an element of \mathcal{B} .

- If $A \in \mathcal{B}^c$ and $B \in \mathcal{B}$, the diagram $\partial_A B \in \mathcal{B}$ is the sum over gluings of all active legs of A on some legs of B . This is an element of \mathcal{B} .
- For $A \in \mathcal{B}$ and $B \in \mathcal{B}^c$ two diagrams, we set $A \cdot B \in \mathcal{B}^c$ to be the sum over all gluings of some active legs of B with some legs of A . The active legs of $A \cdot B$ are the remaining active legs of B . We also define $a \times b = \log(\exp(a) \times \exp(b))$. See figure 2 for both definitions.




Figure 2: The operators ∂ and \cdot .

Lemma 4. These two operators satisfy $\partial_A(BC) = \partial_{C \cdot A} B$.

We suppose all legs of ω^{-1} and ω^0 are active, and for all $n \geq 0$, we set $\omega^{n+1} = (\omega_{pq}^{n-1} - \omega_{pq}^n) \times \omega^n - \omega_{pq}^{n-1} + \omega_{pq}^n$. In this formula, the added term $\omega_{pq}^{n-1} - \omega_{pq}^n$ is considered inert.

Let us prove the following proposition

Proposition 5. For all $n \geq 0$, we have

1. The series $\exp(\frac{pq}{2}\frown)$ acts by derivation on active legs of $\exp(\omega^n)$ by multiplication by $\exp(\frac{pq}{48}\ominus)$. In formulas, we mean that $\partial_{\exp(\frac{pq}{2}\frown)}^{\text{active}} \exp(\omega^n) = \exp(\frac{pq}{48}\ominus) \exp(\omega^n)$. Here we have written ∂^{active} in order to mean that only active legs are glued in this process, contrary to the definition 3.

2.

$$Z^{\boxtimes}(K_{p,q}) = \left(\partial_{\exp(\omega^n)}^{-1} \exp(\frac{pq}{2}\frown + \omega_{pq}^{n-1}) \right) \frac{\exp(-\frac{pq}{2}\frown + \omega_{pq}^{-1} - \omega_{pq}^{n-1} + \frac{pq}{48}\ominus)}{\langle \Omega, \Omega \rangle}$$

3.

$$Z^{\boxtimes}(K_{p,q}) = \partial_{\exp(\omega^n)}^{-1} \left[\partial_{\exp(\omega^n)} \exp(\frac{pq}{2}\frown) \exp(\omega_{pq}^{n-1} - \omega_{pq}^n) \right] \frac{\exp(-\frac{pq}{2}\frown + \omega_{pq}^{-1} - \omega_{pq}^{n-1})}{\langle \Omega, \Omega \rangle}.$$

Moreover the sequence $\omega_{pq}^{-1} - \omega_{pq}^n$ converges to $\log \left(Z^{\boxtimes}(K_{p,q}) \langle \Omega, \Omega \rangle \right)$ with respect to loop degree.

Proof. The first formula is true for $n = 0$ thanks to the relation $\partial_D \Omega = \langle D, \Omega \rangle \Omega$ and is proved by a trivial induction as the derivation $\partial_{\exp(\frac{pq}{2}\frown)}^{\text{active}}$ and the gluing operation defining $\exp(\omega^{n+1})$ from $\exp(\omega^n)$ commute.

Suppose the second formula is true for some $n \geq 0$. To prove the second formula for $n + 1$, we need to show that:

$$\partial_{\exp(\omega^{n+1})}^{-1} (\exp(\frac{pq}{2}\frown + \omega_{pq}^n)) \exp(-\omega_{pq}^n) = \partial_{\exp(\omega^n)}^{-1} (\exp(\frac{pq}{2}\frown + \omega_{pq}^{n-1})) \exp(-\omega_{pq}^{n-1}),$$

which is equivalent to

$$\exp(\frac{pq}{2}\frown + \omega_{pq}^n) = \partial_{\exp(\omega^{n+1})} [\partial_{\exp(\omega^n)}^{-1} (\exp(\frac{pq}{2}\frown + \omega_{pq}^{n-1})) \exp(\omega_{pq}^n - \omega_{pq}^{n-1})]$$

but this is just an application of lemma 4.

The third formula is an application of the formula $\partial_{\exp(\omega^n)} \exp(\frac{pq}{2}\frown) = \exp(\frac{pq}{2}\frown + \frac{pq}{48}\ominus + \omega_{pq}^n)$ which is itself an application of the first proposition.

Let us prove the last property: we check by induction that $\omega^{n-1} - \omega^n$ has loop degree $> n$ for all $n \geq 0$, and that every connected component of ω^n has some active leg. It is also true that the operator $\partial_{\exp(\omega^n)}^{-1}$ has the form $\partial_{\exp(D)}$ for some series of diagrams D having at least one active leg. So, in the third formula, if the operator $\partial_{\exp(\omega^n)}^{-1}$ acts on $\exp(\omega_{pq}^{n-1} - \omega_{pq}^n)$, it creates diagrams with loop degree $> n + 1$ which means that $\omega_{pq}^{-1} - \omega_{pq}^n = \log(Z^{\boxtimes}(K_{p,q}) \langle \Omega, \Omega \rangle)$ modulo terms with loop degree $> n + 1$. This proves the last assertion. \square

2.3 An expression with gluing graphs

We remark that all diagrams appearing in ω^n are made by “gluing wheels”. We make this statement precise by introducing gluing graphs and give a presentation of $\log(Z^{\boxtimes}(K_{p,q}) \langle \Omega, \Omega \rangle)$ with gluing graphs. These graphs will be useful for finding a rational expression of the preceding expression. As a non-trivial application, we will show that only tree-like graphs appear in this expression.

Definition 6. Let P be a set of parameters. We denote by $\mathcal{S}(P)$ the \mathbb{Q} -vector space generated by unoriented finite graphs (V, E, h) with a map $V \rightarrow P$ where V is the set of vertices and E the set of edges and a map $h : P \rightarrow \mathbb{Q}[[x]]$. Elements of this space will be called gluing graphs colored by P . The number of edges is a degree on $\mathcal{S}(P)$. We complete $\mathcal{S}(P)$ with respect to this degree. This space has an obvious Hopf algebra structure whose primitive elements are formed of connected graphs.

We define a substitution map $s : \mathcal{S}(P) \rightarrow \mathcal{B}$ in the following way: if (X, h) is a diagram in $\mathcal{S}(P)$, we define $s(X, h)$ by gluing for all $a \in P$ and vertices v decorated by a the edges adjacent to v to wheels generated by $h(a)$ in all possible ways. Sometimes, we call a -colored the remaining legs generated by $h(a)$. It is clear that s is a map of Hopf algebras.

Let us give some examples: if $P = \{a\}$ $X = \bullet_a$ and $h(a) = f(x)$, then $s(X, h) = \omega$. In the same way, $s(\exp(\bullet_a), h) = \Omega$ and

$$s(a \bullet \bullet_b \bullet_c, h) = 16 \text{---} \bigcirc \text{---} \bigcirc \text{---} \bigcirc \equiv$$

where $h(a) = x^2$, $h(b) = x^2$ and $h(c) = x^4$.

Let us define two operations on $\mathcal{S}(P)$:

Definition 7. Given two connected gluing graphs X and Y with A some set of parameters of X and B some set of parameters of Y , we define $X_A \times_B Y = \log(\exp(X) \cdot \exp(Y))$ where $\exp(X) \cdot \exp(Y)$ is obtained by adding in all ways a finite number of edges from A -colored vertices of $\exp(X)$ to B -colored vertices of $\exp(Y)$.

Let us look at an example: take two formal parameters a and b , then

$$\exp(\bullet_a) \cdot \exp(\bullet_b) = \exp(\bullet_a + \bullet_b + a \bullet \bullet_b + a \bullet \bullet_b + \frac{1}{2} a \bullet \bullet_b \bullet_a + \frac{1}{2} b \bullet \bullet_a \bullet_b + a \bullet \bullet_b + \dots).$$

We have $s(X_A \times_B Y) = s(X)_A \times_B s(Y)$.

Definition 8. Let (X, h) be a diagram in $\mathcal{S}(a)$, $h(x) \in \mathbb{Q}[[x]]$, and $r \in \mathbb{Q} \setminus \{0\}$. We note a^r the operator which divides X by r^N where N is the sum of valences of a -colored vertices of X .

We have $s(X, h)_{ra} = s(a^r X, h(rx))$.

The main result of this section is the following:

Proposition 9. There is an explicit series of gluing graphs $X_{p,q}$ whose substitution is $\log(Z^{\boxtimes}(K_{p,q})\langle\Omega, \Omega\rangle)$.

Proof. We will show by induction that for all $n \geq 0$, $\omega^n = s(X^n)$ for some $X^n \in \mathcal{S}(P)$ where $P = \{*\} \cup \{a, b, c\}$. The first parameter is active and corresponds to $f(x)$, the three last parameters are inert and correspond to $f(px)$, $f(qx)$ and $f(pqx)$.

Firstly, the diagram ω_{pq}^{-1} has only inert legs and is obtained from the substitution of the gluing graph: $X_{pq}^{-1} = a^p b^q \bullet_a \times \bullet_b \in \mathcal{S}(a, b)$ where a and b are respectively associated to $f(px)$ and $f(qx)$.

We start the recursion at $n = 0$ by setting $X^0 = \bullet_*$. We have $s(X^0) = \omega^0$.

Take $n \geq 0$ and suppose we have constructed X^k for all $k \leq n$. Then, we set $X_{pq}^k = *^{pq} X^k|_{* \rightarrow c}$ for all $0 \leq k \leq n$ in such a way that we have $\omega_{pq}^k = s(X_{pq}^k)$ for all $-1 \leq k \leq n$ where c is associated to $f(pqx)$.

Then we set $X^{n+1} = (X_{pq}^{n-1} - X_{pq}^n)_{a,b,c} \times * X^n - (X_{pq}^{n-1} - X_{pq}^n)$. This definition satisfies $\omega^{n+1} = s(X^{n+1}) \in \mathcal{B}^c$. It gives a recursive way for finding all diagrams X^n and then we set $X_{p,q} = \lim_n (X^{-1} - X^n)$. □

Let us give all terms with less than two edges as an example. We use some notations to give a more compact form. Edges may be oriented or not, and vertices are expressed as a sum of integers in a box. To obtain the result, we color a vertex $\boxed{m+n}$ by $f(mx) + f(nx)$. Then, we obtain a sum of diagrams colored by integers. We divide each diagram of vertices x_1, \dots, x_n by the product of colors of each x_i to the power the number of adjacent edges which are not coming to this vertex.

For instance the diagram $\boxed{pq} \rightarrow \boxed{p} \leftarrow \boxed{q}$ is a graphical expression for $\frac{1}{pq^2} c \bullet \bullet_a \bullet_b$.

$$\begin{aligned}
\omega_{pq}^{-1} &= \boxed{p} + \boxed{q} + \boxed{p} \boxed{q} + \boxed{p} \boxed{q} + \frac{1}{2} \boxed{p} \boxed{q} \boxed{p} + \frac{1}{2} \boxed{q} \boxed{p} \boxed{q} + \text{terms with more than two edges} \\
\omega^0 &= \boxed{1} \\
\omega^1 &= \boxed{1} + \boxed{p+q-pq} \boxed{1} + \boxed{p+q-pq} \boxed{1} + \frac{1}{2} \boxed{1} \rightarrow \boxed{p+q-pq} \boxed{1} + \frac{1}{2} \boxed{p+q-pq} \boxed{1} \rightarrow \boxed{p+q-pq} \\
&\quad + \boxed{1} \rightarrow \boxed{p} \boxed{q} + \boxed{1} \rightarrow \boxed{q} \boxed{p} + \dots \\
\omega^2 &= \omega^1 - \boxed{p+q-pq} \boxed{pq} \boxed{1} - \boxed{1} \rightarrow \boxed{p+q-pq} \boxed{pq} + \dots \\
X_{p,q} &= \lim_n (\omega_{pq}^{-1} - \omega_{pq}^n) = \boxed{p} + \boxed{q} - \boxed{pq} + \boxed{p} \boxed{q} - \boxed{p+q-pq} \boxed{pq} + \boxed{p} \boxed{q} - \boxed{p+q-pq} \boxed{pq} \\
&\quad + \frac{1}{2} \boxed{p} \boxed{q} \boxed{p} + \frac{1}{2} \boxed{q} \boxed{p} \boxed{q} - \frac{1}{2} \boxed{pq} \rightarrow \boxed{p+q-pq} \boxed{pq} - \frac{1}{2} \boxed{p+q-pq} \boxed{pq} \rightarrow \boxed{p+q-pq} \\
&\quad - \boxed{pq} \rightarrow \boxed{p} \boxed{q} - \boxed{pq} \rightarrow \boxed{q} \boxed{p} + \boxed{p+q-pq} \boxed{pq} \boxed{pq} + \boxed{pq} \rightarrow \boxed{p+q-pq} \boxed{pq} + \dots
\end{aligned}$$

3 Rationality

The expression we have obtained until now is not rational. When trying to find a rational expression, we will show that expressions coming from non-tree gluing graphs vanish. For example, the graph $\boxed{p} \boxed{q}$ above will not appear in the final expression of the unwheeled Kontsevich integral of torus knots.

3.1 Diagrams with singular colorings

We need to work with diagrams which are colored by some singular algebraic expressions. For that we define two spaces of diagrams \mathcal{B}_s and \mathcal{B}_s^{rat} which fit in the following commutative diagram:

$$\begin{array}{ccc}
\mathcal{B}^{rat} & \longrightarrow & \mathcal{B}_s^{rat} \\
\downarrow \text{Hair} & & \downarrow \text{Hair} \\
\mathcal{B} & \longrightarrow & \mathcal{B}_s
\end{array}$$

Here we define $\mathcal{B}_s = \mathcal{D}(F)$ where $F(H) = (H \setminus \{0\})^{-1} \mathbb{Q}[[H]]$, the localization of $\mathbb{Q}[[H]]$ by elements of $H \setminus \{0\}$. The space \mathcal{B}_s^{rat} is constructed from the functor $F(H) = S^{-1} \mathbb{Q}[\exp(H)]$, where S is made of non zero expressions of the form $P_1(\exp(h_1)) \cdots P_k(\exp(h_k))$ for $P_1, \dots, P_k \in \mathbb{Q}[[h]]$ and $h_1, \dots, h_k \in H$.

The Hair map is again defined by the map induced by the Taylor expansion.

These constructions motivate the definition 1.2. In the article [Mar], we used a different construction using coloring of edges by elements of $\mathbb{Q}[[h]][[h^{-1}]]$. This construction was not convenient because the map from non-singular diagrams to singular ones fails to be injective even for small degrees as it is shown in the following identity:

$$-\bigcirc - \bigcirc - = -\bigcirc \overset{\frac{1}{h}h}{\parallel} \bigcirc - = 0.$$

To avoid this, we have changed the construction of singular diagrams in some way which avoids inverting null-homologous legs.

3.2 A formula for the substitution map

Let X be a connected gluing graph, with vertices x_1, \dots, x_N and power series f_1, \dots, f_N . The aim of this part is to give an explicit formula for $s(X) \in \mathcal{B}_s$.

To compute $s(X)$, we may glue at each vertex x edges incoming to this vertex to wheels generated by $f(x)$. Hence, we can choose a cycling ordering of edges around each vertex, compute the result of the gluings which respect this order, and sum over such orderings.

Fix an ordering $e_1^x, \dots, e_{k_x}^x$ of edges around each vertex x . Let Γ be the trivalent graph obtained by replacing x by a circle attached to the edges in the prescribed order, and $H = H^1(\Gamma, \mathbb{Z})$. Fix the order n_x of the wheel of $f(x)$ we will glue at x . Write x_1, \dots, x_{k_x} the elements of H coming from the edges e_1 to e_2, \dots, e_{k_x} to e_1 (see figure 3).




Figure 3: Diagram obtained by substitution

First, we glue the edge e_1 to any leg, so we have a factor n_x . Then all gluings are listed in the following formula:

$$\sum_{\text{orderings}, x} n_x \sum_{i_1 + \dots + i_{k_x} = n_x - k_x} x_1^{i_1} \dots x_{k_x}^{i_{k_x}}. \quad (3)$$

We explain an algorithmic reduction for this formula. Suppose there are two indices l and m such that $x_l \neq x_m \in H^1(\Gamma, \mathbb{Z})$, write $i_{lm} = i_l + i_m$. Then, $\sum_{i_1 + \dots + i_{k_x} = n_x - k_x} x_1^{i_1} \dots x_{k_x}^{i_{k_x}}$ may be replaced by the sum

$$\sum_{i_1 + \dots + \widehat{i_l} + \dots + \widehat{i_m} + \dots + i_{k_x} + i_{lm} = n_x - k_x} x_1^{i_1} \dots \widehat{x_l^{i_l}} \dots \widehat{x_m^{i_m}} \dots x_{k_x}^{i_{k_x}} \frac{x_l^{i_{lm}} - x_m^{i_{lm}}}{x_l - x_m} \in (H \setminus \{0\})^{-1} \mathbb{Q}[[H]]$$

This is a difference of sums of the same form but where each summand is now a product of $k_x - 1$ monomials, and with a degree one denominator. Observe that the sum of the exponents of the monomials in each summand is unchanged and equal to $n_x - k_x$ as before. We can perform such a reduction again for each one of the sums thus obtained a different number of times. This algorithm stops if all cohomology classes involved in the resulting expression are the same.

Suppose that an expression has been obtained from (3) by reducing it p_x times. Then this expression is a sum of identical terms each of which is equal to a cohomology class y to the power $n_x - k_x$ and divided by a product of elements of H . Counting the number of times this term occurs, the resulting expression is finally equal to

$$\frac{\binom{n_x - p_x - 1}{k_x - p_x - 1} y^{n_x - k_x}}{D_x}$$

where D_x is a product of p_x linear terms.

We can conclude the preceding computation in the following way:

Proposition 10. *Given a gluing graph X with power series f_1, \dots, f_N , its substitution is a finite combination of diagrams obtained by gluing edges to circles at each vertex, and decorated by expressions of the form $\left(\frac{f_i(y)}{y^p}\right)^{(k-p)}/D$ where y is a non zero cohomology class, p and k two integers and D is a product of p linear terms.*

3.3 Application to the integral of torus knots

We recall that in [GKa], the invariant $Z^{rat\boxtimes}$ is defined such that $\text{Hair } Z^{rat\boxtimes}(K) = \langle \Omega, \Omega \rangle Z^{\boxtimes}(K)$ for any knot K .

Our aim would be to compute $Z^{rat\boxtimes}(K_{p,q})$ but we have information only on $\text{Hair } Z^{rat\boxtimes}(K_{p,q})$. Hence we will compute it only up to the kernel of the Hair map. We have shown that $\langle \Omega, \Omega \rangle Z^{\boxtimes}(K) = s(X_{p,q})$. Hence we can apply the preceding section to $X_{p,q}$. We recall that $X_{p,q}$ is a gluing graph whose power series attached to vertices are either $f(px)$, $f(qx)$ or $f(pqx)$, where $f(x) = \frac{1}{2} \log \frac{\sinh(x/2)}{x/2}$. We compute $f'(x) = \frac{1}{4} \coth x/2 - \frac{1}{2x}$.

Thanks to the proposition 10, we can conclude that $s(X_{p,q})$ is obtained by decorating graphs by expressions such as $g(\exp(h))/D$ where $g \in \mathbb{Q}(t)$, $h \in H$ and D is a homogeneous polynomial in H .

Hence it is natural to define the following space:

Let H be a free abelian group of finite rank. We note $G(H)$ the image of $S^{-1}\mathbb{Q}[\exp(H)] \otimes (H \setminus \{0\})^{-1}\mathbb{Q}[H]$ in $(H \setminus \{0\})^{-1}\mathbb{Q}[[H]]$. This is again a \mathcal{C} -module and hence, we can define a space $\mathcal{B}' = \mathcal{D}(G)$.

Lemma 11. *The map $S^{-1}\mathbb{Q}[\exp(H)] \otimes (H \setminus \{0\})^{-1}\mathbb{Q}[H] \rightarrow G(H)$ is an isomorphism of \mathcal{C} -modules. We define the degree of $\frac{P}{Q} \in (H \setminus \{0\})^{-1}\mathbb{Q}[H]$ by $\deg(\frac{P}{Q}) = \deg(P) - \deg(Q)$. This degree extends to $G(H)$ and to \mathcal{B}' .*

Of course, we know from [GKb] that $Z^{\boxtimes}(K_{p,q})$ is in the image of the Hair map, and that the preimage can be obtained by decorating a diagram with expressions such as $\frac{P}{\Delta}$ where Δ is the Alexander polynomial. This rational function is regular at 1, in other words, $Z^{rat\boxtimes}(K_{p,q}) \in \mathcal{B}^{rat}$. Hence, $\text{Hair } Z^{rat\boxtimes}(K_{p,q})$ lies in the degree 0 part of \mathcal{B}' .

But in the process of substitution of a connected gluing graph, we add denominators unless all edges of the graph are null-homologous. In the same way, all terms containing fractions in $f'(px)$, $f'(qx)$ and $f'(pqx)$ will have a negative degree. This motivates the following definition:

Definition 12. *We define the gluing graph $Y_{p,q}$ as the tree part of $X_{p,q}$ decorated by the series*

$$\frac{1}{2} \log \sinh(px/2), \frac{1}{2} \log \sinh(qx/2) \text{ and } \frac{1}{2} \log \sinh(pqx/2) \text{ instead of } f(px), f(qx) \text{ and } f(pqx).$$

This is made of exactly all gluing graphs producing degree 0 diagrams in \mathcal{B}' by substitution and hence, we conclude that $s(Y_{p,q}) = s(X_{p,q}) = \text{Hair } z^{rat\boxtimes}(K_{p,q})$.

Theorem 13. *From $Y_{p,q}$, we define a diagram $Y_{p,q}^{rat}$ in \mathcal{B}_s^{rat} in the following way: at each vertex of $Y_{p,q}$ of valence k decorated by some expression $g(\exp(x))$, we replace the vertex by a circle, glue the edges arbitrarily to it and decorate the circle by the k -th derivative of $g(\exp(x))$ which in our case is a rational expression in $\exp(x)$.*

Then we have $\text{Hair } z^{rat\boxtimes}(K_{p,q}) = \text{Hair } Y_{p,q}^{rat}$.

Question 14. *Do we have $z^{rat\boxtimes}(K_{p,q}) = Y_{p,q}^{rat}$?*

4 Branched coverings

A great interest for rational expression of Kontsevich integral comes from its relation with branched coverings. More precisely, if $K_{p,q}$ is the torus knot of parameters p and q , and r is an integer, let us note $\Sigma^r(K_{p,q})$ be the pair formed of the cyclic branched covering of S^3 of order r over $K_{p,q}$ and the ramification link.

If r is coprime with p and q , the ramification locus is a knot, and the underlying 3-manifold is a rational homology sphere, the Brieskorn manifold $\Sigma(p, q, r)$.

In [GKa], a map Lift_r is described which intertwines rational invariant of the cyclic branched coverings and rational invariant of the initial knot in the following way:

$$Z^{\boxtimes rat}(\Sigma^r(K)) = \exp\left(\frac{\sigma_r(K)}{16}\Theta\right) \text{Lift}_r Z^{\boxtimes rat}(K).$$

We now study this map in the case of torus knots and prove the following proposition:

Proposition 15. *Call Π_r the operator on \mathcal{B}_s^{rat} which multiplies any diagram D by $r^{-\chi(D)}$ where χ is the Euler characteristic. then we have*

$$\text{Lift}_r Y_{p,q}^{rat} = \Pi_r Y_{p,q}^{rat}$$

Proof. The Lift_r map was defined only for diagrams decorated by fractions without poles at r -roots of unity. As we extended the decorations to all fractions, the definition of Lift_r makes sense for any diagram. In the definition of the Lift_r map, we need to express all denominators as polynomials of t^r . Then, we look at the numerators as a coloring by monomials, which is the same as a linear combination of 1-cohomology classes of the underlying graph. We keep only the classes divisible by r and divide them, then we put back denominators replacing t^r by t . Finally we multiply the result by r .

This construction is very easy in our case because we only need to know how the map Lift_r acts on derivatives of the fraction $\frac{t^n+1}{t^n-1}$ where $n = p, q, pq$. Let us write $h(t) = \frac{t^n+1}{t^n-1} \in \mathbb{Q}(t)$. In this space the operator Lift_r acts and also the derivation operator $Dg(t) = tg'(t)$. We can develop $h(t) = -1 - 2 \sum_{k \geq 1} t^{nk}$.

This expression shows that $\text{Lift}_r h(t) = -1 - 2 \sum_{k \geq 1, r|nk} t^{nk/r} = h(t)$ because n is coprime with r . Then, for $i > 0$, $D^i h(t) = -2 \sum_{k \geq 1} (nk)^i t^{nk}$. We check that $\text{Lift}_r D^i h(t) = -2 \sum_{k \geq 1, r|nk} (nk)^i t^{nk/r} = r^i D^i h(t)$.

This shows finally that Lift_r acts on a diagram of $Y_{p,q}^{rat}$ by multiplying it by r to the power $1 + \sum_i (v_i - 1)$ where v_i is the valence of the i -th vertex. This expression is the number of vertices of the diagram minus 1, hence the number of loops minus one.

This ends the proof of the proposition. □

Of course, we have not proved that $\text{Lift}_r z^{rat \boxtimes}(K_{p,q}) = \Pi_r z^{rat \boxtimes}(K_{p,q})$. Although the first diagram we know in the kernel of the Hair map has loop degree 17 (see [PM]), we do not know if the Hair map is injective in degree greater than 3.

References

- [BNGRT00] Dror Bar-Natan, Stavros Garoufalidis, Lev Rozansky, and Dylan P Thurston. Wheels, wheeling, and the kontsevich integral of the unknot. *Israel J. Math.*, 119, 2000.
- [BNL] D. Bar-Natan and R. Lawrence. A rational surgery formula for the lmo invariant. arXiv:math.GT/0007045. to appear in Israel J. Math.
- [BNLT03] Dror Bar-Natan, Thang T Q Le, and Dylan P Thurston. Two applications of elementary knot theory to Lie algebras and Vassiliev invariants. *Geom. Topol.*, 7:1–31, 2003, arXiv:math.QA/0204311.
- [Gar] S. Garoufalidis. Whitehead doubling persists. arXiv:math.GT/0003189.
- [GKa] S. Garoufalidis and A. Kricker. Finite type invariants of cyclic branched covers. arXiv:math.GT/0107220.
- [GKb] S. Garoufalidis and A. Kricker. A rational noncommutative invariant of boundary links. arXiv:math.GT/0105028.
- [Kri] A. Kricker. The lines of the kontsevich integral and rozansky’s rationality conjecture. arXiv:math.GT/0005284.

- [Les99] C. Lescop. Introduction to the kontsevich integral of framed tangles. Technical report, Grenoble Summer School, june 1999.
- [Mar] J. Marché. On kontsevich integral of torus knots. To appear in *Topology and its Applications*, arXiv:math.GT/0310111.
- [Mn] M. Mariño. Chern-simons theory, matrix integrals, and perturbative three-manifold invariants. arXiv:hep-th/0207096.
- [Oht] T. Ohtsuki. A cabling formula for the 2-loop polynomial of knots. arXiv:math.GT/0310216.
- [PM] B. Patureau-Mirand. Non-injectivity of the hair map. arXiv:math.GT/0202065.
- [Roz97] L. Rozansky. Higher order terms in the melvin-morton expansion of the colored jones polynomial. *Comm. Math. Phys.*, 183:291–306, 1997.
- [Roz03] L. Rozansky. A rationality conjecture about kontsevich integral of knots and its implications to the structure of the colored jones polynomial. *Topology Appl.*, 127:47–76, 2003.
- [Thu00] D. Thurston. *Wheeling: a diagrammatic analogue of the Duflo isomorphism*. PhD thesis, UC Berkeley, 2000, arXiv:math.QA/0006083.
- [Vog00] P. Vogel. Vassiliev theory. Technical report, MaPhySto, February 2000.