

ON SYNTOMIC REGULATORS I: CONSTRUCTIONS

WIESŁAWA NIZIOL

ABSTRACT. We show that classical Chern classes from higher (p -adic) K -theory to syntomic cohomology extend to logarithmic syntomic cohomology. These Chern classes are compatible – in a suitable sense – with addition, products, and λ -operations. They are also compatible with the canonical Gysin sequences and, via period maps, with logarithmic étale Chern classes. Moreover, they induce logarithmic crystalline Chern classes. This uses as a critical new ingredient the recent comparison of syntomic cohomology with p -adic nearby cycles [6] and p -adic motivic cohomology [7].

CONTENTS

1. Introduction	1
2. Syntomic cohomology	4
2.1. Syntomic cohomology	4
2.2. Syntomic-étale cohomology	8
2.3. Syntomic cohomology and motivic cohomology	12
3. Cohomology of classifying spaces	12
3.1. Classical computations	12
3.2. Syntomic computations	13
4. K -theory	22
4.1. K -theory of simplicial schemes	22
4.2. Log- K -theory	27
4.3. Operations on log- K -theory	31
5. Syntomic Chern classes	35
5.1. Classical syntomic Chern classes	35
5.2. Truncated syntomic Chern classes	37
5.3. Logarithmic syntomic Chern classes	38
6. Chern class maps and Gysin sequences	42
6.1. Basic properties of syntomic cohomology	42
6.2. Compatibility with Gysin sequences	53
References	60

1. INTRODUCTION

The study of syntomic regulators, a p -adic analogue of Deligne’s regulators, is important in p -adic Hodge Theory [28, 30] and in computations of special values of p -adic L -functions [34, 1]. In this paper we show that syntomic regulators have a well-behaved logarithmic version that is compatible, via the period map, with logarithmic étale regulators. In the sequel to this paper we will use it to extend results from proper schemes to open schemes with normal crossing compactification.

Let p be a prime. Let K be a complete discrete valuation field of mixed characteristic $(0, p)$ with perfect residue field; let \mathcal{O}_K be its ring of integers. Let X be a semistable scheme over \mathcal{O}_K and let D be the canonical horizontal normal crossings divisor on X . We equip X with the log-structure associated

Date: December 20, 2016.

This research was supported in part by the NSF grant DMS0703696 and the grant ANR-14-CE25.

to D and the special fiber. Alternatively, X could be a smooth scheme over \mathcal{O}_K with the log-structure associated to a divisor D as above. Denote by X_0 the special fiber of X and by X_n – the reduction mod p^n of X .

For $n \geq 1, i \geq 0$, let

$$\mathcal{S}'_n(i)_X := (U \rightarrow X) \mapsto \text{Cone}(F^i \text{R}\Gamma_{\text{cr}}(U_n) \xrightarrow{p^i - \varphi} \text{R}\Gamma_{\text{cr}}(U_n))[-1]$$

be the syntomic cohomology complex of sheaves on the étale site of X . The crystalline cohomology used is absolute, i.e. over \mathbf{Z}/p^n . Let N be a constant as in [6, Theorem 1.1]. Recall that, for $p \geq 3$, if K contains enough roots of unity¹ then N depends only on p ; in general it depends on p and e – the absolute ramification index of K .

The main goal of this paper is to prove the following theorem.

Theorem 1.1. *Fix $m \geq N$. Let $U = X \setminus D$. There exists a functorial compatible family of logarithmic syntomic Chern classes²*

$$\bar{c}_{i,j}^{\text{syn}} : K_j(U, \mathbf{Z}/p^n) \rightarrow H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_n(i)), \quad j \geq 2,$$

such that

- (1) they are compatible – in a suitable sense – with addition, products, and λ -operations.
- (2) they are twisted, i.e., in the case of trivial divisor D the Chern class $\bar{c}_{i,j}^{\text{syn}}$ is equal to the p^{mi} -multiple of the classical syntomic Chern class.
- (3) they are compatible with the canonical Gysin sequences, i.e., the Gysin sequences associated to D .
- (4) they are compatible, via the period maps $\alpha_{*,*}$ of Fontaine-Messing, with étale Chern classes, i.e., the following diagram commutes

$$\begin{array}{ccc} K_j(U, \mathbf{Z}/p^n) & \xrightarrow{\bar{c}_{i,j}^{\text{syn}}} & H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_n(i)) \\ \downarrow j^* & & \downarrow \alpha_{2i-j,i} \\ K_j(U_K, \mathbf{Z}/p^n) & \xrightarrow{p^{(m+1)i} \bar{c}_{i,j}^{\text{ét}}} & H_{\text{ét}}^{2i-j}(U_K, \mathbf{Z}/p^n(i)), \end{array}$$

where $j : U_K \hookrightarrow U$ is the natural open immersion.

Similarly, there exists a functorial compatible family of logarithmic syntomic Chern classes

$$c_{i,j}^{\text{syn}} : K_j(U) \rightarrow H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_n(i)), \quad j \geq 0,$$

with the above listed properties.

Remark 1.2. From the above theorem we obtain p -adic logarithmic Chern classes. More specifically, set

$$K_j(U, \mathbf{Q}_p) := \mathbf{Q} \otimes \varprojlim_n K_j(U, \mathbf{Z}/p^n), \quad H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_{\mathbf{Q}}(i)) := \mathbf{Q} \otimes \varprojlim_n H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_n(i)).$$

The limit of Chern classes $\bar{c}_{i,j}^{\text{syn}}$ divided by p^{mi} yields (untwisted) logarithmic Chern classes

$$\bar{c}_{i,j}^{\text{syn}} : K_j(U, \mathbf{Q}_p) \rightarrow H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_{\mathbf{Q}}(i)), \quad j \geq 2.$$

The strategy for proving the above theorem is well-known. In the simplest case, logarithmic cohomology is equal to the cohomology of the open set, where the log-structure is trivial. The definition of the logarithmic Chern classes is then immediate. To treat compatibility with Gysin sequences, one proves a version of Grothendieck-Riemann-Roch to extend the classical universal Chern classes to the logarithmic ones that are compatible (by definition) with the Gysin diagram and then one uses cohomological purity to show uniqueness of such an extension. This works, for example, for logarithmic ℓ -adic étale cohomology.

In general, logarithmic cohomology does not have the above mentioned property. This is the case, for example, for crystalline cohomology. Then one constructs a well-behaved new cohomology that dominates

¹See Theorem 2.2 for what this means.

²They vary with m in the obvious way.

the given cohomology and one constructs the universal logarithmic Chern classes into the new cohomology having all the necessary properties. For crystalline cohomology over a field, the new cohomology is the logarithmic de Rham-Witt cohomology.

Logarithmic syntomic cohomology $\mathcal{S}'_n(*)$ by itself does not behave well enough to employ the above strategy. In its classical form it satisfies a version of the projective space theorem and gives the correct even-degree cohomology groups of the classifying spaces. Hence it has well-behaved classical Chern classes. But it is too weak to allow those to be extended to logarithmic cohomology: it is not equal to syntomic cohomology of the open set, where the log-structure is trivial, nor does it satisfy purity.

To obtain a well-behaved logarithmic cohomology dominating syntomic cohomology one replaces syntomic cohomology by (Nisnevich) syntomic-étale cohomology $\mathcal{E}'_n(*)_{\text{Nis}}$. The latter is defined by gluing étale cohomology of the Tate twist $\mathbf{Z}/p^n(i)$ on the generic fiber with the syntomic cohomology $\mathcal{S}'_n(i)$ of the formal special fiber, projecting the result down to the Nisnevich site, and truncating it at i . We have a natural map $\mathcal{E}'_n(i)_{\text{Nis}} \rightarrow \mathcal{S}'_n(i)_{\text{Nis}}$, where the target is defined by projecting syntomic cohomology $\mathcal{S}'_n(i)$ to the Nisnevich site and truncating it at i . Hence it suffices to construct logarithmic universal Chern classes with values in syntomic-étale cohomology. This cohomology has the properties necessary for the above strategy to work. To get an idea why this could be the case, let us look at the simpler case when X is semistable. Recall that, by p -adic comparison theorems [6], the logarithmic syntomic cohomology $\mathcal{S}'_n(i)$ is approximated (via the Fontaine-Messing period map) by the p -adic nearby cycles $i^* \tau_{\leq j} \mathbf{R}j_* \mathbf{Z}/p^n(i)$, $i : X_0 \hookrightarrow X$, $j : U_K \hookrightarrow X$. It follows that the syntomic-étale cohomology $\mathcal{E}'_n(i)_{\text{Nis}}$ is approximated by the truncated étale cohomology projected onto the Nisnevich site: $\tau_{\leq i} \mathbf{R}(j\varepsilon)_* \mathbf{Z}/p^n(i)_{U_K}$, $\varepsilon : U_{K,\text{ét}} \rightarrow U_{K,\text{Nis}}$. But by the Beilinson-Lichtenbaum conjecture we have a quasi-isomorphism $\tau_{\leq j} \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)_{U_K} \simeq \mathbf{Z}/p^n(j)_{M,U_K}$, where $\mathbf{Z}/p^n(i)_M$ is the motivic cohomology. Hence the syntomic-étale cohomology $\mathcal{E}'_n(i)_{\text{Nis}}$ is approximated by $\tau_{\leq i} \mathbf{R}j_* \mathbf{Z}/p^n(i)_{M,U_K} = \mathbf{R}j_* \mathbf{Z}/p^n(i)_{M,U_K}$. In particular it is equal to syntomic-étale cohomology of the open set, where the log-structure is trivial. Moreover, since motivic cohomology has purity so does syntomic-étale cohomology. We got all the properties we wanted. In the more difficult case of good reduction the role of the motivic cohomology $\mathbf{Z}/p^n(i)_{M,U_K}$ is played by the motivic cohomology $\mathbf{Z}/p^n(i)_{M,U}$; that this can be done follows from the comparison between syntomic cohomology and p -adic motivic nearby cycles proved in [7].

The approximation mentioned above is done up to universal constants (that can be controlled). This results in the twisting of Chern classes in the above theorem. Keeping track of those constants is the most tedious part of the paper. For small Tate twists i these constants can be taken to be trivial.

Remark 1.3. It seems like an overkill to use a very difficult theorem like the Beilinson-Lichtenbaum conjecture to prove properties of such a seemingly simple object as the truncated étale cohomology $\tau_{\leq i} \mathbf{R}j_* \mathbf{Z}/p^n(i)$. Yet, even to prove the projective space theorem to define classical Chern classes with values in this cohomology and to construct Gysin sequences one uses [33, Theorem 4.1] the computations of mod p nearby cycles via symbols due to Bloch-Kato [4] which are closely related to the Bloch-Kato conjecture (and hence to the Beilinson-Lichtenbaum conjecture)³.

Remark 1.4. There is another strategy that we could have employed to construct logarithmic syntomic Chern classes. One starts with the classical Chern classes, proves the Grothendieck-Riemann-Roch theorem for them, and then uses it and Gysin sequences to induce logarithmic Chern classes on the complement of the divisor. This has to be done one irreducible divisor at a time; in particular, the logarithmic Chern classes have to have all the properties necessary for the standard proof of the Grothendieck-Riemann-Roch theorem to work (which, basically means, that they have to be compatible with the action of K_0 -groups in a suitable sense).

In [34], Somekawa tried to make this strategy work. His arguments work for removing one irreducible divisor but fail on the inductive step: he was not able to show that so obtained logarithmic Chern classes have good properties. This seems highly nontrivial. This paper shows that this strategy actually works but with syntomic-étale cohomology in place of syntomic cohomology. Purity of syntomic-étale cohomology is the key property that allows the inductive step. But this property also implies that

³It is interesting that it is much easier to prove projective space theorem and to construct Gysin sequences for syntomic cohomology: it simply reduces to the same for filtered crystalline cohomology, where it is immediate.

logarithmic cohomology is equal to cohomology of the open set, where the log-structure is trivial. Hence we have chosen this (instead of Gysin sequences) as the starting point of the construction of logarithmic syntomic Chern classes. Compatibility with Gysin sequences is then a theorem.

1.0.1. *Structure of the paper.* In Section 2 we review the definitions and basic properties of syntomic and syntomic-étale cohomologies and recall their relationship to p -adic nearby cycles and p -adic motivic cohomology. In Section 3 we study cohomology of classifying spaces: we compute their (Nisnevich) syntomic cohomology and we show that their Nisnevich syntomic-étale and syntomic cohomologies agree in even degrees. In Section 4 we review the basic facts concerning higher K -theory of (simplicial) schemes and operations on K -theory. In Sections 5 we define and study properties of classical Nisnevich syntomic and syntomic-étale Chern classes - by standard arguments this builds on the computations done in Section 3. Then we introduce logarithmic syntomic-étale Chern classes and discuss purity. In Section 6 we study compatibility of these Chern classes with Gysin sequences. We start with proving the properties of syntomic-étale cohomology that are needed for the proof of the Grothendieck-Riemann-Roch theorem. The proof of the theorem itself follows along standard lines.

Acknowledgments. I would like to thank Frédéric Déglise and Marc Levine for answering questions concerning motivic cohomology. Parts of this paper were written during my visits to the Institut Henri Poincaré in Paris, the Institut de Mathématiques de Jussieu, Columbia University, and the IAS at Princeton. I would like to thank these institutions for their support, hospitality, and for the wonderful working conditions they have supplied. Much of this paper was written in Café Nizioł in downtown Kołobrzeg, Poland. I would like to thank Jarosław Nizioł for creating such a pleasant coffee house to work at.

1.0.2. *Notation and Conventions.* Unless stated otherwise we work in the category of fine log-schemes.

Definition 1.5. Let $N \in \mathbf{N}$. For a morphism $f : M \rightarrow M'$ of \mathbf{Z}_p -modules, we say that f is p^N -*injective* (resp. p^N -*surjective*) if its kernel (resp. its cokernel) is annihilated by p^N and we say that f is p^N -*isomorphism* if it is p^N -injective and p^N -surjective. We define in the same way the notion of p^N -*exact sequence* or p^N -*acyclic complex* (complex whose cohomology groups are annihilated by p^N) as well as the notion of p^N -*quasi-isomorphism* (map in the derived category that induces a p^N -isomorphism on cohomology).

2. SYNTOMIC COHOMOLOGY

Let \mathcal{O}_K be a complete discrete valuation ring with fraction field K of characteristic 0 and with perfect residue field k of characteristic p . Let ϖ be a uniformizer of \mathcal{O}_K . Let $\mathcal{O}_{\bar{K}}$ denote the integral closure of \mathcal{O}_K in \bar{K} . Let $W(k)$ be the ring of Witt vectors of k with fraction field F (i.e. $W(k) = \mathcal{O}_F$); let e be the ramification index of K over F . Set $G_K = \text{Gal}(\bar{K}/K)$, and let σ be the absolute Frobenius on $W(\bar{k})$. For an \mathcal{O}_K -scheme X , let X_0 denote the special fiber of X . We will denote by \mathcal{O}_K and \mathcal{O}_K^\times the scheme $\text{Spec}(\mathcal{O}_K)$ with the trivial and the canonical (i.e., associated to the closed point) log-structure, respectively.

In this section we will briefly review the definitions of syntomic and syntomic-étale cohomologies and their basic properties. For details we refer the reader to [39, 2], [7, 2].

2.1. **Syntomic cohomology.** For a log-scheme X we denote by X_{syn} the small log-syntomic site of X . For a log-scheme X log-syntomic over $\text{Spec}(W(k))$, define

$$\mathcal{O}_n^{\text{cr}}(X) = H_{\text{cr}}^0(X_n, \mathcal{O}_{X_n}), \quad \mathcal{J}_n^{[r]}(X) = H_{\text{cr}}^0(X_n, \mathcal{J}_{X_n}^{[r]}),$$

where \mathcal{O}_{X_n} is the structure sheaf of the absolute log-crystalline site (i.e., over $W_n(k)$), $\mathcal{J}_{X_n} = \text{Ker}(\mathcal{O}_{X_n/W_n(k)} \rightarrow \mathcal{O}_{X_n})$, and $\mathcal{J}_{X_n}^{[r]}$ is its r 'th divided power of \mathcal{J}_{X_n} . Set $\mathcal{J}_{X_n}^{[r]} = \mathcal{O}_{X_n}$ if $r \leq 0$. There is a canonical, compatible with Frobenius, and functorial isomorphism

$$H^*(X_{\text{syn}}, \mathcal{J}_n^{[r]}) \simeq H_{\text{cr}}^*(X_n, \mathcal{J}_{X_n}^{[r]}).$$

It is easy to see that $\varphi(\mathcal{J}_n^{[r]}) \subset p^r \mathcal{O}_n^{\text{cr}}$ for $0 \leq r \leq p-1$. This fails in general and we modify $\mathcal{J}_n^{[r]}$:

$$\mathcal{J}_n^{<r>} := \{x \in \mathcal{J}_{n+s}^{[r]} \mid \varphi(x) \in p^r \mathcal{O}_{n+s}^{\text{cr}}\} / p^n,$$

for some $s \geq r$. This definition is independent of s . We can define the divided Frobenius $\varphi_r = \text{"}\varphi/p^{r\text{"}}$: $\mathcal{J}_n^{<r>} \rightarrow \mathcal{O}_n^{\text{cr}}$. Set

$$\mathcal{S}_n(r) := \text{Cone}(\mathcal{J}_n^{<r>} \xrightarrow{1-\varphi_r} \mathcal{O}_n^{\text{cr}})[-1].$$

We will write $\mathcal{S}_n(r)$ for the syntomic sheaves on $X_{m,\text{syn}}$, $m \geq n$, as well as on X_{syn} . We will also need the "undivided" version of syntomic complexes of sheaves:

$$\mathcal{S}'_n(r) := \text{Cone}(\mathcal{J}_n^{[r]} \xrightarrow{p^r-\varphi} \mathcal{O}_n^{\text{cr}})[-1]$$

as well as their twists

$$\mathcal{S}_n^i(r) := \text{Cone}(\mathcal{J}_n^{[r]} \xrightarrow{p^{r+i}-p^i\varphi} \mathcal{O}_n^{\text{cr}})[-1], \quad i \geq 0.$$

We note that the following sequence is exact for $r \geq 0$

$$(2.1) \quad 0 \longrightarrow \mathcal{S}_n(r) \longrightarrow \mathcal{J}_n^{<r>} \xrightarrow{1-\varphi_r} \mathcal{O}_n^{\text{cr}} \longrightarrow 0.$$

So, actually,

$$\mathcal{S}_n(r) := \text{Ker}(\mathcal{J}_n^{<r>} \xrightarrow{1-\varphi_r} \mathcal{O}_n^{\text{cr}}).$$

The natural map $\omega : \mathcal{S}_n^i(r) \rightarrow \mathcal{S}_n(r)$ induced by the maps $p^{r+i} : \mathcal{J}_n^{[r]} \rightarrow \mathcal{J}_n^{<r>}$ and $\text{Id} : \mathcal{O}_n^{\text{cr}} \rightarrow \mathcal{O}_n^{\text{cr}}$ has kernel and cokernel killed by p^{r+i} . So does the map $\tau : \mathcal{S}_n(r) \rightarrow \mathcal{S}'_n(r)$ induced by the maps $\text{Id} : \mathcal{J}_n^{<r>} \rightarrow \mathcal{J}_n^{[r]}$ and $p^{r+i} : \mathcal{O}_n^{\text{cr}} \rightarrow \mathcal{O}_n^{\text{cr}}$. We have $\tau\omega = \omega\tau = p^{r+i}$. There are also natural maps $\omega^a : \mathcal{S}_n^a(r) \rightarrow \mathcal{S}_n^{a+1}(r)$ induced by Id on $\mathcal{J}_n^{[r]}$ and by multiplication by p^{a+1} on $\mathcal{O}_n^{\text{cr}}$. We set $\omega' = \omega^0$. Write $\omega_a : \mathcal{S}_n^{a+1}(r) \rightarrow \mathcal{S}_n^a(r)$ for the map induced by Id on $\mathcal{O}_n^{\text{cr}}$ and by multiplication by p^{a+1} on $\mathcal{J}_n^{[r]}$. We have $\omega^a\omega_a = \omega_a\omega^a = p^{a+1}$.

We have versions of complexes $\mathcal{S}_n(r)$ and $\mathcal{S}'_n(r)$ on the large syntomic sites [40, 4.3]. If it does not cause confusion, we will write $\mathcal{S}_n(r)$, $\mathcal{S}'_n(r)$ for all these complexes as well as for $\text{R}\varepsilon_*\mathcal{S}_n(r)$, $\text{R}\varepsilon_*\mathcal{S}'_n(r)$, respectively, where $\varepsilon : X_{n,\text{syn}} \rightarrow X_{n,\text{ét}}$ is the canonical projection to the étale site (or sometimes to the Nisnevich site)

For X a fine and saturated log-smooth log-scheme over \mathcal{O}_K and $0 \leq r \leq p-2$, the natural map of complexes of sheaves on the étale site of X_0

$$\tau_{\leq r}\mathcal{S}_n(r) \rightarrow \mathcal{S}_n(r)$$

is a quasi-isomorphism. For X semistable over \mathcal{O}_K and $r \geq 0$, the natural map of complexes of sheaves on the étale site of X_0

$$\tau_{\leq r}\mathcal{S}'_n(r) \rightarrow \mathcal{S}'_n(r)$$

is a p^{Nr} -quasi-isomorphism for a universal constant N [6, Prop. 3.12].

2.1.1. Syntomic cohomology and differential forms. Let X be a syntomic scheme over $W(k)$. Recall the differential definition [23] of syntomic cohomology. Assume first that we have an immersion $\iota : X \hookrightarrow Z$ over $W(k)$ such that Z is a smooth $W(k)$ -scheme endowed with a compatible system of liftings of the Frobenius $\{F_n : Z_n \rightarrow Z_n\}$. Let $D_n = D_{X_n}(Z_n)$ be the PD-envelope of X_n in Z_n (compatible with the canonical PD-structure on $pW_n(k)$) and J_{D_n} the ideal of X_n in D_n . Set $J_{D_n}^{<r>} := \{a \in J_{D_{n+s}}^{[r]} \mid \varphi(a) \in p^r\mathcal{O}_{D_{n+s}}\}/p^n$ for some $s \geq r$. For $0 \leq r \leq p-1$, $J_{D_n}^{<r>} = J_{D_n}^{[r]}$. This definition is independent of s . Consider the following complexes of sheaves on $X_{\text{ét}}$.

$$(2.2) \quad \begin{aligned} \mathcal{S}_n(r)_{X,Z} &:= \text{Cone}(J_{D_n}^{<r-\bullet>} \otimes \Omega_{Z_n}^\bullet \xrightarrow{1-\varphi_r} \mathcal{O}_{D_n} \otimes \Omega_{Z_n}^\bullet)[-1], \\ \mathcal{S}'_n(r)_{X,Z} &:= \text{Cone}(J_{D_n}^{[r-\bullet]} \otimes \Omega_{Z_n}^\bullet \xrightarrow{p^{r+i}-p^i\varphi} \mathcal{O}_{D_n} \otimes \Omega_{Z_n}^\bullet)[-1], \end{aligned}$$

where $\Omega_{Z_n}^\bullet := \Omega_{Z_n/W_n(k)}^\bullet$ and φ_r is " $\varphi/p^{r\text{"}}$ (see [39, 2.1] for details). The complexes $\mathcal{S}_n(r)_{X,Z}$, $\mathcal{S}'_n(r)_{X,Z}$ are, up to canonical quasi-isomorphisms, independent of the choice of ι and $\{F_n\}$ (and we will omit the subscript Z from the notation). Again, the natural maps $\omega : \mathcal{S}_n^i(r)_X \rightarrow \mathcal{S}_n(r)_X$ and $\tau : \mathcal{S}_n(r)_X \rightarrow \mathcal{S}'_n(r)_X$ have kernels and cokernels annihilated by p^{r+i} .

In general, immersions as above exist étale locally, and we define $\mathcal{S}_n(r)_X \in \mathbf{D}^+(X_{\text{ét}}, \mathbf{Z}/p^n)$ by gluing the local complexes, and $\mathcal{S}_n(r)_{X_{\mathcal{O}_{\overline{K}}}} \in \mathbf{D}^+((X_{\mathcal{O}_{\overline{K}}})_{\text{ét}}, \mathbf{Z}/p^n)$ as the inductive limit of $\mathcal{S}_n(r)_{X_{\mathcal{O}_{K'}}}$, where

\mathcal{O}_K' varies over the integral closures of \mathcal{O}_K in all finite extensions of K in \overline{K} . Similarly, we define $S_n^i(r)_X$ and $S_n^i(r)_{X_{\mathcal{O}_{\overline{K}}}}$.

Let now X be a log-syntomic scheme over $W(k)$. Using log-crystalline cohomology, the above construction of syntomic complexes goes through almost verbatim (see [39, 2.1] for details) to yield the logarithmic analogs $S_n(r)$ and $S_n^i(r)$ on $X_{\text{ét}}$. There are natural maps

$$\varepsilon : H^i(X_{\text{ét}}, S_n(r)) \rightarrow H^i(X_{\text{ét}}^\times, S_n(r)), \quad \varepsilon : H^i(X_{\text{ét}}, S_n^i(r)) \rightarrow H^i(X_{\text{ét}}^\times, S_n^i(r)),$$

where, for clarity, we wrote X^\times for the log-scheme X with its full log-structure. In this paper we are often interested in log-schemes coming from a regular syntomic scheme X over $W(k)$ and a relative simple (i.e., with no self-intersections) normal crossing divisor D on X . In such cases we will write $S_n(r)_X(D)$ and $S_n^i(r)_X(D)$ for the syntomic complexes and use the Nisnevich topology instead of the étale one. We will write $H^*(X, S_n(*)_X(D))$ and $H^*(X, S_n^i(*)_X(D))$ for the corresponding cohomology groups. We will employ the same convention while talking about log-étale cohomology $H^*(X_K^\times, \mathbf{Z}/p^n(*))$: we will write $H^*(X_K(D_K), \mathbf{Z}/p^n(*))$ instead.

2.1.2. Products. We need to discuss products. Assume that we are in the lifted situation (2.2). Then we have a product structure

$$\cup : S_n^i(r)_{X,Z} \otimes S_n^j(r')_{X,Z} \rightarrow S_n^{i+j}(r+r')_{X,Z}, \quad r, r', i, j \geq 0,$$

defined by the following formulas

$$\begin{aligned} (x, y) \otimes (x', y') &\mapsto (xx', (-1)^a p^{r+i} xy' + yp^j \varphi(x')) \\ (x, y) \in S_n^i(r)_{X,Z}^a &= (J_{D_n}^{[r-a]} \otimes \Omega_{Z_n}^a) \oplus (\mathcal{O}_{D_n} \otimes \Omega_{Z_n}^{a-1}), \\ (x', y') \in S_n^j(r')_{X,Z}^b &= (J_{D_n}^{[r'-b]} \otimes \Omega_{Z_n}^b) \oplus (\mathcal{O}_{D_n} \otimes \Omega_{Z_n}^{b-1}). \end{aligned}$$

Globalizing, we obtain the product structure

$$\cup : S_n^i(r)_X \otimes^{\mathbb{L}} S_n^j(r')_X \rightarrow S_n^{i+j}(r+r')_X, \quad r, r', i, j \geq 0.$$

This product is clearly compatible with the crystalline product.

Similarly, we have the product structures

$$\cup : S_n(r)_{X,Z} \otimes S_n(r')_{X,Z} \rightarrow S_n(r+r')_{X,Z}, \quad r, r' \geq 0,$$

defined by the formulas

$$\begin{aligned} (x, y) \otimes (x', y') &\mapsto (xx', (-1)^a xy' + y\varphi_{r'}(x')) \\ (x, y) \in S_n(r)_{X,Z}^a &= (J_{D_n}^{<r-a>} \otimes \Omega_{Z_n}^a) \oplus (\mathcal{O}_{D_n} \otimes \Omega_{Z_n}^{a-1}), \\ (x', y') \in S_n(r')_{X,Z}^b &= (J_{D_n}^{<r'-b>} \otimes \Omega_{Z_n}^b) \oplus (\mathcal{O}_{D_n} \otimes \Omega_{Z_n}^{b-1}). \end{aligned}$$

Globalizing, we obtain the product structure

$$\cup : S_n(r)_X \otimes^{\mathbb{L}} S_n(r')_X \rightarrow S_n(r+r')_X, \quad r, r' \geq 0.$$

This product is also clearly compatible with the crystalline product.

The above product structures are compatible with the maps ω and the maps ω_0 . On the other hand the maps τ are, in general, not compatible with products.

2.1.3. Syntomic symbol maps. Let X be a regular syntomic scheme over $W(k)$ with a divisor D with relative simple normal crossings. Recall that there are first Chern class maps defined by Kato and Tsuji [40, 2.2]

$$\begin{aligned} c_1^{\text{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] &\rightarrow i_* j_* \mathcal{O}_{(X \setminus D)_{n+1}}^*[-1] \rightarrow S_n(1)_X(D), \\ c_1^{\text{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] &\rightarrow i_* j_* \mathcal{O}_{(X \setminus D)_n}^*[-1] \rightarrow S'_n(1)_X(D), \end{aligned}$$

that are compatible, i.e., the following diagram commutes

$$\begin{array}{ccc} j_*\mathcal{O}_{X \setminus D}^*[-1] & \xrightarrow{c_1^{\text{syn}}} & S'_n(1)_X(D) \\ \downarrow pc_1^{\text{syn}} & \swarrow \omega & \\ S_n(1)_X(D) & & \end{array}$$

Here $j : X \setminus D \hookrightarrow X$ is the natural immersion. In the lifted situation these classes are defined in the following way. Let C_n be the complex

$$(1 + J_{D_n} \rightarrow M_{D_n}^{\text{gp}}) \simeq j_*\mathcal{O}_{(X \setminus D)_n}^*[-1],$$

where, for a log-scheme X , M_X denotes its log-structure. The Chern class maps

$$(2.3) \quad c_1^{\text{syn}} : j_*\mathcal{O}_{(X \setminus D)_n}^*[-1] \rightarrow S'_n(1)_X(D), \quad c_1^{\text{syn}} : j_*\mathcal{O}_{(X \setminus D)_{n+1}}^*[-1] \rightarrow S_n(1)_X(D),$$

are defined by the morphisms of complexes

$$C_n \rightarrow S'_n(1)_{X,Z}, \quad C_{n+1} \rightarrow S_n(1)_{X,Z}$$

given by the formulas

$$\begin{aligned} 1 + J_{D_n} &\rightarrow (S'_n(1)_{X,Z})^0 = J_{D_n}; & a &\mapsto \log a; \\ 1 + J_{D_{n+1}} &\rightarrow (S_n(1)_{X,Z})^0 = J_{D_n}; & a &\mapsto \log a \pmod{p^n}; \end{aligned}$$

and

$$\begin{aligned} M_{D_n}^{\text{gp}} &\rightarrow (S'_n(1)_{X,Z})^1 = (\mathcal{O}_{D_n} \otimes \Omega_{Z_n}^1) \oplus \mathcal{O}_{D_n}; & b &\mapsto (d \log b, \log(b^p \varphi_{D_n}(b)^{-1})); \\ M_{D_{n+1}}^{\text{gp}} &\rightarrow (S_n(1)_{X,Z})^1 = (\mathcal{O}_{D_n} \otimes \Omega_{Z_n}^1) \oplus \mathcal{O}_{D_n}; & b &\mapsto (d \log b \pmod{p^n}, p^{-1} \log(b^p \varphi_{D_{n+1}}(b)^{-1})). \end{aligned}$$

2.1.4. Syntomic cohomology and p -adic nearby cycles. For log-schemes over \mathcal{O}_K^\times , in a stable range, syntomic cohomology tends to compute (via the period morphism) p -adic nearby cycles. We will recall the relevant theorems.

Let X be a log-syntomic scheme over $W(k)$. Let $i : X_{0,\text{ét}} \rightarrow X_{\text{ét}}$ and $j : X_{\text{tr},K,\text{ét}} \rightarrow X_{\text{ét}}$ be the natural maps. Here X_{tr} is the open set of X where the log-structure is trivial. For $0 \leq r \leq p-2$, there is a natural homomorphism on the étale site of X_n [8] (the Fontaine-Messing period map)

$$\alpha_r : \mathcal{S}_n(r) \rightarrow i^*Rj_*\mathbf{Z}/p^n(r)$$

from syntomic complexes to p -adic nearby cycles. It factors through $\tau_{\leq r}i^*Rj_*\mathbf{Z}/p^n(r)$. One checks that α_r is compatible with products. Similarly, for any $r \geq 0$, we get a natural map [8]

$$\tilde{\alpha}_r : \mathcal{S}_n(r) \rightarrow i^*Rj_*\mathbf{Z}/p^n(r)'$$

Composing with the map $\omega : \mathcal{S}'_n(r) \rightarrow \mathcal{S}_n(r)$ we get a natural, compatible with products, morphism

$$\alpha_r : \mathcal{S}'_n(r) \rightarrow i^*Rj_*\mathbf{Z}/p^n(r)'$$

Theorem 2.1. ([40, Theorem 5.1]) *For $i \leq r \leq p-2$ and for a fine and saturated log-scheme X log-smooth over \mathcal{O}_K^\times the period map*

$$(2.4) \quad \alpha_r : \mathcal{S}_n(r)_X \xrightarrow{\sim} \tau_{\leq r}i^*Rj_*\mathbf{Z}/p^n(r)_{X_{\text{tr}}}$$

is an isomorphism.

Theorem 2.2. ([6, Theorem 1.1]) *For $0 \leq i \leq r$ and for a semistable scheme X over \mathcal{O}_K , consider the period map*

$$(2.5) \quad \alpha_r : \mathcal{H}^i(\mathcal{S}'_n(r)_X) \rightarrow i^*R^i j_*\mathbf{Z}/p^n(r)'_{X_{\text{tr}}}.$$

If K has enough roots of unity⁴ then the kernel and cokernel of this map are annihilated by p^{Nr} for a universal constant N depending only on p (and d if $p = 2$) (not depending on X , K , n or r). In general, the kernel and cokernel of this map are annihilated by p^{Nr} for an integer $N = N(p, e)$, which depends on e and p (also d if $p = 2$) but not on X or n .

2.2. Syntomic-étale cohomology. We will now recall the definition and basic properties of syntomic-étale cohomology. More details can be found in [8],[7]. Let X be a log-scheme, log-syntomic over $\text{Spec}(W(k))$. Let $j' : X_{\text{tr},K} \rightarrow X_K$ denote the natural open immersion.

2.2.1. *Syntomic-étale cohomology.* Denote by $\mathcal{E}_n(r)$ and $\mathcal{E}'_n(r)$ the syntomic-étale complexes on $X_{\text{ét}}$ [7, 2.2.2] associated to $\mathcal{S}_n(r)$ and $\mathcal{S}'_n(r)$, respectively. They are obtained by gluing the complexes of sheaves $\mathcal{S}_n(r)$ and $\mathcal{S}'_n(r)$ and the complexes of sheaves $j'_*G\mathbf{Z}/p^n(r)'$, where G denotes the Godement resolution of a sheaf (or a complex of sheaves), by the maps $\tilde{\alpha}_r$ and α_r . We have the distinguished triangles

$$(2.6) \quad j_{\text{ét}}!Rj'_*\mathbf{Z}/p^n(r)' \rightarrow \mathcal{E}_n(r) \rightarrow i_*\mathcal{S}_n(r), \quad j_{\text{ét}}!Rj'_*\mathbf{Z}/p^n(r)' \rightarrow \mathcal{E}'_n(r) \rightarrow i_*\mathcal{S}'_n(r),$$

as well as the natural maps

$$\tilde{\alpha}_r : \mathcal{E}_n(r) \rightarrow Rj_*\mathbf{Z}/p^n(r)', \quad \alpha_r : \mathcal{E}_n(r)' \rightarrow Rj_*\mathbf{Z}/p^n(r)'$$

compatible with the maps $\tilde{\alpha}_r$ and α_r from syntomic complexes. For $a \geq 0$, we have the truncated version of the above - the distinguished triangles

$$(2.7) \quad j_{\text{ét}}!\tau_{\leq a}Rj'_*\mathbf{Z}/p^n(r)' \rightarrow \tau_{\leq a}\mathcal{E}_n(r) \rightarrow i_*\tau_{\leq a}\mathcal{S}_n(r), \quad j_{\text{ét}}!\tau_{\leq a}Rj'_*\mathbf{Z}/p^n(r)' \rightarrow \tau_{\leq a}\mathcal{E}'_n(r) \rightarrow i_*\tau_{\leq a}\mathcal{S}'_n(r).$$

2.2.2. *Syntomic-étale cohomology and étale cohomology of the generic fiber.* For a log-scheme over \mathcal{O}_K^\times , in a stable range, syntomic-étale cohomology tends to compute étale cohomology of the generic fiber.

Theorem 2.3. ([7, Theorem 2.5]) *Let X be a log-scheme log-smooth over \mathcal{O}_K^\times . Then*

- (1) *we have a natural quasi-isomorphism*

$$\tilde{\alpha}_r : \tau_{\leq r}\mathcal{E}_n(r) \simeq \tau_{\leq r}Rj_*\mathbf{Z}/p^n(r), \quad 0 \leq r \leq p-2.$$

- (2) *if X is semistable, there is a constant N as in Theorem 2.2 and a natural morphism*

$$\alpha_r : \mathcal{E}'_n(r) \rightarrow Rj_*\mathbf{Z}/p^n(r)', \quad r \geq 0,$$

such that the induced map on cohomology sheaves in degrees $\leq r$ has kernel and cokernel annihilated by p^{Nr} .

The above theorem implies that the logarithmic syntomic-étale cohomology is close to the logarithmic syntomic-étale cohomology of the complement of the divisor at infinity.

Corollary 2.4. ([7, Cor. 2.6]) *Let X be a semistable scheme over \mathcal{O}_K with a divisor at infinity D_∞ . We treat it as a log-scheme over \mathcal{O}_K^\times . Let $Y := X \setminus D_\infty$ and let $j_1 : Y \hookrightarrow X$.*

- (1) *we have a natural quasi-isomorphism*

$$\tilde{\alpha}_r : \tau_{\leq r}\mathcal{E}_n(r)_X \xrightarrow{\sim} \tau_{\leq r}Rj_{1*}\mathcal{E}_n(r)_Y, \quad 0 \leq r \leq p-2.$$

- (2) *there is a constant N as in Theorem 2.2 and a natural morphism*

$$\alpha_r : \mathcal{E}'_n(r)_X \rightarrow Rj_{1*}\mathcal{E}'_n(r)_Y, \quad r \geq 0,$$

such that the induced map on cohomology sheaves in degrees $\leq r$ has kernel and cokernel annihilated by p^{Nr} .

⁴See Section (2.1.1) of [6] for what it means for a field to contain enough roots of unity. The field F contains enough roots of unity and for any K , the field $K(\zeta_{p^n})$, for $n \geq c(K) + 3$, where $c(K)$ is the conductor of K , contains enough roots of unity.

2.2.3. *Nisnevich syntomic-étale cohomology.* We will pass now to the Nisnevich topos of X . Denote by $\varepsilon : X_{\text{ét}} \rightarrow X_{\text{Nis}}$ the natural projection. For $r \geq 0$, by applying $\mathbf{R}\varepsilon_*$ to the étale period map above and using that $\mathbf{R}\varepsilon_* i^* = i^* \mathbf{R}\varepsilon_*^5$ (cf. [10, 2.2.b]), we obtain a natural map

$$\tilde{\alpha}_r : \mathbf{R}\varepsilon_* \mathcal{S}_n(r) \rightarrow i^* \mathbf{R}j_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)'.$$

Composing with the map $\omega : \mathbf{R}\varepsilon_* \mathcal{S}'_n(r) \rightarrow \mathbf{R}\varepsilon_* \mathcal{S}_n(r)$ we get a natural, compatible with products, morphism

$$\alpha_r : \mathbf{R}\varepsilon_* \mathcal{S}'_n(r) \rightarrow i^* \mathbf{R}j_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)'.$$

Write, for simplicity, $\mathcal{S}_n(r)$ and $\mathcal{S}'_n(r)$ for the derived pushforwards of $\mathcal{S}_n(r)$ and $\mathcal{S}'_n(r)$ from $X_{\text{ét}}$ to X_{Nis} . Same for $\mathcal{E}_n(r)$ and $\mathcal{E}'_n(r)$. Notice that the latter are quasi-isomorphic to the complexes obtained by gluing the complexes of sheaves $\mathcal{S}_n(r)$ and $\mathcal{S}'_n(r)$ on $X_{1, \text{Nis}}$ and the complexes of sheaves $\varepsilon_* j'_* G\mathbf{Z}/p^n(r)'$ on $X_{K, \text{Nis}}$ by the maps $\tilde{\alpha}_r$ and α_r . Hence we have the distinguished triangles

$$(2.8) \quad j_{\text{Nis}!} \mathbf{R}j'_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)' \rightarrow \mathcal{E}_n(r) \rightarrow i_* \mathcal{S}_n(r), \quad j_{\text{Nis}!} \mathbf{R}j'_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)' \rightarrow \mathcal{E}'_n(r) \rightarrow i_* \mathcal{S}'_n(r),$$

as well as the natural maps

$$\tilde{\alpha}_r : \mathcal{E}_n(r) \rightarrow \mathbf{R}j_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)', \quad \alpha_r : \mathcal{E}_n(r)' \rightarrow \mathbf{R}j_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)'$$

compatible with the maps $\tilde{\alpha}_r$ and α_r from syntomic complexes. For $a \geq 0$, we have the truncated version of the above - the distinguished triangles

$$(2.9) \quad j_{\text{Nis}!} \tau_{\leq a} \mathbf{R}j'_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)' \rightarrow \tau_{\leq a} \mathcal{E}_n(r) \rightarrow i_* \tau_{\leq a} \mathcal{S}_n(r), \quad j_{\text{ét}!} \tau_{\leq a} \mathbf{R}j'_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(r)' \rightarrow \tau_{\leq a} \mathcal{E}'_n(r) \rightarrow i_* \tau_{\leq a} \mathcal{S}'_n(r).$$

Define the following complexes of sheaves on X_{Nis}

$$\begin{aligned} \mathcal{S}_n(r)_{\text{Nis}} &:= \tau_{\leq r} \mathcal{S}_n(r), & \mathcal{S}'_n(r)_{\text{Nis}} &:= \tau_{\leq r} \mathcal{S}'_n(r); \\ \mathcal{E}_n(r)_{\text{Nis}} &:= \tau_{\leq r} \mathcal{E}_n(r), & \mathcal{E}'_n(r)_{\text{Nis}} &:= \tau_{\leq r} \mathcal{E}'_n(r). \end{aligned}$$

We will need the following twisted version of the complexes $\mathcal{E}'_n(r)$ and $\mathcal{E}'_n(r)_{\text{Nis}}$. Set the gluing map (on $X_{\text{ét}}$)

$$\alpha_r : \mathcal{S}_n^1(r) \rightarrow i^* \mathbf{R}j_* \mathbf{Z}/p^n(r+1)'(-1)$$

to be equal to the composition

$$\mathcal{S}_n^1(r) \xrightarrow{\omega^0} \mathcal{S}'_n(r) \xrightarrow{\alpha_r} i^* \mathbf{R}j_* \mathbf{Z}/p^n(r)' \rightarrow i^* \mathbf{R}j_* \mathbf{Z}/p^n(r+1)'(-1).$$

And define the complexes $\mathcal{E}_n^1(r)$ by gluing $\mathcal{S}_n^1(r)$ and $j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r+1)'(-1)$ via α_r . We have the distinguished triangle

$$j_{\text{ét}!} \mathbf{R}j'_* \mathbf{Z}/p^n(r+1)'(-1) \rightarrow \mathcal{E}_n^1(r) \rightarrow i_* \mathcal{S}_n^1(r).$$

Write $\mathcal{E}_n^1(r)$ for the derived pushforward of $\mathcal{E}_n^1(r)$ to X_{Nis} and set $\mathcal{E}_n^1(r)_{\text{Nis}} := \tau_{\leq r} \mathcal{E}_n^1(r)$.

2.2.4. *Syntomic-étale cohomology and differential forms.* We will need a differential definition of the syntomic-étale complexes. Let X be a log-syntomic, finite and saturated (fs for short) scheme over $W(k)$. Assume first that X is affine and we have an immersion $X \hookrightarrow Z$ over $W(k)$ such that Z is a log-smooth $W(k)$ -scheme endowed with a compatible system of liftings of the Frobenius $\{F_n : Z_n \rightarrow Z_n\}$. Choose sufficiently large algebraically closed fields Ω and Ω' of characteristic zero and p , respectively. Let C be the set of all isomorphism classes of fs monoids P such that $P^* = \{1\}$. For each isomorphism class $c \in C$ choose a representative P_c of c and define the log-geometric point Ω_c to be $\text{Spec}(\Omega_c)$ with $M_{\Omega_c} = \Omega \oplus_{n \in \mathbb{N}, n \neq 0} 1/n P_c$ and Ω'_c to be $\text{Spec}(\Omega')$ with $M_{\Omega'_c} = \Omega \oplus_{n \in \mathbb{N}, p \nmid n} 1/n P_c$. Let G denote the Godement resolution with respect to all log-geometric points whose sources are Ω_c or Ω'_c for some $c \in C$. The set of points we have chosen is actually larger than what would suffice here but it will turn out useful later.

The map

$$\alpha'_r : \mathcal{S}_n(r) \rightarrow i_* i^* \mathbf{R}j_* \mathbf{Z}/p^n(r)' \xrightarrow{\delta} j_{\text{ét}!} \mathbf{R}j'_* \mathbf{Z}/p^n(r)'[1],$$

⁵This equality fails for the projection to Zariski topology and is the reason we use Nisnevich topology instead of Zariski.

where the map δ is obtained from the distinguished triangle

$$j_{\text{ét}}! \mathbf{R}j_* \mathbf{Z}/p^n(r)' \xrightarrow{\eta} \mathbf{R}j_* \mathbf{Z}/p^n(r)' \rightarrow i_* i^* \mathbf{R}j_* \mathbf{Z}/p^n(r)',$$

can be induced by a map $\alpha'_r : S_n(r) \rightarrow j_{\text{ét}}! j'_* G\mathbf{Z}/p^n(r)'[1]$ that, in turn, can be obtained by shifting the following morphisms of presheaves on $X_{\text{ét}}$ [39, 3.1] (we write θ for the operation $i_* i^* j_* G$ and assume that $U = \text{Spec}(A) \rightarrow X$ is a strict étale map)

$$\begin{aligned} \alpha'_{r,U} : \Gamma(U, S_n(r)_{X,Z}) &\rightarrow \Gamma(U^h, \theta_U \bar{S}_n(r)_{U,Z}) \xleftarrow{\sim} \Gamma(U^h, \theta_U \Lambda_{U^h}) \xleftarrow{\sim} \Gamma(U, \theta \Lambda_U) \\ &\xleftarrow{\sim} \Gamma(U, \text{Cone}(\eta(G\Lambda_U))) \rightarrow \Gamma(U, j_{\text{ét}}! j'_* G\Lambda_U[1]), \end{aligned}$$

where we put $\Lambda_Y = \mathbf{Z}/p^n(r)'_{Y_{K,\text{tr}}}$ and $\eta(G\Lambda_U) : j_{\text{ét}}! j'_* G\Lambda_U \hookrightarrow j_{\text{ét}*} j'_* G\Lambda_U$ is the natural injection. Here U^h denotes the henselization of U with respect to the ideal pU and $\bar{S}_n(r)_{U,Z}$ is an analog of the syntomic complex $S_n(r)_{U,Z}$ [39, 3.1]. This complex is defined using, instead of A , \bar{A}^h - the integral closure of A^h in an algebraic closure of the fraction field $\text{Frac}(A^h)$ that is étale over $A_{\text{tr},K}^h$, and instead of the immersion $X \hookrightarrow Z$, the immersion $\text{Spec}(\bar{A}^h) \hookrightarrow \text{Spec}(\mathbf{A}_{\text{cr}}(\bar{A}^h))$ with the natural Frobenius liftings and log-structures. Unlike in [39, 3.1] we allow here trivial log-structure on the special fiber. The complex $\bar{S}_n(r)_{U,Z}$ is a complex of locally free sheaves on $(U_{\text{tr},K}^h)_{\text{ét}}$. It is a resolution of $\Lambda_{U_{\text{tr},K}^h}$.

In the case of a general X we choose a strict étale affine covering $X' \rightarrow X$ and immersion $X' \hookrightarrow Z'$ with Frobenius liftings $F_{Z'_n} : Z'_n \rightarrow Z'_n$. From this we construct a (Čech) hypercovering $X^\bullet \rightarrow X$ with immersion $X^\bullet \hookrightarrow Z^\bullet$ and Frobenius liftings $F_{Z_n^\bullet} : Z_n^\bullet \rightarrow Z_n^\bullet$. By applying the above construction to each level we obtain a morphism $\alpha'_r : S_n(r)_{X^\bullet, Z^\bullet} \rightarrow j_{\text{ét}}! j'_* G\Lambda_{X^\bullet}[1]$. Our map α'_r is now defined as $\mathbf{R}\varepsilon_* \alpha'_r$, where $\varepsilon : X_{\text{ét}}^\bullet \rightarrow X_{\text{ét}}$ is the change of topoi map. It can be represented by the composition

$$\alpha'_r : S_n(r)_X \rightarrow \varepsilon_* G S_n(r)_{X^\bullet, Z^\bullet} \xrightarrow{\alpha'_r} \varepsilon_* G j_{\text{ét}}! j'_* G\Lambda_{X^\bullet}[1] \xleftarrow{\sim} j_{\text{ét}}! j'_* G\Lambda_X[1].$$

Consider the induced map

$$\alpha'_r : S'_n(r)_X \xrightarrow{\omega} S_n(r)_X \xrightarrow{\alpha'_r} j_{\text{ét}}! j'_* G\Lambda_X[1].$$

This map, a priori a zigzag of maps of complexes, can be "straighten up", that is, we can find a complex $\tilde{S}'_n(r)_X$ and genuine maps of complexes $f : \tilde{S}'_n(r)_X \rightarrow S'_n(r)_X$ and $\tilde{\alpha}_r : \tilde{S}'_n(r)_X \rightarrow j_{\text{ét}}! j'_* G\Lambda_X[1]$ that fit into the following commutative diagram

$$\begin{array}{ccc} \tilde{S}'_n(r)_X & & \\ \downarrow \wr f & \searrow \tilde{\alpha}_r & \\ S'_n(r)_X & \xrightarrow{\alpha'_r} & j_{\text{ét}}! j'_* G\Lambda_X[1] \end{array}$$

This is done by replacing each diagram of maps $A \xrightarrow{a} B \xleftarrow{c} C$ by the map $a' : D \rightarrow C$ from the following commutative diagram of complexes

$$(2.10) \quad \begin{array}{ccc} D = \text{Cone}(a - b)[-1] & \xrightarrow{a'} & C \\ \sim \downarrow b' & & \sim \downarrow b \\ A & \xrightarrow{a} & B \end{array}$$

Set

$$E'_n(r)_X := \text{Cone}(\tilde{S}'_n(r)_X \xrightarrow{\tilde{\alpha}_r} j_{\text{ét}}! j'_* G\Lambda_X[1])[-1], \quad E'_n(r)_{\text{Nis},X} := \tau_{\leq r} \varepsilon'_* G E'_n(r)_X,$$

where $\varepsilon' : X_{\text{ét}} \rightarrow X_{\text{Nis}}$ is the change of topology map. These complexes represent $\mathcal{E}'_n(r)_X$ and $\mathcal{E}'_n(r)_{\text{Nis},X}$, respectively. They are functorial in X . The complexes $E'_n(r)_X$ and $E'_n(r)_{\text{Nis},X}$ inherit functorial product structure that is compatible with the maps

$$E'_n(r)_X \rightarrow \tilde{S}'_n(r)_X \rightarrow S'_n(r)_X.$$

To construct it note that all the maps in $\alpha_{r,U}$ are compatible with products and so are the maps ω and δ . It suffices thus, in the diagram 2.10, to equip the cone D with product structure that is compatible with the projections on A and C . But this is standard and can be done as in [27, 3.1].

The above syntomic-étale complexes have twisted versions. Consider first

$$E_n^1(r)_X := \text{Cone}(\tilde{S}_n^1(r)_X \xrightarrow{\tilde{\omega}_0} \tilde{S}'_n(r)_X \xrightarrow{\tilde{\alpha}'_r} j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r+1)'(-1)[1])[-1], \quad E_n^1(r)_{\text{Nis},X} := \tau_{\leq r} \varepsilon'_* G E_n^1(r)_X,$$

where

$$\tilde{\alpha}'_r : \tilde{S}'_n(r)_X \xrightarrow{\tilde{\alpha}'_r} j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r)'[1] \rightarrow j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r+1)'(-1)[1]$$

and the complex $\tilde{S}_n^1(r)_X$ is defined via the following commutative diagram

$$\begin{array}{ccc} \tilde{S}_n^1(r)_X = \text{Cone}(f - \omega_0)[-1] & \xrightarrow[\sim]{f^1} & S_n^1(r)_X \\ \downarrow \tilde{\omega}_0 & & \downarrow \omega_0 \\ \tilde{S}'_n(r)_X & \xrightarrow[\sim]{f} & S'_n(r)_X \end{array}$$

We also the following twisted version

$$E_n^2(r)_X := \text{Cone}(\tilde{S}_n^2(r)_X \xrightarrow{\tilde{\omega}_{1,0}} \tilde{S}'_n(r)_X \xrightarrow{\tilde{\alpha}'_r} j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r+2)'(-2)[1])[-1],$$

where

$$\tilde{\alpha}'_r : \tilde{S}'_n(r)_X \xrightarrow{\tilde{\alpha}'_r} j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r)'[1] \rightarrow j_{\text{ét}!} j'_* G\mathbf{Z}/p^n(r+2)'(-2)[1],$$

$\omega_{1,0} = \omega_1 \omega_0$, and the complex $\tilde{S}_n^2(r)_X$ is defined via the following commutative diagram

$$\begin{array}{ccc} \tilde{S}_n^2(r)_X = \text{Cone}(f - \omega_{1,0})[-1] & \xrightarrow[\sim]{f^1} & S_n^2(r)_X \\ \downarrow \tilde{\omega}_{1,0} & & \downarrow \omega_{1,0} \\ \tilde{S}'_n(r)_X & \xrightarrow[\sim]{f} & S'_n(r)_X \end{array}$$

Using again the construction of products on cones from [27, 3.1] and the fact that the maps

$$\omega : S'_n(r)_X \rightarrow S_n(r)_X, \quad \omega_0 : S_n^1(r_1)_X \rightarrow S'_n(r)_X$$

as well as the map α_r commute with products, we construct products

$$\cup : E'_n(r_1)_X \otimes^{\mathbb{L}} E_n^1(r_2)_X \rightarrow E_n^1(r_1 + r_2)_X; \quad \cup : E'_n(r_1)_{\text{Nis},X} \otimes^{\mathbb{L}} E_n^1(r_2)_{\text{Nis},X} \rightarrow E_n^1(r_1 + r_2)_{\text{Nis},X}$$

that are compatible with the product $\cup : S'_n(r_1)_X \otimes^{\mathbb{L}} S_n^1(r_2)_X \rightarrow S_n^1(r_1 + r_2)_X$.

In an analogous way we define the complexes $E_n(r)_X$ and $E_n(r)_{\text{Nis},X}$ and the corresponding products.

2.2.5. Syntomic-étale symbol maps. If X is fs then we have Chern class maps

$$(2.11) \quad c_1^{\text{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] \rightarrow E'_n(1)_{\text{Nis},X}(D), \quad c_1^{\text{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] \rightarrow E_n(1)_{\text{Nis},X}(D)$$

that are compatible with the syntomic Chern class maps. To define these maps it suffices, by degree reason, to define maps

$$(2.12) \quad c_1^{\text{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] \rightarrow E'_n(1)_X(D), \quad c_1^{\text{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] \rightarrow E_n(1)_X(D).$$

For that recall that the syntomic Chern classes are compatible with the étale Chern classes [39, Prop. 3.2.4], i.e., that the following diagram commutes

$$\begin{array}{ccc} j_* \mathcal{O}_{X \setminus D}^*[-1] & \xrightarrow{c_1^{\text{syn}}} & S'_n(1)_X(D) \\ & \searrow pc_1 & \downarrow \alpha_1 \\ & & i_* i^* Rj_{\text{ét}*} Rj'_* \mathbf{Z}/p^n(1)' \end{array}$$

where the map c_1 is induced from the Chern class map

$$c_1^{\acute{e}t} : j_* \mathcal{O}_{X \setminus D}^*[-1] \rightarrow \mathrm{R}j_{\acute{e}t}^* \mathrm{R}j'_* \mathbf{Z}/p^n(1).$$

It follows that the composition

$$j_* \mathcal{O}_{X \setminus D}^*[-1] \xrightarrow{c_1^{\mathrm{syn}}} S'_n(1)_X(D) \xrightarrow{\alpha_1} i_* i^* \mathrm{R}j_{\acute{e}t}^* \mathrm{R}j'_* \mathbf{Z}/p^n(1)' \rightarrow j_{\acute{e}t}^* \mathrm{R}j'_* \mathbf{Z}/p^n(1)'[1]$$

is trivial, hence there exists a unique (for degree reason) map

$$c_1^{\mathrm{syn}} : j_* \mathcal{O}_{X \setminus D}^*[-1] \rightarrow E'_n(1)_X(D)$$

compatible with the syntomic Chern class map. The construction for the complex $E_n(1)_X(D)$ is analogous.

2.3. Syntomic cohomology and motivic cohomology. Let X be a smooth scheme over \mathcal{O}_K . Let $\mathbf{Z}(r)_M$ denote the complex of motivic sheaves $\mathbf{Z}(r)_M := X \mapsto z^r(X, 2r - *)$ in the Nisnevich topology of X , where the complex $z^r(X, *)$ is the Bloch's cycles complex [2]. Let $\mathbf{Z}/p^n(r)_M := \mathbf{Z}(r)_M \otimes \mathbf{Z}/p^n$. We have that $H^j(X_{\mathrm{Nis}}, \mathbf{Z}/p^n(i)_M) = H^j \Gamma(X, \mathbf{Z}/p^n(r)_M)$ is the Bloch higher Chow group [10, Prop. 3.6].

Recall that the main theorem of [7] shows that syntomic-étale complexes on smooth schemes over \mathcal{O}_K approximate motivic complexes.

Theorem 2.5. ([7, Theorem 3.10]) *Let X be a semistable scheme over \mathcal{O}_K with a smooth special fiber. Let $j' : X_{\mathrm{tr}} \hookrightarrow X$ be the natural open immersion. Then*

- (1) *there is a natural cycle class map*

$$\mathrm{cl}_r^{\mathrm{syn}} : \mathrm{R}j'_* \mathbf{Z}/p^n(r)_M \rightarrow \mathcal{E}_n(r)_{\mathrm{Nis}}, \quad 0 \leq r \leq p - 2.$$

It is a quasi-isomorphism.

- (2) *there is a natural cycle class map*

$$\mathrm{cl}_r^{\mathrm{syn}} : \mathrm{R}j'_* \mathbf{Z}/p^n(r)_M \rightarrow \mathcal{E}'_n(r)_{\mathrm{Nis}}, \quad r \geq 0.$$

It is a p^{Nr} -quasi-isomorphism for a constant N as in Theorem 2.2.

We have analogous statements in the étale topology. These cycle class maps are compatible (via the localization map and the period map) with the étale cycle class maps.

3. COHOMOLOGY OF CLASSIFYING SPACES

In this section we study cohomology of classifying spaces: we compute their (Nisnevich) syntomic cohomology and we show that their Nisnevich syntomic-étale and syntomic cohomologies agree in even degrees.

3.1. Classical computations. We are interested in cohomology of BGL_a and the related classifying simplicial schemes $B(GL_a \times GL_b)$ and $BGL(a, b)$. Recall that, roughly speaking, if a $\mathbf{Z} \times \mathbf{Z}$ -graded cohomology theory $X/S \mapsto H^*(X, *)$ satisfies homotopy property and projective space theorem (of bidegrees $(2i, i)$), then we have the following isomorphisms

$$\begin{aligned} H^*(BGL_a, *) &\simeq H^*(S, *)[x_1, \dots, x_a], \\ H^*(BGL(a, b), *) &\simeq H^*(BGL_a \times BGL_b, *) \simeq H^*(S, *)[x_1, \dots, x_a; y_1, \dots, y_b], \end{aligned}$$

where the classes $x_i, y_i \in H^{2i}(BGL_a, i)$ are the Chern classes of the universal locally free sheaf on BGL_a (defined via a projective space theorem). This applies, for example, to

- (1) étale cohomology with ℓ -adic coefficients;
- (2) (étale) logarithmic de Rham-Witt cohomology $H^*(X, W_n \Omega_{X, \log}^*[-*])$, for a smooth scheme X over k [15, III, Theorem 2.2.5]:

$$\begin{aligned} H^*(BGL_a, W_n \Omega_{BGL_a, \log}^*[-*]) &\simeq H^*(k, W_n \Omega_{k, \log}^*[-*])[x_1, \dots, x_a], \\ H^*(BGL(a, b), W_n \Omega_{BGL(a, b), \log}^*[-*]) &\simeq H^*(BGL_a \times BGL_b, W_n \Omega_{BGL_a \times BGL_b, \log}^*[-*]) \\ &\simeq H^*(k, W_n \Omega_{k, \log}^*[-*])[x_1, \dots, x_a; y_1, \dots, y_b], \end{aligned}$$

where the classes $x_i, y_i \in H^{2i}(BGL_l, W_n \Omega_{BGL_l, \log}^i[-i])$ are the logarithmic de Rham-Witt Chern classes of the universal locally free sheaf on BGL_a . We have $H^*(k, W_n \Omega_{k, \log}^*[-*]) = H_{\text{ét}}^*(k, \mathbf{Z}/p^n)$, which is trivial in degrees at least 2.

(3) motivic cohomology $\mathbf{Z}/p(*)_M$ over a field F [32, Lemma 7]:

$$\begin{aligned} H^*(BGL_a, \mathbf{Z}/p(*)_M) &\simeq H^*(F, \mathbf{Z}/p(*)_M)[x_1, \dots, x_a] \\ H^*(BGL(a, b), \mathbf{Z}/p(*)_M) &\simeq H^*(BGL_a \times BGL_b, \mathbf{Z}/p(*)_M) \\ &\simeq H^*(F, \mathbf{Z}/p(*)_M)[x_1, \dots, x_a; y_1, \dots, y_b], \end{aligned}$$

where $x_i, y_i \in H^{2i}(BGL_a, \mathbf{Z}/p(i)_M)$ are the motivic Chern classes of the universal locally free sheaf on BGL_a .

(4) Nisnevich logarithmic de Rham-Witt cohomology $H_{\text{Nis}}^*(X, W_n \Omega_{X, \log}^*[-*])$, for smooth schemes X over k :

$$(3.1) \quad \begin{aligned} H_{\text{Nis}}^*(BGL_a, W_n \Omega_{BGL_a, \log}^*[-*]) &\simeq H_{\text{Nis}}^*(k, W_n \Omega_{k, \log}^*[-*])[x_1, \dots, x_a], \\ H_{\text{Nis}}^*(BGL(a, b), W_n \Omega_{BGL(a, b), \log}^*[-*]) &\simeq H_{\text{Nis}}^*(BGL_a \times BGL_b, W_n \Omega_{BGL_a \times BGL_b, \log}^*[-*]) \\ &\simeq H_{\text{Nis}}^*(k, W_n \Omega_{k, \log}^*[-*])[x_1, \dots, x_a; y_1, \dots, y_b], \end{aligned}$$

where $x_i, y_i \in H_{\text{Nis}}^{2i}(BGL_l, W_n \Omega_{BGL_l, \log}^i[-i])$ are the Nisnevich logarithmic de Rham-Witt Chern classes of the universal locally free sheaf on BGL_a . This follows from the quasi-isomorphism $\mathbf{Z}/p(i)_M \simeq W_n \Omega_{X, \log}^i[-i]$. We have $H_{\text{Nis}}^*(k, W_n \Omega_{k, \log}^*[-*]) = \mathbf{Z}/p^n$.

Alternatively, one can proceed as in [15, II], and use projective space theorem plus weak purity to obtain the same formulas. This method of computations applies to de Rham (hence crystalline) and to Hodge cohomologies. We get (we write $BGL_{a, n}$ for $BGL_l/W_n(k)$ if it is not confusing)

$$H_{\text{cr}}^*(BGL_{a, n}) \simeq H_{\text{dR}}^*(BGL_{a, n}) \simeq W_n(k)[x_1, \dots, x_a],$$

where the classes $x_i \in H_{\text{dR}}^{2i}(BGL_{a, n})$ are the de Rham Chern classes of the universal locally free sheaf on $BGL_{a, n}$ (defined via a projective space theorem). Similarly, for Hodge cohomology we have

$$H_{\text{Hdg}}^*(BGL_{a, n}) \simeq W_n(k)[x_1, \dots, x_a].$$

In particular

$$H_{\text{dR}}^{2i}(BGL_{a, n}) = H^i(BGL_{a, n}, \Omega_{BGL_{a, n}}^i) = \bigoplus_I W_n(k) x_I, \quad I \subset \{1, \dots, a\}, |I| = i.$$

Similarly, we get

$$\begin{aligned} H_{\text{cr}}^*(BGL(a, b)_n) &\simeq H_{\text{dR}}^*(BGL(a, b)_n) \simeq H_{\text{dR}}^*((BGL_a \times BGL_b)_n) \\ &\simeq W_n(k)[x_1, \dots, x_a; y_1, \dots, y_b]. \end{aligned}$$

3.2. Syntomic computations.

3.2.1. Syntomic cohomology.

Lemma 3.1. *Let B be one of the classifying simplicial schemes BGL_l , $B(GL_a \times GL_b)$, or $BGL(a, b)$. Then*

$$H^i(B, S_n(j)) = \begin{cases} M & \text{for } i = 2j, \\ H_{\text{dR}}^{i-1}(B) & \text{for } i = 2j - 2m - 1, m \geq 0, \\ 0 & \text{for } i \geq 2j + 2, \end{cases}$$

where M is equal to $\bigoplus_I \mathbf{Z}/p^n x_I$ for BGL_l and to $\bigoplus_{I, J} \mathbf{Z}/p^n x_I y_J$ in the other two cases. Here $I \subset \{1, \dots, l\}$, $|I| = j$; $I \subset \{1, \dots, a\}$, $|I| = c$, $J \subset \{1, \dots, b\}$, $|J| = d$, $c + d = j$, respectively.

Moreover we have the following long exact sequence

$$0 \rightarrow H^{2j}(B, S_n(j)) \rightarrow H^{2j}(B_n, \Omega_{B_n}^{\geq j}) \xrightarrow{1-\varphi_j} H_{\text{dR}}^{2j}(B_n) \rightarrow H^{2j+1}(B, S_n(j)) \rightarrow 0.$$

Proof. First we will prove the lemma for BGL_l . We have

$$S_n(j) = \text{Cone}(\Omega_{BGL_{l,n}}^{\geq j} \xrightarrow{1-\varphi_j} \Omega_{BGL_{l,n}}^\bullet)[-1]$$

This yields the long exact sequence

$$\cdots \rightarrow H^i(BGL_l, S_n(j)) \rightarrow H^i(BGL_{l,n}, \Omega_{BGL_{l,n}}^{\geq j}) \xrightarrow{1-\varphi_j} H_{\text{dR}}^i(BGL_{l,n}) \rightarrow H^{i+1}(BGL_l, S_n(j)) \rightarrow \cdots$$

Using the vanishing of the de Rham cohomology in odd degrees we get the following long exact sequence ($i \geq 0$)

$$0 \rightarrow H^{2i}(B, S_n(j)) \rightarrow H^{2i}(B_n, \Omega_{B_n}^{\geq j}) \xrightarrow{1-\varphi_j} H_{\text{dR}}^{2i}(B_n) \rightarrow H^{2i+1}(B, S_n(j)) \rightarrow 0.$$

Hence the long exact sequence in the lemma.

By considering the maps

$$H^{2j}(B, S_n(j)) \rightarrow H^{2j}(B_n, \Omega_{B_n}^{\geq j})^{\varphi_j=1}, \quad H_{\text{dR}}^{i-1}(B_n) \rightarrow H^i(B, S_n(j))$$

and by devissage we can reduce the computations to $n = 1$. The long exact sequence in our lemma implies that

$$H^{2j}(BGL_l, S_1(j)) \simeq H^{2j}(BGL_{l,1}, \Omega_{BGL_{l,1}}^{\geq j})^{\varphi_j=1} = (\oplus_I W_1(k)x_I)^{\varphi_j=1},$$

where $I \subset \{1, \dots, l\}$, $|I| = j$. But $\varphi(x_i) = p^i x_i$; hence $H^{2j}(BGL_l, S_1(j)) \simeq \oplus_I \mathbf{Z}/px_I$, as wanted.

For $2i \geq 2j + 2$ we claim that the map

$$1 - \varphi_j : H^{2i}(BGL_{l,1}, \Omega_{BGL_{l,1}}^{\geq j}) \rightarrow H_{\text{dR}}^{2i}(BGL_{l,1})$$

is an isomorphism. By the computations of de Rham and Hodge cohomologies above it suffices to show that the map

$$1 - \varphi_j : H^i(BGL_{l,1}, \Omega_{BGL_{l,1}}^i) \rightarrow H^i(BGL_{l,1}, \Omega_{BGL_{l,1}}^i)$$

is an isomorphism. But $\varphi_j = p^{i-j}\varphi_i$ and $1 - \varphi_j = 1$, as wanted. It follows that $H^i(BGL_l, S_1(j)) = 0$ for $i \geq 2j + 2$.

For $i \leq 2j - 2$ and i even we have $H^i(BGL_l, S_1(j)) \hookrightarrow H^i(BGL_{l,1}, \Omega_{BGL_{l,1}}^{\geq j}) = 0$. This implies that, for $i \leq 2j$ and i odd, $H^i(BGL_l, S_1(j)) = H_{\text{dR}}^{i-1}(BGL_{l,1})$, as wanted.

In view of the above computations of crystalline cohomology, the argument for $B(GL_a \times GL_b)$ and $BGL(a, b)$ is practically identical. \square

Similarly, we have the following lemma.

Lemma 3.2. *Let B be as above. Then*

$$H^i(B, S'_n(j)) = \begin{cases} M' & \text{for } i = 2j, \\ H_{\text{dR}}^{i-1}(B_n) & \text{for } i = 2j - 2m - 1, m \geq 0, \\ H_{\text{dR}}^{i-1}(B_j) & \text{for } i = 2j + 2m + 1, m \geq 1, \\ H_{\text{dR}}^i(B_{n-j}) & \text{for } i = 2j + 2m, m \geq 1, \\ 0 & \text{otherwise,} \end{cases}$$

where M' is equal to $(\oplus_I W_n(k)x_I)^{\varphi=p^j}$ for BGL_l and to $(\oplus_{I,J} W_n(k)x_{Iy_J})^{\varphi=p^j}$ in the other two cases (with the index sets I, J as in the previous lemma).

Moreover we have the following long exact sequence

$$0 \rightarrow H^{2j}(B, S'_n(j)) \rightarrow H^{2j}(B_n, \Omega_{B_n}^{\geq j}) \xrightarrow{p^j-\varphi} H_{\text{dR}}^{2j}(B_n) \rightarrow H^{2j+1}(B, S'_n(j)) \rightarrow 0.$$

Proof. We present the proof for BGL_l . The proofs for the other schemes are basically the same. We have the long exact sequence

$$\cdots \rightarrow H^i(BGL_l, S'_n(j)) \rightarrow H^i(BGL_{l,n}, \Omega_{BGL_{l,n}}^{\geq j}) \xrightarrow{p^j-\varphi} H_{\text{dR}}^i(BGL_{l,n}) \rightarrow H^{i+1}(BGL_l, S'_n(j)) \rightarrow \cdots$$

Using the vanishing of the de Rham cohomology in odd degrees we get the following long exact sequence ($i \geq 0$)

$$0 \rightarrow H^{2i}(B, S'_n(j)) \rightarrow H^{2i}(B_n, \Omega_{B_n}^{\geq j}) \xrightarrow{p^j - \varphi} H_{\text{dR}}^{2i}(B_n) \rightarrow H^{2i+1}(B, S'_n(j)) \rightarrow 0.$$

Hence the long exact sequence in the lemma. It implies that

$$H^{2j}(BGL_l, S'_n(j)) \simeq H^{2j}(BGL_{l,n}, \Omega_{BGL_{l,n}}^{\geq j})^{\varphi=p^j} = (\oplus_I W_n(k)x_I)^{\varphi=p^j},$$

where $I \subset \{1, \dots, l\}$, $|I| = j$.

For $2i \geq 2j + 2$ we claim that the following sequence is exact

$$\begin{aligned} 0 \rightarrow H^{2i}(BGL_{l,n-j}, \Omega_{BGL_{l,n-j}}^{\geq j}) \rightarrow H^{2i}(BGL_{l,n}, \Omega_{BGL_{l,n}}^{\geq j}) \xrightarrow{p^j - \varphi} H_{\text{dR}}^{2i}(BGL_{l,n}) \\ \rightarrow H_{\text{dR}}^{2i}(BGL_{l,j}) \rightarrow 0 \end{aligned}$$

Indeed, by the computations of de Rham and Hodge cohomologies above it suffices to show that the following sequence is exact

$$\begin{aligned} 0 \rightarrow H^i(BGL_{l,n-j}, \Omega_{BGL_{l,n-j}}^i) \rightarrow H^i(BGL_{l,n}, \Omega_{BGL_{l,n}}^i) \xrightarrow{p^j - \varphi} H^i(BGL_{l,n}, \Omega_{BGL_{l,n}}^i) \\ \rightarrow H^i(BGL_{l,j}, \Omega_{BGL_{l,j}}^i) \rightarrow 0 \end{aligned}$$

But $p^j - \varphi = p^j(1 - \varphi_j)$ and we have shown in the previous proof that $1 - \varphi_j$ is an isomorphism.

For $i \leq 2j - 2$ and i even we have $H^i(BGL_l, S'_n(j)) \hookrightarrow H^i(BGL_{l,n}, \Omega_{BGL_{l,n}}^{\geq j}) = 0$. This implies that, for $i \leq 2j$ and i odd, $H^i(BGL_l, S'_n(j)) = H_{\text{dR}}^{i-1}(BGL_{l,n})$, as wanted. \square

Remark 3.3. Note that the map

$$\tau : H^{2j}(BGL_l, S_n(j)) \hookrightarrow H^{2j}(BGL_l, S'_n(j)) : \oplus_I \mathbf{Z}/p^n x_I \hookrightarrow (\oplus_I W_n(k)x_I)^{\varphi=p^j},$$

is the canonical injection and that it is compatible with products (since both products are compatible with the de Rham product). An analogous statement is true for the maps

$$\begin{aligned} \tau : H^{2j}(BGL_a \times BGL_b, S_n(j)) \hookrightarrow H^{2j}(BGL_a \times BGL_b, S'_n(j)), \\ \tau : H^{2j}(BGL(a, b), S_n(j)) \hookrightarrow H^{2j}(BGL(a, b), S'_n(j)) \end{aligned}$$

that are both equal to the canonical injection

$$\oplus_{I,J} \mathbf{Z}/p^n x_I y_J \hookrightarrow (\oplus_{I,J} W_n(k)x_I y_J)^{\varphi=p^j}.$$

3.2.2. Truncated syntomic cohomology. We will now compute the truncated syntomic cohomology of BGL_l and the related simplicial schemes $B(GL_a \times GL_b)$ and $BGL(a, b)$.

Proposition 3.4. *Let B denote the simplicial scheme $BGL_l/W(k)$, $B(GL_a \times GL_b)/W(k)$, or $BGL(a, b)/W(k)$. Then*

(1) *the natural morphism $S_n(j)_{\text{Nis}} \rightarrow S_n(j)$ induces an isomorphism*

$$H^{2j}(B, S_n(j)_{\text{Nis}}) \xrightarrow{\sim} H^{2j}(B, S_n(j)), \quad j \geq 0.$$

(2) *the morphism*

$$\tau : H^{2j}(B, S_n(j)_{\text{Nis}}) \rightarrow H^{2j}(B, S'_n(j)_{\text{Nis}}), \quad j \geq 0.$$

is compatible with products.

Proof.

Lemma 3.5. *For a smooth scheme X over $W(k)$ and $n \geq m \geq 0$, the following sequence is exact*

$$0 \rightarrow S_m(j)_{\text{Nis}} \rightarrow S_n(j)_{\text{Nis}} \rightarrow S_{n-m}(j)_{\text{Nis}} \rightarrow 0$$

Proof. We can argue locally so assume that there is a Frobenius lift on X_n , $n \geq 0$. Assume first that $j \geq 1$. Recall [23, 3.5] that then the syntomic complex $S_n(j)$ can be represented by the complex

$$(3.2) \quad 0 \rightarrow \mathcal{O}_{X_n} \xrightarrow{-d} \Omega_{X_n}^1 \xrightarrow{-d} \dots \xrightarrow{-d} \Omega_{X_n}^{j-2} \xrightarrow{(0, -d)} \Omega_{X_n}^j \oplus \Omega_{X_n}^{j-1} \xrightarrow{(1-\varphi_j, -d)} \Omega_{X_n}^j,$$

where \mathcal{O}_{X_n} is in degree 1. On the étale site the last map is surjective and $\tau_{\leq j} S_n(j)_{\text{ét}} \simeq S_n(j)_{\text{ét}}$ [23, 3.6]. However this is not true on the Nisnevich site. The truncated complex $S_n(j)_{\text{Nis}}$ can be represented by the complex

$$0 \rightarrow \mathcal{O}_{X_n} \xrightarrow{-d} \Omega_{X_n}^1 \xrightarrow{-d} \dots \xrightarrow{-d} \Omega_{X_n}^{j-2} \xrightarrow{-d} N_{X_n}(j)_{\text{Nis}},$$

where $N_{X_n}(j)_*$ is the kernel of the map $(1 - \varphi_j, -d) : \Omega_{X_n}^j \oplus \Omega_{X_n}^{j-1} \rightarrow \Omega_{X_n}^j$ taken in topology $*$ and \mathcal{O}_{X_n} is in degree 1.

Let

$$\nu_{X,n}^j = \ker(1 - \varphi_j : \Omega_{X_n}^j \rightarrow \Omega_{X_n}^j / d\Omega_{X_n}^{j-1}), \quad j \geq 0.$$

We easily find that we have the short exact sequence

$$(3.3) \quad 0 \rightarrow Z\Omega_{X_n}^{j-1} \rightarrow N_{X_n}(j)_{\text{Nis}} \rightarrow \nu_{X,n}^j \rightarrow 0,$$

where $Z\Omega_{X_n}^{j-1} = \ker(d : \Omega_{X_n}^{j-1} \rightarrow \Omega_{X_n}^j)$. We note that $Z\Omega_{X_n}^{j-1} \simeq Z\Omega_{X_n}^{j-1} \otimes \mathbf{Z}/p^n$. It follows that we have the short exact sequence

$$(3.4) \quad 0 \rightarrow \tilde{S}_n(j)_{\text{Nis}} \rightarrow S_n(j)_{\text{Nis}} \rightarrow \nu_{X,n}^j \rightarrow 0,$$

where $\tilde{S}_n(j)_{\text{Nis}}$ is the following complex

$$0 \rightarrow \mathcal{O}_{X_n} \xrightarrow{-d} \Omega_{X_n}^1 \xrightarrow{-d} \dots \xrightarrow{-d} \Omega_{X_n}^{j-2} \xrightarrow{-d} Z\Omega_{X_n}^{j-1},$$

We have the short exact sequence

$$0 \rightarrow \tilde{S}_m(j)_{\text{Nis}} \rightarrow \tilde{S}_n(j)_{\text{Nis}} \rightarrow \tilde{S}_{n-m}(j)_{\text{Nis}} \rightarrow 0$$

It follows that it suffices to show that the natural sequence

$$0 \rightarrow \nu_{X,m}^j \rightarrow \nu_{X,n}^j \rightarrow \nu_{X,n-m}^j \rightarrow 0$$

is exact. We will show below that we have a natural isomorphism $\nu_{X,n,\text{ét}}^j \xrightarrow{\sim} W_n \Omega_{X_1,\log}^j$ hence a natural isomorphism $\nu_{X,n}^j \xrightarrow{\sim} W_n \Omega_{X_1,\log}^j$. Since $W_n \Omega_{X_1,\log}^j$ is locally in the Nisnevich topology generated by symbols $d\log\{[a_1], \dots, [a_j]\}$, for local sections $a_1, \dots, a_j \in \mathcal{O}_{X_1}^\times$ [18, p.505], this will give us the exactness we want.

To construct the map $f : \nu_{X,n,\text{ét}}^j \rightarrow W_n \Omega_{X_1,\log}^j$, consider the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \nu_{X,n,\text{ét}}^j & \longrightarrow & \Omega_{X_n}^j & \xrightarrow{1-\varphi_j} & \Omega_{X_n}^j / d\Omega_{X_n}^{j-1} & \longrightarrow & 0 \\ & & & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & W_n \Omega_{X_1,\log}^j & \longrightarrow & W_n \Omega_{X_1}^j / dVW_{n-1} \Omega_{X_1}^{j-1} & \xrightarrow{1-F_j} & W_n \Omega_{X_1}^j / dW_n \Omega_{X_1}^{j-1} & \longrightarrow & 0 \end{array}$$

The vertical maps are induced by the global Frobenius lift. The top sequence is exact by definition, the bottom one by Lemma 4.3 from [3]. We obtained the map f . To show that it is an isomorphism we can reduce by devissage to $n = 1$. But then our claim is clear.

For $j = 0$, we we have

$$S_n(0)_{\text{ét}} : \mathcal{O}_{X_n} \xrightarrow{1-\varphi} \mathcal{O}_{X_n}.$$

Hence $\tau_{\leq 0} S_n(0)_{\text{ét}} \simeq S_n(0)_{\text{ét}}$ and

$$S_n(0)_{\text{Nis}} = \ker(1 - \varphi : \mathcal{O}_{X_n} \rightarrow \mathcal{O}_{X_n}) = \nu_{X,n}^0,$$

which we have shown satisfies devissage. \square

We claim now that to show that the map $H^{2j}(B_n, S_n(j)_{\text{Nis}}) \rightarrow H^{2j}(B_n, S_n(j))$, $j \geq 0$, is an isomorphism we may assume $n = 1$. Indeed, by Lemma 3.5, it suffices to show that $H^{2j+1}(B_n, S_n(j)_{\text{Nis}}) = 0$ and that the map $H^{2j-1}(B_n, S_n(j)) \rightarrow H^{2j-1}(B_m, S_m(j))$, $n \geq m$, is surjective. For the first, by devissage, reduce to $n = 1$. Then we have a global Frobenius. Since $H^a(B_n, \Omega_{B_n}^b)$, $a \neq b$, we reduce to showing that $H^{j+1}(B_n, N_{B_n}(j)_{\text{Nis}}) = 0$ or, by the exact sequence (3.3), to $H^{j+1}(B_n, \nu_{B_n}^j) = 0$ and $H^{j+1}(B_n, Z\Omega_{B_n}^{j-1}) = 0$. But this follows from (3.1) and from (3.6) below. For the second claim, by Lemma 3.1, we need to show that the map $H_{\text{dR}}^{2j-2}(B_n) \rightarrow H_{\text{dR}}^{2j-2}(B_m)$, $n \geq m$, is surjective. But this is clear.

Take then $n = 1$. We want to show that the map $H^{2j}(B_1, S_1(j)_{\text{Nis}}) \rightarrow H^{2j}(B_1, S_1(j))$, $j \geq 0$, is an isomorphism. Consider the natural map $S_n(j)_{\text{Nis}} \rightarrow W_n \Omega_{B_1, \log}^j[-j]$ and its étale analog. Since the map $H^j(B_{1, \text{Nis}}, \Omega_{B_1, \log}^j) \rightarrow H^j(B_{1, \text{ét}}, \Omega_{B_1, \log}^j)$ is an isomorphism (see Section 3.1), it suffices to show that so are the maps

$$(3.5) \quad H^{2j}(B_1, S_n(j)_{\text{Nis}}) \rightarrow H^j(B_1, \Omega_{B_1, \log}^j), \quad H^{2j}(B_1, S_n(j)_{\text{Nis}}) \rightarrow H^j(B_1, \Omega_{B_1, \log}^j).$$

We will argue in the Nisnevich case, the étale case being analogous. We have a global Frobenius. Hence, by the short exact sequence (3.4), it suffices to show that $H^{2j}(B_n, \tilde{S}_n(j)_{\text{Nis}}) = H^{2j+1}(B_n, \tilde{S}_n(j)_{\text{Nis}}) = 0$.

Since $H^i(B_n, \Omega_{B_n}^k) = 0$ unless $i = k$, we have $H^{2j+1}(B_n, \tilde{S}_n(j)_{\text{Nis}}) = 0$. The vanishing of $H^{2j}(B_n, \tilde{S}_n(j)_{\text{Nis}})$ will follow when we show that $H^j(B_n, Z\Omega_{B_n}^{j-1}) = 0$. We will, in fact, show more. Namely that

$$(3.6) \quad H^k(B_n, Z\Omega_{B_n}^j) = H^k(B_n, B\Omega_{B_n}^j) = 0, \quad k > j,$$

where $B\Omega_{B_n}^j = \text{Im}(d : \Omega_{B_n}^{j-1} \rightarrow \Omega_{B_n}^j)$. We will argue by induction on j starting with $j = -1$, which is the trivial case. Assume that the statement is true for j . To prove it for $j+1$ recall that we have the isomorphism

$$C^{-1} : W_n \Omega_{B_1}^* \xrightarrow{\sim} \mathcal{H}^*(W_n \Omega_{B_1}^*).$$

Here C^{-1} is the inverse Cartier isomorphism. It yields the following short exact sequence of abelian sheaves

$$0 \rightarrow B\Omega_{B_n}^k \rightarrow Z\Omega_{B_n}^k \rightarrow \Omega_{B_1}^k \rightarrow 0$$

Applying cohomology we get the exact sequence

$$H^k(B_n, B\Omega_{B_n}^{j+1}) \rightarrow H^k(B_n, Z\Omega_{B_n}^{j+1}) \rightarrow H^k(B_n, \Omega_{B_1}^{j+1}), \quad k > j+1.$$

The last group is trivial, so it remains to show that so is $H^k(B_n, B\Omega_{B_n}^{j+1})$. For that we use the short exact sequence

$$0 \rightarrow Z\Omega_{B_n}^j \rightarrow \Omega_{B_n}^j \xrightarrow{d} B\Omega_{B_n}^{j+1} \rightarrow 0$$

and the induced long exact sequence

$$H^k(B_n, \Omega_{B_n}^j) \xrightarrow{d} H^k(B_n, B\Omega_{B_n}^{j+1}) \rightarrow H^{k+1}(B_n, Z\Omega_{B_n}^j) \rightarrow H^{k+1}(B_n, \Omega_{B_1}^j).$$

Since the first and the last group are trivial we get that $H^k(B_n, B\Omega_{B_n}^{j+1}) \xrightarrow{\sim} H^{k+1}(B_n, Z\Omega_{B_n}^j)$. But, by induction, the last group is trivial and we are done.

For the second statement of the proposition, consider the following commutative diagram

$$\begin{array}{ccccc} H^{2j}(B, S_n(j)_{\text{Nis}}) & \xrightarrow{\tau} & H^{2j}(B, S'_n(j)_{\text{Nis}}) & \hookrightarrow & H_{\text{dR}}^{2j}(B_n) \\ \downarrow \wr & & \downarrow & \nearrow & \\ H^{2j}(B, S_n(j)) & \xrightarrow{\tau} & H^{2j}(B, S'_n(j)) & & \end{array}$$

The top and the bottom injections follow from the fact that $H_{\text{dR}}^{2j-1}(B_n) = 0$; they commute with products. It suffices now to evoke Remark 3.3. \square

Let B denote the simplicial scheme $BGL_l/W(k)$. Let $x_i \in H^{2i}(B, S_n(j)_{\text{Nis}})$, $1 \leq i \leq l$, be the Chern classes of the universal locally free sheaf on B_n obtained from the de Rham classes via the isomorphism in Proposition 3.4.

Corollary 3.6. *The natural morphism*

$$H^*(W(k), S_n(*)_{\text{Nis}})[x_1, \dots, x_l] \xrightarrow{\sim} H^*(B, S_n(*)_{\text{Nis}})$$

is an isomorphism.

Proof. Since we have shown in Proposition 3.4 that the map $H^{2j}(B, S_n(j)_{\text{Nis}}) \rightarrow H^{2j}(B_n, \Omega_{B_n}^{\geq j})$ is an isomorphism, it suffices, by [7, Example 2.7], to show that, for $b \geq 1$, so is the map $H^{2j}(B_n, \Omega_{B_n}^{\bullet}) \rightarrow H^{2j+1}(B, S_n(j+b)_{\text{Nis}})$. But this follows from the fact that $H^{2j}(B_n, \Omega_{B_n}^{\geq b+j}) = H^{2j+1}(B_n, \Omega_{B_n}^{\geq b+j}) = 0$. \square

Analogous statements are true for $B(GL_a \times GL_b)/W(k)$ and $BGL(a, b)/W(k)$.

3.2.3. Truncated syntomic-étale cohomology. We will briefly recall a computation of nearby cycles due to Bloch-Kato [4]. We work in Nisnevich topology. For a smooth scheme X over \mathcal{O}_K , $j \geq 0$, set $L^j = i^*R^j j_* \mathbf{Z}/p(j)_{\text{M}}$. For $m \geq 1$, let $U^m L^j$ be the subsheaf of L^j generated locally by local sections of the form $\{x_1, \dots, x_j\}$ such that $x_1 - 1 \in \pi^m i^* \mathcal{O}_X$, for a uniformizer π of K , and $x_i \in i^* j_* \mathcal{O}_{X_K}^*$, $2 \leq i \leq j$. We set $U^0 L^j := L^j$.

Theorem 3.7. (Bloch-Kato [4])

(1) *For a smooth scheme X over $W(k)$, we have*

$$\text{gr}^m(L^j) \simeq \begin{cases} \Omega_{X_k, \log}^j \oplus \Omega_{X_k, \log}^{j-1} & \text{if } m = 0, \\ \Omega_{X_k}^{j-1} & \text{if } m = 1. \end{cases}$$

Moreover, $U^m L^j = 0$, for $m > 1$.

(2) *For a smooth scheme X over $\mathcal{O}_K = W(k)(\zeta_p)$, we have*

$$\text{gr}^m(L^j) \simeq \begin{cases} \Omega_{X_{k'}, \log}^j \oplus \Omega_{X_{k'}, \log}^{j-1} & \text{if } m = 0, \\ \Omega_{X_{k'}}^{j-1} & \text{if } 1 \leq m < p, \\ \Omega_{X_{k'}}^{j-1}/(1-C)Z\Omega_{X_{k'}}^{j-1} \oplus \Omega_{X_{k'}}^{j-2}/(1-C)Z\Omega_{X_{k'}}^{j-2} & \text{if } m = p. \end{cases}$$

Moreover, for $m > p$, $U^m L^j = 0$.

Proof. Outside of vanishing of $U^m L^j$ both claims are proved in [4, 1.4.1, 6.7, p.135]. For the vanishing of $U^m L^j$ we can argue locally and then, by the Gersten conjecture for étale cohomology, we can pass to the fraction field \tilde{K} of the generic point of the special fiber. It suffices now to show that $U^m H_{\text{ét}}^j(\tilde{K}, \mathbf{Z}/p(j)) = 0$ for $m > e'$, $e' = ep/(p-1)$. But cohomological symbol gives an isomorphism $K_j^M(\tilde{K})/p \xrightarrow{\sim} H_{\text{ét}}^j(\tilde{K}, \mathbf{Z}/p(j))$. Hence it suffices to evoke Lemma 5.1 from [4]. \square

Proposition 3.8. *For B being $BGL_l/W(k)$, $B(GL_a \times GL_b)/W(k)$, or $BGL(a, b)/W(k)$, we have the natural isomorphisms*

$$H^{2j}(B, E_n(j)_{\text{Nis}}) \xrightarrow{\sim} H^{2j}(B, S_n(j)_{\text{Nis}}), \quad H^{2j}(B, E'_n(j)_{\text{Nis}}) \xrightarrow{\sim} H^{2j}(B, S'_n(j)_{\text{Nis}}).$$

Moreover, the morphism

$$\tau : H^{2i}(BGL_l, E_n(i)_{\text{Nis}}) \rightarrow H^{2i}(BGL_l, E'_n(i)_{\text{Nis}})$$

is compatible with products.

Proof. We will start with the first claimed isomorphism. Recall that we have the distinguished triangle in Nisnevich topology (2.9)

$$j_! \tau_{\leq j} R\varepsilon_* \mathbf{Z}/p^n(j)' \rightarrow E_n(j)_{\text{Nis}} \rightarrow S_n(j)_{\text{Nis}},$$

where ε is the projection from the étale to the Nisnevich site. Set $T_n(j) := \tau_{\leq j} R\varepsilon_* \mathbf{Z}/p^n(j)$, $j \geq 0$. It suffices to show that

$$H^{2j}(B, j_! T_n(j)) = H^{2j+1}(B, j_! T_n(j)) = 0, \quad n \geq 1.$$

We will use for that the distinguished triangle

$$(3.7) \quad j_! T_n(j) \rightarrow j_* T_n(j) \rightarrow i_* i^* j_* T_n(j).$$

It suffices to show that

$$(3.8) \quad \begin{aligned} H^{2j-1}(B, j_*T_n(j)) &\rightarrow H^{2j-1}(B, i_*i^*j_*T_n(j)), & H^{2j}(B, j_*T_n(j)) &\xrightarrow{\sim} H^{2j}(B, i_*i^*j_*T_n(j)), \\ H^{2j+1}(B, j_*T_n(j)) &= 0. \end{aligned}$$

We will argue in the case of $B = BGL_l/W(k)$ – the other cases being analogous. We will start with a computation of the cohomology of the complexes $T_n(j)$; this will prove the third equality in (3.8). In the category of smooth schemes over K , by the Beilinson-Lichtenbaum Conjecture, $T_n(j) \simeq \mathbf{Z}/p^n(j)_M$. In particular, the complexes $T_n(j)$ satisfy projective space theorem [22, 4.1] and hence we can define the universal classes

$$x_j \in H^{2j}(BGL_l/K, T_n(j))$$

as the Chern classes of the universal vector bundle over BGL_l/K . We start here with the first Chern class induced by the map $\mathbb{G}_m[-1] \rightarrow \tau_{\leq 1}(\mathbf{R}\varepsilon_*\mathbb{G}_m[-1]) \rightarrow \tau_{\leq 1}\mathbf{R}\varepsilon_*\mathbf{Z}/p^n(1)$ that arises from the étale map $\mathbb{G}_m[-1] \rightarrow \mathbf{Z}/p^n(r)$ of Kummer theory.

These classes are clearly finer than the étale universal classes. In fact we have the following computation.

Lemma 3.9. *We have the isomorphisms*

$$\begin{aligned} H_{\text{ét}}^*(BGL_l/K, \mathbf{Z}/p^n(*)) &\simeq H_{\text{ét}}^*(K, \mathbf{Z}/p^n(*))[x_1, \dots, x_l], \\ H^*(BGL_l/K, T_n(*)) &\simeq H^*(K, T_n(*))[x_1, \dots, x_l]. \end{aligned}$$

They imply the following isomorphisms

$$\begin{aligned} H^{2i+k}(BGL_l/K, T_n(i)) &= 0; \quad k \geq 1; \\ H^{2i}(BGL_l/K, T_n(i)) &\simeq \mathbf{Z}/p^n(i)[x_I], \quad |I| = i, i \leq l; \\ H_{\text{ét}}^{2i}(BGL_l/K, \mathbf{Z}/p^n(i)) &\simeq \mathbf{Z}/p^n(i)[x_I] \oplus \mathbf{Z}/p^n(i)[x_J], \quad |I| = i, |J| = i - 1, i \leq l, \\ H^{2i-1}(BGL_l/K, T_n(i)) &\xrightarrow{\sim} H_{\text{ét}}^{2i-1}(BGL_l/K, \mathbf{Z}/p^n(i)) \simeq H_{\text{ét}}^1(K, \mathbf{Z}/p^n(1))[x_1, \dots, x_l]. \end{aligned}$$

Moreover the natural morphism

$$H^{2i}(BGL_l/K, T_n(i)) \rightarrow H_{\text{ét}}^{2i}(BGL_l/K, \mathbf{Z}/p^n(i)), \quad i \leq l,$$

is the obvious injection.

Proof. The standard computation yields the isomorphism

$$H_{\text{ét}}^*(BGL_l/K, \mathbf{Z}/p^n(*)) \simeq H_{\text{ét}}^*(K, \mathbf{Z}/p^n(*))[x_1, \dots, x_l].$$

For degree $(2i, i)$ only the groups $H_{\text{ét}}^0(K, \mathbf{Z}/p^n) \simeq \mathbf{Z}/p^n$ and $H_{\text{ét}}^2(K, \mathbf{Z}/p^n(1)) \simeq \mathbf{Z}/p^n$ matter. For degree $(2i-1, i)$ only the group $H_{\text{ét}}^1(K, \mathbf{Z}/p^n(1))$ matters. On the other hand, by [32, Lemma 7], we have

$$H^*(BGL_l/K, T_n(*)) \simeq H^*(K, T_n(*))[x_1, \dots, x_l].$$

Since $H^a(K, T_n(b)) = 0$, $a > b$, the groups $H^{2i+k}(BGL_l/K, T_n(i))$, $k \geq 1$, are clearly trivial. Also, for degree $(2i, i)$ (reps. $(2i-1, i)$) only the group $H^0(K, \mathbf{Z}/p^n(0)_M) \simeq H_{\text{ét}}^0(K, \mathbf{Z}/p^n) \simeq \mathbf{Z}/p^n$ ($H^1(K, \mathbf{Z}/p^n(1)_M) \simeq H_{\text{ét}}^1(K, \mathbf{Z}/p^n(1))$) matters. Our lemma follows. \square

To prove the remaining claims in (3.8), we note that, since both $j_*T_n(j)$ and $i_*i^*j_*T_n(j)$ satisfy devissage, we can assume that $n = 1$. For the second map, consider the following commutative diagram

$$\begin{array}{ccc} H^{2j}(B_{\mathcal{O}_{K'}}, j_*\mathbf{Z}/p^n(j)_M) & \longrightarrow & H^{2j}(B_{\mathcal{O}_{K'}}, i_*i^*j_*\mathbf{Z}/p^n(j)_M) \\ \downarrow N_1 & & \downarrow N_2 \\ H^{2j}(B, j_*\mathbf{Z}/p^n(j)_M) & \longrightarrow & H^{2j}(B, i_*i^*j_*\mathbf{Z}/p^n(j)_M) \end{array}$$

Here $\mathcal{O}_{K'}$ is the ring of integers in $K' = K_0(\xi_p)$ and N_1, N_2 denote the maps induced by the norm map (of complexes of étale sheaves)

$$\pi_*\mathbf{Z}/p^n(j)_M \rightarrow \mathbf{Z}/p^n(j)_M,$$

where $\pi : B_{K'} \rightarrow B_F$ is the natural projection. In the case of the map N_2 this uses the isomorphisms

$$\pi_* i_* i^* j_* T_n(j) \simeq \tau_{\leq j} i_* \pi_* i^* R\varepsilon_* Rj_* T_n(j) \simeq \tau_{\leq j} i_* R\varepsilon_* \pi_* i^* Rj_* T_n(j) \simeq \tau_{\leq j} i_* i^* R\varepsilon_* Rj_* \pi_* T_n(j).$$

The second isomorphism follows from the isomorphism $R\varepsilon_* i^* = i^* R\varepsilon_*$ [10, 2.2.b] and the third one from the same isomorphism and the proper base change theorem in étale cohomology.

Since $N_* \pi^* = (p-1)$, to show that the second map in 3.8 is an isomorphism it suffices to show that so is the map

$$H^{2j}(B_{\mathcal{O}_{K'}}, j_* \mathbf{Z}/p(j)_M) \rightarrow H^{2j}(B_{\mathcal{O}_{K'}}, i_* i^* j_* \mathbf{Z}/p(j)_M)$$

Consider the following composition of maps

$$\tau' : H^{2j}(B_{\mathcal{O}_{K'}}, j_* \mathbf{Z}/p(j)_M) \rightarrow H^{2j}(B_{\mathcal{O}_{K'}}, i_* i^* j_* \mathbf{Z}/p(j)_M) \xrightarrow{\sigma_j} H^j(B_{k'}, \Omega_{B_{k'}, \log}^j).$$

The map σ_j is induced by the composition

$$\sigma_j : i^* j_* \mathbf{Z}/p(j)_M \rightarrow i^* \mathcal{H}^j j_* \mathbf{Z}/p(j)_M[-j] \simeq i^* j_* \mathcal{K}_j^M/p[-j] \rightarrow \Omega_{B_{k'}, \log}^j[-j] \oplus \Omega_{B_{k'}, \log}^{j-1}[-j].$$

Here \mathcal{K}_j^M/p is the sheaf of Milnor K-theory groups mod- p and the isomorphism follows from [24, Theorem 7.6.]. The last morphism [4, 6.6] maps symbols $\{\tilde{x}_1, \dots, \tilde{x}_j\}$, for local sections x_i of $\mathcal{O}_{B/k'}$ and \tilde{x}_i lifting of x_i to $i^* \mathcal{O}_{B_{\mathcal{O}_{K'}}}$, to $(\mathrm{dlog}(x_1) \wedge \dots \wedge \mathrm{dlog}(x_j), 0)$ and maps symbols $\{\tilde{x}_1, \dots, \tilde{x}_{j-1}, \pi_{K'}\}$ to $(0, \mathrm{dlog}(x_1) \wedge \dots \wedge \mathrm{dlog}(x_{j-1}))$. First, we claim that the map

$$\sigma_j : H^{2j}(B_{\mathcal{O}_{K'}}, i_* i^* j_* \mathbf{Z}/p(j)_M) \rightarrow H^j(B_{k'}, \Omega_{B_{k'}, \log}^j)$$

is an isomorphism. This will follow from the spectral sequence

$$H^a(B_{\mathcal{O}_{K'}}, i_* i^* \mathcal{H}^b j_* \mathbf{Z}/p(j)_M) \Rightarrow H^{a+b}(B_{\mathcal{O}_{K'}}, i_* i^* j_* \mathbf{Z}/p(j)_M)$$

if we show that

$$H^a(B_{\mathcal{O}_{K'}}, i_* i^* \mathcal{H}^b j_* \mathbf{Z}/p(j)_M) = \begin{cases} 0 & \text{for } a+b=2j-1, 2j, b < j, \\ H^j(B_{k'}, \Omega_{B_{k'}, \log}^j) & \text{for } a=b=j \end{cases}$$

Let us start with $b=j$. By a result of Bloch-Kato [4], the nearby cycles $i_* i^* \mathcal{H}^j j_* \mathbf{Z}/p(j)_M$ have a descending filtration whose graded pieces are listed in Theorem 3.7. It suffices thus to show that (we set $B' = B_{k'}$ and $\Omega_{B'} = \Omega_{B_{k'}, \log}$)

$$H^i(B', \Omega_{B'}^{j-1}/(1-C)Z\Omega_{B'}^{j-1}) = H^i(\Omega_{B'}^{j-2}/(1-C)Z\Omega_{B'}^{j-2}) = 0, \quad i = j, j+1,$$

(note that $H^j(B', \Omega_{B'}^{j-1}) = 0$ and $H^s(B', \Omega_{B', \log}^t) = 0$, for $s \neq t$). But this follows from the exact sequence

$$(3.9) \quad 0 \rightarrow \Omega_{B', \log}^k \rightarrow Z\Omega_{B'}^k \xrightarrow{1-C} \Omega_{B'}^k$$

and the computations from Proposition 3.4 that showed that $H^s(B', Z\Omega_{B'}^t) = 0$, for $s > t$.

For $b < j$, $a+b=2j$, use the isomorphism $\xi_p^{j-b} : i_* i^* \mathcal{H}^b j_* \mathbf{Z}/p(b) \xrightarrow{\sim} i_* i^* \mathcal{H}^b j_* \mathbf{Z}/p(j)$ and once more Theorem 3.7. It suffices thus to show that the groups

$$H^{2j-b}(B', \Omega_{B', \log}^{b-1} \oplus \Omega_{B', \log}^b), \quad H^{2j-b}(B', \Omega_{B'}^{b-1}), \quad H^{2j-b}(B', \Omega_{B'}^{b-1}/(1-C)Z\Omega_{B'}^{b-1}), \quad H^{2j-b}(\Omega_{B'}^{b-2}/(1-C)Z\Omega_{B'}^{b-2})$$

are trivial. But this follows as above from the exact sequence (3.9) and the computations in Proposition 3.4.

The argument for $b < j$, $a+b=2j-1$, is analogous.

It suffices now to show that the composition $\tau' : H^{2j}(B_{\mathcal{O}_{K'}}, j_* \mathbf{Z}/p(j)_M) \rightarrow H^j(B', \Omega_{B', \log}^j)$ is an isomorphism as well. But both groups are isomorphic to $\mathbf{Z}/p[x_I]$, $|I|=j$. Hence it suffices to show that the map τ' is compatible with Chern classes of vector bundles. Since both cohomologies have projective space theorem, the map τ' is functorial and compatible with products, it suffices to show compatibility with the first Chern class maps and this is clear from the definition of the map σ_j .

For the first map of 3.8 we note that injectivity follows from injectivity of the map

$$H_{\text{ét}}^i(B, Rj_* \mathbf{Z}/p(j)) \rightarrow H_{\text{ét}}^i(B, i_* i^* Rj_* \mathbf{Z}/p(j)), \quad i \geq 0.$$

This is because we have Lemma 3.9. But, in fact, the above map is an isomorphism: since the sheaves $i_*i^*Rj_*\mathbf{Z}/p(j)$ satisfy projective space theorem and weak purity this follows from the method of Illusie [15] of computing cohomology of classifying spaces (note that $H_{\text{ét}}^*(W(k), Rj_*\mathbf{Z}/p(j)) \simeq H_{\text{ét}}^*(W(k), i_*i^*Rj_*\mathbf{Z}/p(j))$).

It remains to prove that the first map in (3.8) is surjective. For that consider the localization sequence in motivic cohomology

$$i_*\mathbf{Z}/p(j-1)_{\mathbb{M}}[-2] \rightarrow \mathbf{Z}/p(j)_{\mathbb{M}} \rightarrow j_*\mathbf{Z}/p(j)_{\mathbb{M}}$$

Applying to it the exact functor i_*i^* we obtain the following commutative diagram with short exact sequences as columns

$$\begin{array}{ccccc} H^1(W(k), \mathbf{Z}/p(1)_{\mathbb{M}})[x_I] & \xrightarrow{\gamma} & \ker(\kappa_2) & \xrightarrow{\sim} & H^{j-1}(B_k, \Omega_{B_k}^{j-1}) \\ \downarrow \beta_1 & & \downarrow \beta_2 & & \downarrow \\ H^{2j-1}(B, j_*\mathbf{Z}/p(j)_{\mathbb{M}}) & \longrightarrow & H^{2j-1}(B, i_*i^*j_*\mathbf{Z}/p(j)_{\mathbb{M}}) & \xrightarrow{\sim} & H^{j-1}(B, i_*i^*\mathcal{H}^j j_*\mathbf{Z}/p(j)_{\mathbb{M}}) \\ \downarrow \kappa_1 & & \downarrow \kappa_2 & & \downarrow \sigma_j \\ H^{2j-2}(B_k, \mathbf{Z}/p(j-1)_{\mathbb{M}}) & \xlongequal{\quad} & H^{2j-2}(B_k, \mathbf{Z}/p(j-1)_{\mathbb{M}}) & & \\ \downarrow \wr & & \downarrow \wr & & \downarrow \\ H^{j-1}(B_k, \Omega_{B_k, \log}^{j-1}) & \xlongequal{\quad} & H^{j-1}(B_k, \Omega_{B_k, \log}^{j-1}) & \xrightarrow{\sim} & H^{j-1}(\Omega_{B_k, \log}^{j-1} \oplus \Omega_{B_k, \log}^j). \end{array}$$

The maps κ_1, κ_2 are the boundary maps induced by the above short exact sequence. In the top left term the index I satisfies $|I| = j-1$. For the rightmost column, we note that we have the short exact sequence

$$0 \rightarrow \Omega_{B_k}^{j-1} \rightarrow i^*\mathcal{H}^j j_*\mathbf{Z}/p(j)_{\mathbb{M}} \rightarrow \Omega_{B_k, \log}^{j-1} \oplus \Omega_{B_k, \log}^j \rightarrow 0$$

of Bloch-Kato (quoted in Theorem 3.7). Here the map $\Omega_{B_k}^{j-1} \rightarrow i^*\mathcal{H}^j j_*\mathbf{Z}/p(j)_{\mathbb{M}} \simeq i^*j_*\mathcal{K}_j^M/p$ is defined by sending $x \text{dlog}(y_1) \wedge \cdots \wedge \text{dlog}(y_{j-1})$ to the symbol $\{1 + \tilde{x}p, \tilde{y}_1, \dots, \tilde{y}_{j-1}\}$. This yields the short exact sequences

$$0 \rightarrow H^{j-1}(B_k, \Omega_{B_k}^{j-1}) \rightarrow H^{j-1}(B, i_*i^*\mathcal{H}^j j_*\mathbf{Z}/p(j)_{\mathbb{M}}) \rightarrow H^{j-1}(B_k, \Omega_{B_k, \log}^{j-1} \oplus \Omega_{B_k, \log}^j) \rightarrow 0$$

The fact that the maps κ_1 and κ_2 are surjective, $H^1(W(k), \mathbf{Z}/p(1)_{\mathbb{M}})[x_I] \simeq \ker(\kappa_1)$, and the obvious definitions of the maps β_1 and γ follow from the fact that

$$\begin{aligned} H^*(B, j_*\mathbf{Z}/p(j)_{\mathbb{M}}) &\simeq H^*(F, \mathbf{Z}/p(*)_{\mathbb{M}})[x_1, \dots, x_l]; \\ H^{j-1}(B_k, \Omega_{B_k, \log}^{j-1}) &\simeq \mathbf{Z}/p[x_I], \quad |I| = j-1, \end{aligned}$$

and κ_1 is the natural morphism compatible with the short exact sequence

$$(3.10) \quad 0 \rightarrow H^1(W(k), \mathbf{Z}/p(1)_{\mathbb{M}}) \rightarrow H^1(F, \mathbf{Z}/p(1)_{\mathbb{M}}) \xrightarrow{\kappa_1} H^0(k, \mathbf{Z}/p(0)_{\mathbb{M}}) \rightarrow 0.$$

Note that $H^0(k, \mathbf{Z}/p(0)_{\mathbb{M}}) \simeq \mathbf{Z}/p$.

Consider the natural map

$$H^{2j-1}(B, i_*i^*j_*\mathbf{Z}/p(j)_{\mathbb{M}}) \rightarrow H^{j-1}(B, i_*i^*\mathcal{H}^j j_*\mathbf{Z}/p(j)_{\mathbb{M}}).$$

We claim that it is an isomorphism. This will follow from the fact that $H^{2j-1}(B, \Lambda(j)) = H^{2j}(B, \Lambda(j)) = 0$, for $\Lambda(j) := \tau_{\leq j-1} i_*i^*j_*\mathbf{Z}/p(j)_{\mathbb{M}}$. To see that this is true consider the spectral sequence

$$H^a(B, \mathcal{H}^b \Lambda(j)) \Rightarrow H^{a+b}(B, \Lambda(j)).$$

It suffices to prove that

$$H^a(B, \mathcal{H}^b \Lambda(j)) = 0, \quad \text{for } a+b = 2j-1, a+b = 2j, b < j.$$

We pass to $\mathcal{O}_{K'}$. The norm map $N : H^a(B_{\mathcal{O}_{K'}}, \mathcal{H}^b \Lambda(j)) \rightarrow H^a(B, \mathcal{H}^b \Lambda(j))$ yields that the pullback map $\pi^* : H^a(B, \mathcal{H}^b \Lambda(j)) \rightarrow H^a(B_{\mathcal{O}_{K'}}, \mathcal{H}^b \Lambda(j))$ is injective. Hence it suffices to show that $H^a(B_{\mathcal{O}_{K'}}, \mathcal{H}^b \Lambda(j)) = 0$ for $b < j$ and $a+b = 2j-1, 2j$. But this we have done above. It follows that the map $\ker(\kappa_2) \rightarrow H^{j-1}(B_k, \Omega_{B_k}^{j-1})$ is an isomorphism as well.

It suffices now to show that the map

$$\gamma : H^1(W(k), \mathbf{Z}/p(1)_{\mathbb{M}})[x_I] \rightarrow \ker(\kappa_2)$$

is an isomorphism, or, because it is injective that $H^1(W(k), \mathbf{Z}/p(1)_{\mathbb{M}})[x_I] \simeq H^{j-1}(B_k, \Omega_{B_k}^{j-1})$, $|I| = j-1$, at least abstractly. But

$$H^{j-1}(B_k, \Omega_{B_k}^{j-1}) \simeq k[x_I], \quad |I| = j-1; \quad H^1(W(k), \mathbf{Z}/p(1)_{\mathbb{M}}) \simeq k.$$

The second isomorphism follows from the isomorphisms

$$H^1(W(k), \mathbf{Z}/p^n(1)_{\mathbb{M}}) \simeq K_1^M(W(k))/p^n \simeq W(k)^*/p^n \simeq W_n(k).$$

We have proved the first claim of the proposition.

To show that we have the isomorphism $H^{2j}(B, E'_n(j)_{\text{Nis}}) \xrightarrow{\sim} H^{2j}(B, S'_n(j)_{\text{Nis}})$ we use the distinguished triangle in Nisnevich topology (Lemma 2.9)

$$j_! \tau_{\leq j} R\mathcal{E}_* \mathbf{Z}/p^n(j)' \rightarrow E'_n(j)_{\text{Nis}} \rightarrow S'_n(j)_{\text{Nis}}.$$

It suffices to show that

$$H^{2j}(B, j_! T_n(j)) = H^{2j+1}(B, j_! T_n(j)) = 0, \quad n \geq 1.$$

what we have just done.

The statement about products follows easily from the first part of the proposition and Proposition 3.4. \square

Remark 3.10. It is likely that, in fact, we have a natural isomorphism

$$H^*(B, E_n(*)_{\text{Nis}}) \xrightarrow{\sim} H^*(B, S_n(*)_{\text{Nis}}).$$

Corollary 3.11. *The period map*

$$\alpha_j : H^{2j}(BGL_l, E'_n(j)_{\text{Nis}}) \rightarrow H_{\text{ét}}^{2j}(BGL_{l,K}, \mathbf{Z}/p^n(j)')$$

maps the universal class x_j^{syn} to $p^j x_j^{\text{ét}}$. For $j \leq p-2$ and E -cohomology, we have an analogous statement with no twist necessary.

Proof. We have the following commutative diagram

$$\begin{array}{ccc} H^{2j}(BGL_l, E'_n(j)_{\text{Nis}}) & \xrightarrow{\alpha_j} & H_{\text{ét}}^{2j}(BGL_{l,F}, \mathbf{Z}/p^n(j)') \\ \downarrow \wr & & \downarrow i_* \\ H^{2j}(BGL_l, S'_n(j)) & \xrightarrow{\alpha_j} & H_{\text{ét}}^{2j}(BGL_l, i_* i^* \mathbf{Z}/p^n(j)') \\ \downarrow & \nearrow p^j \tilde{\alpha}_j & \\ H^{2j}(BGL_l, S_n(j)) & & \end{array}$$

The top left arrow is an isomorphism by Propositions 3.4 and 3.8, the right arrow by the proof of Lemma 3.9. It suffices now to show that the bottom period map α_j is compatible with the universal classes as stated. But this was shown in the proof of Theorem 4.10 in [31], i.e., we have

$$\alpha_j(x_j^{\text{syn}}) = p^j \tilde{\alpha}_j(x_j^{\text{syn}}) = p^j x_j^{\text{ét}},$$

as wanted. \square

4. K-THEORY

In this section we review or prove basic facts concerning K -theory of simplicial schemes and log- K -theory.

4.1. K-theory of simplicial schemes. We start with K -theory of simplicial schemes.

4.1.1. *Definition of K-theory.* For a scheme X , let $K(X)$ be the Thomason-Throbaugh spectrum of nonconnective K-theory as defined in Thomason-Throbaugh [37, 6.4]. For a natural number $n \geq 2$, let $K/n(X)$ denote the corresponding spectrum mod n [37, 9.3]. After the usual rigidification [13, 5.1.2], which we assume from now on, these spectra are strictly contravariant in X and covariant on the categories of noetherian schemes and proper maps of finite Tor-dimension and of quasi-compact schemes and perfect projective morphisms [37, 3.16.5, 3.16.5].

Following Thomason [36, 5.6], we define functors from simplicial schemes to the category of spectra

$$X \mapsto K(X) := \operatorname{holim}_i K(X_i), \quad X \mapsto K/n(X) := \operatorname{holim}_i K/n(X_i),$$

and set

$$K_i(X) := \pi_i(K(X)), \quad K_i(X, \mathbf{Z}/n) := \pi_i(K/n(X)).$$

We have

$$K_j(X) = \pi_j(\operatorname{holim}_i K(X_i)_0), \quad j \geq 0; \quad K_j(X, \mathbf{Z}/n) = \pi_j(\operatorname{holim}_i K(X_i)_0, \mathbf{Z}/n), \quad j \geq 2.$$

Here, for any spectrum F , F_0 denotes its 0'th space.

We will also need K -theory with compact support. Let X be a scheme and $i : D \hookrightarrow X$ a simple normal crossing divisor. Let D_1, D_2, \dots, D_n be the irreducible components of D . Let $\tilde{D}^\bullet = \operatorname{cosk}_0^D(D^{(1)})$ be the coskeleton of $D^{(1)} = \coprod_{i=1}^{i=n} D_i$ over D , i.e., the Čech-nerve of the map $D^{(1)} \rightarrow D$. We have

$$\operatorname{cosk}_0^D(D^{(1)})_n = D^{(1)} \times_D \dots \times_D D^{(1)}, \quad (n+1) - \text{times},$$

with the natural boundary and degeneracy maps. We have the following K -theory with compact support spectra

$$K^c(X, D) := \operatorname{fiber}(K(X) \xrightarrow{i^*} K(\tilde{D}^\bullet)), \quad K^c/n(X, D) := \operatorname{fiber}(K/n(X) \xrightarrow{i^*} K/n(\tilde{D}^\bullet)),$$

and the following K -theory groups with compact support

$$K_i^c(X, D) := \pi_i(K^c(X, D)), \quad K_i^c(X, D, \mathbf{Z}/n) := \pi_i(K^c/n(X, D)).$$

Clearly

$$K_j^c(X, D) = \pi_j(\operatorname{fiber}(K(X)_0 \xrightarrow{i^*} K(\tilde{D}^\bullet)_0)), \quad j \geq 0;$$

$$K_j^c(X, D, \mathbf{Z}/n) = \pi_j(\operatorname{fiber}(K(X)_0 \xrightarrow{i^*} K(\tilde{D}^\bullet)_0, \mathbf{Z}/n)), \quad j \geq 2.$$

For a scheme Y with an ample family of line bundles [37, 2.1.1] (for example, for Y regular or quasi-projective), there exists a strictly natural homotopy equivalence $K(Y) \simeq K_Q(Y)$, where $K_Q(Y)$ is the Quillen K-theory spectrum of Y suitably rigidified [13, 5.1.2], [37, 3.10]. Hence, strictly naturally,

$$K(Y)_0 \simeq K_Q(Y)_0 \simeq \Omega BQP(Y),$$

where $\Omega BQP(Y)$ is the loop space of the classifying space of Quillen Q-construction applied to the category of finitely generated locally free sheaves on Y . Recall that for $Y = \operatorname{Spec}(A)$, the strictly natural map

$$K_0(A) \times BGL(A)^+ \xrightarrow{\sim} \Omega BQP(Y)$$

is a homotopy equivalence.

4.1.2. *Operations on K-theory.* Let now X be a simplicial scheme that is degenerate above certain degree. Using the above homotopy equivalences of spaces, the classical definition of λ -operations on K-theory can be extended to $K_j(X)$, $j \geq 0$, and to $K_j(X, \mathbf{Z}/n)$, $j \geq 2$, [14, 3.2], [17, B.2.6, B.2.8]. Let us recall how this is done.

We work with the site C of noetherian B -schemes, for B a field or a local ring, equipped with the Zariski topology. We equip the category of (globally pointed) presheaves of simplicial sets on C with Jardine's model structure [19]. Recall that a map of presheaves of simplicial sets $E \rightarrow F$ is called a weak equivalence if it induces an isomorphism $\tilde{\pi}_*(E) \rightarrow \tilde{\pi}_*(F)$ on the sheaves of homotopy groups. For a presheaf of simplicial sets F we will denote by F^f a fibrant replacement of F . That is, we have a map $F \rightarrow F^f$ to a fibrant presheaf of simplicial sets F^f that is a weak equivalence. Note that $F \rightarrow F^f$ can be chosen to be functorial.

For a presheaf of simplicial sets X , we define the cohomology of X with values in F

$$H^{-m}(X, F) := [S^m X, F] = \pi(S^m X, F^f), \quad m \geq 0.$$

Here $S^m X$ denotes the m 'th suspension of the presheaf X . The bracket $[,]$ denotes maps in the homotopy category of presheaves of simplicial sets and $\pi(,)$ stands for the set of pointed homotopy classes of maps. Similarly, we define the mod p^n cohomology of X with values in F

$$H^{-m}(X, F; \mathbf{Z}/p^n) := [P^m X, F] = \pi(P^m X, F^f), \quad m \geq 2.$$

Here $P^m X = P^m \wedge X$, where P^m denotes the constant presheaf of m -dimensional mod p^n Moore spaces.

We have several important spectral sequences. By filtering X by its skeletons one constructs spectral sequences of Bousfield-Kan type [17, B.1.7].

$$(4.1) \quad \begin{aligned} E_1^{st} &= H^{-t}(X_s, F) \Rightarrow H^{s-t}(X, F), \quad t - s \geq 1, \\ E_1^{st} &= H^{-t}(X_s, F; \mathbf{Z}/m) \Rightarrow H^{s-t}(X, F; \mathbf{Z}/m), \quad t - s \geq 3. \end{aligned}$$

By the same method we get the other hypercohomology spectral sequences, namely, the weight spectral sequences [36, 5.13, 5.48].

$$(4.2) \quad \begin{aligned} E_2^{st} &= H^s(n \mapsto \pi_t(F(X_n))) \Rightarrow H^{s-t}(X, F), \quad t - s \geq 1, \\ E_2^{st} &= H^s(n \mapsto \pi_t(F(X_n), \mathbf{Z}/m)) \Rightarrow H^{s-t}(X, F; \mathbf{Z}/m), \quad t - s \geq 3. \end{aligned}$$

Finally we have the Brown spectral sequences induced by taking the Postnikov tower of F [14, Prop. 2]

$$(4.3) \quad \begin{aligned} E_2^{st} &= H^s(X, \tilde{\pi}_t(F)) \Rightarrow H^{s-t}(X, F), \quad t - s \geq 1, \\ E_2^{st} &= H^s(X, \tilde{\pi}_t(F, \mathbf{Z}/n)) \Rightarrow H^{s-t}(X, F; \mathbf{Z}/n), \quad t - s \geq 3. \end{aligned}$$

In these spectral sequences the r 'th differential is $d_r : E_r^{s,t} \rightarrow E_r^{s+r, t+r-1}$. The spectral sequences converge strongly for X degenerate above certain simplicial degree.

Denote by K the presheaf $\mathbf{Z} \times \mathbf{Z}_\infty BGL$, where $BGL(U) = \text{inj lim}_n BGL_n(U)$. Then, for a simplicial scheme X , we have [14, 3.2.3]

$$K_m(X) = H^{-m}(X, K), \quad m \geq 0; \quad K_m(X, \mathbf{Z}/p^n) = H^{-m}(X, K; \mathbf{Z}/p^n), \quad m \geq 2,$$

where we wrote X for the scheme X as well as for the presheaf of simplicial sets on C represented by X . This presentation of K-theory groups as generalized cohomology of presheaves of the K-theory spaces allowed Gillet-Soulé [14, 4.2] to mimic the classical definition and to define maps

$$\lambda^k : K_m(X) \rightarrow K_m(X), \quad k \geq 1, m \geq 0,$$

that turn $K_m(X)$ into an $H^0(S^0, \mathbf{Z})$ - λ -algebra for any presheaf X that is K -coherent [14, 3.1], i.e., for which

$$\text{inj lim}_N H^{-m}(X, K^N) \xrightarrow{\sim} H^{-m}(X, K),$$

where $K^N = \mathbf{Z} \times \mathbf{Z}_\infty BGL_N$. Note that $H^0(S^0, \mathbf{Z})$ is a λ -ring. Similarly, for any K -coherent presheaf X , we can define maps

$$\lambda^k : K_m(X, \mathbf{Z}/p^n) \rightarrow K_m(X, \mathbf{Z}/p^n), \quad k \geq 1, m \geq 2,$$

that make $K_m(X, \mathbf{Z}/p^n)$, $m \geq 2$, into an $H^0(S^0, \mathbf{Z})$ - λ -algebra.

4.1.3. Examples. Presheaves whose components are representable by schemes of the site C (except for one copy of $*$ in each degree) we will call constructed from schemes. If they are degenerate above a finite simplicial degree and are built from regular schemes they are K -coherent [20, Lemma 2.1]. Loosely, we will call such presheaves (pointed regular) finite simplicial schemes.

Example 4.1. A simplicial scheme gives rise to a presheaf built from schemes. An m -truncated simplicial scheme X gives rise to a finite simplicial scheme $\text{sk}_m X$ that is degenerate above degree m . If X is a regular schemes and $i : D \hookrightarrow X$ a divisor that has normal crossings then the simplicial scheme $\tilde{D}_\bullet = \text{cosk}_0^D(D^{(1)})$

is not degenerate above any simplicial degree if D has more than one irreducible component. To remedy this note that

$$\tilde{D}_n = \coprod_{I \in \{1, \dots, m\}^n} \bigcap_{i \in I} D_i, \quad D = \bigcup_{i=1}^m D_i.$$

And consider the simplicial scheme

$$\tilde{D}'_\bullet : n \mapsto \coprod_{I \in \Delta(m)_n} \bigcap_{i \in I} D_i$$

with the natural face and degeneracy maps. Here $\Delta(m)$ is the simplicial set

$$\Delta(m)_k = \{(i_0, \dots, i_k) | 1 \leq i_0 \leq \dots \leq i_k \leq m\}$$

The simplicial scheme \tilde{D}'_\bullet is degenerate above degree $m-1$ and the natural inclusion of simplicial schemes

$$\tilde{D}'_\bullet \rightarrow \tilde{D}_\bullet$$

is a weak equivalence [17, Lemma 7.1].

In particular, this can be applied to K -theory with compact support. Set

$$C(X, D) := \text{cofiber}(\tilde{D}_\bullet \xrightarrow{i_*} X).$$

Define

$$K_m^c(X, D) = H^{-m}(C(X, D), K), \quad m \geq 0; \quad K_m^c(X, D, \mathbf{Z}/p^n) = H^{-m}(C(X, D), K; \mathbf{Z}/p^n), \quad m \geq 2.$$

Replacing $C(X, D)$ with the weakly equivalent $C'(X, D) = \text{cofiber}(\tilde{D}'_\bullet \xrightarrow{i_*} X)$ that is degenerate above degree m we get λ -operations on K -theory in the case all the irreducible components of D are regular.

Example 4.2. For any map $f : Y \rightarrow X$ of noetherian finite dimensional schemes the cone $C(X, Y)$ is degenerate above finite simplicial degree. The homotopy cofibre sequence

$$Y \xrightarrow{f} X \rightarrow C(X, Y)$$

yields the long exact sequence

$$\rightarrow H^{-m}(C(X, Y), K; \mathbf{Z}/p^n) \rightarrow H^{-m}(X, K; \mathbf{Z}/p^n) \xrightarrow{f^*} H^{-m}(Y, K; \mathbf{Z}/p^n) \rightarrow H^{-m+1}(C(X, Y), K; \mathbf{Z}/p^n) \rightarrow$$

Here $m \geq 3$. In the case X and Y are regular, this sequence is compatible with λ -operations.

4.1.4. γ -filtrations. The construction of the Loday product

$$\mathbf{Z}_\infty BGL_N(U) \wedge \mathbf{Z}_\infty BGL_N(U) \rightarrow \mathbf{Z}_\infty BGL_N(U)$$

is functorial in U . It induces products

$$[S^m X, K] \times [S^n X, K] \rightarrow [S^{n+m} X, K], \quad [P^m X, K] \times [P^n X, K] \rightarrow [P^{n+m} X, K]$$

for all K -coherent presheaves X .

For any K -coherent presheaf X , we have the following γ -filtrations compatible with products:

$$F_\gamma^k K_0(X) = \begin{cases} K_0(X) & \text{if } k \leq 0, \\ \langle \gamma_{i_1}(x_1) \cdots \gamma_{i_n}(x_n) | \varepsilon(x_1) = \dots = \varepsilon(x_n) = 0, i_1 + \dots + i_n \geq k \rangle & \text{if } k > 0, \end{cases}$$

$$F_\gamma^k K_q(X, \mathbf{Z}/p^n) = \langle \gamma_{i_1}(x_1) \cup \dots \cup \gamma_{i_n}(x_n) | x_i \in K_{q_i}(X, \mathbf{Z}/p^n), q_i \geq 2, \\ i_1 + \dots + i_n \geq k \rangle,$$

$$\mathcal{F}_\gamma^k K_q(X, \mathbf{Z}/p^n) = \langle a \gamma_{i_1}(x_1) \cup \dots \cup \gamma_{i_n}(x_n) | a \in F_\gamma^{i_0} K_0(X), x_i \in K_{q_i}(X, \mathbf{Z}/p^n), q_i \geq 2, \\ i_0 + i_1 + \dots + i_n \geq k \rangle,$$

where $p^n > 2$ and ε is the augmentation $\varepsilon : K_0(X) \rightarrow H^0(X, \mathbf{Z})$ obtained by projecting $\mathbf{Z} \times \mathbf{Z}_\infty BGL$ to \mathbf{Z} .

Lemma 4.3. *If X is a regular finite simplicial scheme over a field such that $X \simeq \text{sk}_m X$ then*

$$F_\gamma^{j+d+m+1} K_j(X, \mathbf{Z}/p^n) = 0, \quad j \geq 2,$$

where d is the maximum of dimensions of the schemes appearing in X .

Proof. First, we claim that $F_\gamma^q K_j(X, \mathbf{Z}/p^n) \subset F^{q-j} K_j(X, \mathbf{Z}/p^n)$, where the last filtration is obtained from the Brown spectral sequence (4.3). Since the Brown spectral sequence is multiplicative, it suffices to show that for any $N \geq 1$ we have

$$\gamma^q(H^{-j}(X, K^N; \mathbf{Z}/p^n)) \subset F^{q-j} H^{-j}(X, K; \mathbf{Z}/p^n), \quad j \geq 2.$$

Take $x \in H^{-j}(X, K^N; \mathbf{Z}/p^n)$ and consider the following diagram

$$\begin{array}{ccccccc} P^j X & \xrightarrow{x} & K^N & \xrightarrow{\gamma_N^q} & K & \longrightarrow & K \langle q-1 \rangle \\ \downarrow & & & & \downarrow & & \downarrow \\ S^j X & \xrightarrow{\gamma^q(x)} & K/p^n & \longrightarrow & K/p^n & \longrightarrow & K/p^n \langle q-1 \rangle, \end{array}$$

where K/p^n is the 0'th space of the spectrum K/p^n and, for a presheaf of simplicial sets F , we wrote

$$\{F \langle n \rangle\} := \dots \rightarrow F \langle m+1 \rangle \rightarrow F \langle m \rangle \rightarrow F \langle m-1 \rangle \rightarrow \dots$$

for a Postnikov tower of F that we chose to be functorial. The above diagram commutes (in the homotopy category): the right square commutes by functoriality; the left square commutes by S-duality [22, A.2]

$$[P^j X, K] \simeq [S^j X, K/p^n].$$

Now recall that we have

$$F^{q-j} H^{-j}(X, K; \mathbf{Z}/p^n) = \ker(H^{-j}(X, K/p^n) \rightarrow H^{-j}(X, K/p^n \langle q-1 \rangle)).$$

Hence it suffices to show that the bottom map in the above diagram is nullhomotopic. Since we have an injection

$$[S^j X, K/p^n] \hookrightarrow [P^j X, K/p^n]$$

it suffices to prove that the composition

$$\mathbf{Z}_\infty BGL_N \xrightarrow{\gamma_N^q} \mathbf{Z} \times \mathbf{Z}_\infty BGL_N \rightarrow K \rightarrow K \langle q-1 \rangle$$

is nullhomotopic. But this was done in [14, 5.1].

It remains now to show that

$$F^b K_j(X, \mathbf{Z}/p^n) = 0, \quad b > d + m.$$

First we claim that, for any $b > \text{cd } X$,

$$F^b K_j(X, \mathbf{Z}/p^n) = 0.$$

Indeed, we have

$$F^b H^{-j}(X, K; \mathbf{Z}/p^n) = \ker(H^{-j}(X, K/p^n) \rightarrow H^{-j}(X, K/p^n \langle b+j-1 \rangle)).$$

and the following map of Brown spectral sequences

$$\begin{array}{ccc} H^{-j}(X, K/p^n) & \longrightarrow & H^{-j}(X, K/p^n \langle b+j-1 \rangle) \\ \uparrow \parallel & & \uparrow \parallel \\ H^s(X, \tilde{\pi}_t(K/p^n)) & \longrightarrow & H^s(X, \tilde{\pi}_t(K/p^n \langle b+j-1 \rangle)) \end{array}$$

It follows that $F^b K_j(X, \mathbf{Z}/p^n) = 0$, $b > \text{cd } X$, where $\text{cd } X$ is the cohomological dimension of X . Next we show that $\text{cd } X \leq d + m$. For that, for any sheaf \mathcal{F} of abelian groups over X , it suffices to look at the spectral sequence (4.1)

$$E_1^{st} = H^{-t}(X_s, \mathcal{F}) \Rightarrow H^{s-t}(X, \mathcal{F})$$

and to remember that $X \simeq \text{sk}_m X$ and $\text{cd } X_s \leq d$. □

We will also consider another γ -filtration: $\widetilde{F}_\gamma^i = \langle \gamma^k(x) | k \geq i \rangle$, where $\langle \dots \rangle$ denotes the subgroup generated by the given elements. These filtrations are related. For a scheme X , by [35, 3.4] and Lemma 4.3, we have

$$(4.4) \quad M(d, i, 2j)F_\gamma^i K_j(X, \mathbf{Z}/p^n) \subset \widetilde{F}_\gamma^i K_j(X, \mathbf{Z}/p^n) \subset F_\gamma^i K_j(X, \mathbf{Z}/p^n), \quad j \geq 2,$$

where d is the dimension of X and the integers $M(k, m, n)$ are defined by the following procedure [35, 3.4]. Let l be a positive integer, and let w_l be the greatest common divisor of the set of integers $k^N(k^l - 1)$, as k runs over the positive integers and N is large enough with respect to l . Let $M(k)$ be the product of the w_l 's for $2l < k$. Set $M(k, m, n) = \prod_{2m \leq 2l \leq n+2k+1} M(2l)$. An odd prime p divides $M(d, i, j)$ if and only if $p < (j + 2d + 3)/2$, and divides $M(l)$ if and only if $p < (l/2) + 1$.

More generally we have the following lemma.

Lemma 4.4. *If X is a regular finite simplicial scheme over a field such that $X \simeq \text{sk}_m X$ then*

$$M(d + m, i, 2j)F_\gamma^i K_j(X, \mathbf{Z}/p^n) \subset \widetilde{F}_\gamma^i K_j(X, \mathbf{Z}/p^n) \subset F_\gamma^i K_j(X, \mathbf{Z}/p^n), \quad j \geq 2,$$

where d is the maximum of dimensions of the components of X .

Proof. Since, having Lemma 4.3, the argument in [35, 3.4] goes through almost verbatim (with $d + m$ in place of the cohomological dimension), we refer the interested reader for details to [35]. \square

For a bounded below complex of presheaves of abelian groups \mathcal{F} on the site C equipped with Zariski topology and a simplicial presheaf X on C , we set

$$H^n(X, \mathcal{F}) := H^{-n}(X, \mathcal{K}(\mathcal{F})), \quad n \geq 0,$$

where $\mathcal{K}(\mathcal{F}) := \mathcal{K}(\tau_{\leq 0}\mathcal{F})$ is the Dold-Puppe functor. If \mathcal{F} is built from injectif sheaves then $\mathcal{K}(\mathcal{F})$ is a fibrant presheaf of simplicial sets [14, 1.2.2] and, for X constructed from schemes, we have $H^n(X, \mathcal{F}) = H^{-n}\mathcal{K}(\mathcal{F})(X)$. In particular, if X is an object of the site C and \mathcal{F} is a bounded below complex of sheaves of abelian groups on C , then

$$H^n(X, \mathcal{F}) = H^n(X_{\text{Zar}}, \mathcal{F}).$$

4.2. Log-K-theory. In this section we collect basic facts about log- K -theory.

4.2.1. Definition of log- K -theory. For a scheme X , let $G(X)$ be the Thomason-Throbaugh spectrum of G -theory as defined in Thomason-Throbaugh [37, 3.3]. For a natural number $n \geq 2$, let $G/n(X)$ denote the corresponding spectrum mod n [37, 9.3]. After the usual rigidification [13, 5.1.3], which we assume from now on, these spectra are strictly covariant for noetherian schemes and proper maps as well as for quasi-compact schemes and pseudo-coherent projective morphisms [37, 3.16.1, 3.16.3].

Following Thomason [36, 5.15], we define functors from finite proper simplicial schemes to the category of prespectra

$$X \mapsto \mathbb{G}(X) := \text{hocolim}_i G(X_i), \quad X \mapsto \mathbb{G}/n(X) := \text{hocolim}_i G/n(X_i),$$

and set

$$G_i(X) := \pi_i(\mathbb{G}(X)), \quad G_i(X, \mathbf{Z}/n) := \pi_i(\mathbb{G}/n(X)).$$

Similarly, we have K -theory prespectra and groups.

Let X be a scheme and $i : D \hookrightarrow X$ a simple normal crossing divisor. Define the following prespectra

$$\begin{aligned} G(X(D)) &:= \text{cofiber}(\mathbb{G}(\widetilde{D}\bullet) \xrightarrow{i_*} G(X)), & G/n(X(D)) &:= \text{cofiber}(\mathbb{G}/n(\widetilde{D}\bullet) \xrightarrow{i_*} G/n(X)); \\ K(X(D)) &:= \text{cofiber}(\mathbb{K}(\widetilde{D}\bullet) \xrightarrow{i_*} K(X)), & K/n(X(D)) &:= \text{cofiber}(\mathbb{K}/n(\widetilde{D}\bullet) \xrightarrow{i_*} K/n(X)) \end{aligned}$$

Taking homotopy groups we get the log- G -theory and log- K -theory groups

$$\begin{aligned} G_i(X(D)) &:= \pi_i(G(X(D))), & G_i(X(D), \mathbf{Z}/n) &:= \pi_i(G/n(X(D))); \\ K_i(X(D)) &:= \pi_i(K(X(D))), & K_i(X(D), \mathbf{Z}/n) &:= \pi_i(K/n(X(D))). \end{aligned}$$

We have

$$\begin{aligned} G_j(X(D)) &= \pi_j(\text{cofiber}(\text{hocolim}_i G(\tilde{D}_i)_0 \xrightarrow{i_*} G(X)_0)), \quad j \geq 0; \\ G_j(X(D), \mathbf{Z}/n) &= \pi_j(\text{cofiber}(\text{hocolim}_i G(\tilde{D}_i)_0 \xrightarrow{i_*} G(X)_0), \mathbf{Z}/n), \quad j \geq 2. \end{aligned}$$

Recall that, for a noetherian scheme Y , there exists a homotopy equivalence $G(Y) \simeq G_Q(Y)$, where $G_Q(Y)$ is the Quillen G -theory spectrum of Y (see [37, 3.13]). Hence

$$G(Y)_0 \simeq G_Q(Y)_0 \simeq \Omega BQM(Y),$$

where $\Omega BQM(Y)$ is the loop space of the classifying space of Quillen Q -construction applied to the category of coherent sheaves on Y .

4.2.2. *Localization sequences.* We have the following localization sequence.

Lemma 4.5. *Let Y be a closed subscheme of X such that the scheme $D_Y = Y \cap D$ is a simple normal crossing divisor on Y . Then there is a homotopy cofibre sequence*

$$G(Y(D_Y)) \rightarrow G(X(D)) \rightarrow G(U(D_U)),$$

where $U = X \setminus Y$. In particular, we have the long exact sequence of homotopy groups

$$\rightarrow G_j(Y(D_Y)) \rightarrow G_j(X(D)) \rightarrow G_j(U(D_U)) \rightarrow G_{j-1}(Y(D_Y)) \rightarrow$$

Similarly for G -theory with mod- n coefficients.

Proof. For any $r \geq 0$, \tilde{D}_r is the disjoint union of the schemes D_σ , where σ runs over all maps $\sigma : \{1, \dots, r-1\} \rightarrow \{1, \dots, n\}$ and $D_\sigma = D_{Im(\sigma)}$. For any σ as above we have a homotopy cofibre sequence

$$G(Y \cap D_\sigma) \rightarrow G(D_\sigma) \rightarrow G(U \cap D_\sigma)$$

Since homotopy colimits preserve homotopy cofibre sequences we get the following commutative diagram of homotopy cofibre sequences

$$\begin{array}{ccccc} \text{hocolim}_r G(Y \cap \tilde{D}_r) & \longrightarrow & \text{hocolim}_r G(\tilde{D}_r) & \longrightarrow & \text{hocolim}_r G(U \cap \tilde{D}_r) \\ \downarrow & & \downarrow & & \downarrow \\ G(Y) & \longrightarrow & G(X) & \longrightarrow & G(U) \end{array}$$

Our lemma follows since taking homotopy cofibre preserves homotopy cofibre sequences. \square

The following lemma shows that log- G -theory equals G -theory of the open complement.

Lemma 4.6. (1) *The natural map $s : \tilde{D}_\bullet \rightarrow D$ induces homotopy equivalences*

$$s_* : \mathbb{G}(\tilde{D}_\bullet) \xrightarrow{\simeq} G(D), \quad s_* : \mathbb{G}/n(\tilde{D}_\bullet) \xrightarrow{\simeq} G/n(D).$$

(2) *The natural maps*

$$(4.5) \quad G(X(D)) \xrightarrow{\simeq} G(X \setminus D), \quad G/n(X(D)) \xrightarrow{\simeq} G/n(X \setminus D)$$

are homotopy equivalences.

(3) *If all the irreducible components of D are regular then the natural maps*

$$K(X(D)) \xrightarrow{\simeq} G(X(D)), \quad K/n(X(D)) \xrightarrow{\simeq} G/n(X(D))$$

are homotopy equivalences

Proof. For the first property we will argue by induction on the number m of irreducible components of D ; the case of $m = 1$ being clear. Write $D = \cup_{i=1}^m D_i$, $Y = D_1$, $U = X \setminus Y$, where each D_i is an irreducible component of D . We have the following diagram of maps of homotopy cofiber sequences

$$\begin{array}{ccccc}
\mathrm{hocolim}_r G(Y \cap \tilde{D}_r) & \longrightarrow & \mathrm{hocolim}_r G(\tilde{D}_r) & \longrightarrow & \mathrm{hocolim}_r G(U \cap \tilde{D}_r) \\
\uparrow \wr & & \uparrow \wr & & \uparrow \wr \\
\mathrm{hocolim}_r G(Y) & \longrightarrow & \mathbb{G}(\tilde{D}_\bullet) & \longrightarrow & \mathbb{G}(\tilde{D}_{U,\bullet}) \\
s_* \downarrow \wr & & s_* \downarrow & & s_* \downarrow \wr \\
G(Y) & \longrightarrow & G(D) & \longrightarrow & G(D_U)
\end{array}$$

The top sequence is obtained as in the proof of the previous lemma. The right lower map is a homotopy equivalence by the inductive assumption. The left lower map is induced by an augmentation of a constant simplicial scheme, hence a homotopy equivalence.

For the second property consider the following commutative diagram of prespectra

$$\begin{array}{ccccc}
\mathbb{G}(\tilde{D}_\bullet) & \xrightarrow{i_*} & G(X) & \longrightarrow & G(X(D)) \\
\wr \downarrow s_* & & \parallel & & \downarrow \wr \\
G(D) & \xrightarrow{i_*} & G(X) & \longrightarrow & G(X \setminus D),
\end{array}$$

where the bottom row is a homotopy cofiber sequence. Our property follows immediately. The third property follows from the fact that for a regular schemes Y the natural morphism $K(Y) \rightarrow G(Y)$ is a homotopy equivalence. \square

We get the following localization sequence for log- G -theory.

Corollary 4.7. *Let Y be one of the irreducible components of the divisor D . We have the following homotopy cofiber sequence*

$$G(Y(D_Y)) \xrightarrow{i_*} G(X(D')) \rightarrow G(X(D)),$$

where D' is the divisor D minus the component Y and $D_Y = D' \cap Y$.

Proof. This follows immediately from the above lemma since we have the following map of homotopy cofiber sequences

$$\begin{array}{ccccc}
G(Y(D_Y)) & \xrightarrow{i_*} & G(X(D')) & \longrightarrow & G(X(D)) \\
\downarrow \wr & & \downarrow \wr & & \downarrow \wr \\
G(Y \setminus D_Y) & \longrightarrow & G(X \setminus D') & \longrightarrow & G(X \setminus D).
\end{array}$$

\square

4.2.3. *Pairings, projection formula, base change.* Recall [13, 5.1.3] that the classical pairing of spectra [37, 3.15.3]

$$G(X) \wedge K(X) \rightarrow G(X)$$

after the rigidification of the G -theory and K -theory spectra that we have imposed makes the projection formula strict. That is, for any closed immersion $f : Y \rightarrow X$ the pushforward map $f_* : G(Y) \rightarrow G(X)$ is a map of module spectra over the ring spectrum $K(X)$ in the strict sense, i.e., the following projection

formula diagram of spectra is strictly commutative (not just commutative up to a homotopy)

$$\begin{array}{ccccc}
 & & K(Y) \wedge G(Y) & \xrightarrow{\wedge} & G(Y) \\
 & & \nearrow^{f^* \wedge 1} & & \downarrow f_* \\
 K(X) \wedge G(Y) & & & & \\
 & & \searrow_{1 \wedge f_*} & & \\
 & & K(X) \wedge G(X) & \xrightarrow{\wedge} & G(X)
 \end{array}$$

This allows us to state the following lemma.

Lemma 4.8. *We have pairings of prespectra*

$$G(X(D)) \wedge K(X) \rightarrow G(X(D)), \quad G(X(D)) \wedge K(X(D)) \rightarrow G(X(D))$$

that are compatible with the pairings

$$G(X \setminus D) \wedge K(X) \rightarrow G(X \setminus D), \quad G(X \setminus D) \wedge K(X \setminus D) \rightarrow G(X \setminus D).$$

Proof. First, using the above strict projection formula, define the pairing $\mathbb{G}(\tilde{D}\bullet) \wedge K(X) \rightarrow \mathbb{G}(\tilde{D}\bullet)$ by the formula

$$\begin{aligned}
 \mathbb{G}(\tilde{D}\bullet) \wedge K(X) &= (\text{hocolim}_n G(\tilde{D}_n)) \wedge K(X) \rightarrow \text{hocolim}_n (G(\tilde{D}_n) \wedge K(X)) \\
 &\xrightarrow{\wedge} \text{hocolim}_n G(\tilde{D}_n) = \mathbb{G}(\tilde{D}\bullet),
 \end{aligned}$$

where the first map is induced by the natural projection $\tilde{D}_n \rightarrow X$.

Similarly, set the pairing $G(X(D)) \wedge K(X) \rightarrow G(X(D))$ equal to the map

$$\begin{aligned}
 G(X(D)) \wedge K(X) &= \text{cofiber}(\mathbb{G}(\tilde{D}\bullet) \xrightarrow{i_*} K(X)) \wedge K(X) \rightarrow \text{cofiber}(\mathbb{G}(\tilde{D}\bullet) \wedge K(X) \xrightarrow{i_*} K(X) \wedge K(X)) \\
 &\xrightarrow{i_* \wedge} \text{cofiber}(\mathbb{G}(\tilde{D}\bullet) \xrightarrow{i_*} K(X)) = G(X(D)).
 \end{aligned}$$

The compatibility with the pairing $G(X \setminus D) \wedge K(X) \rightarrow G(X \setminus D)$ follows easily from the projection formula. \square

Let $f : Y \hookrightarrow X$ be a closed immersion of schemes with normal simple crossing divisors D_X, D_Y , such that $D_Y = f^{-1}D_X$. Assume that all the irreducible components of D_X, D_Y are regular. Then we have the following generalization of the projection formula.

Lemma 4.9. *The following diagram commutes up to canonically chosen homotopy.*

$$\begin{array}{ccccc}
 & & G(Y(D_Y)) \wedge K(Y) & \xrightarrow{\wedge} & G(Y(D_Y)) \\
 & & \nearrow^{f^* \wedge 1} & & \downarrow f_* \\
 G(X(D_X)) \wedge K(Y) & & & & \\
 & & \searrow_{1 \wedge f_*} & & \\
 & & G(X(D_X)) \wedge K(X) & \xrightarrow{\wedge} & G(X(D_X))
 \end{array}$$

Proof. The easiest way to see this is to use the natural maps $G(Y(D_Y)) \xrightarrow{\sim} K(Y \setminus D_Y)$ and $G(Y(D_Y)) \xrightarrow{\sim} K(X \setminus D_X)$ to pass to the classical diagram that we know commutes up to canonically chosen homotopy

[37, 3.17].

$$\begin{array}{ccc}
 & K(Y \setminus D_Y) \wedge K(Y) & \xrightarrow{\wedge} K(Y \setminus D_Y) \\
 & \nearrow^{f^* \wedge 1} & \downarrow f_* \\
 K(X \setminus D_X) \wedge K(Y) & & \\
 & \searrow_{1 \wedge f_*} & \\
 & K(X \setminus D_X) \wedge K(X) & \xrightarrow{\wedge} K(X \setminus D_X)
 \end{array}$$

□

Lemma 4.10. *Consider a pullback diagram of regular log-schemes*

$$\begin{array}{ccc}
 Y & \xrightarrow{i} & X \\
 \downarrow j & & \downarrow j' \\
 Y' & \xrightarrow{i'} & X'
 \end{array}$$

where all the maps are regular closed immersions of codimension one and the log-structures are induced by simple normal crossing divisors $D_{X'}$, $D_{Y'}$, D_X , D_Y such that

$$D_{Y'} = (i')^{-1}(D_{X'}), \quad D_Y = j^{-1}D_{Y'}, \quad D_X = (j')^{-1}D_{X'}.$$

Then there is a canonical homotopy between

$$j'^* i'_* \simeq i_* j^* : K(Y'(D_{Y'})) \rightarrow K(X(D_Y)).$$

Proof. Again, pass to the maps

$$j'^* i'_* \simeq i_* j^* : K(Y' \setminus D_{Y'}) \rightarrow K(X \setminus D_Y).$$

That there exists a canonical homotopy between these maps is a classical result [37, 3.18] that follows from Tor-independence of i' and j' . □

4.3. Operations on log- K -theory. In this section we will study the behaviour of certain localization sequences in K -theory under λ -operations.

4.3.1. Adams-Riemann-Roch without denominators. We start with the proof of the Adams-Riemann-Roch without denominators. Let τ be a natural transformation of λ -rings such that $\tau(0) = 0$. Let X be a regular scheme and $i : Y \hookrightarrow X$ a regular divisor. Assume that both schemes are defined over \mathcal{O}_K . Set $U = X \setminus Y$, $j : U \hookrightarrow X$, and

$$K_m^Y(X, \mathbf{Z}/p^n) := H^{-m}(C(X, U), K; \mathbf{Z}/p^n), \quad m \geq 2.$$

For $m \geq 3$, by Example 4.2, the long exact sequence

$$\rightarrow K_m^Y(X, \mathbf{Z}/p^n) \rightarrow K_m(X, \mathbf{Z}/p^n) \rightarrow K_m(U, \mathbf{Z}/p^n) \rightarrow K_{m-1}^Y(X, \mathbf{Z}/p^n) \rightarrow$$

is compatible with the action of τ .

We have a natural isomorphism $i_* : K_m(Y, \mathbf{Z}/p^n) \xrightarrow{\sim} K_m^Y(X, \mathbf{Z}/p^n)$, $m \geq 2$. We need to understand how it behaves with respect to operations. Recall that the group $K_m(Y, \mathbf{Z}/p^n)$ is a $K_0(Y)$ - λ -algebra, i.e., the sum $K_0(Y) \oplus K_m(Y, \mathbf{Z}/p^n)$ is a λ -ring. As in any λ -ring, for every element $x \in K_m(Y, \mathbf{Z}/p^n)$, there exists an element $\tau(Y/X, x) \in K_m(Y, \mathbf{Z}/p^n)$ (cf., [21, 1.1.1]) which is a universal polynomial with integral coefficients in $\lambda(\mathcal{N}_{Y/X}^\vee)$ and $\lambda(x)$, where $\mathcal{N}_{Y/X}^\vee$ is the conormal sheaf of Y in X , such that

$$(4.6) \quad \tau_Y(\lambda_{-1}([\mathcal{N}_{Y/X}^\vee])x) = \lambda_{-1}([\mathcal{N}_{Y/X}^\vee])\tau(Y/X, x).$$

Lemma 4.11. (Adams-Riemann-Roch without denominators) *The diagram*

$$\begin{array}{ccc} K_m(Y, \mathbf{Z}/p^n) & \xrightarrow{\tau(Y/X, \bullet)} & K_m(Y, \mathbf{Z}/p^n) \\ \downarrow i_* & & \downarrow i_* \\ K_m^Y(X, \mathbf{Z}/p^n) & \xrightarrow{\tau_X} & K_m^Y(X, \mathbf{Z}/p^n) \end{array}$$

commutes, i.e.,

$$\tau_X(i_*(y)) = i_*(\tau(Y/X, y)), \quad y \in K_m(Y, \mathbf{Z}/p^n)$$

Proof. Like in the proof of Lemma 6.9 we argue by deformation to the normal cone. Since many computations here are very similar to the ones done in the proof of that lemma we will just describe those that are substantially different.

So we start with a special case of a closed immersion $i : Y \hookrightarrow X$, which is the zero section of a projective bundle $X = \mathbb{P}(\mathcal{N} \oplus \mathcal{O}_Y)$, where \mathcal{N} is an invertible sheaf on Y . The sheaf \mathcal{N} is the conormal sheaf of the immersion. Let $\pi : X \rightarrow Y$ be the projection. Notice that the natural map $K_m^Y(X, \mathbf{Z}/p^n) \rightarrow K_m(X, \mathbf{Z}/p^n)$ is an injection. Hence it suffices to prove the commutativity of the following diagram

$$\begin{array}{ccc} K_m(Y, \mathbf{Z}/p^n) & \xrightarrow{\tau(Y/X, \bullet)} & K_m(Y, \mathbf{Z}/p^n) \\ \downarrow i_* & & \downarrow i_* \\ K_m(X, \mathbf{Z}/p^n) & \xrightarrow{\tau_X} & K_m(X, \mathbf{Z}/p^n) \end{array}$$

Take $y \in K_m(Y, \mathbf{Z}/p^n)$. As in the proof of Lemma 6.9 we compute that there is an equality $i_*(y) = \pi^*(\lambda_{-1}([\mathcal{N}])y)$. From the functoriality of operations it follows that

$$\tau_X i_*(y) = \pi^* \tau_Y([\lambda_{-1}(\mathcal{N})]y).$$

The equality (4.6), the equality $\lambda_{-1}(\pi^*[\mathcal{N}]) = [i_*\mathcal{O}_Y]$ in $K_0(X)$ that follows from the exact sequence (6.13), and the projection formula in K -theory imply now that

$$\begin{aligned} \tau_X i_*(y) &= \pi^*(\tau(Y/X, y)\lambda_{-1}([\mathcal{N}])) = \pi^*(\tau(Y/X, y))\lambda_{-1}(\pi^*[\mathcal{N}]) \\ &= \pi^*(\tau(Y/X, y))[i_*\mathcal{O}_Y] = i_* i^* \pi^*(\tau(Y/X, y)) = i_* \tau(Y/X, y), \end{aligned}$$

as wanted.

In general, we use deformation to the normal cone. We will use freely the notation from the proof of Lemma 6.9. As before, by functoriality of K -theory and [37, 3.18], we find that

$$j_{0*} i_* \tau(Y_0/W_0, y) = j_{0*} j_0^* \bar{i}_* \rho^* \tau(Y_0/W_0, y),$$

where

$$\bar{i}_* : K_m(\mathbb{P}_Y^1, \mathbf{Z}/p^n) \rightarrow K_m^{\mathbb{P}_Y^1}(W, \mathbf{Z}/p^n), \quad j_0^* : K_m^{\mathbb{P}_Y^1}(W, \mathbf{Z}/p^n) \rightarrow K_m^Y(X, \mathbf{Z}/p^n).$$

The projection formula in K -theory from [37, 3.17] and functoriality of K -theory and operations imply now that

$$j_{0*} i_* \tau(Y_0/W_0, y) = \bar{i}_* \tau(\mathbb{P}_Y^1/W, y) \wedge ([t_*\mathcal{O}_X] + [l_*\mathcal{O}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}]).$$

Functoriality, the projection formula in K -theory, the fact that $l^* \bar{i}_* = 0$, and [37, 3.18] imply that

$$j_{0*} i_* \tau(Y_0/W_0, y) = t_* i_{\infty*} \tau(Y/\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y), y).$$

Here

$$i_{\infty*} : K_m(Y, \mathbf{Z}/p^n) \rightarrow K_m^Y(\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y), \mathbf{Z}/p^n), \quad t_* : K_m^Y(\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y), \mathbf{Z}/p^n) \rightarrow K_m^{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}(W, \mathbf{Z}/p^n).$$

Now we apply the computation we did in the special case to the embedding $i_\infty : Y \rightarrow \mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$ to conclude that

$$j_{0*} i_* \tau(Y_0/W_0, y) = t_* \tau_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)} i_{\infty*}(y).$$

Similarly, in Lemma 6.9 we computed that $i_*(y) = j_0^* \bar{i}_* \rho^*(y)$. This, functoriality of K -theory and operations, the projection formula in K -theory, the fact that $l^* \bar{i}_* = 0$, and [37, 3.18] imply that

$$j_{0*} \tau_X i_*(y) = t_* \tau_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)} i_{\infty*}(y).$$

Hence we have that $j_{0*} i_* \tau(Y_0/W_0, y) = j_{0*} \tau_X i_*(y)$.

It suffices now to show that the map

$$j_{0*} : K_m^Y(X, \mathbf{Z}/p^n) \rightarrow K_m^{\mathbb{P}^Y}(W, \mathbf{Z}/p^n)$$

is injective. But this map is isomorphic to the map $j_{0*} : K_m(Y, \mathbf{Z}/p^n) \rightarrow K_m(\mathbb{P}^Y, \mathbf{Z}/p^n)$ that is injective by the projective spaces theorem in K -theory. \square

4.3.2. *γ -filtration and localization sequences.* We need to understand the behaviour of the integral γ -filtrations in localization sequences. We will need the following lemma.

Lemma 4.12. *Let X be a regular scheme over \mathcal{O}_K of dimension d . Then*

- (1) $F_\gamma^{d+j+1} K_j(X, \mathbf{Z}/p^n) = 0$;
- (2) $M(d, i, 2j) F_\gamma^i K_j(X, \mathbf{Z}/p^n) \subset \widetilde{F}_\gamma^i K_j(X, \mathbf{Z}/p^n) \subset F_\gamma^i K_j(X, \mathbf{Z}/p^n)$.

Proof. The property (1) is proved in [26, 14.5]. It is stated there just for K -theory with no coefficients but it holds as well for K -theory modulo p^n . All one needs to do is to paraphrase Corollary 13.11 to the case with coefficients which is immediate. The property (2) follows now as in [35, 3.4]. \square

Consider a triple of regular schemes

$$U \xrightarrow{j} X \xleftarrow{i} Z$$

with Z a closed subscheme of X of codimension one and U its complement. For $j \geq 3$, we have the localization sequence in K -theory

$$(4.7) \quad \rightarrow K_j(Z, \mathbf{Z}/p^n) \xrightarrow{i_*} K_j(X, \mathbf{Z}/p^n) \xrightarrow{j^*} K_j(U, \mathbf{Z}/p^n) \xrightarrow{\partial} K_{j-1}(Z, \mathbf{Z}/p^n) \rightarrow$$

obtained from the long exact sequence

$$\rightarrow K_j^Z(X, \mathbf{Z}/p^n) \rightarrow K_j(X, \mathbf{Z}/p^n) \xrightarrow{j^*} K_j(U, \mathbf{Z}/p^n) \rightarrow K_{j-1}^Z(X, \mathbf{Z}/p^n) \rightarrow$$

and the isomorphism $i_* : K_j(Z, \mathbf{Z}/p^n) \xrightarrow{\sim} K_j^Z(X, \mathbf{Z}/p^n)$.

We claim that, modulo certain universal constants, this sequence of maps behaves well with respect to the γ -filtration.

Lemma 4.13. *Let $j \geq 3$. There exists a natural number $N = N(d, i, j)$ dependent only on d, i, j , where d is the dimension of X , such that we have a long sequence of maps*

$$\rightarrow F_\gamma^{i-1} K_j(Z, \mathbf{Z}/p^n) \xrightarrow{N i_*} F_\gamma^i K_j(X, \mathbf{Z}/p^n) \xrightarrow{N j^*} F_\gamma^i K_j(U, \mathbf{Z}/p^n) \xrightarrow{N \partial} F_\gamma^{i-1} K_{j-1}(Z, \mathbf{Z}/p^n) \rightarrow .$$

If a prime number $p > j + d + 1$ then p does not divide $N(d, i, j)$.

Proof. By functoriality, the restriction map j^* is compatible with γ -filtrations. So is the boundary map ∂ with \widetilde{F}_γ^i -filtration, i.e., $\partial : \widetilde{F}_\gamma^i K_j(U, \mathbf{Z}/p^n) \rightarrow F_\gamma^{i-1} K_{j-1}(Z, \mathbf{Z}/p^n)$. To see that take an element $x = \gamma^k(y) \in \widetilde{F}_\gamma^i K_j(U, \mathbf{Z}/p^n)$ for $y \in K_j(U, \mathbf{Z}/p^n)$, $k \geq i$. Then, by the Adams-Riemann-Roch without denominators (Proposition 4.11) we have that $\partial x = \gamma^k(\mathcal{N}, x)(\partial y)$, for the normal bundle $\mathcal{N} = \mathcal{N}_{Z/X}$ of Z in X . But, for every $z \in K_{j-1}(Z, \mathbf{Z}/p^n)$, we have $\gamma^k(x)(\mathcal{N}, z) \in F_\gamma^{k-1} K_{j-1}(Z, \mathbf{Z}/p^n)$ [16, 0, Appendice, Prop. 1.5]. Since by Lemma 4.12 $M(d, i, 2j) F_\gamma^i K_j(U, \mathbf{Z}/p^n) \subset \widetilde{F}_\gamma^i K_j(U, \mathbf{Z}/p^n)$ this implies that

$$M(d, i, 2j) \partial : F_\gamma^i K_j(U, \mathbf{Z}/p^n) \rightarrow F_\gamma^{i-1} K_{j-1}(Z, \mathbf{Z}/p^n).$$

Consider now $x \in F_\gamma^{i-1} K_j(Z, \mathbf{Z}/p^n)$. We know that $\gamma^i(\mathcal{N}, x) = (-1)^{i-1} (i-1)! x \pmod{F_\gamma^i K_j(Z, \mathbf{Z}/p^n)}$. By Adams-Riemann-Roch without denominators from Proposition 4.11 we have

$$i_*((-1)^{i-1} (i-1)! x) = \gamma^i(i_* x) \pmod{i_*(F_\gamma^i K_j(Z, \mathbf{Z}/p^n))}.$$

By induction, since $F_\gamma^{j+d}K_j(Z, \mathbf{Z}/p^n) = 0$, we get that

$$C(d, i, j)_{i_*} : F_\gamma^{i-1}K_j(Z, \mathbf{Z}/p^n) \rightarrow F_\gamma^i K_j(X, \mathbf{Z}/p^n), \quad C(d, i, j) = (i-1)!i! \cdots (j+d-1)!$$

Set $N(d, i, j) = M(d, i, 2j)C(d, i, j)$. Since an odd prime number p divides $M(d, i, j)$ if and only if $p < (j+2d+3)/2$ we get the last statement of the lemma. \square

Lemma 4.14. *let $j \geq 3$, $N = N(d, i-1, j)N(d, i, j)$ for the constants $N(d, i-1, j)$ and $N(d, i, j)$ from Lemma 4.13. Then the following long sequence is exact up to certain universal constants*

$$\rightarrow F_\gamma^{i-1}/F_\gamma^i K_j(Z, \mathbf{Z}/p^n) \xrightarrow{N_{i_*}} F_\gamma^i/F_\gamma^{i+1} K_j(X, \mathbf{Z}/p^n) \xrightarrow{N_{j^*}} F_\gamma^i/F_\gamma^{i+1} K_j(U, \mathbf{Z}/p^n) \xrightarrow{N\partial} F_\gamma^{i-1}/F_\gamma^i K_{j-1}(Z, \mathbf{Z}/p^n) \rightarrow,$$

where $N = N(d, i-1, j)N(d, i, j)$ for the constants from Lemma 4.13. More precisely, if the element $[x]$ at any level of the above long sequence is a cocycle then $C[x]$ is a coboundary for the following constant C

- (1) $N_{i_*}([x]) = 0$ then $C = (i-1)!i! \cdots (j+d-1)!N(d, i, j)^2 N(d, i, j+1)^2$;
- (2) $N\partial([x]) = 0$ then $C = (i-1)!(i-1)!i! \cdots (j+d-3)!N^2$;
- (3) $N_{j^*}([x]) = 0$ then $C = (i-1)!i! \cdots (j+d-1)!N^2$.

Proof. For the first case, consider an element $[x] \in F_\gamma^{i-1}/F_\gamma^i K_j(Z, \mathbf{Z}/p^n)$ for $x \in F_\gamma^{i-1}K_j(Z, \mathbf{Z}/p^n)$ such that $N_{i_*}[x] = 0$, i.e., $N_{i_*}(x) \in F_\gamma^{i+1}K_j(X, \mathbf{Z}/p^n)$. By Adams-Riemann-Roch without denominators from Lemma 4.11 we have

$$i_*\gamma^{i+1}(\mathcal{N}, Nx) = \gamma^{i+1}(i_*(Nx)) = (-1)^{i!}i!i_*(Nx) \quad \text{mod } F_\gamma^{i+2}K_j(X, \mathbf{Z}/p^n).$$

Since $\gamma^{i+1}(\mathcal{N}, Nx) \in F_\gamma^i K_j(Z, \mathbf{Z}/p^n)$, we get

$$[(-1)^{i!}i!Nx] = [(-1)^{i!}i!Nx - \gamma^{i+1}(\mathcal{N}, Nx)] \quad \text{and} \quad i_*((-1)^{i!}i!Nx - \gamma^{i+1}(\mathcal{N}, Nx)) \in F_\gamma^{i+2}K_j(X, \mathbf{Z}/p^n)$$

Since $F_\gamma^{j+d+1}K_j(X, \mathbf{Z}/p^n) = 0$, by induction, we get that $[i!(i+1)! \cdots (j+d-1)!Nx] = [y]$ for $y \in F_\gamma^{i-1}K_j(Z, \mathbf{Z}/p^n)$ such that $i_*(y) = 0$. By the localization sequence (4.7) $y = \partial(w)$ for $w \in K_{j+1}(U, \mathbf{Z}/p^n)$. We also have $\gamma^i(\mathcal{N}, y) = (-1)^{i-1}(i-1)!y$ modulo $F_\gamma^i K_j(Z, \mathbf{Z}/p^n)$ [16, 0, Appendice, Prop. 1.5]. From Adams-Riemann-Roch without denominators it now follows that

$$[\gamma^i(\mathcal{N}, y)] = [\gamma^i(\mathcal{N}, \partial(w))] = [\partial(\gamma^i(w))].$$

Since clearly $\gamma^i(w) \in F_\gamma^i K_{j+1}(U, \mathbf{Z}/p^n)$, we have that $[(i-1)!y]$ is a coboundary for ∂ . Summing up the above, $[(i-1)!i! \cdots (j+d-1)!N^2(d, i, j)N^2(d, i, j+1)]$ is a coboundary, as wanted.

For the second case, consider an element $[x] \in F_\gamma^i/F_\gamma^{i+1}K_j(U, \mathbf{Z}/p^n)$ for $x \in F_\gamma^i K_j(U, \mathbf{Z}/p^n)$ such that $N\partial[x] = 0$, i.e., $N\partial(x) \in F_\gamma^i K_{j-1}(Z, \mathbf{Z}/p^n)$. As above, Adams-Riemann-Roch without denominators implies that there exists an element $w \in F_\gamma^{i+1}K_j(U, \mathbf{Z}/p^n)$ such that $i!\partial(Nx) = \partial(w)$ modulo $F_\gamma^i K_j(Z, \mathbf{Z}/p^n)$. Since $F_\gamma^{j+d-1}K_{j-1}(Z, \mathbf{Z}/p^n)$, this gives by induction that

$$i! \cdots (j+d-3)!\partial(Nx) = \partial(w), \quad \text{for } w \in F_\gamma^{i+1}K_j(U, \mathbf{Z}/p^n).$$

Set $z = i! \cdots (j+d-3)!\partial(Nx) - w$. We have $[i! \cdots (j+d-3)!\partial(Nx)] = [z]$ and $\partial(z) = 0$. By the localization sequence (4.7) $z = j^*(y)$ for $y \in K_j(X, \mathbf{Z}/p^n)$. We have $[\gamma^i(z)] = [(-1)^{i-1}(i-1)!z]$. Hence $[(-1)^{i-1}(i-1)!z] = [j^*(\gamma^i(y))]$, i.e., $[(-1)^{i-1}(i-1)!z]$ is in the image of $F_\gamma^i K_j(X, \mathbf{Z}/p^n)$ by j^* . Summing it all up, we get that $[(i-1)!i! \cdots (j+d-3)!N^2x]$ is a coboundary, as wanted.

Because the restriction j^* is compatible with γ -operations, the third case is proved in a similar manner. Consider an element $[x] \in F_\gamma^i/F_\gamma^{i+1}K_j(X, \mathbf{Z}/p^n)$ for $x \in F_\gamma^i K_j(X, \mathbf{Z}/p^n)$ such that $N_{j^*}[x] = 0$, i.e., $N_{j^*}(x) \in F_\gamma^{i+1}K_j(U, \mathbf{Z}/p^n)$. We have

$$(-1)^{i!}i!N_{j^*}(x) = \gamma^{i+1}(j^*(Nx)) = j^*(\gamma^{i+1}(Nx)) \quad \text{mod } F_\gamma^{i+2}K_j(U, \mathbf{Z}/p^n).$$

Hence $[Nx] = [Nx - \gamma^{i+1}(Nx)]$ and $j^*(Nx - \gamma^{i+1}(Nx)) \in F_\gamma^{i+2}K_j(U, \mathbf{Z}/p^n)$. Since $F_\gamma^{j+d+1}K_j(U, \mathbf{Z}/p^n)$, inductively this implies that $[i!(i+1)! \cdots (j+d-1)!Nx] = [z]$. and $j^*(z) = 0$. Now, by the localization sequence (4.7), $z = i_*(y)$ for $y \in K_j(Z, \mathbf{Z}/p^n)$. By Adams-Riemann-Roch without denominators we have

$$[i_*\gamma^i(\mathcal{N}, y)] = [\gamma^i(i_*y)] = [\gamma^i(z)] = [(-1)^i(i-1)!z].$$

Hence $[(-1)^i(i-1)!z]$ is a coboundary for i_* . Summing up, we get that $[(i-1)!i! \cdots (j+d-1)!N^2x]$ is a coboundary, as wanted. \square

5. SYNTOMIC CHERN CLASSES

5.1. Classical syntomic Chern classes. We will briefly review the construction and basic properties of classical syntomic Chern classes. For details the reader can consult [28, 2.3], [30, 2.2], and [31, 2.3].

For $i \geq 0$ and a scheme X flat over $W(k)$, there are functorial and compatible families of syntomic Chern classes

$$\begin{aligned} c_{i,j}^{\text{syn}} &: K_j(X) \rightarrow H^{2i-j}(X, S_n(i)) \quad \text{for } j \geq 0, \\ \bar{c}_{i,j}^{\text{syn}} &: K_j(X, \mathbf{Z}/p^n) \rightarrow H^{2i-j}(X, S_n(i)) \quad \text{for } j \geq 2, \end{aligned}$$

that are also compatible with the crystalline Chern classes in $H_{\text{cr}}^{2i-j}(X_n, \mathcal{O}_{X_n})$ via the canonical map $H^{2i-j}(X_n, S_n(i)) \rightarrow H_{\text{cr}}^{2i-j}(X_n, J_{X_n}^{\leq i})$. Similarly, we have syntomic Chern classes in $H^{2i-j}(X, S'_n(i))$.

Recall the construction of the classes $c_{i,j}^{\text{syn}}$ and $\bar{c}_{i,j}^{\text{syn}}$. First, one constructs universal classes $C_{i,l}^{\text{syn}} \in H^{2i}(BGL_l, S_n(i))$, $C_{i,l}^{\text{syn}} \in H^{2i}(BGL_l, S'_n(i))$. For $l \geq i$, $i \geq 0$, one defines

$$C_{i,l}^{\text{syn}} = x_i \in H^{2i}(BGL_l, S_n(i)), \quad C_{i,l}^{\text{syn}} = x_i \in H^{2i}(BGL_l, S'_n(i)).$$

By construction these classes are compatible with the crystalline classes. Recall (see formulas (2.3)) that classes $x_1 \in H^2(BGL_l, S_n(1)) \hookrightarrow H^2(BGL_l, S'_n(1))$ have a direct definition via symbol maps.

The classes $C_{i,l}^{\text{syn}} \in H^{2i}(BGL_l, S_n(i))$, $i \geq 0$, yield compatible universal classes (see [12, p. 221]) $C_{i,l}^{\text{syn}} \in H^{2i}(X, GL_l(\mathcal{O}_X), S_n(i))$, where the last group is the $GL_l(\mathcal{O}_X)$ -cohomology of X with values in $S_n(i)$ (equipped with the trivial action of $GL_l(\mathcal{O}_X)$). Hence a natural map of pointed simplicial presheaves on X ,

$$C_i^{\text{syn}} : BGL(\mathcal{O}_X) \rightarrow \mathcal{K}(2i, \tilde{S}_n(i)_X),$$

where \mathcal{K} is the Dold–Puppe functor of $\tau_{\leq 0} \tilde{S}_n(i)_X[2i]$ and $\tilde{S}_n(i)_X$ is an injective resolution of $S_n(i)_X$. These classes induce Chern class maps

$$C_i^{\text{syn}} : K_X \rightarrow \mathcal{K}(2i, \tilde{S}_n(i)_X), \quad i \geq 0.$$

Here we wrote K_X for the presheaf of simplicial sets $U \mapsto K(U)_0$.

We can now define the total Chern class map

$$C_X^{\text{syn}} : K_X \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_X)$$

by putting the map C_i^{syn} in degree i . Similarly, we get total Chern class maps

$$(5.1) \quad C_X^{\text{syn}} : K_X \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}'_n(i)_X).$$

Notice that the Chern class map C_0^{syn} is a composition of the rank map $rk : K_X \rightarrow \mathbf{Z}$ with the natural map $\mathbf{Z} \rightarrow S_n(0) = S'_n(0) = (\mathcal{O}_n^{\text{cr}})^{\varphi=1}$. The related augmented total Chern class maps \tilde{C}_X^{syn} are defined by replacing C_0^{syn} with the rank map $rk : K_X \rightarrow \mathbf{Z}$.

We list the following properties of the total Chern class maps. They are proved using the embedding $H^{2i}(BGL_l, S_n(i)) \hookrightarrow H_{\text{dR}}^{2i}(BGL_{l,n})$ and the properties of de Rham classes.

Lemma 5.1. ([28, Lemma 2.1])

- (1) *The syntomic total Chern class is functorial, i.e., for a map $f : Y \rightarrow X$ of flat schemes over $W(k)$ the following diagram of presheaves of simplicial sets on X commutes in the homotopy*

category.

$$\begin{array}{ccc} K_X & \xrightarrow{C_X^{\text{syn}}} & \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_X) \\ \downarrow f^* & & \downarrow f^* \\ \mathbf{R}f_* K_Y & \xrightarrow{C_Y^{\text{syn}}} & \mathbf{R}f_* \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_Y) \end{array}$$

- (2) The syntomic total Chern class is compatible with addition, i.e., the following diagram is commutes in the homotopy category.

$$\begin{array}{ccc} K_X \times K_X & \xrightarrow{+} & K_X \\ \downarrow C_X^{\text{syn}} \times C_X^{\text{syn}} & & \downarrow C_X^{\text{syn}} \\ \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_X) \times \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_X) & \xrightarrow{\star} & \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_X). \end{array}$$

- (3) The syntomic augmented total Chern class is compatible with products, i.e., the following diagram commutes in the homotopy category.

$$\begin{array}{ccc} K_X \wedge K_X & \xrightarrow{\wedge} & K_X \\ \downarrow \tilde{C}_X^{\text{syn}} \wedge \tilde{C}_X^{\text{syn}} & & \downarrow \tilde{C}_X^{\text{syn}} \\ \mathbf{Z} \times \prod_{i \geq 1} \mathcal{K}(2i, \tilde{S}_n(i)_X) \wedge \mathbf{Z} \times \prod_{i \geq 1} \mathcal{K}(2i, \tilde{S}_n(i)_X) & \xrightarrow{\star} & \mathbf{Z} \times \prod_{i \geq 1} \mathcal{K}(2i, \tilde{S}_n(i)_X). \end{array}$$

Here \star is the Grothendieck product [12, 2.27] defined via certain universal polynomials with integral coefficients.

Similarly for total Chern classes with values in $S'_n(i)$ -cohomology.

Lemma 5.2. *The augmented total Chern class*

$$\tilde{C}_X^{\text{syn}} : K_0(X) \rightarrow \mathbf{Z} \times \{1\} \times \prod_{i \geq 1} H^{2i}(X, S_n(i))$$

is a morphism of λ -rings.

Proof. This was shown in the proof of Lemma 2.1 in [28]. □

The characteristic classes

$$\begin{aligned} \tilde{c}_{i,j}^{\text{syn}} &: K_j(X, \mathbf{Z}/p^n) \rightarrow H^{2i-j}(X, S_n(i)), \quad j \geq 2, \\ c_{i,j}^{\text{syn}} &: K_j(X) \rightarrow H^{2i-j}(X, S_n(i)), \quad j \geq 0, \end{aligned}$$

are defined [12, 2.22] as the composition

$$\begin{aligned} K_j(X, \mathbf{Z}/p^n) &\rightarrow H^{-j}(X, K_X; \mathbf{Z}/p^n) \rightarrow H^{-j}(X, BGL(\mathcal{O}_X)^+; \mathbf{Z}/p^n) \\ &\xrightarrow{C_i^{\text{syn}}} H^{-j}(X, \mathcal{K}(2i, \tilde{S}_n(i)_X); \mathbf{Z}/p^n) \xrightarrow{f} H^{2i-j}(X, S_n(i)), \end{aligned}$$

where $BGL(\mathcal{O}_X)^+$ is the (pointed) simplicial presheaf on X associated to the $+$ -construction. The map f is defined as the composition

$$\begin{aligned} H^{-j}(X, \mathcal{K}(2i, \tilde{S}_n(i)_X); \mathbf{Z}/p^n) &= \pi_j(\mathcal{K}(2i, \tilde{S}_n(i)(X)), \mathbf{Z}/p^n) \xrightarrow{h_j} H_j(\mathcal{K}(2i, \tilde{S}_n(i)(X)), \mathbf{Z}/p^n) \\ &\rightarrow H_j(\tilde{S}_n(i)(X)[2i]) = H^{2i-j}(X, S_n(i)), \end{aligned}$$

where h_j is the Hurewicz morphism.

This gives mod p^n Chern classes with values in $H^*(X, S_n(*))$. Those with values in $H^*(X, S'_n(*))$ and the ones from integral K -theory are defined in an analogous way. All of the above also works (with basically identical proofs) if we replace X with a finite simplicial scheme X , flat over $W(k)$.

As we have shown in [30, Lemma 2.1] the above Chern classes for schemes have the properties listed below. Again they hold as well for finite simplicial schemes X , flat over $W(k)$.

Lemma 5.3. *Let X be a finite simplicial scheme that is flat over $W(k)$. The syntomic Chern classes are functorial in X and have the following properties.*

- (1) c_{ij}^{syn} , for $j > 0$, is a group homomorphism.
- (2) $c_{\bullet,0}^{\text{syn}}(a+b) = c_{\bullet,0}^{\text{syn}}(a)c_{\bullet,0}^{\text{syn}}(b)$, for $a, b \in K_0(X)$.
- (3) $\bar{c}_{ij}^{\text{syn}}$, for $j \geq 2$ is a group homomorphism unless $j = 2$ and $p = 2$.
- (4) $\bar{c}_{ij}^{\text{syn}}$ are compatible with the reduction maps $S_n(i) \rightarrow S_m(i)$, $n \geq m$.

Moreover, if X is regular, then

- (5) Let p be odd or $p = 2$, $n \geq 2$ and $l, q \neq 2$. If $\alpha \in K_l(X, \mathbf{Z}/p^n)$ and $\alpha' \in K_q(X, \mathbf{Z}/p^n)$, then

$$\bar{c}_{ij}^{\text{syn}}(\alpha\alpha') = - \sum_{r+s=i} \frac{(i-1)!}{(r-1)!(s-1)!} \bar{c}_{rl}^{\text{syn}}(\alpha) \bar{c}_{sq}^{\text{syn}}(\alpha'),$$

assuming that $l, q \geq 2$, $l+q = j$, $2i \geq j$, $i \geq 0$.

- (6) If $\alpha \in F_\gamma^j K_0(X)$, $j \neq 0$, and $\alpha' \in F_\gamma^k K_q(X, \mathbf{Z}/p^n)$, $q \geq 2$, is such that $\bar{c}_{lq}^{\text{syn}}(\alpha') = 0$ for $l \neq k$, then

$$\bar{c}_{j+k,q}^{\text{syn}}(\alpha\alpha') = - \frac{(j+k-1)!}{(j-1)!(k-1)!} c_{j0}^{\text{syn}}(\alpha) \bar{c}_{kq}^{\text{syn}}(\alpha'),$$

assuming that $p \neq 2$ or $q > 2$.

- (7) If X is a scheme, the above multiplication formulas hold also for $p = 2$, $n \geq 4$, $q = 2$ and α' such that $\partial\alpha' \in K_1(X)$ belongs to \mathcal{O}_K^* .
- (8) The integral Chern class maps c_{i0}^{syn} restrict to zero on $F_\gamma^{i+1} K_0(X)$.
- (9) The Chern class maps $\bar{c}_{ij}^{\text{syn}}$ restrict to zero on $F_\gamma^{i+1} K_j(X, \mathbf{Z}/p^n)$, $j \geq 2$, unless $j = 2$, $p = 2$.

Proof. The proof is basically a translation of the proof of Lemma 2.1 in [30] into the language of spaces used in [14]. \square

Remark 5.4. Let X be a scheme over K or \bar{K} . In an analogous way one constructs the étale total Chern classes

$$C_X^{\text{ét}} : K_X \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \text{R}\varepsilon_* \mathbf{Z}/p^n(i)_X),$$

where $\varepsilon : X_{\text{ét}} \rightarrow X_{\text{Zar}}$ is the natural projection, that have all the properties listed in Lemma 5.1. They induce the characteristic classes

$$\begin{aligned} \bar{c}_{i,j}^{\text{ét}} &: K_j(X, \mathbf{Z}/p^n) \rightarrow H^{2i-j}(X, \mathbf{Z}/p^n(i)), \quad j \geq 2, \\ c_{i,j}^{\text{ét}} &: K_j(X) \rightarrow H^{2i-j}(X, \mathbf{Z}/p^n(i)), \quad j \geq 0, \end{aligned}$$

that have all the properties listed in Lemma 5.3.

5.2. Truncated syntomic Chern classes. We will show now that there exists Chern classes into truncated syntomic as well as syntomic-étale cohomology that are compatible with the classical syntomic Chern classes. We define the universal classes $C_{i,a}^{\text{syn}} \in H^{2i}(BGL_a/W(k), S_n(i)_{\text{Nis}})$ as the unique classes mapping to the classical syntomic universal classes. This can be done by Proposition 3.4. They induce Chern class and total Chern class maps

$$C_i^{\text{syn}} : K_X \rightarrow \mathcal{K}(2i, \tilde{S}_n(i)_{\text{Nis},X}), \quad i \geq 0; \quad C_X^{\text{syn}} : K_X \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{S}_n(i)_{\text{Nis},X})$$

As before, we get the characteristic classes

$$\begin{aligned} \bar{c}_{i,j}^{\text{syn}} &: K_j(X, \mathbf{Z}/p^n) \rightarrow H^{2i-j}(X, S_n(i)_{\text{Nis}}), \quad j \geq 2, \\ c_{i,j}^{\text{syn}} &: K_j(X) \rightarrow H^{2i-j}(X, S_n(i)_{\text{Nis}}), \quad j \geq 0, \end{aligned}$$

Via the map $\tau : H^{2i}(BGL_a, S_n(i)_{\text{Nis}}) \rightarrow H^{2i}(BGL_a, S'_n(i)_{\text{Nis}})$ we get induced universal classes, total Chern maps and characteristic classes with values in $S'_n(i)_{\text{Nis}}$.

Lemma 5.5. *The above truncated syntomic total Chern class maps and the induced characteristic classes satisfy analogs of Lemma 5.1, Lemma 5.2, and Lemma 5.3.*

Proof. We start with Lemma 5.1 describing properties of the universal total Chern class maps. Property (1) follows from functoriality of truncated syntomic universal classes. Property (2) follows from a universal Whitney sum formula. More specifically, consider the universal short exact sequence

$$0 \rightarrow E_{\bullet}^a \rightarrow E_{\bullet}^{a+b} \rightarrow E_{\bullet}^b \rightarrow 0$$

of vector bundles over $BGL(a, b)$. Here the middle term is the universal vector bundle and the first and last terms are classified by the natural maps to BGL_a and BGL_b , respectively. A classical argument using only the projective space theorem gives the Whitney sum formula in de Rham cohomology

$$C_{\bullet}(E_{\bullet}^{a+b}) = C_{\bullet}(E_{\bullet}^a)C_{\bullet}(E_{\bullet}^b).$$

By Proposition 3.4 this induces the Whitney sum formula in the truncated syntomic cohomology.

For property (3), again one needs to consider only the universal situation, i.e., the universal tensor product bundle $E_{\bullet}^a \otimes E_{\bullet}^b$ over $B(GL_a \times GL_b)$. Splitting principle gives the formula

$$\tilde{C}_{\bullet}(E_{\bullet}^a \otimes E_{\bullet}^b) = \tilde{C}_{\bullet}(E_{\bullet}^a) \star \tilde{C}_{\bullet}(E_{\bullet}^b).$$

in de Rham cohomology. By Proposition 3.4 it carries over to truncated syntomic cohomology.

The analog of Lemma 5.2 is proved in the same way as the original lemma using Proposition 3.4. Having an analog of Lemma 5.1, the proof of Lemma 5.3 for truncated syntomic cohomology carries verbatim from the untruncated case. \square

Similarly, we construct (using Proposition 3.8) universal Chern classes, total Chern maps, and characteristic classes with values in (truncated) syntomic-étale cohomologies $E_n(i)_{\text{Nis}}$, $E'_n(i)_{\text{Nis}}$ and $E_n(\hat{i})$, $E'_n(\hat{i})$. Clearly, so defined (truncated) syntomic-étale total Chern class maps and the induced characteristic classes satisfy analogs of Lemma 5.1, Lemma 5.2, and Lemma 5.3.

5.3. Logarithmic syntomic Chern classes. Let X be a semistable scheme over \mathcal{O}_K^{\times} or a semistable scheme over \mathcal{O}_K with a smooth special fiber. Let D be the horizontal divisor and set $j : X \setminus D \hookrightarrow X$. In this section, we will construct universal Chern class maps (in the homotopy category of pointed presheaves of simplicial sets on X_{Nis})

$$C_{X \setminus D, i}^{\text{syn}} : j_* K_{X \setminus D} \rightarrow \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D}(D)), \quad i \geq 0,$$

where $\tilde{E}'_n(i)_{X \setminus D}(D)_{\text{Nis}}$ is an injective resolution of $E'_n(i)_{X \setminus D}(D)_{\text{Nis}}$. In what follows we will skip the subscript Nis if confusion does not arise. Up to some universal constants these maps will have all the expected properties.

Consider the map $C_{X \setminus D}^{\text{syn}} : j_* K_{X \setminus D} \rightarrow \text{R}j_* \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D})$ induced by the total syntomic-étale Chern class maps defined in Section 5.2. Lemma 5.6 below shows that a multiple $[p^{mi}]C_{X \setminus D}^{\text{syn}}$, for a universal constant m , of this map lifts, via the map

$$\mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D}(D)) \rightarrow \mathcal{K}(2i, \text{R}j_* \tilde{E}'_n(i)_{X \setminus D}) \rightarrow \text{R}j_* \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D}),$$

to a map

$$C_{X \setminus D}^{\text{syn}}(m) : j_* K_{X \setminus D} \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D}(D)).$$

Lemma 5.6. *Let \mathcal{L} be a complex of pointed presheaves of simplicial sets on X with homotopy presheaves concentrated in nonnegative degrees. The kernel and cokernel of the map*

$$\text{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D}(D))) \rightarrow \text{Hom}_{\mathcal{P}}(\mathcal{L}, \text{R}j_* \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D}))$$

is annihilated by p^{Ni} for a constant $N = c_1 c_2$, where c_1 is the constant⁶ from Theorem 2.2 and c_2 is a constant depending only on the dimension d of X . Here the homomorphisms are taken in the homotopy category \mathcal{P} of pointed presheaves of simplicial sets on X_{Nis} .

⁶We take $c_1 = N$ in the notation of Theorem 2.2.

Proof. For $i = 0$, we have an isomorphism by Corollary 2.4. Assume $i \geq 1$. Since the homotopy presheaves of \mathcal{L} are trivial in negative degrees, we have that

$$\mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \mathrm{R}j_* \tilde{E}'_n(i)_{X \setminus D})) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathrm{R}j_* \mathcal{K}(2i, \tilde{E}'_n(i)_{X \setminus D})).$$

Hence it suffices to show that the above lemma holds for the map

$$\mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \tilde{E}'_n(i)_X(D))) \rightarrow \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \mathrm{R}j_* \tilde{E}'_n(i)_{X \setminus D})).$$

We have,

$$\begin{aligned} \mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \tilde{E}'_n(i)_X(D))) &\simeq \mathrm{Hom}_{\mathcal{S}}(\mathcal{L}^a, \mathcal{K}(2i, \tilde{E}'_n(i)_X(D))) \\ &\simeq \mathrm{Hom}_D(C(\mathcal{L}^a), E_n(i)_X(D)[2i]), \end{aligned}$$

where $\mathrm{Hom}_{\mathcal{S}}(\cdot)$ refers to homomorphisms in the homotopy category of pointed sheaves of simplicial sets, \mathcal{L}^a is the complex of sheaves associated to \mathcal{L} , $C(\mathcal{L}^a)$ is the sheaf of (normalized) chain complexes associated to \mathcal{L}^a that we see as a sheaf of cochain complexes by negating the degrees, and $\mathrm{Hom}_D(\cdot)$ refers to homomorphisms in the derived category of complexes of sheaves of abelian groups on $X(D)_{\mathrm{Nis}}$. Similarly,

$$\mathrm{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \mathrm{R}j_* \tilde{E}'_n(i)_{X \setminus D})) \simeq \mathrm{Hom}_D(C(\mathcal{L}^a), j_* \tilde{E}'_n(i)_{X \setminus D}[2i]).$$

Let $\mathcal{C}(i)$ denote the mapping fiber of the map $\tilde{E}'_n(i)_X(D) \rightarrow j_* \tilde{E}'_n(i)_{X \setminus D}$. We need to show that $\mathrm{Hom}_D(\cdot, \mathcal{C}(i))$ is annihilated by p^{Ni} for a constant N described above. It suffices to show that $\mathcal{C}(i)$ has cohomology annihilated by $p^{c_1 r i}$ for the constant c_1 as above and that $\mathcal{C}(i)$ has cohomological length less than c_2 for a constants c_2 depending only on d .

The first claim follows from Corollary 2.4. For the second claim, since Nisnevich topology of X has cohomological dimension less than d , it suffices to show that $\mathcal{C}(i)$ has cohomological length less than c_3 for a constant c_3 depending only on d . But this is clear since the cohomological dimension of the étale topos of X and the length of the (filtered) crystalline cohomology of $X/W(k)$ are bounded by $2d + 3$. \square

Let $m \geq N$. Set $C_{X \setminus D}^{\mathrm{syn}, m} := [p^{mi}]C_{X \setminus D}^{\mathrm{syn}}(m)$. By the above lemma two different lifts $C_{X \setminus D}^{\mathrm{syn}}(m)$ yield the same class $C_{X \setminus D}^{\mathrm{syn}, m}$. Similarly, we define the total syntomic-étale Chern class maps

$$C_{X \setminus D}^{\mathrm{syn}} : j_* K_{X \setminus D} \rightarrow \prod_{p-2 \geq i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)).$$

No additional constants are needed here.

We record the following properties of these Chern class maps.

- Theorem 5.7.** (1) *The Chern class maps $C_{X \setminus D}^{\mathrm{syn}, m}$ are compatible with the reduction maps $E'_n(i)_X(D) \rightarrow E'_{n_1}(i)_X(D)$, $n \geq n_1$.*
(2) *The Chern class maps $C_{X \setminus D}^{\mathrm{syn}, m}$ are independent of the number m chosen, that is, for two different numbers $m_1 < m_2$ the following diagram commutes*

$$(5.2) \quad \begin{array}{ccc} G_X(D) & \xrightarrow{C_{X \setminus D}^{\mathrm{syn}, m_1}} & \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)) \\ & \searrow C_{X \setminus D}^{\mathrm{syn}, m_2} & \downarrow [p^{2(m_2 - m_1)i}] \\ & & \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)) \end{array}$$

- (3) *The Chern class map $C_{X \setminus D}^{\mathrm{syn}, m}$ is functorial for morphisms of log-schemes as above $\pi : Y \rightarrow X$ such that $\pi^{-1}(D_X) \subset D_Y$, i.e., the following diagram commutes in the homotopy category*

$$\begin{array}{ccc} j_* K_{X \setminus D_X} & \xrightarrow{C_{X \setminus D_X}^{\mathrm{syn}, m}} & \mathcal{K}(2i, \tilde{E}'_n(i)_X(D_X)) \\ \downarrow \pi^* & & \downarrow \pi^* \\ \pi_* j'_* K_{Y \setminus D_Y} & \xrightarrow{C_{Y \setminus D_Y}^{\mathrm{syn}, m}} & \mathrm{R}\pi_* \mathcal{K}(2i, \tilde{E}'_n(i)_Y(D_Y)), \end{array}$$

where $j' : Y \setminus D_Y \hookrightarrow Y$ is the natural open immersion.

- (4) The total Chern class map $C_{X \setminus D}^{\text{syn}, m}$ is compatible with addition. The augmented total Chern class map $\tilde{C}_{X \setminus D}^{\text{syn}, m}$ is compatible with products.
- (5) The total Chern class map

$$C_{X \setminus D}^{\text{syn}, m} : j_* K_{X \setminus D} \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)), \quad n \geq 1,$$

is an extension of the syntomic-étale Chern class map

$$[p^{2mi}]C_X^{\text{syn}} : K_X \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X), \quad n \geq 1.$$

Two such extensions become equal after multiplication by $[p^{mi}]$.

Similarly for the Chern classes with values in $E_n(i)$ -cohomology.

Proof. The first two claims are immediate from construction. The next two follow from Section 5.2 and Lemma 5.6; to control the constants we use the fact that the Whitney sum formula and the product formula involve homogenous polynomials of the right degrees. For the fifth claim it remains to show that an extension of the Chern class maps $[p^{2mi}]C_X^{\text{syn}}$ to $X \setminus D$ is unique up to multiplication by $[p^{mi}]$. To do this, consider the localization homotopy cofiber sequence

$$G(D) \xrightarrow{i_*} G(X) \rightarrow G(X \setminus D)$$

and apply Lemma 5.8 below to $\mathcal{L} = i_* G_D[1]$. \square

Lemma 5.8. *Let \mathcal{L} be a complex of pointed presheaves of simplicial sets on D_1 with homotopy presheaves concentrated in nonnegative degrees.*

- (1) For $0 \leq i \leq p-2$, we have

$$\text{Hom}_{\mathcal{P}}(i_* \mathcal{L}, \mathcal{K}(2i, \tilde{E}_n(i)_X(D))) = 0.$$

- (2) For $0 \leq i$, the group

$$\text{Hom}_{\mathcal{P}}(i_* \mathcal{L}, \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)))$$

is annihilated by p^{Ni} , where N is the constant from Lemma 5.6.

Proof. We start with the first claim. Take $p-2 \geq i \geq 0$. We have

$$\begin{aligned} \text{Hom}_{\mathcal{P}}(i_* \mathcal{L}, \mathcal{K}(2i, \tilde{E}_n(i)_X(D))) &\simeq \text{Hom}_{\mathcal{S}}(i_* \mathcal{L}^a, \mathcal{K}(2i, \tilde{E}_n(i)_X(D))) \\ &\simeq \text{Hom}_D(i_* C(\mathcal{L}^a), E_n(i)_X(D)[2i]) \simeq \text{Hom}_D(C(\mathcal{L}^a), \text{Ri}^1 E_n(i)_X(D)[2i]). \end{aligned}$$

We claim that $\text{Ri}^1 E_n(i)_X(D) = 0$. Indeed, for X semistable over \mathcal{O}_K^\times , by Theorem 2.3, we have a quasi-isomorphism $E_n(i)_X(D)_{\text{Nis}} \xrightarrow{\sim} \tau_{\leq i} \text{R}\varepsilon_* \text{R}j_* \mathbf{Z}/p^n(i)$, where $\varepsilon : X_{\text{ét}} \rightarrow X_{\text{Nis}}$ is the projection and $j : X_K \setminus D_K \hookrightarrow X$ is the natural open immersion. But the Beilinson-Lichtenbaum conjecture implies that

$$\tau_{\leq i} \text{R}\varepsilon_* \text{R}j_* \mathbf{Z}/p^n(i) \simeq \tau_{\leq i} \text{R}j_* \text{R}\varepsilon_* \mathbf{Z}/p^n(i) \simeq \tau_{\leq i} \text{R}j_* \tau_{\leq i} \text{R}\varepsilon_* \mathbf{Z}/p^n(i) \simeq \tau_{\leq i} \text{R}j_* \mathbf{Z}/p^n(i)_M$$

and we have

$$\tau_{\leq i} \text{R}j_* \mathbf{Z}/p^n(i)_M \simeq \tau_{\leq i} j_* \mathbf{Z}/p^n(i)_M \simeq j_* \mathbf{Z}/p^n(i)_M \simeq \text{R}j_* \mathbf{Z}/p^n(i)_M.$$

Hence

$$\text{Ri}^1 E_n(i)_X(D) \simeq \text{Ri}^1 \text{R}j_* \mathbf{Z}/p^n(i)_M = 0,$$

as wanted.

If X is a semistable scheme over \mathcal{O}_K with a smooth special fiber, by Theorem 2.5, we have a quasi-isomorphism

$$E_n(i)_X(D)_{\text{Nis}} \xrightarrow{\sim} \tau_{\leq i} \text{R}\varepsilon_* \text{R}j'_* \mathbf{Z}/p^n(i)_M,$$

where $j' : X \setminus D \hookrightarrow X$ is the natural open immersion. But by the Beilinson-Lichtenbaum conjecture in mixed characteristic [10, Theorem 1.2.2] that states that we have a quasi-isomorphism

$$\tau_{\leq i} \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)_M \simeq \mathbf{Z}/p^n(i)_M$$

and by the fact that $\tau_{\leq i} \mathbf{Z}/p^n(i)_M \xrightarrow{\sim} \mathbf{Z}/p^n(i)_M$ [10, Cor. 4.2], we have

$$\begin{aligned} \tau_{\leq i} \mathbf{R}\varepsilon_* \mathbf{R}j'_* \mathbf{Z}/p^n(i) &\simeq \tau_{\leq i} \mathbf{R}j'_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i) \simeq \tau_{\leq i} \mathbf{R}j'_* \tau_{\leq i} \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i) \simeq \tau_{\leq i} \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M \\ &\simeq \tau_{\leq i} j'_* \mathbf{Z}/p^n(i)_M \simeq j'_* \mathbf{Z}/p^n(i)_M \simeq \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M. \end{aligned}$$

Hence

$$\mathbf{R}i^! E_n(i)_X(D) \simeq \mathbf{R}i^! \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M = 0,$$

as wanted.

For the second claim of the lemma, we can assume that $i \geq 1$ (the case of $i = 0$ being treated above). As above we compute that

$$\mathrm{Hom}_{\mathcal{P}}(i_* \mathcal{L}, \mathcal{K}(2i, \tilde{E}'_n(i)_X(D))) \simeq \mathrm{Hom}_D(C(\mathcal{L}^a), \mathbf{R}i^! E'_n(i)_X(D)[2i]).$$

We need to show that $\mathrm{Hom}_D(C(\mathcal{L}^a), \mathbf{R}i^! E'_n(i)_X(D)[2i])$ is annihilated by p^{Ni} for a constant N as above. It suffices to show that $\mathbf{R}i^! E'_n(i)_X(D)$ has cohomology annihilated by p^{Ni} , N as in Lemma 5.6, and that $\mathbf{R}i^! E'_n(i)_X(D)$ has cohomological length less than c_1 for a constant c_1 depending only on d .

Concerning the cohomological length of $\mathbf{R}i^! E'_n(i)_X(D)$, since Nisnevich topology of X has cohomological dimension less than d , it suffices to show that $E'_n(i)_X(D)$ has cohomological length less than c_1 . By (2.6) we have a distinguished triangle (for $i_0 : X_0 \hookrightarrow X$)

$$j_{\mathrm{Nis}!} \mathbf{R}\varepsilon_* \mathbf{R}j'_* \mathbf{Z}/p^n(i)' \rightarrow \mathbf{R}\varepsilon_* E'_n(i)_X(D) \rightarrow i_{0*} \mathbf{R}\varepsilon_* S'_n(i)_X(D).$$

It suffices thus to show that both $\mathbf{R}\varepsilon_* \mathbf{R}j'_* \mathbf{Z}/p^n(i)'$ and $\mathbf{R}\varepsilon_* S'_n(i)_X(D)$ have cohomological length less than c_1 . But this is clear since the first complex has length bounded by $2d$ - the cohomological dimension of the étale topos of X and the second complex has length bounded by $2d + 3$ ($2d + 2$ being the length of the (filtered) crystalline cohomology of $X/W(k)$).

It remains to show that $\mathbf{R}i^! E'_n(i)_X(D)$ has cohomology annihilated by p^{Ni} . Let us look first at the case when X is semistable over \mathcal{O}_K with a smooth special fiber. By Theorem 2.5 we have a distinguished triangle

$$(5.3) \quad \mathcal{C} \rightarrow E'_n(i)_X(D) \rightarrow \tau_{\leq i} \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M,$$

where \mathcal{C} is a complex whose cohomology is annihilated by p^{Ni} . Arguing as above we have that $\mathbf{R}i^! \tau_{\leq i} \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M = 0$. Hence $\mathbf{R}i^! \mathcal{C} \xrightarrow{\sim} \mathbf{R}i^! E'_n(i)_X(D)$. Since Nisnevich topos of X has cohomological dimension less than d it suffices now to show that the complex \mathcal{C} has cohomological length less than c_1 for a constant c_1 depending only on d . By the distinguished triangle (5.3), since we have shown this for $E'_n(i)_X(D)$, it remains to show it for $\tau_{\leq i} \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M$. Hence it suffices to show that for a log-scheme T , smooth over \mathcal{O}_K the cohomology groups $H^*(T_{\mathrm{tr}, \mathrm{Nis}}, \mathbf{Z}/p^n(i)_M)$ vanish in degree larger than twice the dimension d of T . But this is clear since Nisnevich topos has cohomological dimension d and cohomology sheaves of $\mathbf{Z}/p^n(i)_M$ are trivial above degree d [10, Theorem 1.2].

Let now X be semistable over \mathcal{O}_K^\times . Arguing as above using Theorem 2.3 we reduce to showing that $\tau_{\leq i} \mathbf{R}j'_* \mathbf{Z}/p^n(i)_M$ has cohomological length less than c_1 - a constant depending only on d . This can be treated as above and we have finished the proof of our lemma. \square

Having the universal Chern class maps, as before, we get the (compatible) characteristic classes

$$\begin{aligned} \bar{c}_{i,j}^{\mathrm{syn},m} &: K_j(X \setminus D, \mathbf{Z}/p^n) \rightarrow H^{2i-j}(X, E_n^*(i)_X(D)), \quad j \geq 2, E^* = E, E' \\ c_{i,j}^{\mathrm{syn},m} &: K_j(X \setminus D) \rightarrow H^{2i-j}(X, E_n^*(i)_X(D)), \quad j \geq 0, E^* = E, E'. \end{aligned}$$

By composition with the maps $E'_n(i) \rightarrow S'_n(i)$ and $E_n(i) \rightarrow S_n(i)$ we obtain syntomic Chern classes with values in $S'_n(i)$ and $S_n(i)$. Clearly, all these Chern classes satisfy analogs of Lemma 5.3.

Corollary 5.9. *Set $U = X \setminus D$. The syntomic Chern classes*

$$\bar{c}_{ij}^{\text{syn},m} : K_j(U, \mathbf{Z}/p^n) \rightarrow H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_n(i)), \quad j \geq 2,$$

are compatible, via the period maps $\alpha_{*,*}$ of Fontaine-Messing, with étale Chern classes, i.e., the following diagram commutes

$$\begin{array}{ccc} K_j(U, \mathbf{Z}/p^n) & \xrightarrow{\bar{c}_{ij}^{\text{syn},m}} & H_{\text{syn}}^{2i-j}(X, \mathcal{S}'_n(i)) \\ \downarrow j^* & & \downarrow \alpha_{2i-j,i} \\ K_j(U_K, \mathbf{Z}/p^n) & \xrightarrow{p^{(m+1)i} \bar{c}_{ij}^{\text{ét}}} & H_{\text{ét}}^{2i-j}(U_K, \mathbf{Z}/p^n(i)'), \end{array}$$

where $j : U_K \hookrightarrow U$ is the natural open immersion. We have analogous statement for $j \leq p-2$ and \mathcal{S} -cohomology, where no twist is necessary.

Proof. We will present the argument for \mathcal{S}' -cohomology, the one for \mathcal{S} -cohomology being analogous. It suffices to show that the total universal syntomic-étale and étale Chern classes are compatible, i.e., that the following diagram commutes

$$\begin{array}{ccc} j_* K_{X \setminus D} & \xrightarrow{C_{X \setminus D}^{\text{syn},m}} & \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)) \\ \downarrow & & \downarrow [\alpha_i] \\ j_{\text{Nis},*} j_* K_{X_K \setminus D_K} & \xrightarrow{[p^{(m+1)i}] C_{X_K \setminus D_K}^{\text{ét}}} & \mathbf{R}j_{\text{Nis},*} \mathbf{R}j_* \prod_{i \geq 0} \mathcal{K}(2i, \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)'_{X_K \setminus D_K}). \end{array}$$

It follows from functoriality of syntomic-étale Chern classes and period maps that it suffices to show this in the case when the divisor D is trivial, i.e., in the case of classical Chern classes. But this reduces to showing that the period map $\alpha_i : H^{2i}(BGL_l, E'_n(i)_{\text{Nis}}) \rightarrow H_{\text{ét}}^{2i}(BGL_{l,F}, \mathbf{Z}/p^n(i)')$ maps the syntomic-étale universal class x_i^{syn} to the étale universal class $p^i x_i^{\text{ét}}$. But this we proved in Corollary 3.11. \square

Remark 5.10. Theorem 5.7 allows us to define logarithmic Chern class maps (having all the described above properties) with values in log-crystalline cohomology $H_{\text{cr}}^*(X_n, \mathcal{J}_{X_n}^{[*]})$ for X semistable over \mathcal{O}_K with a smooth special fiber.

6. CHERN CLASS MAPS AND GYSIN SEQUENCES

We will show in this section that the syntomic universal Chern class maps are compatible with the canonical Gysin sequences.

Let X be a semistable scheme over \mathcal{O}_K^\times or a semistable scheme over \mathcal{O}_K with a smooth special fiber. Let D' be the horizontal divisor. Assume that $D' = \cup_{i=1}^m D_i$, $m \geq 1$, is a union of m irreducible components D_i . Note that each scheme D_i with the induced log-structure is of the same type as X (with at most $m-1$ components in the divisor at infinity). Set $D = \cup_{i=2}^m D_i$, $Y = D_1$, $i : Y \hookrightarrow X$. The pairs (X, D) and (Y, D_Y) are of the same type as the pair (X, D') we started with but with at most $m-1$ irreducible divisors at infinity.

6.1. Basic properties of syntomic cohomology. In this section, we will list several basic properties of syntomic and syntomic-étale cohomologies.

6.1.1. Gysin sequences. We start with Gysin sequences. We have the following localization sequences.

Lemma 6.1. *For $r \geq 1$, there exist the following distinguished triangles*

$$\begin{aligned} i_* S_n^{i+1}(r-1)_Y(D_Y)[-2] &\xrightarrow{i^!} S_n^i(r)_X(D) \rightarrow S_n^i(r)_X(Y \cup D), \quad i \geq 0; \\ i_* S_n(r-1)_Y(D_Y)[-2] &\xrightarrow{i^!} S_n(r)_X(D) \rightarrow S_n(r)_X(Y \cup D), \quad r \leq p-1. \end{aligned}$$

There are analogous distinguished triangles for the complexes $E_n(r)$, $r \leq p-2$, $E'_n(r)$, and $E_n^1(r)$. Moreover, for $r \geq 1$, there exist the following distinguished triangles

$$\begin{aligned} i_* F_n^1(r-1)_X(Y \cup D)[-2] &\xrightarrow{i^!} E'_n(r)_{\text{Nis},X}(D) \rightarrow E'_n(r)_{\text{Nis},X}(Y \cup D), \quad i \geq 0; \\ i_* E_n(r-1)_{\text{Nis},Y}(D_Y)[-2] &\xrightarrow{i^!} E_n(r)_{\text{Nis},X}(D) \rightarrow E_n(r)_{\text{Nis},X}(Y \cup D), \quad r \leq p-1. \end{aligned}$$

Here $F_n^1(r-1)_X(Y \cup D)$ is a complex of \mathbf{Z}/p^n -sheaves on Y equipped with a natural map $F_n^1(r-1)_X(Y \cup D) \rightarrow E_n^1(r-1)_{\text{Nis},Y}(D_Y)$ that has an inverse up to $p^{N(r-1)}$, i.e., composition either way is a multiplication by $p^{N(r-1)}$ (in the derived category), where N is the constant from Theorem 2.2.

Proof. (1) *Étale complexes*

We will start the proof in the case of complexes with undivided Frobenius. Recall that we have the following short exact sequence of complexes

$$0 \rightarrow S_n^i(r)_X(D) \rightarrow S_n^i(r)_X(Y \cup D) \xrightarrow{\text{res}_Y} i_* S_n^{i+1}(r-1)_Y(D_Y)[-1] \rightarrow 0$$

This was shown in [41, 4.2.4.3] and follows relatively easily from two facts. First, that for a regular log-scheme X_n log-smooth over $W_n(k)$, an irreducible regular divisor D_1 log-smooth over $W_n(k)$, and a divisor D on X_n such that $D_1 \cup D$ is a relative simple normal crossing divisor on X_n over $W_n(k)$ and $D_1 \cap D$ a relative simple normal crossing divisor on D_1 over $W_n(k)$ the following sequence is exact

$$0 \rightarrow \Omega_{X_n}^q(D_1) \rightarrow \Omega_{X_n}^q(D_1 \cup D) \xrightarrow{\text{res}_{D_1}} \Omega_{D_1, n}^{q-1}(D_1 \cap D) \rightarrow 0, \quad q \geq 1$$

Second, that the corresponding divided power envelopes behave as expected.

In the case of complexes $E'_n(r)$, we want to construct the following distinguished triangle

$$(6.1) \quad E'_n(r)_X(D) \rightarrow E'_n(r)_X(Y \cup D) \xrightarrow{\text{res}_Y} i_* E_n^1(r-1)_Y(D_Y)[-1],$$

where the first map is the natural map and the residue map is yet to be defined. Since we have Gysin sequences for syntomic and étale cohomologies this amounts to checking that they can be glued. Tsuji in [41] verified that these sequences are compatible. This is nontrivial since the period map does not behave well with respect to closed immersions. We will use his constructions to perform the gluing.

To start, we replace the maps $Y(D_Y) \rightarrow X(D) \leftarrow X(Y \cup D)$ with the maps $Y(D_Y) \xrightarrow{\text{Id}} Y(D_Y) \leftarrow Y(D_Y^0)$, where $Y(D_Y^0)$ is the scheme Y endowed with the pullback of the log-structure of $X(Y \cup D)$. Consider the following commutative diagram. All the maps but the bottom rightmost map are genuine maps of complexes.

$$(6.2) \quad \begin{array}{ccccc} E'_n(r)_X(D) & \longrightarrow & E'_n(r)_X(Y \cup D) & & \\ \parallel & & \downarrow \wr & & \\ E'_n(r)_X(D) & \xhookrightarrow{f} & E'_n(r)_X(Y \cup D)_{\text{Két}} & \xrightarrow{\text{res}} & C(f, r-1) \\ \downarrow i^* & & \downarrow i^* & & \downarrow i^* \\ i_* E'_n(r)_Y(D_Y) & \xhookrightarrow{f'} & i_* E'_n(r)_Y(D_Y^0)_{\text{Két}} & \xrightarrow{\text{res}} & i_* C(f', r-1) \xrightarrow{\sim} i_* E_n^1(r-1)_Y(D_Y)[-1] \end{array}$$

The complex $E'_n(r)_X(Y \cup D)_{\text{Két}}$ is defined in an analogous way to $E'_n(r)_X(Y \cup D)$ by (locally) using the map (assume that $U = \text{Spec}(A) \rightarrow X$ is a strict étale map)

$$\begin{aligned} \alpha'_{r,U} : \Gamma(U, S'_n(r)_{X^0, Z^0}) &\rightarrow \Gamma(U^h, \theta_U \tau_* G\overline{S}_n(r)_{U^0, Z^0}) \xrightarrow{\sim} \Gamma(U^h, \theta_U \tau_* G\Lambda_{U^h}) \xrightarrow{\sim} \Gamma(U, \theta_U \tau_* G\Lambda_U) \\ &\xrightarrow{\sim} \Gamma(U, \text{Cone}(\eta(\tau_* G\Lambda_U))) \rightarrow \Gamma(U, j_{\text{ét}!} j'_* \tau_* G\Lambda_U[1]), \end{aligned}$$

where $\tau : X(D)_{K, \text{tr}, \text{Két}}^0 \rightarrow X(D)_{K, \text{tr}, \text{ét}}^0$ is the natural map from the Kummer étale topos to the étale topos and the subscripts 0 refer to the log-structure induced by $Y \cup D$. We have a natural morphism $E'_n(r)_X(Y \cup D) \xrightarrow{\sim} E'_n(r)_X(Y \cup D)_{\text{Két}}$ that is a quasi-isomorphism. Moreover the induced map of complexes $f : E'_n(r)_X(D) \rightarrow E'_n(r)_X(Y \cup D)_{\text{Két}}$ is injective [41, Lemma 4.7.5]. We wrote $C(f, r-1)$ for the cokernel of this map.

The complex $E'_n(r)_Y(D_Y^0)_{\text{Két}}$ is defined in an analogous way to $E'_n(r)_X(Y \cup D)_{\text{Két}}$. The local setting is as follows. We assume that X (with the log-structure defined by D) is affine and we have a closed exact immersion $X \hookrightarrow Z$ over $W(k)$ into a fine log-scheme Z that is log-smooth over $W(k)$ and has a compatible system of liftings of Frobenius $\{F_{Z_n} : Z_n \rightarrow Z_n\}$. Further we assume that Z has a Cartier divisor $Z_1 \hookrightarrow Z$ defined by a global equation $g = 0$ for $g \in \Gamma(Z, \mathcal{O}_Z)$ and such that Y is a pullback of Z_1 and Z_1 equipped with the log-structure pullbacked from Z is log-smooth over $W(k)$. We denote by Z^0 the log-scheme Z equipped with the log-structure induced by Z_1 and M_Z , and by Z_1^0 - the log-scheme equipped with the log-structure pullbacked from Z^0 .

Let $U = \text{Spec}(A) \rightarrow X$ be a strict étale morphism. Write $U_1 = Y \times_X U$. Under certain additional assumptions on Z and U (cf. [41, 4.7]) we get the following map

$$\begin{aligned} \alpha'_{r,U_1} : \Gamma(U_1, S'_n(r)_{Y^0, Z_1^0}) &\rightarrow \Gamma(U_1^h, \theta'_{U_1} \tau'_* G \bar{S}_n(r)_{U_1^0, Z_1^0}) \xleftarrow{\sim} \Gamma(U_1^h, \theta'_{U_1} \tau'_* G \Lambda_{U_1^h}) \xleftarrow{\sim} \Gamma(U_1, \theta'_{U_1} \tau'_* G \Lambda_{U_1}) \\ &\xleftarrow{\sim} \Gamma(U_1, \text{Cone}(\eta'(\tau'_* G \Lambda_{U_1}))) \rightarrow \Gamma(U_1, j_{\text{ét}!} j'_* \tau'_* G \Lambda_{U_1}[1]), \end{aligned}$$

where $\tau' : Y(D_Y)_{K, \text{tr}, \text{Két}}^0 \rightarrow Y(D_Y)_{K, \text{tr}, \text{ét}}^0$ is the natural map of topoi and $\theta' = \iota_* \iota^* j_* G$ for $\iota : Y_0 \hookrightarrow Y$, $j : Y_{K, \text{tr}} \xrightarrow{j'} Y_K \xrightarrow{j_{\text{ét}}} Y$. We set $\eta'(M) = j_{\text{ét}!} j'_* M \hookrightarrow j_{\text{ét}!} j'_* M$. Here the resolution $\Lambda_{U_1} \xrightarrow{\sim} \bar{S}_n(r)_{U_1^0, Z_1^0}$ on the Kummer étale topoi of $U_{1, K, \text{tr}}^0$ is the one defined in [41, 4.7]. By globalizing, straightening, and taking cone of the morphisms α'_{r,U_1} we get the complex $E'_n(r)_Y(D_Y^0)_{\text{Két}}$.

Recall that the complex $E'_n(r)_Y(D_Y)$ is defined locally by an analogous map

$$\begin{aligned} \alpha'_{r,U_1} : \Gamma(U_1, S'_n(r)_{Y, Z_1}) &\rightarrow \Gamma(U_1^h, \theta'_{U_1} \bar{S}_n(r)_{U_1, Z_1}) \xleftarrow{\sim} \Gamma(U_1^h, \theta'_{U_1} \Lambda_{U_1^h}) \xleftarrow{\sim} \Gamma(U_1, \theta'_{U_1} \Lambda_{U_1}) \\ &\xleftarrow{\sim} \Gamma(U_1, \text{Cone}(\eta(G \Lambda_{U_1}))) \rightarrow \Gamma(U_1, j_{\text{ét}!} j'_* G \Lambda_{U_1}[1]). \end{aligned}$$

By globalizing, straightening, and taking cone of the morphisms α'_{r,U_1} we get the complex $E'_n(r)_Y(D_Y)$. The induced map of complexes $f' : E'_n(r)_Y(D_Y) \rightarrow E'_n(r)_Y(D_Y^0)_{\text{Két}}$ is injective [41, Lemma 4.7.8]. We wrote $C(f', r-1)$ for the cokernel of this map. It is supported on Y .

We claim that the induced pullback map $i^* : C(f, r-1) \xrightarrow{\sim} C(f', r-1)$ is a quasi-isomorphism. To verify this notice that locally $C(f, r-1)$ is defined as the cone of the straightening of the following zigzag of maps

$$\begin{aligned} \Gamma(U, i_* S_n^1(r-1)_{Y, Z_1}[-1]) &\rightarrow \Gamma(U^h, \theta_U L_U) \xleftarrow{\sim} \Gamma(U^h, \theta_U C_{U^h}) \xleftarrow{\sim} \Gamma(U, \theta_U C_U) \\ &\xleftarrow{\sim} \Gamma(U, \text{Cone}(\eta(GC_U))) \rightarrow \Gamma(U, j_{\text{ét}!} j'_* GC_U[1]), \end{aligned}$$

where C and L are the cokernels of the injective morphisms $\Lambda_X \rightarrow \tau_* G(\Lambda_{X^0})$ and $\bar{S}_n(r)_{U, Z} \rightarrow \tau_* G \bar{S}_n(r)_{U^0, Z^0}$, respectively.

Similarly, $C(f', r-1)$ is defined as the cone of the straightening of the following zigzag of maps (c.f. [41, 4.7.12])

$$\begin{aligned} \Gamma(U_1, S_n^1(r-1)_{Y, Z_1}[-1]) &\rightarrow \Gamma(U_1^h, \theta'_{U_1} L'_{U_1}) \xleftarrow{\sim} \Gamma(U_1^h, \theta'_{U_1} C'_{U_1^h}) \xleftarrow{\sim} \Gamma(U_1, \theta'_{U_1} C'_{U_1}) \\ &\xleftarrow{\sim} \Gamma(U_1, \text{Cone}(\eta'(GC'_{U_1}))) \rightarrow \Gamma(U_1, j_{\text{ét}!} j'_* GC'_{U_1}[1]), \end{aligned}$$

where C' and L' are the cokernels of the injective morphisms $\Lambda_Y \rightarrow \tau_* G(\Lambda_{Y^0})$ and $\bar{S}_n(r)_{U_1, Z_1} \rightarrow \tau_* G \bar{S}_n(r)_{U_1^0, Z_1^0}$, respectively. There exists a natural map (i^*) from the first zigzag to the second one. Hence the two maps are equal in the derived category and we obtain a genuine map of complexes that yields a quasi-isomorphism $i^* : C(f, r-1) \xrightarrow{\sim} C(f', r-1)$, as wanted.

It remains to construct a quasi-isomorphism $C(f', r-1) \simeq E_n^1(r-1)_Y(D_Y)[-1]$. Consider the following two commutative diagrams of maps of complexes. Here we set $\Gamma^h(M) = \Gamma(U_1^h, \theta'_{U_1} M)$,

$$\Gamma(M) = \Gamma(U_1, \theta'_{U_1} M), \Lambda(a) = \mathbf{Z}/p^n(a)'. \quad (6.3)$$

$$\begin{array}{ccccccc} \Gamma^h(\Lambda(r-1)[-1]) & \xrightarrow{h} & \Gamma^h(\Lambda(r)(-1)[-1]) & \equiv & \Gamma^h(\Lambda(r)(-1)[-1]) & \xleftarrow{\sim} & \Gamma(\Lambda(r)(-1)[-1]) \\ \downarrow \wr & & \uparrow \wr & & \uparrow \wr & & \uparrow \wr \\ \Gamma^h(\mathcal{H}^1(\bar{S}_n(r-1)_{Y,Z_1})[-2]) & \longrightarrow & \Gamma^h(\mathcal{H}^1(L'_{U_1})[-1]) & \xleftarrow{\sim} & \Gamma^h(\mathcal{H}^1(C'_{U_1})[-1]) & \xleftarrow{\sim} & \Gamma(\mathcal{H}^1(C'_{U_1})[-1]) \\ \uparrow \wr & & \uparrow \wr & & \uparrow \wr & & \uparrow \wr \\ \Gamma^h(\tau_{\leq 1} \bar{S}_n(r-1)_{Y,Z_1}[-1]) & \longrightarrow & \Gamma^h(\tau_{\leq 1} L'_{U_1}) & \xleftarrow{\sim} & \Gamma^h(\tau_{\leq 1} C'_{U_1}) & \xleftarrow{\sim} & \Gamma(\tau_{\leq 1} C'_{U_1}) \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \Gamma^h(\bar{S}_n(r-1)_{Y,Z_1}[-1]) & \longrightarrow & \Gamma^h(L'_{U_1}) & \xleftarrow{\sim} & \Gamma^h(C'_{U_1}) & \xleftarrow{\sim} & \Gamma(C'_{U_1}) \end{array}$$

$$(6.4) \quad \begin{array}{ccccc} \Gamma(\Lambda(r)(-1)[-1]) & \xleftarrow{\sim} & \Gamma(\text{Cone}(\eta'(G\Lambda(r)(-1)[-1]))) & \longrightarrow & \Gamma(J_{\text{ét}!} J'_* G\Lambda(r)(-1)) \\ \uparrow \wr & & \uparrow \wr & & \uparrow \wr \\ \Gamma(\mathcal{H}^1(C'_{U_1})[-1]) & \xleftarrow{\sim} & \Gamma(\text{Cone}(\eta'(G\mathcal{H}^1(C'_{U_1})[-1]))) & \longrightarrow & \Gamma(J_{\text{ét}!} J'_* G\mathcal{H}^1(C'_{U_1})) \\ \uparrow \wr & & \uparrow \wr & & \uparrow \wr \\ \Gamma(\tau_{\leq 1} C'_{U_1}) & \xleftarrow{\sim} & \Gamma(\text{Cone}(\eta'(G\tau_{\leq 1} C'_{U_1}))) & \longrightarrow & \Gamma(J_{\text{ét}!} J'_* G\tau_{\leq 1} C'_{U_1}[1]) \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \Gamma(C'_{U_1}) & \xleftarrow{\sim} & \Gamma(\text{Cone}(\eta'(GC'_{U_1}))) & \longrightarrow & \Gamma(J_{\text{ét}!} J'_* GC'_{U_1}[1]) \end{array}$$

The map h is defined to make the upper left corner square of the first diagram commute. The bottom leftmost map in the same diagram is that defined in [41, 4.7.10]. By [41, Prop. 4.8.4] it is equal to the natural map. Take the two maps $\Gamma(U_1, \bar{S}_n(r-1)_{Y,Z_1}[-1]) \rightarrow \Gamma(U_1, \Lambda(-1)[-1])$ obtained by composing the maps from the left and upper sides of the first diagram and the upper side of the second diagram, respectively, from the lower and right sides. Composing these maps with the map $\Gamma(U_1, S_n^1(r-1)_{Y,Z_1}[-1]) \rightarrow \Gamma(U_1^h, \bar{S}_n(r-1)_{Y,Z_1}[-1])$ [41, 4.7.11] we obtain maps that define the complexes $E_n^1(r-1)_Y(D_Y)[-1]$ and $C(f', r-1)$, respectively [41, 4.7.10]. Now our diagrams describe a specific quasi-isomorphism $C(f', r-1) \simeq E_n^1(r-1)_Y(D_Y)[-1]$, as wanted.

We finish by setting the residue map $\text{res}_Y : E_n^1(r)_X(Y \cup D) \rightarrow i_* E_n^1(r-1)_Y(D_Y)[-1]$ to be equal to the composition of the maps from the diagram (6.2)

$$E_n^1(r)_X(Y \cup D) \xrightarrow{\sim} E_n^1(r)_X(Y \cup D)_{\text{Két}} \rightarrow C(f, r-1) \xrightarrow{i^*} i_* C(f', r-1) \simeq i_* E_n^1(r-1)_Y(D_Y)[-1],$$

and by pushing everything down to the Nisnevich site using $\varepsilon_* G$ for $\varepsilon : X_{\text{ét}} \rightarrow X_{\text{Nis}}$. The same diagram shows now that we have the distinguished triangle (6.1). By construction, the above residue map is compatible with the syntomic and étale residue maps

$$(6.5) \quad \begin{array}{l} S_n^1(r)_X(Y \cup D) \rightarrow i_* S_n^1(r-1)_Y(D_Y)[-1], \\ j_{\text{Nis}!} \mathbf{R}^r(j'_\varepsilon)_* \mathbf{Z}/p^n(r)_{X_{\text{tr}}} \rightarrow i_* j_{\text{Nis}!} \mathbf{R}^{r-1} \varepsilon_* \mathbf{Z}/p^n(r-1)_{Y_{\text{tr}}} = j_{\text{Nis}!} i_* \mathbf{R}^{r-1} \varepsilon_* \mathbf{Z}/p^n(r-1)_{Y_{\text{tr}}}. \end{array}$$

The constructions for the complexes $S_n(r)$, $r \leq p-1$, and $E_n(r)$, $r \leq p-2$ are analogous. So is the construction for the complexes $E_n^1(r)$ after one notices that the residue maps are compatible with the maps ω_0 and ω_1 .

(2) *Nisnevich complexes*

(2a) *The case of $r \geq p-2$*

In the case of complexes $E_n(r)_{\text{Nis}}$, $r \leq p-2$, we construct the triangle

$$(6.6) \quad E_n(r)_{\text{Nis},X}(D) \rightarrow E_n(r)_{\text{Nis},X}(Y \cup D) \xrightarrow{\text{res}_Y} i_* E_n(r-1)_{\text{Nis},Y}(D_Y)[-1]$$

by truncating the complexes in the analog of the distinguished triangle (6.1). To show that the obtained triangle is distinguished it suffices to show that the residue map induces a surjection

$$\text{res}_Y : \mathcal{H}^r(E_n(r)_{\text{Nis},X}(Y \cup D)) \rightarrow i_* \mathcal{H}^{r-1}(E_n(r-1)_{\text{Nis},Y}(D_Y)).$$

To simplify the notation we will assume that the divisor D is trivial. Consider the following commutative diagram (see (6.5))

$$\begin{array}{ccc} \mathcal{H}^r(S_n(r)_X(Y)) & \xrightarrow{\text{res}_Y} & i_* \mathcal{H}^{r-1}(S_n(r-1)_Y) \\ \uparrow & & \uparrow \\ \mathcal{H}^r(E_n(r)_{\text{Nis},X}(Y)) & \xrightarrow{\text{res}_Y} & i_* \mathcal{H}^{r-1}(E_n(r-1)_{\text{Nis},Y}) \\ \uparrow & & \uparrow \\ j_{\text{Nis}!} \mathbf{R}^r(j'\varepsilon)_* \mathbf{Z}/p^n(r)_{X_{\text{tr}}} & \xrightarrow{\text{res}_Y} & i_* j_{\text{Nis}!} \mathbf{R}^{r-1} \varepsilon_* \mathbf{Z}/p^n(r-1)_{Y_K} = j_{\text{Nis}!} i_* \mathbf{R}^{r-1} \varepsilon_* \mathbf{Z}/p^n(r-1)_{Y_K} \end{array}$$

It suffices to show that the bottom and the top residue maps in the above diagram are surjective. For that we can assume that X is affine and $Y = (T)$. Assume first that X is semistable over \mathcal{O}_K^\times . Since the period map is compatible with Gysin sequences by [41, Prop. 4.5.3], Theorem 2.1 implies that for the top map it suffices to show surjectivity of the map

$$(6.7) \quad \mathbf{R}^r(j\varepsilon)_* \mathbf{Z}/p^n(r)_{X_{\text{tr}}} \rightarrow i_* \mathbf{R}^{r-1}(j\varepsilon)_* \mathbf{Z}/p^n(r-1)_{Y_K}.$$

For the bottom map, it suffices to show surjectivity of the map

$$(6.8) \quad \mathbf{R}^r(j'\varepsilon)_* \mathbf{Z}/p^n(r)_{X_{\text{tr}}} \rightarrow i_* \mathbf{R}^{r-1} \varepsilon_* \mathbf{Z}/p^n(r-1)_{Y_K}.$$

Set $U := X_{\text{tr}}$. We have the symbol map

$$M_X^{\text{gp}} = j_* \mathcal{O}_U^* \rightarrow \mathbf{R}^1 j_* \mathbf{Z}/p^n(1)_U = \mathbf{R}^1 j_{\text{ét}*} \tau_{\leq 1} \mathbf{R} j'_* \mathbf{Z}/p^n(1)_U$$

and the following commutative diagram (cf. [39, 3.4.6]). We set here $\widetilde{M}_X^{\text{gp}} := j'_* \mathcal{O}_{X_K}^*$.

$$\begin{array}{ccccc} (M_X^{\text{gp}})^{\otimes r} & \longrightarrow & \mathbf{R}^r j_{\text{ét}*} \tau_{\leq r} \mathbf{R} j'_* \mathbf{Z}/p^n(r)_U & \longrightarrow & \mathbf{R}^r j_* \mathbf{Z}/p^n(r)_U \\ \uparrow & \nearrow & \downarrow & & \downarrow \\ M_X^{\text{gp}} \otimes (\widetilde{M}_X^{\text{gp}})^{\otimes (r-1)} & & & & \\ \uparrow \kappa & & & & \\ (\widetilde{M}_X^{\text{gp}})^{\otimes (r-1)} & & \mathbf{R}^{r-1} j_{\text{ét}*} \mathbf{R}^1 j'_* \mathbf{Z}/p^n(r)_U & & \\ \downarrow & & \downarrow \wr & \searrow \text{res}_Y & \\ (\widetilde{M}_Y^{\text{gp}})^{\otimes (r-1)} & \longrightarrow & \mathbf{R}^{r-1} j_{\text{ét}*} \mathbf{Z}/p^n(r-1)_{Y_K} & & \end{array}$$

The map κ is defined by sending the section m to $T \otimes m$. We claim that the symbol map

$$(\widetilde{M}_Y^{\text{gp}})^{\otimes (r-1)} \rightarrow \mathbf{R}^{r-1}(j_{\text{ét}}\varepsilon)_* \mathbf{Z}/p^n(r-1)_{Y_K}$$

is surjective. Since we have Gersten conjecture for Milnor K-theory [24, Theorem 7.1] and étale cohomology, this follows from Bloch-Kato conjecture and [24, Lemma 7.2]. Since the map $\widetilde{M}_X^{\text{gp}} \rightarrow \widetilde{M}_Y^{\text{gp}}$ is surjective in the Zariski topology on X , the above diagram shows surjectivity of the map (6.7) we wanted. The argument for the map (6.8) is analogous.

Assume now that X is semistable over \mathcal{O}_K with a smooth special fiber. Consider the following diagram

$$\begin{array}{ccccc}
\mathcal{H}^r(\tilde{S}_n(r)_X) & \longrightarrow & \mathcal{H}^r(\tilde{S}_n(r)_X(Y)) & \xrightarrow{\text{res}_Y} & i_*\mathcal{H}^{r-1}(\tilde{S}_n(r-1)_Y) \\
\downarrow & & \downarrow & & \downarrow \\
\mathcal{H}^r(S_n(r)_X) & \longrightarrow & \mathcal{H}^r(S_n(r)_X(Y)) & \xrightarrow{\text{res}_Y} & i_*\mathcal{H}^{r-1}(S_n(r-1)_Y) \\
\downarrow \beta & & \downarrow \beta & & \downarrow \beta \\
i_*W_n\Omega_{X_0,\log}^{r-1} & \hookrightarrow & i_*W_n\Omega_{U_0,\log}^{r-1} & \xrightarrow{\text{res}_Y} & i_*W_n\Omega_{Y_0,\log}^{r-2}
\end{array}$$

The horizontal sequences are Gysin sequences. The syntomic cohomology \tilde{S} uses the vertical log-structure, the syntomic cohomology S uses the trivial log-structure. The vertical sequences are exact [25, 1] (the map β is the residue at ϖ map). We claim that the diagram commutes. Indeed, for the top part this follows from functoriality. For the bottom part, the left square commutes by functoriality. For the right square, it suffices to look at symbols. It follows from the definition that the map β sends the symbol $\{f_1, \dots, f_r\}$, $f_i \in i_0^*\mathcal{O}_X^*$, $i_0 : X_0 \hookrightarrow X$, to zero and the symbol $\{f_1, \dots, f_{r-1}, \varpi\}$, $f_i \in i_0^*\mathcal{O}_X^*$, to $\text{dlog}[\bar{f}_1] \wedge \dots \wedge \text{dlog}[\bar{f}_{r-1}]$. The definition of the residue map res_Y is similar with ϖ replaced by T . The commutativity we want is now immediate.

(2b) *The general case.*

In the case of complexes $E_n(r)'_{\text{Nis}}$, we construct the triangle

$$(6.9) \quad E_n(r)'_{\text{Nis},X}(D) \rightarrow E_n(r)'_{\text{Nis},X}(Y \cup D) \xrightarrow{\text{res}_Y} i_*F_n^1(r-1)_X(Y \cup D)[-1]$$

by truncating the complexes in the Nisnevich version of the distinguished triangle (6.1) and by setting $F_n^1(r-1)_X(Y \cup D)$ equal to the image of the composition

$$(\tau_{\leq r}\varepsilon_*GE'_n(r)_X(Y \cup D)_{\text{Két}})[1] \rightarrow (\tau_{\leq r}\varepsilon_*GC(f, r-1))[1].$$

Clearly the obtained triangle is distinguished. It remains to show that the natural injection

$$F_n^1(r-1)_X(Y \cup D) \hookrightarrow (\tau_{\leq r}\varepsilon_*GC(f, r-1))[1]$$

has an inverse up to $p^{N(r-1)}$. Since $C(f, r-1) \simeq i_*E_n^1(r-1)_Y(D_Y)[-1]$, we have

$$\begin{aligned}
\tau_{\leq r-2}F_n^1(r-1)_X(Y \cup D) &\simeq \tau_{\leq r-1}\varepsilon_*GC(f, r-1), \\
\mathcal{H}^r\varepsilon_*GC(f, r-1) &\simeq i_*\mathcal{H}^{r-1}E_n^1(r-1)_Y(D_Y).
\end{aligned}$$

It suffices thus to show that the map

$$\text{res}_Y : \mathcal{H}^r(E'_n(r)_{\text{Nis},X}(Y \cup D)) \rightarrow i_*\mathcal{H}^{r-1}(E_n^1(r-1)_{\text{Nis},Y}(D_Y))$$

has a cokernel annihilated by $p^{N(r-1)}$. But this follows just as in the case $r \leq p-2$ treated above using Theorem 2.2 and, in the good reduction case, [7, Theorem 3.2]. \square

Gysin distinguished triangles are functorial for certain morphisms of log-schemes.

Lemma 6.2. *For Y , X , and D as above, let $f : X' \rightarrow X$ be a flat morphism or a closed immersion such that X' is regular and syntomic over $W(k)$ and $f^{-1}(Y \cup D)$ is a relative simple normal crossing divisor on X' over $W(k)$. Set $Y' = f^{-1}(Y)$ and $D' = f^{-1}(D)$. Then, for $r \geq 1$, $i \geq 0$, we have a map of distinguished triangles*

$$\begin{array}{ccccc}
i_*S_n^{i+1}(r-1)_Y(D_Y)[-2] & \xrightarrow{i_!} & S_n^i(r)_X(D) & \longrightarrow & S_n^i(r)_X(Y \cup D) \\
\downarrow f^* & & \downarrow f^* & & \downarrow f^* \\
\text{R}f_*i_*S_n^{i+1}(r-1)_{Y'}(D'_{Y'})[-2] & \xrightarrow{i_!} & \text{R}f_*S_n^i(r)_{X'}(D') & \longrightarrow & \text{R}f_*S_n^i(r)_{X'}(Y' \cup D')
\end{array}$$

Similarly, for the complexes $S_n(r)_X(D)$, $r \leq p-1$.

There are analogous maps of Gysin distinguished triangles for the complexes $E_n(r)$ and $E_n(r)_{\text{Nis}}$ (and $r \leq p-2$) as well as for the complexes $E'_n(r)$. For the complexes $E'_n(r)_{\text{Nis}}$ we have the following maps of Gysin distinguished triangles

$$\begin{array}{ccccc} i_* F_n^1(r-1)_X(Y \cup D)[-2] & \xrightarrow{i_i} & E'_n(r)_{\text{Nis},X}(D) & \longrightarrow & E'_n(r)_{\text{Nis},X}(Y \cup D) \\ \downarrow f^* & & \downarrow f^* & & \downarrow f^* \\ \text{Rf}_* i_* F_n^1(r-1)_{X'}(Y' \cup D')[-2] & \xrightarrow{i_i} & \text{Rf}_* E'_n(r)_{\text{Nis},X'}(D') & \longrightarrow & \text{Rf}_* E'_n(r)_{\text{Nis},X'}(Y' \cup D') \end{array}$$

Proof. The case of the map f being flat follows from the flat base change. The case of f being a closed immersion follows from the fact that the morphisms $X' \rightarrow X$ and $Y \rightarrow X$ are Tor-independent. \square

6.1.2. *Projection formula.* We have the following projection formula for a closed immersion of codimension one.

Lemma 6.3. *Let X, Y , and D be as above. Then, for $r_1, r_2, i, j \geq 0$, the following diagram commutes.*

$$\begin{array}{ccc} S_n^i(r_1)_X(D) \otimes^{\mathbb{L}} i_* S_n^{j+1}(r_2-1)_Y(D_Y)[-2] & \xrightarrow{1 \otimes i_i} & S_n^i(r_1)_X(D) \otimes^{\mathbb{L}} S_n^j(r_2)_X(D) \\ \downarrow i^* \otimes 1 & & \downarrow \cup \\ i_* S_n^i(r_1)_Y(D_Y) \otimes^{\mathbb{L}} i_* S_n^{j+1}(r_2-1)_Y(D_Y)[-2] & & \downarrow \cup \\ \downarrow \cup & & \downarrow \cup \\ i_* S_n^{i+j+1}(r_1+r_2-1)_Y(D_Y)[-2] & \xrightarrow{i_i} & S_n^{i+j}(r_1+r_2)_X(D) \end{array}$$

Similarly, for the complexes $S_n(r)_X(D)$, $r \leq p-1$.

There are analogous projection formulas for the complexes $E_n(r)$ and $E_n(r)_{\text{Nis}}$ (and $r \leq p-2$) as well as for the complexes $E'_n(r)$. For the complexes $E'_n(r)_{\text{Nis}}$ we have the following commutative diagram (the constant N is the one from Theorem 2.2)

$$(6.10) \quad \begin{array}{ccc} E'_n(r_1)_{\text{Nis},X}(D) \otimes^{\mathbb{L}} i_* F_n^1(r_2-1)_X(Y \cup D)[-2] & \xrightarrow{p^{N(r_1+r_2-1)} \otimes i_i} & E'_n(r_1)_{\text{Nis},X}(D) \otimes^{\mathbb{L}} E'_n(r_2)_{\text{Nis},X}(D) \\ \downarrow i^* \otimes 1 & & \downarrow \cup \\ i_* E'_n(r_1)_{\text{Nis},Y}(D_Y) \otimes^{\mathbb{L}} i_* F_n^1(r_2-1)_X(Y \cup D)[-2] & & \downarrow \cup \\ \downarrow \cup & & \downarrow \cup \\ i_* F_n^1(r_1+r_2-1)_X(Y \cup D)[-2] & \xrightarrow{i_i} & E'_n(r_1+r_2)_{\text{Nis},X}(D) \end{array}$$

Proof. For the first claim of the lemma, it suffices to show that the localization short exact sequence

$$0 \rightarrow S_n^j(r_2)_X(D) \xrightarrow{f} S_n^j(r_2)_X(Y \cup D) \xrightarrow{\text{res}} i_* S_n^{j+1}(r_2-1)_Y(D_Y)[-1] \rightarrow 0$$

is compatible with the action of $S_n^i(r_1)_X(D)$. For the localization map f this follows from the functoriality of products; for the residue map it will follow if we show that the following diagram of complexes commutes.

$$\begin{array}{ccc} S_n^j(r_2)_X(D \cup Y) \otimes S_n^i(r_1)_X(D) & \xrightarrow{1 \otimes f} & S_n^j(r_2)_X(D \cup Y) \otimes S_n^i(r_1)_X(D \cup Y) \\ \downarrow \text{res} \otimes 1 & & \downarrow \cup \\ i_* S_n^{j+1}(r_2-1)_Y(D_Y)[-1] \otimes S_n^i(r_1)_X(D) & & S_n^{i+j}(r_2+r_1)_X(D \cup Y) \\ \downarrow 1 \otimes i^* & & \downarrow \text{res} \\ i_* S_n^{i+j+1}(r_2-1)_Y(D_Y)[-1] \otimes i_* S_n^i(r_1)_Y(D_Y) & \xrightarrow{\cup} & i_* S_n^{i+j+1}(r_1+r_2-1)_Y(D_Y)[-1] \end{array}$$

But this can be checked locally (in the lifted situation) where it follows immediately from the formulas for product. The argument for the complexes $S_n(r)_X(D)$ is basically the same.

For the complexes $E'_n(r)$, it suffices to argue on the étale site. Assume thus that X is equipped with the étale topology. Consider the following commutative diagram of short exact sequences that appears in the proof of Lemma 6.1 (diagram (6.2)). Below we will use the notation from that proof freely.

$$\begin{array}{ccccccc} 0 & \longrightarrow & E'_n(r_2)_X(D) & \xrightarrow{f} & E'_n(r_2)_X(Y \cup D)_{\text{Két}} & \xrightarrow{\text{res}} & C(f, r_2 - 1) \longrightarrow 0 \\ & & \downarrow i^* & & \downarrow i^* & & \downarrow i^* \\ 0 & \longrightarrow & i_* E'_n(r_2)_Y(D_Y) & \xrightarrow{f'} & i_* E'_n(r_2)_Y(D_Y^0)_{\text{Két}} & \xrightarrow{\text{res}} & i_* C(f', r_2 - 1) \longrightarrow 0 \end{array}$$

Proceeding as in the definition of cup product on the complexes $E'_n(r)$, we define an action of $E'_n(r_1)_X(D)$ on the first two terms of the top exact sequence; hence an action on $C(f, r_2 - 1)$. In a similar way we can define a compatible action of $i_* E'_n(r_2)_Y(D_Y)$ on the first two terms of the bottom exact sequence; hence an action on $i_* C(f', r_2 - 1)$. It follows immediately that these two actions are compatible with the above map between exact sequences.

It follows that if we define the product

$$\cup : i_* E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} C(f, r_2 - 1) \xrightarrow{1 \otimes i^*} i_* E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} C(f', r_2 - 1) \xrightarrow{\cup} C(f', r_1 + r_2 - 1) \xrightarrow{\simeq} C(f, r_1 + r_2 - 1),$$

from the action of $E'_n(r_1)_X(D)$ on the top exact sequence, we obtain the following projection formula

$$\begin{array}{ccc} E'_n(r_1)_X(D) \otimes^{\mathbb{L}} C(f, r_2 - 1)[-1] & \xrightarrow{1 \otimes \partial} & E'_n(r_1)_X(D) \otimes^{\mathbb{L}} E'_n(r_2)_X(D) \\ \downarrow i^* \otimes 1 & & \downarrow \cup \\ i_* E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} C(f, r_2 - 1)[-1] & & \\ \downarrow \cup & & \downarrow \partial \\ C(f, r_1 + r_2 - 1)[-1] & \xrightarrow{\partial} & E'_n(r_1 + r_2)_X(D) \end{array}$$

Recall that we have the quasi-isomorphisms

$$\gamma' : C(f, r_2 - 1) \xrightarrow{\simeq} i_* C(f', r_2 - 1) \xrightarrow{\simeq} i_* E'_n(r_2 - 1)_Y(D_Y)[-1]$$

Thus to obtain the projection formula in the statement of the lemma it suffices to show that the following diagram commutes in the derived category.

$$\begin{array}{ccc} i_* E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} C(f, r_2 - 1) & \xrightarrow{\cup} & C(f, r_1 + r_2 - 1) \\ \downarrow 1 \otimes \gamma' & & \downarrow \gamma' \\ i_* E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} i_* E'_n(r_2 - 1)_Y(D_Y)[-1] & \xrightarrow{\cup} & i_* E'_n(r_1 + r_2 - 1)_Y(D_Y)[-1] \end{array}$$

Or, unwinding the action of $i_* E'_n(r_1)_Y(D_Y)$ on $C(f, r_2 - 1)$, that the following diagram commutes in the derived category.

$$(6.11) \quad \begin{array}{ccc} E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} C(f', r_2 - 1) & \xrightarrow{\cup} & C(f', r_1 + r_2 - 1) \\ \downarrow 1 \otimes \gamma & & \downarrow \gamma \\ E'_n(r_1)_Y(D_Y) \otimes^{\mathbb{L}} E'_n(r_2 - 1)_Y(D_Y)[-1] & \xrightarrow{\cup} & E'_n(r_1 + r_2 - 1)_Y(D_Y)[-1] \end{array}$$

To see this consider the commutative diagrams (6.3) and (6.4) that define the map γ . Set $r = r_2$. Consider also their analogs built on top of the map

$$\begin{array}{c} \Gamma^h(\overline{S}_n(r_1)_{Y, Z_1}[-1]) \xrightarrow{\text{Id}} \Gamma^h(\overline{S}_n(r_1)_{Y, Z_1}[-1]) \xleftarrow{\simeq} \Gamma^h(\Lambda(r_1)) \xleftarrow{\simeq} \Gamma(\Lambda(r_1)) \xleftarrow{\simeq} \\ \Gamma(\text{Cone}(\eta' GA(r_1))) \rightarrow \Gamma(\mathcal{J}_{\text{ét}}! j'_* GA(r_1)) \end{array}$$

as well as the extension

$$\Gamma(S'_n(r_1)_{Y,Z_1}[-1]) \rightarrow \Gamma^h(\overline{S}_n(r_1)_{Y,Z_1}[-1])$$

We claim that there exists a pairing of the above two sets of extended diagrams into the extended diagrams (6.3) and (6.4) for $r = r_1 + r_2$ that matches corresponding nodes and all the maps are compatible with pairings. Indeed, by functoriality, it suffices to construct such a pairing for the bottom rows. And there it can be checked easily using the fact that the complexes L' and C' are defined as cokernels of maps that are compatible with certain obvious pairings. It is easy to see that this pairing of diagrams induces a quasi-isomorphism of pairings (6.11) that we wanted.

For the complexes $E'_n(r)_{\text{Nis}}$, consider the following commutative diagram of short exact sequences that is a truncated Zariski version of a diagram that appears in the proof of Lemma 6.1 (diagram (6.2)).

$$\begin{array}{ccccccc} 0 & \longrightarrow & E'_n(r_2)_{\text{Nis},X}(D) & \xrightarrow{f} & \tau_{\leq r_2} \pi_* G E'_n(r_2)_X(Y \cup D)_{\text{Két}} & \xrightarrow{res} & i_* F_n^1(r_2 - 1)_X(Y \cup D) & \longrightarrow & 0 \\ & & \downarrow i^* & & \downarrow i^* & & \downarrow i^* & & \\ 0 & \longrightarrow & i_* E'_n(r_2)_{\text{Nis},Y}(D_Y) & \xrightarrow{f'} & \tau_{\leq r_2} \pi_* G i_* E'_n(r_2)_Y(D_Y^0)_{\text{Két}} & \xrightarrow{res} & i_* \widetilde{F}_n^1(r_2 - 1)_X(Y \cup D) & \longrightarrow & 0 \end{array}$$

Here $i_* \widetilde{F}_n^1(r_2 - 1)_X(Y \cup D)$ is defined as the cokernel of the map f' . Proceeding as above we define an action of $E'_n(r_1)_{\text{Nis},X}(D)$ on the first two terms of the top exact sequence; hence an action on $i_* F_n^1(r_2 - 1)_X(Y \cup D)$. In a similar way we can define a compatible action of $i_* E'_n(r_2)_{\text{Nis},Y}(D_Y)$ on the first two terms of the bottom exact sequence; hence an action on $i_* \widetilde{F}_n^1(r_2 - 1)_X(Y \cup D)$. Clearly these two actions are compatible with the above map between exact sequences and with the action on $i_* E_n^1(r_2 - 1)_{\text{Nis},X}(Y \cup D)[-2]$.

We set the product

$$\cup : i_* E'_n(r_1)_{\text{Nis},Y}(D_Y) \otimes^{\mathbb{L}} i_* F_n^1(r_2 - 1)_X(Y \cup D)[-2] \rightarrow i_* F_n^1(r_1 + r_2 - 1)_X(Y \cup D)[-2]$$

to be equal to the composition

$$\begin{aligned} i_* E'_n(r_1)_{\text{Nis},Y}(D_Y) \otimes^{\mathbb{L}} i_* F_n^1(r_2 - 1)_X(Y \cup D)[-2] &\rightarrow i_* E'_n(r_1)_{\text{Nis},Y}(D_Y) \otimes^{\mathbb{L}} i_* E_n^1(r_2 - 1)_{\text{Nis},X}(Y \cup D)[-2] \\ &\xrightarrow{\cup} i_* E_n^1(r_1 + r_2 - 1)_{\text{Nis},X}(Y \cup D)[-2] \xrightarrow{\beta} i_* F_n^1(r_1 + r_2 - 1)_X(Y \cup D)[-2], \end{aligned}$$

where β is an inverse (in the derived category and up to $p^{N(r_1+r_2-1)}$) of the natural map $F_n^1(r_2 - 1)_X(Y \cup D)[-2] \rightarrow E_n^1(r_1 + r_2 - 1)_{\text{Nis},X}(Y \cup D)[-2]$.

To show that the diagram (6.10) commutes it suffices to show it with the lower right-hand term $E'_n(r_1 + r_2)_{\text{Nis},X}(D)$ replaced by $E'_n(r_1 + r_2)_X(D)$. This is by degree reason: cohomology of $E'_n(r_1)_{\text{Nis},X}(D) \otimes^{\mathbb{L}} i_* F_n^1(r_2 - 1)_X(Y \cup D)[-2]$ is concentrated in degrees $[0, r_1 + r_2 + 1]$ and cohomology of $E'_n(r_1 + r_2)_{\text{Nis},X}(D)$ is concentrated in degrees $[0, r_1 + r_2]$; hence the map

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(E'_n(r_1)_{\text{Nis},X}(D) \otimes^{\mathbb{L}} i_* F_n^1(r_2 - 1)_X(Y \cup D)[-2], E'_n(r_1 + r_2)_{\text{Nis},X}(D)) \\ \rightarrow \text{Hom}_{\mathcal{D}}(E'_n(r_1)_{\text{Nis},X}(D) \otimes^{\mathbb{L}} i_* F_n^1(r_2 - 1)_X(Y \cup D)[-2], E'_n(r_1 + r_2)_X(D)) \end{aligned}$$

between homomorphism groups in the derived category is injective. Now a simple diagram chase reduces this projection formula diagram to the one for $E'_n(r)_X$, which we have just proved. \square

Lemma 6.4. *Let X, Y, D be as above. Then there exists a map of distinguished triangles*

$$\begin{array}{ccccc} j_* \mathcal{O}_{X \setminus D}^*[-1] & \longrightarrow & j'_* \mathcal{O}_{X \setminus (D \cup Y)}^*[-1] & \xrightarrow{\text{ord}_Y} & i_* \mathbf{Z}_Y[-1] \\ \downarrow c_1^{\text{syn}} & & \downarrow c_1^{\text{syn}} & & \downarrow c_0 \\ S'_n(1)_X(D) & \longrightarrow & S'_n(1)_X(Y \cup D) & \xrightarrow{\text{res}_Y} & i_* S_n^1(0)_Y(D_Y)[-1] \end{array}$$

Here $j : X \setminus D \hookrightarrow X$ and $j' : X \setminus (D \cup Y) \hookrightarrow X$ are the natural open immersions and the map c_0 on the right is defined by the natural morphism $\mathbf{Z}_Y \rightarrow \mathcal{H}^0(S^1(0)_Y(D_Y))$. Similarly for the Chern classes with values in $S_n(1)_X(D)$.

We have an analogous map of distinguished triangles for $E_n(1)_X(D)$ and $E'_n(1)_X(D)$ as well as the induced map of distinguished triangles

$$\begin{array}{ccccc} j_* \mathcal{O}_{X \setminus D}^*[-1] & \longrightarrow & j'_* \mathcal{O}_{X \setminus (D \cup Y)}^*[-1] & \xrightarrow{\text{ord}_Y} & i_* \mathbf{Z}_Y[-1] \\ \downarrow c_1^{\text{syn}} & & \downarrow c_1^{\text{syn}} & & \downarrow c_0 \\ E'_n(1)_{\text{Nis}, X}(D) & \longrightarrow & E'_n(1)_{\text{Nis}, X}(Y \cup D) & \xrightarrow{\text{res}_Y} & i_* F_n^1(0)_X(Y \cup D)[-1] \end{array}$$

Proof. The statements for S'_n and S_n follow immediately from the above formulas. Since $\mathcal{H}^0(E_n^1(0)_Y(D_Y)) \hookrightarrow \mathcal{H}^0(S_n^1(0)_Y(D_Y))$ the statements for E'_n and E_n follow as soon as we define the map $c_0 : \mathbf{Z}_Y \rightarrow \mathcal{H}^0(E_n^1(0)_Y(D_Y))$ that is compatible with the map $\mathbf{Z}_Y \rightarrow \mathcal{H}^0(S^1(0)_Y(D_Y))$. For that notice that the composition

$$\mathbf{Z}_Y \xrightarrow{c_0} S^1(0)_Y(D_Y) \xrightarrow{\alpha} i_{Y*} i_Y^* \mathbf{R}j_{Y*} \mathbf{Z}/p^n(1)'(-1)$$

is compatible with the natural map $pc_0^{\text{ét}} : \mathbf{Z}_Y \rightarrow \mathbf{R}j_{Y*} \mathbf{Z}/p^n(1)'(-1)$. Hence the composition

$$\mathbf{Z}_Y \xrightarrow{c_0} S^1(0)_Y(D_Y) \xrightarrow{\alpha} i_{Y*} i_Y^* \mathbf{R}j_{Y*} \mathbf{Z}/p^n(1)'(-1) \rightarrow j_{Y \text{ ét}!} \mathbf{R}j'_{Y*} \mathbf{Z}/p^n(1)'(-1)[1]$$

is homotopic to zero and we obtain its unique factorization $c_0 : \mathbf{Z}_Y \rightarrow E_n^1(0)_Y(D_Y)$. It is clear that this map factors through $F_n^1(0)_X(Y \cup D)$ hence the last statement of the lemma. \square

6.1.3. Projective space theorem and homotopy invariance. We will now discuss certain versions of projective space theorem for syntomic cohomology. For large primes p , we have the following projective space theorem.

Lemma 6.5. *Let X and $D := D'$ be as above. Let \mathcal{E} be a locally free sheaf of rank $d + 1$, $d \geq 0$. Consider the associated projective bundle $\pi : \mathbf{P}(\mathcal{E}) \rightarrow X$. Then, we have the following quasi-isomorphism*

$$\bigoplus_{i=0}^d c_1^{\text{syn}}(\mathcal{O}(1))^i \cup \pi^* : \bigoplus_{i=0}^d S_n(r-i)_X(D)[-2i] \xrightarrow{\sim} \mathbf{R}\pi_* S_n(r)_{\mathbf{P}(\mathcal{E})}(\pi^{-1}(D)), \quad 0 \leq d \leq r \leq p-1.$$

Here, the class $c_1^{\text{syn}}(\mathcal{O}(1)) \in H^2(\mathbf{P}(\mathcal{E}), S_n(1)(\pi^{-1}(D)))$ refers to the class of the tautological bundle on $\mathbf{P}(\mathcal{E})$.

We have an analogous quasi-isomorphism for complexes $E_n(r)$.

Proof. In the absence of the divisor D , for the complexes $S_n(r)$ this follows easily from the (filtered) projective space theorem for crystalline cohomology and the compatibility of the actions of syntomic Chern classes with the cohomology long exact sequence associated to (2.1) [15, I.4.3]. For complexes $E_n(r)$, this follows from the above and the projective space theorem for $j_{\text{ét}!} \mathbf{R}j'_* \mathbf{Z}/p^n(r)$.

In general, write $D = \cup_{i=1}^m D_i$, $Y = D_1$, $D' = \cup_{i=2}^m D_i$, $D_Y = D' \cap Y$, where D_i is an irreducible component of D . We will argue by induction on m ; the case of $m = 0$ being known. Using Lemma 6.1, we get the following map of distinguished triangles

$$\begin{array}{ccc} \bigoplus_{i=0}^d i_* S_n(r-1-i)_Y(D_Y)[-2i-2] & \xrightarrow[\sim]{\bigoplus_{i=0}^d \xi_Y(D_Y)^i \cup \pi^*} & i_* \mathbf{R}\pi_* S_n(r-1)_{\mathbf{P}(\mathcal{E})_Y}(\pi^{-1}(D_Y))[-2] \\ \downarrow i_! & & \downarrow i_! \\ \bigoplus_{i=0}^d S_n(r-i)_X(D)[-2i] & \xrightarrow{\bigoplus_{i=0}^d \xi_X(D)^i \cup \pi^*} & \mathbf{R}\pi_* S_n(r)_{\mathbf{P}(\mathcal{E})}(\pi^{-1}(D)) \\ \downarrow & & \downarrow \\ \bigoplus_{i=0}^d S_n(r-i)_X(D')[-2i] & \xrightarrow[\sim]{\bigoplus_{i=0}^d \xi_X(D')^i \cup \pi^*} & \mathbf{R}\pi_* S_n(r)_{\mathbf{P}(\mathcal{E})}(\pi^{-1}(D')) \end{array}$$

where $i : Y \hookrightarrow X$ is the natural immersion and we wrote $\xi_X(D), \xi_Y(D_Y), \xi_X(D')$ for the Chern classes of the tautological bundle in $H^2(\mathbf{P}(\mathcal{E})_Y, S_n(1)(\pi^{-1}(D_Y)))$, $H^2(\mathbf{P}(\mathcal{E}), S_n(1)(\pi^{-1}(D)))$, and $H^2(\mathbf{P}(\mathcal{E}), S_n(1)(\pi^{-1}(D')))$, respectively. The upper square commutes by the projection formula from Lemma 6.3 and Tor-independence from Lemma 6.2.

By the inductive assumption the top and bottom horizontal arrows are quasi-isomorphisms. So is then the middle one, as wanted.

The argument for $E_n(r)$ is analogous. \square

As a corollary we get the following syntomic cohomology version of homotopy invariance.

Lemma 6.6. *Let X, D be as in Lemma 6.5. Let \mathcal{E} be a locally free sheaf of rank $d + 1$, $d \geq 0$. Consider its projectivization $\pi : \mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X) \rightarrow X$ and its hyperplane at infinity H_∞ . Then, for $a \geq 0$, we have the following quasi-isomorphisms.*

$$\begin{aligned} \pi^* : S_n^a(r)_X(D) &\xrightarrow{\sim} R\pi_* S_n^a(r)_{\mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)}(\pi^{-1}(D) \cup H_\infty), \quad 0 \leq r, \\ \pi^* : S_n(r)_X(D) &\xrightarrow{\sim} R\pi_* S_n(r)_{\mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)}(\pi^{-1}(D) \cup H_\infty), \quad 0 \leq d \leq r \leq p-1. \end{aligned}$$

We have analogous quasi-isomorphisms for the complexes $E_n(r)$, $E'_n(r)$, and $E_n^1(r)$.

Proof. The second quasi-isomorphism follows from the projective space theorem from Lemma 6.5. Indeed, consider the following commutative diagram of distinguished triangles.

$$\begin{array}{ccccc} R\pi_* i_* S_n(r-1)_{H_\infty}(\pi^{-1}(D) \cap H_\infty)[-2] & \xrightarrow{i_!} & R\pi_* S_n(r)_{\mathbf{P}}(\pi^{-1}(D)) & \longrightarrow & R\pi_* S_n(r)_{\mathbf{P}}(\pi^{-1}(D) \cup H_\infty) \\ \uparrow i^* \pi^* & & \parallel & & \parallel \\ Rq_* S_n(r-1)_{\mathbf{P}(\mathcal{E})}(q^{-1}(D))[-2] & \xrightarrow{\xi \cup \tilde{\pi}^*} & R\pi_* S_n(r)_{\mathbf{P}}(\pi^{-1}(D)) & \longrightarrow & R\pi_* S_n(r)_{\mathbf{P}}(\pi^{-1}(D) \cup H_\infty) \\ \uparrow \oplus_i \xi_1^i \cup q^* & & \parallel & & \uparrow \pi^* \\ \oplus_{i=0}^{i=d-1} S_n(r-1-i)_X(D)[-2-2i] & \xrightarrow{\xi^{i+1} \cup \pi^*} & R\pi_* S_n(r)_{\mathbf{P}}(\pi^{-1}(D)) & \xrightarrow{p_1} & S_n(r)_X(D), \end{array}$$

where we put $\xi = c_1^{\text{syn}}(\mathcal{O}(H_\infty))$, $\xi_1 = c_1^{\text{syn}}(\mathcal{O}(1)) \in H^2(\mathbf{P}(\mathcal{E}), S_n(1)(q^{-1}(D)))$, $\mathbf{P} = \mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)$. We have $\xi = \tilde{\pi}^*(\xi)$. We named the maps as in the following diagram

$$\begin{array}{ccc} H_\infty & \xrightarrow{i} & \mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X) \\ & \searrow \sim & \downarrow \tilde{\pi} \\ & & \mathbf{P}(\mathcal{E}) \xrightarrow{q} X \end{array}$$

The top row in the big diagram is the Gysin sequence. The upper left square commutes because by the projection formula from Lemma 6.3 and by Lemma 6.4 we have

$$i_! i^* \pi^* = i_!(1) \cup \pi^* = c_1^{\text{syn}}(\mathcal{O}(H_\infty)) \cup \pi^*.$$

In the bottom row, the left vertical map is a quasi-isomorphism by the projective space theorem for $q : \mathbf{P}(\mathcal{E}) \rightarrow X$ (Lemma 6.5) and the map p_1 comes from the same projective space theorem. The pullback map π^* is its right inverse. The second quasi-isomorphism of the lemma follows now easily.

For the first quasi-isomorphism consider the following commutative diagram of distinguished triangles.

$$\begin{array}{ccccc} R\pi_* S_n^a(r)_{\mathbf{P}}(\pi^{-1}(D) \cup H_\infty) & \longrightarrow & R\pi_* \mathcal{J}_{\mathbf{P},n}^{[r]}(\pi^{-1}(D) \cup H_\infty) & \xrightarrow{p^a(p^r-1)\varphi} & R\pi_* \mathcal{O}_{\mathbf{P},n}^{\text{cr}}(\pi^{-1}(D) \cup H_\infty) \\ \uparrow \pi^* & & \uparrow \pi^* & & \uparrow \pi^* \\ S_n^a(r)_X(D) & \longrightarrow & \mathcal{J}_{X,n}^{[r]}(D) & \xrightarrow{p^a(p^r-1)\varphi} & \mathcal{O}_{X,n}^{\text{cr}}(D). \end{array}$$

Here we put $\mathcal{J}_n^{[r]} = R\varepsilon_* \mathcal{J}_n^{[r]}$ and $\mathcal{O}_n^{\text{cr}} = R\varepsilon_* \mathcal{O}_n^{\text{cr}}$, where ε is the natural projection from the syntomic to the Zariski site. The two marked maps are quasi-isomorphisms because we have the (filtered) projective space theorem for crystalline cohomology and we can use the same argument as we did above to prove the second quasi-isomorphism of the lemma. The first quasi-isomorphism of the lemma follows.

The arguments for the complexes $E_n(r)$, $E'_n(r)$, and $E_n^1(r)$ are analogous (but use also the proper base change theorem and homotopy property of étale cohomology). \square

For a general prime p we still have a cell decomposition of the cohomology of the projective bundle. We will prove a first step of it.

Lemma 6.7. *Let X, D be as in Lemma 6.5. Let \mathcal{E} be a locally free sheaf of rank $d + 1$, $d \geq 0$. Consider its projectivization $\pi : \mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X) \rightarrow X$ and its hyperplane at infinity H_∞ . Then, for $a \geq 0$, we have the following quasi-isomorphism*

$$i_! \oplus \pi^* : \mathbf{R}\pi_* i_* S_n^{a+1}(r-1)_{H_\infty}(\pi^{-1}(D) \cap H_\infty)[-2] \oplus S_n^a(r)_X(D) \xrightarrow{\sim} \mathbf{R}\pi_* S_n^a(r)_{\mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)}(\pi^{-1}(D)), \quad 1 \leq r.$$

Here $i : H_\infty \hookrightarrow \mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)$ is the natural closed immersion.

In particular, we have the following quasi-isomorphism

$$i_!(\pi i)^* \oplus \pi^* : S_n^{a+1}(r-1)_X(D)[-2] \oplus S_n^a(r)_X(D) \xrightarrow{\sim} \mathbf{R}\pi_* S_n^a(r)_{\mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)}(\pi^{-1}(D)), \quad 1 \leq r.$$

Similarly for the complexes $E'_n(r)$ and $E_n^1(r)$.

Proof. We have the following commutative diagram of distinguished triangles.

$$\begin{array}{ccccc} \mathbf{R}\pi_* i_* S_n^{a+1}(r-1)_{H_\infty}(D_\infty)[-2] & \xrightarrow{i_!} & \mathbf{R}\pi_* S_n^a(r)_{\mathbf{P}}(\pi^{-1}(D)) & \longrightarrow & \mathbf{R}\pi_* S_n^a(r)_{\mathbf{P}}(\pi^{-1}(D) \cup H_\infty) \\ \parallel & & \uparrow i_! \oplus \pi^* & & \uparrow \pi^* \wr i \\ \mathbf{R}\pi_* i_* S_n^{a+1}(r-1)_{H_\infty}(D_\infty)[-2] & \longrightarrow & \mathbf{R}\pi_* i_* S_n^{a+1}(r-1)_{H_\infty}(D_\infty)[-2] \oplus S_n^a(r)_X(D) & \longrightarrow & S_n^a(r)_X(D) \end{array}$$

where we put $\mathbf{P} = \mathbf{P}(\mathcal{E} \oplus \mathcal{O}_X)$, $D_\infty = \pi^{-1}(D) \cap H_\infty$. The top triangle is induced by the Gysin sequence. The right vertical arrow is a quasi-isomorphism by Lemma 6.6. The first statement of the lemma follows.

For the second, notice that the natural projection $\pi i : H_\infty \rightarrow X$ is an isomorphism and we have a quasi-isomorphism

$$(\pi i)^* : S_n^{a+1}(r-1)_X(D) \xrightarrow{\sim} \mathbf{R}\pi_* i_* S_n^{a+1}(r-1)_{H_\infty}(D_\infty).$$

The arguments for the complexes $E'_n(r)$ and $E_n^1(r)$ are analogous. \square

6.2. Compatibility with Gysin sequences. We will show in this section that the syntomic universal Chern class maps are compatible with Gysin sequences. We will present the arguments for E' -syntomic-étale cohomology. In the case of E -syntomic-étale cohomology the arguments are very similar.

Let X be a semistable scheme over \mathcal{O}_K^\times or a semistable scheme \mathcal{O}_K with smooth special fiber. Let D be the horizontal divisor. Assume that $D = \cup_{i=1}^m D_i$, $m \geq 1$, is a union of m irreducible components D_i . Note that each scheme D_i with the induced log-structure is of the same type as X (with at most $m - 1$ components in the divisor at infinity). Set $D' = \cup_{i=2}^m D_i$, $i : D_1 \hookrightarrow X$, $D'_1 = D_1 \cap D'$. The pairs (X, D') and (D_1, D'_1) are of the same type as the pair (X, D) we started with but with at most $m - 1$ irreducible divisors at infinity. Fix N as in Lemma 5.6. Consider the following Gysin-type diagram with columns being homotopy cofiber sequences

$$(6.12) \quad \begin{array}{ccc} i_* G_{D_1}(D'_1) & \xrightarrow{[p^{N^i}][p^{N(i+1)}]\gamma'[p^{2N}]P(D_1/X; \tilde{C}_{D_1 \setminus D'_1}^{\text{syn}})} & \{0\} \times \prod_{i \geq 1} \mathcal{K}(2i - 2, i_* \tilde{F}_n^1(i-1)_X(D')) \\ \downarrow i_* & & \downarrow i_! \\ G_X(D') & \xrightarrow{[p^{N^i}][p^{2N^i}]\tilde{C}_{X \setminus D'}^{\text{syn}}} & \mathbf{Z} \times \prod_{i \geq 1} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D')) \\ \downarrow & & \downarrow \\ G_X(D) & \xrightarrow{[p^{N^i}][p^{2N^i}]\tilde{C}_{X \setminus D}^{\text{syn}}} & \mathbf{Z} \times \prod_{i \geq 1} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)). \end{array}$$

Here the Chern classes are associated to N . We skip this index from the notation. The polynomials $P(D_1/X; \tilde{C}_{D_1 \setminus D'_1}^{\text{syn}})$ are given in degree k , $k \geq 0$, by certain polynomials

$$P_k(D_1/X; \tilde{C}_{D_1 \setminus D'_1}^{\text{syn}}) : G_{D_1}(D'_1) \rightarrow \mathcal{K}(2k, \tilde{E}'_n(k)_{D_1}(D'_1))$$

that are equal to 0 for $k = 0$ and, for $k \geq 1$, are obtained by applying the universal polynomials $P_k(U_1; T_0, T_1, \dots, T_{k-1})$ with integer coefficients [21, 1.3], [9, II.4] to the augmented universal Chern classes

$$\tilde{C}_{D_1 \setminus D'_1, i}^{\text{syn}} : G_{D_1}(D'_1) \rightarrow \mathcal{K}(2i, \tilde{E}'_n(i)_{D_1}(D'_1)), \quad 0 \leq i \leq k-1,$$

and to the first Chern class of the conormal sheaf $c_1^{\text{syn}}(\mathcal{N}_{D_1/X}^\vee) \in H^2(D_1, E'_n(1)_{D_1})$. We set $\gamma' = \beta\omega'$, where β is the map defined in the proof of Lemma 6.3.

Theorem 6.8. *Diagram 6.12 commutes in the homotopy category.*

Proof. We start with the upper square.

Lemma 6.9. *(Grothendieck-Riemann-Roch) There exists a homotopy*

$$[p^{2Ni}] \tilde{C}_{X \setminus D', i_*}^{\text{syn}} \simeq [p^{N(i+1)}] i_! \gamma' [p^{2N}] P(D_1/X; \tilde{C}_{D_1 \setminus D'_1}^{\text{syn}}).$$

Proof. The construction of the homotopy is implicit in Gillet's proof of Theorem 3.1 in [12]. Just like in the classical situation one argues by deforming the given closed immersion to a regular zero section of a vector bundle of the right rank.

(1) *The case of zero section*

So we start with a special case of a closed immersion $i : Y \hookrightarrow X$, $Y = D_1$, which is the zero section of a projective bundle $X = \mathbb{P}(\mathcal{N} \oplus \mathcal{O}_Y)$, where \mathcal{N} is an invertible sheaf on Y . The sheaf \mathcal{N} is the conormal sheaf of the immersion. Let $\pi : X \rightarrow Y$ be the projection. Write $D_Y = D' \cap Y$ and $D_X = D'$. We assume that $\pi^{-1}(D_Y) = D_X$. Functoriality of log- K -theory and the projection formula in log- K -theory (Lemma 4.9) yield the following homotopies

$$\tilde{C}_{X \setminus D_X}^{\text{syn}} i_* \simeq \tilde{C}_{X \setminus D_X}^{\text{syn}} i_* i^* \pi^* \simeq \tilde{C}_{X \setminus D_X}^{\text{syn}} ([i_* \mathcal{O}_Y] \wedge \pi^*),$$

where $[i_* \mathcal{O}_Y] \wedge$ refers to the action of the class of $i_* \mathcal{O}_Y$ in $K_0(X)$ on $G_X(D_X)$. The exact sequence

$$(6.13) \quad 0 \rightarrow \pi^* \mathcal{N} \rightarrow \mathcal{O}_X \rightarrow i_* \mathcal{O}_Y \rightarrow 0$$

yields a homotopy $\text{Id} \simeq ([\pi^* \mathcal{N}] + [i_* \mathcal{O}_Y])$. Hence by functoriality and by compatibility of Chern classes with the action of K -theory, as well as by Lemma 5.5 we have

$$\begin{aligned} \tilde{C}_{X \setminus D_X}^{\text{syn}} i_* &\simeq \tilde{C}_{X \setminus D_X}^{\text{syn}} ((\text{Id} - [\pi^* \mathcal{N}]) \wedge \pi^*) \simeq \tilde{C}_{X \setminus D_X}^{\text{syn}} (\pi^* \lambda_{-1}(\mathcal{N}) \wedge \pi^*) \simeq \pi^* \tilde{C}_{Y \setminus D_Y}^{\text{syn}} (\lambda_{-1}(\mathcal{N}) \wedge) \\ &\simeq \pi^* (\tilde{C}_{Y \setminus D_Y}^{\text{syn}} (\lambda_{-1}(\mathcal{N})) \star \tilde{C}_{Y \setminus D_Y}^{\text{syn}}) \simeq \pi^* (\lambda_{-1}(\tilde{C}_{Y \setminus D_Y}^{\text{syn}}(\mathcal{N})) \star \tilde{C}_{Y \setminus D_Y}^{\text{syn}}), \end{aligned}$$

where for a locally free sheaf \mathcal{L} on a scheme Z we wrote $\lambda_{-1}(\mathcal{L}) = [\mathcal{O}_Z] - [\mathcal{L}] \in K_0(Z)$. The equality

$$\tilde{C}_{Y \setminus D_Y}^{\text{syn}} (\lambda_{-1}(\mathcal{N})) \simeq \lambda_{-1}(\tilde{C}_{Y \setminus D_Y}^{\text{syn}}(\mathcal{N}))$$

holds since the classes $\tilde{C}_{Y \setminus D_Y}^{\text{syn}}$ and $[p^{2Ni}] \tilde{C}_Y^{\text{syn}}$ are compatible and we have the Whitney sum formula for the (classical) syntomic-étale classes C_Y^{syn} .

We claim that

$$(6.14) \quad \lambda_{-1}(\tilde{C}_{Y \setminus D_Y}^{\text{syn}}(\mathcal{N})) \star \tilde{C}_{Y \setminus D_Y}^{\text{syn}} \simeq P(Y/X; \tilde{C}_{Y \setminus D_Y}^{\text{syn}}) [p^{2N}] c_1^{\text{syn}}(\mathcal{N}^\vee).$$

This will reduce to a classical result as soon as we put this equation in the right context. Consider the graded ring

$$\prod_{i \geq 1} [G_Y(D_Y), \mathcal{K}(2i, \tilde{E}'_n(i)_Y(D_Y))]$$

with addition and multiplication induced from $\mathcal{K}(2i, \tilde{E}'_n(i)_Y(D_Y))$. Consider the following abelian groups

$$(6.15) \quad \begin{aligned} H^{2i}(Y, G_Y(D_Y), E'_n(i)_Y(D_Y)) &:= [G_Y(D_Y), \mathcal{K}(2i, \tilde{E}'_n(i)_Y(D_Y))], \\ \tilde{H}^*(Y, G_Y(D_Y), E'_n(*)_Y(D_Y)) &:= \mathbf{Z} \times \{1\} \times \prod_{i \geq 1} [G_Y(D_Y), \mathcal{K}(2i, \tilde{E}'_n(i)_Y(D_Y))]. \end{aligned}$$

Recall that [16, Exp.0] $\widetilde{H}^*(Y, G_Y(D_Y), E'_n(*)_Y(D_Y))$ has a structure of a λ -ring with involution, defined as follows. Write an element of $\widetilde{H}^*(Y, G_Y(D_Y), E'_n(*)_Y(D_Y))$ as $[n, x]$ for $x = 1 + x_1 + \dots + x_k + \dots$. Then

$$\begin{aligned} [n, x] + [m, y] &= [n + m, xy], \\ [n, x] \star [m, y] &= [nm, x^m y^n x \star y], \end{aligned}$$

where

$$(1 + \sum_{i \geq 1} x_i) \star (1 + \sum_{i \geq 1} y_i) = 1 + \sum_{k \geq 0} Q_k(x_1, \dots, x_k; y_1, \dots, y_k)$$

for certain universal polynomials Q_k with integral coefficients.

The λ -structure on $\widetilde{H}^*(Y, G_Y(D_Y), E'_n(*)_Y(D_Y))$ is given by the following operations λ^k

$$(6.16) \quad \begin{cases} \lambda^k[0, x] = [0, \lambda^k x], & (\lambda^k x)_n = Q_{k,n}(x_1, \dots, x_n), \\ \lambda^k[n, 1] = [\lambda^k n, 1], \end{cases}$$

where $Q_{k,n}$ are certain universal polynomials with integral coefficients and $\lambda^k n$ on \mathbf{Z} is the canonical one. The involution is $\psi^{-1}[n, 1 + \sum_{i>0} x_i] = [n, 1 + \sum_{i>0} (-1)^i x_i]$.

Now we can rephrase the equality (6.14) as the following equality in the λ -ring $\widetilde{H}^*(Y, G_Y(D_Y), E'_n(*)_Y(D_Y))$ (where we skipped the $Y \setminus D_Y$ subscript).

$$(\lambda_{-1}([1, 1 + C_1^{\text{syn}}(\mathcal{N}^\vee)]) \star [\widetilde{C}_0^{\text{syn}}, 1 + C_1^{\text{syn}} + C_2^{\text{syn}} \dots])_k = -P_k((C_1^{\text{syn}}(\mathcal{N}^\vee)), (\widetilde{C}_0^{\text{syn}}, C_1^{\text{syn}}, \dots, C_k^{\text{syn}})) C_1^{\text{syn}}(\mathcal{N}^\vee).$$

But this is classical [21, 1.3].

By Lemma 6.4 we have $c_1^{\text{syn}}(\pi^* \mathcal{N}^\vee) = c_1^{\text{syn}}(\mathcal{O}(Y)) = i_!(1)$ for $1 \in H^0(Y, F_n^1(0)_X(D_X))$. Note that the projection formula in syntomic-étale cohomology from Lemma 6.3 implies that

$$[p^{N(i-1)}] i_!(1) = i_! \beta \omega' i^* : E'_n(i)_X(D_X) \rightarrow E'_n(i)_X(D_X)[2].$$

Using this and functoriality we get

$$\begin{aligned} [p^{N(i-1)}] \widetilde{C}_{X \setminus D_X}^{\text{syn}} i_* &\simeq \pi^*(P(Y/X; \widetilde{C}_{Y \setminus D_Y}^{\text{syn}})) [p^{2N}] [p^{N(i-1)}] i_!(1) \simeq i_! \gamma' i^* \pi^* [p^{2N}] P(Y/X; \widetilde{C}_{Y \setminus D_Y}^{\text{syn}}) \\ &\simeq i_! \gamma' [p^{2N}] P(Y/X; \widetilde{C}_{Y \setminus D_Y}^{\text{syn}}), \end{aligned}$$

as wanted.

(2) *General case*

We pass to the general case by "deformation to the normal cone". We will now recall some aspects of this construction [9, IV.5]. Consider the blow-up $\pi : W \rightarrow \mathbb{P}_X^1$ along $Y \times \{\infty\}$, $Y = D_1$. Then we have the following deformation diagram.

$$\begin{array}{ccccc} Y_0 = Y & \xrightarrow{i_0=i} & X = W_0 & \xrightarrow{\text{Id}_X} & X \\ \downarrow j_0 & & \downarrow j_0 & & \downarrow j_0 \\ \mathbb{P}_Y^1 & \xrightarrow{\bar{i}} & W & \xrightarrow{\pi} & \mathbb{P}_X^1 \\ \uparrow j_\infty & & \uparrow t & & \uparrow j_\infty \\ Y_\infty = Y & \xrightarrow{i_\infty} & \mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y) & + & \widetilde{X} \xrightarrow{\text{Id}_X} X. \end{array}$$

This diagram is commutative and all the squares are cartesian. Here \widetilde{X} is the blow-up of X at Y , hence $\widetilde{X} = X$. The sheaf $\mathcal{N}_{Y/X}$ is the normal sheaf of Y in X . We write $N_{Y/X}$ for the associated vector bundle which is canonically isomorphic to the normal cone of Y in X . Note that $\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$ is the projectivization of $N_{Y/X}$. The embedding $i_\infty : Y \hookrightarrow W_\infty = \widetilde{X} + \mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$ is the zero-section embedding of Y in $N_{Y/X}$, followed by the canonical open embedding of $N_{Y/X}$ in $\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$. In general, the divisors $\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$ and \widetilde{X} intersect in the scheme $\mathbb{P}(\mathcal{N}_{Y/X}^\vee)$, which is embedded as the hyperplane at infinity in $\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$, and as the exceptional divisor in \widetilde{X} . Since our embedding

$Y \hookrightarrow X$ is of codimension 1, the scheme $\mathbb{P}(\mathcal{N}_{Y/X}^\vee)$ is just $\text{Spec}(W(k))$. Let $\rho : \mathbb{P}_Y^1 \rightarrow Y$ be the projection and $q : W \rightarrow X$ the canonical map. In the above diagram we put log-structures on schemes via the following divisors

$$D_X = D', \quad D_Y = Y \cap D', \quad D_{\mathbb{P}_Y^1} = \rho^{-1}(D_Y) = D_Y \times \mathbb{P}^1, \quad D_W = q^{-1}(D_X), \quad D_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)} = t^{-1}(D_W).$$

They make all the maps in the squares in the above diagram strict, i.e., the log-structures pullback to log-structures.

First, we claim that there is a homotopy

$$(6.17) \quad j_{0!}\gamma' i_! \gamma' [p^{2N}]P(Y_0/W_0; \tilde{C}_{Y_0 \setminus D_Y}^{\text{syn}}) \simeq j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_{X \setminus D_X} i_*.$$

We will show this by deforming both sides to infinity and using the special case treated above. By functoriality of syntomic-étale cohomology and Lemma 6.2 we have

$$j_{0!}\gamma' i_! \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) \simeq j_{0!}\gamma' i_! j_0^* \rho^* \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) \simeq j_{0!}\gamma' j_0^* \bar{i}_! \rho^* \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}).$$

Here and below we skip the divisors from the notation of Chern classes if understood. The projection formula in syntomic-étale cohomology (Lemma 6.3) and functoriality imply that

$$j_{0!}\gamma' i_! \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) \simeq [p^{N(i-1)}] \bar{i}_! \rho^* \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) c_1^{\text{syn}}(j_{0*} \mathcal{O}_{W_0}) \simeq [p^{N(i-1)}] \bar{i}_! \gamma' P(\mathbb{P}_Y^1/W; \tilde{C}_{\mathbb{P}_Y^1}^{\text{syn}}) c_1^{\text{syn}}(j_{0*} \mathcal{O}_{W_0}).$$

Since the divisors W_0 and W_∞ on W are linearly equivalent we have $[j_{0*} \mathcal{O}_{W_0}] = [j_{\infty*} \mathcal{O}_{W_\infty}] = [l_* \mathcal{O}_X] + [t_* \mathcal{O}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}]$ in $K_0(W)$. This implies

$$\begin{aligned} j_{0!}\gamma' i_! \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) &\simeq [p^{N(i-1)}] \bar{i}_! \gamma' P(\mathbb{P}_Y^1/W; \tilde{C}_{\mathbb{P}_Y^1}^{\text{syn}}) c_1^{\text{syn}}(j_{\infty*} \mathcal{O}_{W_\infty}) \\ &\simeq [p^{N(i-1)}] \bar{i}_! \gamma' P(\mathbb{P}_Y^1/W; \tilde{C}_{\mathbb{P}_Y^1}^{\text{syn}}) (c_1^{\text{syn}}(l_* \mathcal{O}_X) + c_1^{\text{syn}}(t_* \mathcal{O}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)})) \end{aligned}$$

The projection formula in syntomic-étale cohomology (Lemma 6.3), the fact that $l^* \bar{i}_! = 0$, Lemma 6.2, and functoriality of Chern classes yield

$$\begin{aligned} j_{0!}\gamma' i_! \gamma' P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) &\simeq l_! \gamma' l^* \bar{i}_! \gamma' P(\mathbb{P}_Y^1/W; \tilde{C}_{\mathbb{P}_Y^1}^{\text{syn}}) \vee t_! \gamma' t^* \bar{i}_! \gamma' P(\mathbb{P}_Y^1/W; \tilde{C}_{\mathbb{P}_Y^1}^{\text{syn}}) \\ &\simeq t_! \gamma' i_{\infty!} j_{\infty}^* \gamma' P(\mathbb{P}_Y^1/W; \tilde{C}_{\mathbb{P}_Y^1}^{\text{syn}}) \simeq t_! \gamma' i_{\infty!} \gamma' P(Y/\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y); \tilde{C}_Y^{\text{syn}}) \end{aligned}$$

We can now apply the computations we did in the special case to the embedding $i_\infty : Y \rightarrow \mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)$ to conclude that

$$j_{0!}\gamma' i_! \gamma' [p^{2N}]P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) \simeq t_! \gamma' [p^{N(i-1)}] \tilde{C}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}^{\text{syn}} i_{\infty,*}$$

Similarly, by functoriality in log- K -theory and Lemma 4.10 we get

$$j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_X^{\text{syn}} i_* \simeq j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_{W_0}^{\text{syn}} i_* j_0^* \rho^* \simeq j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_{W_0}^{\text{syn}} j_0^* \bar{i}_! \rho^*$$

Next, by functoriality of Chern classes and the projection formula in syntomic-étale cohomology from Lemma 6.3 we get

$$\begin{aligned} j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_X^{\text{syn}} i_* &\simeq j_{0!}\gamma' j_0^* [p^{N(i-1)}] \tilde{C}_W^{\text{syn}} \bar{i}_! \rho^* \simeq [p^{2N(i-1)}] \tilde{C}_W^{\text{syn}} \bar{i}_! \rho^* c_1^{\text{syn}}(j_{0*} \mathcal{O}_{W_0}) \\ &\simeq [p^{2N(i-1)}] \tilde{C}_W^{\text{syn}} \bar{i}_! \rho^* c_1^{\text{syn}}(j_{\infty*} \mathcal{O}_{W_\infty}) \simeq [p^{2N(i-1)}] \tilde{C}_W^{\text{syn}} \bar{i}_! \rho^* (c_1^{\text{syn}}(l_* \mathcal{O}_X) \vee c_1^{\text{syn}}(t_* \mathcal{O}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)})) \\ &\simeq t_! \gamma' t^* [p^{N(i-1)}] \tilde{C}_W^{\text{syn}} \bar{i}_! \rho^* \vee l_! \gamma' l^* [p^{N(i-1)}] \tilde{C}_W^{\text{syn}} \bar{i}_! \rho^* \end{aligned}$$

Since $l^* \bar{i}_! = 0$, functoriality of Chern classes and of log- K -theory together with Lemma 4.10 yield

$$\begin{aligned} j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_X^{\text{syn}} i_* &\simeq t_! \gamma' [p^{N(i-1)}] \tilde{C}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}^{\text{syn}} t^* \bar{i}_! \rho^* \vee l_! \gamma' [p^{N(i-1)}] \tilde{C}_X^{\text{syn}} l^* \bar{i}_! \rho^* \\ &\simeq t_! \gamma' [p^{N(i-1)}] \tilde{C}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}^{\text{syn}} i_{\infty*} j_{\infty}^* \rho^* \simeq t_! \gamma' [p^{N(i-1)}] \tilde{C}_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}^{\text{syn}} i_{\infty*} \end{aligned}$$

So we got that $j_{0!}\gamma' i_! \gamma' [p^{2N}]P(Y_0/W_0; \tilde{C}_{Y_0}^{\text{syn}}) \simeq j_{0!}\gamma' [p^{N(i-1)}] \tilde{C}_X^{\text{syn}} i_*$, as wanted.

Lemma 6.10 below and the projective space theorem (Lemma (6.7)) imply that there is a map

$$q_* : \prod_{i \geq 1} \mathcal{K}(2i+2, \mathrm{R}q_* \tilde{E}'_n(i+1)_W(D_W)) \rightarrow \prod_{i \geq 1} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D_X))$$

such that $q_* j_{0!} \beta \simeq [p^{N^i}]$. Hence, applying first q_* then ω_0 to both sides of the equation (6.17) we get

$$[p^{N(i+1)}] i_! \omega' [p^{2N}] P(Y_0/W_0; \tilde{C}_{Y_0}^{\mathrm{syn}}) \simeq [p^{2N^i}] \tilde{C}_X^{\mathrm{syn}} i_*,$$

as wanted. \square

Lemma 6.10. *Set $h_\infty : Y_\infty = Y \times \{\infty\} \hookrightarrow \mathbb{P}_X^1$ and $h_0 : Y_0 = Y \times \{0\} \hookrightarrow \mathbb{P}_X^1$. Then there exists a quasi-isomorphism*

$$E'_n(r)_{\mathbb{P}_X^1}(D_{\mathbb{P}_X^1}) \oplus h_{\infty*} E_n^1(r-1)_{Y_\infty}(D_{Y_\infty})[-2] \simeq \mathrm{R}\pi_* E'_n(r)_W(D_W), \quad r \geq 2,$$

where the map $\pi^* : E'_n(r)_{\mathbb{P}_X^1}(D_{\mathbb{P}_X^1}) \rightarrow \mathrm{R}\pi_* E'_n(r)_W(D_W)$ is the map induced by $\pi : W \rightarrow \mathbb{P}_X^1$ and the complexes E'_n and E_n^1 are not truncated.

Proof. Just to simplify the notation we will assume that the divisor D' is trivial. We claim that we have the following map of distinguished triangles

$$\begin{array}{ccc} (h_{0*} E_n^1(r-1)_{Y_0} \oplus j_{\infty*} E_n^1(r-1)_{X_\infty})[-2] & \longrightarrow & (h_{0*} E_n^1(r-1)_{Y_0} \oplus j_{\infty*} E_n^1(r-1)_{X_\infty} \oplus h_{\infty*} E_n^1(r-1)_{Y_\infty})[-2] \\ \downarrow & & \downarrow \\ E'_n(r)_{\mathbb{P}_X^1} & \xrightarrow{\pi^*} & \mathrm{R}\pi_* E'_n(r)_W \\ \downarrow & & \downarrow \\ E'_n(r)_{\mathbb{P}_X^1}(\mathbb{P}_Y^1 \cup X_\infty) & \xrightarrow[\sim]{\pi^*} & \mathrm{R}\pi_* E'_n(r)_W(\mathbb{P}_Y^1 \cup X_\infty \cup \mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)). \end{array}$$

where the bottom row is a quasi-isomorphism because log-blow-ups do not change syntomic cohomology [29, Theorem 5.10] and étale cohomology.

Indeed, by gluing the Gysin distinguished triangles we get the following two compatible (via the map π_*) weight “exact” sequences.

$$\begin{aligned} h_{\infty*} E_n^2(r-2)_{Y_\infty}[-4] &\xrightarrow{f} i_* E_n^1(r-1)_{\mathbb{P}_Y^1}[-2] \oplus j_{\infty*} E_n^1(r-1)_{X_\infty}[-2] \rightarrow E'_n(r)_{\mathbb{P}_X^1} \xrightarrow{g} E'_n(r)_{\mathbb{P}_X^1}(\mathbb{P}_Y^1 \cup X_\infty) \\ \bar{i}_* j_{\infty*} E_n^2(r-2)_{Y_\infty}[-4] \oplus t_* s_* E_n^2(r-2)_{H_\infty}[-4] &\xrightarrow{f} \bar{i}_* E_n^1(r-1)_{\mathbb{P}_Y^1}[-2] \oplus l_* E_n^1(r-1)_{\bar{X}}[-2] \oplus t_* E_n^1(r-1)_{\mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)}[-2] \\ &\rightarrow E'_n(r)_W \xrightarrow{g} E'_n(r)_W(\mathbb{P}_Y^1 \cup X_\infty \cup \mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)), \end{aligned}$$

where $s : H_\infty \hookrightarrow \mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)$ is the hyperplane at infinity. That is $\mathrm{cofiber}(f) \simeq \mathrm{fiber}(g)$.

To compute the cofiber of the top map f we use the Gysin distinguished triangle

$$j_{\infty*} E_n^2(r-2)_{Y_\infty}[-2] \rightarrow E_n^1(r-1)_{\mathbb{P}_Y^1} \rightarrow E_n^1(r-1)_{\mathbb{P}_Y^1}(Y_\infty)$$

and the projective space theorem from Lemma (6.7)

$$E_n^1(r-1)_{Y_0} \oplus E_n^2(r-2)_{Y_\infty}[-2] \simeq \mathrm{R}p_* E_n^1(r-1)_{\mathbb{P}_Y^1},$$

where $p : \mathbb{P}_Y^1 \rightarrow Y$, to get that

$$\mathrm{cofiber}(f) \simeq h_{0*} E_n^1(r-1)_{Y_0}[-2] \oplus h_{\infty*} E_n^1(r-1)_{X_\infty}[-2].$$

To compute the cofiber of the bottom map f we use the above to get

$$\mathrm{cofiber}(f) \simeq \bar{i}_* j_{0*} E_n^1(r-1)_{Y_0}[-2] \oplus l_* E_n^1(r-1)_{\bar{X}}[-2] \oplus t_* \mathrm{cofiber}(s_* E_n^2(r-2)_{H_\infty}[-2] \rightarrow E_n^1(r-1)_{\mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)})[-2].$$

Next, we use the Gysin distinguished triangle

$$s_* E_n^2(r-2)_{H_\infty}[-2] \rightarrow E_n^1(r-1)_{\mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)} \rightarrow E_n^1(r-1)_{\mathbb{P}(\mathcal{N}_{Y/X}^V \oplus \mathcal{O}_Y)}(H_\infty)$$

and the homotopy quasi-isomorphism

$$E_n^1(r-1)_{Y_\infty} \xrightarrow{\sim} \mathbf{R}p'_* E_n^1(r-1)_{\mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y)}(H_\infty),$$

where $p' : \mathbb{P}(\mathcal{N}_{Y/X}^\vee \oplus \mathcal{O}_Y) \rightarrow Y$, to conclude that

$$\text{cofiber}(f) \simeq \bar{i}_* j_{0*} E_n^1(r-1)_{Y_0}[-2] \oplus l_* E_n^1(r-1)_{\tilde{X}}[-2] \oplus t_* i_{\infty*} E_n^1(r-1)_{Y_\infty}[-2],$$

as wanted. \square

Having Lemma 6.9, we can define a total Chern class map

$$C_{X(D)}^{\text{syn}} : G_X(D) \rightarrow \prod_{i \geq 0} \mathcal{K}(2i, \tilde{E}'_n(i)_X(D))$$

to get a map of homotopy cofiber sequences in the diagram (6.12) but without the $[p^{N^i}]$ factor. We set $C_{X(D),0}^{\text{syn}}$ equal to the composition of the rank map with the map $\mathbf{Z} \rightarrow E'_n(0)$. It suffices now to show that the Chern class map $C_{X(D)}^{\text{syn}}$, multiplied by $[p^{N^i}]$, is homotopic to $[p^{N^i}][p^{2N^i}]C_{X \setminus D}^{\text{syn}}$. Note that, by construction, the map $C_{X(D)}^{\text{syn}}$ makes the lower square in the diagram (6.12) (without the $[p^{N^i}]$ factor) commute and, by functoriality, the same is true of the map $[p^{2N^i}]C_{X \setminus D}^{\text{syn}}$. Since, by Lemma 5.8, the group

$$\text{Hom}_{\mathcal{P}}(i_* G_{D_1}(D'_1)[1], \mathcal{K}(2i, \tilde{E}'_n(i)_X(D)))$$

is annihilated by $[p^{N^i}]$, the maps $[p^{N^i}]C_{X(D)}^{\text{syn}}, [p^{N^i}][p^{2N^i}]C_{X \setminus D}^{\text{syn}}$ are homotopic. This finishes the proof of our theorem. \square

Remark 6.11. Theorem 6.8 is also true for log-étale cohomology $H^*(X(D)_{\text{ét}}, \mathbf{Z}/p^n(*))$. Same proof works. Here $X(D)$ is a Zariski log-scheme over K or \bar{K} . Since $H^*(X(D)_{\text{ét}}, \mathbf{Z}/p^n(*)) \simeq H^*(U_{\text{ét}}, \mathbf{Z}/p^n(*))$, for $U = X \setminus D$, no additional constants are needed. The key Lemma 5.8 follows easily from purity in étale cohomology. Indeed, we have

$$\begin{aligned} \text{Hom}_{\mathcal{P}}(i_* \mathcal{L}, \mathcal{K}(2i, \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)_{X(D)})) &= \text{Hom}_{\mathcal{P}}(i_* \mathcal{L}, \mathcal{K}(2i, \mathbf{R}j_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)_U)) \\ &= \text{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \mathbf{R}i^! \mathbf{R}j_* \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)_U)) = \text{Hom}_{\mathcal{P}}(\mathcal{L}, \mathcal{K}(2i, \mathbf{R}(i^! j_*) \mathbf{R}\varepsilon_* \mathbf{Z}/p^n(i)_U)) = 0, \end{aligned}$$

ε is the projection from the log-étale site to the Nisnevich site, $j : U \hookrightarrow X$, and $i : D_1 \hookrightarrow X$.

6.2.1. Chern classes and Gysin sequences. The above computations imply that Chern classes from motivic cohomology are compatible with Gysin sequences. To state this compatibility we need first to evaluate the twisted Chern classes from the diagram (6.12) on motivic cohomology.

Let $i : Y \hookrightarrow X$ be a closed immersion of regular syntomic schemes over $W(k)$ of codimension one and let D be a divisor on X such that the divisors $D_X = Y \cup D$ and $D_Y = D \cap Y$ have relative simple normal crossings over $W(k)$ and all the irreducible components of D_X, D_Y are regular. Fix m as in Section 5.3. Consider the map

$$\bar{p}_{i,j}^{\text{syn}} : K_j(Y \setminus D_Y, \mathbf{Z}/p^n) \rightarrow H^{2i-j-2}(Y, S'_n(i-1)_Y(D_Y)), \quad j \geq 2,$$

equal to the composition

$$\begin{aligned} [P^j(Y \setminus D_Y), K_{Y \setminus D_Y}] &= [P^j Y, j_* K_{Y \setminus D_Y}] \xrightarrow{P_i(Y/X; \tilde{C}_{Y \setminus D_Y}^{\text{syn}})} [P^j Y, \mathcal{K}(2i-2, \tilde{S}'_n(i-1)_Y(D_Y))] \\ &= H^{-j}(Y, \mathcal{K}(2i-2, \tilde{S}'_n(i-1)_Y(D_Y); \mathbf{Z}/p^n) \xrightarrow{f} H^{2i-j-2}(Y, S'_n(i-1)_Y(D_Y)), \end{aligned}$$

where $j : Y \setminus D_Y \hookrightarrow Y$ is the natural immersion. We skipped the index m in the notation for Chern classes. Similarly, we get the map

$$\bar{p}_{i,j}^{\text{syn}} : K_j(Y \setminus D_Y, \mathbf{Z}/p^n) \rightarrow H^{2i-j-2}(Y, S_n(i-1)_Y(D_Y)), \quad j \geq 2, i \leq p.$$

Lemma 6.12. *Let $j \geq 2$ for p odd and let $j \geq 3$ for $p = 2$.*

- (1) *The map $\bar{p}_{i,j}^{\text{syn}}$ restricts to zero on $F_\gamma^i K_j(Y \setminus D_Y, \mathbf{Z}/p^n)$, $i \geq 2$.*

(2) We have the equality

$$\bar{p}_{i,j}^{\text{syn}} = -(i-1)\bar{c}_{i-1,j}^{\text{syn}} : F_\gamma^{i-1}K_j(Y \setminus D_Y, \mathbf{Z}/p^n) \rightarrow H^{2i-j-2}(Y, S'_n(i-1)_Y(D_Y)).$$

Proof. We have

$$\bar{p}_{i,j}^{\text{syn}} = P_i(c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee); \bar{c}_{0,j}^{\text{syn}}, \bar{c}_{1,j}^{\text{syn}}, \dots, \bar{c}_{i-1,j}^{\text{syn}}),$$

where $P_i(U_1; T_0, T_1, \dots, T_{i-1})$ is a certain universal polynomial with integral coefficients [21, 1.3]. By Lemma 5.3, $\bar{c}_{k,j}^{\text{syn}}(x) = 0$, $k \leq i-1$, for $x \in F_\gamma^i K_j(Y \setminus D_Y, \mathbf{Z}/p^n)$. Hence

$$\bar{p}_{i,j}^{\text{syn}}(x) = P_i(c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee); 0, \dots, 0) = 0, \quad x \in F_\gamma^i K_j(Y \setminus D_Y, \mathbf{Z}/p^n).$$

This proves the first statement of the lemma.

For the second, by the same argument, we have

$$\bar{p}_{i,j}^{\text{syn}}(x) = P_i(c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee); 0, \dots, 0, \bar{c}_{i-1,j}^{\text{syn}}(x)), \quad x \in F_\gamma^{i-1} K_j(Y \setminus D_Y, \mathbf{Z}/p^n).$$

Now the computation is quite formal. By the defining property of the polynomials P_i [21, 1.1.7] and the product formula [16, Exp.V, 6.6.1], we have

$$\begin{aligned} P_i(c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee); 0, \dots, 0, \bar{c}_{i-1,j}^{\text{syn}}(x)) c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee) &= -[(1 + \bar{c}_{i-1,j}^{\text{syn}}(x)) \star \lambda_{-1}(c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee))]^{(i)} \\ &= -[(1 + \bar{c}_{i-1,j}^{\text{syn}}(x)) \star (1 - c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee) + \dots)]^{(i)} \\ &= -(1 + (i-1)/(i-2)! \bar{c}_{i-1,j}^{\text{syn}}(x) c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee) + \dots)^{(i)} \\ &= -(i-1) \bar{c}_{i-1,j}^{\text{syn}}(x) c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee) \end{aligned}$$

Since we can see this as an equality in the ring of polynomials with integral coefficients, we get that

$$P_i(c_1^{\text{syn},m}(\mathcal{N}_{Y/X}^\vee); 0, \dots, 0, \bar{c}_{i-1,j}^{\text{syn}}(x)) = -(i-1) \bar{c}_{i-1,j}^{\text{syn}}(x),$$

as wanted. \square

And here is the compatibility of Chern classes (from motivic cohomology) with Gysin sequences we wanted.

Corollary 6.13. *Let $2b-a \geq 3$, $p^n > 2$. There exists a constant $A = A(d, a, b)$ such that we have the following maps of partial Gysin sequences*

$$\begin{array}{ccccccccccc} K_{2b-a}^{b-1}(Y(D_Y)) & \xrightarrow{A_{i^*}} & K_{2b-a}^b(X(D)) & \xrightarrow{A_{j^*}} & K_{2b-a}^b(X(D_X)) & \xrightarrow{A_\partial} & K_{2b-a-1}^{b-1}(Y(D_Y)) & \xrightarrow{A_{i^*}} & K_{2b-a-1}^b(X(D_X)) \\ \downarrow \gamma' p^{m(2b+3)} (1-b) \bar{c}_{b-1,2b-a}^{\text{syn},m} & & \downarrow p^{3bm} \bar{c}_{b,2b-a}^{\text{syn},m} & & \downarrow p^{3bm} \bar{c}_{b,2b-a}^{\text{syn},m} & & \downarrow \gamma' p^{m(2b+3)} (1-b) \bar{c}_{b-1,2b-a-1}^{\text{syn},m} & & \downarrow p^{3bm} \bar{c}_{b,2b-a-1}^{\text{syn},m} \\ H_1^{a-2}(Y(D_Y), b-1) & \xrightarrow{A_{i_!}} & H^a(X(D), b) & \xrightarrow{A_{j^*}} & H^a(X(D), b) & \xrightarrow{A_\partial} & H_1^{a-1}(Y(D_Y), b-1) & \xrightarrow{A_{i_!}} & H^{a+1}(X(D'), b) \end{array}$$

Here we set

$$\begin{aligned} K_j^i(T(D_T)) &= \text{gr}_\gamma^i K_j(T(D_T), \mathbf{Z}/p^n), \quad H^*(T(D_T), *) = H^*(T, S'_n(*)_T(D_T)), \\ H_1^*(T(D_T), *) &= H^*(T, S_n^1(*)_T(D_T)). \end{aligned}$$

Proof. The top sequence exists because of Lemma 4.13. The constant $A = N(d, b, 2b-a)N(d, b+1, 2b-a)$. The diagram itself is obtained from Theorem 6.8 and Lemma 6.12 (which gives us that $\bar{p}_{b,2b-a}^{\text{syn},m} = (1-b)\bar{c}_{b-1,2b-a}^{\text{syn},m}$ and $\bar{p}_{b,2b-a-1}^{\text{syn},m} = (1-b)\bar{c}_{b-1,2b-a-1}^{\text{syn},m}$). \square

The above lemma and corollary hold for the S , E , and E' cohomologies as well.

REFERENCES

- [1] M. Bertolini, H. Darmon, V. Rotger, *Beilinson-Flach elements and Euler systems II: the Birch and Swinnerton-Dyer conjecture for Hasse-Weil-Artin L-series*, Journal of Algebraic Geometry 24 (2015) 569–604.
- [2] S. Bloch, *Algebraic cycles and higher K-theory*, Adv. in Math. 61 (1986), no. 3, 267–304.
- [3] S. Bloch, H. Esnault, M. Kerz, *p-adic deformation of algebraic cycle classes*, Invent. Math. 195 (2014), 673–722.
- [4] S. Bloch, K. Kato, *p-adic étale cohomology*. Publ. IHES No. 63 (1986), 107–152.
- [5] A.K. Bousfield, D. Kan, *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972.
- [6] P. Colmez, W. Nizioł, *Syntomic complexes and p-adic nearby cycles*, arXiv:1505.06471.
- [7] V. Ertl, W. Nizioł, *Syntomic cohomology and p-adic motivic cohomology*, arXiv:1603.01705.
- [8] J.-M. Fontaine and W. Messing, *p-adic periods and p-adic étale cohomology*, Current Trends in Arithmetical Algebraic Geometry (K. Ribet, ed.), Contemporary Math., vol. 67, Amer. Math. Soc., Providence, 1987, 179–207.
- [9] W. Fulton, S. Lang, *Riemann-Roch algebra*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 277. Springer-Verlag, New York, 1985.
- [10] T. Geisser, *Motivic cohomology over Dedekind rings*, Math. Z. 248 (2004), no. 4, 773–794.
- [11] T. Geisser, M. Levine, *The K-theory of fields in characteristic p*, Invent. Math. **139** (2000), no. 3, 459–493.
- [12] H. Gillet, *Riemann-Roch theorems for higher algebraic K-theory*, Adv. Math. **40** (1981), 203–289.
- [13] H. Gillet, C. Soulé, *Descent, motives and K-theory*. J. Reine Angew. Math. 478 (1996), 127–176.
- [14] H. Gillet, C. Soulé, *Filtrations on higher algebraic K-theory*, Algebraic K-theory (Seattle, WA, 1997), 89–148, Proc. Sympos. Pure Math., **67**, Amer. Math. Soc., Providence, RI, 1999.
- [15] M. Gros, *Régulateurs syntomiques et valeur de fonctions L p-adiques I*, Invent. Math. **99** (1990), 293–320.
- [16] *Théorie des intersections et théorème de Riemann-Roch. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6)*. Dirigé par P. Berthelot, A. Grothendieck et L. Illusie. Avec la collaboration de D. Ferrand, J. P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J. P. Serre. Lecture Notes in Mathematics, Vol. 225. Springer-Verlag, Berlin-New York, 1971.
- [17] A. Huber, J. Wildeshaus, *Classical motivic polylogarithm according to Beilinson and Deligne*. Doc. Math. 3 (1998), 27–133 (electronic).
- [18] L. Illusie, *Complexe de de Rham-Witt et cohomologie cristalline*, Ann. ENS, 4e série, tome 12, no 4 (1979), 501–661.
- [19] J. F. Jardine, *Simplicial presheaves*, J. Pure Appl. Algebra **47** (1987), no. 1, 35–87.
- [20] R. De Jeu, *Zagier’s conjecture and wedge complexes in algebraic K-theory*. Compositio Math. 96 (1995), no. 2, 197–247.
- [21] J.P. Jouanolou, *Riemann-Roch sans dénominateurs*. Invent. Math. 11 (1970), 15–26.
- [22] B. Kahn, *K-theory of semi-local rings with finite coefficients and étale cohomology*. K-Theory 25 (2002), no. 2, 99–138.
- [23] K. Kato, *On p-adic vanishing cycles (application of ideas of Fontaine-Messing)*, Algebraic Geometry, Sendai, 1985, Adv. Stud. Pure Math. **10**, North-Holland, Amsterdam-New York, 1987, 207–251.
- [24] M. Kerz, *The Gersten conjecture for Milnor K-theory*. Invent. Math. 175, (2009), no. 1, 1–33.
- [25] M. Kurihara, *A note on p-adic étale cohomology*. Proc. Japan Acad. Ser. A Math. Sci. 63 (1987), no. 7, 275–278.
- [26] M. Levine, *K-theory and motivic cohomology of schemes*, preprint, 2004.
- [27] W. Nizioł, *Cohomology of crystalline representations*. Duke Math. J. 71 (1993), no. 3, 747–791.
- [28] W. Nizioł, *Crystalline Conjecture via K-theory*, Ann. ENS (4) **31** (1998), no. 5, 659–681.
- [29] W. Nizioł, *Toric singularities: log-blow-ups and global resolutions*, J. Algebraic Geom. 15 (2006), 1–29.
- [30] W. Nizioł, *Semistable Conjecture via K-theory*, Duke Math. J. 141 (2008), no. 1, 151–178.
- [31] W. Nizioł, *On uniqueness of p-adic period morphisms*, Pure Appl. Math. Q. 5 (2009), no. 1, 163–212.
- [32] O. Pushin, *Higher Chern classes and Steenrod operations in motivic cohomology*. K-Theory 31 (2004), no. 4, 307–321.
- [33] K. Sato, *Cycle classes for p-adic étale Tate twists and the image of p-adic regulators*. Doc. Math. 18 (2013), 177–247.
- [34] M. Somekawa, *Log-syntomic regulators and p-adic polylogarithms*. K-Theory 17 (1999), no. 3, 265–294.
- [35] C. Soulé, *Operations on étale K-theory. Applications*, Algebraic K-theory I, Lect. Notes Math. **966**, Springer-Verlag, Berlin, Heidelberg and New York, 1982, 271–303.
- [36] R. Thomason, *Algebraic K-theory and étale cohomology*, Ann. ENS **18** (1985), 437–552.
- [37] R. Thomason, Th. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*. The Grothendieck Festschrift, Vol. III, 247–435, Progr. Math., 88, Birkhäuser Boston, Boston, MA, 1990.
- [38] T. Tsuji, *p-adic Hodge theory in the semi-stable reduction case*, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). Doc. Math. 1998, Extra Vol. II, 207–216 (electronic).
- [39] T. Tsuji, *p-adic étale and crystalline cohomology in the semistable reduction case*, Invent. Math. **137** (1999), no. 2, 233–411.
- [40] T. Tsuji, *On p-adic nearby cycles of log smooth families*. Bull. SMF 128 (2000), no. 4, 529–575.
- [41] T. Tsuji, *On the maximal unramified quotients of p-adic étale cohomology groups and logarithmic Hodge-Witt sheaves*. Kazuya Kato’s fiftieth birthday. Doc. Math. 2003, Extra Vol., 833–890 (electronic).

CNRS, UMPA, ÉCOLE NORMALE SUPÉRIEURE DE LYON, 46 ALLÉE D'ITALIE, 69007 LYON, FRANCE
E-mail address: wieslawa.niziol@ens-lyon.fr