

TWO POLYGRAPHIC PRESENTATIONS OF PETRI NETS

Yves Guiraud¹

Abstract: This document gives an algebraic and two polygraphic translations of Petri nets, all three providing an easier way to describe reductions and to identify some of them. The first one sees places as generators of a commutative monoid and transitions as rewriting rules on it: this setting is totally equivalent to Petri nets, but lacks any graphical intuition. The second one considers places as 1-dimensional cells and transitions as 2-dimensional ones: this translation recovers a graphical meaning but raises many difficulties since it uses explicit permutations. Finally, the third translation sees places as degenerated 2-dimensional cells and transitions as 3-dimensional ones: this is a setting equivalent to Petri nets, equipped with a graphical interpretation.

Outline

In this document, we study Petri nets in order to give two possible polygraphic presentations for them. This work follows Albert Burroni's intuitions: many computer science and proof theory objects have natural translations into polygraphs. These are topology-flavoured objects consisting of collections of directed cells of various dimensions, equipped with a rich algebraic structure.

In section 1, we recall some basic facts about Petri nets, describe their representations and associate them reduction graphs, equipped with a relation that identifies paths that intuitively represent the same sequence of operations.

In section 2, we recall a known algebraic account of Petri nets: they correspond to commutative word rewriting systems (or presentations of commutative monoids) and both objects generate the same reduction graph. Furthermore, in the latter, reductions have a name, which makes easier the definition of a relation between similar paths. We prove a new result concerning stating that this relation is the same as the one defined for Petri nets. All these facts are detailed in theorem 2.7.

In section 3, we craft a 2-dimensional object, a 2-polygraph, in which reductions of a Petri net can be translated. This result is due to Albert Burroni and is formulated as theorem 3.13. We go beyond and study the links between the relation on Petri nets paths and two relations on 2-arrows of the 2-polygraph: the first one corresponds to the relation on the Petri net, while the second one tries to solve the difficulties raised by the presence of explicit permutations in the 2-polygraph. The study of these properties is only started here: much more work will be necessary to totally solve the encountered problems.

Finally, in section 4, we give a new, more natural polygraphic way to faithfully describe Petri nets. We prove that they correspond to 3-polygraphs with one cell in dimension 0 and no cell in dimension 1. Furthermore, both objects generate the same reduction graph, with the same equivalence relation on paths. This is the main result, theorem 4.14.

¹Institut de mathématiques de Luminy, Marseille, France - guiraud@iml.univ-mrs.fr

1 Basic notions on Petri nets

This section briefly recalls the basic notions about Petri nets: the definitions of a net, of its markings and the usual associated graphical representations. It should be noted that there exist many possible definitions of Petri nets, but a simple one has been chosen for this study. More of them can be found in [Murata 1989] for example.


Definition 1.1. A *Petri net* is a quadruple $N = (X, T, w, w')$ made of two finite sets, X and T , and two maps, $w : X \times T \rightarrow \mathbb{N}$ and $w' : T \times X \rightarrow \mathbb{N}$. The elements of X and T are respectively called *places* and *transitions*, while the maps w and w' are the *weights*. Beside this set-theoretic definition, Petri nets are usually encountered as graphical objects. A decorated graph is associated to a given net $N = (X, T, w, w')$ as follows:

0. Its objects are the places and the transitions. Places are pictured as circles, while transitions are represented by double bars.
1. If x is a place and α a transition, there is an arrow from x to α whenever $w(x, \alpha) > 0$ and one from α to x whenever $w'(\alpha, x) > 0$. Such arrows are decorated with the corresponding weight, either $w(x, \alpha)$ or $w'(\alpha, x)$.

Example 1.2. Let us consider the Petri net $N = (X, T, w, w')$ where $X = \{x, y, z\}$, $T = \{\alpha, \beta\}$ and the non-zero values of w and w' are given by:

$$w(x, \alpha) = 1, \quad w(y, \beta) = 2, \quad w'(\alpha, y) = w'(\alpha, z) = w'(\beta, z) = 1.$$


Following the given graph construction recipe, this representation is built for N :



So far, only the hardware part of a Petri net has been represented. On top of this one, the states of the automaton are described:

Definition 1.3. Let $N = (X, T, w, w')$ be a Petri net. A *marking* of N is a map from the set X of places to the set \mathbb{N} of natural numbers. The set of all markings of N is denoted by $\mathcal{M}(N)$. A given marking $\mu : X \rightarrow \mathbb{N}$ on a Petri net $N = (X, T, w, w')$ is represented as an extra decoration on the corresponding graph: inside each place x , one puts $\mu(x)$ token(s), pictured as black dots.

Example 1.4. With the same Petri net as in example 1.2, the marking μ defined by $\mu(x) = \mu(y) = 2$ and $\mu(z) = 0$ is represented as follows (thereafter, the weights equal to 1 are removed, together with places labels, in order to make the representations clearer):



Now, the whole static part of Petri nets has been introduced. Their evolutions are described as follows:


Definition 1.5. Let $N = (X, T, w, w')$ be a Petri net and let α be a transition in T . The *reduction relation associated to α* is the binary relation \rightarrow_α on markings of N , defined by $\mu \rightarrow_\alpha \nu$ if, for every place x in X , both following conditions hold:

$$\begin{cases} \mu(x) \geq w(x, \alpha), \\ \nu(x) = \mu(x) - w(x, \alpha) + w'(\alpha, x). \end{cases}$$

The union of all the relations \rightarrow_α , for all the transitions α , is denoted by \rightarrow_T . The reflexive and transitive closure of \rightarrow_T is denoted by \twoheadrightarrow_T and called the *reachability relation*.

The relation \rightarrow_α associated to a transition α has a graphical interpretation. The first condition checks if the marking μ has at least $w(x, \alpha)$ tokens in each place x . In that case, the second condition tells that ν is entirely determined this way: in each place x , $w(x, \alpha)$ tokens are removed, then $w'(\alpha, x)$ tokens are added.

Example 1.6. Let N be the Petri net of example 1.2 and μ the marking of example 1.4. The graph pictured thereafter displays all the markings of N that can be reached from μ .



2. Petri nets and commutative word rewriting systems

In order to compare Petri nets with the rewriting-flavoured objects to be introduced in the next three sections, the notion of reduction graph appearing in example 1.6 is formalized:

Definition 1.7. Let $N = (X, T, w, w')$ be a Petri net. Its *associated reduction graph* is the graph $G(N)$ defined by:

0. The set of objects of $G(N)$ is the set $\mathcal{M}(N)$ of markings of N .

1. In $G(N)$, there is an arrow from a marking μ to a marking ν for each transition α such that $\mu \rightarrow_\alpha \nu$.

In example 1.6, we have pictured a subgraph of the reduction graph $G(N)$, where N is the Petri net of example 1.2. Let us consider the top-most square. We can see that the two vertical arrows, both labelled by α are "intuitively" the same reduction: indeed, they consume the same tokens and produce the same ones. This is also the case for the two vertical β -labelled arrows. Furthermore, the horizontal and vertical reductions apply on different tokens: there should be some relation between the two sequences α -then- β and β -then- α . Let us define a congruence relation on such reduction paths:

Notation 1.8. Let $N = (X, T, w, w')$ be a Petri net. We denote by \equiv_N the congruence relation on paths of $G(N)$ generated by the identification of subpaths

$$\mu_1 \rightarrow_\alpha \nu_1 \rightarrow_\beta \mu_2 \quad \text{and} \quad \mu_1 \rightarrow_\beta \nu_2 \rightarrow_\alpha \mu_2,$$

such that the following equalities hold for a given marking ρ in $\mathcal{M}(N)$ and for every place x in X :

$$\begin{aligned} \mu_1(x) &= \rho(x) + w(x, \alpha) + w(x, \beta), & \nu_1(x) &= \rho(x) + w'(\alpha, x) + w(x, \beta), \\ \nu_2(x) &= \rho(x) + w(x, \alpha) + w'(\beta, x), & \mu_2(x) &= \rho(x) + w'(\alpha, x) + w'(\beta, x). \end{aligned}$$

One can check that, in the reduction graph of the Petri net of example 1.2, the relation \equiv_N identifies any two paths with same source and same target one can form in the diagram of example 1.6. In each one of the next three sections, we introduce a translation for Petri nets and study how it behaves with respect to this congruence relation.

2 Petri nets and commutative word rewriting systems

In this section, an equivalence between Petri nets and *commutative word rewriting systems* is proved. The underlying idea of the translation is already present in [Caprotti Ferscha Hong 1995] and [Chandler Heyworth 2001] and comes from the following remarks :

- The markings of a Petri net have a commutative monoid structure: the sum is given by addition of the tokens in each place and the empty marking is a neutral element for this operation.
- If α is a transition, then \rightarrow_α is compatible with the commutative monoid structure on markings: if $\mu \rightarrow_\alpha \mu'$, then $\mu + \nu \rightarrow_\alpha \mu' + \nu$ holds for every marking ν .

Definition 2.1. Let X be a set. The *free commutative monoid generated by X* is the set $[X]$ of all finite formal sums of elements of X :

$$a = \sum_{x \in X} a_x \cdot x,$$

2. Petri nets and commutative word rewriting systems

where the a_x are natural numbers that entirely define a . The set $[X]$ is a commutative monoid for the following operation, which admits the empty sum as a neutral element:

$$\sum_{x \in X} a_x \cdot x + \sum_{x \in X} b_x \cdot x = \sum_{x \in X} (a_x + b_x) \cdot x.$$

A (finite) commutative word rewriting system is a pair (X, R) , where X is a (finite) set, called the *alphabet*, and R is a (finite) family of pairs of elements of $[X]$, called the *rules*. If $\alpha = (s(\alpha), t(\alpha))$ is in R , the reduction relation \rightarrow_α it generates is defined by $a \rightarrow_\alpha b$ if there exists some formal sum c such that $a = c + s(\alpha)$ and $b = c + t(\alpha)$. To any commutative word rewriting system (X, R) , one associates a reduction graph $G(X, R)$, defined by:

0. The objects of $G(X, R)$ are the elements of $[X]$.
1. The arrows of $G(X, R)$ are the pairs (c, α) made of an element c of $[X]$ and a rule $\alpha = (s(\alpha), t(\alpha))$ in R . Such an arrow has source $c + s(\alpha)$ and target $c + t(\alpha)$; it can be written $c + \alpha$.

Remark 2.2. The arrows of $G(X, R)$ are contextual applications of rules: indeed, there is an arrow (c, α) in $G(X, R)$ from a to b if and only if $a \rightarrow_\alpha b$. Furthermore, in this case, c is the context of the application of α at a : this is the part that remains unchanged after action of the rule.

Remark 2.3. In [Guiraud 2004(T)], commutative word rewriting systems are seen as presentations by generators and relations of commutative monoids: indeed, such an object defines a commutative monoid which elements are the connected components of its reduction graph. Conversely, every commutative monoid admits a commutative word rewriting system as a presentation: the generators are the elements of the monoid and the relations are given by the "multiplication" table of the sum.

Following the same idea as in section 1, let us define a congruence relation between paths of the reduction graph of a commutative word rewriting system:

Notation 2.4. Let (X, R) be a commutative word rewriting system. The relation $\equiv_{(X, R)}$ is the congruence relation on paths of $G(X, R)$ generated by the identification of squares of the following shape, with α and β in R and c in $[X]$:

$$\begin{array}{ccc} c + s(\alpha) + s(\beta) & \xrightarrow{(c+s(\beta))+\alpha} & c + t(\alpha) + s(\beta) \\ \downarrow (c+s(\alpha))+\beta & & \downarrow (c+t(\alpha))+\beta \\ c + s(\alpha) + t(\beta) & \xrightarrow{(c+t(\beta))+\alpha} & c + t(\alpha) + t(\beta). \end{array}$$

Translations between Petri nets and finite commutative word rewriting systems are defined as follows:

Definition 2.5. Let $N = (X, T, w, w')$ be a Petri net. Its associated finite commutative word rewriting system is denoted by $\Phi(N)$ and defined by:

- The alphabet of $\Phi(N)$ is the set X of places of N .

2. Petri nets and commutative word rewriting systems

- The rules of $\Phi(\mathbb{N})$ are the transitions of \mathbb{N} , seen as pairs $\alpha = (s(\alpha), t(\alpha))$ with:

$$s(\alpha) = \sum_{x \in X} w(x, \alpha) \cdot x \quad \text{and} \quad t(\alpha) = \sum_{x \in X} w'(\alpha, x) \cdot x.$$

Conversely, let (X, R) be a finite commutative word rewriting system. Its *associated Petri net* is denoted by $\Psi(X, R)$ and defined by:

- The places of $\Psi(X, R)$ are the elements of X .
- There is one transition in $\Psi(X, R)$ for each rule in R .
- The weights w and w' are given, on a place x and a transition $\alpha = (s(\alpha), t(\alpha))$, by:

$$w(x, \alpha) = s(\alpha)_x \quad \text{and} \quad w'(\alpha, x) = t(\alpha)_x.$$

Example 2.6. Let us consider the Petri net from example 1.2. The corresponding commutative word rewriting system is the pair (X, R) , where $X = \{x, y, z\}$ and R consists of the two following rewriting rules $\alpha : x \rightarrow y + z$ and $\beta : 2y \rightarrow z$. The marking from example 1.4 corresponds to the formal sum $2x + 2y$. The reduction graph from example 1.6 becomes:

$$\begin{array}{ccc}
 2x + 2y & \xrightarrow{2x+\beta} & 2x + z \\
 \downarrow x+2y+\alpha & & \downarrow x+z+\alpha \\
 x + 3y + z & \xrightarrow{x+y+z+\beta} & x + y + 2z \\
 \downarrow 3y+z+\alpha & & \downarrow y+2z+\alpha \\
 4y + 2z & \xrightarrow{2y+2z+\beta} & 2y + 3z \xrightarrow{3z+\beta} 4z.
 \end{array}$$

One can check that, in this diagram, any two paths with same source and same target are identified by the congruence $\equiv_{(X, R)}$: the translation from Petri nets to commutative word rewriting systems seems to preserve the congruence relation we have defined on Petri nets reduction paths.

The following result proves that, in essence, Petri nets and finite commutative word rewriting systems are the same objects and generate the same reduction graphs:

Theorem 2.7. *For every Petri net \mathbb{N} , the equality $\Psi \circ \Phi(\mathbb{N}) = \mathbb{N}$ holds and the reduction graphs $G(\mathbb{N})$ and $G(\Phi(\mathbb{N}))$ are isomorphic. Furthermore, this isomorphism identifies the congruences $\equiv_{\mathbb{N}}$ and $\equiv_{\Phi(\mathbb{N})}$. Conversely, for every finite commutative word rewriting system (X, R) , the equality $\Phi \circ \Psi(X, R) = (X, R)$ holds and the reduction graphs $G(X, R)$ and $G(\Psi(X, R))$ are isomorphic. Furthermore, this isomorphism identifies the congruences $\equiv_{(X, R)}$ and $\equiv_{\Psi(X, R)}$.*

2. Petri nets and commutative word rewriting systems

Proof. Let us fix $N = (X, T, w, w')$ and prove the equality $\Psi \circ \Phi(N) = N$. The places of $\Psi \circ \Phi(N)$ are the elements of the alphabet of $\Phi(N)$: these are the places of N . The transitions of $\Psi \circ \Phi(N)$ are the rules of $\Phi(N)$: these are the transitions of N . Let us fix a place x in X and a transition α in T . Let us denote by \bar{w} and \bar{w}' the weighting functions of $\Psi \circ \Phi(N)$ and compare them with w and w' . By definition of $\Psi \circ \Phi(N)$:

$$\bar{w}(x, \alpha) = s(\alpha)_x \quad \text{and} \quad \bar{w}'(\alpha, x) = t(\alpha)_x.$$

And by definition of $\Phi(N)$:

$$s(\alpha) = \sum_{y \in X} w(y, \alpha).y \quad \text{and} \quad t(\alpha) = \sum_{y \in X} w'(\alpha, y).y.$$

Invoking the fact that $[X]$ is free, one gets:

$$s(\alpha)_x = w(x, \alpha) \quad \text{and} \quad t(\alpha)_x = w'(\alpha, x).$$

Hence $w = \bar{w}$ and $w' = \bar{w}'$. Now, let us prove that $G(N)$ and $G(\Phi(N))$ are isomorphic graphs. We define a graph morphism φ from the former to the latter. Let μ be a marking of N and let us define an element $\varphi(\mu)$ in $[X]$ this way:

$$\varphi(\mu) = \sum_{x \in X} \mu(x).x.$$

Now, let us consider an arrow $f : \mu \rightarrow \nu$ in $G(N)$. By definition of $G(N)$, this arrow corresponds to a transition α such that $\mu \rightarrow_\alpha \nu$. By definition of the relation \rightarrow_α on markings, this means that:

$$\mu(x) \geq w(x, \alpha) \quad \text{and} \quad \nu(x) = \mu(x) - w(x, \alpha) + w'(\alpha, x).$$

Let us prove that $\varphi(\mu) \rightarrow_\alpha \varphi(\nu)$ is a reduction generated by (X, R) . By definition of φ on markings:

$$\Phi(\mu) = \sum_{x \in X} \mu(x).x \quad \text{and} \quad \Phi(\nu) = \sum_{x \in X} \nu(x).x.$$

Hence, proving $\varphi(\mu) \rightarrow_\alpha \varphi(\nu)$ is equivalent to prove that there exists a c in $[X]$ such that:

$$\sum_{x \in X} \mu(x).x = c + s(\alpha) \quad \text{and} \quad \sum_{x \in X} \nu(x).x = c + t(\alpha),$$

Since $\mu(x) \geq w(x, \alpha)$ for every place x , the following c is well-defined in $[X]$:

$$c = \sum_{x \in X} (\mu(x) - w(x, \alpha)).x.$$

Then:

$$c + s(\alpha) = \sum_{x \in X} (\mu(x) - w(x, \alpha)).x + \sum_{x \in X} w(x, \alpha).x = \sum_{x \in X} \mu(x).x.$$

Furthermore, using the fact that $\nu(x) = \mu(x) - w(x, \alpha) + w'(\alpha, x)$ holds for every x , one gets:

$$c + t(\alpha) = \sum_{x \in X} (\mu(x) - w(x, \alpha)).x + \sum_{x \in X} w'(\alpha, x).x = \sum_{x \in X} \nu(x).x.$$

2. Petri nets and commutative word rewriting systems

Hence $\varphi(\mu) \rightarrow_\alpha \varphi(\nu)$ holds in $[X]$. By definition of $G(\Phi(N))$, this reduction corresponds to an arrow of the form $c + \alpha$, with c in $[X]$, going from $\varphi(\mu)$ to $\varphi(\nu)$ in $G(\Phi(N))$. Let us define $\varphi(f)$ to be this arrow.

Let us define a graph morphism ψ from $G(\Phi(N))$ to $G(N)$ and prove that it is inverse of φ . Let a be an element of $[X]$. Then $\psi(a)$ is defined as the marking $\psi(a)(x) = a_x$ for every place x . Now, let us consider an arrow $c + \alpha$ in $G(\Phi(N))$, which starts at $a = c + s(\alpha)$ and ends at $b = c + t(\alpha)$. Then, for every place x :

$$\psi(a)(x) = a_x = c_x + s(\alpha)_x = c_x + w(x, \alpha).$$

Thus $\psi(a)(x) \geq w(x, \alpha)$. Furthermore:

$$\psi(b)(x) = b_x = c_x + t(\alpha)_x = \psi(a)(x) - w(x, \alpha) + w'(\alpha, x).$$

Hence $\psi(a) \rightarrow_\alpha \psi(b)$ holds in $\mathcal{M}(N)$. This reduction corresponds to an arrow in $G(N)$, which we take as $\psi(c + \alpha)$. Checking that ψ is a left and right inverse for φ is straightforward.

In order to prove that $\varphi(\equiv_N)$ is $\equiv_{\Phi(N)}$, we prove that $\varphi(\equiv_N)$ is included into $\equiv_{\Phi(N)}$ and that $\psi(\equiv_{\Phi(N)})$ is included into \equiv_N . Furthermore, since φ and ψ are graph morphisms, it is sufficient to prove these inclusions on paths of minimal lengths, such as given in the definitions of both congruences.

Hence, let us consider two paths $\mu_1 \rightarrow_\alpha \nu_1 \rightarrow_\beta \mu_2$ and $\mu_1 \rightarrow_\beta \nu_2 \rightarrow_\alpha \mu_2$ in $G(N)$ such that there exists a marking ρ of N that satisfies the following four equalities for every place x :

$$\begin{aligned} \mu_1(x) &= \rho(x) + w(x, \alpha) + w(x, \beta), & \nu_1(x) &= \rho(x) + w'(\alpha, x) + w(x, \beta), \\ \nu_2(x) &= \rho(x) + w(x, \alpha) + w'(\beta, x), & \mu_2(x) &= \rho(x) + w'(\alpha, x) + w'(\beta, x). \end{aligned}$$

Let us denote by c the element $\varphi(\rho)$ of $[X]$. Then φ sends both paths onto the following ones, which are identified by $\equiv_{\Phi(N)}$:

$$c + s(\alpha) + s(\beta) \xrightarrow{(c+s(\beta))+\alpha} c + t(\alpha) + s(\beta) \xrightarrow{(c+t(\alpha))+\beta} c + t(\alpha) + t(\beta)$$

and:

$$c + s(\alpha) + s(\beta) \xrightarrow{(c+s(\alpha))+\beta} c + s(\alpha) + t(\beta) \xrightarrow{(c+t(\beta))+\alpha} c + t(\alpha) + t(\beta).$$

Then, let us consider two paths in $G(\Phi(N))$ written as above, for a given c in $[X]$. Let us denote by ρ the marking $\psi(c)$. Then, if the four markings μ_1, μ_2, ν_1 and ν_2 are defined as above, the graph morphism ψ sends both paths of $G(\Phi(N))$ onto $\mu_1 \rightarrow_\alpha \nu_1 \rightarrow_\beta \mu_2$ and $\mu_1 \rightarrow_\beta \nu_2 \rightarrow_\alpha \mu_2$: these two paths are identified by \equiv_N .

Conversely, let us consider a finite commutative word rewriting system (X, R) and prove that the equality $\Phi \circ \Psi(X, R) = (X, R)$ holds. By definition of the rewriting system $\Phi \circ \Psi(X, R)$, its alphabet is the set of places of $\Psi(X, R)$: this is the alphabet of (X, R) . The rules in $\Phi \circ \Psi(X, R)$ are the pairs $(\bar{s}(\alpha), \bar{t}(\alpha))$ for each transition α in $\Psi(X, R)$, where:

$$\bar{s}(\alpha) = \sum_{x \in X} w(x, \alpha).x \quad \text{and} \quad \bar{t}(\alpha) = \sum_{x \in X} w'(\alpha, x).x.$$

Furthermore, each transition α in $\Psi(X, R)$ comes from a rule $(s(\alpha), t(\alpha))$ in R and:

$$w(x, \alpha) = s(\alpha)_x \quad \text{and} \quad w'(\alpha, x) = t(\alpha)_x.$$

Thus, $\bar{s}(\alpha) = s(\alpha)$ and $\bar{t}(\alpha) = t(\alpha)$, so that the set of rules of $\Phi \circ \Psi(X, R)$ is R . Hence, the two commutative word rewriting systems (X, R) and $\Phi \circ \Psi(X, R)$ are the same.

Let us prove that the two graphs $G(X, R)$ and $G(\Psi(X, R))$ are isomorphic. Since $\Psi(X, R)$ is a Petri net, we already know that $G(\Psi(X, R))$ is isomorphic to $G(\Phi \circ \Psi(X, R))$: this graph is $G(X, R)$ since the equality $\Phi \circ \Psi(X, R) = (X, R)$ holds. Furthermore, this graph isomorphism is defined the same way as φ and ψ in the first part of the proof. Hence $\varphi(\equiv_{\Psi(X, R)})$ is equal to $\equiv_{(X, R)}$. If one applies ψ , one gets the equality of both congruences $\equiv_{\Psi(X, R)}$ and $\psi(\equiv_{(X, R)})$.

◇

Remark 2.8. This equivalence between Petri nets and finite commutative word rewriting systems highlights the underlying algebraic structure of the formers: one immediate usage is that every arrow in the reduction graph has an explicit name, such as $x + 2y + \alpha$, giving the context of application of the rule α .

Remark 2.9. Another more concrete concrete usage of the translation was developed in the aforementioned [Caprotti Ferscha Hong 1995] and [Chandler Heyworth 2001]: there, it was described how Gröbner bases can be used to partially solve the reachability problem for Petri nets, when they are seen as commutative word rewriting systems.

Remark 2.10. If N is a Petri net, the definition of \equiv_N is technical but intuitively simple. The unveiling of the intrinsic algebraic structure of Petri nets makes this definition much simpler. Indeed, let us consider a commutative word rewriting system (X, R) and denote by \circ the composition of paths in the graph $G(X, R)$. Note that this amounts at considering the category $\langle G(X, R) \rangle$ freely generated by $G(X, R)$, as it is defined in section 3. Then, the relation $\equiv_{(X, R)}$ can be defined as the congruence on $\langle G(X, R) \rangle$ generated by the following identifications, for any c in $[X]$:

$$(c + t(\alpha) + \beta) \circ (c + s(\beta) + \alpha) \equiv (c + t(\beta) + \alpha) \circ (c + s(\alpha) + \beta).$$

Let us also note that such equations allow the sum of $[X]$ to be naturally extended to reductions: $\alpha + \beta$ will be any side of the given equation for $c = 0$. This is also the idea developed with polygraphs in sections 3 and 4.

From now on, theorem 2.7 grants us the right to consider that a Petri net *is* a finite commutative word rewriting system. In fact, the results to be proved are not limited to the finite case. Hence, thereafter, the name Petri net stands for a commutative word rewriting system. Let us use this new equivalent definition to give a different graphical account of Petri nets.

3 Petri nets as 2-dimensional objects

The goal of this section is to prove that Petri nets have strong links with a certain class of *two-dimensional polygraphs*. The first result presented here, theorem 3.13, is essentially due to Albert Burroni, who gave the idea of the translation. The behaviour of this translation with respect to the congruence on Petri nets reduction paths is new and described in proposition 3.17. A discussion follows on many issues to be studied in future work.

In order to translate Petri nets into polygraphs, we start by the interpretation of the markings of a Petri net (the formal sums of its places) into 1-dimensional objects. Let us recall the some classical notions about graphs, free categories and monoids.

3. Petri nets as 2-dimensional objects

Notation 3.1. If G is a graph, its set of objects is denoted by G_0 and its set of arrows going from an object x to another object y is denoted by $G(x, y)$; for such an arrow f , $s_0(f)$ is the source x of f and $t_0(f)$ its target y . The set of all arrows of G is denoted by G_1 and G itself is often abusively denoted by (G_0, G_1) only, assuming that the source and target mappings are given with G_1 .

Definition 3.2. Let $G = (G_0, G_1)$ be a graph. The *free category generated by G* , denoted by $\langle G \rangle$, is the following (small) category:

0. The objects of $\langle G \rangle$ are the objects of G .
1. The arrows of $\langle G \rangle$, from x to y , are all the finite paths in G going from x to y . Their composition, denoted by \circ , is the concatenation of paths. The empty paths are local identities for this operation.

Such a category is often denoted by $\langle G \rangle = (\langle G \rangle_0, \langle G \rangle_1)$ or just by $(G_0, \langle G \rangle_1)$, assuming that the source and target mappings are given with the data in $\langle G \rangle_1$, together with the identities and composition operations.

Example 3.3. Let $G = (*, X)$ be a graph with only one object ($*$ denotes any single-element set); the set of arrows can be any set X , with source and target being the only possible map from X to $*$. Then the free category $\langle G \rangle$ is the free monoid $\langle X \rangle$ generated by X : more precisely, the set $\langle G \rangle(*, *)$, containing all the arrows of $\langle G \rangle$, equipped with the composition and the identity of $*$, is isomorphic to the free monoid $\langle X \rangle$. A proof can be found in [MacLane 1998], for example.

Definition 3.4. Let \mathcal{C} be a category. Two arrows in \mathcal{C} are *parallel* when they have same source and same target. A *relation* in \mathcal{C} is a pair of parallel arrows of \mathcal{C} . If R is a family of relations in \mathcal{C} , the *quotient of \mathcal{C} by R* is the category denoted by \mathcal{C}/R built this way:


0. The objects of \mathcal{C}/R are the objects of \mathcal{C} .
1. The arrows from x to y in \mathcal{C}/R are the elements of $\mathcal{C}(x, y)$, *modulo* the reflexive-symmetric-transitive closure \equiv_R of the relation \rightarrow_R defined by: $f \rightarrow_R g$ if there exist a relation (u, v) in R and two arrows h and k in \mathcal{C} such that $f = k \circ u \circ h$ and $g = k \circ v \circ h$. The identities of \mathcal{C}/R are the equivalence classes of the identity of \mathcal{C} . The composition of \mathcal{C}/R is induced by the one in \mathcal{C} .

Remark 3.5. The defined object \mathcal{C}/R is only a graph. One must check, through easy computations, that the composition of \mathcal{C} is compatible with \equiv_R : the result of the composition is independent of any choice of representatives. Furthermore, it must be checked that induced composition satisfies the axioms of associativity and left and right units of the category structure.

Example 3.6. Let $G = (*, X)$ be a graph with one object. On $\langle G \rangle$, one defines R to be the family of all relations $(x \circ y, y \circ x)$, for x and y in X . Then $\langle G \rangle/R$ is the free commutative monoid $[X]$ generated by the set X .


Hence, we have a graphical description of $[X]$. However, the main idea behind higher-dimensional rewriting is to replace any equation between n -dimensional objects by a $(n + 1)$ -dimensional object: equalities are replaced by their proofs - this point of view was developed in both [Burroni 1993] and [Baez Dolan

1998]. Following this leading idea, equalities of the form $x \circ y = y \circ x$ are replaced by 2-dimensional cells, pasted between parallel paths in the graph $(*, X)$, such as the following one:



Remark 3.7. In order to achieve commutativity, one may ask that $\tau_{x,y}$ is an isomorphism, with $\tau_{y,x}$ as inverse: in this case, one gets a *categorified* version of the free commutative monoid. Another point of view would be to replace the equalities $\tau_{y,x} \circ \tau_{x,y} = \text{id}_{x \otimes y}$ and $\tau_{x,y} \circ \tau_{y,x} = \text{id}_{y \otimes x}$ by their proofs: these would be 3-dimensional cells. This issue is discussed at the end of this section.

So far, we have described an object with one 0-cell, as many 1-cells as there are in our set X , together with one 2-cell $\tau_{x,y}$ for each pair (x, y) of distinct elements in X . Now, let us consider the rule $\alpha : x \rightarrow y + z$ from example 2.6. Such a rule is also translated as a 2-dimensional cell:



A choice has been made in order to represent the rule α . Indeed, it could have been seen as transforming x into $z+y$, which is equal to $y+z$ in the commutative monoid $[X]$. This is the arbitrary part of the presented 2-polygraphic interpretation of Petri nets: it assumes that, for every element a in $[X]$, a representative has been chosen in $\langle X \rangle$.

Since we must use the axiom of choice, let us apply the equivalent Zermelo theorem and assume, until the end of this section, that, for every Petri net (X, R) , the set X comes equipped with a total order. Then, every element a of $[X]$ has a unique decomposition $a = n_1 \cdot x_1 + \dots + n_k \cdot x_k$, where the n_i are non-zero natural numbers and the x_i are elements of X such that $x_1 < \dots < x_k$.

Notation 3.8. Let X be a set and a an element of $[X]$. Let us denote by $n_1 \cdot x_1 + \dots + n_k \cdot x_k$ the unique decomposition of a . Then \bar{a} denotes the representative $x_1^{n_1} \dots x_k^{n_k}$ of a in $\langle X \rangle$, where x^n is the product in $\langle X \rangle$ of n copies of x .

Until now, we have constructed a composite object $\Sigma = (\Sigma_0, \Sigma_1, \Sigma_2)$, made of sets Σ_i of i -dimensional cells. On top of these three sets, Σ also contains boundaries informations: for example, the cell $\tau_{x,y}$ has source $x \circ y$ and target $y \circ x$, while α has source x and target $y \circ z$.

Such an object is called a *polygraph*: it is the central structure studied in *higher-dimensional rewriting*. Here, the object Σ is a 2-dimensional polygraph or 2-polygraph for short. Its definition is recalled from [Burrone 1993].

Definition 3.9. A 2-polygraph Σ is given by:

- 0. A set Σ_0 of 0-cells.

3. Petri nets as 2-dimensional objects

1. A set Σ_1 of 1-cells, together with two maps $s_0, t_0 : \Sigma_1 \rightarrow \Sigma_0$, called 0-source and 0-target. The arrows of the free category $(\Sigma_0, \langle \Sigma \rangle_1)$ are called 1-arrows. The composition of f followed by g is denoted by $f \star_0 g$ or $g \circ_0 f$ in the general case and $f \otimes g$ when Σ_0 has only one element.
2. A set Σ_2 of 2-cells, together with two maps $s_1, t_1 : \Sigma_2 \rightarrow \langle \Sigma \rangle_1$, called 1-source and 1-target, and such that $s_0 \circ s_1 = s_0 \circ t_1$ and $t_0 \circ s_1 = t_0 \circ t_1$. The first equality gives a map $s_0 : \Sigma_2 \rightarrow \Sigma_0$ and the second one yields $t_0 : \Sigma_2 \rightarrow \Sigma_0$.

Definition 3.10. Let (X, R) be a Petri net, such that X is equipped with a total order. The 2-polygraph associated with (X, R) is $\Sigma^2(X, R)$ defined this way:

0. There is one 0-cell in $\Sigma^2(X, R)$, denoted by $*$.
1. The 1-cells of $\Sigma^2(X, R)$ are the elements of X , with the only possible 0-source and 0-target maps.
2. The 2-cells of $\Sigma^2(X, R)$ consist of all the $\tau_{x,y}$, for $x \neq y$ in X , together with one 2-cell α for each rule in R . The 1-source and 1-target maps are given by:

$$s_1(\tau_{x,y}) = x \otimes y, \quad t_1(\tau_{x,y}) = y \otimes x, \quad s_1(\alpha) = \overline{s(\alpha)}, \quad t_1(\alpha) = \overline{t(\alpha)}.$$

In order to compare a Petri net to its associated 2-polygraph, we define a notion of reduction graph for these objects. The idea is to see every 2-cell of a 2-polygraph as a rewriting rule on 1-arrow, that can be applied in any context: a 2-cell α can be applied on any 1-arrow of the shape $u \otimes s_1(\varphi) \otimes v$, in order to produce the 1-arrow $u \otimes t_1(\varphi) \otimes v$. Let us formalize this idea.

Definition 3.11. Let $\Sigma = (\Sigma_0, \Sigma_1, \Sigma_2)$ be a 2-polygraph. The reduction graph associated to Σ , denoted by $G(\Sigma)$, is defined this way:

0. The objects of $G(\Sigma)$ are the 1-arrows of Σ .
1. The arrows from f to g in $G(\Sigma)$ are the triples (h, φ, k) where h and k are 1-arrows in $\langle \Sigma \rangle_1$ and φ is a 2-cell in Σ_2 such that the following equalities hold:

$$f = h \star_0 s_1(\varphi) \star_0 k \quad \text{and} \quad g = h \star_0 t_1(\varphi) \star_0 k.$$

A triple (h, φ, k) is denoted $h \star_0 \varphi \star_0 k$, and $h \star_0$ (resp. $\star_0 k$) is dropped when h (resp. k) is an identity (an empty path).

We want to prove that the two graphs $G(X, R)$ and $G(\Sigma^2(X, R))$ have strong links. To begin with, let us note that the objects of the graph $G(\Sigma^2(X, R))$ are the elements of the free monoid $\langle X \rangle$, while the objects of the graph $G(X, R)$ are the ones of the free commutative monoid $[X]$. We define $\pi : \langle X \rangle \twoheadrightarrow [X]$ to be the canonical projection.

Lemma 3.12. Let u and v be two elements in $\langle X \rangle$ such that $\pi(u) = \pi(v)$. Then, there exists an arrow f in $G(\Sigma^2(X, R))$ with source u and target v , such that f has a decomposition of the form:

$$f = (u_n \otimes \tau_{x_n, y_n} \otimes v_n) \circ \cdots \circ (u_1 \otimes \tau_{x_1, y_1} \otimes v_1).$$

Proof. Since $\langle X \rangle$ is freely generated by X , the elements u and v uniquely decompose as:

$$u = z_1 \otimes \dots \otimes z_p \quad \text{and} \quad v = z'_1 \otimes \dots \otimes z'_{p'},$$

with the z_i and z'_i in X . Since $\pi(u) = \pi(v)$, the following equality holds in $[X]$:

$$z_1 + \dots + z_p = z'_1 + \dots + z'_{p'}.$$

Hence, since $[X]$ is freely generated by X , we get that $p = p'$ and that there exists a permutation σ in \mathfrak{S}_p such that, for every $i \in \{1, \dots, p\}$, $z'_{\sigma(i)} = z_i$. Let us consider a decomposition of the permutation σ in n transpositions:

$$\sigma = \tau_{i_n} \circ \dots \circ \tau_{i_1},$$

where each i_j is in $\{1, \dots, p-1\}$ and τ_{i_j} is the transposition that exchanges i_j and i_{j+1} . Let us fix the following notations:

$$u_1 = z_1 \otimes \dots \otimes z_{i_1-1}, \quad x_1 = z_{i_1}, \quad y_1 = z_{i_1+1}, \quad v_1 = z_{i_1+2} \otimes \dots \otimes z_p.$$

Then, the arrow $f_1 = u_1 \otimes \tau_{x_1, y_1} \otimes v_1$ of $G(\Sigma^2(X, R))$ has source u and target:

$$z_1 \otimes \dots \otimes z_{i_1-1} \otimes z_{i_1+1} \otimes z_{i_1} \otimes z_{i_1+2} \otimes \dots \otimes z_p.$$

But this element of $\langle X \rangle$ can also be written as $z_{\tau_{i_1}(1)} \otimes \dots \otimes z_{\tau_{i_1}(p)}$. Hence, if we repeat this construction for each τ_{i_j} , we prove, by induction on the length of the decomposition of σ , that the target of the last arrow $f_n = u_n \otimes \tau_{x_n, y_n} \otimes v_n$, associated with τ_{i_n} , is:

$$v = z_{\sigma(1)} \otimes \dots \otimes z_{\sigma(p)}.$$

In conclusion, $f = f_n \circ \dots \circ f_1$ satisfies the required hypotheses. ◇

Now, the main result of this section can be proved. As mentioned earlier, this result formalizes a construction due to Albert Burroni:

Theorem 3.13. *Let (X, R) be a Petri net. The following equalities extend the canonical map π into a surjective functor from the free category $\langle G(\Sigma^2(X, R)) \rangle$ to the free category $\langle G(X, R) \rangle$:*

$$\pi(u \otimes \tau_{x, y} \otimes v) = \text{id}_{\pi(u)+x+y+\pi(v)} \quad \text{and} \quad \pi(u \otimes \alpha \otimes v) = \pi(u) + \pi(v) + \alpha.$$

Proof. The equalities extend π so that it is now defined on every object and arrow of the reduction graph $G(\Sigma^2(X, R))$ and takes its values into the free category $\langle G(X, R) \rangle$. Hence, a classical categorical argument tells us that π uniquely extends into a functor, still denoted by π , from the free category $\langle G(\Sigma^2(X, R)) \rangle$ to the free category $\langle G(X, R) \rangle$. Now, let us prove that π is surjective, which means that both its restrictions on objects and on arrows are surjective. On objects, π is the canonical morphism from the free monoid $\langle X \rangle$ to the free commutative monoid $[X]$, which is surjective.

Let us consider two objects a and b in $\langle G(X, R) \rangle$: they are elements of the free commutative monoid $[X]$. Let f be an arrow in $\langle G(X, R) \rangle$ from a to b . By definition of $G(X, R)$ and of the free category it generates, this means that f uniquely decomposes as:

$$f = (c_k + \alpha_k) \circ \dots \circ (c_1 + \alpha_1),$$

3. Petri nets as 2-dimensional objects

with the c_i in $[X]$ and the α_i in R , such that the following relations hold in $[X]$:

$$c_1 + s(\alpha_1) = a, \quad c_i + t(\alpha_i) = c_{i+1} + s(\alpha_{i+1}), \quad c_k + t(\alpha_k) = b.$$

Let us denote by f_i the arrow $\overline{c_i} \otimes \alpha_i$ in $G(\Sigma^2(X, R))$: it has source $\overline{c_i} \otimes \overline{s(\alpha_i)}$ and target $\overline{c_i} \otimes \overline{t(\alpha_i)}$. Hence, the equalities $\pi(s(f_1)) = a$ and $\pi(t(f_n)) = b$ hold. There remains to link all the f_i in order to conclude. Indeed, the relation $t(f_i) = s(f_{i+1})$ does not necessarily hold for every i , so that f_i and f_{i+1} are not composable in general.

However, the relation $\pi(t(f_i)) = \pi(s(f_{i+1}))$ holds, by assumption, for every i . By application of lemma 3.12, we know that there exist arrows g_1, \dots, g_{k-1} in $\langle G(\Sigma^2(X, R)) \rangle$ such that each one is a composition of arrows of the form $(u \otimes \tau_{x,y} \otimes v)$ and such that the following diagram is an arrow of $\langle G(\Sigma^2(X, R)) \rangle$:

$$\overline{a} \xrightarrow{f_1} \overline{c_1} \otimes \overline{t(f_1)} \xrightarrow{g_1} \overline{c_2} \otimes \overline{s(f_2)} \xrightarrow{f_2} \dots \xrightarrow{g_{k-1}} \overline{c_k} \otimes \overline{s(f_k)} \xrightarrow{f_k} \overline{b}.$$

Finally, from the definition of the functor π , we conclude that:

$$\pi(g_i) = \text{id}_{c_i + t(f_i)} = \text{id}_{c_{i+1} + s(f_{i+1})} \quad \text{and} \quad \pi(f_i) = c_i + \alpha_i.$$

Hence $\pi(f_k \circ g_k \circ \dots \circ g_1 \circ f_1) = b$, so that π is a surjective functor. ◇

So far, we have built a new graphical object $G(\Sigma^2(X, R))$ in which every path represents a possible evolution of the Petri net (X, R) and in which every possible evolution has a representative.

But $G(\Sigma^2(X, R))$ is not the natural object one would build from the 2-polygraph $\Sigma^2(X, R)$: indeed, such a polygraph is a presentation of a 2-category, which is a quotient of $\langle G(\Sigma^2(X, R)) \rangle$ by some topology-flavoured relations. Furthermore, we will see that these relations are the ones that identify the intuitively equal paths from examples 1.6 and 2.6.

Here we only define the notion of *free 2-category* generated by a 2-polygraph with one 0-cell, while the complete construction is in [Burroni 1993] and [Métayer 2003]. After the formal algebraic definition, we give the topological intuition that underlies it.

Definition 3.14. Let $\Sigma = (*, \Sigma_1, \Sigma_2)$ be a 2-polygraph with one 0-cell. The *free 2-category* generated by Σ , denoted by $\langle \Sigma \rangle$, is the following 2-polygraph:

0. It has one 0-cell.
1. Its 1-cells are the 1-arrows of Σ , which are the elements of $\langle \Sigma \rangle_1$.
2. Its 2-cells, called *2-arrows*, from u to v are the paths in the reduction graph $G(\Sigma)$, *modulo* the congruence \equiv_{o_1} generated by the following *exchange relations* (where $g \circ f$ is written with f on top of g in order to match the graphical representations to be introduced):

$$\begin{array}{ccc} u \otimes \varphi \otimes (v \otimes s_1(\psi) \otimes w) & & (u \otimes s_1(\varphi) \otimes v) \otimes \psi \otimes w \\ \circ & \equiv & \circ \\ (u \otimes t_1(\varphi) \otimes v) \otimes \psi \otimes w & & u \otimes \varphi \otimes (v \otimes t_1(\psi) \otimes w) \end{array}$$

for every 2-cells φ and ψ , every 1-arrows u , v and w and where \circ denotes the composition of paths in $G(\Sigma)$.

The 2-arrows, collectively denoted by $\langle \Sigma \rangle_2$, are equipped with two compositions: the first one is \circ , the operation yielded by the composition of paths in $G(\Sigma)$; the second one is an extension of \otimes , allowed by the exchange relations, which is defined by functorial extension of:


$$(\mathbf{u} \otimes \varphi \otimes \mathbf{v}) \otimes (\mathbf{u}' \otimes \varphi' \otimes \mathbf{v}') = \begin{array}{c} (\mathbf{u} \otimes s_1(\varphi) \otimes \mathbf{v} \otimes \mathbf{u}') \otimes \varphi' \otimes \mathbf{v}' \\ \circ \\ \mathbf{u} \otimes \varphi \otimes (\mathbf{v} \otimes \mathbf{u}' \otimes t_1(\varphi') \otimes \mathbf{v}') \end{array}$$

Remark 3.15. This definition can be quite obscure and the 2-arrows of the free 2-category are hard to represent with the traditional cellular graphical representation. However, they become really easy to handle when using a dual representation, making the 2-dimensional arrows appear as circuits. Let us explain how this representation is built in the case of a 2-polygraph $\Sigma = (*, \Sigma_1, \Sigma_2)$ with one 0-cell.


Each 1-cell x is drawn as a vertical wire, labelled with x (or with any symbol or color associated to the 1-cell x). A 1-arrow is drawn as the horizontal juxtaposition of the wires representing the 1-cells it is made of. Hence, the empty path id_* is pictured as an empty diagram and the 1-arrow $x_1 \otimes \dots \otimes x_n$ as:




A 2-cell $\varphi : \mathbf{u} \rightarrow \mathbf{v}$ is pictured as a circuit component, with the wires corresponding to \mathbf{u} on top, the ones for \mathbf{v} at the bottom, such as:



A 2-arrow is pictured as a circuit built from the circuit components corresponding to the 2-cells it is made of. The two compositions \otimes and \circ are respectively represented as horizontal juxtaposition and vertical branching:




The circuits are identified *modulo* homeomorphic deformation, which exactly corresponds to the equations of the 2-category structure. For example, the exchange relations are pictured this way:




3. Petri nets as 2-dimensional objects


Example 3.16. Let us consider the Petri net (X, R) from example 1.2. Its associated 2-polygraph is made of one 0-cell $*$, three 1-cells x, y and z and eight 2-cells pictured as:



Then one considers the reduction graph from examples 1.6 and 2.6. As we have seen, all the paths in this diagram can be lifted to representatives in the free category $\langle G(\Sigma^2(X, R)) \rangle$. These representatives are organized in a diagram such as the following one:



In this diagram, all parallel paths only differ by the order of application of the same 2-cells in different parts of the same 1-arrows: hence they are identified by the exchange relations, which means that they become equal in the free 2-category generated by $\Sigma^2(X, R)$. For example, the 2-arrow corresponding to any composite from $x^2 \otimes y^2$ to z^4 is written as $(\beta \otimes z^3) \circ (y \otimes \tau_{z,y} \otimes z^2) \circ (\alpha \otimes \alpha \otimes \beta)$ and is pictured as the following more-readable circuit:



From this example, it seems that the congruences $\equiv_{(X, R)}$ in $\langle G(X, R) \rangle$ and \equiv_{01} in $\langle G(\Sigma^2(X, R)) \rangle$ are linked in some way. For that, we denote by $\mathcal{G}(X, R)$ the quotient category $\langle G(X, R) \rangle / \equiv_{(X, R)}$.

Proposition 3.17. *Let (X, R) be a Petri net. The functor $\pi : \langle G(\Sigma^2(X, R)) \rangle \rightarrow \langle G(X, R) \rangle$ induces a functor $\bar{\pi} : \langle \Sigma^2(X, R) \rangle \rightarrow \mathcal{G}(X, R)$.*

Proof. We have to check that, whenever f and g are parallel arrows in $G(\Sigma^2(X, R))$ such that $f \equiv_{01} g$, we have $\pi(f) \equiv_{(X, R)} \pi(g)$. Let u, v, w be 1-arrows and α, β be 2-cells in $\Sigma^2(X, R)$. Then, by definition

of the functor π , the following four equalities hold:

$$\begin{cases} \pi(\mathbf{u} \otimes \alpha \otimes \mathbf{v} \otimes s_1(\beta) \otimes \mathbf{w}) &= (\mathbf{u} + \mathbf{v} + \mathbf{w} + s_1(\beta)) + \alpha, \\ \pi(\mathbf{u} \otimes \alpha \otimes \mathbf{v} \otimes t_1(\beta) \otimes \mathbf{w}) &= (\mathbf{u} + \mathbf{v} + \mathbf{w} + t_1(\beta)) + \alpha, \\ \pi(\mathbf{u} \otimes s_1(\alpha) \otimes \mathbf{v} \otimes \beta \otimes \mathbf{w}) &= (\mathbf{u} + \mathbf{v} + \mathbf{w} + s_1(\alpha)) + \beta, \\ \pi(\mathbf{u} \otimes t_1(\alpha) \otimes \mathbf{v} \otimes \beta \otimes \mathbf{w}) &= (\mathbf{u} + \mathbf{v} + \mathbf{w} + t_1(\alpha)) + \beta. \end{cases}$$

Thus, the functor π satisfies:

$$\pi \left(\begin{array}{c} \mathbf{u} \otimes \varphi \otimes (\mathbf{v} \otimes s_1(\psi) \otimes \mathbf{w}) \\ \circ \\ (\mathbf{u} \otimes t_1(\varphi) \otimes \mathbf{v}) \otimes \psi \otimes \mathbf{w} \end{array} \right) \equiv_{(X,R)} \pi \left(\begin{array}{c} (\mathbf{u} \otimes s_1(\varphi) \otimes \mathbf{v}) \otimes \psi \otimes \mathbf{w} \\ \circ \\ \mathbf{u} \otimes \varphi \otimes (\mathbf{v} \otimes t_1(\psi) \otimes \mathbf{w}) \end{array} \right).$$

Since π is a functor, we get that $\pi(f) \equiv_{(X,R)} \pi(g)$ for any two parallel f and g such that $f \equiv_{01} g$.


◇

For the moment, we have seen that Petri nets can be translated as 2-polygraphs $\Sigma = (*, \Sigma_1, \Sigma_2 \amalg S_{\Sigma_1})$ where Σ_1 and Σ_2 are finite sets and where S_X denotes the set of all 2-cells $\tau_{x,y}$, with x and y distinct elements in X .

Conversely, given any 2-polygraph of the form $\Sigma = (*, \Sigma_1, \Sigma_2 \amalg S_{\Sigma_1})$ with Σ_1 and Σ_2 finite, one can build a Petri net with alphabet Σ_1 and rules given by the projection through $\pi : \langle \Sigma_1 \rangle \rightarrow [\Sigma_1]$ of the 2-cells of Σ_2 . Furthermore, it can be proved that the two transformations between Petri nets and 2-polygraphs of this form are inverse to each other.

Hence, we could state that Petri nets *are* 2-polygraphs of the form $\Sigma = (*, \Sigma_1, \Sigma_2 \amalg S_{\Sigma_1})$. However, this would be quite excessive since there are much more 2-arrows in $\langle \Sigma \rangle$ than rewriting paths in the corresponding Petri net.

Example 3.18. Once again, let us consider the Petri net from example 1.2 and the path in $\mathcal{G}(X, R)$ given in examples 1.6 and 2.6. In example 3.16, we have already seen a 2-arrow of $\langle \Sigma^2(X, R) \rangle$ representing this reduction path. The following parallel 2-arrows are also possible representatives for this path:




Hence, even if there is a correspondance between Petri nets and 2-polygraphs $\Sigma = (*, \Sigma_1, \Sigma_2 \amalg S_{\Sigma_1})$, both objects do not naturally generate the same reduction graphs since $\langle \Sigma^2(X, R) \rangle$ is bigger than $\mathcal{G}(X, R)$. There are many possible solutions to this problem. One possibility is to add relations between parallel 2-arrows of Σ that represent the same path in the Petri net reduction graph: we are going to sketch such a study in the rest of this section. Another really different solution is studied in section 4, where we use the fact that commutative monoids correspond to a special class of 2-polygraphs.

For the moment, let us consider a 2-polygraph $\Sigma = (*, \Sigma_1, \Sigma_2 \amalg S_{\Sigma_1})$, but where S_{Σ_1} now also contains explicit permutations $\tau_{x,x}$ for every 1-cell x in Σ_1 . This extension does not change the properties studied so far if we extend the functor π with $\pi(\tau_{x,x}) = \text{id}_{x+x}$. We denote by (Σ_1, Σ_2) the corresponding

3. Petri nets as 2-dimensional objects

Petri net. The following result gives a family of relations for some parallel 2-arrows corresponding to the same Petri net reduction. Its proof is straightforward and uses the facts that $\bar{\pi}$ is a functor and maps each $\tau_{x,y}$ onto an identity.

Lemma 3.19. *The functor $\bar{\pi}$ is compatible with the congruence \equiv generated by the following relations, given for all 1-cells x, y and z and every 2-cell α :*



where the generalized explicit permutations used in the third relation are defined inductively from the permutation 2-cells in a graphically intuitive way.

Remark 3.20. The first relation states that, in a given marking of a Petri net, two tokens in the same place are totally indiscernible: for example, one cannot tell if a given transition has consumed one given token or another one in the same place.

Remark 3.21. We conjecture that the congruence \equiv also satisfies the converse property: if f and g are two parallel 2-arrows in $\langle \Sigma \rangle$ such that $\bar{\pi}(f) = \bar{\pi}(g)$, then $f \equiv g$. However, we do not yet have a proof of this fact.

So far, we have a set of equations relating 2-arrows we wish to identify. However this raises a 2-dimensional word problem [Burroni 1993]: given two parallel 2-arrows in $\langle \Sigma \rangle$, are they equal modulo the congruence \equiv or not? One way to build a decision procedure for such a problem is to follow the methodology developed in [Lafont 2003] and [Guiraud 2004] and build a convergent 3-polygraph equivalent to the given equational presentation.

Remark 3.22. Here, we do not recall basic notions about rewriting: they can be found in [Baader Nipkow 1998] for example. Let us say that, for this section, a 3-polygraph is specified by a 2-polygraph equipped with rewriting rules between parallel 2-arrows. These rules are in fact 3-cells, but we postpone all definitions until section 4 since we only need the intuition of it being a "circuit rewriting system" here.

We would like to craft a convergent 3-polygraph for the congruence \equiv on the 2-category $\langle \Sigma \rangle$. However, the fact that 2-cells may have several inputs and several outputs at the same time makes the rewriting study much different than in the already-encountered cases. We give here a possible starting point for future work.


Remark 3.23. For this introduction, we limit ourselves on several points:

- First of all, we only consider the congruence \equiv_0 generated by the last third families: we remove the relations $\tau_{x,x} \equiv \text{id}_{x \otimes x}$ since we still do not know how to handle them. This must be seen as a first step towards the study of \equiv .
- The second limitation is that we assume that Σ_2 does not contain any 2-cell with an empty output: the corresponding Petri net cannot have any transition that do not produce any token.

- Finally, we suppose that every 2-cell in Σ_2 with an empty input has only one output. This is not a real limitation since, in a Petri net, we can replace a transition $\alpha : * \rightarrow y_1 + \dots + y_n$ by two transitions $* \rightarrow z$ and $z \rightarrow y_1 + \dots + y_n$, with z a new place. The Petri net one gets fully simulates the original one.


The idea is the following one: instead of giving an answer to the question $f \equiv g$ directly in $\langle \Sigma \rangle$, we translate 2-arrows of Σ into a 3-polygraph in which we know a decision procedure and such that the translation preserves the congruence \equiv .

Notation 3.24. We denote by $\bar{\Sigma}$ the 2-polygraph with one cell in dimension 0, with Σ_1 as its set of 1-cells and with the following families of 2-cells:



The first family $(\tau_{x,y})$ is indexed by all possible 1-cells x and y ; the second family (δ_x) by every 1-cell x ; the last one (α_i) by every 2-cell $\alpha : x_1 \dots x_m \rightarrow y_1 \dots y_n$ and every i in $\{1, \dots, n\}$.

On top of the 2-category $\langle \bar{\Sigma} \rangle$, we denote by R the family made of the following 3-cells, given for every possible coloration of the wires by 1-cells:



The generalized duplication in the topmost-rightmost family is inductively built from local duplications and local permutations in an inductive way described in [Guiraud 2004] for example. We denote by \equiv_R the congruence relation generated by R on parallel 2-arrows of $\langle \bar{\Sigma} \rangle$.

Following the same method as the one presented in [Guiraud 2004] and using the coloration technique sketched in [Guiraud 2005], one proves that the 3-polygraph $\bar{\Sigma}$ is convergent. Hence, given parallel 2-arrows f and g in $\langle \bar{\Sigma} \rangle$, one can decide whether $f \equiv_R g$ holds or not.

Furthermore, we conjecture here that it is possible to define a 2-functor $\Phi : \langle \Sigma \rangle \rightarrow \langle \bar{\Sigma} \rangle$ such that $f \equiv_o g$ holds if and only if $\Phi(f) \equiv_R \Phi(g)$ holds. Here we define a 2-functor Φ which is a good candidate for this rôle and check the easy part of the claim.

Notation 3.25. We define a 2-functor $\Phi : \langle \Sigma \rangle \rightarrow \langle \bar{\Sigma} \rangle$ by giving its values on the cells of Σ :

0. It sends the only 0-cell of Σ onto itself.
1. It sends each 1-cell x of Σ onto itself.

3. Petri nets as 2-dimensional objects

2. It sends each $\tau_{x,y}$ onto itself and, for every 2-cell $\alpha : x_1 \dots x_m \rightarrow y_1 \dots y_n$ in Σ_2 with $n \geq 2$, we define:


$$\Phi(\alpha) = (\alpha_1 \otimes \dots \otimes \alpha_n) \circ \delta_{x_1 \dots x_m}^n$$

where $\delta_{x_1 \dots x_m}^n$ is the only generalized duplication from $x_1 \dots x_m$ to $(x_1 \dots x_m)^n$ that is in normal form with respect to R .

Then we have:


Proposition 3.26. *The congruence $\Phi(\equiv_0)$ is included into \equiv_R .*

Proof. We check that, for every relation $f \equiv g$ defining \equiv_0 , we have $\Phi(f) \equiv_R \Phi(g)$. This is immediate for the two relations that only involve local permutations. And for the third family of equations:




◇

Example 3.27. Let us consider the Petri net from example 1.2. Its associated 2-polygraph $\bar{\Sigma}$ has the following 2-cells, beside the nine explicit permutations $(\tau_{\xi, \xi'})_{\xi, \xi' \in \{x, y, z\}}$:



Once translated into $\bar{\Sigma}$, the four representatives we have seen of the Petri net reduction of example 1.6 have the following respective normal forms:



If the announced conjecture is true, then this will prove that the first and the third representatives are identified by \equiv_0 and hence by \equiv .

The 2-polygraphic translation of Petri nets we have built in this section has the advantage of having graphical representations that are easy to draw and interpret. However, as we have seen, the explicit way in which it handles the intrinsic commutativity of the net raises many issues we have only started to study here. The non distinction of tokens might be even worse since relations $\tau_{x,x} \equiv \text{id}_{x \otimes x}$ will create many nasty critical pairs when added to a rewriting system. However, future work will be devoted to a thorough study of these polygraphs.

The next section is devoted to a much more natural translation of Petri nets that unveils their intrinsic 3-dimensional nature.

4 Petri nets as 3-dimensional objects

In this section, we prove that Petri nets are exactly 3-dimensional polygraphs with one cell of dimension 0 and no cell of dimension 1. The 2-cells are the places of the net, while the 3-cells are its transitions: there is no need of extra explicit permutation cells. This is due to a topological properties of this class of polygraphs which comes from the folkloric result of algebra, attributed to Hilton:

Lemma 4.1. *Let M be a set equipped with two monoid structures (\bullet, e) and $(\star, 1)$ such that, for every elements x, y, z and t in M , the relation $(x \bullet y) \star (z \bullet t) = (x \star z) \bullet (y \star t)$ holds. Then the two monoid structures are equal and commutative, which means that $e = 1$ and that $x \bullet y = x \star y = y \bullet x = y \star x$.*

Proof. Let us start by proving the equality $e = 1$. Let us apply the hypothesis with $x = t = e$ and $y = z = 1$, which gives $(e \bullet 1) \star (1 \bullet e) = (e \star 1) \bullet (1 \star e)$. On one hand, we have $(e \bullet 1) \star (1 \bullet e) = 1 \star 1 = 1$, since e is a bilateral unit for \bullet and since 1 is a left (or right) unit for \star . But, on the other hand, $(e \star 1) \bullet (1 \star e) = e \bullet e = e$, since 1 is a bilateral unit for \star and since e is a left (or right) unit for \bullet . Hence $e = 1$.

In order to prove that both operations \bullet and \star are the same, let us fix two elements x and y in M . We have the following chain of equalities, using the hypothesis together with the facts that 1 is a bilateral unit for \star and for \bullet :

$$x \bullet y = (x \star 1) \bullet (1 \star y) = (x \bullet 1) \star (1 \bullet y) = x \star y.$$

Finally, we prove that the operation \star is commutative, using the same arguments:

$$x \star y = (1 \bullet x) \star (y \bullet 1) = (1 \star y) \bullet (x \star 1) = y \bullet x = y \star x.$$

◇

Remark 4.2. The proof does not use the associativity of \bullet nor \star . It works with a set with two binary relations such that each one admits a bilateral unit.

Let us translate the lemma 4.1 in our setting:

Corollary 4.3. *Let $\Sigma = (*, \Sigma_1, \Sigma_2)$ be a 2-polygraph with one 0-cell. Then the two compositions \otimes and \circ are equal and commutative on the set $\langle \Sigma \rangle_2(\text{id}_*, \text{id}_*)$, which is the set of all the 2-arrows $\text{id}_* \rightarrow \text{id}_*$ of the free 2-category $\langle \Sigma \rangle$.*

Proof. On $\langle \Sigma \rangle_2(\text{id}_*, \text{id}_*)$ both compositions \otimes and \circ induce a monoid structure. We already know that both structures have the same neutral element, id_{id_*} . Furthermore, the exchange relation gives, for any four f, g, h and k in $\langle \Sigma \rangle_2(\text{id}_*, \text{id}_*)$:

$$(f \otimes g) \circ (h \otimes k) = (f \circ h) \otimes (g \circ k).$$

Then, one applies lemma 4.1 to conclude.

◇

Notation 4.4. Let $\Sigma = (*, \Sigma_1, \Sigma_2)$ be a 2-polygraph with one 0-cell. The 1-arrow id_* is denoted by 0 and, by a slight abuse, so is the 2-arrow id_{id_*} . The common restriction of \circ and \otimes to $\langle \Sigma \rangle_2(0, 0)$ is denoted by $+$.

4. Petri nets as 3-dimensional objects

Remark 4.5. A 2-arrow with source and target equal to 0 is represented as a circuit with no input wire and no output wire. The proof that both compositions are equal and commutative on this kind of 2-arrows corresponds to the following moves:

$$\boxed{f} \boxed{g} = \boxed{f} \quad \boxed{g} = \boxed{g} \quad \boxed{f} = \boxed{g}$$

Explicitly:

$$f \otimes g = (f \otimes 0) \circ (0 \otimes g) = f \circ g = (0 \otimes f) \circ (g \otimes 0) = g \otimes f = (g \otimes 0) \circ (0 \otimes f) = g \circ f.$$

This means that such a special 2-arrow can turn around another one: there is no wire, hence no limitation to their homeomorphic movement.

Using corollary 4.6, we give a polygraphic description of the free commutative monoid generated by a given set:

Proposition 4.6. *Let Σ be a 2-polygraph of the form $(*, \emptyset, \Sigma_2)$. Then the set $\langle \Sigma \rangle_2(0, 0)$ contains all the 2-arrows of $\langle \Sigma \rangle$ and, equipped with the structure $(+, 0)$, is isomorphic to the free commutative monoid $[\Sigma_2]$ generated by Σ_2 .*

Proof. Since there is one 0-cell and no 1-cell in the 2-polygraph Σ , the only 1-arrow of the free 2-category $\langle \Sigma \rangle$ is $\text{id}_* = 0$: indeed, there is only one path in the graph $(*, \emptyset)$ with one object and no arrow, the empty one. Hence, every 2-arrow of $\langle \Sigma \rangle$ starts and ends at 0.

By application of corollary 4.3, we know that both compositions \circ and \otimes are equal and commutative, so that $(\langle \Sigma \rangle_2, +, 0)$ is a commutative monoid. Furthermore, each element of Σ_2 is represented in $\langle \Sigma \rangle_2$: this inclusion induces a unique monoid morphism from $[\Sigma_2]$ to $\langle \Sigma \rangle_2$. This morphism is surjective, since every 2-arrow of $\langle \Sigma \rangle$ is built from 2-cells (elements of Σ_2) using only the operations \otimes and \circ , both equal to $+$. Hence, every 2-arrow f of $\langle \Sigma \rangle$ admits a decomposition:

$$f = \sum_{x \in \Sigma_2} f(x).x,$$

where the $f(x)$ are natural numbers. In order to conclude the proof, one must prove that this decomposition is unique. Let us assume that f has another decomposition:

$$f = \sum_{x \in \Sigma_2} f'(x).x.$$

Let us fix a 2-cell $x \in \Sigma_2$ and assume that $f(x) = f'(x) + k$, with k a natural number. Then:

$$f - f'(x).x = \sum_{y \neq x} f(y).y + k.x = \sum_{y \neq x} f'(y).y.$$

Hence, in the first decomposition of $f - f'(x).x$, there are k copies of the 2-cell x , but there are no in the other. However, in a free 2-category, two arrows are equal if and only if they differ only by a limited number of applications of the rules of associativity, units and exchange for \circ and \otimes : all these operations leave the number of generating 2-cells x unchanged. Hence $k = 0$ and $f(x) = f'(x)$. Finally, there are only a finite number of x such that $f(x) \neq 0$: an induction on this number concludes the proof. \diamond

Now, we have a correspondance between the elements of $[X]$ and the 2-arrows of the free 2-category generated by $(*, \emptyset, X)$. Then, transitions of a Petri net, through their rewriting representation, are translated as 3-cells in a 3-polygraph.

Definition 4.7. A 3-polygraph is a family $\Sigma = (\Sigma_0, \Sigma_1, \Sigma_2, \Sigma_3)$ of sets, equipped with an additional structure of 2-polygraph on $(\Sigma_0, \Sigma_1, \Sigma_2)$ and with a graph structure $s_2, t_2 : \Sigma_3 \rightarrow \langle \Sigma \rangle_2$ such that:

$$s_1 \circ s_2 = s_1 \circ t_2 \quad \text{and} \quad t_1 \circ s_2 = t_1 \circ t_2.$$

Remark 4.8. Usually, the 3-cells are seen as *directed volumes* between parallel circuits (circuits with the same 1-source and the same 1-target).

Let us formalize the translation from Petri nets into 3-polygraphs:

Definition 4.9. Let (X, R) be a Petri net. The 3-polygraph associated to (X, R) , denoted by $\Sigma^3(X, R)$, is the 3-polygraph $(*, \emptyset, X, R)$, where each rewriting rule $\alpha = (a, b)$ is seen as a 3-cell with 2-source the circuit representing a and 2-target the circuit representing b .

Conversely, let $\Sigma = (*, \emptyset, \Sigma_2, \Sigma_3)$ be a 3-polygraph with one 0-cell and no 1-cell. Its associated Petri net is the pair $\mathcal{N}(\Sigma) = (\Sigma_2, \Sigma_3)$.

In order to compare a Petri net and its associated 3-polygraph, a notion of reduction graph is defined, which conveys the idea of reduction under a context - see [Guiraud 2004(T)] for a study of contexts for circuits:

Definition 4.10. Let $\Sigma = (*, \Sigma_1, \Sigma_2, \Sigma_3)$ be a 3-polygraph with one 0-cell. Its *associated reduction graph* is the graph $G(\Sigma)$ defined this way:

0. The objects of $G(\Sigma)$ are the 2-arrows of $\langle \Sigma \rangle_2$.
1. The arrows of $G(\Sigma)$ from u to v are all the triples (f, α, g) , made of two 2-arrows f and g of $\langle \Sigma \rangle_2$ and one 3-cell α of Σ_3 , such that the two following equalities are defined and hold:



These triples are considered *modulo* the following *deformation* equations, given for every possible 2-arrows f, g and h and 3-cell α :



4. Petri nets as 3-dimensional objects

A triple (f, α, g) is denoted by $g \circ \alpha \circ f$, with $(\circ f)$ and/or $(g \circ)$ dropped when f and/or g is an identity. We denote by \star the composition of the free category $\langle G(\Sigma) \rangle$, with $A \star B$ standing A followed by B .


Once again, the reduction graph is not the natural object one associates to a 3-polygraph: we prefer the 3-category it generates. We give a formal definition and, then, its underlying graphical intuition.

Definition 4.11. Let $\Sigma = (*, \Sigma_1, \Sigma_2, \Sigma_3)$ be a 3-polygraph with one 0-cell. The free 3-category generated by Σ is denoted by $\langle \Sigma \rangle$ and is made of the 0, 1 and 2-arrows of Σ , together with a family of 3-arrows which are the paths of the reduction graph $G(\Sigma)$ modulo the congruence \equiv_{Σ} generated by the following exchange relations:


$$\begin{aligned} (A \otimes s_2(B)) \star (t_2(A) \otimes B) &\equiv_{02} (s_2(A) \otimes B) \star (A \otimes t_2(B)), \\ (B \circ s_2(A)) \star (t_2(B) \circ A) &\equiv_{12} (s_2(B) \circ A) \star (B \circ t_2(A)). \end{aligned}$$

These equations allow one to extend the two compositions \otimes and \circ on equivalence classes of paths in the graph $G(\Sigma)$, with $A \otimes B$ being given by either side of the relation \equiv_{02} and $B \circ A$ by either side of \equiv_{12} .


Remark 4.12. Let us give a more graphical account of the free 3-category $\langle \Sigma \rangle$ generated by a 3-polygraph Σ . Its 3-arrows are generated by the 3-cells of Σ seen as blocks:




On these generators, one can use the three following constructors, called compositions:



If they are sliced, these compositions appear this way:



All the constructions are identified modulo the following moves:



This picture contains three families of moves, one for each exchange relation \equiv_{02} , \equiv_{12} and \equiv_{01} , where the relation \equiv_{01} is induced by the deformation relations and the other two exchange relations.

Remark 4.13. In the case of a 3-polygraph Σ with one 0-cell and no 1-cell, there are only two ways to compose 3-arrows, namely $+$ and \star , since \circ and \otimes are the same and denoted by $+$. As a consequence, there is only one family of exchange relations:

$$(A + s_2(B)) \star (t_2(A) + B) \equiv_{\Sigma} (s_2(A) + B) \star (A + t_2(B)).$$


We prove that the reduction graphs of a Petri net and of its associated 3-polygraph are the same. Moreover, the 3-arrows of the 3-category generated by the latter are exactly the equivalence classes of Petri net reductions *modulo* the congruence relation we have defined on them.

Theorem 4.14. *Let (X, R) be a commutative word rewriting system. Then $\mathcal{N}(\Sigma^3(X, R)) = (X, R)$ and the graphs $G(\Sigma^3(X, R))$ and $G(X, R)$ are isomorphic. Furthermore, this isomorphism identifies the congruences $\equiv_{(X, R)}$ and $\equiv_{\Sigma^3(X, R)}$. Conversely, given any 3-polygraph $\Sigma = (*, \emptyset, \Sigma_2, \Sigma_3)$, the equality $\Sigma^3(\mathcal{N}(\Sigma)) = \Sigma$ holds and the graphs $G(\Sigma)$ and $G(\Sigma^3(\mathcal{N}(\Sigma)))$ are isomorphic. Furthermore, this isomorphism identifies the congruences \equiv_{Σ} and $\equiv_{\mathcal{N}(\Sigma)}$.*

Proof. Let us fix a Petri net (X, R) . The equality $\mathcal{N}(\Sigma^3(X, R)) = (X, R)$ is immediate. The objects of both graphs $G(X, R)$ and of $G(\Sigma^3(X, R))$ are the same: the elements of the free commutative monoid $[X]$.

Then, the arrows from u to v in $G(X, R)$ are the $c + \alpha$, made of an element c of $[X]$ and a rule α in R , such that $u = c + s(\alpha)$ and $v = c + t(\alpha)$. To such an arrow $c + \alpha$, we associate the arrow $\varphi(c + \alpha) = (c, \alpha, 0)$ in $G(\Sigma^3(X, R))$.

Conversely, let us consider an arrow (f, α, g) in $G(\Sigma^3(X, R))$. Let us prove graphically that $(f, \alpha, g) = (f + g, \alpha, 0)$, using the fact that all the 2-arrows of $\langle \Sigma^3(X, R) \rangle_2$ have source and target 0:



4. Petri nets as 3-dimensional objects

Let us denote by ψ the map that sends each (f, α, g) onto $f + g + \alpha$ and let us check that ψ is an inverse for φ :

$$\psi \circ \varphi(c + \alpha) = \psi(c, \alpha, 0) = c + 0 + \alpha = c + \alpha.$$

And:

$$\varphi \circ \psi(f, \alpha, g) = \varphi(f + g + \alpha) = (f + g, \alpha, 0) = (f, \alpha, g).$$

Let us prove that $\varphi(\equiv_{(X, R)})$ is included into $\equiv_{\Sigma^3(X, R)}$. For that, we fix c in $[X]$ and α, β in Σ_3 . Then φ sends the following square of $G(X, R)$

$$\begin{array}{ccc} c + s(\alpha) + s(\beta) & \xrightarrow{(c+s(\beta))+\alpha} & c + t(\alpha) + s(\beta) \\ \downarrow (c+s(\alpha))+\beta & & \downarrow (c+t(\alpha))+\beta \\ c + s(\alpha) + t(\beta) & \xrightarrow{(c+t(\beta))+\alpha} & c + t(\alpha) + t(\beta) \end{array}$$

onto the following square of $G(\Sigma^3(X, R))$:

$$\begin{array}{ccc} c + s(\alpha) + s(\beta) & \xrightarrow{(c+s(\beta), \alpha, 0)} & c + t(\alpha) + s(\beta) \\ \downarrow (c+s(\alpha), \beta, 0) & & \downarrow (c+t(\alpha), \beta, 0) \\ c + s(\alpha) + t(\beta) & \xrightarrow{(c+t(\beta), \alpha, 0)} & c + t(\alpha) + t(\beta). \end{array}$$

Using the already-known properties of $G(\Sigma^3(X, R))$, we get the following two equations:

$$\begin{cases} (c + s(\beta), \alpha, 0) \star (c + t(\alpha), \beta, 0) & = & c + ((\alpha + s(\beta)) \star (t(\alpha) + \beta)), \\ (c + s(\alpha), \beta, 0) \star (c + t(\beta), \alpha, 0) & = & c + ((s(\alpha) + \beta) \star (\alpha + t(\beta))). \end{cases}$$

Thus, two paths in $G(X, R)$ identified by $\equiv_{(X, R)}$ are sent by φ on two paths in $G(\Sigma^3(X, R))$ identified by $\equiv_{\Sigma^3(X, R)}$. The inclusion of $\psi(\equiv_{\Sigma^3(X, R)})$ into $\equiv_{(X, R)}$ is proved similarly, starting from the last two equations, in the case $c = 0$, and moving upwards to a square whose paths are identified by $\equiv_{(X, R)}$.

Now, let us fix a 3-polygraph $\Sigma = (\star, \emptyset, \Sigma_2, \Sigma_3)$. The equality $\Sigma^3(\mathcal{N}(\Sigma)) = \Sigma$ is once again immediate. Since $\mathcal{N}(\Sigma)$ is a Petri net, we know that $G(\mathcal{N}(\Sigma))$ is isomorphic to $G(\Sigma^3(\mathcal{N}(\Sigma)))$, which is the same as $G(\Sigma)$. Furthermore, this graph isomorphism is defined the same way as φ and ψ in the first part of the proof. Hence $\varphi(\equiv_{\mathcal{N}(\Sigma)})$ is equal to \equiv_{Σ} . We apply ψ to get the equality between $\equiv_{\mathcal{N}(\Sigma)}$ and $\psi(\equiv_{\Sigma})$. \diamond

This result allows the informal statement "Petri nets *are exactly* the 3-polygraphs with one 0-cell and no 1-cell" for the following reasons:

- There is a correspondance between the presentations, given by the interpretation of places as 2-cells and of transitions as 3-cells.

- Both presentations generate the same reduction graph, so that each one can simulate the evolutions of the other one.
- There is a correspondance between the congruences that identify, in each graph, the paths that only differ by the order of application of the same transitions/3-cells.

Another, more categorical way to formulate this correspondance is to say that the category $\mathcal{G}(X, R)$ generated by a Petri net is isomorphic to the category whose objects and arrows are respectively the 2-arrows and 3-arrows of $\langle \Sigma^3(X, R) \rangle$.

Comments and future directions

We have proved that Petri nets have two natural interpretations in terms of polygraphs. Let us informally compare them.

The first one, using a 2-polygraph, is really convenient to use, since the circuit-like representation is now well-understood and user-friendly. The only difficulty comes with the explicit permutations: one has to choose a way to identify two paths that only differ by permutations. We have discussed possible starting points in order to reach a solution for this issue. And, as we have seen, this is non trivial and is postponed to further work. Nonetheless, this is an important new challenge for 3-dimensional rewriting, since the polygraphs involved provide a new class of rather different examples.

The second polygraphic interpretation we have studied, using a 3-dimensional polygraph, provides, at least theoretically, a better description of the intrinsic algebraic structure of Petri nets: they do not require any extra cell, apart from the ones given with the Petri nets. However, these objects are hard to handle for the moment and this mainly comes from the lack of graphical representations: indeed, the first ones have been constructed in [Guiraud 2005] to represent classical proofs, but they remain hard to produce and handle in a convenient way. For that reason, part of the future work will concern these 3-dimensional representations: the goals are to improve the ones already known, to automatize their production and, maybe, to search for other ones. In the case of Petri nets, the representations should be really interesting since their shape will strangely be close to diagrams used in superstring theory to represent interactions between superstrings.

Let us finish by a more general comment on polygraphs. The results presented here constitute another clue of the expressive power of polygraphs in theoretical computer science, proof theory and universal algebra. Indeed, it is already known that polygraphs generalize word and term rewriting systems, equational presentations of algebraic structures, Reidemeister moves on knots and tangles, formal proofs of classical logic. The interested reader can find more information about the translations of all these objects into polygraphs in the following documents: [Burroni 1993], [Lafont 2003], [Métayer 2003], [Guiraud 2004(T), 2004, 2005].

I wish to thank Albert Burroni and Yves Lafont for many discussions and advices and the referees for their comments that have helped to improve this document.

References

FRANZ BAADER, TOBIAS NIPKOW

Term rewriting and all that, Cambridge University Press, 1998.

JOHN CARLOS BAEZ, JAMES DOLAN

Categorification, ArXiv preprint, 1998.

ALBERT BURRONI

Higher-dimensional word problems with applications to equational logic,
Theoretical Computer Science 115(1), 1993.

OLGA CAPROTTI, ALOIS FERSCHA, HOON HONG

Reachability test in Petri nets by Gröbner bases, RISC report series 95(03), 1995.

ANGIE CHANDLER, ANNE HEYWORTH

Gröbner bases as a tool for Petri net analysis, Proceedings SCI 2001.

YVES GUIRAUD

Présentations d'opérades et systèmes de réécriture, Thèse de doctorat, 2004(T).

Termination orders for 3-dimensional rewriting,

To appear in Journal of Pure and Applied Algebra (2004).

The three dimensions of proofs, To appear in Annals of Pure and Applied Logic (2005).

YVES LAFONT

Towards an algebraic theory of boolean circuits, Journal of Pure and Applied Algebra 184, 2003.

SAUNDERS MACLANE

Categories for the working mathematician, Springer, second edition 1998.

FRANÇOIS MÉTAYER

Resolutions by polygraphs, Theory and Applications of Categories 11(7), 2003.

TADAO MURATA

Petri nets: properties, analysis and applications, Proceedings IEEE 77(4), 1989.